Long-read only assembly of *Drechmeria coniospora* genomes reveals widespread chromosome plasticity and illustrates the limitations of current nanopore methods.

Damien Courtine[1], Jan Provaznik[2], Jerome Reboul[1*], Guillaume Blanc[3] Vladimir Benes[2] and Jonathan J. Ewbank[1+]

[1]Aix Marseille Univ, CNRS, INSERM, CIML, Turing Centre for Living Systems, Marseille, France

[2]European Molecular Biology Laboratory (EMBL), GeneCore, Heidelberg, Germany.

[3]Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille, France

*Current address, Institut de Génetique Moléculaire de Montpellier, Montpellier, France

+ ewbank@ciml.univ-mrs.fr

Abstract

Long read sequencing is increasingly being used to determine eukaryotic genomes. We used nanopore technology to generate chromosome-level assemblies for 3 different strains of *Drechmeria coniospora*, a nematophagous fungus used extensively in the study of innate immunity in *Caenorhabditis elegans*. One natural geographical isolate demonstrated high stability over decades, whereas a second isolate, not only had a profoundly altered genome structure, but exhibited extensive instability. We conducted an in-depth analysis of sequence errors within the 3 genomes and

established that even with state-of-the-art tools, nanopore methods alone are

insufficient to generate eukaryotic genome sequences of sufficient accuracy to merit

inclusion in public databases.

Background

Drechmeria coniospora is an obligate parasitic fungus belonging to the order of

Hypocreales. This fungus forms spores that adhere to the cuticle of a range of different

nematodes to infect them [1]. We adopted *D. coniospora* strain ATCC-96282, derived

from a strain isolated in Sweden, as a model pathogen for *Caenorhabditis elegans* 20

years ago [2]. We have cultured this strain, referred to here as Swe1, continuously

since then, using it to understand innate immune mechanisms in its nematode host

[3,4].

As part of our characterization of the interaction between *D. coniospora* and *C.*

elegans, in 2013, we extracted DNA from our laboratory strain of the time (referred to

here as Swe2), and determined its genome. Despite attempts to complete the

assembly, the Swe2 genome remained fragmented, with an N50 of 3.86 Mb [5]. In

addition to the genome of Swe2, a second *D. coniospora* genome is available (referred

to here as Dan2) [6], derived from a strain related to a Danish isolate (Dan1; Figure 1).

Although corresponding to a chromosome level assembly, this latter genome still

contains large stretches (up to 500 kb) of undetermined sequence. In this study, we

used Oxford Nanopore Technology (ONT) long-read sequencing to assemble

complete fungal genomes. This revealed that the 2 isolates (Swe1 and Dan1) display

strikingly different levels of genomic stability. We provide a detailed analysis that

illustrates the continuing challenges to using only ONT long-read sequencing for

genome assembly. As the genome sequences were of insufficient quality to allow

accurate gene prediction, we polished the genomes using short DNA reads to generate

high-quality sequences, providing a resource for future comparative studies.

Result

An all-against-all *in silico* genome comparison of the 2 publicly available *D. coniospora*

genome sequences, for Dan2 [6] and Swe2 [5], indicated the presence of extensive

genomic rearrangements (Figure 2A). These could reflect real differences or assembly

errors in one or both genomes. We directly confirmed one major rearrangement by

PCR (Figure 2B, C), suggesting that the differences could be real. To characterise this

genomic plasticity, we determined the genomes of 3 strains related to the 2 that had

been sequenced previously (Figure 1). We used ONT nanopore sequencing to

generate long reads and current assembly tools to construct chromosome level

assemblies for all 3 strains (Supplementary Fig. S1, Supplementary Table S1). Manual

curation allowed complete ca. 30 kb mitochondrial genomes to be predicted from the

assemblies generated by Canu [7].

All 3 nuclear genomes were divided in 3 similarly sized chromosomes, an unusual

arrangement for such a fungus, as previously noted by Zhang *et al.* for Dan2 [6]. For

the 2 strains related to Swe2, there was almost complete synteny of their nuclear

genomes. Inspection of the one anomalous region in Swe1 where synteny broke down

revealed that it was supported by only one long (215 kb) read and corresponded to a

local discontinuity in the read coverage, as well as a break in the alignment between

Canu-generated contigs and unitigs. All these factors indicated that this was an

assembly artefact with a contig misassembled on the basis of an individual very long

chimeric read (Supplementary Fig. S2). The same was true for the distinct unique non-

syntenic region of the Swe3 assembly (Supplementary Fig. S3).

These were exceptional cases since the overwhelming majority of chimeric reads were

identified and either trimmed or excluded from the assembly process by Canu

(Supplementary Fig. S4, Supplementary Fig. S5). An in-depth analysis of the Swe1

chimeric reads revealed that a large proportion was in fact the consequence of

sequencing errors. In almost 40% of cases (1010 / 2566), the two regions flanking the

presumptive site of chimerism mapped to within 50 nucleotides of each other on the

corresponding single scaffold. There was no discernible pattern to the distribution of

this interval in the remaining candidate chimeric reads (Supplementary Fig. S6A-B),

nor where there any regions that were more likely to be the site of chimeric junctions

(Supplementary Fig. S6C).

Notably the single chimeric read that escaped censoring, leading to a misassembly of

Swe1, was not identified by the dedicated tool YACRD, but was flagged as anomalous

in reads recalled by Guppy (see Methods). This is an indication of the continuing

improvement to base-calling tools. Also, these specific Swe1 and Swe3 misassemblies

were absent from the corresponding chromosome assemblies produced by the *de*

novo assembler Flye [8] (Figure 3A). This latter, however, introduced other assembly

artefacts, including an erroneous fusion of contigs for the Dan1 assembly. This could

not be ascribed to the inclusion of chimeric reads, but rather appeared to result from

the incorrect treatment of repeat sequences, including telomeric repeats at the

extremity of one of the fused contigs (Figure 3B-D). These results illustrate the interest

of using more than one tool to aid in genome assembly. Therefore, starting with the

Canu-generated sequences, we manually corrected anomalous regions and thereby

produced assemblies for Swe1 and Swe3 that were entirely collinear (Figure 4A).

These 2 genomes have 3 large chromosomes (8.5 Mb, 11.6 Mb, 11.6 Mb), each with

identifiable telomeric [9] and centromeric regions, indicating that the overall genome

structure has remained constant over 20 years of laboratory culture. This allowed us

then to use the Swe1 sequence to scaffold the fragmented Swe2 genome (Figure 4B).

To our great satisfaction, we were able to produce an entirely collinear chromosome

scale assembly. Thus, it appears that there were no assembly errors in the published

Swe2 genome, it was simply incompletely scaffolded. This applies equally to the

genomic regions containing copies of some mitochondrial genes that we previous

suggested might indicate assembly errors [5]. They were revealed to be accurate; *D.*

coniospora has nuclear paralogous copies of 10 mitochondrial protein-coding and 15

tRNA genes (so called *numts* sequences [10]). These results give further support to

the existence of long-term stability of the genome of the Swe2 related strains. A whole

genome comparison between Swe1 and Dan2, however, revealed multiple and

extensive genome rearrangements, involving intra- and inter-chromosomal

translocations and inversions (Figure 4C).

Using the same strategy described above, we assembled and polished the Dan1

genome to give chromosome-level sequences. When we compared Dan1 and Dan2,

we were surprised to find 2 major events of reciprocal exchange of chromosome ends,

and an intra-chromosomal inversion (Figure 4D). These events were supported in a

coherent and consistent manner by all the available data (Supplementary Fig. S7). In

other fungal species, such chromosomal rearrangements have been reported to be the

result of ectopic recombination between non-allelic homologous sequences, including

repeated DNA elements [11,12]. A search of the 50 kb regions flanking each break

point for transposable elements [13] and repetitive DNA families [14], failed to reveal

any significant repeat sequence signature (see Supplementary Methods). As the Dan2

assembly is of high confidence, supported by long reads and optical mapping [6], given

the short time of *in vitro* culture that separates it from Dan1, this suggests that the

genome of the Dan1 isolate is not stable.

In alignments of the sequence of Swe1, generated using only nanopore reads, with

that of Swe2, there were stretches of complete nucleotide identity extending over more

than 25 kb. This is a testament to the general reliability of nanopore sequencing. We

therefore identified the complete set of proteins identical in Swe2 and Dan2

corresponding to single copy, single exon genes (see Methods). These would be

expected to be present in the newly assembled Swe1, Swe3 and Dan1 genomes.

Indeed, using these 305 genes as a query, we could identify homologous sequences

for each in all 3 genomes. Less than 1/6 of the corresponding genes, however, were

predicted to encode full-length proteins in any of the 3 new genomes (Figure 5A). While

nanopore reads are very useful for genome assembly, they suffer from a high error

rate, especially in homopolymer stretches. Sequence quality can be improved using

polishing tools that aim to ameliorate consensus sequences generally by going back

to raw reads and applying integrative algorithms [15]. In our case, applying current best

practices, while providing a very substantial improvement (up to 5-fold in the best

case), did not take the prediction level beyond 82% accuracy. The quality of the

prediction seen with the Dan1 genome was strikingly lower than the other 2 genomes

(Figure 5A, Supplementary Table S1).

Inspection suggested that the majority of errors were in homopolymer sequences, as

expected, with nucleotide insertions and deletions leading to alterations of the reading

frame. To investigate this poor homopolymer predictive performance systematically,

we computed the number of G/C or A/T homopolymer stretches of at least 4

nucleotides for each of the 305 genes. We plotted these values, indicating the

proportion of genes that encoded the expected full-length predicted protein for each of

the 3 genomes. While there was the expected inverse relationship between accuracy

and the number of homopolymer stretches, there were striking exceptions. Curiously

some of these exceptions were specific to a single genome (Figure 5B-D). Further,

and unexpectedly, polishing introduced more nucleotide insertion errors than

deletions, frequently on the basis of tenuous read support. Overall, however, there was

no obvious pattern to explain why errors were introduced, given the underlying reads

used to build the consensus sequence (Supplementary Table S2).

During the inspection of the assembled and polished genomes, we found two other

types of anomalies. The first concerned the regions flanking the nuclear genomic

copies of mitochondrial genes (*numts*), where polishing added short extraneous low

complexity sequences (average length 15 nt, mainly As or Ts), for which, surprisingly

there was no sequence support from the reads used by the assembler (Figure 6A).

This probably arose because of the very high nucleotide similarity between regions of

the nuclear and mitochondrial genomes that extended across more than 25 kb,

including a repeat of 9.8 kb (Supplementary Fig. S8A-B). Notably, despite using high

coverage ONT long reads, we could not establish with absolute certainty the precise

copy number for the unit sequence in the Swe genomes (Supplementary Fig. S8B).

In the second case, for the Swe3 genome, a large (ca. 10 kb) region, with a complex

sequence, well supported by the Canu corrected and trimmed reads, was inexplicably

excluded from the initial Canu assembly and only imprecisely restored by polishing

(Figure 6B-C, Supplementary Fig S8C). Here, while there was no evidence for

repeated DNA elements on both sides of the point of sequence discontinuity, there

was a single such 1.2 kb duplication (Supplementary Fig S8C-D). These few regions

were identified because of discontinuities in the depth of read coverage, which

otherwise was remarkably constant across the complete genomes. With the resolution

of these assembly errors, we were able to generate complete genomes of high overall

structural quality using ONT long reads only.

As explained above, however, these assemblies were not of sufficient sequence

quality to allow accurate gene prediction. Therefore, to extend our analysis, we used

Illumina sequencing to generate very deep short read coverage for the Swe1, Swe3

and Dan1 genomes. This allowed high quality final sequences to be generated for all

3 strains. While short-read-based polishing did not alter the global structure, it allowed

homopolymer length errors to be corrected and the generation of entirely contiguous

chromosome sequences (Supplementary Table S1).

To confirm the correctness of the short-read polished assemblies, we returned to our

305 single copy orthologues. After the short-read polishing, all 305 genes could be

identified in each of the 3 genomes (Supplementary Table S1). We also benchmarked

our successive assemblies using BUSCO that searches for a set of universal single-

copy orthologues (USCO) by sequence similarity. While the initial genome assemblies

gave low scores, with roughly 65% of complete USCOs and 35% fragmented or

missing (Table 1), after long-read polishing the score for complete USCOs increased

up to as high as 97%. Given the demonstrably low quality of the genome sequences

(Figure 5), we investigated the basis of this disparity. We identified among the USCOs

those that corresponded to single exon genes in the Dan2 and Swe2 reference

genomes. These genes were then used as queries for high-stringency searches of the

Dan1, Swe1 and Swe3 genomes at successive steps of assembly and polishing and

the results compared to the results of the corresponding BUSCO analysis. While

BUSCO gave no false negatives, it gave a large number of false positives, except in

the analysis of the short-read polished genomes (Figure 7A). These arose because

BUSCO was not sufficiently sensitive to the presence of short indels. As an example,

the Swe1 gene corresponding to RJ55_06485 had the expected sequence after short-

read polishing. Two errors in homopolymer sequences led to 2 frameshifts in the

unpolished assembly. One of these was corrected by long-read polishing, but for the

other there was an over-compensation, leading to a different frameshift (Figure 7B). In

both assemblies, these errors were compatible with open-reading frames that

collectively reconstituted a close ortholog of RJ55_06485 leading to the erroneous

BUSCO result. As discussed below, this analysis highlights the fact that BUSCO

scores based on sequence alignments are not an appropriate measure for ONT-only

eukaryotic genomes. The BUSCO score rose to nearly 99% after the short-read

polishing. In this case, the figures accurately reflect genome completeness and quality

(Figure 7A). These figures are comparable to those for the previous Dan2 and Swe2

assemblies. The new Swe1, Swe3 and Dan1 genomes therefore represent the starting

point for future detailed analysis to characterise the molecular evolution of *D.*

coniospora.

Table 1: BUSCO results

Strain	Assembly	Complete	Complete: single	Complete: duplicated	Fragmented	Missing
Dan1	Canu curated	820 (62.4%)	820 (62.4%)	0 (0%)	259 (19.7%)	236 (17.9%)
Dan1	Long-read polish	1187 (90.3%)	1187 (90.3%)	0 (0%)	62 (4.7%)	66 (5%)
Dan1	Short-read polish	1297 (98.6%)	1296 (98.6%)	1 (0.1%)	9 (0.7%)	9 (0.7%)
Dan2	[6]	1298 (98.7%)	1297 (98.6%)	1 (0.1%)	8 (0.6%)	9 (0.7%)
Swe1	Canu curated	869 (66.1%)	868 (66%)	1 (0.1%)	243 (18.5%)	203 (15.4%)
Swe1	Long-read polish	1266 (96.6%)	1266 (96.6%)	0 (0%)	21 (1.6%)	28 (2.1%)
Swe1	Short-read polish	1296 (98.6%)	1295 (98.5%)	1 (0.1%)	8 (0.6%)	11 (0.8%)
Swe2	[5]	1296 (98.6%)	1294 (98.4%)	2 (0.2%)	9 (0.7%)	10 (0.8%)
Swe3	Canu curated	859 (65.3%)	858 (65.2%)	1 (0.1%)	243 (18.5%)	213 (16.2%)
Swe3	Long-read polish	1274 (96.9%)	1274 (96.9%)	0 (0%)	17 (1.3%)	24 (1.8%)
Swe3	Short-read polish	1295 (98.5%)	1294 (98.4%)	1 (0.1%)	9 (0.7%)	11 (0.8%)

Percentage of each category of the expected 1315 USCOs for different genome

assemblies. Of the 11 USCOs missing in Swe1 and Swe3, 10 are also absent from

Swe2, and 9 from Dan1 (and Dan2). These are therefore likely to be real gene losses

in *D. coniospora*, so that only 2 USCOs (0.2%) at most are missing.

Discussion and conclusion

Previous genome assemblies for *D. coniospora* required a combination of sequencing

approaches [5,6]. Here, using only long reads and Canu, we produced the first

complete circular mitochondrial genome for *D. coniospora* and were able to generate

chromosome-scale assemblies for the nuclear genome. The rare misassembled

contigs, formed by Canu because of single very long chimeric reads, as previously

described [16], could be detected by read coverage anomalies and comparisons with

unitigs, suggesting that solutions to avoid their creation could be implemented within

Canu. The majority of reads that were flagged as chimeric arose from sequencing or

polishing errors. They reflected a short (<50 bp) discrepancy between the individual

reads and the final sequence. There was no indication of any sequence bias at the

break points of the remaining chimeric reads supporting the notion that these reads

arise from too rapid reloading of the sequencing pore [17].

The use of other genome assembly tools, and the comparison of assembly

discrepancies is an additional method to produce high confidence genomes. Here, we

used Flye that for these genomes required run times that were ten-fold shorter than

Canu. A comparison of the assemblies highlighted ambiguous regions in the genome

that could then be resolved by manual inspection. On the other hand, Flye was

confounded by telomeric repeats. Since telomeres can be identified on the basis of

their sequence, there is also clear room for algorithmic improvement to Flye through

the explicit definition of chromosome ends.

One clear and well-established advantage of using long reads is the possibility of

resolving very extended stretches of complex tandem repeats (VeCTRs) [18] and other

repetitive sequences including centromeres. These correspond to most of the breaks

in the continuity of the published Swe2 genome. In addition to acrocentric regional

centromeres, Zhang *et al*. reported the presence of a vestigial centromere from a

putative chromosomal fusion event [6]. These were also found in the fully assembled

Swe1 and Swe3 genomes, indicating that chromosomal fusions were present in the

common ancestor of the Swe1 and Dan1 strains.

For Swe1, Swe3 and Dan1 we were able to reconstruct complete mitochondrial

genomes, with features typical of fungi of the order Hypocreales. On the other hand,

unlike Dan1 (and Dan2), the nuclear genomes of Swe1 and its derivatives Swe2 and

Swe3, contained different numbers of copies of sequence very similar to parts of their

own mtDNA. This type of event, and more generally repeated regions with long and

nearly identical sequences are more readily detectable with long reads [19], and are

particularly challenging for polishing even with short reads [20].

The duplication of mitochondrial genes in the nuclear genome has been described in

other fungal genomes [10] and must have occurred after the divergence of Dan1 and

Swe1. Despite this genome plasticity, even after 20 years of continuous laboratory

culture the Swe1 and Swe3 genomes were entirely collinear. This contrasts with the

rearrangements seen between the Dan1 and Dan2 genomes that in principal should

be from strains that have had little opportunity to diverge (L. Castrillo, Curator, ARS

Collection of Entomopathogenic Fungal Cultures, personal communication). It will be

interesting in the future to characterize the reasons for the marked difference in

genomic stability between Dan1 and Swe1.

The accuracy of ONT long read sequencing is increasing because of improvements in

the chemistry used, signal detection, as well as base-calling [21]. Despite good read

depth, however, our assemblies were not of sufficient quality at the nucleotide level to

allow accurate gene prediction. Further, we noted that although polishing using only

long reads dramatically increased overall sequence accuracy, it introduced errors

around the *numts*. Similar errors during polishing of near identical sequences has been

noted in ONT-based metagenomic studies [22]. Despite these limitations, research

groups are publishing and submitting to public sequence databases genomes for fungi,

plants and animals based on nanopore sequencing alone (86 for Eukaryotes in

addition to the 134 Bacterial genomes in "Assembly" from GenBank release 236 from

the 2020/02/15). This is problematic as low-quality genome sequences compromise

the accuracy of sequence similarity searches in public databases. On the basis of our

results, a re-analysis of the completeness of these "nanopore-only" genomes is

merited, to confirm that they are indeed low quality. Similar concerns do not apply to

fungal genomes assembled using only long reads generated with Pacific Bioscience

technology [23] as these do not suffer from the intrinsic problem of homopolymer length

errors that we found to be the most significant quality barrier when using ONT reads.

On the basis of our detailed analysis and in line with the consensus regarding *de novo*

assembly with ONT long reads (e.g. [24]), we polished our 3 assemblies with short

reads. This greatly improved their quality.

Regarding the homopolymer sequence errors, as noted above, they were not

consistent across the sequenced genomes; even between Swe1 and Swe3 there were

instances of widely differing rates of errors in orthologous genes, despite very similar

underlying reads. Indeed there was no clear pattern in the inaccuracies, which will

render bioinformatics approaches to remedy this problem more difficult. On the other

hand, the errors were more often over-prediction of homopolymer length, despite

having a majority of reads supporting the correct sequence. It is possible that polishing

tools have not kept pace with improvements in base-calling, leading to an over-

compensation in the inference of homopolymer length.

It is standard practice to check the completeness of *de novo* genome assemblies with

a strategy based on the detection of predicted groups of conserved orthologous

proteins. One popular and much cited tool is BUSCO [25] which was developed before

ONT-based sequencing became prevalent. Since BUSCO relies on *in silico* translation,

small indels can be overlooked as the resulting virtual sequence can be recapitulated

despite a frameshift. This explains the disparity between the BUSCO results and our

own analyses that were deliberately restricted to mono-exonic genes. Contrary to

BUSCO, our analysis indicated that about 1/5 of the genes after long-read polishing

had an incorrect sequence. Current BUSCO-type approaches, based on sequence

similarity and not excluding genes with improbably short introns, cannot be used as a

quality metric for ONT-only assemblies, and are appropriate only after short-read

correction.

In conclusion, nanopore long read sequencing provides a powerful way to assemble

complex genomes with limited manual curation but still fall short of the quality required

to produce publishable eukaryotic genomes. In our case, it has revealed new

information about genome plasticity in *D. coniospora* and provided a backbone that will

permit future detailed study to characterize gene evolution in this important model

fungal pathogen.

Methods

<u>DNA extraction:</u>

D. coniospora spores were cultured in liquid NGMY medium [26] at 37ºC for 5 days.

Fungal DNA was extracted according to a published protocol (from p13 onwards of

[27]) [28], with the following modifications: instead of centrifugation to collect DNA after

precipitation with isopropanol, we recovered the DNA filaments with a glass hook,

washed and dried them as described [29] and resuspended the DNA without agitation

in Tris-EDTA buffer.

<u>Nanopore sequencing library preparation:</u>

Libraries were prepared for sequencing on GridION with the ligation sequencing kit

SQK-LSK109. The GridION sequencing was run on flowcell FLO-MIN106 for 47, 48

and 48 hours using 972, 660 and 610 ng of DNA (for Swe3, Swe1, Dan1 respectively)

and MinKNOW 2.1 v18.06.2.

<u>Illumina sequencing library preparations:</u>

The same DNA samples were used to prepare paired-end libraries with insert size of

circa 680 bp, following the manufacturer's instructions for the kit NEBNext® Ultra™ II

DNA (New England Biolabs Inc. Ipswich, MA, USA). The libraries were sequenced

using an Illumina NextSeq500 system (s/n NB501764).

<u>Basecalling, adaptor trimming and chimeric read detection:</u>

For a first assembly, reads were basecalled at the EMBL using Guppy v1.5.1 (Oxford

Nanopore Technologies). For subsequent polishing, we used Guppy v3.0.3 (with

parameters -c dna_r9.4.1_450bps_hac.cfg), then adaptors were trimmed with

Porechop v0.2.4 [30] with default parameters. YACRD v0.5.1 [31] with the

subcommand chimeric and the option --filter was used to remove chimeric reads.

Whole genome alignments:

Genomes were aligned using LAST v979 [32]. A database was first generated (last-db

-cR01), and then lastal and last-dotplot with default parameters were used to generate

respectively an alignment file and a dot-plot. For the circular visualization of genome

alignments, we used the command lastal with -f BlastTab parameter, then parsed the

alignment to filter out short alignments and generate the links file needed by Circos

[33].

Mapping of long reads:

Validation of genomes during and after assembly involved rounds of read mapping.

Reads were aligned with Minimap2 v2.16r922 [34] (with parameters -ax map-ont). The

resulting mapping file was processed with Samtools v1.9 [35] to obtain a sorted BAM

file (samtools view -bS -q 1 -F 4; samtools sort; samtools index). Mapping results were

visualized with IGV v2.5.0 [36].

Genome assembly:

Assemblies were performed with Canu v1.7 [37] and the parameters useGrid=False,

genomeSize=30m, correctedErrorRate=0.16 with reads basecalled by Guppy v1.5.1.

For the manual curation of the assemblies, we generated whole assembly alignments

and dot-plots of Swe1, Swe2 and Swe3 two by two. For Swe1 and Swe3, Canu contigs

were ordered by synthesizing the results from the 3 possible all-against-all alignments.

To confirm a link between two contigs, we employed the following strategy: when a

contig of the Swe1 assembly spanned two contigs of Swe3, long reads of Swe1 present

in this spanning area were extracted from the Swe1 corrected and trimmed reads

provided by Canu. Then this set of reads was mapped on Swe2 and Swe3 assemblies.

The two targeted contigs of Swe3 were considered 'linked' if different parts of several

unique reads mapped on the two Swe3 contigs ends. If the reads that supported the

link had different mapping orientation (forward or reverse), one contig was

complemented before the last step (see *Solving links between contigs*) to ensure a

correct orientation of the final chromosome.

To guide correct assembly, we also searched for centromeres in the contigs. They

were identified as highly duplicated regions in the all-against-all alignment dot-plots

produced by LAST. The identification of the repeated canonical telomeric sequence

$(TTAGGG)_n$ [9] and its reverse complement $(CCCTAA)_n$ at the beginning or end of

certain contigs allowed the identification of chromosome ends. The Dan1 assembly

was manually curated using a similar strategy with the Dan2 genome as a reference.

<u>Solving links between contigs:</u>

Overlaps between linked contigs were identified by a BLASTN [38] alignment of their

last 100 kb. Any duplicate sequence was trimmed out from one contig and both contigs

were joined. The inferred junction was then validated by verification of the underlying

read support. For the linked contigs that did not overlap, the sequence in the gap was

extrapolated from the reads that matched and extended the ends of contigs, on the

basis of alignments at the last 1 kb of each contig. These sequences were aligned with

MAFFT v7.427 [39]. The alignment was visualized with SeaView [40], and only the

portion of the alignment strictly between the two contigs sequences was kept. Seaview

also generated a consensus sequence (on the basis of 60 % sequence identify by

default). The resulting sequence was inserted between the two contigs to link them

and the supposed continuity verified by a further cycle of read mapping.

<u>Assembly polishing with long-reads:</u>

Genome polishing was carried out with 2 or 4 iterative runs of Racon v1.4.2 [41] and

parameters -m 8 -x -6 -g -8 -w 500, and a run of Medaka v0.8.1 (Oxford Nanopore

Technologies) with the parameter -m r941_min_high.

<u>Mitochondrial genome circularization:</u>

Canu assembles small circular elements as contigs with tandem duplications of the

element. We resolved the mitochondrial genomes as recommended by Canu's authors

[7]. MUMmer suite v4.0.0.beta2 [42] was used to align the contig identified as the

putative mitochondria on itself with NUCmer and parameters --maxmatch --nosimplify.

Coordinates of a full copy were identified with the show-coords command and -lrcT

parameters.

<u>PCR:</u>

PCR was carried out to test a genome rearrangement between Swe2 and Dan2

genomes, with primers P1F (GAGATATCGAACGTCGCATGG), P1R

(ACATCAAGCCTTTGTCGAGGA), and P3F (GCTCAGGACCGACGTACAAG). PCR

reactions were run according to the GoTaq® G2 Flexi DNA polymerase instructions

(Promega), with 50 ng of template DNA, 1 mM of each forward and reverse primer, in

a final volume of 25 μL. The reaction started by initial denaturation at 95°C for 2 min,

followed by 30 amplification cycles (95°C for 30 sec, 60°C for 30 sec and 72°C for 30

sec), and a final elongation for 5 min at 72°C.

Defining a set of 305 identical proteins:

Identical proteins shared by the two *D.* coniospora genomes available (Swe2 and

Dan2) were recovered using a reciprocal best BLAST [38] hit strategy on the two

proteomes. Proteins that were duplicated in one or both genomes were filtered out.

The set was further refined by only retaining proteins corresponding to mono-exonic

genes.

Assessment of gene sequence in ONT-only assemblies:

TBLASTN searches were run using the amino-acid sequence of the set of 305 identical

proteins against the different nanopore only assemblies. A gene was considered as

correct if the query coverage, *i.e* the ratio of alignment length over the query length,

was equal to 1.

Short read polishing:

Shorts reads were trimmed using Trimmomatic v0.39 [43] with the parameters

LEADING:3 TRAILING:3 SLIDINGWINDOW:4:30 MINLEN:36. Then, only paired

reads were mapped on assemblies with bwa v0.7.17 [44] and default parameters (bwa

index, then bwa mem). The resulting mapping file was converted in BAM, sorted and

indexed with samtools. This latter file was used to polish the assembly with Pilon v1.23

[45] with the parameters --fix bases --vcf --mindepth 10 --minmq 20 --minqual 15 --

changes --diploid. Several iterations were conducted for each strain, until the number

of changes was less than 5.

Flye assembly:

An additional *de novo* assembly was performed with Flye v2.4.2 [8], and the parameter

--genome-size 32m, using the ONT reads recalled by Guppy v3.0.3.

Assessing the genome integrity:

The genome integrity was assessed with BUSCO v3.1.0 and the curated set

ascomycota_odb9 version 2016-02-13 [25]. A BLASTP search enabled Swe2

monoexonic genes present among USCOs to be identified. This list of 219 Swe2 genes

was then used as a TBLASTN query against the different assemblies of Swe1 and

Swe3. A gene was considered correct when it matched the corresponding Swe2 gene

perfectly in length. An analogous analysis was carried out for Dan1, on the basis of the

273 Dan2 monoexonic genes that are USCOs.

Characterisation of chimeric reads:

Swe1 reads identified as chimeric by YACRD were aligned on the final (short-read

polished) Swe1 assembly. The main alignment was identified using samtools view -F

2308. The CIGAR string was then parsed to determine whether the longest residual

part of the read was 5' or 3' to the main alignment, thereby giving an orientation to the

putative chimeric read and localising the potential chimeric break point. The 500 bp of

sequence 5' and 3' of this point were extracted and individually mapped back on the

Swe1 final assembly and the number of unique reads in a 10 kb non-overlapping sliding

window was calculated. For the reads for which both 500 bp fragments mapped on the

same chromosome, the smallest distance between the two fragments was calculated.

Data availability

Genomes of the strains Swe1, Swe3 and Dan1 are available on our institute website

(http://www.ciml.univ-mrs.fr/applications/DC/Genome.htm). All the reads used in this

work can be found at the European Nucleotide Archive (ENA) under the study numbers

PRJEB35969, PRJEB35970 and PRJEB35971. The raw signal runs are available

under the accessions ERR3774158, ERR3774162 and ERR3774163; the FASTQ files

of basecalled reads (Guppy v3.0.3) are available under the accessions ERR3997391,

ERR3997394 and ERR3997483; the FASTQ files of Illumina paired-end reads are

available under the accessions ERR3997389, ERR3997392, ERR3997395. Accession

numbers are given in the order Swe1, Swe3 and Dan1.

Funding

Supported by institutional grants from the Institut national de la santé et de la recherche

médicale, Centre National de la Recherche Scientifique and Aix-Marseille University

to the CIML, and the Agence Nationale de la Recherche program grant (ANR-16-

CE15-0001-01), and "Investissements d'Avenir" ANR-11-LABX-0054 (Labex

INFORM), ANR-16-CONV-0001 and ANR-11-IDEX-0001-02, and funding from the

Excellence Initiative of Aix-Marseille University - A*MIDEX.

Acknowledgments

The authors thank Yuquan Xu and Liwen Zhang for providing access to the raw

sequencing data for Dan2, Xing Zhang for help in preparing DNA samples, Lionel

Spinelli for informatic support and Nathalie Pujol and Laurent Tichit for comments.

Supplementary data

supplementary_figures.pdf contains 8 supplementary figures.

supplementary_table_1.xls contains the read coverage of the genomes. This table also

contains the results for the TBLASTN on the 305 candidate identical proteins.

supplementary_table_2.xls is a table recording read support, in Swe1, Swe3 and Dan1

assemblies, for the predicted correct sequence for each homopolymer stretch in the

genes corresponding to 10 protein of the 305 candidate identical proteins.

supplementary_methods.pdf contains additional methodological details.

References

1. Jansson HB, Jeyaprakash A, Zuckerman BM. Differential Adhesion and Infection of
Nematodes by the Endoparasitic Fungus *Meria coniospora* (Deuteromycetes). Appl Environ
Microbiol. 1985;49:552–5.

2. Pujol N, Link EM, Liu LX, Kurz CL, Alloing G, Tan MW, et al. A reverse genetic analysis
of components of the Toll signalling pathway in *Caenorhabditis elegans*. Curr Biol.
2001;11:809–21.

3. Dierking K, Polanowska J, Omi S, Engelmann I, Gut M, Lembo F, et al. Unusual regulation
of a STAT protein by an SLC6 family transporter in *C. elegans* epidermal innate immunity.
Cell Host Microbe. 2011;9:425–35.

4. Labed SA, Omi S, Gut M, Ewbank JJ, Pujol N. The pseudokinase NIPI-4 is a novel regulator
of antimicrobial peptide gene expression. PloS One. 2012;7:e33887.

5. Lebrigand K, He LD, Thakur N, Arguel M-J, Polanowska J, Henrissat B, et al. Comparative
Genomic Analysis of *Drechmeria coniospora* Reveals Core and Specific Genetic Requirements
for Fungal Endoparasitism of Nematodes. PLoS Genet. 2016;12:e1006017.

6. Zhang L, Zhou Z, Guo Q, Fokkens L, Miskei M, Pócsi I, et al. Insights into Adaptations to a
Near-Obligate Nematode Endoparasitic Lifestyle from the Finished Genome of *Drechmeria*
coniospora. Sci Rep. 2016;6:23122.

7. Canu 1.8 documentation. https://canu.readthedocs.io/en/latest/faq.html#my-circular-
element-is-duplicated-has-overlap. Accessed 15 Nov 2019

8. Kolmogorov M. Fast and accurate de novo assembler for single molecule sequencing reads:
fenderglass/Flye. https://github.com/fenderglass/Flye. Accessed 3 June 2019

9. Schechtman MG. Characterization of telomere DNA from *Neurospora crassa*. Gene.
1990;88:159–65.

10. Hazkani-Covo E, Zeller RM, Martin W. Molecular Poltergeists: Mitochondrial DNA
Copies (numts) in Sequenced Nuclear Genomes. PLoS Genet. 2010;6:e1000834.

11. Argueso JL, Westmoreland J, Mieczkowski PA, Gawel M, Petes TD, Resnick MA. Double-
strand breaks associated with repetitive DNA can reshape the genome. Proc Natl Acad Sci U S
A. 2008;105:11845–50.

12. Sun S, Yadav V, Billmyre RB, Cuomo CA, Nowrousian M, Wang L, et al. Fungal genome
and mating system transitions facilitated by chromosomal translocations involving
intercentromeric recombination. PLoS Biol. 2017;15:e2002527.

13. TransposonPSI: An Application of PSI-Blast to Mine (Retro-)Transposon ORF
Homologies. http://transposonpsi.sourceforge.net. Accessed 1 Apr 2020

14. Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, et al. The Dfam database of
repetitive DNA families. Nucleic Acids Res. Oxford Academic; 2016;44:D81–9.

15. Senol Cali D, Kim JS, Ghose S, Alkan C, Mutlu O. Nanopore sequencing technology and
tools for genome assembly: computational analysis of the current state, bottlenecks and future
directions. Brief Bioinform. 2018;20:1542–59.

16. Scheunert A, Dorfner M, Lingl T, Oberprieler C. Can we use it? On the utility of de novo
and reference-based assembly of Nanopore data for plant plastome sequencing. PloS One.
2020;15:e0226234.

17. White R, Pellefigues C, Ronchese F, Lamiable O, Eccles D. Investigation of chimeric reads
using the MinION. F1000Res. 2017;6:631.

18. Eccles D, Chandler J, Camberis M, Henrissat B, Koren S, Le Gros G, et al. De novo
assembly of the complex genome of *Nippostrongylus brasiliensis* using MinION long reads.
BMC Biol. 2018;16:6.

19. Schmid M, Frei D, Patrignani A, Schlapbach R, Frey JE, Remus-Emsermann MNP, et al.
Pushing the limits of de novo genome assembly for complex prokaryotic genomes harboring
very long, near identical repeats. Nucleic Acids Res. 2018;46:8953–65.

20. Watson M, Warr A. Errors in long-read assemblies can critically affect protein prediction.
Nature Biotechnology. 2019;37:124–6.

21. Wick RR, Judd LM, Holt KE. Performance of neural network basecalling tools for Oxford
Nanopore sequencing. Genome Biol. 2019;20:129.

22. Somerville V, Lutz S, Schmid M, Frei D, Moser A, Irmler S, et al. Long-read based de novo
assembly of low-complexity metagenome samples results in finished genomes and reveals
insights into strain diversity and an active phage system. BMC Microbiol. 2019;19:143.

23. Dal Molin A, Minio A, Griggio F, Delledonne M, Infantino A, Aragona M. The genome
assembly of the fungal pathogen *Pyrenochaeta lycopersici* from Single-Molecule Real-Time
sequencing sheds new light on its biological complexity. PLoS One. 2018;13:e0200217

24. Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, et al. Nanopore sequencing and
assembly of a human genome with ultra-long reads. Nat Biotechnol. 2018;36:338–45.

25. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31:3210–2.

26. He LD, Ewbank JJ. Polyethylene Glycol-mediated Transformation of *Drechmeria*
coniospora. Bio-Protoc. 2017;7:e2157.

27.   Fungal   DNA   extraction   protocol.
https://www.pnas.org/content/pnas/suppl/2018/01/08/1715954115.DCSupplemental/pnas.171
5954115.sapp.pdf. Accessed 16 Apr 2020.

28. Kjærbølling I, Vesth TC, Frisvad JC, Nybo JL, Theobald S, Kuo A, et al. Linking secondary
metabolites to gene clusters through genome sequencing of six diverse *Aspergillus* species.
Proc Natl Acad Sci. 2018;115:E753–61.

29. Quick J. Ultra-long read sequencing protocol for RAD004 v3.
https://www.protocols.io/view/ultra-long-read-sequencing-protocol-for-rad004-mrxc57n.
Accessed 25 Oct 2019

30. Wick R. Porechop. https://github.com/rrwick/Porechop. Accessed 3 Dec 2019

31. Marijon P, Chikhi R, Varré J-S. yacrd and fpa: upstream tools for long-read genome
assembly. bioRxiv. 2019;674036.

32. Kiełbasa SM, Wan R, Sato K, Horton P, Frith MC. Adaptive seeds tame genomic sequence
comparison. Genome Res. 2011;21:487–93.

33. Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, et al. Circos: An
information aesthetic for comparative genomics. Genome Res. 2009;19:1639–45.

34. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34:3094–100.

35. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinforma Oxf Engl. 2009;25:2078–9.

36. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. Brief Bioinform. 2013;14:178–92.

37. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res.
2017;27:722–36.

38. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool.
J Mol Biol. 1990;215:403–10.

39. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30:772–80.

40. Gouy M, Guindon S, Gascuel O. SeaView Version 4: A Multiplatform Graphical User
Interface for Sequence Alignment and Phylogenetic Tree Building. Mol Biol Evol.
2010;27:221–4.

41. Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from
long uncorrected reads. Genome Res. 2017;27:737–46.

42. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile
and open software for comparing large genomes. Genome Biol. 2004;5:R12.

43. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. Oxford Academic; 2014;30:2114–20.

44. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
ArXiv 2013;1303.3997

45. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An
Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly
Improvement. PLoS One. 2014;9:e112963.

604
605

Figure legends

Figure 1. An overview of *D. coniospora* strain isolation and culture history.

A strain of *D. coniospora* collected from Denmark in 1982 at the latest was deposited at the CBS-KNAW culture collection, now held by the Westerdijk Fungal Biodiversity Institute as CBS615.82. It was transferred in 1987 to the ARS Collection of Entomopathogenic Fungal Cultures (as ARSEF 2468) and then re-isolated in 2001 as ARSEF 6962. A second strain collected from Sweden was deposited at the American Type Culture Collection as ATCC 96282. It has been cultured through serial passage in *C. elegans* continuously since 1999.

Figure 2. Inter-chromosomal rearrangements between strains Swe2 and Dan2.

A. Circos plot representing regions >6 kb that are very similar between Dan2 (left – olive) and Swe2 (right – light blue) assemblies as determined by an all-against-all LAST analysis. Swe2 contig numbers are the last two digits of the accession ID (shown in B), preceding the suffix. Red and dark blue rectangles represent rearrangement junctions probed by PCR. B. Conceptual design of the PCR primers. C. Amplicons from the PCR were visualized after electrophoresis. Each pair gave one specific band of the expected size. The colour code is the same for the 3 panels.

Figure 3. Comparisons between Canu and Flye assemblies.

A, B. Dot plots of the non-congruent assemblies generated by Canu (x-axis) against those generated by Flye (y-axis) for the Swe3 (A) and Dan1 (B) genomes. The orange triangle (A) marks the position where the Canu contig *tig00000004* was split during the manual curation because of its chimeric nature. The green arrow (B) marks the position

of a Flye scaffolding error. C. Schematic representation of the Dan1 Flye assembly, showing the mapping of chimeric reads close to the scaffolding error (green triangle). The coordinates in brackets are the mapping positions of the clipped part of the reads (dash line) on another contig of the assembly. Notably, this error was eliminated when these chimeric reads were excluded from the input data. D. Mapping of long-reads close to the scaffolding error (green triangle) on the Dan1 Flye assembly. The green bar marks the telomeric tandem repeat motif. The grey bar indicates the 100 Ns inserted by Flye to unite the scaffold.

Figure 4. Synteny among the genomes of 5 *D. coniospora* strains.

Circos plot representing regions >20 kb that are very similar between assemblies as determined by all-against-all LAST analyses. Each assembly is shown at the same scale and in the same order and orientation across panels.

Figure 5. Evaluation of sequence errors in the 3 new genomes.

A. Percentage of correct genes (based on length of the corresponding predicted protein) among 305 conserved genes, for the 3 new genomes, in the initial assembly and after two different polishing strategies. B, C, D. Scatter plots of homopolymer composition (A/T or C/G) and accuracy among the same 305 conserved genes for Dan1 (B), Swe1 (C) and Swe3 (D). The dot size is proportional to the number of genes, and the colour indicates the proportion of genes predicted to be correct. Red and purple arrows highlight two particular cases, among many, where homopolymer errors are only present in one genome.

Figure 6. Sequence anomalies introduced by assembly and/or polishing tools.

A. A comparison of one small region of the Swe3 sequence before (top) and after polishing (Racon x4 and Medaka; bottom). As indicated by the orange line, long stretches of A and T homopolymers are introduced by polishing, in the absence of coherent read support. B. From top to bottom, the assembly produced by Canu excludes a region of around 10 kb, despite strong read support. After 2 and 4 iterations, Racon progressively filled the gap. Medaka then introduced an insert of roughly the correct size, but of aberrant sequence composition. For each panel, the height of the boxes in the top line indicates the read coverage for each base. A grey box indicates full agreement with the consensus sequence, otherwise the colour indicates the proportion of read support for each nucleotide (G, tan; C, blue; A, green; T, red). Below this, the ONT reads that align in forward (pink) and reverse (blue) orientation are shown as lines. A coloured letter or purple rectangle show a difference (nucleotide variant or insertion in reads, respectively) in the read's sequence compared to the genome sequence. C. The 10 kb sequence introduced by polishing is of aberrant composition as illustrated by the region immediately surrounding the 5' breakpoint (yellow arrowhead). There are single nucleotide errors introduced despite coherent read support for the "Before" sequence (light blue dots), and then a continuous stretch, exemplified by A and T homopolymers that lack any sequence support at all (light blue line).

Figure 7. Example of sequence errors introduced during assembly and polishing.

A. Stacked bar plot of USCO status for the orthologues of selected mono-exonic Swe2 or Dan2 genes classified according to the result of a TBLASTN search against the

indicated assembly (Canu: from Canu; Racon: after long-read polish; Pilon: after short-read polishing). B. Detailed view of 2 parts of RJ55_06485 from the Swe2 reference genome each containing a homopolymer sequence (underlined) and the corresponding positions in successive Swe1 assemblies. For each, the predicted protein sequence, highlighted in turquoise, with the other open reading frames in grey, is shown above the corresponding nucleotide sequence. The red arrow heads highlight the missing nucleotides, the extraneous nucleotide is boxed in red.

Figure 1

Figure 2

A

Dan2

CM004174.1

04
02
07
01

Swe2

CM004175.1

CM004176.1

11 10
03
06
08 09 05

[gradient bar] = 1 Mb

B

Swe2 JYHR01000001.1 → P1R
                        P1F ←

Dan2 CM004174.1 → P1R
                   P3F ←

C

P1F + P1R   P3F + P1R
Swe2 Dan2   Swe2 Dan2

2 kb
1 kb
0.5 kb

Figure 3

A Swe3 Canu assembly

Swe3 Flye assembly

tig00000003 tig00000004 tig00000023 tig00000034

contig_1

contig_2

B Dan1 Canu assembly

Dan1 Flye assembly

tig00000004 tig00000007

scaffold_3

C Dan1 scaffold_3
(Flye assembly)

Chimeric reads {

D

Telomeric repeat (TTAGGG)$_n$ Scaffold insert

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

Supplementary Figure 1. Distribution of the size of the reads and the assembly statistics.

Distributions in 5 kb bins of the size of the set of reads basecalled by Guppy v1.5.1(left-hand panels), and of reads corrected and trimmed by Canu for the initial assemblies (righthand panels).

Supplementary Figure 2

Supplementary Figure 2. Detection of an error in the initial Swe1 assembly introduced by a long chimeric read.

A. Circos plot representing regions >20 kb that are very similar between Swe3 and Swe1 Canu assemblies as determined by an all-against-all LAST analysis. The red arrow indicates a break in the synteny for the largest Swe1 contig. B. Dot-plot of an all-against-all comparison of the Swe1 contigs and unitigs produced by Canu. Contigs are contiguous sequences present in the primary assembly, including both unique and repetitive elements. Unitigs are contigs split at alternate paths in the assembly graph. The red arrow indicates the discontinuity in the alignment between Swe1 contigs and unitigs. This occurs on the same contigs and at the same coordinates as in A. C. Mapping of the Swe1 reads, corrected and trimmed by Canu, on the Swe1 Canu assembly (detail of around 70 kb on contig tig00002308 flanking the synteny break). Moving into the central 5 kb region, read support progressively drops from 40 to just one, corresponding to a very long read of about 215 kb, which was shown to be chimeric.

Supplementary Figure 3

A

B

C

Supplementary Figure 3. Detection of an error in the Swe3 initial assembly introduced by a long chimeric read.

A. Mapping of the Swe3 reads, corrected and trimmed by Canu, on the Swe3 Canu assembly. The putative 175 kb chimeric read was identified by a break of synteny (not shown) and because of a sharp (ca. 2-fold) increase in coverage, spanning only 2 nucleotides, (purple arrow) on the contig tig0000004. B. Conceptual design of the PCR primers used to verify the assembly. Three pairs of primers were designed on the tig0000004: the Up pair (red), the Down pair (green) and the Span pair (blue). Two other primers were designed on the basis of the corrected assembly sequence (dotted coloured lines): SpanF in turquoise on the contig tig00000027 and SpanR in light purple on the contig tig00000695. C. PCR result of the different pairs used. Amplicons had the expected sizes.

Supplementary Figure 4

Supplementary Figure 4. Canu correction of non-chimeric and chimeric reads.

A. Scatter plots of the size of reads in the initial dataset (y-axis) and after Canu correction (x-axis) for non-chimeric (top) and chimeric (bottom) reads. Subsets of non-chimeric reads of a similar number (as indicated) and having the equivalent size distribution (before Canu correction, +/- 1% in length) as the chimeric read were used to allow a valid comparison. B. Percentage of read length reduction after Canu correction, in 5 kb bins. The bin size corresponds to the read size in the raw dataset. The red arrows highlight the rare chimeric reads that are longer after the correction step.

Supplementary Figure 5

Supplementary Figure 5. Size distribution of chimeric reads.

Distributions in 5 kb bins of the size of the set of reads identified as chimeric by YACRD (see Methods), in the original dataset (left panels) and amongst reads basecalled by Guppy v3.0.3 (right panels).

Supplementary Figure 6

Supplementary Figure 6. Analysis of the Swe1 chimeric reads.

For each chimeric read, the 500 bp of sequence on each side of the putative break point was mapped onto the final Swe1 genome. A, Distribution in 500 equal bins of the interval between the two parts of each chimeric read. The inserts highlight reads that map within 1 kb of each other (in 200 bins). B, Cumulative distribution of the interval between the two parts of each chimeric read (for those separated by > 1 Mb). The distance separating the two parts is broadly spread. C. Distribution of mapped sequences from each side of the putative breakpoint for each chimeric read that mapped to a single chromosome. The peak on chromosome 2 corresponds to the site where the nuclear genome matches the mitochondrial DNA, and that on chromosome 3 to a highly repeated region containing tandem copies of rDNA. In both cases, these therefore reflect erroneous attribution of chimerism. There are thus no true hot spots for chimeric read breakpoints on any chromosome.

Supplementary Figure 7

Supplementary Figure 7. Identification of intra- and inter-chromosomal rearrangements between Dan1 and Dan2.

A. Mapping of long reads from Dan1 (left panels) and Dan2 (right panels) on Dan1 chromosome 1 (top panels) and Dan2 chromosome 1 (CM004174.1; bottom panels). The arrowheads highlight points of discontinuity in the read coverage, consistent with chromosomal rearrangements between Dan1 and Dan2. B. Alignment of selected chromosomes of the final assemblies between Dan1 and Swe1 (left) and Dan2 and Swe1 (right). The unique difference in the orientation of part of Swe1 chromosome 2 between Dan1 and Dan2 is highlighted by the green circles.

Supplementary Figure 8

Supplementary Figure 8. Analysis of the pattern of repeated sequences in the neighbourhood of polishing error in the Swe3 genome.

A. Dot-plot based on k-mer identity (k = 11; See Supplementary Methods) of the neighbourhood of the *numts vs* itself, from the left to the right, before any polishing, after long-read polishing and after short-read polishing. The vertical dashed line at the centre of each plot indicates the start of the *numts*. B. A schematic representation of the large-scale genomic organization in the same region. The rectangles (red and dark red) represent *numts* (identified by BLASTN against the Swe3 mitochondrial genome) and the ribbons show the regions of sequence similarity (>99.5%) between them. For the Swe1 assembly, one dark-red rectangle is absent, while in the Swe2 assembly, the two dark-red rectangles are absent. C. An alignment of the area before any polishing (x-axis) and after short-read polishing (y-axis) reveals the re-inclusion of this missing sequence in the final assembly. The arrow highlights a 1.2 kb region that is duplicated on both sides of the sequence discontinuity (vertical dashed line). D. Dot-plots based on k-mers (k = 11) of the same region. The blue square reflects the presence of a region of low-complexity repeated sequences. The pattern at the top left hand of the plots reflect a region of low-complexity sequence that is not shared across the sequence discontinuity (vertical dashed line).

Supplementary table 1: Coverage, genome length and results of the tBLASTn

	Set of reads	Swe1	Swe3	Dan1
	Long read - Guppy 1.5.1	136	94	150
Coverage (fold)	Long read - Canu corrected	36	36	36
	Long read - Guppy 3.0.3	140	97	155
	Short read	406	430	482
	Canu after curation	31971578	32076624	31919747
Genome length (bp)	Long-read polishing	32121045	32271010	32120031
	Short-read polishing	32117370	32271155	32095731
	Canu after curation	43 (14%)	48 (16%)	42 (14%)
TBLASTN result for the 305 benchmarking genes	Long-read polishing	236 (77%)	248 (81%)	152 (50%)
	Short-read polishing	305 (100%)	305 (100%)	305 (100%)

The coverage for each set of reads was computed as the mean depth for the assemblies after short-read polishing, using Samtools coverage v1.10.

Supplementary Table 2: Examples of read support for different homopolymer sequences in 10 representative genes.

gene	reference	Swe2	Swe1	Swe1 read	Swe1 guppy remark	Swe3	Swe3 read	Swe3 guppy remark	Dan2	Dan1	Dan1 read	Dan1_guppy_remark
ODA80475.1	c	5	5	36 (2)	supported	5	34 (0)	supported	5	6	28 (0)	weak support (<10% read)
ODA80475.1	t	4	4	35 (0)	supported	4	34 (3)	supported	4	4	28 (12)	supported
ODA80475.1	t	NA	NA	NA (NA)	NA	NA	NA (NA)	NA	4	4	28 (3)	supported
ODA80475.1	c	4	4	37 (23)	supported	4	35 (33)	supported	4	5	27 (2)	supported
ODA80475.1	g	7	8	36 (0)	weak support (<10% read)	6	35 (2)	supported	7	7	28 (0)	supported
ODA80133.1	c	4	4	33 (33)	supported	4	41 (41)	supported	4	4	37 (36)	supported
ODA80133.1	c	4	4	33 (26)	supported	4	41 (38)	supported	4	4	36 (18)	supported
ODA80133.1	g	4	4	33 (32)	supported	4	42 (40)	supported	4	4	35 (35)	supported
ODA80133.1	c	7	6	32 (0)	weak support (<10% read)	6	41 (0)	supported	7	7	37 (0)	supported
ODA83792.1	c	4	4	33 (33)	supported	4	38 (38)	supported	4	4	28 (28)	supported
ODA83792.1	c	4	4	33 (30)	supported	4	38 (36)	supported	4	4	28 (24)	supported
ODA83792.1	c	4	4	33 (32)	supported	4	38 (34)	supported	4	4	28 (28)	supported
ODA83792.1	c	5	6	33 (0)	weak support (<10% read)	5	38 (17)	supported	5	6	28 (0)	weak support (<10% read)
ODA83792.1	c	4	4	33 (33)	supported	4	38 (36)	supported	4	4	28 (27)	supported
ODA83792.1	c	5	5	32 (0)	supported	5	38 (36)	supported	5	5	28 (1)	supported
ODA83792.1	c	5	5	32 (12)	supported	6	38 (0)	weak support (<10% read)	5	5	28 (11)	supported
ODA83792.1	c	4	4	32 (32)	supported	4	38 (37)	supported	4	4	28 (28)	supported
ODA83792.1	c	4	4	32 (31)	supported	4	38 (36)	supported	4	4	28 (28)	supported
ODA83792.1	a	4	4	33 (28)	supported	4	37 (35)	supported	4	4	28 (27)	supported
ODA83792.1	a	4	4	33 (28)	supported	4	37 (37)	supported	4	4	28 (21)	supported
ODA83792.1	g	4	4	32 (32)	supported	4	38 (37)	supported	4	4	28 (28)	supported
ODA83792.1	t	4	4	32 (32)	supported	4	38 (38)	supported	4	4	28 (28)	supported
ODA83726.1	c	4	5	25 (0)	weak support (<10% read)	4	37 (12)	supported	4	5	37 (0)	supported
ODA83726.1	c	4	4	25 (23)	supported	4	37 (35)	supported	4	4	37 (36)	supported
ODA83726.1	c	5	5	25 (17)	supported	5	37 (27)	supported	5	5	37 (14)	supported
ODA83726.1	c	5	6	25 (0)	weak support (<10% read)	5	37 (12)	supported	5	6	37 (0)	supported
ODA83726.1	c	7	7	25 (1)	weak support (<10% read)	7	37 (0)	weak support (<10% read)	7	7	37 (2)	supported
ODA83726.1	c	5	5	25 (24)	supported	5	36 (28)	supported	5	5	37 (26)	supported
ODA83726.1	c	5	5	25 (6)	supported	5	36 (13)	supported	5	5	37 (14)	supported
ODA83726.1	g	4	4	25 (24)	supported	4	36 (29)	supported	4	4	37 (20)	supported
ODA84322.1	c	5	5	38 (1)	supported	5	41 (0)	supported	5	5	41 (11)	supported
ODA84322.1	g	4	4	38 (29)	supported	4	41 (29)	supported	4	4	41 (21)	supported
ODA84322.1	g	4	4	38 (13)	supported	4	40 (7)	supported	4	5	41 (1)	supported
ODA84322.1	c	5	5	38 (6)	supported	5	42 (3)	supported	5	5	41 (3)	supported
ODA84322.1	c	5	5	38 (6)	supported	5	42 (8)	supported	5	5	41 (1)	supported
ODA84322.1	c	5	6	38 (0)	weak support (<10% read)	6	42 (3)	weak support (<10% read)	5	5	41 (7)	supported
ODA84322.1	g	4	4	37 (37)	supported	4	42 (42)	supported	4	5	41 (1)	supported
ODA82449.1	c	5	5	39 (4)	supported	5	46 (2)	supported	5	5	26 (8)	supported
ODA82449.1	c	5	6	39 (1)	weak support (<10% read)	6	46 (1)	weak support (<10% read)	5	8	26 (0)	weak support (<10% read)
ODA82449.1	c	5	5	39 (16)	supported	6	46 (2)	weak support (<10% read)	5	5	26 (8)	supported
ODA82449.1	c	5	6	39 (0)	weak support (<10% read)	6	46 (0)	weak support (<10% read)	5	6	26 (0)	weak support (<10% read)
ODA82449.1	a	5	5	38 (21)	supported	5	47 (34)	supported	5	5	26 (6)	supported
ODA82449.1	c	6	7	38 (0)	weak support (<10% read)	7	47 (2)	weak support (<10% read)	6	6	26 (2)	supported
ODA81732.1	c	5	5	38 (25)	supported	5	47 (33)	supported	5	5	26 (15)	supported
ODA81732.1	g	4	5	19 (0)	supported	5	40 (0)	supported	4	5	39 (0)	supported
ODA81732.1	c	7	7	20 (1)	weak support (<10% read)	7	40 (2)	weak support (<10% read)	7	8	40 (1)	weak support (<10% read)
ODA76966.1	c	5	5	40 (1)	supported	5	31 (1)	supported	5	6	39 (2)	weak support (<10% read)
ODA76966.1	g	4	4	40 (40)	supported	4	31 (30)	supported	4	4	39 (39)	supported
ODA76966.1	c	4	4	40 (32)	supported	4	31 (30)	supported	4	4	39 (29)	supported

#gene	#reference	#Swe2	#Swe1	#Swe1 read	#Swe1 guppy remark	#Swe3	#Swe3 read	#Swe3 guppy remark	#Dan2	#Dan1	#Dan1 read	#Dan1 guppy remark
ODA76966.1	g	4	4	39 (37)	supported	4	31 (30)	supported	4	4	39 (37)	supported
ODA76966.1	c	4	4	39 (31)	supported	4	31 (28)	supported	4	4	39 (29)	supported
ODA76966.1	g	8	7	38 (0)	weak support (<10% read)	7	31 (1)	weak support (<10% read)	8	7	39 (0)	weak support (<10% read)
ODA76653.1	g	4	4	39 (37)	supported	4	38 (36)	supported	4	4	39 (39)	supported
ODA76653.1	c	4	4	39 (39)	supported	4	38 (38)	supported	4	4	40 (39)	supported
ODA76653.1	a	4	4	39 (39)	supported	4	38 (37)	supported	4	4	40 (40)	supported
ODA76653.1	g	4	4	39 (37)	supported	4	38 (38)	supported	4	4	40 (40)	supported
ODA76653.1	a	4	4	24 (22)	supported	4	38 (8)	supported	4	4	40 (38)	supported
ODA78969.1	g	4	4	23 (23)	supported	4	29 (28)	supported	4	4	42 (38)	supported
ODA78969.1	c	4	4	23 (17)	supported	4	29 (29)	supported	4	4	42 (42)	supported
ODA78969.1	c	4	4	23 (22)	supported	4	29 (20)	supported	4	4	42 (19)	supported
ODA78969.1	c	4	4	NA (NA)	NA	4	29 (26)	supported	5	6	42 (2)	supported
ODA78969.1	c	NA	NA	23 (23)	supported	NA	NA (NA)	NA	6	6	42 (2)	supported
ODA78969.1	c	4	4	23 (0)	supported	4	29 (29)	supported	4	4	42 (41)	supported
ODA78969.1	c	7	6	23 (22)	supported	6	29 (0)	supported	6	7	42 (3)	supported
ODA78969.1	c	4	4	23 (22)	supported	4	29 (29)	supported	4	4	42 (36)	supported

Header descriptions

#gene	Gene involved, refers to Swe2 proteinGenebank ID
#reference	nucleotide of the homopolymer
#Swe2	Homopolymer length in Swe2 (reference for Swe1 and Swe3)
#Swe1	Homopolymer length in Swe1
#Swe1 read	Total number of reads (number of reads that support the homopolymer length in Swe1) - Canu correct and trimmed reads
#Swe3 guppy remark	Support of this homopolymer in the full set of reads that was used to polish the genome - Guppy 3.0.3
#Swe3	Homopolymer length in Swe3
#Swe3 read	Total number of reads (number of reads that support the homopolymer length in Swe3) - Canu correct and trimmed reads
#Swe3 guppy remark	Support of this homopolymer in the full set of reads that was used to polish the genome - Guppy 3.0.3
#Dan2	Homopolymer length in Dan2 (Reference for Dan1)
#Dan1	Homopolymer length in Dan1
#Dan1 read	Total number of reads (number of reads that support the homopolymer length in Dan1) - Canu correct and trimmed reads
#Dan1 guppy remark	Support of this homopolymer in the full set of reads that was used to polish the genome - Guppy 3.0.3

Counts were made manually, with an estimated accuracy of +/- 10%.

Supplementary methods

<u>GenBank query:</u>

We queried the Assembly database on the NCBI website with:

("minion"[Sequencing Technology] OR "nanopore"[Sequencing Technology] OR "nanopore minion"[Sequencing Technology] OR "nanopore technologies"[Sequencing Technology] OR "nanopore technology"[Sequencing Technology] OR "nanpore"[Sequencing Technology] OR "ont"[Sequencing Technology] OR "ont minion"[Sequencing Technology] OR "oxford"[Sequencing Technology] OR "oxford nanopore"[Sequencing Technology] OR "oxford nanopore minion"[Sequencing Technology] OR "oxford nanopore technologies"[Sequencing Technology] OR "oxford nanopore technology"[Sequencing Technology]) NOT ("illumina" OR "bgi" OR "pacbio" OR "pacbio rs" OR "pacbio rs ii" OR "pacbio rsii" OR "pacbio sequel" OR "pacbio smrt" OR "pacific biosciences" OR "sequel" OR "hybrid" OR "hybrid assembly") AND (latest[filter] OR "latest genbank"[filter]) AND (all[filter] NOT "derived from surveillance project"[filter]) AND (all[filter] NOT anomalous[filter])

The accuracy of this query depends on the assembly metadata in GenBank.

<u>K-mer based dot-plots:</u>

The dot-plots based on k-mers were computed with the script *repaver.r* (available at https://gitlab.com/gringer/bioinfscripts) The version used here was the commit 0854f3af76cb5bc5a7e0a2b4a032518f51e210fa, with the parameters *-k 11 -style dotplot*.

<u>Search for Dan1 / Dan2 transposable element at the breakpoints:</u>

A first analysis was conducted with the Dfam (https://dfam.org/home) to search for transposable elements (TE). The parameter *organism* was set to *other*. No TE was present in the neighbourhood of the different breakpoints. We also mined the two genomes for TE with transposonPSI (http://transposonpsi.sourceforge.net/) with default parameters. Here also, no TE was present within the 50kb surrounding each breakpoint.

PCR:

PCR was performed to test the putative chimeric nature of a Canu contig in the Swe3 assembly, with primers UpF (AACTGTGTCTAACTAGCCCG), UpR (AGGGTCCTCATAAACTTGGC), DownF (TGTATCAGGTTCCCGAATGG), DownR (CTAGGCTGGGGAATCTTCTG), SpanF (CCATCAACTTCAGCTGCTC), SpanR (CTCCTCAATCTCCCTCTCGG), ConfF (ATCGGCGACTACCTGCAC), ConfR (CGTTCCATCGTTACCACAGC). PCR reactions were run according to the GoTaq® G2 Flexi DNA polymerase instructions (Promega), with 50 ng of template DNA, 1 mM of each forward and reverse primers, in a final volume of 25 μL. The reaction started by initial denaturation at 95°C for 2 min, followed by 30 amplification cycles (95°C for 30 sec, 60°C for 30 sec and 72°C for 30 sec), and a final elongation for 5 min at 72°C.