Neural speech restoration at the cocktail party: Auditory cortex recovers masked speech of both attended and ignored speakers

Christian Brodbeck[1,*], Alex Jiao[2], L. Elliot Hong[3] & Jonathan Z. Simon[1,2,4]

1) Institute for Systems Research, University of Maryland, College Park, Maryland 20742, U.S.A
2) Department of Electrical and Computer Engineering, University of Maryland, College Park, Maryland 20742, U.S.A
3) Department of Psychiatry, Maryland Psychiatric Research Center, University of Maryland School of Medicine, Baltimore, Maryland 21201, U.S.A
4) Department of Biology, University of Maryland, College Park, Maryland 20742, U.S.A

* christianbrodbeck@me.com

Short title: Neural speech restoration at the cocktail party

Abstract

Humans are remarkably skilled at listening to one speaker out of an acoustic mixture of several speech sources. Two speakers are easily segregated, even without binaural cues, but the neural mechanisms underlying this ability are not well understood. One possibility is that early cortical processing performs a spectro-temporal decomposition of the acoustic mixture, allowing the attended speech to be reconstructed via optimally weighted recombinations that discount spectro-temporal regions where sources heavily overlap. Using human magnetoencephalography (MEG) responses to a two-talker mixture, we show evidence for an alternative possibility, in which early, active segregation occurs even for strongly spectro-temporally overlapping regions. Early (~70 ms) responses to non-overlapping spectro-temporal features are seen for both talkers. When competing talkers' spectro-temporal features mask each other, the individual representations persist, but they occur with a ~20 ms delay. This suggests that the auditory cortex recovers acoustic features that are masked in the mixture, even if they occurred in the ignored speech. The existence of such noise-robust cortical representations, of features present in attended as well as ignored speech, suggests an active cortical stream segregation process, which could explain a range of behavioral effects of ignored background speech.

Keywords

Auditory streaming, stream segregation, continuous speech, temporal response function

Introduction

When listening to an acoustic scene, the signal that arrives at the ears is an additive mixture of the different sound sources. Listeners trying to selectively attend to one of the sources face the task of determining which spectro-temporal features belong to that source [1]. When multiple speech sources are involved, as in the classic cocktail party problem [2], this is a nontrivial problem because the spectrograms of the different sources often have strong overlap. Nevertheless, human listeners are remarkably skilled at focusing on one out of multiple talkers [3,4]. Binaural cues can facilitate segregation of different sound sources based on their location [5], but are not necessary for this ability, since listeners are able to selectively attend even when two speech signals are mixed into a monophonic signal and presented with headphones [6]. Here we are specifically interested in the fundamental ability to segregate and attend to one out of multiple speakers even without such external cues.

The neural mechanisms involved in this ability are not well understood, but previous research suggests at least two separable cortical processing stages. In magnetoencephalographic (MEG) responses to multiple talkers [7], the early (~50 ms) cortical component is better described as a response to the acoustic mixture than as the sum of the responses to the individual (segregated) source signals, consistent with an early unsegregated representation of the mixture. In contrast, the later (> 85 ms) response component is dominated by the attended (segregated) source signal. Recent direct cortical recordings largely confirm this picture, suggesting that early responses in Heschl's Gyrus (HG) reflect a spectro-temporal decomposition of the acoustic mixture that is largely unaffected by selective attention, whereas later responses in the superior temporal gyrus (STG) dynamically change to represent the attended speaker [8]. In general, cortical regions further away from core auditory cortex tend to mainly reflect information about the attended speaker [9]. Together, these results suggest a cortical mechanism which, based on a detailed representation of the acoustic input, detects and groups features belonging to the attended source.

A long-standing question is whether early cortical processing of the acoustic mixture is restricted to passive spectro-temporal filtering, or whether it involves active grouping of acoustic features leading to the formation of auditory object representations. The filter theory of attention suggests early representations of physical stimulus characteristics independent of attention, with attention selecting a subset of these for further processing and semantic identification [10,11]. Consistent with this, electroencephalography (EEG) and MEG results suggest that time-locked processing of higher order linguistic features, such as words and meaning, is restricted to the attended speech source [12,13]. However, it is not known whether, in the course of recovering the attended source, the auditory cortex also segregates acoustic features of the ignored source from the mixture. Individual intracranially recorded HG responses to a two-speaker mixture can be selective for either one of the speakers, but this selectivity can be explained merely by spectral response characteristics favoring the spectrum of a given speaker over the other [8]. A conservative hypothesis is thus that early auditory cortical responses represent acoustic features of the mixture based on stable (possibly pre-defined) spectro-temporal receptive fields, allowing the attended speech to be segregated through an optimally weighted combination of these responses. Alternatively, the auditory cortex could employ more active mechanisms to dynamically recover potential speech features, regardless of what stream they belong to.

Selective attention could then rely on these local auditory (proto-) objects to recover the
attended speech [14]. This hypothesis critically predicts the existence of representations of
acoustic features from an ignored speech source, even when those features are not apparent in
the acoustic mixture, i.e., when those features are masked by acoustic energy from another
source. Here we report evidence for such representations in human MEG responses.

Cortical responses to speech reflect a strong representation of the envelope (or spectrogram) of
the speech signal [15,16]. Prior work has also shown that acoustic onsets are prominently
represented in auditory cortex, both in naturalistic speech [17,18] and in non-speech stimuli
[19,20]. Previous studies using paradigms similar to the one used here often predicted brain
responses from only envelopes or only onsets [16,21–24], but more recent studies show that
both representations explain non-redundant portion of the responses [12,18]. Behaviorally,
acoustic onsets are also specifically important for speech intelligibility [25,26]. Here we consider
both envelope and onset features, but focus on onset features in particular because of their
relevance for stream segregation, as follows [1]. If acoustic elements in different frequency
regions are co-modulated over time, they likely stem from the same physical source [27]. A
simultaneous onset in distinct frequency bands thus provides sensory evidence that these cross-
frequency features originate from the same acoustic source, and should be processed as an
auditory object. Accordingly, shared acoustic onsets promote perceptual grouping of acoustic
elements into a single auditory object, such as a complex tone and, vice versa, separate onsets
lead to perceptual segregation [28,29]. For example, the onset of a vowel is characterized by a
shared onset at the fundamental frequency of the voice and its harmonics. Correspondingly, if
the onset of a formant is artificially offset by as little as 80 ms, it can be perceived as a separate
tone rather than as a component of the vowel [30]. This link to object perception thus makes
acoustic onsets particularly relevant cues, which might be represented distinctly from envelope
cues and used to detect the beginning of local auditory objects, and thus to aid segregation of
the acoustic input into different, potentially overlapping auditory objects.

We analyzed human MEG responses to a continuous two-talker mixture to determine to what
extent the auditory cortex reliably tracks acoustic onset or envelope features of the ignored
speech, above and beyond the attended speech and the mixture. Participants listened to 1-
minute long continuous audiobook segments, spoken by a male or a female speaker. Segments
were presented in two conditions: a single talker in quiet ("clean speech"), and a two-talker
mixture, in which a female and a male speaker were mixed at equal perceptual loudness. MEG
responses were analyzed as additive, linear response to multiple concurrent stimulus features
(see Figure 1). First, cross-validated model comparisons were used to determine which
representations significantly improved prediction of the responses. Then, the resultant spectro-
temporal response functions (STRFs) were analyzed to gain insight into the nature of the
representations.

Figure 1. **Additive linear response model based on spectro-temporal response functions (STRFs).**
A) MEG responses recorded during stimulus presentation were source localized with distributed
minimum norm current estimates. A single virtual source dipole is shown for illustration, with its
physiologically measured response and the response prediction of a model. Model quality was
assessed by the correlation between the measured and the predicted response. B) The model's
predicted response is the sum of tonotopically separate response contributions generated by
convolving the stimulus envelope at each frequency (C) with the estimated temporal response
function (TRF) of the corresponding frequency (D). TRFs quantify the influence of a predictor
variable on the response at different time lags. The stimulus envelopes at different frequencies
can be considered a collection of parallel predictor variables, as shown here by the gammatone
spectrogram (8 spectral bins); the corresponding TRFs as a group constitute the spectro-
temporal response function (STRF). Physiologically, the component responses (B) can be thought
of as corresponding to responses in neural subpopulations with different frequency tuning, with
MEG recording the sum of those currents.

Results and Discussion

Auditory cortex represents acoustic onsets

MEG responses to clean speech were predicted from the gammatone spectrogram of the
stimulus and, simultaneously, from the spectrogram of acoustic onsets (Figure 2-A). Acoustic
onsets were derived from a neural model of auditory edge detection [19]. The two predictors
were each binned into 8 frequency bands, such that the MEG responses were predicted from a
model of the acoustic stimulus encompassing 16 time series in total. Each of the two predictors
was assessed based on how well (left-out) MEG responses were predicted by the full model,
compared to a null model in which the relevant predictor was omitted. Both predictors
significantly improve predictions (onsets: t_{max} = 12.00, $p \le .001$; envelopes: t_{max} = 9.39, $p \le .001$),
with an anatomical distribution consistent with sources in HG and STG bilaterally (Figure 2-B).
Since this localization agrees with findings from intracranial recordings [8,17], results were
henceforth analyzed in an auditory region of interest (ROI) restricted to these two anatomical
landmarks (Figure 2-C). When averaging the model fits in this ROI, almost all subjects showed
responses associated with both predictors (Figure 2-D).

Figure 2. **MEG responses to clean speech**. A) Schematic illustration of the neurally inspired
acoustic edge detector model, which was used to generate onset representations. The signal at
each frequency band was passed through multiple parallel pathways, with increasing delays, so
that an "edge detector" receptive field could detect the occurrence of fast increases. Final half-
wave rectification (HWR) removed the negative sections. An excerpt from a gammatone
spectrogram ("envelope") and the corresponding onset representation is shown for illustration
purposes. B) Regions of significant explanatory power of onset and envelope representations,
consistent with a main source in bilateral auditory cortex ($p \leq .05$, corrected for whole brain
analysis). C) Region of interest (ROI) used for the analysis of response functions, including
superior temporal gyrus and Heschl's gyrus. An arrow indicates the average dominant current
direction in the ROI (upward current), determined through the first principal component of
response power. D) Individual subject model fit improvements, averaged over the ROI in the left
and right hemisphere (LH/RH). E) Spectro-temporal response functions (STRFs) corresponding to
onset and envelope representations in the ROI; the onset STRF exhibits a clear pair of positive
and negative peaks, while peaks in the envelope STRF are less well-defined. Different color
curves reflect the frequency bins, as indicated next to the onset and envelope spectrograms in
panel A. Shaded areas indicate the within-subject standard error (SE) [31]. Regions in which
STRFs differ significantly from 0 are marked with more saturated (less faded) colors ($p \leq .05$,
corrected for time/frequency).

Auditory cortical STRFs were summarized for each subject and hemisphere using a subject-
specific spatial filter based on principal component analyses of overall STRF power in the ROI.
The average direction of that spatial filter replicates the direction of the well-known auditory
MEG response (Figure 2-C, arrows). This current vector is consistent with activity in core auditory
cortex and the superior temporal plane. Since MEG sensors are less sensitive to any radial
currents, expected from lateral STG areas implicated by intracranial recordings [8], we focus here
on the temporal information in STRFs rather than the spatial distribution of sources. STRFs can
be interpreted as reflecting different processing stages associated with different latencies,
possibly involving multiple regions in the superior temporal lobe. STRFs were initially separately

analyzed by hemisphere, but since none of the reported results interact significantly with
hemisphere the results shown are collapsed across hemisphere to simplify presentation.

STRFs to acoustic onsets exhibit a well-defined two-peaked shape, consistent across frequency
bands (Figure 2-E). An early, positive peak (average latency 65 ms) is followed by a later, negative
peak (126 ms). This structure closely resembles previously described auditory response functions
to envelope representations when estimated without consideration of onsets [16]. In
comparison, envelope STRFs in the present results are diminished and exhibit a less well-defined
structure. This is consistent with acoustic onsets explaining a large portion of the signal usually
attributed to the envelope; indeed, when the model was refitted with only the envelope
predictor, excluding the onset predictor, the envelope STRFs exhibited that canonical pattern
and with larger amplitudes (not shown).

STRFs have disproportionately higher amplitudes at lower frequencies (Figure 2-E), which is
consistent with previous tonotopic mapping of speech areas and may follow from the spectral
distribution of information in the speech signal [32,33]. This explanation is also supported by
simulations, where responses were generated using equal TRFs for each band, and yet estimated
STRFs exhibited higher amplitudes in lower frequency bands (Figure S1).

Auditory cortex represents ignored speech
MEG responses to a two-speaker mixture were then used to test for a neural representation of
ignored speech. Participants listened to a perceptually-equal loudness mixture of a male and a
female talker and were instructed to attend to one talker and ignore the other. The speaker to
be attended was counterbalanced across stimuli and subjects. Responses were predicted using
both onset and envelope representations for: the acoustic mixture, the attended speech source,
and the ignored source (Figure 3-A). The underlying rationale is that, since the brain does not
have direct access to the individual speech sources, if there is neural activity corresponding to
either source separately (above and beyond the mixture), this indicates that cortical responses
have reconstructed features of that source. Both predictors representing the ignored speech
significantly improve predictions of the responses in the ROI (both $p < .001$, onsets: $t_{max} = 6.70$,
envelopes: $t_{max} = 6.28$; group level statistics were evaluated with spatial permutation tests;
subject-specific model fits, averaged in the ROI are shown in Figure 3-B). This indicates that
acoustic features of the ignored speech are represented neurally even after controlling for
features of the mixture and the attended source. The remaining four predictors also significantly
increased model fits (all $p < .001$; mixture onsets: $t_{max} = 8.61$, envelopes: $t_{max} = 5.70$; attended
onsets: $t_{max} = 6.32$, envelopes: $t_{max} = 7.37$).

Figure 3. **Responses to the two-speaker mixture, using the stream-based model**. A) The envelope
and onset representations of the acoustic mixture and the two speech sources were used to
predict MEG responses. B) Individual subject model fit improvement due to each predictor,
averaged in the auditory cortex ROI. Each predictor explains neural data not accounted for by
the others. C) Auditory cortex STRFs to onsets are characterized by the same positive/negative
peak structure as STRFs to a single speaker. The early, positive peak is dominated by the mixture,
but also contains speaker-specific information. The second, negative peak is dominated by
representations of the attended speaker, followed, to a lesser extent, by those of the mixture. As
with responses to a single talker, the envelope STRFs have lower amplitudes, but they do show a
strong and well-defined effect of attention. Explicit differences between the attended and
ignored representations are shown in the bottom row. Details as in Figure 2. D) The major onset
STRF peaks representing individual speech sources are delayed compared to corresponding
peaks representing the mixture. To determine latencies, mixture-based and individual-speaker-
based STRFs were averaged across frequency (lines with shading for mean ±1 SE). Dots represent
the largest positive and negative peak for each subject between 20 and 200 ms. Note that the y-
axis is scaled by an extra factor of 4 beyond the indicated break points at y = 14 and -6.

Onset STRFs exhibit the same characteristic positive-negative pattern as for responses to a single
talker, but with reliable distinctions between the mixture and the individual speech streams
(Figure 3-C, D). The early, positive peak occurs earlier and has a larger amplitude for onsets in

the mixture than for onsets in either of the sources (latency mixture: 72 ms; attended: 81 ms, t_{25}
= 4.47, $p < .001$; ignored: 89 ms, t_{25} = 6.92, $p < .001$; amplitude mixture > attended: t_{25} = 8.41, p
$< .001$; mixture > ignored: t_{25} = 7.66, $p < .001$). This positive peak is followed by a negative peak
only in responses to the mixture (136 ms) and the attended source (150 ms; difference t_{25} =
3.20, $p = .004$). The amplitude of these negative peaks is statistically indistinguishable (t_{25} = 1.56,
$p = .132$).

The mixture predictor is not completely orthogonal to the source predictors. This might raise a
concern that a true response to the mixture might cause spurious responses to the sources.
Simulations using the same predictors as used in the experiment suggest, however, that such
contamination is unlikely to have occurred (see Figures S1 and S2).

Envelope processing is strongly modulated by selective attention
While the envelope STRFs seem to be generally less structured than those of the onsets, a
comparison of the STRFs to the attended and the ignored source revealed a strong and well-
defined effect of attention (Figure 3-C, right column). The attended-ignored difference wave
exhibits a negative peak at ~100 ms, consistent with previous work [16], and an additional
positive peak at ~200 ms. In contrast to previous work, however, a robust effect of attention on
the envelope representation starts almost as early as the very earliest responses. Thus,
accounting for responses to onset features separately reveals that envelope processing is
thoroughly influenced by attention. The reason for this might be that onsets often precede
informative regions in the spectrogram, such as the spectral detail of voiced segments. The
onsets might thus serve as cues to direct attention to specific regions in the spectrogram [28],
which would allow early attentional processing of the spectrogram, or envelope features.

Auditory cortex "un-masks" masked onsets
The above analysis using the stream-based predictors suggests that the auditory cortex
represents acoustic onsets in both speech sources separately, in addition to onsets in the
acoustic mixture. This is particularly interesting because, while envelopes combine mostly in an
additive manner, acoustic onsets may experience *negative* interference. This can be seen in the
spectrograms in Figure 3-A: the envelope mixture representation largely looks like a sum of the
two stream envelope representations. In contrast, the onset mixture representation has several
features that have a *lower* amplitude than the corresponding feature in the relevant source.
Neural representations of the source onsets thus suggest that the auditory cortex reconstructs
features that are masked in the mixture. This possibility might be related to instances of neural
filling-in, where cortical representations show evidence of filling in missing information to repair
degraded input signals [34–36]. The latency difference between mixture and source onsets
might then reflect additional processing for the recovery of underlying features that are not
directly accessible in the sensory input. However, these neural representations of stream-
specific onsets could also be the result of an acoustic representation that the mixture is simply
transparent to (unlike our gammatone representation). An example could be a secondary stage
of onset extraction after spectral sharpening; such spectral sharpening might resolve some of
the overlap among the two speakers, and thus reveal additional source specific onsets.

While the two possibilities could both explain the results described so far, they make different
predictions regarding responses to masked onsets. A passive mechanism, based on features to

which the mixture is transparent, should apply to overt and masked onsets equally (because they
appear masked only in the gammatone representations), and should thus be exhaustively
explained by the model used above. On the other hand, an active mechanism that specifically
processes masked onsets might generate an additional response modulation associated with
masked onsets. To test for such a modulation, we subdivided the stream-based onset predictors
to allow for separate responses to overt and masked onsets. The new predictors were
implemented as element-wise operations on the onset spectrograms (Figure 4-A). Specifically,
for each speech source, the new "masked onsets" predictor models the degree to which an
onset in the source is attenuated (masked) in the mixture, i.e. the amount by which a segregated
speech source exceeds the (physically presented) resultant acoustic mixture, or zero when it
does not: (*max(source – mixture, 0)*). The new "overt onsets" predictor models all other time-
frequency points, where an onset in the source is also seen as a comparable onset in the mixture
(element-wise *min(mixture, source)*). Note that with this definition the sum of overt and masked
onsets exactly equals the original speech source onset representation, i.e., the new predictors
model the same onsets, but allow response to differ depending on whether an onset is masked
or not. Replacing the two speech-source based onset predictors with the four overt/masked
onset predictors significantly improves the model fit (t_{max} = 6.81, p < .001), suggesting that
cortical responses indeed distinguish between overt and masked onsets. Each of the four new
predictors individually improved MEG response predictions, although masked onsets did so more
robustly (attended: t_{max} = 8.42, p < .001; ignored: t_{max} = 5.23, p < .001) than overt onsets
(attended: t_{max} = 3.34, p = .027; ignored: t_{max} = 3.82, p = .016; Figure 4-B); this difference could
be due to overt source onsets being more similar to the mixture onsets predictor. Critically, the
significant effect for masked onsets in the ignored source confirms that the auditory cortex
recovers masked onsets even when they occur in the ignored source.

Figure 4. **Responses to overt and masked onsets**. A) Spectrograms (note that in this figure, the onset representations are placed below the envelope representations, to aid visual comparison of the different onset representations) were transformed using element-wise operations to distinguish between overt onsets, i.e., onsets in a source that are apparent in the mixture, and masked onsets, i.e., onsets in a source that are masked in the presence of the other source. Two examples are marked by rectangles: The light blue rectangle marks a region with an overt (attended) onset, i.e., an onset in the attended source that also corresponds to an onset in the mixture. The dark blue rectangle marks a masked (attended) onset, i.e., an onset in the attended source which is not apparent in the mixture. B) All predictors significantly improve the cross-validated model fit (note that improvements were statistically tested with a test sensitive to spatial variation, whereas these plots show single-subject ROI average fits). C) The corresponding overt/masked STRFs exhibit the previously described positive-negative two peaked structure. The first, positive peak is dominated by a representation of the mixture, but also contains segregated features of the two talkers. For overt onsets, only the second, negative peak is modulated by attention. For masked onsets, even the first peak exhibits a small degree of attentional modulation. D) Responses to masked onsets are consistently delayed compared to responses to overt onsets. Details are analogous to Figure 3-D, except that the time window for

finding peaks was extended to 20 – 250 ms to account for the longer latency of masked onset
response functions. E) Direct comparison of the frequency-averaged onset TRFs highlights the
amplitude differences between the peaks. For overt onsets, the negative deflection due to
selective attention starts decreasing the response magnitude even near the maximum of the
first, positive peak. For masked onsets, the early peak reflecting attended onsets is increased
despite the subsequent enhanced negative peak. Results for envelope predictors are omitted
from this figure because they are practically indistinguishable from those in Figure 3.

Masked onsets are processed with a delay and an early effect of attention

Model comparison thus indicates that the neural representation of masked onsets is significantly
different from that of overt onsets. The analysis of STRFs suggests that this is for at least two
reasons (Figure 4-C-E): response latency differences, and a difference in the effect of selective
attention.

First, responses to masked onsets are systematically delayed compared to overt onsets (Figure 4-
D). This is true for the early, positive peak (mixture: 72 ms; attended/overt: 72 ms, masked: 91
ms, t_{25} = 2.85, p = .009; ignored/overt: 83 ms, masked: 97 ms, t_{25} = 6.11, p < .001), as well as the
later, negative peak (mixture: 138 ms; attended/overt: 133 ms, masked: 182 ms, t_{25} = 4.45, p <
.001). Thus, at each major peak, representations of masked onsets lag behind representations of
overt onsets by at least 15 ms.

Second, for overt onsets, the early representations at the positive peak appear to be
independent of the target of selective attention (Figure 4-C, top right). In contrast, for masked
onsets, even these early representations are enhanced by attention (Figure 4-C, bottom right).
This difference is confirmed in a stream (attended, ignored) by masking (overt, masked) ANOVA
on peak amplitudes with a significant interaction ($F_{(1,25)}$ = 24.45, p < .001). For overt onsets,
Figure 4-E might suggest that the early peak is actually enhanced for the ignored speaker;
however, this difference can be explained by the early onset of the second, negative response to
attended onsets, which overlaps with the earlier peak. This observation makes the early effect of
attention for masked onsets all the more impressive, because the early peak is larger *despite* the
onset of the subsequent, negative peak (note the steeper slope between positive and negative
peak for attended masked onsets). Also note that we consider the effect of attention early
relative to the peak structure of the TRFs; in terms of absolute latency, the onset of the effect of
attention is actually more similar between masked and overt onsets (cf. Figure 4-E).

Delayed response to masked onsets

Previous research has found that the latency of responses to speech increases with increasing
levels of stationary noise [37,38], suggesting a cost to recovering acoustic source information
from noise. Our results indicate that this is not a uniform delay for a given perceptual stream,
but that the delay varies for each acoustic element based on whether this element is overt or
locally masked by the acoustic background. That the delay applies to the individual onsets, rather
than the stream as a whole, implies that at this level, the stream is not processed as a
homogeneous entity, but that the acoustic elements related to the onsets constitute separate
units of analysis.

The addition of background noise is associated with increased response latencies to auditory
stimuli at least as early as the brainstem response wave-V [39]. This observed shift in latency is in
the sub-millisecond range and may have a mechanistic explanation in terms of different
populations of auditory nerve fibers: Background noise saturates high spontaneous rate fibers,
and the response is now dominated by somewhat slower, low spontaneous rate fibers [40]. In
cortical responses to simple stimuli, like tones, much larger delays are observed in the presence
of static noise, in the order of tens of milliseconds [41]. Latency shifts due to absolute signal
intensity [42] might be additive with shifts due to noise [43]. Such a non-linear increase in
response latency with intensity might be a confounding factor in our analysis, which is based on
linear methods: Compared to overt onsets, masked onsets in the ignored talker should generally
correspond to weaker onsets in the mixture. Splitting overt and masked onsets might thus
improve the model fit because it allows modeling different response latencies for different
intensity levels of onsets in the mixture, rather than reflecting a true response to the background
speech. In order to control for this possibility, we compared the model fit of the background-
aware model with a model allowing for 3 different intensity levels of onsets in the mixture (and
without an explicit representation of onsets in the ignored speaker). The background speaker-
aware model out-performed the level-aware model (t_{max} = 9.21, p < .001), suggesting that the
present results are not explained by this level-based non-linear response to the mixture.
Furthermore, a background-unaware non-linearity would not explain the difference in the effect
of attention between overt and masked onsets. Together, this suggests that the observed delay
is related to recovering acoustic source information, rather than a level-based non-linearity in
the response to the mixture. Furthermore, a study of single units in primary auditory cortex
found that neurons with delayed, noise-robust responses exhibited response properties
suggestive of network effects [44]. This is consistent with the effect of attention found here on
the early peak of masked, but not of overt onsets, i.e., the delayed responses to masked onsets
also exhibit more evidence of goal-driven processing than the corresponding responses to overt
onsets.

In sum, masking causes latency increases at different stages of the auditory system. These
latency shifts increase at successive stages of the ascending auditory pathway, as does the
preponderance of noise-robust response properties [45]. It is likely that different levels of the
auditory system employ different strategies to recover auditory signals of interest, and do so at
different time scales. Together with these considerations, our results suggest that the auditory
cortex actively recovers masked speech features, and not only of the attended, but also of the
ignored speech source.

Early effect of selective attention
Besides the shift in latency, response functions to overt and masked onsets differed in a more
fundamental way: While the early, positive response peak to overt onsets did not differentiate
between attended and ignored onsets, the early peak to masked onsets contained significantly
larger representations of attended onsets (cf. Figure 4-C). Thus, not only do early auditory
cortical responses represent masked onsets, but these representations are substantively
affected by whether the onset belongs to the attended or the ignored source. This distinction
could have several causes. In the extreme, it could indicate that the two streams are completely
segregated and represented as two distinct auditory objects. However, it might also be due to a

weighting of features based on their likelihood of belonging to the attended source. This could
be achieved, for example, through modulation of excitability based on spectro-temporal
prediction of the attended speech signal [46]. Thus, onsets that are more likely to belong to the
attended source might be represented more strongly, without yet being ascribed to one of the
sources exclusively.

One discrepancy in previous studies using extra- and intracranial recordings is that the former
were unable to detect any early effect of selective attention [7,16], whereas the latter showed a
small but consistent enhancement of feature representations associated with the attended
acoustic source signal [8]. Furthermore, an early effect of attention would also be expected
based on animal models that show task-dependent modulations of A1 responses [47,48]. Our
results show that this discrepancy may depend, in part, on which acoustic features are analyzed:
while overt acoustic onsets were not associated with an effect of selective attention, masked
onsets were.

Overall, the early difference between the attended and ignored source suggests that acoustic
information from the ignored source is represented to a lesser degree than information from the
attended source. This is consistent with evidence from psychophysics suggesting that auditory
representations of background speech are not as fully elaborated as those of the attended
foreground [49]. More generally, it is consistent with results that suggest an influence of
attention early on in auditory stream formation [50].

Stages of speech segregation through selective attention
Regardless of whether listening to a single talker or two concurrent talkers, response functions
to acoustic onsets are characterized by a prominent positive-negative two peak structure. Brain
responses to two concurrent talkers allow separating these response functions into components
related to different representations, and thus reveal different processing stages. Figure 5
presents a plausible model incorporating these new findings. Early on, the response is
dominated by a representation of the acoustic mixture, with a preliminary segregation, possibly
implemented through spectral filters [8]. This is followed by restoration of speech features that
are masked in the mixture, regardless of speaker, but with a small effect of selective attention
suggesting a more active mechanism. Finally, later responses are dominated by selective
attention, suggesting a clearly separated representation of the attended speaker as would be
expected in successful streaming.

Figure 5. **Model of onset-based stream segregation**. A model of cortical processing stages compatible with the results reported here. *Left*: The auditory scene, with additive mixture of the waveforms from the attended and the ignored speakers (red and blue, respectively). *Right*: Illustration of cortical representations at different processing stages. *Passive filtering*: At an early stage, onsets are extracted from the acoustic mixture and representations are partially segregated, possibly based on frequency. This stage corresponds to the early positive peak in onset TRFs. *Active Restoration*: A subsequent stage also includes representations of onsets in the underlying speech sources that are masked in the mixture, corresponding to the first peak in TRFs to masked onsets. At this stage, a small effect of attention suggests a preliminary selection of onsets with a larger likelihood of belonging to the attended speaker. *Streaming*: Finally, at a third stage, the response to onsets from the ignored speaker is suppressed, suggesting that now the two sources are clearly segregated (see also [8]). This stage corresponds to the second, negative peak, which is present in TRFs to mixture and attended onsets but not to ignored onsets.

An open question concerns how the overt and masked feature representations are temporally integrated. Representations of masked onsets were consistently delayed compared to those of overt onsets by approximately 20 ms (see Figure 4-D). This latency difference entails that upstream speech processing mechanisms may receive different packages of information about the attended speech source with some temporal desynchronization. While this might imply a need for a higher order corrective mechanism, it is also possible that upstream mechanisms are tolerant to this small temporal distortion. A misalignment of 20 ms is small compared to the normal temporal variability encountered in speech (although phonetic contrasts do exist where a distortion of a few tens of milliseconds would be relevant). Indeed, in audio-visual speech perception, temporal misalignment up to 100 ms between auditory and visual input can be tolerated [51].

Broadly, the new results are consistent with previous findings that early cortical responses are dominated by the acoustic mixture, rather than receiving pre-segregated representations of the individual streams [7,8]. However, the new results do show evidence of an earlier, partial segregation, in the form of representations of acoustic onsets which are segregated from the mixture, though not grouped into separate streams. Since these early representations do not

459 strictly distinguish between the attended and the ignored speaker they likely play the role of an
460 intermediate step in extracting the information needed to selectively attend to one of the two
461 speakers. Overall, these results are highly consistent with object-based models of auditory
462 attention, in which perception depends on an interplay between bottom-up analysis and
463 formation of local structure, and top-down selection and global grouping, or streaming [14,52].

Implications for processing of "ignored" acoustic sources

The interference in speech perception from a second talker can be very different from the
interference caused by non-speech sounds. For instance, music is cortically segregated from
speech even when both signals are unattended, consistent with a more automatic segregation,
possibly due to distinctive differences in acoustic signal properties [22]. In contrast, at moderate
signal to noise ratios (SNRs), a second talker causes much more interference with speech
perception than a comparable non-speech masker. Interestingly, this interference manifests not
just in the inability to hear attended words, but in intrusions of words from the ignored talker
[53]. The latter fact in particular has been interpreted as evidence that ignored speech might be
segregated and processed to a relatively high level. On the other hand, listeners seem to be
unable to process words in more than one speech source at a time, even when the sources are
spatially separated [54]. Furthermore, demonstrations of lexical processing of ignored speech
are rare and usually associated with specific perceptual conditions such as dichotic presentation
[55]. Consistent with this, recent EEG/MEG evidence suggests that unattended speech is not
processed in a time-locked fashion at the lexical [12] or semantic [13] level. The results described
here, showing systematic recovery of acoustic features from the ignored speech source, suggest
a potential explanation for the increased interference from speech compared to other maskers.
Representing onsets in two speech sources could be expected to increase cognitive load
compared to detecting onsets of a single source in stationary noise. These representations of
ignored speech might also act as bottom-up cues and cause the tendency for intrusions from the
ignored talker. They might even explain why a salient and overlearned word, such as one's own
name [56], might sometimes capture attention, which could happen based on acoustic rather
than lexical analysis [57]. Finally, at very low SNRs the behavioral pattern can invert, and a
background talker can be associated with better performance than stationary noise maskers
[53]. In such conditions, there might be a benefit of being able to segregate the ignored speech
source and use this information strategically [21].

An open question is how the auditory system deals with the presence of multiple background
speakers. When there are multiple background speakers, does the auditory system attempt to
unmask the different speakers all separately, or are they represented as a unified background
[7]? An attempt to isolate speech features even from multiple background talkers might
contribute to the overly detrimental effect of babble noise with a small number of talkers [58].

Conclusions

How do listeners succeed in selectively listening to one of two concurrent talkers? Our results
suggest that active recovery of acoustic onsets plays a critical role. Early responses in the
auditory cortex represent not only overt acoustic onsets, but also reconstruct acoustic onsets in
the speech sources that are masked in the mixture, even if they originate from the ignored
speech source. This suggests that early responses, in addition to representing a spectro-temporal

decomposition of the mixture, actively reconstruct acoustic features that could originate from
either speech source. Consequently, these early responses make comparatively complex
acoustic features from both speech sources available for downstream processes, thus enabling
both selective attention and bottom-up effects of salience and interference.

Acknowledgements

506 This work was supported by a National Institutes of Health grant R01-DC014085 (to J.Z.S.) and by
507 a University of Maryland Seed Grant (to L.E.H. and J.Z.S.). We would like to thank Shihab
508 Shamma for several fruitful discussions and Natalia Lapinskaya for her help in collecting data and
509 for excellent technical support.

Author contributions

511 J.Z.S. and L.E.H. designed experiment and secured funding. C.B. and J.Z.S. analyzed data and
512 wrote the manuscript. A.J., C.B. and J.Z.S. performed simulations.

Materials and Methods

Participants

The data analyzed here have been previously used in an unrelated analysis [12] and can be
retrieved from the Digital Repository at the University of Maryland
(http://hdl.handle.net/1903/21109). MEG responses were recorded from 28 native speakers of
English, recruited by media advertisements from the Baltimore area. Participants with medical,
psychiatric or neurological illnesses, head injury, and substance dependence or abuse were
excluded. All participants provided written informed consent in accordance with the University
of Maryland Baltimore Institutional Review Board and were paid for their participation. Data
from two participants were excluded, one due to corrupted localizer measurements, and one
due to excessive magnetic artifacts associated with dental work, resulting in a final sample of 18
male and 8 female participants with mean age 45.2 (range 22 - 61).

Stimuli

Two chapters were selected from an audiobook recording of A Child's History of England by
Charles Dickens, one chapter read by a male and one by a female speaker (https://librivox.org/a-
childs-history-of-england-by-charles-dickens/, chapters 3 and 8, respectively). Four 1-minute
long segments were extracted from each chapter (referred to as male-1 through 4 and female 1
through 4). Pauses longer than 300 ms were shortened to an interval randomly chosen between
250 and 300 ms, and loudness was matched perceptually (such that either speaker was deemed
equally easy to attend to). Two-talker stimuli were generated by additively combining two
segments, one from each speaker, with an initial 1 s period containing only the to-be attended
speaker (mix-1 through 4 were constructed by mixing male-1 and female-1, through 4).

Procedure

During MEG data acquisition, participants lay supine and were instructed to keep their eyes
closed to minimize ocular artifacts and head movement. Stimuli were delivered through foam
pad earphones inserted into the ear canal at a comfortably loud listening level, approximately 70
dB SPL.

Participants listened four times to mix-1, while attending to one speaker and ignoring the other
(which speaker they attended to was counterbalanced across participants), then 4 times to mix-2
while attending to the other speaker. After each segment, participants answered a question
relating to the content of the attended stimulus. Then, the four segments just heard were all
presented once each, as single talkers. The same procedure was repeated for stimulus segments
3 and 4.

Data acquisition and preprocessing

Brain responses were recorded with a 157 axial gradiometer whole head MEG system (KIT,
Kanazawa, Japan) inside a magnetically shielded room (Vacuumschmelze GmbH & Co. KG,
Hanau, Germany) at the University of Maryland, College Park. Sensors (15.5 mm diameter) are
uniformly distributed inside a liquid-He dewar, spaced ~25 mm apart, and configured as first-
order axial gradiometers with 50 mm separation and sensitivity better than 5 fT·Hz$^{-1/2}$ in the
white noise region (> 1 KHz). Data were recorded with an online 200 Hz low-pass filter and a 60
Hz notch filter at a sampling rate of 1 kHz.

Recordings were pre-processed using mne-python [59]. Flat channels were automatically
detected and excluded. Extraneous artifacts were removed with temporal signal space
separation [60]. Data were filtered between 1 and 40 Hz with a zero-phase FIR filter (mne-
python 0.15 default settings). Extended infomax independent component analysis [61] was then
used to remove ocular and cardiac artifacts. Responses time-locked to the onset of the speech
stimuli were extracted and resampled to 100 Hz. For responses to the two-talker mixture, the
first second of data, in which only the to-be attended talker was heard, was discarded.

Five marker coils attached to participants' head served to localize the head position with respect
to the MEG sensors. Two measurements, one at the beginning and one at the end of the
recording were averaged. The FreeSurfer [62] ''fsaverage'' template brain was coregistered to
each participant's digitized head shape (Polhemus 3SPACE FASTRAK) using rotation, translation,
and uniform scaling. A source space was generated using four-fold icosahedral subdivision of the
white matter surface, with source dipoles oriented perpendicularly to the cortical surface.
Minimum ℓ2 norm current estimates [63,64] were computed for all data. Initial analysis was
performed on the whole brain as identified by the FreeSurfer "cortex" label. Subsequent
analyses were restricted to sources in the STG and Heschl's gyrus as identified in the ''aparc''
parcellation [65].

Predictor variables

Predictor variables were based on gammatone spectrograms sampled at 256 frequencies,
ranging from 20 to 5000 Hz in ERB space [66], resampled to 1 kHz and scaled with exponent 0.6
[67].

Acoustic onset representations were computed by applying an auditory edge detection model
[19] independently to each frequency band of the spectrogram. The model was implemented
with a delay layer with 10 delays ranging from τ_2 = 3 to 5 ms, a saturation scaling factor of $C =$
30, and a receptive field based on the derivative of a Gaussian window with $SD = 2$ ms.
Negative values in the resulting onset spectrogram were set to 0.

Onset representations of the two speakers were split into masked and overt onsets using
element-wise operations on the onset spectrograms. Masked onsets were defined by the extent
to which onsets were larger in the source than in the mixture:

583
$$o_{masked} = \max\left(o_{source} - o_{mixture}, 0\right)$$

584   Overt onsets were onsets that were not masked, i.e., speech source onsets that were also visible
585   in the mixture:

586
$$o_{overt} = o_{source} - o_{masked} \equiv min\left(o_{source}, o_{mixture}\right)$$

Using this procedure, approximately 67% of total onset magnitudes ($\ell 1$ norm) was assigned to
overt onsets, and 33% to masked onsets.

For model estimation, envelope and onset spectrograms were then binned into 8 frequency
bands equally spaced in ERB space (omitting frequencies below 100 Hz because the female
speaker had little power below that frequency) and resampled to match the MEG data. As part
of the reverse correlation procedure, each predictor time series (i.e., each frequency bin) was
scaled by its $\ell 1$ norm over time.

For testing an intensity-based nonlinear response (see *Delayed response to masked onsets*), the
onset predictor was split into three separate predictors, one for each of three intensity levels.
For each of the 8 frequency bins, individual onsets were identified as contiguous non-zero
elements; each onset was assigned an intensity based on the sum of its elements, and the onsets
were then assigned to one of three predictors based on intensity tertiles (calculated separately
for each band). This resulted in three 8-band onset spectrograms modeling low, medium and
high intensity onsets.

Reverse correlation
Spectro-temporal response functions (STRFs) were computed independently for each virtual
current source [see 68]. The neural response at time t, y_t was predicted from the sum of N
predictor variables x_n convolved with a corresponding response function h_n of length T:

605
$$\hat{y}_t = \sum_n^N \sum_\tau^T h_{n,\tau} \cdot x_{i,t-\tau}$$

STRFs were generated from a basis of 50 ms wide Hamming windows and were estimated using
an iterative coordinate descent algorithm [69] to minimize the $\ell 1$ error.

For model evaluation, left-out data were predicted using 4-fold cross-validation. Folds were
created by assigning successive trials to the different folds in order (1, 2, 3, 4, 1, 2, …). In an outer
loop, the responses in each fold were predicted with STRFs estimated from the remaining 3
folds. These predictions, combined, served to calculate the correlation between measured and
predicted responses used for model tests. In an inner loop, each of the 3 estimation folds was in
turn used as validation set for STRFs trained on the two remaining folds. STRFs were iteratively
improved based on the maximum error reduction in the training set (the steepest coordinate
descent) and validated in the validation set. Whenever a predictor time series (i.e., one
spectrogram bin) would have caused an increasing in the error in the validation set, the kernel
for this predictor was frozen, continuing until all predictors were frozen (see [70] for further

details). The 3 STRFs from the inner loop were averaged to predict responses in the left-out
testing data.

Model tests

621   Each spectrogram comprising of 8 time series (frequency bins) was treated as an individual
622   predictor. Speech in quiet was modeled using the (envelope) spectrogram and acoustic onsets:

623
$$MEG \sim o + e$$

Where o = onsets and e = envelope. Models were estimated with STRFs with $T = [0, \dots, 500)$
ms. Model quality was quantified as the Fisher z-scored Pearson correlation between actual and
predicted response. Maps of z-values were smoothed with a Gaussian kernel (SD = 5 mm). In
order to test the predictive power of each predictor, a corresponding null model was generated
by removing that predictor. For each predictor, the model quality of the full model was
compared with the model quality of the corresponding null model using a mass-univariate
related measures t-test with threshold-free cluster enhancement [71] and a null distribution
based on 10,000 permutations [72 for further details]. In addition to p-values we report t_{max} for
each comparison, the largest t-value in the significant area, for effect size comparison. For single
talker speech (Figure 2) this test included the whole cortex (as labeled by FreeSurfer). For
subsequent tests of the two talker condition the same tests were used, but the test area was
restricted to the auditory ROI comprising the STG and transverse temporal gyrus in each
hemisphere.

Initially, responses to speech in noise (Figure 3) was predicted from:

638
$$MEG \sim o_{mix} + o_{att} + o_{ign} + e_{mix} + e_{att} + e_{ign}$$

639   Where mix = mixture, att = attended, ign = ignored. Masked onsets (Figure 4) were analyzed
640   with:

641
$$MEG \sim o_{mix} + o_{att,overt} + o_{att,masked} + o_{ign,overt} + o_{ign,masked} + e_{mix} + e_{att} + e_{ign}$$

642   In order to test for a level-dependent nonlinear response to onsets in the mixture, this model
643   was compared with the following:

644
$$MEG \sim o_{mix-low} + o_{mix-mid} + o_{mix-high} + o_{att,overt} + o_{att,masked} + e_{mix} + e_{att} + e_{ign}$$

645   Where mix-low, -mid and -$high$ = mixture low, mid and high intensity. This model has the same
646   number of predictors, but assumes no awareness of onsets in the ignored speaker.

STRF analysis

To evaluate STRFs, the corresponding model was refit with $T = [-100, \dots, 500)$ ms to include
an estimate of baseline activity (due to occasional edge artifacts, STRFs are displayed between -
50 to 450 ms). Using the same 4-fold split of the data as for model fits, 4 STRF estimates were
averaged, each using one fold of the data for validation and the remaining 3 for training. Since
predictors and responses were $\ell 1$ normalized for the reverse correlation, and STRFs were
analyzed in this normalized space, STRFs provide an SNR-like measure of response strength at
different latencies for each subject.

Auditory STRFs were computed for each subject and hemisphere as a weighted sum of STRFs in
the auditory ROI encompassing the STG and transverse temporal (Heschl's) gyrus. Weights were

computed separately for each subject and hemisphere. First, each source point was assigned a
vector with direction orthogonal to the cortical surface, and length equal to the total TRF power
for responses to clean speech (sum of squares over time, frequency and predictor). The ROI
direction was then determined as the first principal component of these vectors, with the sign
adjusted to be positive on the inferior-superior axis. A weight was then assigned to each source
as the dot product of this direction with the source's direction, and these weights were
normalized within the ROI.

In order to make STRFs more comparable across subjects, they were smoothed on the frequency
axis with a Hamming window of width 7 bins. STRFs were statistically analyzed in the time range
$[0, …, 450)$ ms using mass-univariate t-tests and ANOVAs, with p-values calculated from null
distributions based on the maximum statistic (t, F) in 10,000 permutations [73].

For visualization and peak analysis, STRFs were upsampled to 500 Hz. Peak latencies were
computed by first averaging auditory STRFs along the frequency axis, and then finding the largest
or smallest value in each subject's TRF in a window of $[20, 200)$ ms for single speaker and
stream-based analysis (Figures 2 and 3) or $[20, 250)$ ms for the masked onset analysis (Figure 4).
Reported peak latencies are always average latencies across subject.

References

1. Bregman AS. Auditory scene analysis: the perceptual organization of sound. Cambridge,
Mass: MIT Press; 1990.

2. Cherry EC. Some Experiments on the Recognition of Speech, with One and with Two Ears. J
Acoust Soc Am. 1953;25: 975–979. doi:10.1121/1.1907229

3. McDermott JH. The cocktail party problem. Curr Biol. 2009;19: R1024–R1027.
doi:10.1016/j.cub.2009.09.005

4. Middlebrooks JC, Simon JZ, Popper AN, Fay RR, editors. The Auditory system at the cocktail
party. Cham: Springer International Publishing; 2017.

5. Brungart DS, Simpson BD. The effects of spatial separation in distance on the informational
and energetic masking of a nearby speech signal. J Acoust Soc Am. 2002;112: 664–676.
doi:10.1121/1.1490592

6. Kidd G, Mason CR, Swaminathan J, Roverud E, Clayton KK, Best V. Determining the
energetic and informational components of speech-on-speech masking. J Acoust Soc Am.
2016;140: 132–144. doi:10.1121/1.4954748

7. Puvvada KC, Simon JZ. Cortical Representations of Speech in a Multitalker Auditory Scene. J
Neurosci. 2017;37: 9189–9196. doi:10.1523/JNEUROSCI.0938-17.2017

8. O'Sullivan J, Herrero J, Smith E, Schevon C, McKhann GM, Sheth SA, et al. Hierarchical
Encoding of Attended Auditory Objects in Multi-talker Speech Perception. Neuron. 2019;
S0896627319307809. doi:10.1016/j.neuron.2019.09.007

9.   Zion Golumbic EM, Ding N, Bickel S, Lakatos P, Schevon CA, McKhann GM, et al.
Mechanisms Underlying Selective Neuronal Tracking of Attended Speech at a "Cocktail
Party." Neuron. 2013;77: 980–991. doi:10.1016/j.neuron.2012.12.037

10.   Broadbent DE. Perception and communication. London: Pergamon Press; 1958.

11.   Lachter J, Forster KI, Ruthruff E. Forty-five years after Broadbent (1958): Still no
identification without attention. Psychol Rev. 2004;111: 880–913. doi:10.1037/0033-
295X.111.4.880

12.   Brodbeck C, Hong LE, Simon JZ. Rapid Transformation from Auditory to Linguistic
Representations of Continuous Speech. Curr Biol. 2018;28: 3976-3983.e5.
doi:10.1016/j.cub.2018.10.042

13.   Broderick MP, Anderson AJ, Liberto GMD, Crosse MJ, Lalor EC. Electrophysiological
Correlates of Semantic Dissimilarity Reflect the Comprehension of Natural, Narrative
Speech. Curr Biol. 2018;28: 803-809.e3. doi:10.1016/j.cub.2018.01.080

14.   Shinn-Cunningham BG. Object-based auditory and visual attention. Trends Cogn Sci.
2008;12: 182–186. doi:10.1016/j.tics.2008.02.003

15.   Lalor EC, Foxe JJ. Neural responses to uninterrupted natural speech can be extracted with
precise temporal resolution. Eur J Neurosci. 2010;31: 189–193. doi:10.1111/j.1460-
9568.2009.07055.x

16.   Ding N, Simon JZ. Emergence of neural encoding of auditory objects while listening to
competing speakers. Proc Natl Acad Sci U S A. 2012;109: 11854–9.
doi:10.1073/pnas.1205381109

17.   Hamilton LS, Edwards E, Chang EF. A Spatial Map of Onset and Sustained Responses to
Speech in the Human Superior Temporal Gyrus. Curr Biol. 2018;28: 1860-1871.e4.
doi:10.1016/j.cub.2018.04.033

18.   Daube C, Ince RAA, Gross J. Simple Acoustic Features Can Explain Phoneme-Based
Predictions of Cortical Responses to Speech. Curr Biol. 2019;29: 1924-1937.e9.
doi:10.1016/j.cub.2019.04.067

19.   Fishbach A, Nelken I, Yeshurun Y. Auditory Edge Detection: A Neural Model for Physiological
and Psychoacoustical Responses to Amplitude Transients. J Neurophysiol. 2001;85: 2303–
2323. doi:10.1152/jn.2001.85.6.2303

20.   Zhou Y, Wang X. Cortical Processing of Dynamic Sound Envelope Transitions. J Neurosci.
2010;30: 16741–16754. doi:10.1523/JNEUROSCI.2016-10.2010

21.   Fiedler L, Wöstmann M, Herbst SK, Obleser J. Late cortical tracking of ignored speech
726   facilitates neural selectivity in acoustically challenging conditions. NeuroImage. 2019;186:
727   33–42. doi:10.1016/j.neuroimage.2018.10.057

728   22.   Hausfeld L, Riecke L, Valente G, Formisano E. Cortical tracking of multiple streams outside
729   the focus of attention in naturalistic auditory scenes. NeuroImage. 2018;181: 617–626.
730   doi:10.1016/j.neuroimage.2018.07.052

731   23.   Petersen EB, Wöstmann M, Obleser J, Lunner T. Neural tracking of attended versus ignored
732   speech is differentially affected by hearing loss. J Neurophysiol. 2017;117: 18–27.
733   doi:10.1152/jn.00527.2016

734   24.   Fiedler L, Wöstmann M, Graversen C, Brandmeyer A, Lunner T, Obleser J. Single-channel in-
735   ear-EEG detects the focus of auditory attention to concurrent tone streams and mixed
736   speech. J Neural Eng. 2017;14: 036020. doi:10.1088/1741-2552/aa66dd

737   25.   Stilp CE, Kluender KR. Cochlea-scaled entropy, not consonants, vowels, or time, best
738   predicts speech intelligibility. Proc Natl Acad Sci U S A. 2010;107: 12387–12392.
739   doi:10.1073/pnas.0913625107

740   26.   Koning R, Wouters J. The potential of onset enhancement for increased speech intelligibility
741   in auditory prostheses. J Acoust Soc Am. 2012;132: 2569–2581. doi:10.1121/1.4748965

742   27.   Elhilali M, Ma L, Micheyl C, Oxenham AJ, Shamma SA. Temporal Coherence in the
743   Perceptual Organization and Cortical Representation of Auditory Scenes. Neuron. 2009;61:
744   317–329. doi:10.1016/j.neuron.2008.12.005

745   28.   Bregman AS, Ahad P, Kim J, Melnerich L. Resetting the pitch-analysis system: 1. Effects of
746   rise times of tones in noise backgrounds or of harmonics in a complex tone. Percept
747   Psychophys. 1994;56: 155–162. doi:10.3758/BF03213894

748   29.   Bregman AS, Ahad PA, Kim J. Resetting the pitch-analysis system. 2. Role of sudden onsets
749   and offsets in the perception of individual components in a cluster of overlapping tones. J
750   Acoust Soc Am. 1994;96: 2694–2703. doi:10.1121/1.411277

751   30.   Hukin RW, Darwin CJ. Comparison of the effect of onset asynchrony on auditory grouping in
752   pitch matching and vowel identification. Percept Psychophys. 1995;57: 191–196.
753   doi:10.3758/BF03206505

754   31.   Loftus GR, Masson MEJ. Using confidence intervals in within-subject designs. Psychon Bull
755   Rev. 1994;1: 476–490. doi:10.3758/BF03210951

756   32.   Moerel M, De Martino F, Formisano E. Processing of Natural Sounds in Human Auditory
757   Cortex: Tonotopy, Spectral Tuning, and Relation to Voice Sensitivity. J Neurosci. 2012;32:
758   14205–14216. doi:10.1523/JNEUROSCI.1388-12.2012

33.　Hullett PW, Hamilton LS, Mesgarani N, Schreiner CE, Chang EF. Human Superior Temporal
Gyrus Organization of Spectrotemporal Modulation Tuning Derived from Speech Stimuli. J
Neurosci. 2016;36: 2014–2026. doi:10.1523/JNEUROSCI.1779-15.2016

34.　Cervantes Constantino F, Simon JZ. Dynamic cortical representations of perceptual filling-in
for missing acoustic rhythm. Sci Rep. 2017;7: 17536. doi:10.1038/s41598-017-17063-0

35.　Leonard MK, Baud MO, Sjerps MJ, Chang EF. Perceptual restoration of masked speech in
human cortex. Nat Commun. 2016;7: 13619. doi:10.1038/ncomms13619

36.　Cervantes Constantino F, Simon JZ. Restoration and Efficiency of the Neural Processing of
Continuous Speech Are Promoted by Prior Knowledge. Front Syst Neurosci. 2018;12: 56.
doi:10.3389/fnsys.2018.00056

37.　Ding N, Simon JZ. Adaptive Temporal Encoding Leads to a Background-Insensitive Cortical
Representation of Speech. J Neurosci. 2013;33: 5728–5735. doi:10.1523/JNEUROSCI.5297-
12.2013

38.　Billings CJ, McMillan GP, Penman TM, Gille SM. Predicting Perception in Noise Using Cortical
Auditory Evoked Potentials. J Assoc Res Otolaryngol. 2013;14: 891–903.
doi:10.1007/s10162-013-0415-y

39.　Burkard RF, Sims D. A Comparison of the Effects of Broadband Masking Noise on the
Auditory Brainstem Response in Young and Older Adults. Am J Audiol. 2002;11: 13–22.
doi:10.1044/1059-0889(2002/004)

40.　Mehraei G, Hickox AE, Bharadwaj HM, Goldberg H, Verhulst S, Liberman MC, et al. Auditory
Brainstem Response Latency in Noise as a Marker of Cochlear Synaptopathy. J Neurosci.
2016;36: 3755–3764. doi:10.1523/JNEUROSCI.4460-15.2016

41.　Billings CJ, Tremblay KL, Stecker GC, Tolin WM. Human evoked cortical activity to signal-to-
noise ratio and absolute signal level. Hear Res. 2009;254: 15–24.
doi:10.1016/j.heares.2009.04.002

42.　Drennan DP, Lalor EC. Cortical Tracking of Complex Sound Envelopes: Modeling the
Changes in Response with Intensity. eneuro. 2019;6: ENEURO.0082-19.2019.
doi:10.1523/ENEURO.0082-19.2019

43.　Teschner MJ, Seybold BA, Malone BJ, Hüning J, Schreiner CE. Effects of Signal-to-Noise Ratio
on Auditory Cortical Frequency Processing. J Neurosci. 2016;36: 2743–2756.
doi:10.1523/JNEUROSCI.2079-15.2016

44.　Malone BJ, Heiser MA, Beitel RE, Schreiner CE. Background noise exerts diverse effects on
the cortical encoding of foreground sounds. J Neurophysiol. 2017;118: 1034–1054.
doi:10.1152/jn.00152.2017

45.   Rabinowitz NC, Willmore BDB, King AJ, Schnupp JWH. Constructing Noise-Invariant
794   Representations of Sound in the Auditory Pathway. Zador AM, editor. PLoS Biol. 2013;11:
795   e1001710. doi:10.1371/journal.pbio.1001710

796   46.   Lakatos P, Musacchia G, O'Connel MN, Falchier AY, Javitt DC, Schroeder CE. The
797   Spectrotemporal Filter Mechanism of Auditory Selective Attention. Neuron. 2013;77: 750–
798   761. doi:10.1016/j.neuron.2012.11.034

799   47.   Fritz J, Shamma S, Elhilali M, Klein D. Rapid task-related plasticity of spectrotemporal
800   receptive fields in primary auditory cortex. Nat Neurosci. 2003;6: 1216–1223.
801   doi:10.1038/nn1141

802   48.   Atiani S, Elhilali M, David SV, Fritz JB, Shamma SA. Task Difficulty and Performance Induce
803   Diverse Adaptive Patterns in Gain and Shape of Primary Auditory Cortical Receptive Fields.
804   Neuron. 2009;61: 467–480. doi:10.1016/j.neuron.2008.12.027

805   49.   Shinn-Cunningham BG, Lee AKC, Oxenham AJ. A sound element gets lost in perceptual
806   competition. Proc Natl Acad Sci. 2007;104: 12223–12227. doi:10.1073/pnas.0704641104

807   50.   Carlyon RP. How the brain separates sounds. Trends Cogn Sci. 2004;8: 465–471.
808   doi:10.1016/j.tics.2004.08.008

809   51.   van Wassenhove V, Grant KW, Poeppel D. Temporal window of integration in auditory-
810   visual speech perception. Neuropsychologia. 2007;45: 598–607.
811   doi:10.1016/j.neuropsychologia.2006.01.001

812   52.   Elhilali M, Xiang J, Shamma SA, Simon JZ. Interaction between Attention and Bottom-Up
813   Saliency Mediates the Representation of Foreground and Background in an Auditory Scene.
814   Griffiths TD, editor. PLoS Biol. 2009;7: e1000129. doi:10.1371/journal.pbio.1000129

815   53.   Brungart DS. Informational and energetic masking effects in the perception of two
816   simultaneous talkers. J Acoust Soc Am. 2001;109: 1101–1109. doi:10.1121/1.1345696

817   54.   Kidd G, Arbogast TL, Mason CR, Gallun FJ. The advantage of knowing where to listen. J
818   Acoust Soc Am. 2005;118: 12.

819   55.   Rivenez M, Darwin CJ, Guillaume A. Processing unattended speech. J Acoust Soc Am.
820   2006;119: 4027–4040. doi:10.1121/1.2190162

821   56.   Wood N, Cowan N. The cocktail party phenomenon revisited: How frequent are attention
822   shifts to one's name in an irrelevant auditory channel? J Exp Psychol Learn Mem Cogn.
823   1995;21: 255–260. doi:10.1037/0278-7393.21.1.255

824   57.   Woods KJP, McDermott JH. Schema learning for the cocktail party problem. Proc Natl Acad
825   Sci. 2018;115: E3313–E3322. doi:10.1073/pnas.1801614115

58.   Simpson SA, Cooke M. Consonant identification in N-talker babble is a nonmonotonic
function of N. J Acoust Soc Am. 2005;118: 2775–2778. doi:10.1121/1.2062650

59.   Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MNE
software for processing MEG and EEG data. NeuroImage. 2014;86: 446–460.
doi:10.1016/j.neuroimage.2013.10.027

60.   Taulu S, Simola J. Spatiotemporal signal space separation method for rejecting nearby
interference in MEG measurements. Phys Med Biol. 2006;51: 1759. doi:10.1088/0031-
9155/51/7/008

61.   Bell AJ, Sejnowski TJ. An Information-Maximization Approach to Blind Separation and Blind
Deconvolution. Neural Comput. 1995;7: 1129–1159. doi:10.1162/neco.1995.7.6.1129

62.   Fischl B. FreeSurfer. NeuroImage. 2012;62: 774–781.
doi:10.1016/j.neuroimage.2012.01.021

63.   Hämäläinen MS, Ilmoniemi RJ. Interpreting magnetic fields of the brain: minimum norm
estimates. Med Biol Eng Comput. 1994;32: 35–42. doi:10.1007/BF02512476

64.   Dale AM, Sereno MI. Improved Localizadon of Cortical Activity by Combining EEG and MEG
with MRI Cortical Surface Reconstruction: A Linear Approach. J Cogn Neurosci. 1993;5:
162–176. doi:10.1162/jocn.1993.5.2.162

65.   Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated
labeling system for subdividing the human cerebral cortex on MRI scans into gyral based
regions of interest. NeuroImage. 2006;31: 968–980. doi:10.1016/j.neuroimage.2006.01.021

66.   Heeris J. Gammatone Filterbank Toolkit. 2018. Available:
https://github.com/detly/gammatone

67.   Biesmans W, Das N, Francart T, Bertrand A. Auditory-Inspired Speech Envelope Extraction
Methods for Improved EEG-Based Auditory Attention Detection in a Cocktail Party Scenario.
IEEE Trans Neural Syst Rehabil Eng. 2017;25: 402–412. doi:10.1109/TNSRE.2016.2571900

68.   Brodbeck C, Presacco A, Simon JZ. Neural source dynamics of brain responses to continuous
stimuli: Speech processing from acoustics to comprehension. NeuroImage. 2018;172: 162–
174. doi:10.1016/j.neuroimage.2018.01.042

69.   David SV, Mesgarani N, Shamma SA. Estimating sparse spectro-temporal receptive fields
with natural stimuli. Netw Comput Neural Syst. 2007;18: 191–212.
doi:10.1080/09548980701609235

70.   Brodbeck C, Das P, Brooks TL, Reddigari S. Eelbrain 0.31. Zenodo; 2019.
doi:10.5281/ZENODO.3564850

71.   Smith SM, Nichols TE. Threshold-free cluster enhancement: Addressing problems of
smoothing, threshold dependence and localisation in cluster inference. NeuroImage.
2009;44: 83–98. doi:10.1016/j.neuroimage.2008.03.061

72.   Brodbeck C, Brooks TL, Das P, Reddigari S. Eelbrain 0.30. Zenodo; 2019.
doi:10.5281/zenodo.2653785

73.   Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. J Neurosci
Methods. 2007;164: 177–190. doi:10.1016/j.jneumeth.2007.03.024

866