Boldness and exploration vary between shell morphs but not environmental

contexts in the snail *Cepaea nemoralis*

Running title: Personality and shell morph in snails

Maxime Dahirel*[1,2] ; Valentin Gaudu[1] ; Armelle Ansart[1]

*corresponding author: maxime.dahirel@yahoo.fr

[1] Univ Rennes, CNRS, ECOBIO (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35042 Rennes,

France

[2] INRAE, Université Côte d'Azur, CNRS, ISA, F-06903 Sophia-Antipolis, France

Acknowledgments

We thank Youn Henry and Kévin Tougeron for helpful pointers on the link between behavior and

thermal tolerance.

Conflicts of interest

The authors have no conflict of interest to declare.

Data accessibility

Data and code to reproduce all analyses are available on GitHub

(https://github.com/mdahirel/cepaea-personality-2017) and archived in Zenodo (doi:

10.5281/zenodo.3899042 ; version 1.1).

Abstract

Understanding the maintenance of individual behavioral variation (or animal personalities) in

populations, and predicting its consequences, are key challenges in behavioral ecology. Studying the

association between repeatable behavior and other traits under selection may shed light on these

selective pressures. We used the model snail *Cepaea nemoralis* to examine whether individual

behavioral is associated with shell morph, a key trait that has been extensively studied in relation to

thermal tolerance and predator avoidance, and which is known to be under strict genetic control in

this species. We quantified proxies of boldness and exploration in snails of three morphs coming

from two contrasted thermal environments. We show that both behaviors were repeatable at the

individual level in *C. nemoralis*. Behavior was associated with shell morph, with the darker morph

(five-banded) being consistently shyer and slower to explore. There was no evidence that

experienced thermal environment influenced behavior at the individual level, whether we

considered past environment or experimental temperature. Boldness and exploration were

correlated at the individual level in a common syndrome. We discuss what these results may tell us

about the type of selection exerted by predators. We also detail how our results hint to a genetic link

between shell morph and behavior, and the evolutionary implications of such a link. Finally, we

discuss how our results combined with decades of evolutionary research make *C. nemoralis* a very

valuable model to study the evolution of behavior in response to environmental changes.

Keywords animal personality; behavioral syndromes; Gastropoda; multivariate multilevel model;

shell color; temperature

Introduction

A key question in behavioral ecology, and more broadly in evolutionary ecology, is how to explain the persistence of variation in phenotypic traits. Although behavior is often seen as highly labile and dynamically adjustable to experienced conditions, individuals of many animal species actually exhibit "personalities", i.e. behave consistently across time and contexts, and differ consistently from each other (Kralj-Fišer & Schuett, 2014; Réale, Reader, Sol, McDougall, & Dingemanse, 2007; Sih, Bell, Johnson, & Ziemba, 2004). This inter-individual variation persists even when better adjustments of behaviors to environmental conditions would be adaptive, and the ability to tune behavior to conditions may itself also vary among individuals (variation in "behavioral reaction norms"; Dingemanse, Kazem, Réale, & Wright, 2010). Moreover, behaviors are often correlated together and with other traits in multivariate syndromes (Réale et al., 2007; Royauté, Berdal, Garrison, & Dochtermann, 2018; Santostefano, Wilson, Niemelä, & Dingemanse, 2017), further constraining the range of behavioral phenotypes that are on display in populations (Dochtermann & Dingemanse, 2013).

State-dependent behavior is often invoked as one of the key mechanisms/frameworks potentially explaining both adaptive correlations/feedbacks between behaviors and other traits, and the maintenance of individual behavioral variation (Sih et al., 2015; Wolf & McNamara, 2012; Wolf & Weissing, 2010). If costs and benefits of behaviors vary depending on individual state variables, including morphology, size, past experienced environment... which may be less labile or even fixed at the individual level, then we should expect individuals differing in state to adaptively differ in behavior as well (Wolf & Weissing, 2010). The pace-of-life hypothesis, which ties several axes of behavioral variation to underlying differences in life history and metabolism along a common fast-slow axis (Réale et al., 2010; Wolf & McNamara, 2012; Wright, Bolstad, Araya-Ajoy, & Dingemanse, 2019), can be seen under this lens. Other examples include cases of phenotypic compensation, where predation risk can either be mitigated by behavioral changes or morphological defenses,

leading to a positive association between risk-taking behavior and morphological defenses (e.g.

Ahlgren, Chapman, Nilsson, & Brönmark, 2015). Quantitative genetics and/or experimental evolution

approaches may provide evidence of the evolution of state-behavior associations (e.g. Kern,

Robinson, Gass, Godwin, & Langerhans, 2016). In some species in which this may be difficult, we

believe a fruitful alternative may be to study the association between personality and state traits

already known to be fully genetically determined with little to no plasticity, and to study it across a

range of environments known to select on the state variable.

The grove snail *Cepaea nemoralis* (Linnaeus 1758) (family Helicidae) has a long history as a model in

evolutionary biology, in relation to its conspicuous shell polymorphism (reviewed by Jones, Leith, &

Rawlings, 1977; Ożgo, 2009)(Fig. 1A-B). Genetic variation in terms of shell background color (from

pale yellow to brown, but usually divided in yellow, pink and brown; Davison, Jackson, Murphy, &

Reader, 2019) and in the number or width of dark bands has been well described in Mendelian terms

(Jones et al., 1977). Modern genomics studies now aim to pinpoint the actual

molecular/physiological underpinnings of shell color (Kerkvliet, Boer, Schilthuizen, & Kraaijeveld,

2017; Richards et al., 2013).

In *C. nemoralis*, lighter colored shells absorb less heat and allow snails to maintain a lower body

temperature (Heath, 1975) and higher water content (Chang, 1991). Many studies have shown that

lighter (darker) snails are as a consequence selectively advantaged in hotter/sunnier (colder/shaded)

environments, whether one looks at continental-scale latitudinal clines (Jones et al., 1977; Silvertown

et al., 2011), local-scale habitat comparisons (Kerstes, Breeschoten, Kalkman, & Schilthuizen, 2019;

Ozgo & Kinnison, 2008; Schilthuizen, 2013), or historical comparisons in the context of climate

change (Ożgo, Liew, Webster, & Schilthuizen, 2017; Ożgo & Schilthuizen, 2012). Local variations in

morph frequencies have also been tied to predation pressure, with both frequency-dependent and

crypsis-based visual selection thought to play a role (Jones et al., 1977; Surmacki, Ożarowska-

Nowicka, & Rosin, 2013, and references therein; but see Cook, 2008). In addition to visual selection,

there are indications that shell color morphs vary in the physical resistance they oppose to predators,

possibly because pigments (e.g. melanins) also influence the structural properties of shells (Kerkvliet

et al., 2017; Williams, 2017). For instance, shells with more dark bands are more resistant to crushing

in the areas of the shell attacked by rodent predators (Rosin, Kobak, Lesicki, & Tryjanowski, 2013).

Although this has been much less studied, shell morphs also vary in their behavior (Chang, 1991;

Jones, 1982; Ożgo & Kubea, 2005; Rosin et al., 2018). Part of this variation is likely the direct

consequence of differences in shell thermal properties and thus effective body temperature, as

demonstrated by experiments that created "artificial" morphs by painting shells (Tilling, 1983).

However, they probably also reflect, at least partly, intrinsic physiological differences: snail preferred

temperatures can be altered using opioid agonists or antagonists, but banded snails are less

responsive to this pharmaceutical manipulation (Kavaliers, 1992). Existing studies however have

several major shortfalls for our understanding of the association between *Cepaea* morphology and

behavioral syndromes. In particular, individuals are generally assayed once, meaning separating

within- from-among individual variation, and individual variation in average behavior versus variation

in plasticity is impossible. This also means the level of individual variation that may exist within

morphs, and how it compares with among-morph variation, has remained to our knowledge

unstudied. Additionally, all snails are often sampled from the same habitat, or habitat information is

not used in behavioral analyses, meaning there is often no way to determine how behavior responds

to selection pressures on shell color.

In this context, we investigated the existence and magnitude of personality variation and behavioral

syndromes in *C. nemoralis*, how behavior is linked to shell variation and how it is influenced by

environment of origin (sun-exposed or shaded) and currently experienced conditions. More

specifically, we make the following hypotheses:

(i) Exploration and boldness (risk-taking behavior) are both repeatable in this species, and positively

correlated in a common behavioral syndrome (Réale et al., 2010).

(ii) As snails are ectotherms, activity/exploration should increase with temperature due to increased

metabolism (over the range of temperatures suitable to movement; Abram, Boivin, Moiroux, &

Brodeur, 2017; Cloyed, Dell, Hayes, Kordas, & O'Gorman, 2019). We expect this temperature-

activity/exploration reaction norm should vary both in its slope and average value according to shell

morph and habitat of origin. Populations having evolved in sun-exposed habitats, and lighter

(unbanded) snails should be better adapted to maintain activity in the face of high temperatures (e.g.

Cloyed et al., 2019), at the possible costs of lower activity at lower temperatures (Tilling, 1983).

(iii) As boldness may increase predation risk (Hulthén et al., 2017; Smith & Blumstein, 2008), we can

expect boldness to be counter-selected in habitat × morph combinations more exposed to predation,

or an association between boldness and non-behavioral defense traits compensating for these costs.

Predictions here differ depending on the importance of frequency-dependency versus crypsis in

shaping predation: we should expect the rarer morphs to be bolder, if frequency-dependent

predation drives the evolution of shell morphs, while we should expect the less conspicuous snails to

be bolder if crypsis is prominent. Unbanded snails are expected to be more cryptic to avian

predators, especially on dry vegetation (Surmacki et al., 2013). On the other hand, the phenotypic

compensation hypothesis predicts that bolder morphs should have better defended shells (Ahlgren

et al., 2015). In *Cepaea nemoralis*, this leads to the prediction that snails with increased number of

bands, which have on average thicker shells (Rosin et al., 2013), would be bolder.

Methods

Sampled sites and snail maintenance under laboratory conditions

Snails were sampled in fall 2016 in and close to the village of Arçais, France (Fig. 1C), located roughly

in the middle of the recorded latitudinal range of *Cepaea nemoralis* (GBIF Secretariat, 2020). We

studied two sites located about 2 km apart and differing in terms of vegetation cover. One was a

garden with few isolated trees, and thus under relatively direct sun exposure all year long (hereafter

the "open habitat"; approximate location: 46° 17' 50"N, 0° 41' 30" W, Fig. 1D). The other was a 200

by 150 m deciduous forested lot, and thus fully shaded a large part of the year, especially the hottest spring and summer months ("shaded habitat"; approximately 46° 18' 01" N, 0° 42' 56" W, Fig. 1E). We only sampled snails with the three most abundant shell banding patterns: shells with no bands, three bands on the lower side of the shell, and five bands (Fig. 1B). Following previous authors (e.g. Kavaliers, 1992), we here focused for simplicity solely on band presence, and thus sampled only snails with yellow background shells, which are the most common in the study region (Silvertown et al., 2011; personal observations) and on which the contrast between shell background and dark bands is the strongest. Only adult snails were selected (recognizable by a reflected "lip" on their shell opening). Sampling for the present study was targeted and adjusted in the field to obtain roughly equal numbers of each banding pattern from both landscapes; it therefore did not allow us to make inferences on their relative abundances. The same sites were however sampled again in 2018 for a separate experiment, this time with random sampling with respect to banding pattern. As in previous studies (e.g. Schilthuizen, 2013) and reflecting potential thermal selection, the darker five-banded snails were more frequent in the shaded habitat than in the open habitat (22.3 vs. 13.5 %; see Supplementary Material S1).

We then kept snails under winter dormancy conditions (6 ± 1°C, no food or water sources) until March 2017, about 3 weeks before the start of the experiment. We then divided them in groups of 15 individuals from the same landscape, 5 (randomly chosen) of each shell phenotype. Groups were kept under controlled conditions (20 ± 1°C, L:D 16:8) in 8.5 × 15 × 12 cm polyethylene boxes lined with 1-2 cm of soil kept humid at the bottom. Snails had *ad libitum* access to prepared snail food (cereal flour supplemented with calcium, Hélinove, Saint Paul en Pareds, France) in a Petri dish. We gave each snail a unique ID written on the side of their shell with a paint marker (uni Posca, Mitsubishi Pencil Co., Ltd, Tokyo, Japan; Henry & Jarne, 2007). A total of 360 snails (60 for each habitat × shell phenotype combination) were used in the experiments described below. By necessity, the observer (see below) was not blind to individual habitat of origin/phenotype; note that the analyst (MD) did not contribute to the actual observations.

 Behavioral tests: boldness

We studied boldness using simulated predator attacks as in Dahirel et al. (2017). All tests were done

by the same operator (V.G.) to avoid effects of inter-experimenter variability. Snails were assayed

during the last four hours of the photophase. To stimulate activity, we first placed them in a Petri

dish with water for 5 minutes, before putting them on individual clean glass plates. After snails had

moved at least one shell length (≈20 mm) from their starting position, the operator used a pipette tip

to pinch them for 5 seconds on the right side of the foot. Preliminary tests confirmed that this was

the shortest time needed to ensure all snails retracted fully in their shell. We then recorded the time

snails took to exit the shell and resume activity after the attack (from retraction to the full extension

of all tentacles out of the shell), as our measure of boldness (snails with shorter latencies being

considered bolder). We stopped observations after 20 min. Snails from the same test box were

tested on the same day, and replaced in their box after testing. To estimate the repeatability of

boldness, snails were tested a second time after seven days, using the same protocol. The initial

order in which groups were tested within a sequence was random; this order was conserved for all

subsequent tests.

Behavioral tests: exploration/speed

We studied snail movement at four temperatures within the activity range of *C. nemoralis* (Cameron,

1970): 15, 18, 22 and 25 °C. All tests were again performed by the same operator (V.G.). Movement

tests started 7 days after the last boldness test for a given individual, successive movement tests

were separated by 24h. Half of the boxes, equally distributed between landscapes of origin, were

tested in increasing temperature order (from 15 to 25 °C), the other half in decreasing order (25°C to

15°). Twenty-four hours before a given test, we placed snails and their box at the testing

temperature for habituation, using temperature-controlled cabinets (ET 619-4, Lovibond, Dortmund,

Germany). For testing, each snail was placed individually at the center of a clean 25 × 25 cm

polyethylene box (height: 9 cm) and left free to move. Snails were deemed active once they had

moved more than 2 cm away from their starting point. We used the time snails took to move more

than 10 cm from their starting point, minus the time taken to start activity, as our exploration metric,

with lower times for snails that moved away faster. We stopped observations after 20 min post-

activity initiation. This metric was chosen for its ease of implementation; we acknowledge that it

conflates exploration of the environment *stricto sensu* with movement speed (as both slow-moving

individuals and thorough explorers would have higher first-passage times).

Ethical note

This study complies with all relevant national and international laws, and the ASAB/ABS Guidelines

for the use of animals (2020) were adhered to as closely as possible. Potentially stressful

experimental treatments (boldness experiment) were limited to the shortest possible time to elicit

the behaviors of interest. No ethical board recommendation or administrative authorization was

needed to work on or sample *Cepaea nemoralis*. The marking method used is non-invasive and has

minimal to no documented effects on life-history traits (Henry & Jarne, 2007).

Statistical analyses

We analyzed snail behavioral data in a Bayesian multilevel/mixed model framework, using the Stan

language (Carpenter et al., 2017), with R (version 4.0; R Core Team, 2020) and the *brms* R package

(Bürkner, 2017) as frontends. Scripting, analysis and plotting relied on the *tidybayes*, *bayesplot*, and

patchwork packages, as well as the *tidyverse* family of packages (Gabry, Simpson, Vehtari,

Betancourt, & Gelman, 2019; Kay, 2019; Pedersen, 2019; Wickham et al., 2019).

We used a bivariate generalized linear multilevel model to estimate the effect of shell phenotype and

environment on behavior, quantify behavioral (co)variances and partition them across hierarchical

levels (inter-box and inter-individual variation) (Dingemanse & Dochtermann, 2013; Houslay &

Wilson, 2017). We did not estimate within-individual trait correlations, as exploration and boldness

were tested independently at the within-individual level (that is, boldness measure 1 had no stronger

"link" to speed measure 1 than boldness measure 2; scenario 4 of table 2 in Dingemanse &

Dochtermann, 2013). We present a full write-up of the model as Supplementary Material S2; a verbal

description follows below.

Boldness and exploration were analyzed assuming a lognormal distribution to account for the

skewed distribution of time to event data. We accounted for the fact that monitoring was stopped

before some individuals could express the behavior of interest by including a censored data indicator

in the model. Fixed effects for both behaviors included shell banding (three-level categorical

variable), landscape of origin (binary variable), and their interaction, as well as test order (1 or 2 for

boldness, 1 to 4 for exploration / speed). The model for exploration additionally included a test

temperature effect as well as interactions between this effect, shell banding and landscape effects.

Categorical variables (shell banding, landscape of origin) were converted to centered dummy

variables, and numeric variables (test order, temperature) were centered, following Schielzeth

(2010)(temperature was additionally scaled to unit 1SD). This has two benefits. First, it makes main

effect coefficients directly interpretable even in the presence of interactions (Schielzeth, 2010).

Second, for categorical variables, having the intercept on an "average" rather than on one arbitrary

default category avoids the problem of putting a more precise prior on the reference category (which

would be defined by the intercept only) than on the others (which would be defined by the intercept

and one or several other coefficients)(McElreath, 2020). Random effects included box-level and

individual-level intercepts as well as, in the case of exploration, the associated slopes for

temperature. This allowed us to estimate among-box and among-individual variation in mean

behavior and thermal behavioral reaction norms as well as the box- and individual-level covariances

among them (Dingemanse & Dochtermann, 2013).

We used a Normal(μ = ln(400), σ = 0.5) prior for the fixed effects intercepts (mean log- latencies), so

that ~99% of the probability mass was within the range of latencies that could actually be observed

by design during the experiment (i.e. 0 to 1200 sec, see above), but not totally excluding larger

values, because of censoring. We set the other priors to be weakly informative and follow some

suggestions by McElreath (2020): a Normal(0,1) prior for the other fixed effects, a half-Normal(0, 1)

prior for both random effect and distributional standard deviations. For the random effects

correlation matrices, we use a LKJ(η = 3) prior, as it helps reach convergence faster than McElreath

(2020)'s η = 2 default. Note that our choice here is more skeptical of high correlations and thus

penalizes against our hypotheses of interest (there are detectable correlations).

We partitioned total phenotypic variation V_P for each behavior into the following components: V_P =

$V_F + V_I + V_B + V_D$, where V_F is the fixed effect variation, including $V_{F(state)}$ the portion of fixed-effect

variance attributable to known individual state (banding pattern, environment of origin) to the

exclusion of experimental effects (test order, temperature) (estimated following P. de Villemereuil,

Morrissey, Nakagawa, & Schielzeth, 2018); V_I the random effect inter-individual variation (including

the effect of random temperature slope, estimated following Johnson, 2014), with $V_{I(intercept)}$ the inter-

individual variation at the average test temperature ($V_I = V_{I(Intercept)}$ for boldness); V_B and $V_{B(Intercept)}$ are

the equivalent box-level variances; and V_D is the distributional, or residual, variation. As pointed by

Wilson (2018) and de Villemereuil et al. (2018), there is in most cases no one "true" repeatability

estimate just as there is no one "true" way of partitioning the phenotypic variance pie; several

estimates with differing interpretations can be presented. Therefore, both absolute variance

components and analytical choices regarding repeatabilities should be made explicit. We estimated

the following two unadjusted repeatabilities (i.e. including the entirety of V_P in the denominator;

Nakagawa & Schielzeth, 2010): within-state repeatability $R_{(within-state)} = V_{I(Intercept)}/V_P$, and what we

term total repeatability, $R_{(total)} = (V_{I(Intercept)} + V_{F(state)})/V_P$. The proportion of persistent inter-individual

variation that is attributable to individual state (banding and landscape of origin) is then denoted by

$V_{F(state)} / (V_{I(Intercept)} + V_{F(state)})$. Variance components and repeatabilities are presented on the

observed data scale (sensu Pierre de Villemereuil, Schielzeth, Nakagawa, & Morrissey, 2016).

Variance components on the latent log scale (i.e. using model coefficients) led to qualitatively and

quantitatively similar results.

We ran four chains for 12000 iterations, with the first 2000 iterations of each chain used for warmup.

We checked mixing graphically and confirmed chain convergence using the improved $\widehat{R}$ statistic by

Vehtari et al. (2020). The chains were run longer than the default number of iterations (2000), to

ensure effective sample size was satisfactory for all parameters (both bulk- and tail-effective sample

sizes sensu Vehtari et al., 2020 > 1000). All credible intervals given are highest posterior density

intervals.

Results

Exploration was influenced by shell morph (Table 1, Fig. 2). Five-banded snails were on average

slower to explore their surroundings than either unbanded or three-banded snails (Fig.2). Snails also

became slower as tests went on (Table 1). There was no clear evidence of a landscape of origin

effect. Snails explored faster with increasing temperature (Table 1, Fig. 2). There was however no

clear evidence that the slope of the temperature reaction norm varied between the three morphs, or

between snails coming from different landscapes (Table 1; credible intervals for all interactions

largely overlap 0).

Morphs also varied in average boldness (Table 1, Fig. 2). Unbanded snails were bolder than five-

banded snails, with three-banded snails presenting intermediate values. Again, there was no

evidence for landscape of origin, or landscape × morph effects.

Both exploration and boldness were repeatable at the individual level, with average repeatabilities in

the same range for both behaviors (Table 2). Including fixed effect variation due to individual state

(morph and landscape of origin) in the calculation only slightly increased repeatabilities. Indeed, the

proportion of persistent individual variation attributable to fixed effects was different from zero but

small, with over 90% of individual-level variation attributable to other, unmeasured sources (Fig. 3,

Table 2). Variation among boxes was small but non negligible, in the same range as the proportion of

variation explained by fixed effects for both behaviors. Average exploration and boldness were

positively correlated at the individual level (Table 3, Fig. 4). There was no evidence that individual-

level variation in responses to temperature was correlated with either mean exploration times or

mean boldness (Table 3). There was no evidence for box-level correlations among traits (Table 3).

Discussion

We showed that behaviors linked to boldness and exploration/activity are repeatable in *Cepaea*

nemoralis, confirming once again that personality variation is common even in invertebrates (Kralj-

Fišer & Schuett, 2014). This result has implications for the evolutionary potential of behaviors in *C.*

nemoralis, given repeatability sets the higher limit for the heritability of behaviors (Ned A.

Dochtermann, Schwab, & Sih, 2015). Additionally, we demonstrated that these traits are positively

correlated in a common syndrome, and that their expression varies depending on shell banding, a

Mendelian trait that has been the focus on much research in this species in relation to predation and

thermal selection (Ożgo, 2009). Given how behavior can shape effective thermal tolerance (Abram et

al., 2017) or vulnerability to predation (e.g. Hulthén et al., 2017), we believe these behavioral

differences must be taken in account when discussing the evolution of shell color in this model

species.

Unbanded snails were both bolder and explored faster than five-banded snails (Table 1, Fig. 2).

Three-banded snails behaved similarly to unbanded snails with respect to boldness (but were

intermediate between unbanded and five-banded snails for exploration). This shows the "effectively

unbanded" category sometimes used in *Cepaea* studies (Cain & Sheppard, 1954) has at least some

behavioral relevance (that category groups together snails with little to no banding on the side of

their shell exposed to the sun). Exploration and boldness were positively correlated both at the

among-individual (Table 2) and among-morph levels (the shyest morph was also the slowest, Fig. 2).

At the individual level, while some clutches were laid during the experiments, we were not able to

test if this behavioral syndrome was integrated into a broader pace of life syndrome *sensu* Réale et

al. (2010) by linking behavioral and life history variation. Indeed, we were unable to ascertain the

maternal and especially paternal origin of most clutches, and were not able to follow snail fecundity

or longevity over their entire life. There is however some indications in the literature that more

active/ mobile snails are faster-growing (Oosterhoff, 1977), as predicted by the pace-of-life syndrome

hypothesis. At the morph level, it appears that contrary to behaviors, shell morphs do not differ

much in terms of average fecundity or growth rate (Oosterhoff, 1977; Wolda, 1967). They however

differ markedly in their physiology, based on their response to pharmaceutical assays: the morphs

we show here to be faster/bolder also have more responsive endogeneous opioid systems (Kavaliers,

1992). Further research is needed to understand the drivers of these correlations and their

consequences (Royauté et al., 2018; Wright et al., 2019).

Five-banded snails, which have been shown to have on average more resistant shells against rodent

attacks (Rosin et al., 2013), were shyer than unbanded snails (Fig. 2). At first glance, these results go

against the phenotypic compensation hypothesis which predicts boldness should be associated with

better defenses against predators, in particular thicker shells, mitigating the costs of risk taking

behavior (Ahlgren et al., 2015; De Winter, Ramalho Martins, Trovo, & Chapman, 2016; Hulthén,

Chapman, Nilsson, Hollander, & Brönmark, 2014). However, birds, in particular thrushes (genus

Turdus) are likely more important predators than rodents (Rosin, Olborska, Surmacki, & Tryjanowski,

2011); given they are visual predators, shell visual characteristics may be more important than

structural resistance in influencing behavioral response to predation. Both frequency-dependent

predation and direct visual selection have historically been invoked as explanations for predator-

dependent morph variation in *Cepaea* (Jones et al., 1977; Ożgo, 2009), but discussions often used

human vision as a baseline. More recently, visual selection explanations have received increased

support from an experiment using models of avian vision to rigorously test how thrushes actually see

different shell morphs (Surmacki et al., 2013). In both our test sites, the boldest morph (unbanded

shell) is the least conspicuous (based on Surmacki et al., 2013), not the rarest. This supports crypsis-

based explanations over frequency-dependent selection, under the assumption that phenotypes

already more at risk of predation should be shyer given boldness increases predation risk (Hulthén et

al., 2017). We must nonetheless remember that (i) our knowledge of how thrushes actually perceive

snails is very limited (Surmacki et al., 2013), (ii) we only tested a small set of the available morphs,

which do not include the rarest background colors (pink and brown), and (iii) shell banding is a trait

under multiple selection pressures, including thermal selection (see below).

Exploration speed was temperature-dependent: as expected from an ectothermic species, snails

were on average faster at higher temperatures (Fig. 2). The temperature reaction norm of

exploration was remarkably conserved among individuals (most of the inter-individual variance V_i

was due to differences in average behavior, rather than in temperature slopes; Table 2, Fig. 3). In

addition, there was surprisingly no evidence that behavioral differences among morphs are

influenced by the thermal environment, whether we consider the previously experienced

environment (no habitat × morph interaction) or the current environment (no effect of morph

identity on thermal reaction norms) (Table 1). This is despite abundant evidence in the literature for

thermal selection on shell morphs, based on field comparisons (Kerstes et al., 2019; for this study,

see Methods; e.g. Richardson, 1974; Schilthuizen, 2013), but also experimental evidence of both

survival (Lamotte, 1959; Tilling, 1983) and fecundity (Wolda, 1967) thermal selection. In addition,

studies giving snails a choice between multiple temperatures show snail morphs do have different

thermal preferences that align with expectations based on thermal selection (Kavaliers, 1992). Some

studies suggest that snails use shade and humidity just as much (and potentially more) as

temperature as cues to adjust their behavior to microclimate (Ożgo & Kubea, 2005; Rosin et al.,

2018). Our exploration tests were short, under standardized lighting conditions and with no water,

and snails were brought back to favorable humidity soon after. It is possible longer experiments, or

experiments comparing the responses of snails from different habitats to realistic climate variation

(including shade and/or humidity) would yield different responses. Maybe more importantly, we only

tested temperatures favorable for activity, i.e. the limited part of the thermal niche closer to the

optimum. Morph differences in behavior might be stronger closer to critical minimal or maximal

temperature thresholds (Tilling, 1983). This can be investigated by using a wider range of

temperatures and expanding the reaction norm approach used here to either a character state

approach (e.g. Houslay, Earley, Young, & Wilson, 2019) or a non-linear reaction norm approach

(Arnold, Kruuk, & Nicotra, 2019); both would account for the typical non-linearity of complete

thermal performance curves (Arnold et al., 2019). It is very important to note, however, that these

results do not mean populations from landscapes differing in sun exposure are identical in behavior,

even for the range of situations we tested. Indeed, because morphs differ in their behavior, and

because morph frequencies differ among landscapes (see Methods), the average snail from a sun

exposed population may well be bolder and more active than its counterpart from a shaded

population.

In any case, the links between behaviors and morphs we observed are conserved across contexts,

despite (apparent) selection on shell morph. While this is not a definite proof by itself, we consider

this a first hint in favor of a genetic association between morphs and behaviors that cannot be easily

broken by environmental changes, in this model species. In addition to studies aiming to confirm

these behavioral traits are heritable, further research into the physiological underpinnings of

behavioral differences between morphs (building on e.g. Kavaliers, 1992) and of shell color and

pattern determination (Kerkvliet et al., 2017) should help confirm (or infirm) this putative genetic

correlation and elucidate its proximate basis.

Assuming this genetic link is confirmed, any discussion about how selection on morph may influence

the evolution of behavior (or vice versa) must be tempered by the fact that the greater part of the

repeatable inter-individual variation in behavior was not actually explained by shell morph (see Fig. 3,

Table 1). It is in a way unsurprising, as we did not expect a single discrete trait to entirely constrain

individual behavioral variation. Indeed, the expression of animal personalities can be influenced by

many unobserved drivers and state variables which should have a priori limited links to shell morph

and its own drivers (Burns et al., 2012; Petelle, Martin, & Blumstein, 2019; Sih et al., 2015; Wright et

al., 2019). Furthermore, some level of stochastic behavioral individuality is inevitable even in the

total absence of meaningful genetic and environmental variation (Bierbach, Laskowski, & Wolf,

2017). What must be noted, though, is that some of this "remaining" individual variation may still,

actually, relate to shell morph. Indeed, because of dominance within loci and especially epistatic

relationships among loci, individuals that share the same shell phenotype may actually vary greatly in

terms of underlying genotype at morph-determining loci (e.g. having one dominant allele at the

« band presence » locus leads to total band absence and masks variation at all other banding genes;

Jones et al., 1977). However, while much is known of among-morph variation in thermal tolerance,

life history, physiology (Kavaliers, 1992; Kerstes et al., 2019; Lamotte, 1959; Oosterhoff, 1977;

Richardson, 1974; Tilling, 1983; Wolda, 1967), nothing as far as we can tell is known about within-

morph, but among-genotype variation. Investigations using individuals of known genotype obtained

through repeated crosses or, as our knowledge of the actual molecular underpinnings increases,

through direct genotyping (Gonzalez, Aramendia, & Davison, 2019; Kerkvliet et al., 2017), may shed

light on this "hidden" genetic variation and whether it contributes to the persistence of morph-

related behavior differences.

Increased boldness and activity have been tied to higher probability of dispersal in many species

(Cote, Clobert, Brodin, Fogarty, & Sih, 2010), including land snails (Dahirel et al., 2017), and non-

random dispersal is now acknowledged as a potentially widespread force behind population

phenotypic divergence (Edelaar & Bolnick, 2012; Jacob, Bestion, Legrand, Clobert, & Cote, 2015).

Bolder animals are thought to trade increased success against a greater predation risk (Hulthén et al.,

2017; but see Moiron, Laskowski, & Niemelä, 2020; Smith & Blumstein, 2008); predation is generally

considered a key driver of morphological differences in *Cepaea*, and plays a key role in dispersal

across taxa (Fronhofer et al., 2018). Although active dispersal can safely be dismissed as a driver of

continental-scale differences in morph frequencies, our results point to *Cepaea* as a good model to

understand how existing behavioral differences may drive local-scale morphological differences (and

vice versa). By building on, and complementing, a decades-long history of evolutionary research, this

will help us better understand the role of behavior, and constraints on behavioral variation, in

shaping responses to rapid environmental changes (Candolin & Wong, 2012), including landscape

alteration and climate change.

References

Abram, P. K., Boivin, G., Moiroux, J., & Brodeur, J. (2017). Behavioural effects of temperature on

ectothermic animals: Unifying thermal physiology and behavioural plasticity. *Biological*

Reviews, *92*(4), 1859–1876. doi: 10.1111/brv.12312

Ahlgren, J., Chapman, B. B., Nilsson, P. A., & Brönmark, C. (2015). Individual boldness is linked to

protective shell shape in aquatic snails. *Biology Letters*, *11*(4), 20150029. doi:

10.1098/rsbl.2015.0029

Arnold, P. A., Kruuk, L. E. B., & Nicotra, A. B. (2019). How to analyse plant phenotypic plasticity in

response to a changing climate. *New Phytologist*, *222*(3), 1235–1241. doi:

10.1111/nph.15656

ASAB, & ABS. (2020). Guidelines for the treatment of animals in behavioural research and teaching.

Animal Behaviour, *159*, I–XI. doi: 10.1016/j.anbehav.2019.11.002

Bierbach, D., Laskowski, K. L., & Wolf, M. (2017). Behavioural individuality in clonal fish arises despite

near-identical rearing conditions. *Nature Communications*, *8*, 15361. doi:

10.1038/ncomms15361

Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal of*

Statistical Software, *80*(1), 1–28. doi: 10.18637/jss.v080.i01

Burns, J. G., Svetec, N., Rowe, L., Mery, F., Dolan, M. J., Boyce, W. T., & Sokolowski, M. B. (2012).

Gene–environment interplay in *Drosophila melanogaster*: Chronic food deprivation in early

life affects adult exploratory and fitness traits. *Proceedings of the National Academy of*

Sciences of the United States of America, *109*(Suppl 2), 17239–17244. doi:

10.1073/pnas.1121265109

Cain, A. J., & Sheppard, P. M. (1954). Natural selection in *Cepaea*. *Genetics*, *39*(1), 89–116.

Cameron, R. A. D. (1970). The effect of temperature on the activity of three species of helicid snail

(Mollusca⬚: Gastropoda). *Journal of Zoology*, *162*(3), 303⬚315. doi: 10.1111/j.1469-

7998.1970.tb01267.x

Candolin, U., & Wong, B. B. M. (Éds.). (2012). *Behavioural Responses to a Changing World⬚:*

Mechanisms and Consequences (1st edition). Oxford: Oxford University Press.

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., … Riddell, A. (2017).

Stan⬚: A probabilistic programming language. *Journal of Statistical Software*, *76*(1), 1⬚32.

doi: 10.18637/jss.v076.i01

Chang, H.-W. (1991). Activity and weight loss in relation to solar radiation in the polymorphic land

snail *Cepaea nemoralis. Journal of Zoology*, *225*(2), 213⬚225. doi: 10.1111/j.1469-

7998.1991.tb03812.x

Cloyed, C. S., Dell, A. I., Hayes, T., Kordas, R. L., & O'Gorman, E. J. (2019). Long-term exposure to

higher temperature increases the thermal sensitivity of grazer metabolism and movement.

Journal of Animal Ecology, *0*(0). doi: 10.1111/1365-2656.12976

Cook, L. M. (2008). Variation with habitat in *Cepaea nemoralis⬚*: The Cain & Sheppard diagram.

Journal of Molluscan Studies, *74*(3), 239⬚243. doi: 10.1093/mollus/eyn011

Cote, J., Clobert, J., Brodin, T., Fogarty, S., & Sih, A. (2010). Personality-dependent dispersal⬚:

Characterization, ontogeny and consequences for spatially structured populations.

Philosophical Transactions of the Royal Society of London B: Biological Sciences, *365*(1560),

4065⬚4076. doi: 10.1098/rstb.2010.0176

Dahirel, M., Vong, A., Ansart, A., & Madec, L. (2017). Individual boldness is life stage-dependent and

linked to dispersal in a hermaphrodite land snail. *Ecological Research*, *32*, 751⬚755. doi:

10.1007/s11284-017-1484-x

Davison, A., Jackson, H. J., Murphy, E. W., & Reader, T. (2019). Discrete or indiscrete? Redefining the

colour polymorphism of the land snail *Cepaea nemoralis. Heredity*, 1. doi: 10.1038/s41437-

019-0189-z

De Winter, G., Ramalho Martins, H., Trovo, R. A., & Chapman, B. B. (2016). Knights in shining armour

are not necessarily bold : Defensive morphology correlates negatively with boldness, but

positively with activity, in wild threespine stickleback, *Gasterosteus aculeatus. Evolutionary*

Ecology Research, *17*, 1 12.

Dingemanse, N. J., & Dochtermann, N. A. (2013). Quantifying individual variation in behaviour :

Mixed-effect modelling approaches. *Journal of Animal Ecology*, *82*(1), 39 54. doi:

10.1111/1365-2656.12013

Dingemanse, N. J., Kazem, A. J., Réale, D., & Wright, J. (2010). Behavioural reaction norms : Animal

personality meets individual plasticity. *Trends in Ecology & Evolution*, *25*(2), 81 89. doi:

10.1016/j.tree.2009.07.013

Dochtermann, N. A., & Dingemanse, N. J. (2013). Behavioral syndromes as evolutionary constraints.

Behavioral Ecology, *24*(4), 806 811. doi: 10.1093/beheco/art002

Dochtermann, Ned A., Schwab, T., & Sih, A. (2015). The contribution of additive genetic variation to

personality variation : Heritability of personality. *Proceedings of the Royal Society B:*

Biological Sciences, *282*(1798), 20142201. doi: 10.1098/rspb.2014.2201

Edelaar, P., & Bolnick, D. I. (2012). Non-random gene flow : An underappreciated force in evolution

and ecology. *Trends in Ecology & Evolution*, *27*(12), 659 665. doi:

10.1016/j.tree.2012.07.009

Fronhofer, E. A., Legrand, D., Altermatt, F., Ansart, A., Blanchet, S., Bonte, D., … Cote, J. (2018).

Bottom-up and top-down control of dispersal across major organismal groups. *Nature*

Ecology & Evolution, *2*(12), 1859 1863. doi: 10.1038/s41559-018-0686-0

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian

workflow. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, *182*(2),

389 402. doi: 10.1111/rssa.12378

GBIF Secretariat. (2020). Cepaea nemoralis *(Linnaeus, 1758)* in *GBIF Backbone Taxonomy. Checklist*

dataset accessed via GBIF.org on 2020-08-11. Consulté à l'adresse

https://doi.org/10.15468/39omei

Gonzalez, D. R., Aramendia, A. C., & Davison, A. (2019). Recombination within the *Cepaea nemoralis*

supergene is confounded by incomplete penetrance and epistasis. *Heredity*, 1. doi:

10.1038/s41437-019-0190-6

Heath, D. J. (1975). Colour, sunlight and internal temperatures in the land-snail *Cepaea nemoralis*

(L.). *Oecologia*, *19*(1), 29⊡38. doi: 10.1007/BF00377587

Henry, P.-Y., & Jarne, P. (2007). Marking hard-shelled gastropods⊡: Tag loss, impact on life-history

traits, and perspectives in biology. *Invertebrate Biology*, *126*(2), 138⊡153. doi:

10.1111/j.1744-7410.2007.00084.x

Houslay, T. M., Earley, R. L., Young, A. J., & Wilson, A. J. (2019). Habituation and individual variation in

the endocrine stress response in the Trinidadian guppy (*Poecilia reticulata*). *General and*

Comparative Endocrinology, *270*, 113⊡122. doi: 10.1016/j.ygcen.2018.10.013

Houslay, Thomas M., & Wilson, A. J. (2017). Avoiding the misuse of BLUP in behavioural ecology.

Behavioral Ecology, *28*(4), 948⊡952. doi: 10.1093/beheco/arx023

Hulthén, K., Chapman, B. B., Nilsson, P. A., Hansson, L.-A., Skov, C., Brodersen, J., ... Brönmark, C.

(2017). A predation cost to bold fish in the wild. *Scientific Reports*, *7*(1), 1239. doi:

10.1038/s41598-017-01270-w

Hulthén, K., Chapman, B. B., Nilsson, P. A., Hollander, J., & Brönmark, C. (2014). Express yourself⊡:

Bold individuals induce enhanced morphological defences. *Proceedings of the Royal Society*

B: Biological Sciences, *281*(1776), 20132703. doi: 10.1098/rspb.2013.2703

Jacob, S., Bestion, E., Legrand, D., Clobert, J., & Cote, J. (2015). Habitat matching and spatial

heterogeneity of phenotypes⊡: Implications for metapopulation and metacommunity

functioning. *Evolutionary Ecology*, *29*(6), 851⊡871. doi: 10.1007/s10682-015-9776-5

Johnson, P. C. D. (2014). Extension of Nakagawa & Schielzeth's R^2_{GLMM} to random slopes models.

Methods in Ecology and Evolution, *5*(9), 944⬚946. doi: 10.1111/2041-210X.12225

Jones, J. S. (1982). Genetic differences in individual behaviour associated with shell polymorphism in

the snail *Cepaea nemoralis*. *Nature*, *298*(5876), 749⬚750. doi: 10.1038/298749a0

Jones, J. S., Leith, B. H., & Rawlings, P. (1977). Polymorphism in *Cepaea*⬚: A Problem with Too Many

Solutions? *Annual Review of Ecology and Systematics*, *8*, 109⬚143. JSTOR. Consulté à

l'adresse JSTOR.

Kavaliers, M. (1992). Opioid systems, behavioral thermoregulation and shell polymorphism in the

land snail, *Cepaea nemoralis*. *Journal of Comparative Physiology B*, *162*(2), 172⬚178. doi:

10.1007/BF00398344

Kay, M. (2019). *tidybayes⬚: Tidy Data and Geoms for Bayesian Models*. doi: 10.5281/zenodo.1308151

Kerkvliet, J., Boer, T. de, Schilthuizen, M., & Kraaijeveld, K. (2017). Candidate genes for shell colour

polymorphism in *Cepaea nemoralis*. *PeerJ*, *5*, e3715. doi: 10.7717/peerj.3715

Kern, E. M. A., Robinson, D., Gass, E., Godwin, J., & Langerhans, R. B. (2016). Correlated evolution of

personality, morphology and performance. *Animal Behaviour*, *117*, 79⬚86. doi:

10.1016/j.anbehav.2016.04.007

Kerstes, N. A. G., Breeschoten, T., Kalkman, V. J., & Schilthuizen, M. (2019). Snail shell colour

evolution in urban heat islands detected via citizen science. *Communications Biology*, *2*(1),

264. doi: 10.1038/s42003-019-0511-6

Kralj-Fišer, S., & Schuett, W. (2014). Studying personality variation in invertebrates⬚: Why bother?

Animal Behaviour, *91*, 41⬚52. doi: 10.1016/j.anbehav.2014.02.016

Lamotte, M. (1959). Polymorphism of natural populations of *Cepaea nemoralis*. *Cold Spring Harbor*

Symposia on Quantitative Biology, *24*(0), 65⬚86. doi: 10.1101/SQB.1959.024.01.009

McElreath, R. (2020). *Statistical Rethinking⬚: A Bayesian Course with Examples in R and Stan* (2nd

edition). Boca Raton: Chapman and Hall/CRC.

Moiron, M., Laskowski, K. L., & Niemelä, P. T. (2020). Individual differences in behaviour explain

variation in survival : A meta-analysis. *Ecology Letters*, *23*(2), 399 408. doi:

10.1111/ele.13438

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data : A practical

guide for biologists. *Biological Reviews*, *85*(4), 935 956. doi: 10.1111/j.1469-

185X.2010.00141.x

Oosterhoff, L. M. (1977). Variation in growth rate as an ecological factor in the landsnail *Cepaea*

nemoralis (L.). *Netherlands Journal of Zoology*, *27*(1), 1 132. doi: 10.1163/002829677X00072

Ożgo, M. (2009). Current problems in the research of *Cepaea* polymorphism. *Folia Malacologica*,

16(2), 55 60. doi: 10.12657/folmal.016.009

Ozgo, M., & Kinnison, M. T. (2008). Contingency and determinism during convergent contemporary

evolution in the polymorphic land snail, *Cepaea nemoralis*. *Evolutionary Ecology Research*,

10, 721 733.

Ożgo, M., & Kubea, A. (2005). Humidity and the effect of shell colour on activity of *Cepaea nemoralis*

(Linnaeus, 1758). *Folia Malacologica*, *13*(3).

Ożgo, M., Liew, T.-S., Webster, N. B., & Schilthuizen, M. (2017). Inferring microevolution from

museum collections and resampling : Lessons learned from *Cepaea*. *PeerJ*, *5*, e3938. doi:

10.7717/peerj.3938

Ożgo, M., & Schilthuizen, M. (2012). Evolutionary change in *Cepaea nemoralis* shell colour over 43

years. *Global Change Biology*, *18*(1), 74 81. doi: 10.1111/j.1365-2486.2011.02514.x

Pedersen, T. L. (2019). *patchwork : The Composer of Plots*. Consulté à l'adresse https://CRAN.R-

project.org/package=patchwork

Petelle, M. B., Martin, J. G. A., & Blumstein, D. T. (2019). Mixed support for state maintaining risky

personality traits in yellow-bellied marmots. *Animal Behaviour*, *150*, 177 188. doi:

10.1016/j.anbehav.2019.02.008

R Core Team. (2020). R: a language and environment for statistical computing (Version 4.0.0). Vienna,

Austria: R Foundation for Statistical Computing. Consulté à l'adresse https://www.R-

project.org/

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P.-O. (2010).

Personality and the emergence of the pace-of-life syndrome concept at the population level.

Philosophical Transactions of the Royal Society of London B: Biological Sciences, *365*(1560),

4051�4063. doi: 10.1098/rstb.2010.0208

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating animal

temperament within ecology and evolution. *Biological Reviews*, *82*(2), 291�318. doi:

10.1111/j.1469-185X.2007.00010.x

Richards, P. M., Liu, M. M., Lowe, N., Davey, J. W., Blaxter, M. L., & Davison, A. (2013). RAD-Seq

derived markers flank the shell colour and banding loci of the *Cepaea nemoralis* supergene.

Molecular Ecology, *22*(11), 3077�3089. doi: 10.1111/mec.12262

Richardson, A. M. M. (1974). Differential climatic selection in natural population of land snail *Cepaea*

nemoralis. *Nature*, *247*(5442), 572. doi: 10.1038/247572a0

Rosin, Z. M., Kobak, J., Lesicki, A., & Tryjanowski, P. (2013). Differential shell strength of *Cepaea*

nemoralis colour morphs—Implications for their anti-predator defence.

Naturwissenschaften, *100*(9), 843�851. doi: 10.1007/s00114-013-1084-8

Rosin, Z. M., Kwieciński, Z., Lesicki, A., Skórka, P., Kobak, J., Szymańska, A., … Tryjanowski, P. (2018).

Shell colour, temperature, (micro)habitat structure and predator pressure affect the

behaviour of *Cepaea nemoralis*. *The Science of Nature*, *105*(5), 35. doi: 10.1007/s00114-018-

1560-2

Rosin, Z. M., Olborska, P., Surmacki, A., & Tryjanowski, P. (2011). Differences in predatory pressure

on terrestrial snails by birds and mammals. *Journal of Biosciences*, *36*(4), 691�699. doi:

10.1007/s12038-011-9077-2

Royauté, R., Berdal, M. A., Garrison, C. R., & Dochtermann, N. A. (2018). Paceless life? A meta-

analysis of the pace-of-life syndrome hypothesis. *Behavioral Ecology and Sociobiology*, *72*(3),

64. doi: 10.1007/s00265-018-2472-z

Santostefano, F., Wilson, A. J., Niemelä, P. T., & Dingemanse, N. J. (2017). Behavioural mediators of

genetic life-history trade-offs: A test of the pace-of-life syndrome hypothesis in field

crickets. *Proc. R. Soc. B*, *284*(1864), 20171567. doi: 10.1098/rspb.2017.1567

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.

Methods in Ecology and Evolution, *1*(2), 103–113. doi: 10.1111/j.2041-210X.2010.00012.x

Schilthuizen, M. (2013). Rapid, habitat-related evolution of land snail colour morphs on reclaimed

land. *Heredity*, *110*(3), 247–252.

Sih, A., Bell, A. M., Johnson, J. C., & Ziemba, R. E. (2004). Behavioral Syndromes: An Integrative

Overview. *The Quarterly Review of Biology*, *79*(3), 241–277. doi: 10.1086/422893

Sih, A., Mathot, K. J., Moirón, M., Montiglio, P.-O., Wolf, M., & Dingemanse, N. J. (2015). Animal

personality and state–behaviour feedbacks: A review and guide for empiricists. *Trends in*

Ecology & Evolution, *30*(1), 50–60. doi: 10.1016/j.tree.2014.11.004

Silvertown, J., Cook, L., Cameron, R., Dodd, M., McConway, K., Worthington, J., … Juan, X. (2011).

Citizen science reveals unexpected continental-scale evolutionary change in a model

organism. *PLOS ONE*, *6*(4), e18927. doi: 10.1371/journal.pone.0018927

Smith, B. R., & Blumstein, D. T. (2008). Fitness consequences of personality: A meta-analysis.

Behavioral Ecology, *19*(2), 448–455. doi: 10.1093/beheco/arm144

Surmacki, A., Ożarowska-Nowicka, A., & Rosin, Z. M. (2013). Color polymorphism in a land snail

Cepaea nemoralis (Pulmonata: Helicidae) as viewed by potential avian predators. *Die*

Naturwissenschaften, *100*(6), 533–540. doi: 10.1007/s00114-013-1049-y

Tilling, S. M. (1983). An experimental investigation of the behaviour and mortality of artificial and

natural morphs of *Cepaea nemoralis* (L.). *Biological Journal of the Linnean Society*, *19*(1),

35–50. doi: 10.1111/j.1095-8312.1983.tb00775.x

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., & Bürkner, P.-C. (2020). Rank-normalization,

folding, and localization: An improved \widehat{R} for assessing convergence of MCMC.

Bayesian Analysis. doi: 10.1214/20-BA1221

Villemereuil, P. de, Morrissey, M. B., Nakagawa, S., & Schielzeth, H. (2018). Fixed-effect variance and

the estimation of repeatabilities and heritabilities: Issues and solutions. *Journal of*

Evolutionary Biology, *31*(4), 621–632. doi: 10.1111/jeb.13232

Villemereuil, Pierre de, Schielzeth, H., Nakagawa, S., & Morrissey, M. (2016). General methods for

evolutionary quantitative genetic inference from Generalized Mixed Models. *Genetics*,

204(3), 1281–1294. doi: 10.1534/genetics.115.186536

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., … Yutani, H. (2019).

Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. doi:

10.21105/joss.01686

Williams, S. T. (2017). Molluscan shell colour: Molluscan shell colour. *Biological Reviews*, *92*(2),

1039–1058. doi: 10.1111/brv.12268

Wilson, A. J. (2018). How should we interpret estimates of individual repeatability? *Evolution Letters*,

2(1), 4–8. doi: 10.1002/evl3.40

Wolda, H. (1967). The effect of temperature on reproducton in some morphs of the landsnail *Cepaea*

nemoralis (l.). *Evolution*, *21*(1), 117–129. doi: 10.1111/j.1558-5646.1967.tb00135.x

Wolf, M., & McNamara, J. M. (2012). On the Evolution of Personalities via Frequency-Dependent

Selection. *The American Naturalist*, *179*(6), 679–692. doi: 10.1086/665656

Wolf, M., & Weissing, F. J. (2010). An explanatory framework for adaptive personality differences.

Philosophical Transactions of the Royal Society B: Biological Sciences, *365*(1560), 3959–3968.

doi: 10.1098/rstb.2010.0215

Wright, J., Bolstad, G. H., Araya-Ajoy, Y. G., & Dingemanse, N. J. (2019). Life-history evolution under

fluctuating density-dependent selection and the adaptive alignment of pace-of-life

syndromes. *Biological Reviews*, *94*(1), 230–247. doi: 10.1111/brv.12451

Tables

Table 1: Estimated fixed effect parameters of the model explaining exploration and boldness

latencies (mean and 95% credible intervals). Continuous explanatory variables are centered and

scaled, and categorical variables converted to centered dummy variables; the intercept then refers to

the behavior of a hypothetical "average" snail.

	Mean effect (95% credible interval)	
	Exploration	Boldness
Fixed effects: main effects		
Intercept (log-latency of an "average" individual)	6.670 (6.630 ; 6.700)	2.980 (2.830 ; 3.140)
banding = 3 bands	-0.002 (-0.064 ; 0.057)	0.149 (-0.054 ; 0.358)
banding = 5 bands	**0.066 (0.004 ; 0.125)**	**0.260 (0.059 ; 0.472)**
landscape of origin = shaded	0.056 (-0.017 ; 0.128)	0.196 (-0.108 ; 0.483)
test temperature	**-0.112 (-0.150 ; -0.072)**	--
test order	**0.036 (0.006 ; 0.067)**	0.079 (-0.033 ; 0.198)
Fixed effects: interactions		
banding = 3bands × landscape = shaded	-0.071 (-0.191 ; 0.048)	0.157 (-0.247 ; 0.561)
banding = 5bands × landscape = shaded	-0.008 (-0.130 ; 0.114)	0.239 (-0.164 ; 0.644)
banding = 3bands × temperature	0.002 (-0.039 ; 0.042)	--
banding = 5bands × temperature	0.013 (-0.028 ; 0.053)	--
landscape = shaded × temperature	-0.053 (-0.130 ; 0.023)	--
banding = 3bands × landscape = shaded × temperature	-0.023 (-0.107 ; 0.056)	--
banding = 5bands × landscape = shaded × temperature	0.020 (-0.063 ; 0.101)	--

655 **Table 2.** Variance partitioning and repeatabilities. Estimated variances and 95% credible interval are

656 obtained from a multivariate mixed model. Variances are estimated on the observed scale *sensu* de

657 Villemereuil et al (2016). Variances are rounded to the nearest unit. Readers looking at the data may

658 note that the total variances V_P are higher than the empirically observed variances. This is because

659 the latter are underestimated, due to censoring.

	Posterior mean (95% credible interval)	
	Exploration	Boldness
Variance partitioning (observed scale, in sec²)		
Fixed effects V_F	13596 (6340 ; 21088)	49 (6 ; 108)
- explained by state	2179 (337 ; 4332)	46 (4 ; 103)
Box-level V_B	8439 (2782 ; 15010)	128 (12; 310)
- random intercept only	3085 (0 ; 7028)	128 (12; 310)
Individual V_I	25748 (18111 ; 33850)	542 (260; 867)
- random intercept only	24400 (17412 ; 32270)	542 (260; 867)
Residual V_D	73943 (63481 ; 84587)	1795 (970 ; 2851)
Total variance V_P	121726 (102528 ; 141858)	2514 (1356 ; 4010)
Repeatabilities (observed scale)		
$R_{(within\text{-}state)}$	0.200 (0.150 ; 0.253)	0.218 (0.147 ; 0.290)
$R_{(total)}$	0.218 (0.165 ; 0.271)	0.236 (0.166 ; 0.307)
$R_{(within\text{-}state)}$ / $R_{(total)}$	0.918 (0.842 ; 0.984)	0.923 (0.841 ; 0.989)

662 **Table 3.** Random effect correlation matrices (latent scale, mean and 95% credible intervals). Box-

663 level correlations are above the diagonal, individual-level correlations below the diagonal.

	Boldness, intercept	Exploration, intercept	Exploration, temperature slope
Boldness, intercept		0.119 (-0.402 ; 0.653)	0.384 (-0.059 ; 0.780)
Exploration, intercept	**0.279 (0.096 ; 0.457)**		-0.184 (-0.688 ; 0.326)
Exploration, temperature slope	-0.202 (-0.681 ; 0.312)	0.286 (-0.207 ; 0.785)	

Figure legends

Figure 1. Study species and sites. (A) An unbanded yellow *Cepaea nemoralis*, showing the position of

individual paint marks on the shell (B) Representative shells of the three studied morphs as seen

from above: yellow unbanded, three-banded, and five-banded snails (C) Study sites location in

France; the open habitat (D) and the shaded habitat (E) are separated by about 2 km. Photographs

were taken during winter, when snails were collected.

Figure 2. (A;B) posterior distributions of median boldness (A) and exploration (B) in relation to shell

morph. The model estimates the mean log-latencies, which correspond to the medians on the

observed latencies scale. (C) Mean and 95% credible band for the relationship between exploration

latency and test temperature. Grey lines connect trials from the same individual. Values are plotted

on a log-transformed axis. N = 360 individuals (60 per landscape × morph combination)

Figure 3. Mean (points) and posteriors for the proportion of variance explained by the different

variance components. See Table 2 or Methods for the names of the variance components.

Figure 4. Correlation between individual-level random effects for boldness and exploration,

illustrated by plotting their respective Best Linear Unbiased Predictors and 95% credible intervals.

Inset: posterior distribution of the correlation coefficient. BLUPs are plotted, and correlation

estimated, on the latent log scale.