1   FULL PAPER

3   Antagonistic plants preferentially target arbuscular mycorrhizal fungi that are highly

4   connected to mutualistic plants

6   Sofia IF Gomes[1], Miguel A Fortuna[2], Jordi Bascompte[2], Vincent SFT Merckx[3,4*]

[1] Institute of Biology, Leiden University, Leiden, the Netherlands

[2] Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Switzerland.

[4] Naturalis Biodiversity Center, Leiden, the Netherlands

[3] Department of Evolutionary and Population Biology, Institute for Biodiversity and Ecosystem

Dynamics, University of Amsterdam, Amsterdam, the Netherlands

* corresponding author: vincent.merckx@naturalis.nl

Summary

•   How antagonists – mycoheterotrophic plants that obtain carbon and soil nutrients from

fungi – are integrated in the usually mutualistic arbuscular mycorrhizal networks is

unknown. Here, we compare mutualistic and antagonistic plant associations with

arbuscular mycorrhizal fungi and use network analysis to investigate fungal association

preferences in the tripartite network.

•   We sequenced root tips from mutualistic and antagonistic plants in a tropical forest to

assemble the combined tripartite network between mutualistic plants, mycorrhizal fungi,

and antagonistic plants. We compared the fungal ecological similarity between

mutualistic and antagonist networks, and searched for modules (an antagonistic and a

mutualistic plant interacting with the same pair of fungi) to investigate whether pairs of

fungi simultaneously linked to plant species from each interaction type were

overrepresented throughout the network.

•   Antagonistic plants interacted with approximately half the fungi detected in mutualistic

plants. Antagonists were indirectly linked to any of the detected mutualistic plants, and

fungal pairwise ecological distances were correlated in both network types. Moreover,

pairs of fungi sharing the same antagonistic and mutualistic plant species occurred more

often than expected by chance.

•   We hypothesize that the maintenance of antagonistic interactions is maximized by

targeting well-linked mutualistic fungi, thereby minimizing the risk of carbon supply

shortages.

Keywords: mycoheterotrophy, mycorrhizal networks, arbuscular mycorrhizal fungi,

antagonism, mutualism

Introduction

Since the early history of life, interspecific mutualisms have been paramount in the functioning of ecosystems (Thompson, 2005; Bascompte & Jordano, 2013). Mutualisms can form complex networks of interdependence between dozens or even hundreds of species. A prime example of this 'web of life' is the 450-million-year-old mutualism between the majority of land plants and arbuscular mycorrhizal (AM) fungi (Strullu-Derrien *et al.*, 2018). In this interaction plants supply the arbuscular mycorrhizal Glomeromycotina and Mucoromycotina fungi with carbohydrates, essential for fungal survival and growth. In return, the fungi provide their host plants with mineral nutrients and water from the soil (Smith & Read, 2008; Bidartondo *et al.*, 2011). One of the key characteristics of the arbuscular mycorrhizal interaction is its low interaction specificity: a mycorrhizal plant typically associates simultaneously with multiple fungi and a mycorrhizal fungus often associates simultaneously with multiple plants (Lee *et al.*, 2013). This creates complex underground networks in which plants of different species are indirectly linked through shared arbuscular mycorrhizal fungi (Toju *et al.*, 2015; Chen *et al.*, 2017). Despite this low specificity, there is evidence that networks of plants and arbuscular mycorrhizal fungi do not assemble randomly, and effects of plant functional group (Davison et al. 2011; Sepp *et al.*, 2019), and plant or fungal evolutionary relationships (Montesinos-Navarro *et al.*, 2015; Chen *et al.*, 2017) on arbuscular mycorrhizal interaction networks have been reported.

Considerable progress has been made in dissecting the exchange of resources between plants and AM fungi, and its regulation, in recent years. Experimental work on low-diversity systems has demonstrated that control is bidirectional, and partners offering the best rate of exchange are rewarded, suggesting that AM networks consist of 'fair trade' interactions (Bever *et al.*, 2009; Kiers *et al.*, 2011). However, the importance of reciprocally regulated resource exchange is questioned, as mycorrhizas also affect plant health, interactions with other soil organisms, host-defense reactions, and suppression of non-mycorrhizal competitor plants (Walder & Van Der Heijden, 2015). Also, strictly reciprocal regulation of resource exchange does not seem to apply to all AM interactions. Some exceptional plants, for example, behave as antagonists (Selosse & Rousset, 2011; Walder & Van Der Heijden, 2015): mycoheterotrophic plants can obtain carbon from root-associated fungi and some species

have replaced photosynthesis by carbon uptake from arbuscular mycorrhizal fungi (Leake,

1994; Merckx, 2013). The mechanism underpinning carbon transfer from arbuscular

mycorrhizal fungi to mycoheterotrophic plants remains unclear, but it is unlikely that resource

exchange in these mycoheterotrophic associations relies on reciprocal dynamics. Hence,

mycoheterotrophic plants are considered cheaters of the mycorrhizal symbiosis because they

exploit the arbuscular mycorrhizal symbiosis for soil nutrients and carbon without

reciprocating, to our current knowledge (Selosse & Rousset, 2011), or without apparently

being sanctioned by the fungal partners (Walder & Van Der Heijden, 2015). Moreover, it has

been suggested that mycoheterotrophic plants may display a truly biotrophic parasitic mode,

digesting the fungus colonizing their roots (Imhof *et al.*, 2013).

Phylogenetic studies demonstrate that these cheaters evolved from mutualistic ancestors

(Merckx *et al.*, 2013). Within obligate mutualisms, the critical barrier to mutualism breakdown

and to the evolutionary stability of the resulting cheater species is thought to be a

requirement for three-species coexistence: a cheater plant relies on a mutualistic partner – a

mycorrhizal fungus – which simultaneously interacts with a mutualistic plant (Pellmyr &

Leebens-Mack, 1999). In species-rich mutualisms such as the arbuscular mycorrhizal

symbiosis, where multi-species coexistence is the rule, a high potential for the occurrence of

these tripartite linkages is expected (Merckx & Bidartondo, 2008). Indeed, while evolution of

cheating in specialized obligate mutualisms is relatively rare (Sachs & Simms, 2006), cheating

in the arbuscular mycorrhizal symbiosis evolved in more than a dozen of plant clades,

including over 250 species which together occur in nearly all tropical and subtropical forests

(Merckx, 2013; Gomes *et al.*, 2019a).

Previous work has shown that mycoheterotrophic plants – hereafter 'antagonists' – target a

subset of the mycorrhizal fungi available in the local community (e.g Bidartondo *et al.* (2002);

Gomes *et al.* (2017a); Sheldrake *et al.* (2017)). Their associated fungal communities can vary

in specificity (Merckx *et al.*, 2012), and in communities of co-occurring antagonists the

phylogenetic diversity of their fungal associations appears to increase proportionally to their

fungal overlap, a pattern which potentially responds to an ecological mechanism driven by

maximizing co-occurrence and avoiding competitive exclusion among antagonistic plants.

(Gomes *et al.*, 2017b). However, whether partner choice of these antagonists is affected by mutualistic mycorrhizal interactions is currently unknown. Here, we hypothesize that antagonists preferentially associate with 'keystone' (Mills & Doak, 1993) fungi that are connected to many different mutualistic plants, since these fungi are potentially more resilient to perturbations (Bascompte & Jordano, 2007) and may be the most reliable source of carbon (Waterman *et al.*, 2013). In addition, since fungal traits play an important role in arbuscular mycorrhizal interactions – phylogenetically related AM fungi (assumed to have similar functional traits), preferentially interact with similar plant species (Chagnon *et al.*, 2015) – we hypothesize that if partner selection in tripartite networks is trait-driven we will be able to detect an influence of the phylogenetic relationships of the fungi. Here, we test these hypotheses on a combined tripartite mycorrhizal network of co-occurring antagonistic and surrounding mutualistic plants linked by shared arbuscular mycorrhizal fungi compiled by high-throughput DNA sequencing.

Materials and methods

Sampling

Since mycoheterotrophic plants are relatively rare and often have patchy distributions (Gomes *et al.*, 2019b), we sampled two 4 x 4 m subplots a few meters apart in a coastal low land plain rainforest in French Guiana (5°28'25"N 53°34'51"W) on 28 July 2014, with overlapping antagonistic species. In both subplots, roots of antagonistic plants and surrounding mutualistic plants were sampled, cleaned with water, and stored on CTAB buffer at -20°C until further processing. We found in total five antagonistic plant species: *Dictyostega orobanchoides, Gymnosiphon breviflorus* (Burmanniaceae)*, Voyria aphylla, Voyriella parviflora* (Gentianaceae)*,* and *Soridium spruceanum* (Triuridaceae)*.* Complete individuals of antagonistic plants were dug out, and around each, 3 root tips of mutualistic plants were collected. In addition, we sampled five additional mutualistic plant root tips randomly from each quadrant of each plot aiming to better represent the local belowground mutualistic community. A detailed taxon list of antagonistic and mutualistic plant species collected can be found in Supporting Information, Table **S1**.

Plant identification and fungal communities sequencing

DNA was extracted from the CTAB-preserved roots with the NucleoMag96 Plant Kit (Macherey-Nagel Gmbh & Co., Düren, Germany), using the KingFisher Flex Magnetic Particle Processor (Thermo Fisher Scientific, Waltham, MA, USA). Mutualistic plant species were identified by sequencing the markers *matK or trnL* as described in (Gomes *et al.*, 2017a). Plant identification to species, genus, or family level based on BLAST against GenBank was reviewed by consulting the checklist of plants in French Guiana (Berry & Weitzman, 2007). Fungal communities associated with the roots of mutualistic and antagonistic plants were amplified using the primers fITS7 (Ihrmark *et al.*, 2012) and ITS4 (White *et al.*, 1990) and sequenced with a Personal Genome Machine (Ion Torrent; Life Technologies, Guildford, CT, USA), as described in Gomes *et al.* (2017b) in two separate runs. Negative controls from the extraction and the PCR reactions were included (and had zero reads). The same bioinformatics methods from Gomes et al. (2017b) were used to process the raw reads from the two runs combined until clustering into 97% Operational Taxonomic Units (OTUs), using USearch v.7.0 (Edgar, 2010). OTUs represented by less than six reads in each sample were excluded to avoid spurious OTUs (Lindahl *et al.*, 2013). The taxonomical assignment of each OTU was done by querying against the UNITE database (Kõljalg *et al.*, 2013). Because the antagonistic plants in this study are presently only known to associate with fungi that belong to the sub-phylum Glomeromycotina (Merckx *et al.*, 2012), we only retained fungal OTUs from this sub-phylum in the subsequent analysis. In these analyses, we accounted for the phylogenetic relatedness between fungal OTUs, due to the uncertainty of correspondence of individual OTUs with taxon diversity of arbuscular mycorrhizal fungi (Flynn *et al.*, 2015), despite the fact that delimitation of OTUs is not likely to interfere with ecological interpretations (Lekberg *et al.*, 2014). We inferred the phylogenetic relationships between the fungal OTUs following the strategy of Gomes *et al.* (2017b). Briefly, we aligned the OTUs with partial reference sequences and performed a phylogenetic inference in which the relationships between these references were enforced based on Krüger *et al.*, (2012). Hereafter, we refer to OTUs as 'fungi'. The fungal communities obtained from the root samples are not necessarily similar representations of the total communities of antagonists and mutualists because the sampling coverage of their root systems varied: antagonistic plants are small herbs, and thus large parts of their root system were collected and extracted, while for mutualistic plants, mostly forest

trees (see Results), root samples represents only a small fragment of their root system and
thus the detected fungal communities associated with mutualistic plants are likely a subset
of their total fungal communities.

Plant-fungal interactions

Fungal communities at the plant individual level were significantly structured by plant species
identity with a minor influence of the subplot where they were collected (see Supporting
Information, Methods **S1**, Figure **S1**). To contrast the interactions of antagonistic and
mutualistic plants, we considered their plant-fungal interactions as a single tripartite network
(Fig **1a**) by combining the fungal communities associated with individual plants from the two
subplots into overall communities per plant species. By combining the data from the two
subplots, we aimed to reconstruct a more robust picture of the interactions in this highly
diverse rainforest while compensating for the relatively low sampling intensity (for 77 out of
220 root tips of mutualistic plants both plant identification and AM fungal community data
with sufficient reads could be obtained). Plant species for which Glomeromycotina fungi were
represented by less than 500 reads were excluded, resulting in removing 11 mutualistic plant
species (see details in Supporting Information, Table **S1**). Moreover, because rarefying the
OTU matrix has been shown to greatly increase the false positive rate of OTUs per sample
(McMurdie & Holmes, 2014), we performed the subsequent analyses on 100 matrices
rarefied to 844 reads per plant species, based on lowest number of reads obtained for all
plant species in the observed dataset. Results are presented with mean and standard
deviation values obtained from running the analyses on the rarefied matrices. To investigate
fungal association patterns of antagonistic and mutualistic plants, we used incidence (binary)
data, because our main interest was to determine which interactions can be established and
not how abundant they are. Also, due to the difference in root sampling (see above) DNA read
abundance data unlikely correspond to fungal abundances at the plant species level, in
particular between plant types (antagonist and mutualists). We considered the simultaneous
presence of a fungus in the roots of an antagonistic and a mutualistic plant as a potential link
between both (Southworth *et al.*, 2005).

We tested for the effect of plant and fungal phylogenetic relatedness on the observed
interactions in both networks. We used the fungal phylogeny described above, and the

phylogenetic relationships of the plants as derived from *TimeTree* (Kumar *et al.*, 2017) (see

details in Supporting Information, Methods **S2**). We computed Mantel test correlations

between the phylogenetic distance matrix and the community dissimilarity matrix in each

instance for antagonistic and mutualistic species individually. The phylogenetic distance

matrices were extracted from the phylogenetic trees of plants and fungi, and the community

dissimilarity matrices were calculated as the Jaccard distance on the binary interaction

matrices. Phylogenetic signal was calculated for the 100 rarefied matrices, and consistency of

significant results was assessed across multiple rarefaction depths (Supporting Information,

Fig. **S2**).

Mutualistic and antagonistic plant-plant interactions

To compare the range of fungal interactions between mutualistic and antagonistic plants, we

calculated their normalized degree and the phylogenetic species variability of their associated

fungal communities. The normalized degree of a plant species is the proportion of its

associated fungi out of the total possible fungi in the network (Martín González *et al.*, 2010),

and was calculated with the *ND* function of the *bipartite* R package (Dormann *et al.*, 2018).

The phylogenetic species variability (*psv*) of the fungal community associated with a plant

species summarizes the level to which the fungi in this community are phylogenetically

related (Helmus *et al.*, 2007a). When a community consists of unrelated fungi, the index

equals 1, indicating maximum phylogenetic variability. As relatedness increases, the index

approaches 0, indicating high phylogenetic specificity (Helmus *et al.*, 2007b). To explore how

the number of interactions relates to the phylogenetic relationships among the detected

fungi, we tested for a linear correlation between normalized degree and *psv* by calculating

the Pearson's correlation coefficient. In addition, we calculated the fungal overlap between

each pair of plant species as the number of fungi they share to infer plant–plant preferences

(Fig **1b**).

Mutualistic and antagonistic fungal interactions

To investigate whether fungi have similar number of interactions with plants in both

mutualistic and antagonistic networks, we calculated the Pearson correlation between the

normalised degrees of the shared fungi within each interaction type. We compared the

ecological similarity of the fungi shared between the two interaction type networks (i.e.

mutualistic and antagonistic interactions). The ecological similarity of a pair of fungi

represents their similarity in interactions through shared plants. We calculated similarity

matrices between the fungi linked to mutualistic and antagonistic plants separately, using two

different measures. The first measure was the Jaccard index, which corresponds to the

number of plant species with which both fungi interact divided by the total number of plant

species with which they interact, and the second was the Overlap measure: $C_{ij}/\min(d_i, d_j)$,

where C_{ij} is the number of shared plants between fungus i and j, and $\min(d_i, d_j)$ is the smallest

number of associated plants between fungus i and j (Saavedra *et al.*, 2013). The overlap

measure corresponds to the number of shared plant species relative to the maximum number

of plant species that can be shared, and takes into account the possibility that fungi have

differential limits in their maximum number of plant partners. We computed the Mantel test

correlation between the similarity matrices for the mutualistic and antagonistic interactions

using the two different measures. We computed the partial Mantel test correlation between

the similarity matrices calculated using the Jaccard and Overlap measures, controlling for the

phylogenetic relatedness of the shared fungi.

Network analysis

We searched for diamond-shape modules in the tripartite network (reduced to only include

fungi shared between both interaction types) representing a mutualistic and an antagonistic

plant interacting with the same pair of fungi (Fig. **2**). If the module is overrepresented

throughout the entire network, it can be considered a motif (Milo *et al.*, 2002; Bascompte &

Melián, 2005; Stouffer *et al.*, 2007). This specific motif does not consider the identity of the

partners *per se*, but searches for this particular topological structure. To assess whether the

diamond-shaped module was overrepresented relative to random expectations, we

randomized the original antagonistic plants – mycorrhizal fungi – mutualistic plants

community matrix and calculated the frequency of this diamond-shaped module in 1000

generated random networks. We used a null model that draws an interaction between a plant

and a fungal species which is proportional to the generalization level of both

species. Specifically, this probability is defined as the arithmetic mean of the fraction of

interactions of the plant and that of the fungi (null model 2 in (Fortuna & Bascompte, 2006)),

initially proposed in (Olesen *et al.*, 2003). Because there was an imbalanced number of plant

species (five antagonistic and 21 mutualistic plants), each randomized matrix resulted from

randomizing the antagonistic and mutualistic interactions separately, and then combined into

a single matrix before module search. To exclude the effects of sampling effort both in the

number of individuals collected (leading to roughly half of fungal reads belonged to the five

antagonistic species while the other half represented 21 mutualistic species) and in

representation of whole (antagonists) or partial (autotrophic) root systems, we repeated this

procedure in a set of 100 rarefied matrices where each set was resampled to multiple depths.

To evaluate whether the diamond-shaped module was overrepresented throughout the

empirical network, we calculated the standard z-score = (obs − $\overline{rand}$) / sd($\overline{rand}$), where *obs*

is the observed network modules and $\overline{rand}$ is the mean of the same type of modules across

the 1000 random networks, using a 95% confidence interval. An empiric result above the null

confidence interval indicates that pairs of fungi are shared between particular antagonistic

and mutualistic plants more often than expected by chance, reflecting a preference of

antagonistic plants to link to pairs of fungi that are generalists in their interactions with

mutualistic plants. An empirical result below the null confidence interval indicates that

antagonistic plants target pairs of fungi that are specialists in their interactions with

mutualistic plants.

Results

Plant-fungal interactions

We successfully extracted DNA from root tips of 32 mutualistic (123 root tips) and five

antagonistic (60 individuals) plant species (see Table **S1** for detailed species list). After

retaining samples that contained a minimum of 500 reads identified as Glomeromycotina, we

reduced our dataset to 21 mutualistic (77 root tips) and five antagonistic (27 individuals) plant

species, with a total of 365,135 reads. Mutualistic plants belonged to 24 families, including 16

trees, two shrubs, one herb, one vine, and one liana. We obtained 115 arbuscular mycorrhizal

fungi, identified as Glomeromycotina within three families Gigasporaceae (4 fungi),

Acaulosporaceae (14) and Glomeraceae (97). Of these, 96 OTUs (49% of total reads) were

present in the antagonistic network.

The 100 rarefied binary matrices included 26 plant species (21 mutualists and 5 antagonists)

and 110 ± 1.47 fungi. Of these, mutualistic networks had 100 ± 1.51 fungi and antagonistic

networks had 61 ± 3.14 fungi. In all the generated matrices, mutualistic and antagonistic

plants shared 51 ± 3.16 fungi, which represents 55% and 92% of total fungi present in

mutualistic and antagonistic plants, respectively (Fig. **1a**).

We measured a relatively low but significant phylogenetic signal of the fungal phylogeny on

the antagonistic (r = 0.20 ± 0.06, p = 0.003 ± 0.005) and mutualistic (r = 0.21, p = 0.001)

networks (see effect of rarefaction depth on phylogenetic signal in Supporting Information,

Fig. **S2**). We found no significant effects of plant phylogeny on any of the networks.

Mutualistic and antagonistic plant-plant interactions

The number of fungal interactions per plant species varied, with *Aspidosperma* sp. showing

the highest normalised degree and Sapindaceae sp. the lowest amongst the mutualistic plants

(Fig. **3a**). Ranked over all the plants, three antagonistic species, namely *V. aphylla*, *V.*

parviflora and *D. orobanchoides,* presented a normalised degree in the lower half of the

spectrum, while *G. breviflorus* and *S. spruceanum* are amongst the plants with the highest

number of associated fungi. We also observed a wide range of phylogenetic species variability

(psv) for mutualistic plants, with *Acacia* sp. and *Aspidorperma* sp. with the highest *psv* and

Sapindaceae sp. the lowest, throughout which, the antagonistic plants are distributed (Fig.

3b). There was a significant correlation between the normalised degree and *psv* of the

mutualistic plants (Pearson's correlation: r = 0.54, t = 2.80, df = 19, p = 0.011), while for the

antagonistic plants the number of interactions (normalised degree) does not reflect the fungal

diversity (*psv*) in their roots (Pearson's correlation: r = -0.29, t = -0.53, df = 3, p = 0.632).

Within the antagonistic plants, we observed that *S. spruceanum* shared most fungal

interactions with *G. breviflorus*, followed by *S. spruceanum* with *D. orobanchoides* and *V.*

aphylla, and also *G. breviflorus* with *D. orobanchoides*. We observed that antagonistic plants

as a group associate with fungi that are simultaneously linked to all mutualistic plants (Fig.

1b), indicating that antagonistic plants are indirectly linked to any of the mutualistic plants

detected in this study, which could therefore be the ultimate source of carbon for the

antagonistic plants. Of all the possible connections between antagonistic and mutualistic

plant species, *S. spruceanum* had the highest fungal overlap with *Clusia* sp., and then with *Tapirira* sp. and *Protium* sp.; *G. breviflorus* with *Aspidosperma* sp. and *Protium* sp., then with *Schadera* sp. and *Clusia* sp.; *D. orobanchoides* with *Tapirira* sp. and then with Fabaceae sp. 1; *V. parviflora* with *Aspidosperma* sp. and then with Araceae sp., Malpighiaceae sp., and *Clusia* sp; and *V. aphylla* with *Aspidosperma* sp. and then with *Paloue* sp.. These mutualistic plants are also those with highest number of fungal interactions overall (Fig. **3a**). Among the mutualistic plants, *Aspidosperma* sp. had the highest fungal overlap with *Protium* sp., Malpighiaceae sp. and Metteniusaceae sp., which are also among the highest ranked species in terms of number of associated fungi and phylogenetic diversity (Fig. **3**).

Mutualistic and antagonistic plant-fungal interactions

We observed a significant correlation (Pearson correlation $r = 0.63$, $p < 0.001$, $t = 8.27$, $df = 105$) between the normalised degrees of the fungi in the antagonistic and mutualistic networks (Fig. **4,** Supporting Information Fig. **S3**). Fungi present in the mutualistic network but absent from the antagonistic network were generally characterized by a low normalized degree, except for two *Rhizophagus* taxa which were present in the majority of mutualistic plants (Fig. **4**). Moreover, we found a significant correlation of fungal ecological similarity between the mutualistic and antagonistic interactions (Mantel tests: Jaccard $r = 0.20 \pm 0.04$, $p = 0.002 \pm 0.002$; overlap $r = 0.24 \pm 0.04$, $p = 0.001$), and also when accounting for the phylogenetic relatedness among the shared fungi (partial Mantel tests: Jaccard $r = 0.18 \pm 0.003$, $p = 0.018 \pm 0.04$; overlap $r = 0.22 \pm 0.04$, $p = 0.002 \pm 0.003$). We also observed that the fungi with the highest normalized degrees, both in the mutualistic and antagonistic network, are all members of the Glomeraceae (Fig. **4**).

Network analysis

The network analysis indicated that the presence of 2,560.39 ± 296.36 instances of the diamond-shaped module in the empirical networks across the rarefied matrices was overrepresented in relation to random expectations (*z-score* = 3.45 ± 0.40, p = 0.002 ± 0.003). The same analysis repeated for multiple rarefaction depths indicates that despite the number of diamond-shaped modules increases with increasing rarefaction depth, it does not impact

the result that pairs of fungi share an antagonistic and a mutualistic plant more often than

expected by chance (Supporting Information, Fig. **S4**).

Discussion

We found that antagonistic plants as a group target approximately half of the fungi that are

potentially available, yet this subset of fungi is associated with all mutualistic plants detected

in this study. Thus, despite associating only with a subset of the local pool of fungi, the

antagonists are still indirectly linked to every mutualistic plant. Fungi not detected in the roots

of antagonists were generally connected to few mutualistic plants. Within the fungi that are

shared between the mutualistic and antagonistic networks, we detected a significant

ecological symmetry between the mutualistic and antagonistic interactions of the fungi: pairs

of fungi that interact with similar sets of mutualistic plants also interact with overlapping

antagonistic plants. The network analysis indicates that this pattern occurs more often than

expected by chance. Based on the highest fungal overlap between each antagonistic species

with subsets of highly connected mutualistic species, this pattern is mainly driven by an

antagonist preference for fungi that are well-linked to specific mutualistic plants. These

mutualistic plants are the ultimate sources of the carbon which antagonistic plants take up

from the fungi shared between mutualists and antagonists. Therefore, we suggest that the

observed pattern reflects a strategy in which the maintenance of antagonistic interactions is

maximized by targeting well-linked mutualistic fungi, thereby minimizing the risk of carbon

supply shortages.

Fungal preferences of antagonistic plants

We found that plant species identity had a significant influence on the fungal community

composition, regardless of the plant type (mutualist or antagonist), which indicates that these

communities are non-random subsets of the local fungal taxon pool. This supports previous

evidence that co-occurring plant species show differences in selectivity towards available

arbuscular mycorrhizal fungi (Davison *et al.*, 2011). Antagonistic plant species in particular,

are known to select particular groups of fungi, often a narrower range than surrounding

mutualistic plants (Bidartondo et al. 2002; Gomes et al. 2017). Here we observed that five co-
occurring antagonistic plant species collectively associate with about half of the available
fungal taxa (Fig. **4**). Antagonistic interactions can thus be supported by a relatively wide array
of arbuscular mycorrhizal taxa, as shown previously (Merckx et al. 2012, Gomes *et al.*, 2017b;
Sheldrake *et al.*, 2017). Although fungi from three different fungal families were detected in
the roots of antagonistic plants, a clear preference for Glomeraceae taxa, and *Rhizophagus*
irregularis relatives in particular, was observed. The taxa of this clade were the most
frequently encountered in the roots of the autotrophic plants as well (Fig. **4**). *Rhizophagus*
contains some of the most globally widespread and common arbuscular mycorrhizal fungi
(Kivlin *et al.*, 2011; Moora *et al.*, 2011; Davison *et al.*, 2015; Gomes *et al.*, 2018) although this
is mostly derived from studies in temperate areas. Our results indicate that the tropical
rainforest offers no exception to this pattern. Glomeraceae are usually not only the most
dominant clade in natural arbuscular mycorrhizal communities, often accounting for c. 70%
of all species (Montesinos-Navarro *et al.*, 2012), but they also have consistently been found
to include the most generalist arbuscular mycorrhizal fungi in other network studies (e.g.
Montesinos-Navarro *et al.,* 2012; Chagnon *et al.*, 2015; Chen *et al.*, 2017). The ability to
interact with many mutualistic plant species can be a potential reason for why antagonists
generally target Glomeraceae fungi (Merckx *et al.*, 2012; Renny *et al.*, 2017). Ecological theory
predicts that generalist species tend to have large distribution ranges (Brown, 1984) and,
consequently, are less vulnerable to (local) extinction than specialized species (Schleuning *et*
al., 2016). Therefore, associations to generalist fungi may be advantageous for the
evolutionary persistence of antagonists. Furthermore, associations with multiple mutualistic
plant partners may increase fungal resilience to disturbance, while mediating temporal
fluctuations in carbon flow and interaction dynamics (Bennett *et al.*, 2013) guaranteeing
continuous carbon supply to the entire network without pronounced negative effects even in
the presence of antagonists. In addition, in the context of mycorrhizal fungi, which can be
linked to different plant species simultaneously (Montesinos-Navarro *et al.*, 2012), generalist
fungi are therefore likely to be more reliable carbon sources for antagonists. An alternative
and perhaps not mutually exclusive explanation for why antagonistic plants preferentially
target well-connected fungi may be that these fungi are less effective in detecting and
excluding non-mutualistic plant partners (Bruns *et al.*, 2002; Egger & Hibbett, 2004;
Bidartondo, 2005; Walder & Van Der Heijden, 2015).

 Fungal links between mutualistic and antagonistic networks

We detected that pairs of fungi that interact with similar sets of mutualistic plants share links

with overlapping antagonistic plants. Thus, there is a high level of interaction symmetry

between mutualistic and antagonistic mycorrhizal networks. Also, we measured a significant

influence of the fungal phylogenetic relationships on both the mutualistic and antagonistic

interactions, showing that closely related fungi interact with similar mutualistic and

antagonistic plants respectively. Because, biotic interactions are mediated by functional

traits, and most functional traits are evolutionarily conserved, shared evolutionary history of

fungi can serve as a proxy for functional similarity. We therefore hypothesize that both

mutualistic and antagonistic interactions are shaped by evolutionary conserved functional

traits of the fungi. In addition, the network analysis indicated that pairs of fungi share an

antagonistic and a mutualistic plant more often than expected by chance. This analysis solely

indicates that the diamond-shaped module is overrepresented in the empirical network (Fig.

2), without reference to species degree nor to species identity. Yet, our results support that

the observed pattern is driven by the tendency of antagonistic plants to target fungi that are

well-linked to mutualistic plants (Fig. **1b**). The mutualistic plants with highest fungal overlap

in relation to the antagonistic plants are among those with the highest ranked degree and

phylogenetic species variability from the pool of detected mutualistic plant species (Fig. **3**).

Moreover, fungi with the highest number of interactions in the mutualistic network are also

among the best-connected fungi in the antagonistic network (Fig. **4**). Our findings thus reveal

that antagonistic plants preferentially associate with fungi that are simultaneously linked to

a wide range of mutualistic plants. Although many antagonistic plants share a large number

of fungi with the mutualistic tree *Aspidosperma* sp., the mutualistic plants with highest

normalised degree, each antagonistic plant species also indirectly associates with non-

overlapping sets of mutualistic plants, as indicated by their divergent positions in the plant-

plant interaction network (Fig. **2b**). This pattern suggests a potential strategy to avoid

competition among antagonistic plant species while targeting well connected mutualistic

plant species through shared fungi.

Our results indicate that antagonistic and mutualistic mycorrhizal networks are not linked

randomly but are symmetrically influenced by mutualistic and antagonistic plant identity and

fungal connectance. Whether these correlations are potentially influenced by spatial patterns

in the distribution of mutualists, antagonists, and fungi remain to be determined. Our earlier

work suggests that antagonistic mycorrhizal interactions can be better explained by

understanding plant–plant interactions through sharing fungal associations (Gomes *et al.*,

2017b), and indicates that antagonistic interactions may respond to an ecological mechanism

driven by maximizing co-occurrence and avoiding competitive exclusion among antagonists.

The results obtained here expand this view and suggest that not only plant-plant interactions

drive patterns of co-occurrence of antagonistic plants, but also the indirect interactions

between mutualistic and antagonistic plants through shared arbuscular mycorrhizal fungi.

Potential sampling biases

Considering that rainforests are species-rich ecosystems (e.g. (Ter Steege *et al.*, 2013)), it is

possible that, despite our efforts, the sampling of the belowground diversity of arbuscular

mycorrhizal fungi, and their plant partners remained incomplete. Furthermore, the

representation of roots from antagonistic and mutualistic plants was necessarily imbalanced,

as whole and partial root systems, respectively, were investigated. This made the use of read

abundances to estimate interaction strengths impossible. While other studies have addressed

the importance of sampling intensity, (i.e. the number of possible interactions per node which

directly impacts the normalised degree per species), and network size (i.e. number of nodes

involved) (Blüthgen *et al.*, 2007; Dormann *et al.*, 2009) on network metrics, the effect of

species relationships is often elusive. In our approach, we accounted for the phylogenetic

relationships between the fungi, as other studies have accounted for taxonomic diversity

(Morris *et al.*, 2014), and also we verified the consistency of our results across multiple

rarefactions depths, by the use of multiple rarefied matrices at each depth (Gotelli & Colwell,

2001). This allows us to separate the statistical patterns in our data from the influence of

sampling effort, both in terms of the plant species detected in the sampled roots, and in their

potentially incomplete number of fungal associations. Furthermore, the number of

interactions for mutualistic plants is moderately correlated with the phylogenetic diversity of

the arbuscular mycorrhizal fungi associated with these tropical trees, while for antagonistic

plants this correlation is not found. This further suggests that bias due to sampling effort is

limited. It also indicates that antagonistic plants have specialised interactions, occupying

particular fungal niches, which could have consequence for their occurrence and coexistence
(Gomes *et al.*, 2017b).

Conclusions and future perspectives

To our knowledge, our study is the first to assess how antagonistic plant species are
embedded in mutualistic mycorrhizal networks. We find that antagonistic plants as a group
interact with half of the available fungal partners, and generally target fungi that are well-
connected to mutualistic plants. Although antagonistic species show overlap in their fungal
associations, we found that they are indirectly linked to different sets of mutualistic plants
suggesting a potential mechanism to avoid competition by preferentially relying on different
carbon sources (Gomes *et al.*, 2017b). The phylogenetic relationships between the fungi,
likely a proxy for fungal traits, have a significant influence on these non-random tripartite
interactions. Therefore, we conclude that the persistence of antagonists in arbuscular
mycorrhizal networks is dependent on particular well-connected 'keystone' mycorrhizal
fungi, which provide the antagonists with carbon from a wide range of plants. Our observation
that the way fungi connect mutualistic and antagonistic networks is not random and that well-
connect fungal nodes in arbuscular mycorrhizal networks are more prone to be targeted by
antagonists are similar to those of (Sauve *et al.*, 2016) for a plant-pollinator-herbivore
network when considering binary interactions. But further research is needed to assess
whether this is a general feature of diffuse interactions, also when taking interaction strength
into account. Our study emphasises the raising awareness of considering multiple interaction
types simultaneously (e.g. antagonistic and mutualistic) to deepen our understanding of
complex biodiversity patterns (Losapio *et al.*, 2021).

In ectomycorrhizal systems mutualistic and achlorophyllous antagonistic plant species
represent opposite extremes of a continuum of interaction outcomes and several plant
species are able to combine photosynthesis and carbon uptake from fungi, a strategy known
as 'partial mycoheterotrophy' (Selosse & Roy, 2009). Recent discoveries now indicate that
partial mycoheterotrophy may be common among arbuscular mycorrhizal plants as well, and
potentially many photosynthetic understory plants are able to take up carbon from
associated arbuscular mycorrhizal fungi (Giesemann *et al.*, 2021). It remains to be tested

whether carbon uptake by these partially mycoheterotrophic plants occurs through sets of

fungi, as suggested for fully mycoheterotrophic plant species.

Acknowledgements

This research was supported by a Veni grant from the Netherlands Organisation of Scientific

Research (NWO) to VSFTM (863.11.018) and field work was funded by the Royal Academy of

Arts and Sciences of the Netherlands (KNAW) Ecology fund, grant J1606/Eco/G437 to SIFG. JB

is supported by the Swiss National Science Foundation (Grant 31003A-169671 3). The authors

thank the field assistance of Mélanie Roy, Vincent, Mathieu Gerard, the molecular lab

assistance of Elza Duijm, and Hans ter Steege for verification of plant species occurrence in

French Guiana. The authors also thank the valuable comments of four anonymous reviewers

and editor Marc-André Selosse, which greatly improved the quality of the manuscript.

Author contributions

SG, MF, JB, and VM designed the study and collected the materials. SG performed the

laboratory work and data analysis. All authors contributed to the data analysis and the writing

of the manuscript.

References

Bascompte J, Jordano P. **2007**. Plant-Animal Mutualistic Networks: The Architecture of

Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**: 567+593.

Bascompte J, Jordano P. **2013**. *Mutualistic networks*. Princeton: Princeton University Press.

Bascompte J, Melián C. **2005**. Simple trophic modules for complex food webs. *Ecology* **86**:

2868–2873.

Bennett AE, Daniell TJ, Öpik M, Davison J, Moora M, Zobel M, Selosse MA, Evans D. **2013**.

Arbuscular mycorrhizal fungal networks vary throughout the growing season and between

successional stages. *PLoS ONE* **8**.

Berry PE, Weitzman AL. **2007**. Checklist of the Plants of the Guiana Shield (Venezuela:

Amazonas, Bolivar, Delta Amacuro; Guyana, Surinam, French Guiana). In: Funk, Vicki Ann,

Hollowell, Thomas H., Berry, P. E., Kelloff, Carol L., and Alexander SN, ed. Checklist of the

Plants of the Guiana Shield (Venezuela: Amazonas, Bolivar, Delta Amacuro; Guyana,

Surinam, French Guiana). Washington D. C.: Smithsonian Institution, 535–536.

Bever JD, Richardson SC, Lawrence BM, Holmes J, Watson M. **2009**. Preferential allocation

to beneficial symbiont with spatial structure maintains mycorrhizal mutualism. *Ecology*

Letters **12**: 13–21.

Bidartondo MI. **2005**. The evolutionary ecology of myco-heterotrophy. *New Phytologist*

167: 335–352.

Bidartondo MI, Read DJ, Trappe JM, Merckx V, Ligrone R, Duckett JG. **2011**. The dawn of

symbiosis between plants and fungi. *Biology letters* **7**: 574–577.

Bidartondo MI, Redecker D, Hijri I, Wiemken A, Bruns TD, Domínguez L, Sérsic A, Leake JR,

Read DJ. **2002**. Epiparasitic plants specialized on arbuscular mycorrhizal fungi. *Nature* **419**:

389–392.

Blüthgen N, Menzel F, Hovestadt T, Fiala B, Blüthgen N. **2007**. Specialization, Constraints,

and Conflicting Interests in Mutualistic Networks. *Current Biology* **17**: 341–346.

Brown JH. **1984**. On the Relationship between Abundance and Distribution of Species. *The*

American Naturalist **124**: 255–279.

Bruns TD, Bidartondo MI, Taylor DL. **2002**. Host specificity in ectomycorrhizal communities:

what do the exceptions tell us? *Integrative and comparative biology* **42**: 352–359.

Chagnon PL, Bradley RL, Klironomos JN. **2015**. Trait-based partner selection drives

mycorrhizal network assembly. *Oikos* **124**.

Chen L, Zheng Y, Gao C, Mi XC, Ma KP, Wubet T, Guo LD. **2017**. Phylogenetic relatedness

explains highly interconnected and nested symbiotic networks of woody plants and

arbuscular mycorrhizal fungi in a Chinese subtropical forest. *Molecular Ecology* **26**: 2563–

2575.

Davison J, Moora M, Öpik M, Adholeya A, Ainsaar L, Bâ A, Burla S, Diedhiou AG, Hiiesalu I,

Jairus T, *et al.* 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals

very lowendemism. *Science* **349**: 970–973.

Davison J, Öpik M, Daniell TJ, Moora M, Zobel M. **2011**. Arbuscular mycorrhizal fungal

communities in plant roots are not random assemblages. *FEMS Microbiology Ecology* **78**:

103–115.

Dormann C, Fruend J, Gruber B. **2009**. Indices, graphs and null models: analysing bipartite

ecological networks. *The Open Ecology Journal* **2**: 7–24.

Dormann CF, Gruber B, Fruend J. **2018**. Introducing the bipartite Package: Analysing

Ecological Networks. *R news* **8**: 8–11.

Edgar RC. **2010**. Search and clustering hundreds of times faster than BLAST. *Bioinformatics*

26: 2460–2461.

Egger KN, Hibbett DS. **2004**. The evolutionary implications of exploitation in mycorrhizas.

Canadian Journal of Botany **82**: 1110–1121.

Flynn JM, Brown EA, Chain FJJ, Macisaac HJ, Cristescu ME. **2015**. Toward accurate

molecular identification of species in complex environmental samples: Testing the

performance of sequence filtering and clustering methods. *Ecology and Evolution*.

Fortuna MA, Bascompte J. **2006**. Habitat loss and the structure of plant-animal mutualistic

networks. *Ecology Letters* **9**: 281–286.

Giesemann P, Rasmussen HN, Gebauer G. **2021**. Partial mycoheterotrophy is common

among chlorophyllous plants with Paris-type arbuscular mycorrhiza. *Annals of botany* **127**:

645–653.

Gomes SIF, Aguirre-Gutiérrez J, Bidartondo MI, Merckx VSFT. **2017a**. Arbuscular

mycorrhizal interactions of mycoheterotrophic Thismia are more specialized than in

autotrophic plants. *New Phytologist* **213**: 1418–1427.

Gomes SIFF, van Bodegom PM, Merckx VSFTFT, Soudzilovskaia NA. **2019a**. Global

distribution patterns of mycoheterotrophy. *Global Ecology and Biogeography* **28**: 1133–

1145.

Gomes SIF, van Bodegom PM, Merckx VSFT, Soudzilovskaia NA. **2019b**. Environmental

drivers for cheaters of arbuscular mycorrhizal symbiosis in tropical rainforests. *New*

Phytologist **223**: 1575–1583.

Gomes SIFF, Merckx VSFTFT, Hynson NA. **2018**. Biological invasions increase the richness of

arbuscular mycorrhizal fungi from a Hawaiian subtropical ecosystem. *Biological Invasions*

20: 2421–2437.

Gomes SIF, Merckx VSFT, Saavedra S. **2017b**. Fungal-host diversity among

mycoheterotrophic plants increases proportionally to their fungal-host overlap. *Ecology and*

Evolution **7**: 3623–3630.

Gotelli NJ, Colwell RK. **2001**. Quantifying biodiversity: Procedures and pitfalls in the

measurement and comparison of species richness. *Ecology Letters* **4**: 379–391.

Helmus MR, Bland TJ, Williams CK, Ives AR. **2007a**. Phylogenetic Measures of Biodiversity.

The American Naturalist **169**: E68–E83.

Helmus MR, Bland TJ, Williams CK, Ives AR. **2007b**. Phylogenetic measures of biodiversity.

The American naturalist **169**: E68-83.

Ihrmark K, Bödeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y,

Stenlid J, Brandström-Durling M, Clemmensen KE, *et al.* **2012**. New primers to amplify the

fungal ITS2 region - evaluation by 454-sequencing of artificial and natural communities.

FEMS Microbiology Ecology **82**: 666–677.

Imhof S, Massicotte HB, Melville LH, Peterson RL. **2013**. Subterranean morphology and

mycorrhizal structures. In: Mycoheterotrophy: The Biology of Plants Living on Fungi. 157–

214.

Kiers ET, Duhamel M, Beesetty Y, Mensah J a, Franken O, Verbruggen E, Fellbaum CR,

Kowalchuk G a, Hart MM, Bago A, *et al.* **2011**. Reciprocal Rewards Stabilize in the

Mycorrhizal Symbiosis. *Science* **333**: 880–882.

Kivlin SN, Hawkes C V., Treseder KK. **2011**. Global diversity and distribution of arbuscular

mycorrhizal fungi. *Soil Biology and Biochemistry* **43**: 2294–2303.

Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns TD,

Bengtsson-Palme J, Callaghan TM, *et al.* **2013**. Towards a unified paradigm for sequence-

based identification of fungi. *Molecular Ecology* **22**: 5271–5277.

Krüger M, Krüger C, Walker C, Stockinger H, Schüßler A. **2012**. Phylogenetic reference data

for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species

level. *New Phytologist* **193**: 970–984.

Kumar S, Stecher G, Suleski M, Hedges SB. **2017**. TimeTree: a resource for timelines,

timetrees, and divergence times. *Molecular Biology and Evolution* **34**: 1812–1819.

Leake JR. **1994**. The biology of myco-heterotrophic ('saprophytic') plants. *New Phytologist*

127: 171–216.

Lee EH, Eo JK, Ka KH, Eom AH. **2013**. Diversity of arbuscular mycorrhizal fungi and their

roles in ecosystems. *Mycobiology* **41**: 121–125.

Lekberg Y, Gibbons SM, Rosendahl S. **2014**. Will different OTU delineation methods change

interpretation of arbuscular mycorrhizal fungal community patterns? *New Phytologist* **202**:

1101–1104.

Lindahl BD, Nilsson RH, Tedersoo L, Abarenkov K, Carlsen T, Kjøller R, Kõljalg U, Pennanen

T, Rosendahl S, Stenlid J, *et al.* **2013**. Fungal community analysis by high-throughput

sequencing of amplified markers - a user's guide. *New Phytologist* **199**: 288–299.

Losapio G, Schöb C, Staniczenko PPA, Carrara F, Palamara GM, de Moraes CM, Mescher

MC, Brooker RW, Butterfield BJ, Callaway RM, *et al.* **2021**. Network motifs involving both

competition and facilitation predict biodiversity in alpine plant communities. *Proceedings of*

the National Academy of Sciences of the United States of America **118**: 1–6.

Martín González AM, Dalsgaard B, Olesen JM. **2010**. Centrality measures and the

importance of generalist species in pollination networks. *Ecological Complexity* **7**: 36–43.

McMurdie PJ, Holmes S. **2014**. Waste not, want not: why rarefying microbiome data is

inadmissible. *PLoS Computational Biology*.

Merckx VSFT. **2013**. *Mycoheterotrophy: The biology of plants living on fungi*. New York, NY,

USA: Springer.

Merckx V, Bidartondo MI. **2008**. Breakdown and delayed cospeciation in the arbuscular

mycorrhizal mutualism. *Proceedings of the Royal Society B* **275**: 1029–1035.

Merckx VSFT, Janssens SB, Hynson NA, Specht CD, Bruns TD, Smets EF. **2012**.

Mycoheterotrophic interactions are not limited to a narrow phylogenetic range of

arbuscular mycorrhizal fungi. *Molecular Ecology* **21**: 1524–1532.

Merckx VSFT, Mennes CB, Peay KG, Geml J. **2013**. Evolution and diversification. In:

Mycoheterotrophy: The Biology of Plants Living on Fungi.

Mills LS, Doak DF. **1993**. The Keystone-Species Concept in Ecology and Conservation.

BioScience **43**: 219–224.

Milo R, Shen-Orr S, Itzkovitz S, Kashtan N, Chklovskii D, Alon U. **2002**. Network motifs:

Simple building blocks of complex networks. *Science* **298**: 824–827.

Montesinos-Navarro, A. S-M, J. G., Valiente-Banuet A, Verdu M. **2015**. Evidence for

phylogenetic correlation of plant-AMF assemblages? *Annals of Botany* **115**: 171–177.

Montesinos-Navarro A, Segarra-Moragues JG, Valiente-Banuet A, Verdú M. **2012**. The

network structure of plant-arbuscular mycorrhizal fungi. *New Phytologist* **194**: 536–547.

Moora M, Berger S, Davison J, Öpik M, Bommarco R, Bruelheide H, Kühn I, Kunin WE,

Metsis M, Rortais A, *et al.* **2011**. Alien plants associate with widespread generalist

arbuscular mycorrhizal fungal taxa: Evidence from a continental-scale study using massively

parallel 454 sequencing. *Journal of Biogeography* **38**: 1305–1317.

Morris RJ, Gripenberg S, Lewis OT, Roslin T. **2014**. Antagonistic interaction networks are

structured independently of latitude and host guild. *Ecology Letters* **17**: 340–349.

Olesen JM, Bascompte J, Jordano P, Melia CJ. **2003**. The nested assembly of plant–animal

mutualistic networks ´. *Proceedings of the National Academy of Sciences of the United*

States of America **100**: 9383–9387.

Pellmyr O, Leebens-Mack J. **1999**. Forty million years of mutualism: Evidence for eocene

origin of the yucca-yucca moth association. *Proceedings of the National Academy of*

Sciences of the United States of America **96**: 9178–9183.

Renny M, Acosta MC, Cofré N, Domínguez LS, Bidartondo MI, Sérsic AN. **2017**. Genetic

diversity patterns of arbuscular mycorrhizal fungi associated with the mycoheterotroph

Arachnitis uniflora Phil. (Corsiaceae). *Annals of Botany* **119**: 1279–1294.

Saavedra S, Rohr RP, Dakos V, Bascompte J. **2013**. Estimating the tolerance of species to

the effects of global environmental change. *Nature Communications* **4**.

Sachs JL, Simms EL. **2006**. Pathways to mutualism breakdown. *Trends in Ecology and*

Evolution **21**: 585–592.

Sauve AMC, Thébault E, Pocock MJO, Fontaine C. **2016**. How plants connect pollination and

herbivory networks and their contribution to community stability. *Ecology* **97**: 908–917.

Schleuning M, Fründ J, Schweiger O, Welk E, Albrecht J, Albrecht M, Beil M, Benadi G,

Blüthgen N, Bruelheide H, *et al.* 2016. Ecological networks are more sensitive to plant than

to animal extinction under climate change. *Nature Communications* **7**: 13965.

Selosse MA, Rousset F. **2011**. The plant-fungal marketplace. *Science* **333**: 828–829.

Selosse MA, Roy M. **2009**. Green plants that feed on fungi: facts and questions about

mixotrophy. *Trends in Plant Science* **14**: 64–70.

Sepp S, Davison J, Jairus T, Vasar M, Moora M, Zobel M, Öpik M. **2019**. Non-random

association patterns in a plant–mycorrhizal fungal network reveal host–symbiont specificity.

Molecular Ecology **28**: 365–378.

Sheldrake M, Rosenstock NP, Revillini D, Olsson PA, Wright SJ, Turner BL. **2017**. A

phosphorus threshold for mycoheterotrophic plants in tropical forests. *Proceedings of the*

Royal Society B: Biological Sciences **284**: 20162093.

Smith SE, Read DJ. **2008**. *Mycorrhizal Symbiosis*. Cambridge: Academic Press, San Diego, CA.

Southworth D, He X-H, Swenson W, Bledsoe CS, Horwath WR. **2005**. Application of

network theory to potential mycorrhizal networks. *Mycorrhiza* **15**: 589–595.

Ter Steege H, Pitman NCA, Sabatier D, Baraloto C, Salomão RP, Guevara JE, Phillips OL,

Castilho C V., Magnusson WE, Molino JF, *et al.* 2013. Hyperdominance in the Amazonian

tree flora. *Science* **342**: 1243092.

Stouffer DB, Camacho J, Jiang W, Amaral LAN. **2007**. Evidence for the existence of a robust

pattern of prey selection in food webs. *Proceedings of the Royal Society B: Biological*

Sciences **274**: 1931–1940.

Strullu-Derrien C, Selosse M-A, Kenrick P, Martin FM. **2018**. The origin and evolution of

mycorrhizal symbioses: from palaeomycology to phylogenomics. *New phytologist* **172**: 589–

519.

Thompson JN. **2005**. *The Geographic Mosaic of Coevolution*. University of Chicago Press,

Chicago.

Toju H, Guimaraes PR, Olesen JM, Thompson JN. **2015**. Below-ground plant-fungus network

topology is not congruent with above-ground plant-animal network topology. *Science*

Advances **1**: e1500291–e1500291.

Walder F, Van Der Heijden MGA. **2015**. Regulation of resource exchange in the arbuscular

mycorrhizal symbiosis. *Nature Plants* **1**.

Waterman RJ, Klooster MR, Hentrich H, Bidartondo MI. **2013**. Species interactions of

mycoheterotrophic plants: Specialization and its potential consequences. In:

Mycoheterotrophy: The Biology of Plants Living on Fungi. 267–296.

White TJ, Bruns T, Lee S, Taylor J. **1990**. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds.

PCR Protocols: A guide to methods and applications. San Diego: Academic Press, 315–322.

Figures

a

b

Figure 1. Tripartite arbuscular mycorrhizal interactions. (a) Visualization of the tripartite network between fungi (grey) and antagonist (yellow), and mutualist (green) plants, where edges represent a connection between a plant and a fungus. (b) Plant-plant fungal overlap network, in which edges represent a link between plant species through shared arbuscular mycorrhizal fungus. The thickness of the lines represents the interaction strength between the plants (the thicker, the more fungi are shared). Yellow lines link antagonistic and mutualistic plants; light grey lines link mutualistic to mutualistic plants; and dark grey lines link antagonistic to antagonistic plants. Identification of autotrophic plants is indicated by the first three letters of their name (see full names in Fig. 3); mycoheterotrophic plants are *S. spruceanum* (SS), *G. breviflorus* (GB), *D. orobanchoides* (DO), *V. aphylla* (VA), and *V. parviflora* (VP). In both network representations (a, b), one of the 100 rarefied matrices to a depth of 844 reads was used; and the Fruchterman-Reingold layout was used, in which nodes are evenly distributed through the graph, where plants that share more connections are closer to each other.

748 **Figure 2.** Diamond-shape module investigated in this paper. Representation of the module

749 linking pairs of fungi to the same antagonistic and mutualistic plant species (a). Examples of

750 under- (b) and overrepresentation (c) of the module in a network. When the module is

751 overrepresented it is considered a network motif.

Figure 3: Mean ± SD plant normalised degree and PSV for the 21 autotrophic and five mycoheterotrophic plant species, resulted from the 100 rarefactions to a depth of 844 reads.

Figure 4: Plant-fungal interactions of mutualistic (green) and antagonistic (yellow) plants using a matrix rarefied to a depth of 844 reads. Phylogenetic relationships between the fungi are shown on top. Plant species are listed on the left; hierarchy clustered dendrogram based on the Bray-Curtis distance of their fungal communities is shown on the right. The intensity of the orange dots on the tips of the fungal phylogeny depicts the normalised degree (representing their interaction strength) of each fungus in each of the plant-fungal networks (i.e. in the antagonistic plants-fungi and mutualistic plants-fungi networks).