# Differential gene expression elicited by ZIKV infection in trophoblasts from congenital Zika syndrome discordant twins

Murilo Sena Amaral[1¶], Ernesto Goulart[2¶], Luiz Carlos Caires-Júnior[2¶], David Abraham Morales-Vicente[1,3], Alessandra Soares-Schanoski[4], Roselane Paiva Gomes[5], Giovanna Gonçalves de Oliveira Olberg[1], Renato Mancini Astray[5], Jorge E. Kalil[6], Mayana Zatz[2], Sergio Verjovski-Almeida[1,3*]

[1]Laboratório de Parasitologia, Instituto Butantan, São Paulo, Brazil

[2]Departmento de Genética e Biologia Evolutiva, Centro de Pesquisas sobre o Genoma Humano e Células Tronco, Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil

[3]Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, Brazil

[4]Laboratorio de Bacteriologia, Instituto Butantan, São Paulo, Brazil

[5]Laboratorio de Imunologia Viral, Instituto Butantan, São Paulo, Brazil

[6]Laboratório de Imunologia, Faculdade de Medicina e Instituto do Coração, Universidade de São Paulo, São Paulo, Brazil

* Corresponding author:
Email: verjo@iq.usp.br (SVA)

¶These authors contributed equally to this work.

**Short title: Molecular signatures from ZIKV-infected discordant twins' trophoblasts**

## Abstract

Zika virus (ZIKV) causes congenital Zika syndrome (CZS), which is characterized by fetal demise, microcephaly and other abnormalities. ZIKV in the pregnant woman circulation must cross the placental barrier that includes fetal endothelial cells and trophoblasts, in order to reach the fetus. CZS occurs in ~1-40% of cases of pregnant women infected by ZIKV, suggesting that mothers' infection by ZIKV during pregnancy is not deterministic for CZS phenotype in the fetus. Therefore, other susceptibility factors might be involved, including the host genetic background. We have previously shown that in three pairs of dizygotic twins discordant for CZS, neural progenitor cells (NPCs) from the CZS-affected twins presented differential *in vitro* ZIKV susceptibility compared with NPCs from the non-affected. Here, we analyzed human-induced-pluripotent-stem-cell-derived (hiPSC-derived) trophoblasts from these twins and compared by RNA-Seq the trophoblasts from CZS-affected and non-affected twins. Following *in vitro* exposure to a Brazilian ZIKV strain (ZIKV$^{BR}$), trophoblasts from CZS-affected twins were significantly more susceptible to ZIKV$^{BR}$ infection when compared with trophoblasts from the non-affected. Transcriptome profiling revealed no differences in gene expression levels of ZIKV candidate attachment factors, IFN receptors and IFN in the trophoblasts, either before or after ZIKV$^{BR}$ infection. Most importantly, ZIKV$^{BR}$ infection caused, only in the trophoblasts from CZS-affected twins, the downregulation of genes related to extracellular matrix organization and to leukocyte activation, which are important for trophoblast adhesion and immune response activation. In addition, only trophoblasts from non-affected twins secreted significantly increased amounts of chemokines RANTES/CCL5 and IP10 after infection with ZIKV$^{BR}$. Overall, our results showed that trophoblasts from non-affected twins have the ability to more efficiently activate genes that are known to play important roles in cell adhesion and in triggering the immune response to ZIKV infection in the placenta, and this may contribute to predict protection from ZIKV dissemination into fetuses' tissues.

**Author summary**

The Zika virus (ZIKV) infection in adults is usually characterized by mild flu-like symptoms, with most cases remaining asymptomatic. However, in the last years, widespread ZIKV infection was shown for the first time to be associated with congenital Zika syndrome (CZS) and death of neonates. CZS is a very debilitating condition that includes microcephaly and mental retardation, leading to a strong social and health impact. This dramatic condition calls for a careful evaluation of the molecular mechanisms involved in ZIKV infection in the maternal-fetal interface. It is estimated that CZS occurs in ~1-40% of cases of pregnant women infected by ZIKV, which suggests that different susceptibility factors might be involved, including the host genetic background. By analyzing trophoblast cells that recapitulate the placenta from three pairs of dizygotic twins discordant for CZS, we were able to show that trophoblasts from CZS-affected twins were significantly more susceptible to ZIKV infection when compared with trophoblasts from the non-affected twins. We also provide a detailed picture of genes differentially expressed by trophoblasts from the discordant twins after infection with ZIKV. These genes can be further investigated as possible therapeutic targets to avoid viral dissemination into developing fetus' tissues.

**Introduction**

Zika virus (ZIKV) is a flavivirus with sporadic outbreaks reported in several countries, causing an infection usually characterized by mild symptoms, where up to 80% of cases remain asymptomatic [1-3]. However, most likely due to the mutation acquired during the large outbreak recorded in French Polynesia in 2013–2014 [4-6], for the first time, widespread ZIKV infection was shown to be associated with congenital Zika syndrome (CZS) and death of neonates [3,7-13].

CZS is characterized by variable clinical presentations, including fetal demise, microcephaly and other abnormalities (hearing and ocular loss, mental retardation, epilepsy, muscle weakness,

learning disabilities and behavioral abnormalities) [14-16]. It is well established now that ZIKV

shows tropism for a wide range of host tissues [17-19], especially neuronal cell types, including

neural progenitor cells, mature neurons and astrocytes (reviewed by Christian et al. [20]), and

replicates in *ex vivo* slices from adult human cortical tissues [21]. Also, Zika virus infection

reprograms global transcription in the host cells [22]. ZIKV in the maternal circulation needs to cross

the placental barrier that includes fetal endothelial cells and trophoblasts in order to reach the fetus

[23,24]. Several placenta related cells have been shown to be infected by ZIKV, including placental

macrophages and trophoblasts [25-29].

It has been estimated that CZS occurs in ~1-40% of cases of pregnant women infected by

ZIKV [3,13,30,31]. This suggests that mothers' infection by ZIKV during pregnancy is not the only

factor determining CZS phenotype in the fetus, and other susceptibility factors might be involved.

Indeed, neural progenitor cells (NPCs) from different individuals have been shown to respond

differently to ZIKV infection [32,33]. In this scenario, discordant twins represent a good case–control

sample to test for the genetic contribution determining the fetuses' discordant outcome of gestational

infection with ZIKV, as they are supposed to have been exposed to ZIKV under similar conditions in

the uterus during gestation.

We have previously shown that twins discordant for CZS outcome show differential neural

progenitor cells (NPCs) *in vitro* viral susceptibility to a Brazilian ZIKV strain (ZIKV[BR]) [32], but it

remains to be determined if discordant CZS twins also show differential placental susceptibility to

viral infection. Here, we compared the susceptibility and molecular signatures associated with *in*

*vitro* ZIKV[BR] infection of trophoblasts from the same CZS-affected and non-affected twins, using a

well-established trophoblasts model that recapitulates the primitive placenta formed during

implantation [34,35]. We show here that hiPSC-derived trophoblasts from CZS-affected twins were

significantly more susceptible to *in vitro* ZIKV[BR] infection when compared with trophoblasts from

non-affected twins. In this context, we had previously shown that, before infection, ESC-derived

trophoblasts express a wide range of attachment factors for ZIKV entry and lack the components of a

robust antiviral response system [35]; in contrast, cells from term placentas, which resist infection, do

not express genes encoding attachment factors implicated in ZIKV entry and do express many genes

associated with antiviral defense [35]. However, no ZIKV infection assays were performed with

these ESC-derived trophoblasts [35]. Here, transcriptome profiling of hiPSC-derived trophoblasts

revealed that ZIKV[BR] infection elicited different responses in hiPSC-derived trophoblasts from CZS-

affected and non-affected twins, highlighting that genes involved with extracellular matrix

organization as well as with immune response activation in the placental tissue may contribute to

modulate ZIKV infection outcome.

**Results**

**Infection with ZIKV[BR] of hiPSC-derived trophoblasts from discordant twins**

We obtained blood from three pairs of dizygotic twins discordant for CZS (non-affected:

#10608-4, #10763-4, and #10788-4; CZS-affected: #10608-1, #10763-1, and #10788-1) for

generation of hiPSC-derived trophoblasts and for phenotypic and gene expression analysis after *in*

*vitro* infection with ZIKV[BR] (**Fig 1A**). Erythroblasts from the three pairs of DZ twins were

reprogrammed towards hiPSCs. All hiPSC lines were previously shown by immunofluorescence

staining to express markers of pluripotency (TRA-1-60 and OCT4) and by RT-qPCR to express

endogenous pluripotent transcription factors including *NANOG* and *OCT4* [32].

Then, the hiPSCs originated from the three pairs of twins were differentiated into primitive

trophoblasts using an established protocol [34,35] and further characterized to confirm their

differentiation *in vitro*. Under these conditions, we found that all hiPSC-derived trophoblast lines

robustly expressed chorionic gonadotropin subunit beta 3 (β-CG/ hCGB3) and keratin 7 (CK7/KRT7)

(**Fig 1B**), two of the most commonly used trophoblast markers [36].

Subsequently, we infected hiPSC-derived trophoblasts with ZIKV$^{BR}$ using multiplicity of

infection (MOI) of 0.3 and 3, and at 96 hpi (hours post-infection) we investigated the viral titer by

measuring the number of plaque-forming units (PFU) in cell culture supernatants (**Fig 1C**). Virus

titers were significantly higher (2.8-fold) in the supernatant of CZS-affected twins' trophoblasts

infected with MOI 0.3 (**Fig 1C** and **1D**), indicating that trophoblasts from CZS-affected twins were

significantly more susceptible to ZIKV infection or at least more virus productive when compared

with trophoblasts from non-affected twins. Infection with the higher MOI (MOI = 3) obliterated this

difference due to the high susceptibility of trophoblasts in general to the Zika infection. In order to

have a better resolution of the differences between cells derived from CZS-affected and non-affected

twins, subsequent studies were performed with MOI of 0.3.

**Potential flavivirus attachment factors and IFN receptor genes were not differentially**

**expressed between trophoblasts from discordant twins**

To evaluate the possible differences in molecular signatures associated with ZIKV$^{BR}$ infection

in trophoblasts from CZS-affected and non-affected twins, we performed RNA-Seq analysis in

hiPSC-derived trophoblasts before and after *in vitro* infection with ZIKV$^{BR}$ for 96 h (MOI = 0.3).

With this approach, we could analyze possible differences in gene expression of potential attachment

factors for ZIKV and of genes related to antiviral response.

We first confirmed the efficiency of hiPSC differentiation into trophoblasts by looking at the

expression levels of a set of over100 genes which have been associated with the trophoblast lineage

of mammals [35,36]. When compared with hiPSCs, most of these genes showed significant

upregulation in trophoblasts after differentiation (**S1 Table**). We validated by RT-qPCR the

differential expression of *NANOG*, a hiPSC marker, and of *HCGA*, *HCGB* and *KRT7*, three of the

genes upregulated in the trophoblasts upon differentiation, as compared with the hiPSCs (**S1 Fig**).

Consistent with this and with previous experiments that used the same differentiation protocol [35],

proliferation-related genes were downregulated in the trophoblasts relative to the hiPSCs (**S2 Fig**). In

addition, four genes encoding transcription factors – *CDX2, ELF5, EOMES* and *ASCL1,* which are

generally regarded as markers of trophoblast stem cells [37] – were found to be barely expressed

(TPM < 1) in all six hiPSC-derived trophoblast samples from the twins analyzed here. This suggests

that all trophoblasts from all the twins differentiated beyond the trophoblast stem-cell stage.

We then looked for differences in the expression levels of potential flavivirus attachment

factor genes, of IFN genes and of IFN receptor genes between the trophoblasts from CZS-affected

and non-affected twins. These differences could be related to the greater susceptibility to ZIKV[BR]

infection of the trophoblasts from CZS-affected twins.

Many potential flavivirus attachment factors have been proposed as ZIKV candidate

receptors, including glycosaminoglycans (CD209/DC-SIGN and HSPG*2*) and TAM receptors

(TYRO3, AXL, MERTK) [38-40]. None of these receptor genes was found to be differentially

expressed between the trophoblasts from CZS-affected and non-affected twins (**S3 Fig**). *CD209/DC-*

*SIGN* gene was barely expressed in the hiPSC-derived trophoblasts either before or after ZIKV[BR]

infection (**S3 Fig**). *HSPG2* gene had the highest expression levels in hiPSC-derived trophoblasts

among the genes encoding potential ZIKV candidate receptors, however no differences in the

expression levels of *HSPG2* gene between trophoblasts from CZS-affected and non-affected twins

were found either before or after ZIKV[BR] infection (**S3 Fig**). Recently, AXL has been reported as the

receptor involved in ZIKV entry in many cell types, including neural stem cells [41], human

umbilical vein endothelial cells (HUVECs) [42], primary human astrocytes [43] and Sertoli cells

[44]. *AXL* gene expression showed a marked increase upon differentiation from hiPSC towards

trophoblasts (> 5-fold on average), but infected and non-infected hiPSC-derived trophoblasts from

non-affected or CZS-affected twins show similar expression levels of *AXL* gene (**S3 Fig**). TAM-

family receptors bind phosphatidylserine indirectly, through the soluble intermediates GAS6 (growth

arrest-specific 6) and PROS1 [45].  Recently, it was shown that ZIKV infects HUVECs much more

efficiently than other flaviviruses because it binds GAS6 more avidly, which in turn facilitates its

interaction with AXL [42]. Infected and non-infected hiPSC-derived trophoblasts from non-affected

or CZS-affected twins show similar expression levels of *GAS6* and *PROS1* genes (**S3 Fig**). HAVCR1

(TIM1; hepatitis C receptor) was shown to have an important role in the entry of ZIKV into primary

cell types from mid- and late-gestation placentas and explants from first-trimester chorionic villi

[26]. Interestingly, *HAVCR1* gene was barely expressed in all the hiPSC-derived trophoblasts (**S3**

**Fig**). Expression levels of all the above receptor genes measured here in non-infected hiPSC-derived

trophoblasts from the three pairs of twins confirmed previous data from ESC-derived non-infected

trophoblasts [35].

Production of interferons has been reported as a key step in the antiviral immune response to

ZIKV [25,46,47]. In agreement with Sheridan et al. [35], primitive trophoblasts expressed low levels

of mRNAs from all the IFN genes (**S4 Fig**) when compared with ZIKV potential attachment factors

(**S3 Fig**). Upon *in vitro* infection with ZIKV$^{BR}$, a marked induction of mRNA expression of *IFNB1*,

*IFNL1*, *IFNL2* and *IFNL3* genes but not of *IFNE* was observed (**S4 Fig**). Of note, IFN genes were

not differentially expressed between the trophoblasts from CZS-affected and non-affected twins (**S4**

**Fig**). *IFNA*, *IFNG, IFNK* and *IFNW1* genes were not detectable in any hiPSC-derived trophoblast

cells.

Interestingly, *IFNGR1* and *IFNGR2* (type II IFN receptor genes) were highly upregulated in

hiPSC-derived trophoblasts when compared with the hiPSCs that originated them (**S5 Fig**). These

two IFN receptor genes were not differentially expressed between trophoblasts from CZS-affected

and non-affected twins, and their expression was not changed upon ZIKV$^{BR}$ infection (**S5 Fig**).

Compared with the expression of type II IFNR genes, the expression levels of the other interferon

receptor genes, namely type I IFNR (*IFNAR1* and *IFNAR2*) and the first (*IFNLR1*) and second

(*IL10RB*) subunits of type III IFNR gene (*IFNL*) were lower in both hiPSCs and hiPSC-derived

trophoblasts (**S5 Fig**), and again none of them had their expression affected by ZIKV[BR].

**Induction of interferon-stimulated genes (ISGs) expression and of IFN secretion in hiPSC-**

**derived trophoblasts upon ZIKV[BR] infection**

It is known that interferons participate as mediators of the response to ZIKV infection in the

maternal-fetal interface [25,48,49] and that ESC-derived trophoblasts lack components of a robust

antiviral response system before infection with ZIKV [35]. However, it is not known if hiPSC-

derived primitive trophoblasts are able to induce an antiviral response after infection with ZIKV. To

determine this, we analyzed our RNA-Seq data and looked for differentially expressed genes (DEGs)

in the hiPSC-derived primitive trophoblasts after *in vitro* infection with ZIKV[BR].

An important gene expression response was induced in both hiPSC-derived trophoblasts from

CZS-affected and non-affected twins following *in vitro* infection with ZIKV[BR], and a set of 471

DEGs were upregulated in hiPSC-derived trophoblasts after infection with ZIKV[BR] (FDR < 0.05,

Edge R exact test) (**Fig 2A**). These genes (**S2 Table**) include interferon-stimulated genes (ISG) and

genes related to cytokine secretion. We found a significant (FDR < 0.05, cumulative hypergeometric

distribution) enrichment of different Gene Ontology (GO) terms among these 471 upregulated DEGs

detected in the RNA-Seq experiment, being the top categories "response to type I interferon",

"defense response to virus" and "negative regulation of viral process" (**Fig 2B, S3 Table**). Consistent

with the GO analysis, Ingenuity Pathway Analysis (IPA) also pointed to an enriched network of

interferon-stimulated genes upregulated in hiPSC-derived trophoblasts after ZIKV infection

(upregulated ISGs are colored in pink in **Fig 3A**).

We also quantified in the hiPSC-derived trophoblasts the secreted levels of type I (IFNA2),

type II (IFNG) and type III (IFNL1) IFNs produced by these cells in the absence of virus or after

infection with ZIKV[BR] and culture for 48 h (**Fig 3B, upper panel**) or 96 h (**Fig 3B, lower panel**).

After 48 h in culture without infection, hiPSC-derived trophoblasts from both CZS-affected and non-

affected twins were able to secrete IFNA2 and IFNG, whereas IFNL1 was not detectable. There was

no statistically significant increase in any IFN secretion at 48 h after ZIKV$^{BR}$ infection of

trophoblasts (**Fig 3B, upper panel**). After 96 h in culture, hiPSC-derived trophoblasts from both

CZS-affected and non-affected twins secreted higher levels of IFNA2, IFNG and IFNL1 compared

with those at 48 h, although high variability in the secreted IFN levels was observed among the

trophoblasts from different individuals (**Fig 3B, lower panel**). More importantly, a statistically

significant increase in the secretion of IFNL1 by trophoblasts from non-affected twins was observed

at 96 h after infection with ZIKV$^{BR}$, but not by trophoblasts from CZS-affected twins (**Fig 3B, lower**

**panel**).

Taken together, these results indicate that hiPSC-derived trophoblasts from both CZS-

affected and non-affected twins were able to respond to ZIKV$^{BR}$ infection by secreting IFNA2, IFNG

and IFNL1, which induced the upregulation of a set of ISG genes potentially involved in the response

to ZIKV$^{BR}$ infection. In addition, only trophoblasts from non-affected twins showed a significant

increase in secreted IFNL1 at 96 hpi with ZIKV$^{BR}$ (**Fig 3B, lower panel**).

**A set of genes involved with extracellular matrix organization and leukocyte activation was**

**downregulated in trophoblasts from CZS-affected twins after ZIKV$^{BR}$ infection**

We next investigated if there were differences in gene expression between trophoblasts from

CZS-affected and non-affected twins after ZIKV$^{BR}$ infection. In total, 44 genes were downregulated

after ZIKV$^{BR}$ infection in trophoblasts from CZS-affected when compared with non-affected twins

(FDR < 0.05, Edge R exact test) (**Fig 4A and S4 Table**). Different Gene Ontology (GO) terms were

found to be significantly (FDR < 0.05, cumulative hypergeometric distribution) enriched among

these 44 downregulated DEGs, including "extracellular matrix" and "regulation of leukocyte

activation" (**Fig 4B, S5 Table**). Significantly enriched Gene Ontology (GO) terms that were found

among the upregulated DEGs are shown in **S6 Fig** and **S5 Table**, and the most significantly enriched

categories are "amino acid biosynthetic process" and "acid secretion".

The levels of a panel of cytokines and chemokines secreted by the hiPSC-derived

trophoblasts were quantified in the supernatants of 48 h or 96 h cell cultures in the absence of virus

or after infection with ZIKV$^{BR}$. Interestingly, from all the analytes tested, the chemokines

RANTES/CCL5 and IP10 showed a consistent (both at 48 h and 96 h post-infection) and significant

increase in secretion by trophoblasts from non-affected twins after infection with ZIKV$^{BR}$, but not by

trophoblasts from CZS-affected twins (**Fig 4C**). RANTES/CCL5 secretion levels by trophoblasts

from non-affected twins increased 2.4- and 4.6-fold at 48 h or 96 h after infection with ZIKV$^{BR}$,

respectively, while IP10 secretion levels increased 16- and 96-fold (**Fig 4C**). The other tested

cytokines and chemokines did not show statistically significant differences in the levels produced by

infected trophoblasts from CZS-affected or non-affected twins.

**Discussion**

Here, we were able to analyze, for the first time, the *in vitro* viral susceptibility and the gene

expression patterns after *in vitro* ZIKV$^{BR}$ infection of hiPSC-induced trophoblasts from dizygotic

(DZ) twins discordant for the presence of microcephaly. Because these are DZ twins whose mothers

were infected with ZIKV$^{BR}$ during pregnancy, the two fetuses in each of the three twin pairs were

exposed to the virus at the same time, thus representing a rare and unique cohort to test whether the

host genetic background plays any role in determining CZS outcome. Indeed, we have previously

shown that hiPSC-induced NPCs from the same subjects exhibit differential *in vitro* ZIKV$^{BR}$

susceptibility, as NPCs from CZS-affected twins had significantly higher ZIKV$^{BR}$ replication and

reduced cell growth when compared with NPCs from non-affected twins [32]. No rare Mendelic

potentially pathogenic variant was assigned in this cohort that could explain NPCs susceptibility to

CZS [32], suggesting that CZS caused by ZIKV may be a multifactorial disorder. Herein, using a

well-established model that recapitulates the primitive placenta formed during fetal implantation

[34,35], we sought to investigate if hiPSC-induced trophoblasts from the same twins also show

differential responses to *in vitro* ZIKV[BR] infection.

Many studies have established causal effects between ZIKV infection during pregnancy and

microcephaly developed in the fetus [15], using *in vitro* as well as mice and non-human primates *in*

*vivo* models [50-54]. Cells localized in the epidermis and dermis were primarily considered as

targets for ZIKV infection [55], as mosquito bite remains the major transmission route [56]. ZIKV

infection exhibits broad distribution and persistence in body tissues and fluids, and the presence of

ZIKV in the amniotic fluid of pregnant women and in semen [57-61] suggests the additional

possibility of sexual and perinatal transmissions. Trophectoderm cells of pre-implantation human

embryos can be infected with ZIKV [62]. Also, endometrial stromal cells are highly permissive to

ZIKV infection, and likely represent a crucial cell target of ZIKV reaching them, either via the

uterine vasculature in the viremic phase of the infection or by sexual viral transmission, being a

potential source of virus spreading to placental trophoblasts during pregnancy [63]. Indeed, recent

studies suggest that the placenta is the key mediator for vertical transmission of ZIKV from infected

mothers to fetal brains; several placental cells have been shown to be infected by ZIKV, including

placental macrophages, trophoblasts and fibroblasts of the maternal *decidua basalis* [25-29,35,58].

Here, we show that hiPSC-induced trophoblasts from DZ twins discordant for CZS were

differentially susceptible to infection with ZIKV[BR]. Interestingly, 96 hpi with ZIKV[BR], virus titers in

culture supernatants were significantly higher in the CZS-affected twins' trophoblasts when

compared with non-affected (**Fig 1C**), indicating that ZIKV[BR] can replicate more efficiently in the

trophoblasts from CZS-affected twins, potentially facilitating virus dissemination into fetal tissues.

Noteworthy, our RNA-seq results indicate that hiPSC-derived trophoblasts express the ZIKV

candidate receptor genes (*HSPG2* and *TAM* receptor genes) and the IFN receptor genes, and at lower

levels the IFN genes; however none of these genes were differentially expressed between the

trophoblasts from CZS-affected and non-affected twins either before or after ZIKV[BR] infection.

295             Interestingly, ZIKV[BR] infection caused a significant increase in IFNL1 secretion by

trophoblasts from non-affected twins (**Fig 3B**), whereas in trophoblasts from CZS-affected twins no

significant increase was observed (**Fig 3B**). IFNL1 is a type III IFN constitutively released by

primary human trophoblasts from full-term placentas [25], which are known to be refractory to ZIKV

infection [25,35]. The increased secretion of IFNL1 by trophoblasts from non-affected twins, already

at 48 hpi by ZIKV[BR], but especially at 96 hpi, may protect against ZIKV[BR] dissemination, as shown

in the female reproductive tract of mice [47,64,65]. Our results indicate a possible role of IFNL1 in

the control of ZIKV infection by primitive trophoblasts. In addition, after ZIKV[BR] infection,

trophoblasts from the non-affected twins were able to significantly induce the secretion of immune

mediator chemokines RANTES/CCL5 [66,67] and IP10 [68], while trophoblasts from the CZS-

affected twins were not (**Fig 4C**).  RANTES/CCL5 promotes trophoblasts cell migration and can

recruit immune cells [69-71], whereas IP10 plays a role as a chemotactic molecule implicated in the

migration of trophoblast cells [72]. Thus, lower secretion of RANTES/CCL5 and IP10 indicates that

CZS-affected twins' trophoblasts have a lower ability of migration, immune cell recruitment and

viral control.

310             The placenta plays a critical role in immunological protection and can undergo major

structural and functional adaptations in order to protect the fetus from environmental stressors

[48,49]. When the placental function is impaired [73], the intrauterine environment might be

perturbed and the placental defenses involved in fetal protection compromised. In the RNA-Seq

analysis, we found 79 differentially expressed genes in common among the three CZS-affected

compared with non-affected twins, 44 of which were downregulated after ZIKV[BR] infection in the

trophoblasts from CZS-affected compared with non-affected twins. The top GO category associated

with these downregulated differentially expressed genes is "extracellular matrix". It is known that for

successful fetus development, one of the critical steps is proper invasion of the maternal decidua by

trophoblasts [74] and that many molecules, including galectins, are involved in this process [75,76].

One of the downregulated genes in the trophoblasts from CZS-affected twins after ZIKV[BR]

infection was *COL3A1* (**Fig 4A**). Collagens are the main structural proteins in the extracellular

matrix, and they have been related to pregnancy and/or placental pathological conditions including

gestational diabetes and pre-eclampsia [77,78]. *ITGA1*, another gene downregulated in trophoblasts

from CZS-affected twins after ZIKV[BR] infection, interacts with the extracellular matrix, particularly

with collagen and laminin [79].  Importantly, extravillous trophoblasts (EVT) have been shown to

express ITGA1 as they invade from the anchoring villi deeply into the maternal endometrium and

myometrium in weeks 8–13 of gestation [80,81]. EVTs fail to express ITGA1 in preeclampsia, which

is associated with both poor trophoblast invasion and oxidative stress [81,82]. In addition, *LGALS3*

gene (encoding galectin-3) was downregulated in trophoblasts from CZS-affected twins after ZIKV[BR]

infection. LGALS3 is a galectin that has been described as involved in the process of trophoblast cell

migration and invasion, significant for human embryo implantation [83,84]. Overall, downregulation

of genes involved in trophoblast adaptation to the intrauterine environment, including *ITGA1,*

*COL3A1* and *LGALS3* (**S7 Fig**)*,* in the trophoblasts of CZS-affected twins after ZIKV[BR] infection,

may affect trophoblast migration, implantation, homeostasis and possibly impair the control of

ZIKV[BR] infection. Noteworthy, all the above differentially expressed genes were found in common

among the three CZS-affected compared with non-affected twins, in spite of the genetic background

variability of our cohort. Importantly, as all experiments were done under the same conditions, these

differences in genetic background underlie both possible differences in progression of trophoblasts

differentiation and the resulting difference in susceptibility to ZIKV infection. Thus, the set of 79

differentially expressed genes found here represents the minimum core of genes significantly altered

in common among the three pairs of twins. This study provides genetic targets to be further explored

as possible infection susceptibility factors in the placenta.

Our gene expression results indicate that ZIKV[BR] infection has caused, only in the

trophoblasts from CZS-affected twins, the downregulation of genes important for trophoblast

adhesion as well as immune response activation. Noteworthy, when gene expression differences

between the NPCs from these non-affected and CZS-affected twins were analyzed [32], another set

of 64 genes was found to be differentially expressed, including *FOXG1* and *LHX2*, two transcription

factors important for neural development [85,86], which were down-regulated in NPCs from CZS-

affected compared with non-affected twins [32]. Overall, development of congenital Zika syndrome

might result, among other factors, from a concomitant decreased ability of the placenta to respond to

ZIKV infection in the CZS-affected neonates, along with a deregulation of neural development genes

in ZIKV-infected NPCs of these CZS-affected neonates.

Based on our gene expression analyses, we suggest that the ability to respond more efficiently

to ZIKV infection in the placenta may be a key parameter to predict the success of ZIKV

dissemination into fetuses' tissues. Moreover, further understanding of the participation of immune

mediators, such as the chemokines RANTES/CCL5 and IP10, in the trophoblast response to ZIKV

infection may open a path for drug development or repurposing to possibly inhibit viral replication or

avoid viral dissemination into fetus' tissues.

**Materials and methods**

**Human subjects**

Three pairs of DZ-D (dichorionic and diamniotic) twins discordant for the presence of

microcephaly (#10608-1 and #10608-4; #10763-1 and #10763-4; #10788-1 and #10788-4) whose

peripheral blood mononuclear cells had been previously collected and isolated [32] were examined in

this study. Zygosity had been previously confirmed by whole-exome sequencing (WES) and by

microsatellite analysis [32]. All babies were born from mothers negative for previous STORCH

infections and in each affected baby (#10608-1, #10763-1 and #10788-1) the head circumference was

three standard deviations (SD) below the mean for the given age, sex, and gestation stage at birth

[32,87]. Diagnosis of microcephaly due to ZIKV infection (CZS) in all the three affected subjects

was confirmed by neuroimaging, serology and by the mother reporting ZIKV infection symptoms

during pregnancy (**S6 Table** and Caires-Junior et al. [32]). The protocol used in this study was

approved by the Human Research Ethics Committee from Biosciences Institute, University of São

Paulo (protocol # 184/2016). All mothers gave written informed consent in accordance with the

Declaration of Helsinki.

**Cell lines and maintenance of hiPSCs**

hiPSCs were generated according to Caires-Junior et al. [32]. Briefly, CD71+ cells were

isolated from the three pairs of DZ-D twins' peripheral blood samples (#10608-1 and #10608-4;

#10763-1 and #10763-4; #10788-1 and #10788-4). CD71-positive cells were sorted using magnetic

labeled antibody (Miltenyi) following the manufacturer's instructions. All hiPSC lines were tested

for ZIKV infection by RT-qPCR using primers described in **S7 Table** and the results were negative.

The reprogramming protocol was performed with episomal vectors system (Addgene plasmids

27077, 27078 and 27080), and using the Amaxa human CD34+ cells Nucleofection Kit (Lonza),

following the manufacturer's recommendations. Three days after nucleoporation, cells were seeded

on irradiated murine embryonic fibroblasts (Millipore, A24903) in embryonic stem cell (ESC)

medium (Dulbecco's modified Eagle's medium (DMEM)/F12 supplemented with 2 mM GlutaMAX-

I, 0.1 mM non-essential amino acids, 100 μM 2-mercaptoethanol, 20% knockout serum replacement

(all provided by Life Technologies), 10 ng/mL bFGF (Peprotech), 0.25 mM NaB, 0.5 mM VPA, 2

μM thiazovivin, 0.5 μM PD 0325901 and 2 μM SB 431542; all provided by Tocris Bioscience). The

typical hiPSC colonies were transferred to hESC-qualified Matrigel (Corning)-coated 60 mm petri

dishes (Corning) and cultured in Essential 8 Medium (Gibco) with 100 μg/mL normocin

(InvivoGen). All derived cell lines were checked for mycoplasma contamination periodically.

**Differentiation of human iPSCs into trophoblasts**

Differentiation of human iPSCs into trophoblasts was performed according to Amita et al.

[34]. Briefly, hiPSCs were seeded at 15,000 cells/cm$^2$ on Matrigel (Corning) coated plates and

cultured in mTESR1 medium (Stem Cell Technologies) for 3 days with daily medium changes. Next,

basal medium was changed to hES medium supplemented with 4 ng/mL FGF-2 for 24 h. For the next

9 days, hES-BAP medium was used, which consisted of hES medium supplemented with BMP4 (10

ng/mL), A83-01 (1 μM) and PD173074 (0.1 μM), with daily medium changes. hES medium was

composed of DMEM/F-12 supplemented with 20 % KO-serum replacement, 1 % NEAA, 1 %

glutamine, 0.1 mM 2-mercaptoethanol, all from Gibco.

**In situ immunofluorescence**

hiPSCs and trophoblast cultures were fixed with 4 % PFA followed by permeabilization with

0.01 % Triton X-100 and then blocked with 5 % bovine serum albumin (BSA) for 1 h. The cells were

incubated overnight with primary antibodies at 4 °C (Anti-hCG beta/β-CG/hCGB3, Abcam #ab53087

and Anti-Cytokeratin 7/CK7/KRT7, Abcam #ab9021), washed with PBS and subsequently incubated

with secondary fluorescent antibodies for 1 h at room temperature. Finally, cells were stained with

DAPI (4',6-diamidino-2-phenylindole) and or Phalloidin for 2 min at room temperature. Confocal

analysis was performed using Zeiss LSM 800.

**Zika virus**

410       ZIKV[BR] was a courtesy of Dr. Pedro Vasconcelos [88], Instituto Evandro Chagas, Brazil.

411    Viral stock was established after viral propagation for two serial passages in VERO cells (ATCC®

412    CCL-81™) in serum-free medium (VP SFM, Thermo scientific).

**Infection of the hiPSC-derived trophoblasts**

414       hiPSC-derived day-9 trophoblasts were seeded into 6-well or 24-well plates (Corning) and 2-

415    well chamber slides (Nunc; Thermo Fisher Scientific) to a confluence of $3 \times 10^4$ cells/cm$^2$.

416    Trophoblasts (monolayer) were exposed to ZIKV[BR] (MOI: 0.3, 3 and Mock). Monolayer cells were

417    exposed to the virus for 1 h at 37 °C and 5 % $CO_2$ (g), washed with hES-BAP medium, and then

418    maintained by up to 96 h (end point).

**Measurement of viral burden**

For ZIKV titration, plaque assay was performed with the supernatants of cell cultures. For

plaque assay, an amount of $6 \times 10^4$ VERO cells/well were seeded in 24-well plates 48 h before the

assay. Samples were serially diluted in DMEM culture medium from $10^{-1}$ to $10^{-6}$, applied in

duplicates of 100 µL to each well, and incubated for 30 min at 37 °C. After virus adsorption, wells

were overlaid with culture medium containing carboxymethyl cellulose (1 %) and incubated at 37 °C.

After 5 days, plates were drained, washed with PBS, and stained with 0.1 % naphthol blue-black, 1.6

% sodium acetate in 6 % glacial acetic acid for 30 min. Plaque formation units were visually

determined in the most appropriate viral dilution and expressed as PFU/mL.

**RNA-Seq assay**

Total RNA from hiPSC-derived trophoblasts was extracted using the RNeasy Micro Kit

(Qiagen, 74004), treated with TURBO DNase (Ambion, AM2238) for 30 min at 37 °C, and then re-

purified with the Qiagen RNeasy Micro Kit. RNA samples were quantified using the Qubit RNA HS

Assay Kit (Thermo Fisher Scientific, Q32852); purity was evaluated using NanoDrop ND-1000

Spectrophotometer (NanoDrop Technologies) and the integrity was verified using the Agilent RNA

6000 Pico Kit (Agilent Technologies, 5067-1513) in the 2100 Bioanalyzer Instrument (Agilent

Technologies). Stranded tagged cDNA libraries were prepared using the KAPA Stranded mRNA-Seq

Kit (Illumina, KK8421) and cluster generation was performed using the Illumina HiSeq 4000 PE

Cluster Kit (Illumina, PE-410-1001). Tagged libraries were pooled and sequenced (300 cycles,

paired-end sequencing) in the Illumina HiSeq 4000 instrument using a HiSeq 4000 SBS Kit

(Illumina, FC-410- 1003). Raw reads were preprocessed using the standard Illumina pipeline to

segregate multiplexed reads.

**RNA-seq data processing and analysis**

Paired-end adapters and low quality reads were removed by fastp version 0.20.0 [89] using

the  quality-filtering parameters -l 20 -5 3 -3 3. Filtered paired-end reads were mapped and quantified

at gene level by the STAR-RSEM pipeline [90,91], using an index made from GRCh38.p12

GencodeV.28 [92].  All differential expression (DE) analyses used the bioconductor package edgeR

[93]. To call differentially expressed genes a general linear models (glm) was fitted, and likelihood

ratio tests (lrt) were performed using twins' covariates as blocking groups in all cases. P-values were

adjusted using FDR, all genes with FDR lower than 0.05 were considered differentially expressed

genes. To identify up-regulated or down-regulated genes, the logCPM data from edgeR were used.

Heatmaps were plotted using the R package pheatmap. Enriched gene ontology analysis was perform

using the bioconductor package clusterProfiler [94], based on the annotation of the bioconductor

package org.Hs.eg.db. All plots were generated with the R packages gg plot2, cowplot and the

bioconductor package DOSE.

**Reverse transcription - quantitative PCR (RT-qPCR)**

Total RNA from hiPSCs and hiPSC-derived trophoblasts was extracted as described above.

The reverse transcription (RT) reaction was performed with 150 ng of each total RNA sample using

the SuperScript IV First-Strand Synthesis System (Life Technologies, cat. #18091050) and random

hexamer primers in a 20 μL final volume. The obtained cDNAs were diluted 10 times in water and

quantitative PCR was performed using 2.5 μL of each diluted cDNA in a total volume of 10 μL

containing 1× LightCycler 480 SYBR Green I Master Mix (Roche Diagnostics, cat. #04707516001)

and 800 nM of each primer in a LightCycler 480 System (Roche Diagnostics). RT-qPCR was run in

two biological replicates with three technical replicates each and primers are shown in **S7 Table**. The

GAPDH gene (NM_002046) was used as the reference for internal normalization.

**Cytokine and chemokine protein quantification**

Supernatants from trophoblasts' cell cultures were collected after 48 h or 96 h of infection

with ZIKV[BR] and then evaluated for 33 cytokines/chemokines for the following analytes: IFNalfa2,

IFNbeta, IFNg, IFN-lambda1/ IL-29, IFN-lambda 2/ IL28a, IL6, IL10, IL12p40, IL12p70, IL13,

IL15, IL17A, IL1RA, IL1a, IL1b, IL2, IL3, IL4, IL5, IL7, IL8, IP10, MCP1, MIP1a, MIP1b,

RANTES, TNFa, TNFb, VEGF, EGF, EOTAXIN, GCSF and GMCSF. The Human

Cytokine/Chemokine Magnetic Bead Panel kit (Millipore) was used according to the manufacturer's

recommendations. Samples were analyzed in the MagPix instrument (Luminex) and the data was

analyzed with the Milliplex Analyst software (Millipore).

**Quantification and statistical analysis**

Two-tailed unpaired t test was used for pairwise comparisons. Graphpad Prism software was

used to perform statistical analysis (version 7.0). Quantification of data are represented as mean ±

SEM and p-value threshold was * 0.05, ** 0.01, *** 0.001 and **** 0.0001.

**Data availability**

The RNA-Seq data that support the findings of this study have been deposited in the

Sequence Read Archive (SRA) NCBI repository under Accession number PRJNA565997. All other

data supporting the findings of this study are available within the article and its Supporting

Information files or are available from the authors upon request.

**Acknowledgments**

We thank Dr. R. Michael Roberts and Dr. Toshihiko Ezashi, University of Missouri, Columbia,

MO, USA for advice on trophoblast cells cultivation protocols. We thank Dr. J.C. Setubal for access

to the computational facilities of the Bioinformatics Laboratory of Instituto de Química,

Universidade de São Paulo, São Paulo, Brazil.

**Author contributions**

Conceptualization: Murilo Sena Amaral, Sergio Verjovski-Almeida

Data Curation: Murilo Sena Amaral, David Abraham Morales-Vicente, Sergio Verjovski-Almeida

Formal Analysis: Murilo Sena Amaral, Ernesto Goulart, Luiz Carlos Caires-Júnior, Alessandra

Soares-Schanoski, Jorge E. Kalil, Mayana Zatz, Sergio Verjovski-Almeida

Funding Acquisition: Mayana Zatz, Sergio Verjovski-Almeida

Investigation: Murilo Sena Amaral, Ernesto Goulart, Luiz Carlos Caires-Júnior, Alessandra Soares-

Schanoski, Roselane Paiva Gomes, Giovanna Gonçalves de Oliveira Olberg

Methodology:  Murilo Sena Amaral, Ernesto Goulart, Luiz Carlos Caires-Júnior, David Abraham

Morales-Vicente, Alessandra Soares-Schanoski, Sergio Verjovski-Almeida

Project Administration: Mayana Zatz, Sergio Verjovski-Almeida

Resources: Renato Mancini Astray, Mayana Zatz, Sergio Verjovski-Almeida

Software: David Abraham Morales-Vicente

Supervision:  Jorge E. Kalil, Mayana Zatz, Sergio Verjovski-Almeida

Validation: Murilo Sena Amaral, Ernesto Goulart, Luiz Carlos Caires-Júnior, David Abraham

Morales-Vicente

Visualization: Murilo Sena Amaral, David Abraham Morales-Vicente

Writing – Original Draft Preparation: Murilo Sena Amaral, Sergio Verjovski-Almeida

Writing – Review & Editing: Murilo Sena Amaral, Ernesto Goulart, Luiz Carlos Caires-Júnior,

Alessandra Soares-Schanoski, Renato Mancini Astray, Jorge E. Kalil, Mayana Zatz, Sergio

Verjovski-Almeida

## Competing interests

510    The authors have declared that no competing interests exist.

## Financial disclosure statement

This work was supported by grants from Fundação de Amparo à Pesquisa do Estado de São

Paulo (FAPESP) (Thematic grant number 2014/03620-2 and 2018/23693-5 to SVA and CEPID

number 2013/08028-1 and INCT number 14/50931-3 to MZ) and from Conselho Nacional de

Desenvolvimento Científico e Tecnológico (CNPq) INCT number 465355/2014-5 to MZ. DAMV

received a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES),

Brazil – Finance Code 001; LCCJ is a fellow of FAPESP (2017/16283-2); EG is a fellow of FAPESP

(2015/14821-1); SVA laboratory was also supported by institutional funds from Fundação Butantan;

SVA received an established investigator fellowship award from CNPq, Brazil. The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

## References

1. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, et al. (2009) Zika virus outbreak on Yap Island, Federated States of Micronesia. N Engl J Med 360: 2536-2543.

2. Cao-Lormeau VM, Roche C, Teissier A, Robin E, Berry AL, et al. (2014) Zika virus, French polynesia, South pacific, 2013. Emerg Infect Dis 20: 1085-1086.

3. Johansson MA, Mier-y-Teran-Romero L, Reefhuis J, Gilboa SM, Hills SL (2016) Zika and the Risk of Microcephaly. N Engl J Med 375: 1-4.

4. Liu Y, Liu J, Du S, Shan C, Nie K, et al. (2017) Evolutionary enhancement of Zika virus infectivity in Aedes aegypti mosquitoes. Nature 545: 482-486.

5. Yuan L, Huang XY, Liu ZY, Zhang F, Zhu XL, et al. (2017) A single mutation in the prM protein of Zika virus contributes to fetal microcephaly. Science 358: 933-936.

6. Rossi SL, Ebel GD, Shan C, Shi PY, Vasilakis N (2018) Did Zika Virus Mutate to Cause Severe Outbreaks? Trends Microbiol 26: 877-885.

7. Ferraris P, Yssel H, Misse D (2019) Zika virus infection: an update. Microbes Infect.

8. Pierson TC, Diamond MS (2018) The emergence of Zika virus and its new clinical syndromes. Nature 560: 573-581.

9. Triunfol M (2016) Microcephaly in Brazil: confidence builds in Zika connection. Lancet Infect Dis 16: 527-528.

10. Schuler-Faccini L, Ribeiro EM, Feitosa IM, Horovitz DD, Cavalcanti DP, et al. (2016) Possible Association Between Zika Virus Infection and Microcephaly - Brazil, 2015. MMWR Morb Mortal Wkly Rep 65: 59-62.

11. Chimelli L, Melo ASO, Avvad-Portari E, Wiley CA, Camacho AHS, et al. (2017) The spectrum of neuropathological changes associated with congenital Zika virus infection. Acta Neuropathol 133: 983-999.

12. Del Campo M, Feitosa IM, Ribeiro EM, Horovitz DD, Pessoa AL, et al. (2017) The phenotypic spectrum of congenital Zika syndrome. Am J Med Genet A 173: 841-857.

13. Franca GV, Schuler-Faccini L, Oliveira WK, Henriques CM, Carmo EH, et al. (2016) Congenital Zika virus syndrome in Brazil: a case series of the first 1501 livebirths with complete investigation. Lancet 388: 891-897.

14. Wen Z, Song H, Ming GL (2017) How does Zika virus cause microcephaly? Genes Dev 31: 849-861.

15. Rasmussen SA, Jamieson DJ, Honein MA, Petersen LR (2016) Zika Virus and Birth Defects--Reviewing the Evidence for Causality. N Engl J Med 374: 1981-1987.

16. Mlakar J, Korva M, Tul N, Popovic M, Poljsak-Prijatelj M, et al. (2016) Zika Virus Associated with Microcephaly. N Engl J Med 374: 951-958.

17. Morris G, Barichello T, Stubbs B, Kohler CA, Carvalho AF, et al. (2018) Zika Virus as an Emerging Neuropathogen: Mechanisms of Neurovirulence and Neuro-Immune Interactions. Mol Neurobiol 55: 4160-4184.

18. Li H, Saucedo-Cuevas L, Shresta S, Gleeson JG (2016) The Neurobiology of Zika Virus. Neuron 92: 949-958.

19. Miner JJ, Diamond MS (2017) Zika Virus Pathogenesis and Tissue Tropism. Cell Host Microbe 21: 134-142.

20. Christian KM, Song H, Ming GL (2019) Pathophysiology and Mechanisms of Zika Virus Infection in the Nervous System. Annu Rev Neurosci 42: 249-269.

21. Figueiredo CP, Barros-Aragao FGQ, Neris RLS, Frost PS, Soares C, et al. (2019) Zika virus replicates in adult human brain tissue and impairs synapses and memory in mice. Nat Commun 10: 3890.

22. Tiwari SK, Dang J, Qin Y, Lichinchi G, Bansal V, et al. (2017) Zika virus infection reprograms global transcription of host cells to allow sustained infection. Emerg Microbes Infect 6: e24.

23. Arora N, Sadovsky Y, Dermody TS, Coyne CB (2017) Microbial Vertical Transmission during Human Pregnancy. Cell Host Microbe 21: 561-567.

24. Khaiboullina SF, Ribeiro FM, Uppal T, Martynova EV, Rizvanov AA, et al. (2019) Zika Virus Transmission Through Blood Tissue Barriers. Front Microbiol 10: 1465.

25. Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, et al. (2016) Type III Interferons Produced by Human Placental Trophoblasts Confer Protection against Zika Virus Infection. Cell Host Microbe 19: 705-712.

26. Tabata T, Petitt M, Puerta-Guardo H, Michlmayr D, Wang C, et al. (2016) Zika Virus Targets Different Primary Human Placental Cells, Suggesting Two Routes for Vertical Transmission. Cell Host Microbe 20: 155-166.

27. Quicke KM, Bowen JR, Johnson EL, McDonald CE, Ma H, et al. (2016) Zika Virus Infects Human Placental Macrophages. Cell Host Microbe 20: 83-90.

28. Aagaard KM, Lahon A, Suter MA, Arya RP, Seferovic MD, et al. (2017) Primary Human Placental Trophoblasts are Permissive for Zika Virus (ZIKV) Replication. Sci Rep 7: 41389.

29. Weisblum Y, Oiknine-Djian E, Vorontsov OM, Haimov-Kochman R, Zakay-Rones Z, et al. (2017) Zika Virus Infects Early- and Midgestation Human Maternal Decidual Tissues, Inducing Distinct Innate Tissue Responses in the Maternal-Fetal Interface. J Virol 91: e01905-01916.

30. Brasil P, Pereira JP, Jr., Moreira ME, Ribeiro Nogueira RM, Damasceno L, et al. (2016) Zika Virus Infection in Pregnant Women in Rio de Janeiro. N Engl J Med 375: 2321-2334.

31. Shapiro-Mendoza CK, Rice ME, Galang RR, Fulton AC, VanMaldeghem K, et al. (2017) Pregnancy Outcomes After Maternal Zika Virus Infection During Pregnancy - U.S. Territories, January 1, 2016-April 25, 2017. MMWR Morb Mortal Wkly Rep 66: 615-621.

32. Caires-Junior LC, Goulart E, Melo US, Araujo BHS, Alvizi L, et al. (2018) Discordant congenital Zika syndrome twins show differential in vitro viral susceptibility of neural progenitor cells. Nat Commun 9: 475.

33. McGrath EL, Rossi SL, Gao J, Widen SG, Grant AC, et al. (2017) Differential Responses of Human Fetal Brain Neural Stem Cells to Zika Virus Infection. Stem Cell Reports 8: 715-727.

34. Amita M, Adachi K, Alexenko AP, Sinha S, Schust DJ, et al. (2013) Complete and unidirectional conversion of human embryonic stem cells to trophoblast by BMP4. Proc Natl Acad Sci U S A 110: E1212-1221.

35. Sheridan MA, Yunusov D, Balaraman V, Alexenko AP, Yabe S, et al. (2017) Vulnerability of
primitive human placental trophoblast to Zika virus. Proc Natl Acad Sci U S A 114: E1587-
E1596.

36. Lee CQ, Gardner L, Turco M, Zhao N, Murray MJ, et al. (2016) What Is Trophoblast? A
Combination of Criteria Define Human First-Trimester Trophoblast. Stem Cell Reports 6:
257-272.

37. Roberts RM, Fisher SJ (2011) Trophoblast stem cells. Biol Reprod 84: 412-421.

38. Perera-Lecoin M, Meertens L, Carnec X, Amara A (2013) Flavivirus entry receptors: an update.
Viruses 6: 69-88.

39. Lee I, Bos S, Li G, Wang S, Gadea G, et al. (2018) Probing Molecular Insights into Zika Virus(-
)Host Interactions. Viruses 10: 233.

40. Hamel R, Dejarnac O, Wichit S, Ekchariyawat P, Neyret A, et al. (2015) Biology of Zika Virus
Infection in Human Skin Cells. J Virol 89: 8880-8896.

41. Nowakowski TJ, Pollen AA, Di Lullo E, Sandoval-Espinosa C, Bershteyn M, et al. (2016)
Expression Analysis Highlights AXL as a Candidate Zika Virus Entry Receptor in Neural
Stem Cells. Cell Stem Cell 18: 591-596.

42. Richard AS, Shim BS, Kwon YC, Zhang R, Otsuka Y, et al. (2017) AXL-dependent infection of
human fetal endothelial cells distinguishes Zika virus from other pathogenic flaviviruses. Proc
Natl Acad Sci U S A 114: 2024-2029.

43. Ojha CR, Rodriguez M, Karuppan MKM, Lapierre J, Kashanchi F, et al. (2019) Toll-like receptor
3 regulates Zika virus infection and associated host inflammatory response in primary human
astrocytes. PLoS One 14: e0208543.

44. Strange DP, Jiyarom B, Pourhabibi Zarandi N, Xie X, Baker C, et al. (2019) Axl Promotes Zika
Virus Entry and Modulates the Antiviral State of Human Sertoli Cells. MBio 10.

45. Stitt TN, Conn G, Gore M, Lai C, Bruno J, et al. (1995) The anticoagulation factor protein S and
its relative, Gas6, are ligands for the Tyro 3/Axl family of receptor tyrosine kinases. Cell 80:
661-670.

46. Pardy RD, Valbon SF, Richer MJ (2019) Running interference: Interplay between Zika virus and
the host interferon response. Cytokine 119: 7-15.

47. Caine EA, Scheaffer SM, Arora N, Zaitsev K, Artyomov MN, et al. (2019) Interferon lambda
protects the female reproductive tract against Zika virus infection. Nat Commun 10: 280.

48. Yockey LJ, Iwasaki A (2018) Interferons and Proinflammatory Cytokines in Pregnancy and Fetal
Development. Immunity 49: 397-412.

49. Ander SE, Diamond MS, Coyne CB (2019) Immune responses at the maternal-fetal interface. Sci
Immunol 4.

50. Wu KY, Zuo GL, Li XF, Ye Q, Deng YQ, et al. (2016) Vertical transmission of Zika virus
targeting the radial glial cells affects cortex development of offspring mice. Cell Res 26: 645-
654.

51. Miner JJ, Cao B, Govero J, Smith AM, Fernandez E, et al. (2016) Zika Virus Infection during
Pregnancy in Mice Causes Placental Damage and Fetal Demise. Cell 165: 1081-1091.

52. Hirsch AJ, Smith JL, Haese NN, Broeckel RM, Parkins CJ, et al. (2017) Zika Virus infection of rhesus macaques leads to viral persistence in multiple tissues. PLoS Pathog 13: e1006219.

53. Dudley DM, Aliota MT, Mohr EL, Weiler AM, Lehrer-Brey G, et al. (2016) A rhesus macaque model of Asian-lineage Zika virus infection. Nat Commun 7: 12204.

54. Tang H, Hammack C, Ogden SC, Wen Z, Qian X, et al. (2016) Zika Virus Infects Human Cortical Neural Progenitors and Attenuates Their Growth. Cell Stem Cell 18: 587-590.

55. Kim JA, Seong RK, Son SW, Shin OS (2019) Insights into ZIKV-Mediated Innate Immune Responses in Human Dermal Fibroblasts and Epidermal Keratinocytes. J Invest Dermatol 139: 391-399.

56. Lee JK, Shin OS (2019) Advances in Zika Virus(-)Host Cell Interaction: Current Knowledge and Future Perspectives. Int J Mol Sci 20: 1101.

57. Counotte MJ, Kim CR, Wang J, Bernstein K, Deal CD, et al. (2018) Sexual transmission of Zika virus and other flaviviruses: A living systematic review. PLoS Med 15: e1002611.

58. El Costa H, Gouilly J, Mansuy JM, Chen Q, Levy C, et al. (2016) ZIKA virus reveals broad tissue and cell tropism during the first trimester of pregnancy. Sci Rep 6: 35296.

59. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, et al. (2011) Probable non-vector-borne transmission of Zika virus, Colorado, USA. Emerg Infect Dis 17: 880-882.

60. D'Ortenzio E, Matheron S, Yazdanpanah Y, de Lamballerie X, Hubert B, et al. (2016) Evidence of Sexual Transmission of Zika Virus. N Engl J Med 374: 2195-2198.

61. Calvet G, Aguiar RS, Melo ASO, Sampaio SA, de Filippis I, et al. (2016) Detection and sequencing of Zika virus from amniotic fluid of fetuses with microcephaly in Brazil: a case study. Lancet Infect Dis 16: 653-660.

62. Tan L, Lacko LA, Zhou T, Tomoiaga D, Hurtado R, et al. (2019) Pre- and peri-implantation Zika virus infection impairs fetal development by targeting trophectoderm cells. Nat Commun 10: 4155.

63. Pagani I, Ghezzi S, Ulisse A, Rubio A, Turrini F, et al. (2017) Human Endometrial Stromal Cells Are Highly Permissive To Productive Infection by Zika Virus. Sci Rep 7: 44286.

64. Wells AI, Coyne CB (2018) Type III Interferons in Antiviral Defenses at Barrier Surfaces. Trends Immunol 39: 848-858.

65. Chen J, Liang Y, Yi P, Xu L, Hawkins HK, et al. (2017) Outcomes of Congenital Zika Disease Depend on Timing of Infection and Maternal-Fetal Interferon Action. Cell Rep 21: 1588-1599.

66. Levy JA (2009) The unexpected pleiotropic activities of RANTES. J Immunol 182: 3945-3946.

67. Appay V, Rowland-Jones SL (2001) RANTES: a versatile and controversial chemokine. Trends Immunol 22: 83-87.

68. Vazirinejad R, Ahmadi Z, Kazemi Arababadi M, Hassanshahi G, Kennedy D (2014) The biological functions, structure and sources of CXCL10 and its outstanding part in the pathophysiology of multiple sclerosis. Neuroimmunomodulation 21: 322-330.

69. Arima K, Nasu K, Narahara H, Fujisawa K, Matsui N, et al. (2000) Effects of lipopolysaccharide
and cytokines on production of RANTES by cultured human endometrial stromal cells. Mol
Hum Reprod 6: 246-251.

70. Fujiwara H, Higuchi T, Sato Y, Nishioka Y, Zeng BX, et al. (2005) Regulation of human
extravillous trophoblast function by membrane-bound peptidases. Biochim Biophys Acta
1751: 26-32.

71. Kayisli UA, Mahutte NG, Arici A (2002) Uterine chemokines in reproductive physiology and
pathology. Am J Reprod Immunol 47: 213-221.

72. Dominguez F, Martinez S, Quinonero A, Loro F, Horcajadas JA, et al. (2008) CXCL10 and IL-6
induce chemotaxis in human trophoblast cell lines. Mol Hum Reprod 14: 423-430.

73. Cao B, Diamond MS, Mysorekar IU (2017) Maternal-Fetal Transmission of Zika Virus: Routes
and Signals for Infection. J Interferon Cytokine Res 37: 287-294.

74. Red-Horse K, Zhou Y, Genbacev O, Prakobphol A, Foulk R, et al. (2004) Trophoblast
differentiation during embryo implantation and formation of the maternal-fetal interface. J
Clin Invest 114: 744-754.

75. Kolundzic N, Bojic-Trbojevic Z, Kovacevic T, Stefanoska I, Kadoya T, et al. (2011) Galectin-1 is
part of human trophoblast invasion machinery--a functional study in vitro. PLoS One 6:
e28514.

76. Kolundzic N, Bojic-Trbojevic Z, Radojcic L, Petronijevic M, Vicovac L (2011) Galectin-8 is
expressed by villous and extravillous trophoblast of the human placenta. Placenta 32: 909-
911.

77. He P, Shao D, Ye M, Zhang G (2015) Analysis of gene expression identifies candidate markers
and pathways in pre-eclampsia. J Obstet Gynaecol 35: 578-584.

78. Vambergue A, Fajardy I (2011) Consequences of gestational and pregestational diabetes on
placental function and birth weight. World J Diabetes 2: 196-203.

79. Fazleabas AT, Bell SC, Fleming S, Sun J, Lessey BA (1997) Distribution of integrins and the
extracellular matrix proteins in the baboon endometrium during the menstrual cycle and early
pregnancy. Biol Reprod 56: 348-356.

80. Damsky CH, Fitzgerald ML, Fisher SJ (1992) Distribution patterns of extracellular matrix
components and adhesion receptors are intricately modulated during first trimester
cytotrophoblast differentiation along the invasive pathway, in vivo. J Clin Invest 89: 210-222.

81. Zhou Y, Damsky CH, Chiu K, Roberts JM, Fisher SJ (1993) Preeclampsia is associated with
abnormal expression of adhesion molecules by invasive cytotrophoblasts. J Clin Invest 91:
950-960.

82. Kim MS, Yu JH, Lee MY, Kim AL, Jo MH, et al. (2016) Differential Expression of Extracellular
Matrix and Adhesion Molecules in Fetal-Origin Amniotic Epithelial Cells of Preeclamptic
Pregnancy. PLoS One 11: e0156038.

83. Bojic-Trbojevic Z, Jovanovic Krivokuca M, Vilotic A, Kolundzic N, Stefanoska I, et al. (2019)
Human trophoblast requires galectin-3 for cell migration and invasion. Sci Rep 9: 2136.

84. Hutter S, Knabl J, Andergassen U, Hofmann S, Kuhn C, et al. (2016) Placental Expression
Patterns of Galectin-1, Galectin-2, Galectin-3 and Galectin-13 in Cases of Intrauterine
Growth Restriction (IUGR). Int J Mol Sci 17: 523.

85. Hou PS, Chuang CY, Kao CF, Chou SJ, Stone L, et al. (2013) LHX2 regulates the neural
differentiation of human embryonic stem cells via transcriptional modulation of PAX6 and
CER1. Nucleic Acids Res 41: 7753-7770.

86. Kumamoto T, Hanashima C (2017) Evolutionary conservation and conversion of Foxg1 function
in brain development. Dev Growth Differ 59: 258-269.

87. Barbier A, Boivin A, Yoon W, Vallerand D, Platt RW, et al. (2013) New reference curves for
head circumference at birth, by gestational age. Pediatrics 131: e1158-1167.

88. Faria NR, Azevedo R, Kraemer MUG, Souza R, Cunha MS, et al. (2016) Zika virus in the
Americas: Early epidemiological and genetic findings. Science 352: 345-349.

89. Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34: i884-i890.

90. Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics 12: 323.

91. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, et al. (2013) STAR: ultrafast universal
RNA-seq aligner. Bioinformatics 29: 15-21.

92. Frankish A, Diekhans M, Ferreira A-M, Johnson R, Jungreis I, et al. (2018) GENCODE reference
annotation for the human and mouse genomes. Nucleic Acids Research 47: D766-D773.

93. McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-
Seq experiments with respect to biological variation. Nucleic Acids Research 40: 4288-4297.

94. Yu G, Wang L-G, Han Y, He Q-Y (2012) clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. OMICS: A Journal of Integrative Biology 16: 284-
287.

**Figure captions**

**Fig 1. Experimental design and ZIKV[BR] infection in hiPSC-derived trophoblasts.**
**(A).** Schematic: generation of trophoblasts from congenital Zika syndrome affected and non-affected
discordant twins' hiPSCs followed by ZIKV[BR] infection and analysis. Silhouettes are courtesy of
www.vecteezy.com (mother) and Yulia Ryabokon (babies). **(B).** Immunofluorescence for (β-CG,
human chorionic gonadotropin β) and KRT7 (CK7, cytokeratin 7, a pan trophoblast marker) in
hiPSC-derived trophoblasts. Scale bar: 20 μm. **(C).** ZIKV PFU/mL in trophoblasts' supernatant at
MOI = 0.3 and MOI = 3; mean ± SEM of the three twins; *p < 0.05; Student's t test. **(D).**
Representative plaque forming assay wells with stained VERO cells exposed to ZIKV collected at 96
hpi from the culture supernatants of affected or non-affected #10608 twins' hiPSC-derived
trophoblasts infected at a MOI of 0.3.
762

**Fig 2. Gene expression analyses of RNA-Seq data from hiPSC-derived trophoblasts from non-**
**affected and CZS-affected twins in culture after ZIKV$^{BR}$ infection.**
**(A).** Heatmap representation and clusterization of differentially expressed genes (DEGs) (FDR <
0.05; edgeR exact test), one in each line, in trophoblasts in culture after ZIKV$^{BR}$ infection (purple bar
at top) compared with control non-infected trophoblasts (blue bar at top), in cells derived from both
non-affected (#10608-4, #10763-4, and #10788-4) and CZS-affected (#10608-1, #10763-1, and
#10788-1) twins (one in each column, as indicated at the bottom). Color scale bar at right = Z score.
**(B).** Gene Ontology terms enrichment analysis of upregulated genes in hiPSC-derived trophoblasts
after ZIKV$^{BR}$ *in vitro* infection. The three major GO term categories, namely Biological Process,
Cellular Component and Molecular Function are separately represented in each panel. The size of the
circles is proportional to the number of genes in each significantly enriched category, as indicated at
the lower part scales; the colors show the statistical significance of the enrichment, as indicated by
the -log10 FDR values that appear in the color-coded scales at the bottom. A GO enrichment
significance cutoff of FDR ≤ 0.05 was used.
777
**Fig 3. Interferon responses in hiPSC-derived trophoblast after ZIKV$^{BR}$ infection.**
**(A).** Pathway enrichment of genes detected as differentially expressed (FDR < 0.05) between hiPSC-
derived trophoblasts from non-affected and CZS-affected twins at 96 h after ZIKV$^{BR}$ *in vitro*
infection; analysis was carried out with the ingenuity pathway analysis (IPA) tool. Gene upregulation
is depicted in shades of red, from white (not significantly changed), to dark red (highly upregulated).
**(B).** Luminex quantitation of IFNA, IFNG and IFNL1 detected in the supernatants of hiPSC-derived
trophoblasts in culture, from non-affected or CZS-affected twins at 48 h or 96 h after ZIKV$^{BR}$
infection. Data are represented as mean ± SEM and p-value * < 0.05 (Paired Student's t test).
786
**Fig 4. Differential gene expression between hiPSC-derived trophoblast from CZS-affected and**
**non-affected twins after ZIKV$^{BR}$ infection.**
**(A).** Heatmap representation and clusterization of differentially expressed genes (DEGs) (FDR <
0.05; edgeR exact test), one in each line, after ZIKV$^{BR}$ *in vitro* infection of trophoblasts from non-
affected twins (green bar at top; #10608-4, #10763-4, and #10788-4) as compared with trophoblasts
from CZS-affected twins (pink bar at top; #10608-1, #10763-1, and #10788-1) (one in each column,
as indicated at the bottom). Color scale bar at right = Z score. **(B).** Gene Ontology terms enrichment
analysis of downregulated genes in hiPSC-derived trophoblasts from CZS-affected twins when
compared with trophoblasts from non-affected twins after ZIKV$^{BR}$ *in vitro* infection. The three major
GO term categories, namely Biological Process, Cellular Component and Molecular Function are
separately represented in each panel. The size of the circles is proportional to the number of genes in
each significantly enriched category, as indicated at the scales in the lower part; the colors show the
statistical significance of the enrichment, as indicated by the -log10 FDR values that appear in the
color-coded scales at the bottom. A GO enrichment significance cutoff of FDR ≤ 0.05 was used. **C.**
Luminex quantitation of RANTES/CCL5 and IP10 detected in the culture supernatants of hiPSC-
derived trophoblasts from non-affected twins or from CZS-affected twins at 48 h (left panels) or 96 h
(right panels) after ZIKV$^{BR}$ infection. Data are represented as mean ± SEM and p-value * < 0.05
(Paired Student's t test).

# Supporting information

**S1 Fig**. **Expression measured by RT-qPCR of marker genes of hiPSCs and trophoblasts**.
Expression measured by RT-qPCR of *NANOG*, a hiPSC marker, and of *HCGA*, *HCGB* and *KRT7*,
three of the genes upregulated in the trophoblasts as compared with the hiPSCs. Twins from each
family are represented with a different color: red, #10608 twins; blue, #10763 twins; green, #10788
twins. (mean ± SEM; n = 2 biological replicates, except for #10763 due to sample loss during
culture)

**S2 Fig. Expression levels measured by RNA-Seq of proliferation-related genes in the hiPSCs
and in the hiPSC-derived trophoblasts from non-affected or CZS-affected twins. Related to
Figure 2.** The bars represent expression levels (in TPM) of selected genes associated with cellular
proliferation in hiPSCs from non-affected (light blue, hiPSC NA) or CZS-affected (dark blue, hiPSC
Aff) twins, in the hiPSC-derived trophoblasts from non-affected twins' mock (yellow, Troph – NA-
Mock) or ZIKV-infected cells (orange, Troph – NA-MOI 0.3), and in the hiPSC-derived trophoblasts
from CZS-affected twins' mock (red, Troph – Aff-Mock) or ZIKV-infected cells (brown, Troph –
Aff-MOI 0.3). The levels of expression of the genes were compared between non-affected (both non-
infected and infected) trophoblasts and non-affected hiPSCs; and from CZS-affected (both non-
infected and infected) trophoblasts and CZS-affected hiPSCs. Genes significantly down-regulated in
trophoblast cells when compared with hiPSCs are shown (one-away ANOVA, p-value threshold was
* 0.05, ** 0.01, *** 0.001 and **** 0.0001). Error bars show SEM.

**S3 Fig. Expression levels measured by RNA-Seq of genes encoding candidate attachment
factors implicated in ZIKV infection of human cells. Related to Figure 2.** The bars represent
expression levels (in TPM) of eight candidate ZIKV attachment factor genes in hiPSCs from non-
affected (light blue, hiPSC NA) or CZS-affected (dark blue, hiPSC Aff) twins, in the hiPSC-derived
trophoblasts from non-affected twins' mock (yellow, Troph – NA-Mock) or ZIKV-infected cells
(orange, Troph – NA-MOI 0.3), and in the hiPSC-derived trophoblasts from CZS-affected twins'
mock (red, Troph – Aff-Mock) or ZIKV-infected cells (brown, Troph – Aff-MOI 0.3). None of these
genes was significantly differentially expressed in hiPSC-derived trophoblasts from CZS-affected
twins, when compared with hiPSC-derived trophoblasts from non-affected twins in pairwise
comparisons (two-tailed t-test, equal variance). Error bars show SEM.

**S4 Fig. Expression levels measured by RNA-Seq of genes encoding representatives of Type I,
Type II, and Type III IFNs. Related to Figure 2.** The bars represent expression levels (in TPM) of
genes encoding interferons in hiPSCs from non-affected (light blue, hiPSC NA) or CZS-affected
(dark blue, hiPSC Aff) twins, in the hiPSC-derived trophoblasts from non-affected twins' mock
(yellow, Troph – NA-Mock) or ZIKV-infected cells (orange, Troph – NA-MOI 0.3), and in the
hiPSC-derived trophoblasts from CZS-affected twins' mock (red, Troph – Aff-Mock) or ZIVK-
infected cells (brown, Troph – Aff-MOI 0.3). None of these genes was significantly differentially
expressed in hiPSC-derived trophoblasts from CZS-affected twins when compared with hiPSC-
derived trophoblasts from non-affected twins in pairwise comparisons (two-tailed t-test, equal
variance). None of the 13 recognized human *IFNA* (*IFNA1, 2, 4, 5, 6, 7, 8, 10, 13, 14, 16, 17, 21*),

*IFNG*, *IFNK* and *IFNW1* genes were significantly expressed in any of the data sets. Error bars show
SEM.

**S5 Fig. Expression levels measured by RNA-Seq of genes encoding receptors for Type I, Type**
**II, and Type III IFNs. Related to Figure 2**. The bars represent expression levels (in TPM) of genes
encoding interferon receptors in hiPSCs from non-affected (light blue, hiPSC NA) or CZS-affected
(dark blue, hiPSC Aff) twins, in the hiPSC-derived trophoblasts from non-affected twins' mock
(yellow, Troph – NA-Mock) or ZIKV-infected cells (orange, Troph – NA-MOI 0.3), and in the
hiPSC-derived trophoblasts from CZS-affected twins' mock (red, Troph – Aff-Mock) or ZIKV-
infected cells (brown, Troph – Aff-MOI 0.3). None of these genes was significantly differentially
expressed in hiPSC-derived trophoblasts from CZS-affected twins when compared with hiPSC-
derived trophoblasts from non-affected twins in pairwise comparisons (two-tailed t-test, equal
variance). Error bars show SEM.

**S6 Fig. Differential gene expression between hiPSC-derived trophoblast from CZS-affected and**
**non-affected twins after ZIKV$^{BR}$ infection. Related to Figure 4.** Gene Ontology terms enrichment
analysis of upregulated genes in hiPSC-derived trophoblasts from CZS-affected compared with non-
affected twins after ZIKV$^{BR}$ *in vitro* infection. The major GO term categories, namely Biological
Process and Molecular Function are separately represented in each panel. The size of the circles is
proportional to the number of genes in each significantly enriched category, as indicated at the lower
part scales; the colors show the statistical significance of the enrichment, as indicated by the -log10
FDR values that appear in the color-coded scales at the bottom. A GO enrichment significance cutoff
of FDR ≤ 0.05 was used.

**S7 Fig. Expression measured by RT-qPCR of genes found in the RNA-Seq analysis**
**downregulated after ZIKV$^{BR}$ infection in trophoblasts from CZS-affected when compared with**
**non-affected twins**. Expression measured by RT-qPCR of *COL3A1*, *ITGA1* and *LGALS3*, three of
the genes downregulated after ZIKV$^{BR}$ infection in trophoblasts from CZS-affected (Aff) when
compared with non-affected (NA) twins. Twins from each family are represented with a different
color: red, #10608 twins; blue, #10763 twins; green, #10788 twins. Mean ± SEM is shown. (n = 2
biological replicates, except for #10763 due to sample loss during culture; One-tailed t-test, *
p<0.05).

**Fig 1**

880    **Fig 2**

885    **Fig 3**

**A**

**B**

**48h in culture**

**96h in culture**

889    **Fig 4**

894    **Supporting information**

895    **S1 Fig**

898   **S2 Fig**

903 **S3 Fig**

907    **S4 Fig**

911    **S5 Fig**

914    **S6 Fig**

919    **S7 Fig**

923