Differential tolerance of *Zymoseptoria tritici* to altered optimal moisture conditions during the early stages of wheat infection

Anne-Lise Boixel[1], Sandrine Gélisse[1], Thierry C. Marcel[1], Frédéric Suffert[1*]

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850, Thiverval-Grignon, France

* Corresponding author: Frédéric Suffert (frederic.suffert@inrae.fr)

Abstract

Foliar plant pathogens require liquid or vapour water for at least part of their development, but their response and adaptive tolerance to moisture conditions has been much less studied than other meteorological factors to date. Here, we examine the impact on the wheat-*Zymoseptoria tritici* interaction of altering optimal moisture conditions conducive to infection. We assessed the responses *in planta* of 48 *Z. tritici* strains collected in two climatologically distinct locations (Ireland and Israel) to four high moisture regimes differing in the duration of uninterrupted exposure to saturated relative humidity (100% RH) during the first three days of infection. Individual- and population-level moisture reaction norms expressing how the sporulating area of a lesion change with the RH conditions were established based on visual assessments of lesion development at 14, 17 and 20 days post-inoculation (dpi). Our findings highlighted: (i) a critical time-dependent effect on lesion development of interrupted periods of exposure to 100% RH during these earliest infection stages; (ii) a marked interindividual variation in the sensitivity to RH conditions both in terms of strain average moisture response and plasticity; (iii) a higher tolerance at 14 dpi of the Israeli population to interrupted periods of exposure to 100% RH. By indicating that sensitivity to sub-optimal moisture conditions

may vary greatly between *Z. tritici* individuals and populations, this study provides the first evidence of moisture adaptation signature in a fungal plant pathogen. This suggests that understanding such variation will be critical to predict their response to changing climatic conditions.

Keywords

Moisture sensitivity, relative humidity, phenotypic plasticity, reaction norm, individual variation, Septoria tritici blotch

Introduction

Micrometeorological factors within the plant canopy define the physical environment in which foliar crop pathogens develop (Campbell and Norman, 1998; Chelle, 2005). Temperature and moisture are key factors determining the outcome of host-pathogen interactions with critical effects highlighted at each epidemiological stage of foliar diseases. The presence, abundance and persistence of moisture determine whether the biological processes of infection (Yarwood, 1992), and colonization accelerate, slow or stop altogether, with consequences for the incubation and latent periods (Huber, 1992), spore production (Rotem *et al.*, 1978), and release (Ingold, 1978). The effects of temperature have been quantified in detail, but much less is known about the precise effects of moisture levels and thresholds, whether considered in terms of relative humidity (RH) or leaf wetness (presence of water films and droplets on aerial plant surfaces), on pathogen and disease development.

This gap in our knowledge may reflect the experimentally greater challenge of controlling, measuring and reporting moisture conditions with sufficient resolution and fidelity (Rowlandson *et al.*, 2015). Indeed, moisture conditions display a high degree of spatio-temporal variability in the field as they result from the dynamic equilibrium between water interception and evaporation (physical energy balance; Nobel, 2009). The accurate control of moisture conditions requires controlled environment plant growth chambers in which fluctuations in other meteorological variables are minimized (Butler *et al.*, 1995). For instance, RH is temperature-dependent as it is defined as the ratio of the partial pressure of water vapor to the equilibrium vapor pressure of water at a given temperature. Leaf wetness duration (LWD), by contrast, is even less easy to define (no single meteorological definition; Dawson & Goldsmith, 2018), lacks of a standard for its measurement (Sentelhas *et al.*, 2004) and exhibits heterogeneous spatio-temporal patterns as different parts of the leaves may be

wet and dry at different times, both in the field and in controlled conditions (Huber & Gillespie, 1992).

Several epidemiological studies have identified prolonged wetness as an important factor, and several disease-forecasting models contain the variable LWD (Rowlandson *et al.*, 2015). Although fewer studies have focused on RH requirements for optimal infection, some have established that the RH threshold (the level below which infection does not occur) differs between types of pathogens, up to very high RH requirements for some fungal species (e.g. 90% for *Magnaporthe oryzae*, Li *et al.*, 2014; 95% for *Didymella rabiei*, Jhorar *et al.*, 1998).

Studies of the response of a fungal pathogen to moisture conditions at the leaf scale during the earliest stages of infection (incubation period) are particularly relevant, because this period, together with the spore release and dispersal stages, is one of the most sensitive to LWD and RH in the vast majority of pathogens. The causal agent of Septoria tritici blotch, *Zymoseptoria tritici,* is a highly relevant biological model in this context because of: (i) its long latent period, spanning from 10 to 20 dpi (days post-inoculation) on seedlings and 15 to 30 on adult plants under controlled conditions at an average daily temperature of 18°C (Suffert & Thomson, 2018) *vs*. 15 to 35 dpi in field conditions, depending on temperature (Shaw, 1990); (ii) the crucial impact of moisture at all stages of the disease cycle in field conditions (Shaw, 1990).

The impacts of the duration and interruption of the wet period on *Z. tritici* (post-inoculation LWD) have already been investigated in controlled conditions (Holmes, 1974; Shearer & Zadoks, 1972; Eyal, 1977; Hess *et al.*, 1987; Chungu *et al.*, 2001; Magboul *et al.*, 1992; Shaw, 1990; Shaw, 1991; Fones *et al.*, 2017). An increase in RH favors an increase in overall infection rate (penetration of leaf tissues via the stomata; Kema *et al.*, 1996), hyphal growth and pycnidiation intensity (Shaw, 1991; Fones *et al.*, 2017). RH has been inferred to be optimal between 3 and 4 days, at 100% (Chungu *et al.*, 2001; Suffert *et al.*, 2013), and

pycnidiation has been reported to occur at a RH of 35-100% (Pachinburavan, 1981). However, threshold RH levels have not been experimentally tested, except for 50%-75%-100% comparisons (Shaw, 1991). The RH requirement threshold for *Z. tritici* development therefore remains unknown. LWD has also been shown to have critical effects on infection, pycnidiation and, in some cases, the duration of the latent period (Chungu *et al.*, 2001; Shaw, 1990). These experimental results are supported by the importance of the presence and duration of moisture (particularly LWD) for the prediction of Septoria tritici blotch epidemics in wheat (Hess & Shaner, 1987; Markham, 1992).

Reviews on the impact of climate change on plant diseases have stressed the importance of focusing on changes in temperature and rainfall patterns (Garrett *et al.*, 2006). Global climate models (GCMs) predict more frequent and extreme rainfall events and higher atmospheric water vapor concentrations with increasing temperature (Kirk-Davidoff *et al.*, 1999; Huntingford *et al.*, 2003). Until recently, it was difficult to obtain LWD and RH, which critically influence plant pathogen infection and disease development, from GCM outputs (Chakraborty *et al.*, 2000). It had been suggested that higher levels of canopy moisture promoted the development of a range of foliar pathogens, but this factor was never considered essential and was rarely taken into account in discussions on the effects of climate change on plant diseases (e.g. West *et al.*, 2012). However, the changes in RH are likely to be at least as marked as those in temperature in semi-arid zones, such as the area around the Mediterranean basin (Vicente-Serrano, 2007). The predicted slight decrease in the duration of wet periods and slightly warmer conditions will probably have opposite effects, decreasing and promoting infection, respectively, and detailed modeling approaches are therefore required to predict the overall outcome. This paradigm shift will require an understanding of the diversity of responses and levels of adaptation of pathogen populations to contrasting RH conditions. This understanding can be developed by identifying differences in moisture responses between

pathogen genotypes within the same species, and, more generally, by characterizing the intra-*vs.* inter-population phenotypic variability connected to genetic differentiation at neutral marker loci to reveal local adaptation (Merilä & Crnokrak, 2001).

RH may have been much less considered than temperature to date because there are more convenient methods for thermal phenotyping (e.g. Boixel *et al.*, 2019a) than for moisture phenotyping (Li *et al.*, 2014; Xu *et al.*, 2016). Assessments of the differences in response to temperature *in vitro* (tests conducted without interaction with the plant) are biologically meaningful and can be reasonably extrapolated (Boixel *et al.*, 2019a). By contrast, differences in the response to humidity must necessarily be performed *in planta* (tests conducted in interaction with the plant), particularly for the early stages of infection.

In this study, we investigated the diversity in moisture responses in *Z. tritici,* at the individual and population levels, for two field populations collected from contrasting climatic zones (humid *vs*. dry environment). We evaluated the aggressiveness of the individuals of these populations on wheat seedlings subjected to four moisture regimes. These regimes were defined by increasingly altering the inferred optimal RH conditions for *Z. tritici* over the first three days after inoculation that were reported as critical in the literature. This was done by sequentially incorporating interrupted periods of exposure to 100% RH resulting in the study of diurnal changes in humidity (reflecting what is happening in the field) rather than different levels of constant humidity.

Methods

Fungal material

Two *Z. tritici* populations were collected from contrasting Köppen-Geiger Euro-Mediterranean climatic zones (Peel *et al.*, 2007): (i) an Irish population (IR) sampled on cultivar JB Diego from a single field at Carlow (52°83'65'' N - 6°93'41'' E) in July 2016

(Cfb, oceanic climate); (ii) an Israeli population (IS) sampled on cultivar Galil from a single field at Kiryat-Tivon (32°71'62'' N - 35°12'74'' E) in March 2017 (Csa, hot-summer Mediterranean climate). These two populations, which have already been characterized for thermal adaptation and genotyped for 12 neutral genetic markers (Boixel *et al.*, 2019b), originate from environments with different average air moisture conditions, mainly due to marked differences in annual temperature and rainfall regimes (Figure 1). We randomly selected 24 IR and 24 IS *Z. tritici* isolates from these populations. These isolates were phenotyped in three sequential randomized batches of 16 strains at a time under controlled conditions in a growth chamber experiment with four treatments (different moisture regimes but identical temperature conditions). In total, each isolate × RH condition interaction was tested on three leaves and replicated twice at opposite ends of the growth chamber. These two replicates were statistically treated as two experimental blocks.

Plant material

The 48 isolates were phenotyped on the highly susceptible bread wheat cultivar, 'Taichung 29', commonly used as a susceptible check cultivar in the screening of wheat germplasm (Makhdoomi *et al.*, 2015) or in tests of global panels of *Z. tritici* isolates (Zhong *et al.*, 2017). Wheat seeds were sown in 0.4-liter pots, so as to obtain three seedlings per pot. The pots were kept in a growth chamber, under a controlled light/dark cycle (16 hours of light at 300 $\mu mol.m^{-2}.s^{-1}$ - Osram Lumilux L58W/830 - at 22°C and 80 % RH / 8 hours of dark at 18 °C and 100 % RH).

Inoculation procedure

Isolates were grown in Petri dishes containing PDA (potato dextrose agar, 39 g L^{-1}) at 18°C, in the dark, for five days. Spores suspensions were prepared by flooding the surface of the

five-day-old cultures with sterile distilled water and then scraping the agar surface with a sterilized glass rod to release the spores. The concentration of spores in the suspension was adjusted to 10^5 spores.mL^{-1} with a Malassez counting chamber. This concentration, previously used to assess Z. *tritici* aggressiveness and thermal adaptation (Suffert *et al*., 2013; 2015) is more relevant than much higher concentrations (e.g. 10^7 spores.mL^{-1}) because it maximises differences in disease expression and facilitates the detection of the effect of an experimental factor. We added one drops of surfactant (Tween 20, Sigma) per 15 mL suspension to ensure adequate coverage of the inoculated leaf surface. The spore suspensions were paintbrush-inoculated evenly over a precise area segment marked with indelible pen, the 7.5 cm mid-length portion of the first true leaf of 16-day-old seedling plants. Each of the isolate suspensions was tested in six biological replicates (six leaves in total) generating parallel measurements of biologically distinct samples.

Differential exposure to moisture regimes during the incubation period

After inoculation, the six plants from each isolate × RH condition interaction were split into two identical groups and assigned to two experimental blocks of the growth chamber compartment to account for inherent patterns of heterogeneity within growth chambers (Potvin *et al.*, 1990). Plants were subjected to four different moisture regimes (R0, R1, R2, and R3) defined based on the alteration of the reported optimal conditions for infection over the first three days after inoculation (the time required for pycnidiospore germination, epiphytic hyphal growth and stomatal penetration; Kema *et al.*, 1996; Duncan and Howard, 2000). Indeed, most Z. *tritici* inoculation protocols include a bagging time of three days at near 100 % RH (Chungu *et al.*, 2001; Suffert *et al.*, 2013; Fones *et al.*, 2015). As such, moisture conditions were modulated by varying the duration of time after inoculation (0, 1, 2 and 3 days, respectively) at which infected plants were continuously subjected to saturated

RH (near 100%). Saturated RH was reached when plants were covered with transparent polyethylene bags previously moistened with distilled water (maintenance of water condensation on the bags over the experiment as favored by the plant transpiration). Uncovered plants were exposed to the diurnal RH changes of the growth chamber (RH of 100% in the dark and of 80% during light periods; average automatic recording, every 15 minutes, with a ventilated sensor). Concretely, moisture regimes consisted in four durations of time at which infected plants were bagged (Figure 2): (R0) no-bagging treatment; (R1) one-day-bagging treatment; (R2) two-day-bagging treatment; (R3) three-day-bagging treatment. To establish moisture reaction norms for characterization and comparison of the RH sensitivity of the 48 isolates (see '*Data analyses*' section below), we summarized the lowest to highest moisture regimes in terms of the mean RH prevailing during the three-day-post-inoculation period for ease of calculation: 88.3% (R0), 92.2% (R1), 96.1% (R2) and 100% (R3). We chose not to characterize each moisture regime by the duration of time for which RH = 100 %. Indeed, there is no evidence to suggest that a RH of 80 %, which is higher than the RH value below which infection is thought to be impossible (RH = 50%; Shaw, 1991) but below the lowest humid conditions tested here, can prevent fungal growth and stop infection.

Disease assessment

Disease severity was assessed by eye, by the same assessor, at 14, 17 and 20 dpi, as the percentage of the inoculated leaf surface (1, 2, 3 and 5%, then increments of 5% up to 100%) displaying chlorosis (chlorotic area), necrosis (necrotic area, including both sporulating and non-sporulating area) and visible fruiting bodies or pycnidia (sporulating area).

Data analysis

Effect of moisture conditions on disease severity – The experimental and biological main effects on the variability of sporulating area (SPO, expressed as a percentage of the inoculated leaf surface) and their interactions were assessed by a statistical modeling approach. We investigated the partitioning of the variability into eight different components of variation (deviance): series (S), block (B), RH conditions (H), population (P), isolate I(P), RH conditions × population interaction (H×P), RH conditions × isolate interaction (H×I(P)), residuals (ε_{sbpih}). We analyzed these deviance components, by fitting a generalized linear model (GLM) with a log-link function and quasi-Poisson errors to account for overdispersion, according to the following model (R *glm* function; R Core Team, 2018): $SPO_{sbpih} = S + B + P + I(P) + H + H×P + H×I(P) + \varepsilon_{sbpih}$. *Post-hoc* comparisons of differences in SPO means (multivariate analysis of variance) and homoscedasticity (Levene's test) between IR and IS populations were performed to investigate differences in inter- and intra-population responses to moisture conditions.

Establishment of individual moisture reaction norms – Individual moisture reaction norms, describing the pattern of SPO as a function of relative humidity, were established for each isolate. Reaction norms were assumed to be linear within the investigated 80-100% RH range and were therefore estimated by a linear regression $y = ax + b$. Individual reaction norm properties were summarized by three parameters accounting for sensitivity to RH: (i) the intercept '*b*' corresponding to the elevation or the sensitivity to the driest environment; (ii) the steepness of the slope '*a*' corresponding to the response to variation, *i.e.* whether phenotypic responses increased or decreased in extreme environments and to what extent; (iii) the midpoint '$y_{94.2\%}$' corresponding to the average response or the response at the midpoint of the RH range, to consider shifts in the entire reaction norm. The heterogeneity of RH sensitivity across isolates at the individual level and the difference between IR and IS population-level moisture reaction norms were assessed on the basis of these three parameters.

Differentiation in individual and population responses – Phenotypic differentiation in terms of the intercept, slope and midpoint was compared to between-population neutral genetic differentiation, to determine the potential for local adaptation of moisture responses (P_{ST} - F_{ST} comparisons performed with the '*Pstat*' R package). Neutral genetic variability, and the structure and distribution of diversity between and within the IR and IS populations were characterized in a previous study (Boixel *et al.*, 2019b) based on microsatellite genotyping data acquired for 12 SSRs (ST1, ST2, ST3A, ST3B, ST3C, ST4, ST5, ST6, ST7, ST9, ST10, ST12 neutral microsatellites; Gautier *et al.*, 2014). In parallel, components of phenotypic variation in the response to the four moisture regimes (R0, R1, R2, R3) were extracted by principal component analysis. We projected all four-fold phenotypes onto a two-dimensional space and clustered the isolates (hierarchical classification on principal components, HCPC) on the basis of their sensitivities to the four moisture regimes ('*FactoMineR*' and '*factoextra*' R packages).

Results

Overall time-dependence effect of the experimental conditions

The block (B) and series (S) effects were significant despite the strong control over RH conditions of our experimental design (saturation-level application). These experimental effects are probably due to the spatial and temporal variability of other micrometeorological conditions reported to occur within growth chambers, even under controlled conditions (e.g. gradient in irradiance intensity due to artificial lighting; Poorter *et al.*, 2012). Mean RH had a significant effect on sporulating area at all times of disease assessment (14, 17 or 20 dpi; GLM, *p*-value < 0.01; Table 1, Figure 3). The significant differences in sporulating area induced by different *Z. tritici* isolates (I(P)) confirmed the genetic origin of the variability in aggressiveness. The population effect (P) was significant at 14 dpi, when sporulating area was

greater for IS than for IR isolates under the R0, R1 and R2 regimes (Figure 4). Significant H×I(P) and H×P interactions highlighted differential individual and population-level tolerance to the four moisture regimes (Table 1). At 14 dpi, differences in the mean sporulating area (GLM *post-hoc* analysis of the H×P interaction, *i.e.* the interaction of moisture regimes and differential population-level response to these conditions) and its intrapopulation variance (Levene's test for homogeneity of variance) between the IR and IS populations were significant (p-value < 0.05; Figure 4). At 17 dpi, these differences were small and did not differ significantly between the two populations; they had completely disappeared by 20 dpi.

Significant effect of moisture during the early stages of infection

The temporal dynamics of the effect of moisture regimes justified the sequential fitting of GLMs at 14, 17, and 20 dpi (Figure 3): at 14 dpi, the differences in sporulating area between R0-R1 and R2-R3 highlighted a dichotomy between unfavorable and favorable RH conditions. At 17 dpi, the differences between the four moisture regimes R0, R1, R2, and R3 were larger and significant. At 20 dpi, there were no longer any differences, due to saturation being reached for sporulating area (100%). In summary, from 0 to 3 dpi, the longer the duration of time at which infected wheat seedlings were exposed to a continuous RH 100%, the more intense the development of lesions. Indeed, a slight difference in high moisture regime during the very early stages of *Z. tritici* infection, despite the relatively high RH value (80%) of the air saturation interruption period, had a strong impact on the dynamics of lesion development.

Comparisons of tolerance to drier conditions are best performed at 14 dpi

Sporulating areas at 14 dpi were significantly larger for IS isolates on wheat plants subjected to the R0-R1-R2 regimes, whereas no significant difference was observed for R3 (Figure 4), highlighting the greater tolerance of drier conditions in the IS population. No difference was

observed between the populations at 17 and 20 dpi (no statistically significant H×P interaction, although this interaction was close to significance at 17 dpi; GLM, p-value < 0.1), suggesting that the driest regimes delayed infection processes but did not prevent *Z. tritici* development. These results led us to perform further analyses at 14 dpi, through comparisons of 'moisture reaction norms' or 'response curves', to assess the differences between isolates within these two pathogen populations in more detail.

Interindividual differences in moisture adaptation within the two populations at 14 dpi

The comparison of individual reaction norms (Figure S1), particularly for density distributions, and the ranges of values for the three parameters capturing the general characteristics of the response (intercept, slope, midpoint), highlighted high levels of interindividual variation in sensitivity to moisture regimes (Figure 5). Mean phenotypes did not differ significantly between the two populations (Wilcoxon rank sum test; differences in mean response assessed for the intercept: p-value $= 0.33$; slope: p-value $= 0.39$; midpoint: p-value $= 0.92$), but individual variation was greater for the midpoint coefficient in the IS population (Bartlett test of homogeneity of variance; p-value < 0.01). For the population-level linear moisture reaction norm, a significant difference in sporulating area was observed between the IS and IR populations for R1 (Wilcoxon rank sum test, p-value $= 0.03$), but not for the other three moisture regimes (p-value > 0.05). This suggest that RH conditions may have been too restrictive or detrimental for the expression of biological differences and variance for the trait under the R0 regime, whereas they were not limiting for R2 and R3. Neutral markers highlighted no differences in genetic structure between the IR and IS populations (Figure S2). P_{ST} values (computed at the critical c/h^2 ratio of 1) and their confidence intervals at $c/h^2 = 1$ indicated a robust difference between P_{ST} and F_{ST} (Figure S3). Thus, any phenotypic difference suggests that interindividual differences in moisture adaptation within the two populations may conceal signatures of local adaptation.

Classification of *Z. tritici* sensitivity to the four moisture regimes

Based on a PCA of phenotypic variation, we were able to classify the 48 *Z. tritici* isolates according to their sensitivity to the four moisture regimes (Figure 6). At 14 dpi, the R2 and R3 regimes did not discriminate between individual responses in terms of sporulating area. Cluster 4 consisted of isolates with a particularly large sporulating area (extreme phenotypes of is16, is19 and is34; Figure S1) that were less affected by the R0 and R1 regimes.

Discussion

Effect of moisture conditions during the early stages of *Z. tritici* infection

In this experimental study we established individual- and population-level moisture reaction norms, expressed as the area of *Z. tritici* sporulation on inoculated leaves for a RH averaged over the three days following inoculation. By this approach, we were able to quantify the critical effects of moisture conditions during the earliest stages of infection on the development of Septoria tritici blotch despite the relatively high RH value of the lowest humid regime tested (R0). It also made it possible to refine the findings of several previous studies performed in our range of experimental conditions (Holmes, 1974; Shearer & Zadoks, 1972; Eyal, 1977; Hess *et al.*, 1987; Chungu *et al.*, 2001; Magboul *et al.*, 1992; Shaw, 1990; Shaw, 1991; Fones *et al.*, 2017). The effects of moisture conditions were more pronounced at 14 dpi than at 17 and 20 dpi. Based on these results, we analyzed the data acquired at 14 dpi in more detail, to determine moisture sensitivity at both the individual and population levels. These findings also raise the question as to why the effects of moisture conditions are less pronounced at 17 and 20 dpi. Fones *et al.* (2017) previously showed that increasing leaf moisture is associated with increasing disease severity (number of pycnidia per leaf), because it decreases the time required for *Z. tritici* to penetrate into leaf tissues. This finding is

consistent with our results, suggesting that moisture conditions have a greater impact on the rate at which symptoms appear than on their final intensity. We may hypothesize that suboptimal moisture regimes have slowed epiphytic hyphal growth and penetration into the leaf tissues via the stomata, but that they did not ultimately reduce the germination rate or infection efficiency, at least for the RH values which were tested here. Epiphytic development was probably slower during the daytime periods, when wheat leaves were not bagged, resulting in a 'delay' in the infection process rather than an 'irreversible' ending of this process. This hypothesis should be tested with direct experimental approaches based on cytological observations or indirect approaches measuring infection efficiency, defined as the proportion of pathogen spores able to infect susceptible plant tissues once they have landed on them, which could be assessed by methods based on the time‑resolved imaging of disease progress (Karisto *et al.*, 2019).

Technical limitations: moving towards *in vitro* tests for minimizing background noise?

Even under controlled conditions, growth chambers are subject to spatiotemporal fluctuations in within-chamber conditions, particularly as concerns the distribution of light intensity (Potvin *et al.*, 1990; Poorter *et al.*, 2012). Such spatial patterns of heterogeneity may have contributed to the block effect we detected in our experimental design, whereas differences in inoculation conditions (e.g. abiotic factors, spore virulence levels) might also explain the differences in symptom expression between series (Kay *et al.*, 2019). One possible improvement to experimental methods, making it possible to screen larger numbers of isolates in a more standardized way (making it easier to separate environmental and physiological responses), would be to control and test moisture conditions *in vitro*, with the experimental devices proposed by Li *et al.* (2014; different RH conditions formed within humidity chambers obtained with different glycerol solution concentrations) or Xu *et al.* (2016; discrete RH gradient formed in spatially separated wells of a modified multiwell plate), for example.

This would have the advantage of countering the effect of the host resistance × environmental response interaction on the expression of aggressiveness in isolates, although, in our experiment, the two *Z. tritici* populations were similarly aggressive on the cultivar 'Taichung 29'. *In vitro* approaches may be particularly relevant for *Z. tritici,* as they have been shown to be a reliable *proxy* for the assessment of differences in thermal sensitivity (Boixel *et al*., 2019a).

First evidence of moisture adaptation in a fungal plant pathogen

Our results highlight strong individual variations in the phenotypic plasticity of *Z. tritici* with respect to sensitivity to high moisture regimes. Notwithstanding the relatively limited sample size (48 isolates) and number of populations ($n = 2$) investigated here, this study revealed, for the first time in a fungal plant pathogen, the existence of individual variation in responses to RH conditions. Moreover, the Israeli population was shown to be more tolerant to interrupted periods of optimal moisture conditions at 14 dpi. Together with the absence of genotypic differentiation for neutral microsatellite loci between the two populations, this result reveals signature of adaptation to moisture conditions which is consistent with the climatic conditions of the areas from which the two populations were collected.

The interaction between RH and isolate effects was significant, whenever disease was assessed, highlighting the importance of taking the interindividual variation in response to moisture conditions into account. Differences in moisture sensitivity were analyzed at both the population and individual levels. The moisture performance curves highlighted sensitivity variations between the two populations. The differences in the mean values of the parameters characterizing this sensitivity were not significant. However, the interindividual analysis revealed significant differences in moisture sensitivity between isolates, suggesting that it would be possible to quantify such differences more accurately in future experiments. This would require the study of larger numbers of individuals and more stringent moisture

conditions. For instance, lower moisture regimes could be tested to provide an accurate analysis of moisture adaptation to lower humidity levels which are similar to the annual air relative humidity conditions experienced by the Israeli population (69.7%).

The estimation of interactions between moisture and temperature adaptation at both population and individual levels is ambitious, but technically possible

The Israeli and Irish *Z. tritici* populations studied here are part of a Euro-Mediterranean set of eight populations included in a broader study on thermal adaptation (Boixel *et al.*, 2019b). General ecological concepts, knowledge and methods have been developed to a greater extent for thermal biology than for moisture biology. The estimation of interactions between moisture and temperature adaptation would also be relevant at both population and individual level. For this purpose, *Z. tritici* is an interesting case, both because a physical relationship between the two microclimatic variables, temperature and moisture, has been established (Shaw & Royle, 1993; Pietravalle *et al.*, 2003), and because the development of Septoria tritici blotch is known to be strongly influenced by both variables (Markham, 1992). The interactions between temperature and moisture effects could be studied from a mechanistic angle (for instance, testing pleiotropy *vs*. co-selection hypotheses in the genetic determinism of such a dual adaptation) but also from an epidemiological angle (predicting the consequences of climate changes affecting plant disease development in a more holistic way). The comparison of differences between temperature and moisture responses in different pathogen populations, starting with the Israeli and Irish populations, is particularly interesting because moisture adaptation is already interconnected with thermal adaptation in some models (e.g. Shinozaki & Yamaguchi-Shinozaki, 2000; Mutamiswa *et al.*, 2019).

How best to characterize individual variation of phenotypic plasticity and sensitivity to RH conditions in *Z. tritici* populations?

This study did not aim to determine the impact of mean RH value (constant moisture regime) nor the RH threshold below which *Z. tritici* isolates cannot infect wheat, but we can unambiguously conclude that a "moisture regime as close as possible to air saturation is best for *Z. tritici*". The minimal RH threshold for *Z. tritici* remains unknown. The challenge of answering this question is however quite limited since the moisture conditions to which a foliar pathogen is actually exposed in the field fluctuate justifying the relevance of focusing on responses to fluctuant rather than constant regimes. For instance, Shaw (1991) showed that breaks at 50% relative humidity had large effects but nevertheless enabled infection to occur. We can assume that there is a threshold moisture level (a lethal condition definitively blocking the infectious process), like that for temperature, but previous studies in *Z. tritici* and general knowledge about the epidemiology of fungal disease suggest that a minimum threshold, rather than a maximum, must be taken into account (as RH, by definition, cannot exceed 100%), together with a time period during which RH remains below this minimum threshold. Conversely, for temperature, the lethal threshold that should be considered is a maximum value, because *Z. tritici* can survive the thermal conditions of winter and freezing at -80°C in laboratory conditions.

Comparing the effects of the duration of uninterrupted periods of exposure to optimal moisture conditions as it was done here (see also Shaw, 1991; Magboul *et al.*, 1992; Chungu *et al.*, 2001) rather than effects of mean RH values (Fones *et al.*, 2017) makes the comparison of results from different experimental studies more complex. Therefore, one might be tempted to compare directly constant moisture conditions (e.g. optimal 100% RH *vs.* limiting 50% RH). However, this poorly represents what is happening in the field where fluctuating rather than constant moisture conditions prevail. Even in the driest areas where wheat is grown, the level of moisture in a crop canopy can be very high during or just after a rain period regardless of the smoothing average monthly air RH recorded at a site (e.g. 78-92% of RH in

Carlow, Ireland; 67-73% of RH in Kiryat-Tivon, Israel; Figure 1). In that respect, the moisture regimes tested in this study are a step closer to better understand the response to actual field conditions for which canopy RH is maximal during the night and lower during the day. More generally, the question of how best to describe RH conditions remains unresolved: averaging, intermittent favorable/unfavorable conditions, etc. How can the effect of the putative minimum moisture threshold (still unknown) be linked to the effect of the duration of dry periods? To which microclimatic variables are the individuals and populations really (mal)adapted? These questions and the spatio-temporal resolution of measurements should be addressed. For instance, it has been experimentally demonstrated that *Z. tritici* responds to 'leaf temperature' (mirroring with 'body temperature' for animals) rather than 'air temperature' (Bernard *et al.*, 2012), and that the time step for measurement is important (Bernard, 2012). These conclusions can also be likely extended to moisture.

Additional studies on adaptation to very suboptimal moisture regimes (e.g. 50-60% RH) rather than regimes close to the optimum (e.g. 90-100% RH) should be considered: (i) for methodological reasons, because this would probably maximize the expression of adaptation, making it easier to characterize experimentally, with populations contrasting less than the Israeli and Irish populations studied here; (ii) for epidemiological reasons, because it would provide knowledge to improve predictions of the adaptation of *Z. tritici* populations on bread and durum wheat in dryland farming areas (e.g. in Middle East and North Africa) in response to climate change. Such aims are ambitious and the studies will need to take into account the difficulties involved in experimental studies investigating the effects of moisture.

The exploitation of relevant analyses based on biological data requires attention to the issue of 'moisture stress tolerance' and an extension of reflections to the ecology of communities (Sheik *et al.*, 2011). Our study provides preliminary insight into such moisture stress tolerance and the diversity of its responses across individuals and populations when considering the

effects of climate change on plant disease epidemics. A knowledge of the microclimatic requirements for fungal pathogen development is essential for prospective studies of the impact of climate change with a solid experimental basis. To date, such studies have mostly focused on temperature, but climate change in the global context will also include large changes in moisture conditions with pronounced domino effects (Siepielski *et al.*, 2017).

Data Availability Statement

The data that support the findings of this study are openly available in the INRA Dataverse online data repository (https://data.inra.fr/) at https://doi.org/10.15454/FK7WHW.

Author Contributions

FS conceived the study and was in charge of overall direction and planning, with substantial input from ALB. SG performed the experiments according to a protocol developed jointly by FS and TM. ALB and FS analyzed the data, prepared the figures and wrote the manuscript. TM provided critical feedback. All authors approved the final version of the manuscript.

Funding

This work was supported by a grant from the French National Research Agency (ANR) as part of the "*Investissements d'Avenir*" programme ('SEPTOVAR' project; LabEx BASC; ANR-11-LABX-0034).

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

We are grateful to Dr. Stephen Kildea (Teagasc, Agriculture and Food Development Authority) and Dr. Hanan Sela (Institute for Cereal Crops Improvement, Tel Aviv University) for their help in sampling the Irish and Israeli *Z. tritici* populations, to Martin Willigsecker (INRAE BIOGER) for technical support throughout the experiment, and to Dr. Julie Sappa for her editorial advice in our English usage.

Supplementary Material

Supplementary material is available in the online version of this article.

References

Bashi, E., Rotem, J. (1974). Adaptation of four pathogens to semiarid habitats as conditioned by penetration rate and germinating spore survival. *Phytopathology* 64, 1035-1039.

Bernard, F. (2012). The development of a fungal foliar pathogen does react to temperature, but to which temperature? PhD thesis, AgroParisTech, Paris, France.

Bernard, F., Sache, I., Suffert, F., Chelle, M. (2013). The development of a foliar fungal pathogen does react to leaf temperature! *New Phytologist* 198, 232-240.

Boixel, A.-L., Chelle, M., Suffert, F. (2019b). Patterns of thermal adaptation in a worldwide plant pathogen: local diversity and plasticity reveal two-tier dynamics. *bioRxiv* https://doi.org/10.1101/867572.

Boixel, A.-L., Delestre, G., Legeay, J., Chelle, M., Suffert, F. (2019a). Phenotyping thermal responses of yeasts and yeast-like microorganisms at the individual and population levels: proof-of-concept, development and application of an experimental framework to a plant pathogen. *Microbial Ecology* 78, 42-56.

Butler, D.R., Reddy, R.K., Wadia, K.D.R. (1995). Single□plant chambers to control humidity for pathological studies. *Plant pathology* 44, 1-9.

Campbell, G.S., Norman, J.M. (1998). An introduction to environmental biophysics, 2nd ed. Springer-Verlag, New York, NY, USA.

Chakraborty, S., Tiedemann, A.V., Teng, P.S. (2000). Climate change: potential impact on plant diseases. *Environmental Pollution* 108, 317-326.

Chelle, M. (2005). Phylloclimate or the climate perceived by individual plant organs: What is it? How to model it? What for? *New Phytologist* 166, 781-790.

Chungu, C., Gilbert, J., Townley-Smith, F. (2001). *Septoria tritici* blotch development as affected by temperature, duration of leaf wetness, inoculum concentration, and host. *Plant Disease* 85, 430-435.

Dawson, T.E., Goldsmith, G.R. (2018). The value of wet leaves. *New Phytologist* 219, 1156-1169.

Duncan, K.E., Howard, R.J. (2000). Cytological analysis of wheat infection by the leaf blotch pathogen *Mycosphaerella graminicola*. *Mycological Research* 104, 1074-1082.

Eyal, Z., Brown, J.F., Krupinsky, J.M., Scharen, A.L. (1977). The effect of postinoculation periods of leaf wetness on the response of wheat cultivars to infection by *Septoria nodorum*. *Phytopathology* 67, 874-878.

Fones, H.N., Steinberg, G., Gurr, S.J. (2015). Measurement of virulence in *Zymoseptoria tritici* through low inoculum-density assays. *Fungal Genetics and Biology* 79, 89-93.

Fones, H.N., Eyles, C.J., Kay, W., Cowper, J., Gurr, S.J. (2017). A role for random, humidity-dependent epiphytic growth prior to invasion of wheat by *Zymoseptoria tritici*. *Fungal Genetics and Biology* 106, 51-60.

Garrett, K.A., Dendy, S.P., Frank, E.E., Rouse, M.N., Travers, S.E. (2006). Climate change effects on plant disease: genomes to ecosystems. *Annual Review of Phytopathology* 44, 489-509.

Hess, D.E., Shaner, G. (1987). Effect of moisture and temperature on development of septoria tritici blotch in wheat. *Phytopathology* 77, 215-219.

Holmes, S.J.I., Colhoun, J. (1974). Infection of wheat by *Septoria nodorum* and *S. tritici* in relation to plant age, air temperature and relative humidity. *Transactions of the British Mycological Society* 63, 329-338.

Huber, L., Gillespie, T.J. (1992). Modeling leaf wetness in relation to plant-disease epidemiology. *Annual Review of Phytopathology* 30, 553-577.

Huntingford, C., Jones, R.G., Prudhomme, C., Lamb, R., Gash, J.H., & Jones, D.A. (2003). Regional climate model predictions of extreme rainfall for a changing climate. *Quarterly Journal of the Royal Meteorological Society* 129, 1607-1621.

Ingold, C.T. (1978). Water and spore liberation. In: Water deficits and plant growth. V - Water and Plant Disease, ed. Kozlowski TT. Academic Press, London, UK, 119-140.

Jhorara, O.P., Butlerb, D.R., Mathaudaa, S.S. (1998). Effects of leaf wetness duration, relative humidity, light and dark on infection and sporulation by *Didymella rabiei* on chickpea. *Plant Pathology* 47, 586-594.

Karisto, P., Dora, S., Mikaberidze, A. (2019). Measurement of infection efficiency of a major wheat pathogen using time☐resolved imaging of disease progress. *Plant Pathology* 68, 163-172.

Kay, W.T., Fones, H.N., & Gurr, S.J. (2019). Rapid loss of virulence during submergence of *Z. tritici* asexual spores. *Fungal Genetics and Biology* 128, 14-19.

Kema, G.H., Yu, D., Rijkenberg, F.H., Shaw, M.W., Baayen, R.P. (1996). Histology of the pathogenesis of *Mycosphaerella graminicola* in wheat. *Phytopathology* 86, 777-786.

Kirk-Davidoff, D.B., Hintsa, E.J., Anderson, J.G., & Keith, D.W. (1999). The effect of climate change on ozone depletion through changes in stratospheric water vapour. *Nature* 402, 399.

Li, Y., Uddin, W., Kaminski, J.E. (2014). Effects of relative humidity on infection, colonization and conidiation of *Magnaporthe orzyae* on perennial ryegrass. *Plant Pathology* 63, 590-597.

Magboul, A.M, Geng, S., Gilchrist, D.G., Jackson, L.F. (1992). Environmental influence on the infection of wheat by *Mycosphaerella graminicola*. *Phytopathology* 82, 1407-1413.

Makhdoomi, A., Mehrabi, R., Khodarahmi, M., Abrinbana, M. (2015). Efficacy of wheat genotypes and Stb resistance genes against Iranian isolates of *Zymoseptoria tritici*. *Journal of General Plant Pathology* 81, 5-14.

Markham, P. (1992). Stress management: filamentous fungi as exemplary survivors. *FEMS Microbiology Letters* 100, 379-385.

Merilä, J., Crnokrak, P. (2001). Comparison of genetic differentiation at marker loci and quantitative traits. *Journal of Evolutionary Biology* 14, 892-903.

Mutamiswa, R., Machekano, H., Chidawanyika, F., & Nyamukondiwa, C. (2019). Life-stage related responses to combined effects of acclimation temperature and humidity on the thermal tolerance of *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae). *Journal of Thermal Biology* 79, 85-94

New, M.G., Hulme, M., Jones, P.D. (1999). Representing 20^{th} century space-time climate variability. I: Development of a 1961-1990 mean monthly terrestrial climatology. *Journal of Climate* 12, 829-856.

Nobel, S.P. (2009). Physicochemical and environmental plant physiology (4th Ed). Elsevier, London, UK.

Pachinburavan, A. (1981). Pycnidiospore germination, penetration, and pycnidial formation of *Septoria tritici* Rob. ex Desm. PhD thesis. Washington State University, Washington, USA.

Peel, M.C., Finlayson, B.L., & McMahon, T.A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4, 439-473.

Pietravalle, S., Shaw, M.W., Parker, S.R., Van Den Bosch, F. (2003). Modeling of relationships between weather and *Septoria tritici* epidemics on winter wheat: a critical approach. *Phytopathology* 93, 1329-1339.

Poorter, H., Fiorani, F., Stitt, M., Schurr, U., Finck, A., Gibon, Y., Usadel, B., Munns, R., Atkin, O.K., Tardieu, F., Pons, T.L. (2012). The art of growing plants for experimental purposes: a practical guide for the plant biologist. *Functional Plant Biology* 39, 821-838.

Potvin, C., Lechowicz, M.J., Bell, G., Schoen, D. (1990). Spatial, temporal, and species-specific patterns of heterogeneity in growth chamber experiments. *Canadian Journal of Botany* 68, 499-504.

R Core Team. (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rotem, J., Cohen, Y., Bashi, E. (1978). Host and environmental influences on sporulation in vivo. *Annual Review of Phytopathology* 16, 83-101

Rowlandson, T., Gleason, M., Sentelhas, P., Gillespie, T., Thomas, C., Hornbuckle, B. (2015). Reconsidering leaf wetness duration determination for plant disease management. *Plant Disease* 99, 310-319.

Sentelhas, P.C., Gillespie, T.J., Gleason, M.L., Monteiro, J.E.B., Helland, S.T. (2004). Operational exposure of leaf wetness sensors. *Agricultural and Forest Meteorology* 126, 59-72.

Shaw, M.W. (1990). Effects of temperature, leaf wetness and cultivar on the latent period of *Mycosphaerella graminicola* on winter wheat. *Plant Pathology* 39, 255-268.

Shaw, M.W. (1991). Interacting effects of interrupted humid periods and light on infection of wheat leaves by *Mycosphaerella graminicola* (*Septoria tritici*). *Plant Pathology* 40, 595-607.

Shaw, M.W., Royle, D.J. (1993). Factors determining the severity of epidemics of *Mycosphaerella graminicola* (*Septoria tritici*) on winter wheat in the UK. *Plant Pathology* 42, 882-899.

Shearer, B.L., Zadoks, J.C. (1972). The latent period of *Septoria nodorum* in wheat. 1. The effect of temperature and moisture treatments under controlled conditions. *Netherlands Journal of Plant Pathology* 78, 231-241.

Sheik, C.S., Beasley, W.H., Elshahed, M.S., Zhou, X., Luo, Y., Krumholz, L.R. (2011). Effect of warming and drought on grassland microbial communities. *The ISME Journal* 5, 1692-1700.

Shinozaki, K., Yamaguchi-Shinozaki, K. (2000). Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Current Opinion in Plant Biology* 3, 217-223.

Siepielski, A. M., Morrissey, M. B., Buoro, M., Carlson, S.M., Caruso, C.M., Clegg, S.M., Coulson, T., DiBattista, J., Gotanda, K.M., Francis, C.D., Hereford, J., Kingsolver, J.G., Augustine, K.E., Kruuk, L.E., Martin, R.A., Sheldon, B.C., Sletvold, N., Svensson, E.I., Wade, M.J., MacColl, A.D. (2017). Precipitation drives global variation in natural selection. *Science* 355, 959-962.

Suffert, F., Sache, I., Lannou, C. (2013). Assessment of quantitative traits of aggressiveness in *Mycosphaerella graminicola* on adult wheat plants. *Plant Pathology* 62, 1330-1341.

Suffert, F., Ravigné, V., Sache, I. (2015). Seasonal changes drive short-term selection for fitness traits in the wheat pathogen *Zymoseptoria tritici*. *Applied and Environmental Microbiology* 81, 6367-6379.

Suffert, F., Thompson, R. (2018). Some reasons why the latent period should not always be considered constant over the course of a plant disease epidemic. *Plant Pathology* 67, 1831-1840.

Vicente-Serrano, S.M. (2007). Evaluating the impact of drought using remote sensing in a Mediterranean, semi-arid region. *Natural Hazards* 40, 173-208.

West, J.S, Townsend, J.A., Stevens, M., Fitt, B.D.L. (2012). Comparative biology of different plant pathogens to estimate effects of climate change on crop diseases in Europe. *European Journal of Plant Pathology* 133, 315-331.

Xu, Z., Jiang, H., Sahu, B.B., Kambakam, S., Singh, P., Wang, X., Wang, G., Bhattacharyya, M.K., Dong, L. (2016). Humidity assay for studying plant-pathogen interactions in miniature controlled discrete humidity environments with good throughput. *Biomicrofluidics* 10, 034108.

Yarwood, C.E. (1978). Water and the infection process. In: Water deficits and plant growth. V - Water and Plant Disease, ed. Kozlowski TT. Academic Press, London, UK, 141-173.

Table 1 - **Analysis of the deviance of sporulating area (SPO in %) induced by 48** *Zymoseptoria tritici* **isolates on wheat seedlings exposed to four high moisture regimes (see Figure 2) at 14, 17 and 20 dpi**. A quasi-Poisson generalized linear model (GLM) with a log-link function was fitted to experimental data, to assess the relative importance of the factors considered for the observed variation: series (S), block (B), RH conditions (H), population (P), isolate I(P), RH conditions × population interaction (H×P), RH conditions × isolate interaction (H×I(P)). The values reported in the table correspond to the components of deviance and their corresponding *p*-values (*P*).

	SPO_{14dpi}		SPO_{17dpi}		SPO_{20dpi}	
	Deviance	*P*	Deviance	*P*	Deviance	*P*
Series (S)	2480.1	< 0.01	4899.5	< 0.01	3917.4	< 0.01
Block (B)	49.5	< 0.01	87.5	< 0.01	94.9	< 0.01
Population (P)	87.5	< 0.01	2.5	0.56	0.2	0.83
Isolate I(P)	7368.6	< 0.01	10033.2	< 0.01	7772.5	< 0.01
RH conditions (H)	5283.0	< 0.01	7426.8	< 0.01	3851.1	< 0.01
H×P	224.0	< 0.01	51.4	0.08	2.8	0.91
H×I(P)	2241.0	< 0.01	4047.8	< 0.01	4091.8	< 0.01

Fig. 1 - Relative humidity (RH) conditions and precipitation at the two sampling sites.
(a) Average annual air RH in the Euro-Mediterranean area (data from the Atlas of the Biosphere, http://atlas.sage.wisc.edu/; New *et al.*, 1999); (b) average monthly air RH and rainfall distribution (1982-2012 data from the World Climate Database, https://en.climate-data.org/) at the sites from which the two *Zymoseptoria tritici* populations of 24 isolates each were collected (Carlow in Ireland for the IR population; Kyriat-Tivon in Israel for the IS population). The horizontal gray bar indicates the standard wheat-growing period (from sowing to harvest).

Fig. 2 - **Moisture regimes exerted over the three first days post-inoculation (dpi).** (a) Overview of the experimental steps. (b) Solid (from 0 to 3 dpi) and dotted (after 3 dpi) lines depict the four post-inoculation moisture regimes to which the wheat seedlings were subjected in the early stages of infection: R0 = no-bagging treatment; R1 = one-day-bagging treatment; R2 = two-day-bagging treatment; R3 = three-day-bagging treatment. White and gray areas indicate successive light and dark periods, respectively. Horizontal arrows indicate the period during which wheat plants were enclosed within polyethylene bags to maintain 100% HR.

Fig. 3 - Effect of moisture regime (R0, R1, R2, R3; see Figure 2) on the percentage of sporulating area (SPO) induced by *Zymoseptoria tritici* (*n* = 48 isolates), by disease assessment time (14, 17, 20 dpi). Medians are indicated by the lines at which the notches converge, and different letters indicate significant difference in SPO between experimental conditions in *post-hoc* test comparisons (multivariate analysis of variance).

Fig. 4 - **Dynamics of the effect of moisture regime (R0, R1, R2, R3; see Figure 2) on the percentage of sporulating area (SPO expressed as the mean ± SEM) in the early stages of *Zymoseptoria tritici* infection, by isolate origin - Ireland (IR; solid line) or Israel (IS; dashed line) - and the timing of disease assessment (14, 17, 20 dpi).** Significant differences in the mean (*post-hoc* analysis of the GLM H×P interaction, *i.e.* the interaction of moisture regimes and differential population-level response to these conditions) and the intrapopulation variance (Levene's test for homogeneity of variance) of SPO between IR and IS populations are highlighted by black and gray stars, respectively (*p*-value < 0.05). Each data point corresponds to the mean of 24 isolates (six biological replicates per isolate) composing the population.

Fig. 5 - Variation of *Zymoseptoria tritici* response to moisture conditions at the population (IR: Irish population, IS: Israeli population) and individual levels, at 14 dpi. (a) Population-level moisture reaction norms (mean RH prevailing over the first three days after inoculation: 88.3% for R0, 92.2% for R1, 96.1% for R2, 100% for R3) for the IR ($n = 24$ isolates; black circles/solid line) and IS ($n = 24$ isolates; gray triangles/dashed line) populations. The corresponding linear regression lines fitted to the percentage of sporulating area (SPO, expressed as the mean ± SEM) are displayed with their 95% confidence intervals. (b) Interindividual variation of the three chosen descriptors of individual moisture reaction norms: intercept (*b*), slope (*a*) and midpoint ($y_{94.2\%}$). Crude individual values (open circles), means (horizontal black thick lines), distributions (smoothed density curves) and 95% Bayesian highest density intervals (central rectangular boxes enclosing the means) are shown for each descriptor for which interindividual variation is displayed by population (IR, IS; six biological replicates per isolate).

Fig. 6 - Classification of *Zymoseptoria tritici* sensitivity to the four moisture regimes (R0, R1, R2, R3; see Figure 2) at 14 dpi. (a) Principal component analysis (PCA) biplot showing Irish (IR, *n* = 24 isolates, black points) and Israeli (IS, *n* = 24 isolates, gray points) isolates plotted in two dimensions, using their projections onto the first two principal components to summarize their response (SPO, *i.e.* percentage of sporulating area) under the four moisture regimes. HCPC clusters of individual responses are shown as colored areas on the factorial plane. (b) Relative proportions of the four identified clusters in the IR and IS populations. (c) Reading grid for the clustering of isolates based on their responses to the four moisture regimes (six biological replicates per isolate): SPO significantly lower (-) or higher (+) than the dataset mean (0).

a

b

Fig. S1 - Individual moisture reaction norms at 14 dpi for (a) the 24 Irish (IR) and the (b) 24 Israeli (IS) *Zymoseptoria tritici* **isolates.** Sporulating areas (SPO) are expressed as a percentage of the inoculated leaf area for each isolate (except ir7, due to unsuccessful inoculation), in response to four moisture conditions characterized by the mean RH prevailing during the three-day period immediately after inoculation (see Figure 2). Linear regression lines (in blue) and their 95% confidence intervals were fitted independently to experimental observations (dark points corresponding to six seedling wheat leaves per isolate × moisture conditions, *i.e.* six biological replicates).

Fig. S2 - Genetic subdivision of the 48 *Zymoseptoria tritici* isolates composing the Irish (IR) and Israeli (IS) populations. Bar plot displaying the Bayesian genetic clustering of IR and IS populations according to 12 SSR markers. The STRUCTURE algorithm was applied under the admixture and correlated allele frequencies model (500,000 iterations of the Markov chain followed by a run phase of 1,000,000 iterations with 10 independent replicates for each tested number of clusters). Each isolate is represented by a single vertical bar broken into two color segments, the lengths of which are proportional to the probability of to the isolate being assigned to the inferred genetic clusters ($K = 2$). No population structure was detected because each individual was affected to the three genetic clusters with similar probabilities.

Fig. S3 - Phenotypic divergence in moisture reaction norms between the IR and IS populations (*n* = 48 isolates). Sensitivity analyses exploring the robustness of P_{ST}–F_{ST} comparisons over variations in the c/h^2 ratio, which determines the accuracy of the approximation of Q_{ST} by P_{ST}. Phenotypic divergence in individual moisture reaction norms between populations (P_{ST}) was investigated for the intercept, slope and midpoint parameters. On each plot, estimates of P_{ST} (solid line in red) and F_{ST} (dashed line in green) are shown, to reveal the contrasting patterns of phenotypic responses to moisture conditions and neutral genetic divergence. The P_{ST} values displayed (computed at the critical c/h^2 ratio of 1) and their confidence intervals at $c/h^2 = 1$ indicate the occurrence of a robust difference between P_{ST} and F_{ST}, suggesting the existence of signatures of local adaptation.