1

Diffusion MRI free water is a sensitive marker of age-related changes in the cingulum

Manon Edde[1,2*], Guillaume Theaud[6*], François Rheault[6], Bixente Dilharreguy[2], Catherine Helmer[4], Jean-François Dartigues [4,5], Hélène Amieva[4], Michèle Allard[1,2,5], Maxime Descoteaux[6†], Gwénaëlle Catheline[1,2,3†]

[1] EPHE, PSL, F-33000, Bordeaux, France

[2] CNRS, INCIA, UMR 5287, F-33000 Bordeaux, France

[3] Université de Bordeaux, INCIA, UMR 5287, F-33000 Bordeaux, France

[4] Université de Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR 1219, F-33000 Bordeaux, France

[5] CHU de Bordeaux, Bordeaux, France

[6] Sherbrooke Connectivity Imaging Lab (SCIL), Université de Sherbrooke, Sherbrooke, QC, Canada

Corresponding author:

Manon Edde

Bât. 2A - 2ème étage - Case 22,

146 rue Léo Saignat,

33076 Bordeaux cedex

eddemanon@gmail.com

*Co-authors. These authors have contributed equally.

[†] Co-senior authors. These senior authors have contributed equally.

Running title:

Free Water: a marker of age-related white matter changes

Abstract

Diffusion MRI is extensively used to investigate changes in white matter microstructure. However, diffusion measures within white matter tissue can be affected by partial volume effects due to cerebrospinal fluid and white matter hyperintensities, especially in the aging brain. In previous aging studies, the cingulum bundle that plays a central role in the architecture of the brain networks supporting cognitive functions has been associated with cognitive deficits. However, most of these studies did not consider the partial volume effects on diffusion measures. The aim of this study was to evaluate the effect of free water elimination on diffusion measures of the cingulum in a group of 68 healthy elderly individuals. We first determined the effect of free water elimination on conventional DTI measures and then examined the effect of free water elimination on verbal fluency performance over 12 years. The cingulum bundle was reconstructed with a tractography pipeline including a white matter hyperintensities mask to limit the negative impact of hyperintensities on fiber tracking algorithms. We observed that free water elimination improved the sensitivity of conventional DTI measures to detect associations between tissue-specific diffusion measures of the cingulum and changes in verbal fluency in older individuals. Moreover, free water content measured along the cingulum was independently strongly associated with changes in verbal fluency. These observations suggest the importance of using free water elimination when studying brain aging and indicate that free water itself could be a relevant marker for age-related cingulum white matter modifications and cognitive decline.

Keywords: aging, tractography, free water, white matter hyperintensity, cingulum

1. Introduction

Aging is associated with widespread brain structural modifications including neurodegeneration of white and grey matter [1–3]. In the last few years, diffusion tensor imaging (DTI) has been widely used to indirectly explore microstructural properties of white matter and constitutes a sensitive technique to describe age-related white matter microstructural changes [4,5]. Parameters quantified by DTI such as Fractional Anisotropy (FA), Mean Diffusivity (MD) and Radial Diffusivity (RD) can be used to indirectly infer changes in axonal integrity and myelination [6]. In older adults, previous DTI studies reported a decrease in FA and an increase in MD and RD in major white matter tracts [6–8] that correlated with cognitive impairment [9–12].

However, diffusion measures within brain tissues can be affected by partial volume effects due to cerebrospinal fluid [13], especially in aging individuals with atrophied brains and ventriculomegaly [14–17]. When voxels contain cerebrospinal fluid (e.g. partial volume), diffusion measures can be biased towards a pattern of high diffusivity (MD, Axial Diffusivity (AD), RD) and reduced FA [18,19]. This effect has been particularly observed in the fornix and the corpus callosum because of their close proximity to periventricular regions [18,20,21]. To overcome this problem, the Free Water Elimination (FWE) method was used to differentiate the diffusion properties of brain tissues such as white matter bundles from the surrounding free water composed of cerebrospinal fluid [13,22]. Elimination of extracellular free water contamination improves the sensitivity and specificity of most measures derived from conventional DTI [16,22–24]. Although such correction has been used in pathological conditions [25–30], the majority of diffusion MRI studies do not consider free water effects of aging. Aging is associated with grey and white matter tissue loss, and the resulting enlargement of interstitial space could lead to increase in free water [31–34]. In addition, free water content in white matter fibers was recently associated with the presence of white matter hyperintensities (WMHs) in clinical studies [28,35–38]. In older adults, the prevalence and severity of WMH burden increases after age 60 [39,40]. Previous studies reported that the presence of WMH lesions was not only associated with age-related white matter changes evaluated with DTI (Maillard et al., 2014; Maniega et al., 2015; Pelletier et al., 2015) or tractography (Reginold et al., 2016, 2018; Svärd et al., 2017), but also with cognitive impairments mainly in memory and executive functions [5,46,47].

The cingulum bundle is one of the most studied white matter tracts running from the anterior to the posterior part of the brain that interconnects frontal, parietal, and temporal areas (Catheline et al., 2010; Jones et al.,2013; Bubb et al., 2018). Due to this central location, the cingulum plays a central role in white matter architecture and functional networks that support and contribute to optimal cognitive function (Bubb et al., 2018). In previous studies, changes within this bundle due to aging were correlated with executive and memory deficits (Catheline et al., 2010; Ezzati et al., 2016; Kantarci et al., 2011; Metzler-Baddeley et al., 2012; Seiler et al., 2018;

Sexton et al., 2010). However, these observations still remain controversial considering that most of these studies

did not consider CSF partial effect due to atrophy or WMH burden. Recent tractography studies consistently

observed that atrophy-related partial volume effects [16,18] and age-related ventricular enlargement [15] or WMH

burden (Reginold et al., 2016) affect diffusion measures of the cingulum in healthy older individuals. In this regard,

Reginold and colleagues reported higher RD in cingulum tracts that crossed WMH than those outside WMH

regions (Reginold et al., 2016; see also Kurki & Heiskanen, 2018). Atrophy and WMH burden might therefore

impact cingulum diffusion measures suggesting a source of significant bias in studies exploring the integrity of

this bundle and its association with cognitive functions in aging.

In this study, we hypothesized that free water elimination would provide more accurate white matter

diffusion measures of the cingulum, which in turn would be better correlated with verbal fluency changes in 68

elderly individuals. To do so, the cingulum bundle was reconstructed with a tractography pipeline that includes a

WMH mask to limit the negative impact of hyperintensities on fiber tracking algorithms. We first described the

effect of FW-correction on diffusion measures of the cingulum derived from the conventional tensor DTI model.

Then, we explored the association between FW-corrected diffusion measures and cognitive changes as evaluated

using the Isaacs Set Test over 12 years. The Isaacs Set Test was chosen since it specifically assesses semantic

memory and executive functioning.

2. Methods

2.1. Dataset

Participants were selected from the Bordeaux sample of the Three-City study [55], an ongoing longitudinal

multicenter population-based elderly cohort designed to evaluate risk factors for dementia. The study protocol was

approved by the ethics committee of Kremlin-Bicêtre University Hospital (Paris, France), and all participants

provided written informed consent. At baseline, subjects were non-institutionalized, randomly recruited from

electoral lists, and were older than 65 years. Since the 1999-2000 baseline inclusion, an extended

neuropsychological assessment was administered by trained psychologists at each follow-up occurring at 2, 4, 7,

10 and 12 years. An MRI scan was performed at the 10-year follow-up for every subject. All 239 subjects were

screened, and individuals were excluded if they had dementia (n=8), brain pathologies (n=27) or if MRI images

were either unavailable (n= 42), distorted by artefacts (n=24), or failed pre-processing (n=15). In addition,

cingulum reconstruction failed in 16 subjects and tractography quality control revealed suboptimal data in 19

additional subjects. Out of the 239 initially screened subjects, 68 subjects were included in the study. All

participants were right-handed and had a Mini Mental State Examination (MMSE) score greater than or equal to

24.

2.2. Cognitive and clinical variables

The Isaacs Set Test (IST, Isaacs & Kennie, 1973) chosen for the study consists of a test on verbal fluency, where

subjects are asked to cite the highest number of words in four semantics categories. Three scores of the IST were

used, corresponding to the number of words cited by the subject at 15-seconds (IST 15s), 30-seconds (IST 30s)

and 60-seconds (IST 60s). Verbal fluency was evaluated over 12 years, using a subject-specific slope for each IST

score computed using a linear mixed model with time as a continuous variable, random intercept and slope. Over

the 12-year follow-up period, the negative mean annual slope indicated a decrease in the number of given words.

Some clinical variables were also collected: depressive symptoms evaluated using the Center for Epidemiologic

Studies-Depression scale (CESD, Radloff, 1977), vascular risk factors including hypertension as defined in

patients with a blood pressure > 140/90 mm Hg or taking anti-hypertensive medication, diabetes as defined in

patients with a blood glycemic levels > 7 mmol/l or taking hypoglycemic medication, and body mass index (BMI).

2.3. MRI acquisition

MRI scanning was performed using a 3T Achieva system (Philips Medical Systems, The Netherlands) equipped

with a 8-channel SENSE head coil. Head motions were minimized by using tightly padded clamps attached to the

head coil. Anatomical MRI volumes were acquired using a 3D magnetization-prepared rapid gradient-echo

(MPRAGE) T1-weighted sequence with the following parameters: repetition time (TR)/ echo time (TE) = 8.2

ms/3.5 ms, flip angle 7°, field of view (FOV) 256x256 mm^2, 180 slices of 1 mm of thickness, voxel size 1x1x1

mm^3. Fluid-attenuated inversion recovery (FLAIR) images were obtained with the following parameters:

TR=11000 ms; TE=140 ms, inversion time = 2800 ms, 90-degree flip angle, FOV 230x172 mm^2, 24 slices of 5

mm of thickness, voxel size 0.72x1.20x5 mm^3. Diffusion weighted images were acquired using a spin echo single

shot EPI sequence composed of one b0 map (b-value=0 s/mm^2) followed by 21 diffusion gradients maps (b-

value=1000s/mm^2) homogenously spread over a half sphere and 21 opposite directions spread over the opposite

half sphere. To increase signal-to-noise ratio, a second series of two b0 and 42 DWI volumes was acquired. Sixty

axial slices were acquired with the following parameters: TR= 9700 ms, TE= 60 ms, flip angle 90°, FOV 224×224

mm^2, 60 slices, no gap and voxel size 2x2x2 mm^3. All acquisitions were aligned on the anterior commissure-

posterior commissure plan.

2.4. MRI preprocessing

2.4.1. Diffusion MRI preprocessing

Diffusion MRI (dMRI) images were pre-processed using FMRIB's Diffusion toolbox [58,59] in order to produce individual FA, MD, AD and RD maps in native space. For each subject, dMRI images were co-registered to the b0 reference image with an affine transformation and were corrected for motion and eddy current distortions. Brain Extraction Tool (BET, Smith, 2002) was applied to eliminate non-brain voxels and resulting dMRI images were denoised using the non-local mean denoising method [61]. To increase signal-to-noise ratio, the two reversed phases were then combined into a dMRI image using FSL tools. Finally, the fiber Orientation Distribution Functions (fODF) map was computed using the spherical constrained deconvolution (Descoteaux et al., 2007; Tournier et al., 2007; Theaud et al., 2019). Visual quality check was performed and did not reveal any processing failure for the included subjects.

2.4.2. White Matter Hyperintensity segmentation

White matter hyperintensity volumes were automatically segmented by the lesion growth algorithm implemented in the Lesion Segmentation Tool (LST, v2.0; Schmidt et al., 2012) of SPM12. For each participant, the T1-weighted image was used to generate partial volume estimation and three tissue probability maps (grey matter, white matter and cerebrospinal fluid). Then, each FLAIR image was co-registered on the corresponding T1 image to compute lesion belief maps for the three tissue classes. These maps were finally summed up and a lesion growth model with a pre-chosen initial threshold ($\kappa = 0.3$) was applied to create lesion maps in native space. Visual inspection of the lesion probability maps was performed and manually corrected when inaccuracies were found. Finally, WMH volumes were extracted, normalized by white matter volume and log transformed (total WMH). The volume of WMH within the cingulum was estimated by crossing the WMH mask with the cingulum bundle (cingulum WMH).

2.4.3. Free Water Elimination

Free Water Elimination (FWE, Pasternak et al., 2009) was used to estimate and remove the free water components from diffusion images. To isolate the fractional volume of freely diffusing extracellular water molecules from tissue compartments (i.e. reflecting intracellular processes), FWE fits a bi-tensor model within each voxel: a first one models free water diffusion, defining as an isotropic diffusion, and a second one models the tissue compartment [24]. The isotropic compartment models extracellular water which is characterized by freely and not hindered diffusion. A fixed diffusivity of 3.10^{-3} mm^2/s (i.e. diffusion coefficient of water at body temperature) is

used for this compartment. The fraction of free water content per voxel provided a native free water (FW) map for

each subject and varied between 0 to 1. In contrast, the tissue compartment models water molecules close to cellular

membranes of brain tissue, which are defined by restricted or hindered diffusion using a diffusion tensor. Thereby,

the volume of freely diffusing extracellular water molecules is removed from tissue compartments. Consequently,

the FW-corrected measures were expected to be more sensitive and specific to tissue changes than the measures

derived from the single tensor model [23,27,29]. The FW-corrected DTI maps were called FA tissue (FAt), RDt,

ADt and MDt (Chad et al., 2018).

2.4.4. Cingulum tractography and measures

For each participant, a local tracking using probabilistic algorithm based on fODF maps was performed. The

tractography was performed using twenty tracking seeds per voxels included in the white matter mask. The seeding

and tracking masks were modified to include WMH masks (see supplementary material). Next, the left and right

cingulum bundles were extracted using the RecoBundles approach (Garyfallidis et al., 2018, Figure 1).

RecoBundles is a model-based algorithm performing a registration of a Cingulum model on each subject to extract

the subject specific cingulum tract from the whole tractogram. Five different Cingulum models were used in this

study (RecoBundles). Finally, after visual inspection of cingulum bundles using dmriqc tool

(https://github.com/GuillaumeTh/dmriqcpy), 13 subjects were excluded due to the absence of right or left

cingulum bundles and 6 subjects were excluded because of miss-segmentation of hippocampal and anterior parts

of the cingulum bundle, mainly because of the age range of our population (older than 85 years of age) at the time

of MRI sessions. Therefore, the cingulum bundle was considered as a whole bundle and the different subdivisions

of the cingulum were not investigated in this study.

Cingulum diffusion measures were extracted before (FA, MD, RD and AD) and after FW-correction (FAt, MDt,

RDt and ADt). The diffusion measures without free water correction are defined as conventional DTI measures

and reflect the weighted average of all compartments including free water, whereas diffusion measures after

correction of free water are defined as FW-corrected measures. Measures of the left and right cingulum were

averaged for the analysis.

A percentage of measure changes (% change) for each participant was determined by computing the difference

between the measure before and after FW-correction divided by the measure before FW-correction [*i.e.* %

change = ((FW-corrected measure – DTI measure)/DTI measure) × 100].

Figure 1. Left and right cingulum templates used to extract the cingulum tract for each participant (A); Example of the cingulum bundle obtained for one subject displayed on the corresponding T1-weighted image (B). Along the fibers, color represents the RGB scale.

2.5. Statistical analysis

Because of non-normality of the continuous and categorical variables, Mann-Whitney U-tests and Spearman's correlations were used to evaluate the associations between cingulum diffusion measures as well as verbal fluency with continuous variables (age, global cognition score, depressive symptoms score, body mass index and WMH volume) and categorical variables (gender, education level, presence/absence of diabetes, of hypertension) respectively. To compare the effects of FW-correction to conventional DTI on cingulum diffusion measures, a Paired t-Test was performed. Linear regression models were then computed to describe the extent to which the FW-correction affects the association between cingulum diffusion measures and changes in verbal fluency. Diffusion measures were included as independent variables (conventional and FW-corrected FA, MD, AD and RD in separate models) and the verbal fluency slope set as the dependent variable in model adjusted for age and WM volume of the cingulum. Similar additional linear regression models were performed to examine the relationship between free water content and changes in verbal fluency. Total WMH volume was added in previous models to evaluate to the extent to which WMH burden affects the observed associations. In a supplementary analysis, specific WMH burden of the cingulum bundle was added to the models (see S1 Table 3).

A false discovery rate (FDR) multiple-comparison correction method was systematically applied for each analysis. Results were considered significant for $p<0.05$, FDR corrected [67]. All statistical analyses were performed using IBM SPSS Statistics v.23 software (IBM Corporation, Armonk, NY, USA).

3. Results

3.1. Sample characteristics

Characteristics of participants are presented in Table 1. Significant age effects on verbal fluency changes were observed for all IST scores (IST 15s, r= -0.212, p=0.022; IST 30s, r= -0.207, p=0.038 and IST 60s, r= -0.252, p=0.019). WMH volumes correlated with age (total WMH: r=0.447, p<0.001; cingulum WMH: r=0.201, p=0.038) and changes in verbal fluency scores (r=-0.218, p=0.038; r=-0.293, p=0.016 and r=-0.328 p= 0.008 respectively). None of clinical variables were related to age, verbal fluency or WMH volumes (p > 0.05). Finally, no association was observed between demographic or clinical variables and cingulum diffusion measures (see S1 Table 1).

Table 1. Characteristics of participants,12-year verbal fluency decline and volumetric variables

Sample n=68	
Sociodemographic variables	Mean ± SD or %
Age	81.2 ± 0.48
Male gender	36.8%
High level of education	37%
Clinical variables	
MMSE	27.3 ± 0.3
CES-D	8.3 ± 0.9
BMI (kg/m^2)	25.2 ± 0.5
Diabetes	10.3%
Hypertension	35.3 %
Verbal fluency decline	
IST at 15 seconds	-0.489 ± 0.02
IST at 30 seconds	-0.697 ± 0.04
IST at 60 seconds	-0.924 ± 0.06
Volumetric variables	
Total WMH volume ml	15.9 ± 4.1
Cingulum bundle volume (% TIV)	0.46 ± 0.06
Cingulum WMH volume (%)	3.2 ± 0.65

MMSE, Mini Mental State Examination; BMI, Body Mass Index; CES-D, Center for Epidemiologic Studies-Depression scale; IST, Isaacs Set Test; WMH, White Matter Hyperintensity.

3.2. Effect of FW-correction on conventional DTI parameters

Cingulum diffusion measures are presented in Table 2. All FW-corrected measures were significantly different

from their conventional counterparts (paired t-test, p<0.05 FDR-corrected, Table 2 and Fig 2). After FW-

correction, a mean increase of 1.52 % of FAt and a mean decrease of 1.61 % of MDt, 2.5% of RDt and 1.08% of

ADt were observed.

Table 2. Cingulum diffusion measures before (conventional DTI) and after FW-correction and % of change

between both measures for each group

Cingulum diffusion measures					
	DTI mean ± SD	**FW-corrected** mean ± SD	**t**	**p-FDR**	**% Change** mean ± SD
FA	0.543 ± 0.049	0.567 ± 0.054	-3.367	0.019*	1.52 ± 0.50
MD (10^{-3} mm^2/s)	0.768 ± 0.0041	0.755 ± 0.0034	-3.196	0.024*	-1.61 ± 0.22
RD (10^{-3} mm^2/s)	0.506 ± 0.0047	0.493 ± 0.0049	-5.315	0.010*	-2.50 ± 0.19
AD (10^{-3} mm^2/s)	1.29 ± 0.0683	1.27 ± 0.0638	-2.975	0.044*	-1.08 ± 0.34
FW	-	0.141 ± 0.017			

* p<0.05 FDR corrected,

Paired Student's t-test

Figure 2. Boxplot charts of the difference in cingulum diffusion measures before (light grey) and after (dark grey)

FW-correction for (a) fractional anisotropy, (b) mean diffusivity, (c) radial diffusivity, (d) axial diffusivity. The

blue line represents a decrease and the red line represents an increase.

3.3. Effect of FW-correction on relationships between cingulum diffusion measures and verbal fluency

3.3.1. Associations with DTI conventional measures

A lower MD value was related to IST decline at 15 and 30 seconds in a model adjusted for age and white matter

volume of the cingulum ($p < 0.05$ FDR-corrected, Table 3). No association was observed with FA, RD and AD.

Table 3. Correlations between cingulum diffusion measures and verbal fluency score (IST) before (conventional

DTI) and after FW-correction

		DTI					
		IST 15s		IST 30s		IST 60s	
		β	R^2	β	R^2	β	R^2
Model unadjusted for total WMH[a]	FA	0.015	0.001	0.047	0.002	0.067	0.004
	MD	**-0.431***	**0.17**	**-0.335***	**0.12**	-0.174	0.07
	AD	-0.186	0.036	-0.135	0.018	-0.060	0.004
	RD	-0.114	0.013	-0.128	0.016	-0.129	0.017
Model adjusted for total WMH[b]	FA	0.022	0.023	0.040	0.052	0.069	0.067
	MD	-0.267	0.020	-0.218	0.010	-0.147	0.09
	AD	-0.172	0.054	-0.109	0.062	-0.029	0.062
	RD	-0.119	0.036	-0.110	0.062	-0.121	0.075

		FW-corrected					
		IST 15s		IST 30s		IST 60s	
		β	R^2	β	R^2	β	R^2
Model unadjusted for total WMH[a]	FAt	0.047	0.003	0.098	0.01	0.095	0.009
	MDt	**-0.443***	**0.19**	**-0.361***	**0.13**	-0.226	0.08
	ADt	-0.198	0.04	-0.134	0.018	-0.061	0.004
	RDt	**-0.293***	**0.11**	**-0.248***	**0.13**	-0.199	0.039
	FW	**-0.353***	**0.12**	**-0.241***	**0.10**	-0.196	0.061
Model adjusted for total WMH[b]	FAt	0.035	0.044	0.072	0.061	0.071	0.07
	MDt	**-0.401***	**0.20**	**-0.341***	**0.17**	-0.189	0.11
	ADt	-0.067	0.10	-0.185	0.10	-0.097	0.08
	RDt	-0.224	0.11	-0.213	0.11	-0.173	0.09
	FW	**-0.333***	**0.15**	**-0.275***	**0.14**	-0.213	0.10

274 [a] β, standardized coefficient regression adjusted for age and cingulum white matter volume

275 [b] β, standardized coefficient regression adjusted for age, cingulum white matter volume and total WMH volume

276 R^2, R square value

277 * $p < 0.05$ FDR corrected

3.3.2. Associations with FW-corrected measures

After FW-correction, high values of MDt and RDt were strongly associated with IST decline at 15 and 30 seconds in a model adjusted for age and white matter volume of the cingulum (p < 0.05, Table 3). No association was observed with FAt and ADt.

3.3.3. Association with cingulum free water content

High free water content was associated with changes in IST score at 15 and 30 seconds in a model adjusted for age and white matter volume of the cingulum (p < 0.05 FDR-corrected, Table 3).

3.4. Impact of WMH burden on the association between cingulum diffusion measures and verbal fluency

In regression models adjusted for total WMH burden, correlations between conventional DTI measures (MD) and IST decline were not significant anymore. In contrast, MDt and free water content correlations were preserved when total WMH volume (Table 3) as well as WMH volume within the cingulum (see S1 Table 3) were added as covariate in regression models (p < 0.05 FDR-corrected, Table 3). In a global model, both diffusion measures remained significantly associated with IST at 15 (FW: β = -0.262 and MDt: β = -0.310) and 30 seconds (FW: β = -0.216 and MDt: β = -0.249, p < 0.05) when adjusted for total WMH volume, indicating that both free water content and MDt independently contributed to the cognitive variances.

4. Discussion

In this study, we examined, over a 12-year period, the effect of free water elimination on conventional DTI measures of white matter within the cingulum tract and the effect of such correction on verbal fluency changes in elderly subjects. In a group of 68 older participants, FW-correction significantly impacted all conventional DTI measures and measure following such correction correlated with decline in verbal fluency performances at 15 and 30 seconds. In addition, free water content was also associated with changes in verbal fluency. Finally, in models adjusted for WMH volumes, correlations between MDt and free water content were preserved.

4.1. Free water elimination effect

In accordance with recent findings in young adults [22], older adults (Chad et al., 2018) and clinical studies [28,68], FW-correction was associated with an increase in FA, and a decrease in diffusivity measures (MD, RD and AD). These changes in DTI measures after elimination of the free water compartment suggested not only a non-null volume of the extracellular water [6,13,28,69], but also indicate that white matter microstructural

changes were less pronounced than previously suggested by conventional DTI measures (Bennett & Madden,

2014; Bennett et al., 2017; Lockhart & DeCarli, 2014; Pelletier et al., 2015).

Based on conventional DTI, our study revealed that only MD showed significant correlations with

changes in verbal fluency, especially at 15 and 30 seconds. The elimination of the free water content confirmed

such strong association for MDt and revealed additional correlations for RDt that could not be fully observed

without considering the free water content. The observed association is in line with previous studies reporting the

role of the cingulum bundle in executive functioning in older adults [50,53,73–76]. Even if the underlying

neurobiological properties of these parameters remain controversial, a high RDt, without any changes in ADt, has

been described as predominantly reflecting myelin changes in animal studies (Song et al., 2003; Sun et al., 2005)

and demyelination severity in human post mortem studies on multiple sclerosis (Klawiter et al., 2011; Schmierer

et al., 2008). This suggests that changes in verbal fluency in our population might be related to myelin changes

within the cingulum, rather than axonal damage (Madden et al., 2012; M. W. Vernooij et al., 2008; Burzynska et

al., 2010). Therefore, our results support previous studies reporting that the elimination of free water improves the

estimation of tissue-specific indices which were more strongly predictive of cognitive changes than conventional

DTI-derived parameters [6,23,26,38,69,77].

4.2. Free water content

In our group of elderly participants, a non-null value of free water content was observed, suggesting that

despite its distance from the ventricles, some free water is likely present in the extracellular space of the cingulum.

This is consistent with recent studies on whole brain white matter reporting a fraction of free water in the elderly

(Albi et al., 2017; Chad et al., 2018; Papadaki et al., 2019). In addition, no association between the free water

content within the cingulum and CSF volumes was observed in our participants (data not shown). However, we

observed that a high content of free water was related to a low cingulum volume, suggesting that in our population

the high free water content may be due to atrophy-related processes rather than CSF contamination (Oestreich et

al., 2016; Wang et al., 2011). In line with previous older adults and clinical studies, we reported that a high content

of free water within the cingulum was strongly associated with changes in verbal fluency performances at 15 and

30 seconds (Bergamino et al., 2017; Ji et al., 2017, 2019; Maillard et al., 2017, 2019). In this regard, in a recent

longitudinal study on 224 elderly subjects, Maillard and colleagues showed that an increase in free water was not

only associated with reduced performances in executive function assessments but was the strongest predictor of

cognitive decline. Taken together, these results support the idea that beyond the elimination of CSF contamination,

free water could provide additional structural information that could constitute a physiological index reflecting

tissue changes in white matter [27,69,79]. However, the underlying microcellular events accounting for the

observed free water content are far from being fully understood; its increase could be related to different pathophysiological processes such as a microvascular degeneration [28,37,80], a reduction in myelin content (Oestreich et al., 2016; Papadaki et al., 2019) or a modulation in the permeability of the blood-brain barrier (Maillard et al., 2017).

4.3. White matter hyperintensities burden effect

The present study showed that when adjusting for WMH volumes, the associations between MDt and free water content and verbal fluency performances at 15 and 30 seconds remained significant, suggesting that both could contribute independently to cognitive impairment in aging. In accordance with these results, recent tractography studies reported that cingulum tracts crossing WMH exhibited significant changes in diffusion measures (Reginold et al., 2016), and suggested that these modifications could extend beyond the WMH lesions (Maillard et al., 2014; Promjunyakul et al., 2016; Reginold et al., 2018). Previous DTI studies also reported that WMH were associated with higher diffusion measures in the normal-appearing white matter [41,43,46,81–83]. Taken together, these results suggest that small focal WMH may lead to both local and distant effects that are large enough to impact white matter diffusion properties of the cingulum tract.

Some methodological limitations should be taken into account when interpreting our findings. First, this study was based on a moderate sample size of healthy elderly participants. However, our population exhibited sufficient inter-individual variability in structural measures and cognitive performances to detect associations between these parameters. Second, the cingulum was analyzed as a single entity despite the fact that it consists of a complex structure containing not only long association fibers, but also short tracts connecting adjacent cortices [84–86]. Finally, the current study was based on diffusion MRI data that was acquired with a single b-value. The algorithm used to fit the free water imaging model involved spatial regularization of data which decreases intra-group variability (Pasternak et al., 2009), and may hide subtle spatial features. Even though comparable results using single- and multi-shell acquisitions have been previously reported [87], advanced acquisitions that include multiple b-values could further increase the accuracy of the free water model [69,87].

To conclude, we reported that FW-correction improves the sensitivity to detect associations between tissue specific diffusion measures of the cingulum and changes in verbal fluency in elderly individuals. Moreover, free water content *per se* appears to be a relevant parameter to describe age-related modifications of the white matter and its association with executive functioning. These observations suggest the importance of considering the free water compartment for DTI measures in aging.

Funding

This research was funded by the FONDATION VAINCRE ALZHEIMER (#14733, www.vaincrealzheimer.org). This study was considered an emerging project and therefore partly funded as part of the laboratory of Excellence TRAIL ANR-10-LABX-57 (trail.labex.u-bordeaux.fr). The Three-City study is conducted under a partnership agreement between the Institut National de la Santé et de la Recherche Médicale (INSERM, www.inserm.fr), the University Bordeaux 2 Victor Segalen (www.u-bordeaux.fr/) and Sanofi-Aventis (www.sanofi.fr/). The Fondation pour la Recherche Médicale funded the planning and initiation of the study (www.frm.org/). The Three-City study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés (www.ameli.fr/), Direction Générale de la Santé MGEN (solidarites-sante.gouv.fr/), Institut de la Longévité (www.ilvv.fr), Conseils Régionaux d'Aquitaine et Bourgogne, Fondation de France (www.fondationdefrance.org), Ministry of Research-INSERM Programme "Cohortes et collections de données biologiques", Agence Nationale de la Recherche ANR PNRA 2006 and LongVie 2007 (www.anr.fr), the "Fondation Plan Alzheimer" (FCS 2009-2012) and the Caisse Nationale de Solidarité pour l'Autonomie (CNSA, www.cnsa.fr). Part of this research was also supported by the Fonds de recherche du Québec – Nature et technologies (FRQNT, www.frqnt.gouv.qc.ca), the NSERC Discovery grant (www.nserc-crsng.gc.ca), the Université de Sherbrooke institutional chair in neuroinformatics from Pr Descoteaux (www.usherbrooke.ca) and the Mitacs Accelerate program (www.mitacs.ca).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Studies involving human participants All procedures were conducted with approval from the ethics committee of Kremlin Bicêtre Hospital (Paris, France)

Informed consent All participants provided written informed consent for participation in this study.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author contributions

Conceptualization: ME, GC; Formal analysis: ME; Funding acquisition: BD, CH, JFD, HA, MA, GC; Methodology: ME, GT, BD; Software: GT, FR, MD; Supervision: MD, GC; Visualization: ME, GT; Writing -

original draft: ME, GT; Writing - review & editing: all authors review the article. All authors provided critical

feedback and approved the final manuscript.

10. References

1. Damoiseaux JS. Effects of aging on functional and structural brain connectivity. Neuroimage. 15 oct 2017;160:32-40.

2. Ikram MA, Vrooman HA, Vernooij MW, van der Lijn F, Hofman A, van der Lugt A, et al. Brain tissue volumes in the general elderly population: The Rotterdam Scan Study. Neurobiology of Aging. 1 juin 2008;29(6):882-90.

3. Yang Z, Wen W, Jiang J, Crawford JD, Reppermund S, Levitan C, et al. Age-associated differences on structural brain MRI in nondemented individuals from 71 to 103 years. Neurobiology of Aging. 1 avr 2016;40:86-97.

4. Sexton CE, Walhovd KB, Storsve AB, Tamnes CK, Westlye LT, Johansen-Berg H, et al. Accelerated changes in white matter microstructure during aging: a longitudinal diffusion tensor imaging study. J Neurosci. 12 nov 2014;34(46):15425-36.

5. Vernooij MW, Ikram MA, Vrooman HA, Wielopolski PA, Krestin GP, Hofman A, et al. White matter microstructural integrity and cognitive function in a general elderly population. Arch Gen Psychiatry. mai 2009;66(5):545-53.

6. Madden DJ, Bennett IJ, Burzynska A, Potter GG, Chen N, Song AW. Diffusion tensor imaging of cerebral white matter integrity in cognitive aging. Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease. 1 mars 2012;1822(3):386-400.

7. de Groot M, Ikram MA, Akoudad S, Krestin GP, Hofman A, van der Lugt A, et al. Tract-specific white matter degeneration in aging: the Rotterdam Study. Alzheimers Dement. mars 2015;11(3):321-30.

8. Vinke EJ, de Groot M, Venkatraghavan V, Klein S, Niessen WJ, Ikram MA, et al. Trajectories of imaging markers in brain aging: the Rotterdam Study. Neurobiology of Aging. 1 nov 2018;71:32-40.

9. Arvanitakis Z, Fleischman DA, Arfanakis K, Leurgans SE, Barnes LL, Bennett DA. Association of white matter hyperintensities and gray matter volume with cognition in older individuals without cognitive impairment. Brain Struct Funct. 2016;221(4):2135-46.

10. Bendlin BB, Fitzgerald ME, Ries ML, Xu G, Kastman EK, Thiel BW, et al. White matter in aging and cognition: a cross-sectional study of microstructure in adults aged eighteen to eighty-three. Dev Neuropsychol. 2010;35(3):257-77.

11. Borghesani PR, Madhyastha TM, Aylward EH, Reiter MA, Swarny BR, Schaie KW, et al. The association between higher order abilities, processing speed, and age are variably mediated by white matter integrity during typical aging. Neuropsychologia. juill 2013;51(8):1435-44.

12. Wiseman SJ, Booth T, Ritchie SJ, Cox SR, Muñoz Maniega S, Valdés Hernández MDC, et al. Cognitive abilities, brain white matter hyperintensity volume, and structural network connectivity in older age. Hum Brain Mapp. 2018;39(2):622-32.

13. Pasternak O, Sochen N, Gur Y, Intrator N, Assaf Y. Free water elimination and mapping from diffusion MRI. Magn Reson Med. sept 2009;62(3):717-30.

14. Jbabdi S, Johansen-Berg H. Tractography: where do we go from here? Brain Connect. 2011;1(3):169-83.

15. Kurki TJI, Heiskanen LAA. Diffusion parameters of the core of cingulum are associated with age-related ventricular enlargement: a diffusion tensor tractography study. Neuroradiology. 1 oct 2018;60(10):1013-8.

16. Salminen LE, Conturo TE, Bolzenius JD, Cabeen RP, Akbudak E, Paul RH. REDUCING CSF PARTIAL
VOLUME EFFECTS TO ENHANCE DIFFUSION TENSOR IMAGING METRICS OF BRAIN
MICROSTRUCTURE. Technol Innov. avr 2016;18(1):5‑20.

17. Vos SB, Jones DK, Viergever MA, Leemans A. Partial volume effect as a hidden covariate in DTI analyses.
NeuroImage. 15 avr 2011;55(4):1566‑76.

18. Metzler-Baddeley C, Jones DK, Steventon J, Westacott L, Aggleton JP, O'Sullivan MJ. Cingulum
Microstructure Predicts Cognitive Control in Older Age and Mild Cognitive Impairment. J Neurosci. 5 déc
2012;32(49):17612‑9.

19. Pfefferbaum A, Sullivan EV. Increased brain white matter diffusivity in normal adult aging: Relationship to
anisotropy and partial voluming. Magnetic Resonance in Medicine. 2003;49(5):953‑61.

20. Concha L, Gross DW, Beaulieu C. Diffusion Tensor Tractography of the Limbic System. American Journal
of Neuroradiology. 1 oct 2005;26(9):2267‑74.

21. Jones DK, Cercignani M. Twenty-five pitfalls in the analysis of diffusion MRI data. NMR Biomed. août
2010;23(7):803‑20.

22. Sepehrband F, Cabeen RP, Choupan J, Barisano G, Law M, Toga AW, et al. Perivascular space fluid
contributes to diffusion tensor imaging changes in white matter. Neuroimage. 15 2019;197:243‑54.

23. Albi A, Pasternak O, Minati L, Marizzoni M, Bartrés-Faz D, Bargalló N, et al. Free water elimination
improves test-retest reproducibility of diffusion tensor imaging indices in the brain: A longitudinal multisite
study of healthy elderly subjects. Hum Brain Mapp. 2017;38(1):12‑26.

24. Pasternak O, Westin C-F, Bouix S, Seidman LJ, Goldstein JM, Woo T-UW, et al. Excessive extracellular
volume reveals a neurodegenerative pattern in schizophrenia onset. J Neurosci. 28 nov
2012;32(48):17365‑72.

25. Bergamino M, Kuplicki R, Victor TA, Cha Y-H, Paulus MP. Comparison of two different analysis
approaches for DTI free-water corrected and uncorrected maps in the study of white matter microstructural
integrity in individuals with depression. Hum Brain Mapp. 2017;38(9):4690‑702.

26. Dumont M, Roy M, Jodoin P-M, Morency FC, Houde J-C, Xie Z, et al. Free water in white matter
differentiates MCI and AD from control subjects. bioRxiv. 31 janv 2019;537092.

27. Hoy AR, Ly M, Carlsson CM, Okonkwo OC, Zetterberg H, Blennow K, et al. Microstructural white matter
alterations in preclinical Alzheimer's disease detected using free water elimination diffusion tensor imaging.
PLoS ONE. 2017;12(3):e0173982.

28. Ji F, Pasternak O, Liu S, Loke YM, Choo BL, Hilal S, et al. Distinct white matter microstructural
abnormalities and extracellular water increases relate to cognitive impairment in Alzheimer's disease with
and without cerebrovascular disease. Alzheimers Res Ther. 17 août 2017;9(1):63.

29. Ofori E, Krismer F, Burciu RG, Pasternak O, McCracken JL, Lewis MM, et al. Free water improves
detection of changes in the substantia nigra in parkinsonism: A multisite study. Mov Disord. oct
2017;32(10):1457‑64.

30. Tanner JJ, Amin M, Hardcastle C, Parvataneni H, Vaillancourt DE, Mareci TH, et al. Better Brain and
Cognition Prior to Surgery Is Associated With Elevated Postoperative Brain Extracellular Free-Water in
Older Adults. Front Aging Neurosci [Internet]. 2019 [cité 29 mai 2019];11. Disponible sur:
https://www.frontiersin.org/articles/10.3389/fnagi.2019.00117/full?utm_source=F-
AAE&utm_medium=EMLF&utm_campaign=MRK_999212_55_Neuros_20190528_arts_A

31. Aboitiz F, Rodriguez E, Olivares R, Zaidel E. Age-related changes in fibre composition of the human corpus
callosum. Neuroreport. 1 juill 1996;7(11):1761‑4.

32. Barkhof F, Fox NC, Bastos-Leite AJ, Scheltens P. Normal Ageing. In: Barkhof F, Fox NC, Bastos-Leite AJ,
Scheltens P, éditeurs. Neuroimaging in Dementia [Internet]. Berlin, Heidelberg: Springer Berlin Heidelberg;
2011 [cité 5 oct 2019]. p. 43‑57. Disponible sur: https://doi.org/10.1007/978-3-642-00818-4_4

33.   Meier-Ruge W, Ulrich J, Brühlmann M, Meier E. Age-related white matter atrophy in the human brain. Ann
N Y Acad Sci. 26 déc 1992;673:260-9.

34.   Tang Y, Nyengaard JR, Pakkenberg B, Gundersen HJG. Age-Induced White Matter Changes in the Human
Brain: A Stereological Investigation. Neurobiology of Aging. 1 nov 1997;18(6):609-15.

35.   Alber J, Alladi S, Bae H-J, Barton DA, Beckett LA, Bell JM, et al. White matter hyperintensities in vascular
contributions to cognitive impairment and dementia (VCID): Knowledge gaps and opportunities.
Alzheimers Dement (N Y). 2019;5:107-17.

36.   Duering M, Finsterwalder S, Baykara E, Tuladhar AM, Gesierich B, Konieczny MJ, et al. Free water
determines diffusion alterations and clinical status in cerebral small vessel disease. Alzheimers Dement. juin
2018;14(6):764-74.

37.   Maillard P, Mitchell GF, Himali JJ, Beiser A, Fletcher E, Tsao CW, et al. Aortic Stiffness, Increased White
Matter Free Water, and Altered Microstructural Integrity: A Continuum of Injury. Stroke.
2017;48(6):1567-73.

38.   Maillard P, Fletcher E, Singh B, Martinez O, Johnson DK, Olichney JM, et al. Cerebral white matter free
water: A sensitive biomarker of cognition and function. Neurology. 7 mai 2019;92(19):e2221-31.

39.   Habes M, Erus G, Toledo JB, Zhang T, Bryan N, Launer LJ, et al. White matter hyperintensities and imaging
patterns of brain ageing in the general population. Brain. avr 2016;139(4):1164-79.

40.   Leeuw F-E de, Groot JC de, Achten E, Oudkerk M, Ramos LMP, Heijboer R, et al. Prevalence of cerebral
white matter lesions in elderly people: a population based magnetic resonance imaging study. The Rotterdam
Scan Study. Journal of Neurology, Neurosurgery & Psychiatry. 1 janv 2001;70(1):9-14.

41.   Maillard P, Fletcher E, Lockhart SN, Roach AE, Reed B, Mungas D, et al. White Matter Hyperintensities
and their Penumbra Lie Along a Continuum of Injury In The Aging Brain. Stroke. juin 2014;45(6):1721-6.

42.   Maniega SM, Valdés Hernández MC, Clayden JD, Royle NA, Murray C, Morris Z, et al. White matter
hyperintensities and normal-appearing white matter integrity in the aging brain. Neurobiol Aging. févr
2015;36(2):909-18.

43.   Pelletier A, Periot O, Dilharreguy B, Hiba B, Bordessoules M, Chanraud S, et al. Age-Related Modifications
of Diffusion Tensor Imaging Parameters and White Matter Hyperintensities as Inter-Dependent Processes.
Front Aging Neurosci. 2015;7:255.

44.   Reginold W, Sam K, Poublanc J, Fisher J, Crawley A, Mikulis DJ. Impact of white matter hyperintensities
on surrounding white matter tracts. Neuroradiology. sept 2018;60(9):933-44.

45.   Reginold W, Itorralba J, Luedke AC, Fernandez-Ruiz J, Reginold J, Islam O, et al. Tractography at 3T MRI
of Corpus Callosum Tracts Crossing White Matter Hyperintensities. American Journal of Neuroradiology.
1 sept 2016;37(9):1617-22.

46.   Svärd D, Nilsson M, Lampinen B, Lätt J, Sundgren PC, Stomrud E, et al. The effect of white matter
hyperintensities on statistical analysis of diffusion tensor imaging in cognitively healthy elderly and
prodromal Alzheimer's disease. PLOS ONE. 21 sept 2017;12(9):e0185239.

47.   Rizvi B, Lao PJ, Colón J, Hale C, Igwe KC, Narkhede A, et al. Tract-defined regional white matter
hyperintensities and memory. NeuroImage: Clinical. 1 janv 2020;25:102143.

48.   Jones DK, Christiansen KF, Chapman RJ, Aggleton JP. Distinct subdivisions of the cingulum bundle
revealed by diffusion MRI fibre tracking: Implications for neuropsychological investigations.
Neuropsychologia. janv 2013;51(1):67.

49.   Amieva H, Le Goff M, Millet X, Orgogozo JM, Pérès K, Barberger-Gateau P, et al. Prodromal Alzheimer's
disease: successive emergence of the clinical symptoms. Ann Neurol. nov 2008;64(5):492-8.

50.   Catheline G, Periot O, Amirault M, Braun M, Dartigues J-F, Auriacombe S, et al. Distinctive alterations of
the cingulum bundle during aging and Alzheimer's disease. Neurobiology of Aging. 1 sept
2010;31(9):1582-92.

51. Ezzati A, Katz MJ, Lipton ML, Zimmerman ME, Lipton RB. Hippocampal volume and cingulum bundle
fractional anisotropy are independently associated with verbal memory in older adults. Brain Imaging
Behav. sept 2016;10(3):652-9.

52. Kantarci K, Senjem ML, Avula R, Zhang B, Samikoglu AR, Weigand SD, et al. Diffusion tensor imaging
and cognitive function in older adults with no dementia. Neurology. 5 juill 2011;77(1):26-34.

53. Seiler S, Fletcher E, Hassan-Ali K, Weinstein M, Beiser A, Himali JJ, et al. Cerebral tract integrity relates
to white matter hyperintensities, cortex volume, and cognition. Neurobiology of Aging. 1 déc
2018;72:14-22.

54. Sexton CE, Mackay CE, Lonie JA, Bastin ME, Terrière E, O'Carroll RE, et al. MRI correlates of episodic
memory in Alzheimer's disease, mild cognitive impairment, and healthy aging. Psychiatry Res. 30 oct
2010;184(1):57-62.

55. 3C Study Group. Vascular factors and risk of dementia: design of the Three-City Study and baseline
characteristics of the study population. Neuroepidemiology. déc 2003;22(6):316-25.

56. Isaacs B, Kennie AT. The Set test as an aid to the detection of dementia in old people. Br J Psychiatry. oct
1973;123(575):467-70.

57. Radloff LS. The CES-D Scale: A Self-Report Depression Scale for Research in the General Population.
Applied Psychological Measurement. 1 juin 1977;1(3):385-401.

58. Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. Probabilistic diffusion tractography with
multiple fibre orientations: What can we gain? Neuroimage. 1 janv 2007;34(1):144-55.

59. Behrens TEJ, Woolrich MW, Jenkinson M, Johansen-Berg H, Nunes RG, Clare S, et al. Characterization
and propagation of uncertainty in diffusion-weighted MR imaging. Magn Reson Med. nov
2003;50(5):1077-88.

60. Smith SM. Fast robust automated brain extraction. Hum Brain Mapp. nov 2002;17(3):143-55.

61. Descoteaux M, Wiest-Daesslé N, Prima S, Barillot C, Deriche R. Impact of Rician adapted Non-Local
Means filtering on HARDI. Med Image Comput Comput Assist Interv. 2008;11(Pt 2):122-30.

62. Descoteaux M, Angelino E, Fitzgibbons S, Deriche R. Regularized, fast, and robust analytical Q-ball
imaging. Magn Reson Med. sept 2007;58(3):497-510.

63. Tournier J-D, Calamante F, Connelly A. Robust determination of the fibre orientation distribution in
diffusion MRI: Non-negativity constrained super-resolved spherical deconvolution. NeuroImage. 1 mai
2007;35(4):1459-72.

64. Theaud G, Houde J-C, Boré A, Rheault F, Morency F, Descoteaux M. TractoFlow: A robust, efficient and
reproducible diffusion MRI pipeline leveraging Nextflow & Singularity. bioRxiv. 9 mai 2019;631952.

65. Schmidt P, Gaser C, Arsic M, Buck D, Förschler A, Berthele A, et al. An automated tool for detection of
FLAIR-hyperintense white-matter lesions in Multiple Sclerosis. NeuroImage. 15 févr 2012;59(4):3774-83.

66. Garyfallidis E, Côté M-A, Rheault F, Sidhu J, Hau J, Petit L, et al. Recognition of white matter bundles
using local and global streamline-based registration and clustering. NeuroImage. 15 avr 2018;170:283-95.

67. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to
Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological). 1995;57(1):289-300.

68. Bergamino M, Pasternak O, Farmer M, Shenton ME, Paul Hamilton J. Applying a free-water correction to
diffusion imaging data uncovers stress-related neural pathology in depression. NeuroImage: Clinical. 1 janv
2016;10:336-42.

69. Hoy AR, Kecskemeti SR, Alexander AL. Free water elimination diffusion tractography: A comparison with
conventional and fluid-attenuated inversion recovery, diffusion tensor imaging acquisitions. Journal of
Magnetic Resonance Imaging. 2015;42(6):1572-81.

70. Bennett IJ, Madden DJ. Disconnected aging: Cerebral white matter integrity and age-related differences in
cognition. Neuroscience. 12 sept 2014;276:187-205.

71. Bennett IJ, Greenia DE, Maillard P, Sajjadi SA, DeCarli C, Corrada MM, et al. Age-related white matter
integrity differences in oldest-old without dementia. Neurobiology of Aging. 1 août 2017;56:108-14.

72. Lockhart SN, DeCarli C. Structural imaging measures of brain aging. Neuropsychol Rev. sept
2014;24(3):271-89.

73. Bettcher BM, Mungas D, Patel N, Elofson J, Dutt S, Wynn M, et al. Neuroanatomical substrates of executive
functions: Beyond prefrontal structures. Neuropsychologia. 1 mai 2016;85:100-9.

74. Charlton RA, Barrick TR, Lawes INC, Markus HS, Morris RG. White matter pathways associated with
working memory in normal aging. Cortex. 1 avr 2010;46(4):474-89.

75. Chiang H-L, Chen Y-J, Shang C-Y, Tseng W-YI, Gau SS-F. Different neural substrates for executive
functions in youths with ADHD: a diffusion spectrum imaging tractography study. Psychological Medicine.
avr 2016;46(6):1225-38.

76. Sasson E, Doniger GM, Pasternak O, Tarrasch R, Assaf Y. White matter correlates of cognitive domains in
normal aging with diffusion tensor imaging. Front Neurosci [Internet]. 2013 [cité 7 nov 2018];7. Disponible
sur: https://www.frontiersin.org/articles/10.3389/fnins.2013.00032/full

77. Lyall AE, Pasternak O, Robinson DG, Newell D, Trampush JW, Gallego JA, et al. Greater Extracellular
Free Water in First-Episode Psychosis Predicts Better Neurocognitive Functioning. Mol Psychiatry. mars
2018;23(3):701-7.

78. Ji F, Pasternak O, Ng KK, Chong JSX, Liu S, Zhang L, et al. White matter microstructural abnormalities
and default network degeneration are associated with early memory deficit in Alzheimer's disease
continuum. Sci Rep. 18 mars 2019;9(1):4749.

79. Chad JA, Pasternak O, Salat DH, Chen JJ. Re-examining age-related differences in white matter
microstructure with free-water corrected diffusion tensor imaging. Neurobiol Aging. nov 2018;71:161-70.

80. Rydhög AS, Szczepankiewicz F, Wirestam R, Ahlgren A, Westin C-F, Knutsson L, et al. Separating blood
and water: Perfusion and free water elimination from diffusion MRI in the human brain. Neuroimage. 01
2017;156:423-34.

81. Lange RT, Shewchuk JR, Heran MKS, Rauscher A, Jarrett M, Brubacher JR, et al. To exclude or not to
exclude: further examination of the influence of white matter hyperintensities in diffusion tensor imaging
research. J Neurotrauma. 15 janv 2014;31(2):198-205.

82. Leritz EC, Shepel J, Williams VJ, Lipsitz LA, McGlinchey RE, Milberg WP, et al. Associations between
T1 white matter lesion volume and regional white matter microstructure in aging. Hum Brain Mapp. mars
2014;35(3):1085-100.

83. Promjunyakul N-O, Lahna DL, Kaye JA, Dodge HH, Erten-Lyons D, Rooney WD, et al. Comparison of
cerebral blood flow and structural penumbras in relation to white matter hyperintensities: A multi-modal
magnetic resonance imaging study. J Cereb Blood Flow Metab. 2016;36(9):1528-36.

84. Bubb EJ, Metzler-Baddeley C, Aggleton JP. The cingulum bundle: Anatomy, function, and dysfunction.
Neurosci Biobehav Rev. 10 mai 2018;

85. Catani M, Thiebaut de Schotten M. A diffusion tensor imaging tractography atlas for virtual in vivo
dissections. Cortex. sept 2008;44(8):1105-32.

86. Lawes INC, Barrick TR, Murugam V, Spierings N, Evans DR, Song M, et al. Atlas-based segmentation of
white matter tracts of the human brain using diffusion tensor tractography and comparison with classical
dissection. Neuroimage. 1 janv 2008;39(1):62-79.

87. Pasternak O, Shenton ME, Westin C-F. Estimation of extracellular volume from regularized multi-shell
diffusion MRI. Med Image Comput Comput Assist Interv. 2012;15(Pt 2):305-12.

13. Figures

Figure 1

Figure 2