Quantitative Structure-Mutation-Activity Relationship Tests (QSMART) Model for Protein Kinase Inhibitor Response Prediction

Liang-Chin Huang[1], Wayland Yeung[1], Ye Wang[2], Huimin Cheng[2], Aarya Venkat[3], Sheng Li[4], Ping Ma[2], Khaled Rasheed[4], Natarajan Kannan[1,3*]

1 Institute of Bioinformatics, University of Georgia, Athens, GA, USA
2 Department of Statistics, University of Georgia, Athens, GA, USA
3 Department of Biochemistry and Molecular Biology, University of Georgia, Athens, GA, USA
4 Department of Computer Science, University of Georgia, Athens, GA, USA

* nkannan@uga.edu

Abstract

Predicting how mutations impact drug sensitivity is a major challenge in personalized medicine. Although several machine learning models have been developed to predict drug sensitivity from gene expression and genomic profiles, these methods do not explicitly incorporate the structural properties of drug-mutation interactions to understand the molecular mechanisms of drug resistance/sensitivity. To facilitate the understanding of how the drug-mutation interactions quantitatively contribute to drug response, we developed a framework that not only estimates IC_{50} with high accuracy ($R^2 = 0.861$ and RMSE = 0.818) but also identifies features contributing to the accuracy, thereby enhancing explainability. Our framework uses a multi-component approach that includes (1) collecting drug fingerprints, cancer cell line's multi-omics features, and drug responses, (2) testing the statistical significance of interaction effects, (3) selecting features by Lasso with Bayesian information criterion, and (4) using neural networks to predict drug response. We validate each component in the proposed framework and explain the biological relevance and mathematical interpretation of pertinent features, including afatinib- and lapatinib-EGFR L858R interactions, in a non-small cell lung cancer case study. This is the first study to systematically explain drug response in cancer cell lines by investigating the contribution of interaction effects, such as protein-protein interactions and drug-mutation interactions. The concept of our proposed framework can also be applied to other prediction models with the interaction effects of interest, such as drug-drug interaction and agent-host interaction.

Author summary

In recent years, artificial intelligence (AI) has been successfully used in image analysis, natural language processing, and to solve strategy games. People are also interested in implementing AI in the medical field, such as personalized medicine and recommender system, the goals of which are respectively to customize the treatment based on the patient's genomic profile and to support doctors in making a proper decision for drug prescription. However, AI's "black box" issue impedes doctors and pharmaceutical scientists from accepting results from an unexplainable model. To this end, we proposed

a framework to facilitate increasing the explainability of predicting drug response in cancer cells. This framework combines neural networks with traditional statistical tests outside the black box to achieve high prediction accuracy while also identifying informative multi-omics predictors and drug-target interactions, thereby increasing the model's explainability. Compared to previous studies, our framework is one of the most accurate methods to predict drug response. Moreover, in this study, we illustrate several examples of how the predictors' biological relevance and their interactions impact drug response in non-small cell lung cancer cells, which reflect both the novelty and utility of this approach.

Introduction

Protein kinases are a class of signaling proteins, greatly valued as therapeutic targets for their key roles in human diseases, such as cancer [1]. For decades, chemotherapy has served as part of a standard set of cancer treatments; however, the resistance of cancer cells to chemotherapy is still a major clinical problem and remains a challenging task [2]. Protein kinase mutations are known to play important roles not only in drug resistance [3] but also in drug sensitivity [4]; even mutations occurring in the same protein kinase can have diverse drug responses. For example, non-small cell lung cancer (NSCLC) cells with EGFR T790M or L858R mutation are respectively resistant or hypersensitive to both gefitinib and erlotinib [5,6], while those with EGFR T790M/L858R double mutants are resistant to both gefitinib and erlotinib [7]. As the efficacy of different cancer drugs is affected by these mutations, there is a need to systematically explain how drug-mutation associations quantitatively contribute to drug response in cancer cells.

To facilitate the understanding of the molecular mechanisms that cause drug sensitivity and drug resistance in cancer cells, the Genomics of Drug Sensitivity in Cancer (GDSC) Project [8] recently screened the drug responses of 266 anti-cancer drugs against ∼1,000 human cancer cell lines and provided the largest publicly available drug response dataset. Moreover, to broaden the pharmacologic annotation for human cancers, the Cancer Cell Line Encyclopedia [9] (CCLE) provided pharmacologic profiles for 24 drugs across 504 cancer cell lines. By utilizing these datasets, several prediction models were built to pursue a more precise drug response estimation by different types of approaches, from traditional statistical models, network-based models, to the recent machine learning methods and state-of-the-art neural networks (Table 1). These approaches include (1) statistical models: MANOVA [10] and generalized linear models (regularization: ridge [11–14], elastic net [11–13,15,16], Lasso [11–13], and mixture [17]), (2) network-based models [18–24], (3) random forests [25,26], (4) support vector machine (SVM) [22,27,28] and other kernelized methods [29–31], and (5) neural networks: artificial neural network (ANN) [32], convolutional neural network (CNN) [33–35], recurrent neural network (RNN) [35], and other deep neural networks (DNN) [15,36].

Over the years, new techniques continue to emerge and the samples of drug response have increased constantly; nevertheless, existing prediction models still cannot achieve high performance to realize "precision" medicine goals. Their prediction performances measured by the coefficient of determination (R^2) are in the range from 0.25 to 0.78. Until very recently, CDRscan [33], tCNNS [34], and MCA [35] achieve R^2 higher than 0.8 (R^2 = 0.84, 0.83, and 0.86, respectively) by using complicated deep neural networks with considerable hidden layers. Although they achieve high prediction performance, all of them hinder the explanation of detailed drug-cancer cell interactions by using convolutional drug and cell line features before performing "virtual docking", the hidden layer where both types of features converge [33]. Moreover, most of the cancer cell line features used in previous studies were gene-level or higher-level features, instead

Table 1. Current drug response prediction approaches.

Date	Author	Model (Comparative model)	Cancer cell line feature	Drug response		Validation	Performance
				GDSC	CCLE		
2013.04.30	Menden et al. [32]	ANN (RF)	MUT, CNV	✓		8-fold CV	$R^2 = 0.72$
2014.03.03	Geeleher et al. [11]	GLM	EXP	✓		LOOCV	AUC = 0.81
2015.01.01	Jang et al. [12]	GLM (PLS, SVM, PCA, RF)	MUT, EXP, CNV, CLS	✓	✓	5-fold CV	r = ~0.5
2015.06.30	Dong et al. [27]	SVM	EXP		✓	10-fold CV	Accuracy = ~0.8
2015.09.29	Zhang et al. [18]	Network (EN)	EXP	✓	✓	LOOCV	r = 0.6
2016.03.31	Gupta et al. [28]	SVM	MUT, EXP, CNV		✓	LOOCV	r = 0.78
2016.09.01	Ammad-ud-din et al. [29]	Kernel (GLM)	PWY	✓		5-fold CV	ρ = ~0.22
2016.12.28	Nguyen et al. [10]	MANOVA (RF)	EXP	✓		10-fold CV	MCC = 0.18
2017.01.09	Stanfield et al. [19]	Network (Kernel)	MUT, PPI	✓	✓	LOOCV	AUC = 0.881
2017.07.15	Ammad-ud-din et al. [13]	GLM (PLS, SGL, RF, SVM)	EXP, PWY	✓		LOOCV	ρ = 0.375
2017.08.28	Geeleher et al. [14]	Ridge	EXP	✓		10-fold CV	ρ = 0.48
2017.09.12	Rahman et al. [25]	RF	EXP	✓	✓	3-fold CV	AUC = ~0.3
2017.11.13	Ding et al. [15]	EN, DNN (SVM)	MUT, EXP, CNV	✓	✓	25-fold CV	AUC = 0.87
2018.03.08	He et al. [30]	Kernel (EN, Ridge, RF)	EXP	✓		3-fold CV	Precision = ~0.35
2018.06.11	Chang et al. [33]	CNN (RF, SVM)	SNP	✓		5% leave-out	$R^2 = 0.843$
2018.07.01	Cichonska et al. [31]	Kernel	SNP, MET, EXP, CNV	✓		10-fold CV	r = 0.858
2018.09.14	Le et al. [20]	Network (Kernel)	MUT, EXP	✓	✓	5-fold CV	r = 0.804
2018.09.14	Juan-Blanco et al. [21]	Network	MUT, EXP		✓	LOOCV	AUC = ~0.72
2018.10.10	Yang et al. [22]	Network, SVM (Kernel)	MUT, MET, CNV, PPI	✓		5-fold CV	AUC = 0.788
2018.12.07	Liu et al. [23]	Network	EXP	✓	✓	10-fold CV	r = 0.73
2019.01.22	Wei et al. [24]	Network	EXP	✓	✓	LOOCV	r = 0.63
2019.01.31	Wang et al. [16]	EN	EXP, PWY		✓	10-fold CV	MSE = ~2.8
2019.01.31	Chiu et al. [36]	DNN (LR, SVM, PCA)	MUT, EXP	✓		10% leave-out	r = ~0.86
2019.02.27	Li et al. [17]	Mixture (GLM, RF)	EXP		✓	20% leave-out	r = 0.882
2019.07.11	Lind et al. [26]	RF (SVM, ANN)	MUT	✓		5-fold CV	r = 0.86
2019.07.29	Liu et al. [34]	CNN (ANN)	MUT, CNV	✓		10% leave-out	$R^2 = 0.826$
2019.10.16	Manica et al. [35]	CNN, RNN (RF, SVM)	EXP, CNV, PPI	✓		5-fold CV	$R^2 = 0.86$

ANN: artificial neural network; AUC: area under the ROC curve; CCLE: Cancer Cell Line Encyclopedia; CLS: cancer classification; CNN: convolutional neural network; CNV: copy number variation; CV: cross-validation; EN: elastic net; EXP: gene expression; GDSC: Genomics of Drug Sensitivity in Cancer; GLM: generalized linear model, including ridge, elastic net, and lasso regression; DNN: deep neural networks; LOOCV: leave-one-out cross-validation; LR: linear regression; MCC: Matthews correlation coefficient; MET: methylation; MSE: mean squared error; MUT: gene-level mutation (i.e. whether the gene is mutated or not); PCA: principal component analysis; PLS: partial least squares; PPI: protein-protein interaction; PWY: pathway; r: Pearson correlation coefficient; R^2: coefficient of determination; RF: random forests; ρ: Spearman's rank correlation coefficient; RNN: recurrent neural network; SGL: sparse group lasso; SNP: single nucleotide polymorphism; SVM: support vector machine.

of residue-level features, such as single nucleotide polymorphisms (Table 1). Therefore, the impact of drug target mutation on detailed drug-target binding mechanisms is not available from their prediction models.

The trade-off between prediction performance and explainability is an issue not only for CDRscan, tCNNS, and MCA but also for other existing machine learning approaches, thus the Defense Advanced Research Projects Agency (DARPA) recently launched the Explainable Artificial Intelligence (XAI) program [37] to facilitate building explainable models while maintaining prediction performance. In recognition of the interest in building explainable AI models, we built the Quantitative Structure-Mutation-Activity Relationship Tests (QSMART) model by (1) introducing more explainable drug-mutation interaction effects to the quantitative structure-activity relationship (QSAR) model, (2) using traditional statistical tests to identify significant interactions, and (3) utilizing a feature selection method to obtain highly informative features (Fig 1). This is equivalent to moving two hidden layers outside the neural networks "black box" for increasing the prediction model's explainability. Combining with neural networks, our proposed framework also kept prediction performance for precisely predicting protein kinase inhibitors (PKIs) response in cancer cells (overall $R^2 = 0.861$, AUC = 0.981, and RMSE = 0.818 based on 10-fold cross-validation). MCA [35] also

achieves the same level of prediction performance, but its performance of PKI response prediction is $R^2 = 0.823$ (Table 2 and S1 Data). Although building fully explainable models is not the goal of this study, our framework can not only provide researchers with more opportunities to explain potential mechanisms of drug resistance/sensitivity from statistically significant drug-mutation interaction effects but also improve drug response prediction for applications in precision medicine and drug discovery.

Fig 1. The framework of using the QSMART model with neural networks to predict protein kinase inhibitor response in cancer cell lines. Four main components of this framework: (1) drug features, cancer cell line features, and drug responses, (2) statistics tests for interaction effects, (3) a feature selection method for identifying highly informative features, and (4) a machine learning method for predicting drug response.

Results

The framework for protein kinase inhibitor response prediction

The overall objective of this study is to emphasize the contribution of adding drug-mutation interaction terms to a drug response prediction model and to show how these interaction terms could help explain the mechanism of drug resistance/sensitivity. The framework we proposed in this study includes four main components: (1) PKIs' chemical descriptors, cancer cell line's multi-omics data, and PKI responses, (2) F-test for identifying significant drug-mutation interaction effects, (3) a feature selection method: Lasso with Bayesian information criterion (BIC) control, and (4) a machine learning method to predict PKI response: neural networks (Fig 1). This framework has flexibility in adapting different materials and methods in each component. To implement

this framework, we collected ∼0.2 million drug response (IC_{50} in a logarithmic scale; "IC_{50}" thereafter) dataset from GDSC, and then split them into 23 sub-datasets for building cancer-centric models. The overall prediction performance of our proposed framework and the evaluation of each component's contribution are described below.

The overall performance of QSMART model with neural networks

The number of PKI responses, the total number of features (including drug features, cancer cell line features, and interaction features) in the prediction model, the number of nodes in the first and second hidden layers of neural networks, and prediction performance (R^2) for each cancer type are shown in Table 2. More measurements of prediction performance (RMSE and AUC) and detailed numbers of cancer cell line features at seven feature levels, five types of interaction effects, and tours (training iterations) are shown in S1 Table. By using the features from the QSMART model and neural networks, we have the ability to precisely predict PKI response in 23 cancer types ($R^2 = 0.805$ to 0.880). Fig 2a presents an actual IC_{50} vs. predicted IC_{50} plot for all types of cancer cell lines (overall RMSE = 0.818 and $R^2 = 0.861$, which means these prediction models can explain 86% of the variation of PKI responses). Although we designed three types of neural network architectures in this study: single dense layer (SDL), simple double dense layers (SDDL), and complex double dense layers (CDDL) (see Materials and methods), we found that the prediction models for all the 23 cancer types can achieve $R^2 > 0.8$ by using either SDL or SDDL models. Based on Occam's razor principle [38], we chose the architecture as simple as possible and thus we did not implement CDDL models.

Residual analysis was then performed to assess the appropriateness of our trained prediction models. The residual plot (Fig 2b) shows that there is no specific U shape, inverted U shape, or funnel shape, which means these prediction models need no more higher-order features to capture the variation of drug responses (S1 Fig shows residual plots for 23 cancer types). To further confirm the prediction model's ability to classify drug responses into two categories (sensitive vs. non-sensitive), we chose thresholds to define actual IC_{50} as sensitive or non-sensitive. Comparing to the single threshold used in a previous study [33] ($IC_{50} = -2$), we set multiple thresholds (-4, -3, -2, -1, and 0) and averaged the results to avoid overestimating the prediction performance. The result ROC curves of 23 cancer types and the overall curve are shown in Fig 2c. The overall AUC is 0.981, similar to the performance in the previous study [33] (AUC > 0.98). AUC for each cancer type is available in S1 Table.

For more information about the prediction performance for different PKI target groups, see Supporting information.

The contribution of different feature groups

In the QSMART with neural network models, to approximately estimate the contribution of different feature groups, we split the features into drug features, cancer cell line features, and interaction features, used the same neural network architecture (parameters and the number of nodes in the first and second hidden layers) of each cancer type, and then evaluated the prediction performances of using different feature sets. As a result, Fig 2d shows the approximate contribution of each feature category to prediction performance (the detailed number of features and performances are shown in Table 3). Across different cancer types, the result showed that the contribution from drug features (overall $R^2 = 0.661$) outperformed those from cancer cell line features and interaction features (overall $R^2 = 0.126$ and 0.152, respectively), and the contribution from interaction features was higher than that from cancer cell line features (p-value =

Table 2. Prediction performances of using QSMART model with different machine learning methods.

Cancer type	#IC$_{50}$	#All Features	#Drug Features	#Cancer features Residue	Others	#Interactions DxM	Others	#Nodes 1st	2nd	NN	RF	SVM	Lasso	ANOVA	MCA
AG	2,971	62	38	0	5	9	10	8	38	**0.815**	0.362	0.243	0.293	0.672	0.656
Bone	3,410	84	52	0	13	4	15	10	0	**0.856**	0.483	0.316	0.370	0.693	0.819
Breast	4,706	129	70	5	26	12	16	6	26	**0.880**	0.527	0.452	0.496	0.702	0.814
CNS	4,250	114	65	0	23	11	15	11	0	**0.858**	0.548	0.399	0.465	0.774	0.851
Cervix	1,044	37	29	0	3	1	4	7	0	**0.864**	0.552	0.389	0.455	0.809	0.824
Endometrium	1,073	33	21	0	4	4	4	4	11	**0.878**	0.358	0.279	0.328	0.769	0.832
Haematopoietic	4,204	119	58	3	24	28	6	11	0	**0.858**	0.518	0.378	0.429	0.679	0.807
Kidney	2,458	73	51	0	3	0	19	9	0	**0.836**	0.537	0.347	0.415	0.794	0.820
Large intestine	4,628	141	53	10	14	50	14	12	0	**0.814**	0.468	0.449	0.495	0.736	0.794
Liver	1,348	48	35	0	4	2	7	7	0	0.836	0.575	0.301	0.377	0.730	**0.859**
Lung (NSCLC)	9,205	207	72	7	35	47	46	15	0	**0.854**	0.466	0.470	0.513	0.728	0.819
Lung (others)	7,206	162	58	2	16	46	40	6	30	**0.859**	0.381	0.428	0.470	0.725	0.791
Lymphoid	13,302	291	72	54	30	86	49	18	0	**0.873**	0.449	0.448	0.495	0.758	0.834
Oesophagus	3,337	91	58	0	17	4	12	10	0	**0.841**	0.509	0.391	0.452	0.771	0.838
Ovary	3,502	113	64	2	18	9	20	11	0	**0.844**	0.532	0.471	0.522	0.741	0.810
Pancreas	2,421	84	60	0	7	0	17	10	0	**0.833**	0.591	0.419	0.492	0.784	0.816
Pleura	1,431	36	23	0	5	0	8	4	11	0.805	0.263	0.243	0.303	0.776	**0.837**
Skin	5,732	132	64	9	21	15	23	12	0	**0.875**	0.602	0.398	0.458	0.754	0.800
Soft tissue	1,938	63	45	0	10	2	6	8	0	**0.818**	0.540	0.333	0.404	0.758	0.786
Stomach	2,327	83	49	0	13	16	5	5	20	0.836	0.490	0.319	0.392	0.720	**0.842**
Thyroid	1,352	33	25	0	5	0	3	6	0	0.830	0.538	0.359	0.398	0.798	**0.853**
UAT	3,856	126	74	1	13	13	25	12	0	**0.869**	0.653	0.545	0.600	0.792	0.841
Urinary tract	1,454	68	47	0	5	9	7	9	0	**0.863**	0.558	0.344	0.433	0.754	0.847
Overall	87,155									**0.861**	0.496	0.429	0.460	0.755	0.823

The best performance for each cancer type is highlighted in bold. The performance of each machine learning method, except for ANOVA and MCA [35], is based on 10-fold cross-validation. The performance of MCA is based on its prediction for PKI response. AG: autonomic ganglia; ANOVA: analysis of variance; CNS: central nervous system; DxM: drug-mutation interaction; MCA: multiscale convolutional attentive; NN: neural networks; NSCLC: non-small cell lung cancer; R^2: coefficient of determination; RF: random forests; SVM: support vector machine; UAT: upper aerodigestive tract; #IC$_{50}$: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks.

0.0081, Wilcoxon signed-rank test). Although it was partially due to the number of selected drug features was more than those of the other two feature categories, the main reason was that drug features were more informative. Since the entire training dataset was split into 23 cancer-centric datasets, the similarity among cancer cell lines in one dataset was higher than the similarity among PKIs and thus the drug features had higher variation and higher entropy.

Assuming that the features from different categories were independent and could explain the variation of drug response from different aspects, the summation of the respective R^2 of split models (the R^2_{Split} shown in Table 3) would ideally be the upper limit of a full model. However, Table 3 shows that there were 14 cancer-centric models having prediction performance R^2_{Full} even higher than R^2_{Split}, which implies that the synergistic prediction performance (R^2_{Full} - R^2_{Split}) was potentially from the higher-order interactions performed by neural networks. Interestingly, we found that the neural network architectures of the models with the top four synergistic effects were all double-hidden-layer neural networks, instead of single-hidden-layer neural networks, which also supported our hypothesis that the synergistic prediction performance was from higher-order interactions. On the other hand, the three cancer types (large intestine, cervix, and lymphoid) with the least synergistic effects had the top three $R^2_{Interaction}$. It implied that for these three cancer types, the contribution from the higher-order interactions performed by neural networks was limited because those informative interaction features had been captured by the QSMART model.

Fig 2. The prediction performances of different datasets and different prediction models. Wilcoxon signed-rank test is performed to compare prediction performances and the p-value is shown in each box plot. (a) Comparison between actual IC_{50} (x-axis) and the IC_{50} predicted by using QSMART with neural networks across all cancer types (y-axis); a regression line is shown. (b) Residual analysis for the models using QSMART with neural networks across all cancer types. X-axis: predicted IC_{50}; y-axis: residuals, defined as actual IC_{50} minus predicted IC_{50}. (c) AUC curves of 23 cancer-centric models and an overall AUC. (d) The prediction performances of split QSMART models. (e) The prediction performances of using different datasets (multi-omics, genomics fingerprints, and NoX: no interaction terms) and different feature selection methods (random and Rand10X: randomly select 10 times of the feature number in the QSMART model). (f) The prediction performances of using different statistical or machine learning methods. NN: neural networks; ANOVA: analysis of variance; RF: random forests; SVM: support vector machine.

More informative features for predicting PKI response: multi-omics data

To evaluate the first component of the framework in this study – drug features and cancer cell line's multi-omics data – referring to the features used in a previous study [33], we used PaDEL-descriptor [39] (a software to calculate molecular descriptors and fingerprints) to generate PKI's fingerprints, extended fingerprints, and graph-only fingerprints (3,072 drug features in total) and obtained cancer cell line's genomic fingerprints (mutation genome positions) from COSMIC Cell Lines Project [40] (44,364 cancer cell line features, illustrated in S2 Fig). To make them comparable with our models, we used the same feature selection method to prioritize all the drug fingerprints

Table 3. Approximate contribution of each feature category to prediction performance by using QSMART model with neural networks.

Cancer type	#Nodes		Split QSMART models						Performance comparison		
			Drug		Cancer cell line		Interaction		Full model	Split models	Difference
	1st	2nd	#Features	R^2_{Drug}	#Features	R^2_{Cancer}	#Features	$R^2_{Interaction}$	R^2_{Full}	R^2_{Split}	$R^2_{Full}-R^2_{Split}$
AG	8	38	38	0.611	5	0.044	19	0.041	0.815	0.696	0.119
Stomach	5	20	49	0.611	13	0.053	21	0.062	0.836	0.726	0.110
Breast	6	26	70	0.629	31	0.070	28	0.073	0.880	0.771	0.109
Pleura	4	11	23	0.614	5	0.043	8	0.061	0.805	0.718	0.088
Liver	7	0	35	0.652	4	0.020	9	0.078	0.836	0.751	0.086
Haematopoietic	11	0	58	0.599	27	0.092	34	0.098	0.858	0.789	0.070
Oesophagus	10	0	58	0.699	17	0.027	16	0.050	0.841	0.776	0.066
Soft tissue	8	0	45	0.561	10	0.100	8	0.104	0.818	0.765	0.053
CNS	11	0	29	0.683	3	0.072	5	0.055	0.858	0.810	0.048
Urinary tract	9	0	47	0.673	5	0.105	16	0.048	0.863	0.826	0.037
Lung (NSCLC)	15	0	72	0.610	42	0.084	93	0.128	0.854	0.822	0.031
Skin	12	0	64	0.685	30	0.041	38	0.122	0.875	0.848	0.027
Bone	10	0	52	0.607	13	0.111	19	0.112	0.856	0.830	0.026
Lung (others)	6	30	58	0.610	18	0.121	86	0.104	0.859	0.834	0.024
Pancreas	10	0	60	0.717	7	0.058	17	0.061	0.833	0.835	-0.002
Thyroid	6	0	25	0.713	5	0.067	3	0.053	0.830	0.833	-0.003
UAT	12	0	74	0.732	14	0.061	38	0.080	0.869	0.873	-0.004
Endometrium	4	11	21	0.709	4	0.076	8	0.099	0.878	0.884	-0.006
Ovary	11	0	64	0.648	20	0.092	29	0.122	0.844	0.861	-0.017
Kidney	9	0	51	0.666	3	0.074	19	0.126	0.836	0.866	-0.030
Lymphoid	18	0	72	0.661	84	0.097	135	0.149	0.873	0.907	-0.034
Cervix	7	0	65	0.669	23	0.033	26	0.244	0.864	0.946	-0.081
Large intestine	12	0	53	0.574	24	0.160	64	0.209	0.814	0.943	-0.129
Overall				0.661		0.126		0.152	0.861	0.940	-0.079

R^2_{Full}: the performance of using full QSMART model with neural networks shown in Table 2; R^2_{Split}: the summation of the performances of split models ($R^2_{Split} = R^2_{Drug} + R^2_{Cancer} + R^2_{Interaction}$). AG: autonomic ganglia; CNS: central nervous system; NSCLC: non-small cell lung cancer; UAT: upper aerodigestive tract; #Nodes: number of nodes in the first and second hidden layers of neural networks.

and genomic fingerprints, selected the same total number of features for each cancer type in our model (shown in Table 2), and then used the same neural network architectures. The number of selected features and prediction performances are shown in S2 Table. The box plot in Fig 2e shows that the performance distribution of 23 cancer-centric models using multi-omics data is significantly higher than that of the models using genomic fingerprints (p-value < 2.9e-05, Wilcoxon signed-rank test). Although the performance in the previous study [33] achieved $R^2 = 0.843$ by using genomic fingerprints as features for the neural networks with 17 to 31 hidden layers, this comparison result implies that using these informative multi-omics data is more efficient.

More explainable features for predicting PKI response: interaction effects

To evaluate the second component of the framework in this study – statistical tests for interaction effects – we removed the interaction terms in the models, directly moved forward to the third component (feature selection) to select the same number of features for each cancer type in the original model, and then used the same neural network architectures to train the new models. The number of selected features and prediction performances are shown in S3 Table. The box plot in Fig 2e shows the performance of using full QSMART models is significantly higher than that of the models without interaction effects (p-value = 0.033, Wilcoxon signed-rank test). Comparing to the overall performance of full QSMART models, using the models without interaction effects decreased the overall performance to $R^2 = 0.823$. Interestingly, compared to the full QSMART models, we found that the prediction models of some cancer types, such

as upper aerodigestive tract and breast, achieved higher performance without using interaction effects. We conjectured that some informative high-order interactions were captured inside the neural network black box and compensated the lack of interaction effects in the input layer. However, using neural networks cannot guarantee that these informative but unexplainable high-order interactions will be captured under the limited number of samples and the training iteration we used. This fact is reflected in Fig 2e, which shows the prediction performances of using no interaction effects are not stable (R^2 = 0.581 to 0.912).

More efficient feature selection method: Lasso with BIC control

To evaluate the third component of the framework in this study – a feature selection method – after the first two components, we randomly selected the same number of features in the original models and then used the same neural network architectures to make the performances comparable. For each cancer type, the number of randomly selected features along with prediction performances are shown in S4 Table. It was not surprising that the prediction performances dropped to R^2 = 0.031 to 0.138 (overall R^2 = 0.125). To further evaluate the feature selection method we used, we increased the number of randomly selected features to 10 times the original number. The performances increased to R^2 = 0.052 to 0.707 (S5 Table; overall R^2 = 0.378). As the number of selected features increased to 10 times, we saw the performances were increased. If a prediction process has no feature selection at all, we would definitely expect that the prediction performance is better than that of a reduced model; however, regardless of the degree of freedom and overfitting issues, the huge number of chemical and biological properties, including considerable redundant and trivial information, will reduce training efficiency, and they require more complex models, deeper neural networks, or more training iterations to achieve high accuracy. Therefore, we performed these two random selection experiments to validate that Lasso with BIC control efficiently provided highly informative feature sets.

The best performing machine learning method for the QSMART model: neural networks

To evaluate the last component of the framework in this study – a machine learning method – we chose random forests, SVM, and Lasso regression to compare with neural networks for each comparative dataset/feature set mentioned above. Based on the same feature set as inputs, neural networks significantly outperformed other machine learning approaches (Table 2; overall R^2 = 0.496, 0.429, and 0.460 for random forests, SVM, and Lasso regression, respectively). Furthermore, based on the feature sets used to validate the contribution of previous components in the framework, neural networks also outperformed random forests, SVM, and Lasso regression (S2 Table-S5 Table). In a previous study [33], it also showed the phenomenon that neural networks have better drug response prediction performance than random forests and SVM (R^2 = 0.843, 0.698, and 0.562 for DNN, random forests, and SVM, respectively). Interestingly, neural networks were only slightly better than Lasso in overall performance when randomly selected features were used as inputs (R^2 = 0.125 vs. 0.116, p-value = 0.015, Wilcoxon signed-rank test). It further validated the importance of the feature selection method we chose. Overall, neural networks had shown better ability to utilize multi-omics features and their interaction effects.

In addition to machine learning approaches, we compared our models with two-way ANOVA analyses and MCA [35]. Two-way ANOVA analyses were used to assess how much the two factors, drug and cancer cell line, can explain the variation of drug response. Drug IDs and cancer cell line IDs represented different levels of drug and

cancer cell line, respectively. The result of two-way ANOVA showed that these two ²²⁸
factors could explain 67.2% to 80.9% of the drug response variation in different cancer ²²⁹
types (Table 2; overall $R^2 = 0.755$), meaning the datasets we collected and cleaned had ²³⁰
limited noise or other uncertain factors responsible for the variation. Although the ²³¹
result seems decent, using no drug features nor cancer cell line features (only using their ²³²
IDs) loses the predictive power of the drug responses for new drugs or new cancer ²³³
samples which were respectively not included in the drug levels or cancer cell line levels ²³⁴
used in the ANOVA analyses. Comparing to ANOVA and MCA, using the multi-omics ²³⁵
features from QSMART model with neural networks had significantly higher ability to ²³⁶
explain the PKI response variation in 23 cancer types (p-value $< 2.9e\text{-}05$ and p-value $=$ ²³⁷
0.0011 based on Wilcoxon signed-rank test, respectively; Fig 2f). ²³⁸

Case study: non-small cell lung cancer ²³⁹

Above, we have validated the contribution of multi-omics data and interaction effects in ²⁴⁰
the models by comparing the prediction performances. Now, we will discuss how these ²⁴¹
features and models are explainable. We chose one of the largest datasets, non-small cell ²⁴²
lung cancer (NSCLC), as a case study to exemplify how the selected features explain ²⁴³
drug response and the potential mechanism of drug resistance. All 207 features selected ²⁴⁴
by NSCLC's QSMART model and their descriptions were listed in S2 Data. We chose ²⁴⁵
several pertinent features and explain their biological relevance in this case study to ²⁴⁶
demonstrate how scientists may use our prediction model and explain their findings. ²⁴⁷

Drug feature ²⁴⁸

"From_Sanger". This feature was introduced into the model to distinguish the assays ²⁴⁹
done by Massachusetts General Hospital (0) or Wellcome Sanger Institute (1). This ²⁵⁰
feature represents the batch effects among the laboratory experiments performed by ²⁵¹
these two institutes. On average, the PKI responses obtained from Massachusetts ²⁵²
General Hospital showed lower drug sensitivity (higher IC_{50} value) than those from the ²⁵³
Wellcome Sanger Institute in the NSCLC dataset (average actual $IC_{50} = 2.88$ vs. 2.41). ²⁵⁴
To investigate these experimental batch effects, we increased one unit to this feature and ²⁵⁵
held other features constant. Although holding other features constant is not possible in ²⁵⁶
reality, from the mathematical point of view, the result showed that if we replace 0 with ²⁵⁷
1 for From_Sanger, the average IC_{50} predicted by our pre-trained model will reduce 0.65 ²⁵⁸
(S2 Data; average predicted $IC_{50} = 2.87$ vs. 2.22). Interestingly, this feature was ²⁵⁹
selected not only in the NSCLC model but also in other 22 cancer-centric models, ²⁶⁰
meaning the batch effects were significant across the assays done by these two institutes. ²⁶¹

Biological processes interaction ²⁶²

"GO_0030324_X_GO_0048675". This feature represents the multiplication of the number ²⁶³
of mutations that occurred in the proteins associated with the biological process "lung ²⁶⁴
development" (Gene Ontology ID: GO:0030324) and the number of mutations that ²⁶⁵
occurred in the proteins associated with "axon extension" (Gene Ontology ID: ²⁶⁶
GO:0048675). Axon initiation, extension, and guidance are known to play some roles in ²⁶⁷
cancer invasion and metastasis [41]. In the NSCLC dataset, there were eight cell lines ²⁶⁸
with mutations in protein kinases associated with axon extension: CAL-12T, EKVX, ²⁶⁹
LCLC-97TM1, SK-LU-1, NCI-H1793, NCI-H1944, NCI-H2030, and NCI-H2087; the last ²⁷⁰
two were from patients with metastatic NSCLC. On average, the NSCLC cell lines with ²⁷¹
this interaction showed higher PKI responses than those without this interaction ²⁷²
(average actual $IC_{50} = 4.32$ vs. 2.69) and those involved in "lung development" or ²⁷³
"axon extension" alone (average actual $IC_{50} = 3.20$ or 2.07, respectively). Based on our ²⁷⁴

prediction model, every one unit increase in this interaction term was associated with a ^{275} 0.45 unit increase in IC_{50} on average (average predicted IC_{50} = 2.73 vs. 3.18). ^{276}

Protein-protein interaction ^{277}

Instead of explaining a single protein-protein interaction (PPI), in this paragraph, we ^{278} will represent a PPI network consisting of the PPIs selected as features in the PKI ^{279} response prediction model for NSCLC and other interactions among the proteins ^{280} involved in those PPIs. In the 207 features selected by NSCLC's QSMART model, there ^{281} were 27 PPIs weighted by gene expression level. Every one unit of gene expression level ^{282} increase in these PPIs was associated with -0.089 to 0.061 unit increase in IC_{50} on ^{283} average (Fig 3). Taking the 27 genes in this subnetwork to perform a gene list analysis ^{284} by using PANTHER [42], we found that they were significantly (FDR < 0.05) ^{285} over-represented in 11 PANTHER pathways, including angiogenesis, inflammation, ^{286} apoptosis, and axon guidance (S6 Table). MAP4K4, one of the genes involved in the ^{287} apoptosis signaling pathway, is an emerging therapeutic target in cancer [43], and its ^{288} over-expression is a prognostic factor for lung adenocarcinoma, one of the major ^{289} subtypes of NSCLC [44]. MAP4K4 expression is up-regulated upon binding by p53, a ^{290} tumor suppressor gene, and it will then activate the JNK signaling pathway to drive ^{291} apoptosis [45]. In the NSCLC dataset, when the expression of MAP4K4-TP53 ^{292} interaction increase, average IC_{50} is slightly decreased (Pearson correlation = -0.10); in ^{293} our PKI response prediction model, every one unit of gene expression level increase in ^{294} MAP4K4-TP53 PPI was associated with 0.012 unit decrease in IC_{50} on average ^{295} (average predicted IC_{50} = 2.727 vs. 2.715). ^{296}

Although CDK13, classified as an understudied protein kinase by NIH Illuminating ^{297} the Druggable Genome (IDG) program [46] (S3 Data, last updated on June 11, 2019), is ^{298} not involved in the enriched pathways shown in S6 Table, it participates in the pathway ^{299} "TP53 Regulates Transcription of DNA Repair Genes" (Reactome ID: R-HSA-6796648) ^{300} and a 4-clique PPI module in the TP53-centric subnetwork (Fig 3). Its three PPIs in ^{301} this module were all selected as features in the PKI response prediction model. One of ^{302} CDK13's PPI partners, AKAP4, is a biomarker for NSCLC [47], and its expression ^{303} increase was associated with tumor stage. In addition to NSCLC, AKAP4 is also a ^{304} potential therapeutic target of colorectal cancer [48] and ovarian cancer [49], and it ^{305} regulates the expression of CDK family, which plays an important role in cellular ^{306} proliferation [50]. In the NSCLC dataset, the expression of CDK13-AKAP4 interaction ^{307} had a weak positive correlation with IC_{50} (Pearson correlation = 0.07); in the ^{308} prediction model, every one unit of gene expression level increase in CDK13-AKAP4 ^{309} PPI was associated with 0.017 unit increase in IC_{50} on average (average predicted IC_{50} ^{310} = 2.727 vs. 2.744). ^{311}

Drug-mutation interaction ^{312}

In this paragraph, we will illustrate drug-mutation interaction hot spots on a reference ^{313} protein kinase A (PKA) structure (PDB ID: 1ATP, chain E). In total, there were 47 ^{314} drug-mutation interactions in the NSCLC's QSMART model, and they were located in ^{315} 22 PKA positions represented by spheres in Fig 4a. Note that these interactions were ^{316} statistical terms that might not be directly interpreted as physical interactions. The ^{317} drug-mutation interactions located in canonical ATP-binding pocket (highlighted by a ^{318} dashed rectangle in Fig 4a), such as PKA_123 (at the hinge region) and PKA_187 (right ^{319} next to the DFG motif), could be formed by type I or type II protein kinase inhibitors ^{320} according to the protein structure's active or inactive conformation, respectively [51]. ^{321} The interactions adjacent to the ATP-binding pocket, such as PKA_73 (right next to ^{322} the lysine of the K-E salt bridge) and PKA_197 (at the activation loop), could be ^{323}

Fig 3. A protein-protein interaction network constructed by the interaction features for predicting PKI response in NSCLC cell lines. Green node: protein kinase; dark green node: dark/understudied protein kinase; yellow node: other protein; the node with a thick border: known PKI target; red edge: the PPI with positive impacts on IC_{50}; light red edge: the PPI with weak positive impacts on IC_{50}; blue edge: the PPI with negative impacts on IC_{50}; light blue edge: the PPI with weak negative impacts on IC_{50}; gray edge: the PPI not in the prediction model.

formed by type III inhibitors that bind to an allosteric pocket near the ATP-binding   324

pocket [51]. The rest interactions could be formed by type IV inhibitors that bind to an allosteric pocket remote from the ATP-binding pocket [52]. Taking PKA_187 for example, we further investigated how the interactions contribute to drug responses. In our NSCLC dataset, there were four cell lines, NCI-H2087, H3255, NCI-H1975, and NCI-H345, having mutations located in this position: BRAF L597V, EGFR L858R, EGFR L858R, and STK32C I237V, respectively.

Fig 4. Drug-mutation interaction hot spots on the reference protein kinase A structure and examples of the interactions located in ATP-binding pocket. (a) Interaction hot spots are labeled and represented by larger spheres on the reference PKA structure (PDB ID: 1ATP). Medan impact on IC$_{50}$ was chosen to represent a residue involved in multiple drug-mutation interactions. Red sphere: the drug-mutation interaction with positive impacts on IC$_{50}$; light red sphere: the interaction with weak positive impacts on IC$_{50}$; blue sphere: the interaction with negative impacts on IC$_{50}$; light blue sphere: the interaction with weak negative impacts on IC$_{50}$. (b) and (c): Examples of two PKIs (afatinib and lapatinib) with different binding modes in the active (PDB ID: 4G5J) and inactive (PDB ID: 1XKK) conformations of EGFR crystal structures, respectively. The residue corresponding to PKA_187 – EGFR L858 – is labeled in each example; its arginine mutant form simulated by PyMol is illustrated. (d) and (e): Statistical interaction analyses for Fingerprint_791 vs. PKA_187_CHA and Fingerprint_826 vs. PKA_187_VOL in the NSCLC dataset, respectively.

Fig 4b and Fig 4c respectively illustrate different binding modes of two EGFR inhibitors in our dataset, afatinib, and lapatinib, which brought diverse drug responses to the EGFR L858R mutation. Compared to erlotinib and gefitinib (first-generation EGFR inhibitors), afatinib (a second-generation EGFR inhibitor) was associated with longer progression-free survival for the patients with EGFR L858R mutation [53]. Molecular dynamics simulations illustrated that the hydrophobic leucine replaced with a large, positively charged arginine at this position helps to bring additional electrostatic interactions with negatively charged residues at the αC-helix and stabilize the active conformation [54]. Moreover, the EGFR L858R mutation in the active conformation compacted the ATP-binding pocket, increased inter-atomic contacts between afatinib and αC-helix, and thus improved afatinib's binding affinity [54].

In our NSCLC dataset, the drug response of treating H3255 with afatinib, having the drug features involved in the drug-mutation interactions at PKA_187, was one of the lowest (IC$_{50}$ = -4.35) across all the NSCLC cell lines treated with afatinib (average IC$_{50}$ = 2.03, standard deviation = 2.10). Comparing to the afatinib showing no direct interaction with active EGFR L858 in Fig 4b, lapatinib has hydrophobic interaction with the L858 residue in EGFR inactive conformation (Fig 4c). Ones this residue is

substituted with a large, positively charged arginine, the original hydrophobic ⁠349 interaction will be lost and cause several Van der Waals clashes in the binding pocket ⁠ (based on the mutagenesis simulation performed by PyMol [55]), and thus the L858R ⁠ mutation cannot be accommodated in the EGFR inactive conformation with ⁠ lapatinib [56]. In the NSCLC dataset, although the drug response of treating H3255 ⁠ with lapatinib was relatively high ($IC_{50} = 4.88$), the responses across all the NSCLC ⁠ cell lines treated with lapatinib were also high (average $IC_{50} = 4.20$, standard deviation ⁠ $= 1.46$).

Interaction analyses of two drug-mutation interactions, ⁠ "PKA_187_CHA_X_Fingerprint_791" and "PKA_187_VOL_X_Fingerprint_826", located in ⁠ PKA_187 are shown in Fig 4d and Fig 4e, respectively. ⁠ PKA_187_CHA_X_Fingerprint_791 represents the interaction between Fingerprint_791 ⁠ (the drug substructure "NC1CCC(N)CC1") and the charge difference caused by the ⁠ mutation aligned to PKA_187, while PKA_187_VOL_X_Fingerprint_826 means the ⁠ interaction between Fingerprint_826 (the drug substructure "OC1C(N)CCCC1") and ⁠ the side chain volume change caused by the mutation aligned to PKA_187. By ⁠ comparing the average IC_{50}, we see Fig 4d presents significant interactions between ⁠ PKA_187_CHA and Fingerprint_791 (p-value = 0.043, F-test) and Fig 4e shows ⁠ significant interactions between PKA_187_VOL and Fingerprint_826 (p-value = 0.035, ⁠ F-test). Comparing to the blue line in Fig 4d or Fig 4e (the group that lapatinib ⁠ belongs to), the orange line (the group that afatinib belongs to) shows a significant drop ⁠ in average IC_{50} value when both factors are positive. Based on our prediction model, ⁠ every one unit increase in PKA_187_CHA_X_Fingerprint_791 was associated with a 0.46 ⁠ unit decrease in IC_{50} on average (average predicted $IC_{50} = 2.73$ vs. 2.27), while every ⁠ one unit increase in PKA_187_VOL_X_Fingerprint_826 was associated with a 0.01 unit ⁠ decrease in IC_{50} on average (average predicted $IC_{50} = 2.73$ vs. 2.72).

For more information about the biological relevance and mathematical interpretation ⁠ of the features in the NSCLC case study, see Supporting information.

Discussion

To facilitate the understanding of drug response in cancer cell lines from microscopic to ⁠ macroscopic levels, we proposed a PKI response prediction framework to precisely ⁠ estimate IC_{50} values with a more explainable AI model. This framework includes four ⁠ components: (1) drug features, cancer cell line's multi-omics data, and PKI responses, ⁠ (2) statistical tests for interaction effects, (3) feature selection, and (4) neural networks. ⁠ In this study, we validated the contribution of each component, showed high prediction ⁠ performances, and used NSCLC dataset as a case study to explain several features. We ⁠ systematically investigate the previously unknown contributions of various interaction ⁠ effects (such as protein-protein, pathway-pathway, and drug-mutation interactions) on ⁠ drug response.

The intrinsic limitation of any study about drug response prediction should be ⁠ disclosed: the unexplainable variation of drug response caused by different experimental ⁠ environments, assays, and human error. Currently, GDSC and CCLE are the two main ⁠ sources for studying cancer drug response. Several previous studies about predicting ⁠ drug response used data not only from GDSC but also from CCLE (Table 1). However, ⁠ a previous study [21] pointed out that although GDSC and CCLE datasets shared 343 ⁠ cancer cell lines and 15 drugs, the drug responses from these two datasets were poorly ⁠ correlated. Thus, we chose to only use a single source in this study to minimize the ⁠ unexplainable effect from different experimental environments. Nevertheless, this ⁠ situation impeded us from finding appropriate independent testing set outside the ⁠ GDSC data. Even the drug response data we used were only from GDSC, our feature ⁠

selection process showed that the drug feature "From_Sanger" was selected for all the 23 cancer-centric prediction models, meaning the batch effects were significant across the assays done by Wellcome Sanger Institute and Massachusetts General Hospital. Recently, we noticed that GDSC 8.0 was released. Compared with release 7.0, it contains 160 thousand more drug responses. However, this dramatic increase did not provide us a syncretic testing set since the old drug response dataset (called GDSC1 in release 8.0) and the new drug response dataset (called GDSC2) were generated based on different types of assays. Although the drug responses measured by different assays seemed to have high correlation (R = 0.838 in Pearson correlation coefficient), unfortunately, it implied that even if we train a perfect model for GDSC1, the performance of predicting the drug responses in GDSC2 as an independent testing set would only be $R^2 = 0.838^2 = 0.702$ (S3 Fig panel a). Moreover, if we only focus on PKI responses between the two datasets, the correlation is reduced to 0.774 and $R^2 = 0.599$ (S3 Fig panel b). Furthermore, if we use our pre-trained models to predict the PKI response in GDSC2, the overall performance drops to $R^2 = 0.556$ (S3 Fig panel c).

In the case study section, we illustrated the possibility of interpreting statistical interaction terms into potential physical interactions. When we investigated the contribution of protein-protein interactions to drug response prediction, the original purpose of utilizing biological knowledge (known PPIs from STRING [57]) was to narrow down the huge search space (a matrix of 30,000 proteins by 30,000 proteins). Consequently, this additional information also enabled us to explain the biological role of these statistical interaction terms much easier. On the contrary, when we investigated the contribution of drug-mutation interactions to drug response prediction, we explored the entire interactions between all the non-redundant drug features and the mutations at all reference positions. Although limiting the mutations to be in the region around ATP-binding pocket (from PKA_47 to PKA_188, defined by the Kinase-Ligand Interaction Fingerprints and Structures (KLIFS) database [58]) would increase the probability of finding physical interactions among those statistical interaction terms, we would lose the opportunity to explore potential allosteric binding sites and their interactions with PKIs.

In conclusion, by integrating multi-omics data, utilizing the innovative QSMART model, and employing neural networks, we not only can accurately predict PKI responses in cancer cell lines but also increase the explainability behind our prediction models. Comparing to traditional QSAR models, the QSMART model proposed in this study further introduces different types of interaction effects. These interaction effects are universal. While we demonstrate our model in protein kinase binding, the QSMART model can be applied to other protein families, such as G protein-coupled receptors (GPCRs) and ion channels. Moreover, the concept of QSMART model can also be broadly applied to other types of interactions, such as the protein-protein interaction that we had demonstrated, drug-drug interaction, glycosyltransferase-donor analog interaction, gene-environment interaction, and agent-host interaction.

Materials and methods

Protein kinase inhibitor

We define small-molecule (molecular weight < 900 daltons) protein kinase inhibitors (PKIs) in GDSC (release 7.0) [8] from a variety of publicly available, manually curated drug target databases, and experimental data. The list of human protein kinases in this study is defined by ProKinO (version 2.0) [59]. Drug-kinase associations were extracted from DrugBank (version 5.1.0) [60], Therapeutic Target Database (TTD, last accessed on September 15th, 2017) [61], Pharos (last accessed on May 15th, 2018) [62], and

LINCS Data Portal (last accessed on May 15th, 2018) [63]. We define a drug as a PKI ⁠447
if it is annotated as an "inhibitor", "antagonist", or "suppressor" in the drug-kinase ⁠448
associations. We also include the PKIs in LINCS Data Portal if their controls are less ⁠449
than 5% in KINOMEscan$^{\textregistered}$ assays. Based on these criteria, we define 143 ⁠450
small-molecule PKIs out of the 252 unique screened compounds in GDSC (S4 Data). ⁠451

Drug response ⁠452

GDSC provides the half-maximal inhibitory concentration values (IC_{50}, on a ⁠453
logarithmic scale) for 224,202 drug-cancer cell line pairs of drug sensitivity assays. ⁠454
These assays were performed by either the Wellcome Trust Sanger Institute or ⁠455
Massachusetts General Hospital Cancer Center. In this drug response dataset, there are ⁠456
12,509 duplicated drug-cancer cell line pairs derived from 16 duplicated drugs. We ⁠457
measured the Pearson correlation coefficient between the IC_{50} values of each duplicated ⁠458
drug. Only afatinib and refametinib showed a strong positive correlation ($r > 0.7$); their ⁠459
IC_{50} values were merged by their respective weighted means [64]. Drug responses of all ⁠460
other duplicated drugs were excluded from our study as they may have been assayed ⁠461
under different experimental conditions. The resulting dataset of 197,459 non-redundant ⁠462
drug responses consists of 236 drugs and 1,065 cancer cell lines. After filtering out ⁠463
non-PKIs, 109,856 non-redundant drug responses consisting of 135 PKIs and 1,064 ⁠464
cancer cell lines remained. ⁠465

Drug features ⁠466

Drug structures were obtained from PubChem in SDF format. The CDK Descriptor ⁠467
Calculator GUI (version 1.4.6) [65] generated 881 PubChem fingerprints and 286 ⁠468
chemical descriptors including constitutional, topological, electronic, geometric, and ⁠469
bridge descriptors. Observing high multicollinearity within features, we removed ⁠470
redundant features and implemented the variance inflation factor (VIF) criterion [66] to ⁠471
reduce multicollinearity (for more details, see the Feature screening section below). ⁠472
After filtering, 92 PubChem fingerprints and 0 chemical descriptors remained. ⁠473
 To compare our prediction performances with those in a previous study [33], we used ⁠474
the same methods to generate (1) fingerprints, (2) extended fingerprints, and (3) ⁠475
graph-only fingerprints by PaDEL-descriptor (version 2.21) [39] for each drug. In total, ⁠476
there are 3,072 binary descriptors as drug features in comparison models. The ⁠477
comparison models used all features without filtering, as described in the previous ⁠478
study [33]. The relatively large, unfiltered set of drug features are only used for ⁠479
comparison purposes in our study. ⁠480

Cancer cell line features ⁠481

Using mutation profiles for each cancer cell line sample provided by COSMIC Cell Lines ⁠482
Project (v87) [40], we incorporate 7 categories of multi-omics data to quantify ⁠483
differences between wild type and mutants: (1) residue-level: reference protein kinase A ⁠484
(PKA) position (from ProKinO), mutant type, charge, polarity, hydrophobicity, ⁠485
accessible surface area, side-chain volume, energy per residue [67], and substitution ⁠486
score (BLOSUM62 [68]); (2) motif-level: sequence and structural motifs of protein ⁠487
kinase (from ProKinO); (3) domain-level: subdomain in protein kinase (from ProKinO) ⁠488
and functional domain (from Pfam v31.0 [69]); (4) gene-level: the number of mutations ⁠489
in genes, gene expression (from GDSC), and copy number variation (from COSMIC); ⁠490
(5) family-level: protein kinase family and group (from ProKinO); (6) pathway-level: ⁠491
reaction, pathway (from Reactome [70], last accessed on May 15th, 2018), and biological ⁠492
process (from AmiGO [71], last accessed on May 15th, 2018); and (7) sample-level: ⁠493

microsatellite instability, average ploidy, age, cancer originated tissue type, and ⁴⁹⁴ histological classification (from COSMIC and Cellosaurus [72]). ⁴⁹⁵

The formula for generating all cancer cell line features is shown in S7 Table. ⁴⁹⁶ Residue-level features of a cancer cell line were extracted from COSMIC mutants ⁴⁹⁷ labeled as "Substitution - Missense". These features were calculated if the mutation ⁴⁹⁸ position could be aligned to the reference PKA position. This choice is based on an ⁴⁹⁹ assumption that, for all protein kinases, mutations at equivalent positions will have ⁵⁰⁰ similar effects on drug response. An example of this is the gatekeeper residue ⁵⁰¹ (PKA_120). We further used two different types of weights, conservation score ⁵⁰² (KinView [73] with Jensen-Shannon divergence calculation [74]) and gene expression, to ⁵⁰³ estimate the different effects of the same mutant type occurred at the same PKA ⁵⁰⁴ position from different protein kinases. ⁵⁰⁵

Based on mutation position, the values of motif-level or domain-level features were ⁵⁰⁶ calculated if it occurs in a specific motif or domain and its mutation description is ⁵⁰⁷ "Substitution - Missense" or in-frame INDELs (insertions and deletions) in COSMIC. All ⁵⁰⁸ mutation types, except for "Substitution - coding silent" and "Unknown", were taken ⁵⁰⁹ into account for calculating the values of gene-level or higher-level features. For missing ⁵¹⁰ data, we assigned "Neutral" for copy number variation and "Unknown" for ⁵¹¹ microsatellite instability and gender. No imputation was implemented for missing age. ⁵¹²

QSMART model ⁵¹³

The Quantitative Structure-Mutation-Activity Relationship Tests (QSMART) model ⁵¹⁴ was developed based on the QSAR model with interaction effects. Because the ⁵¹⁵ residue-level features of a cancer cell line represent the mutation status in the reference ⁵¹⁶ PKA structure and we are interested in their interactions with the substructures of a ⁵¹⁷ drug, we first built a basic model for estimating IC_{50}: ⁵¹⁸

$$IC_{50} = \beta_0 + \sum_{i=1}^{I} \beta_{1i} D_i + \sum_{k=1}^{K} \beta_{2k} M_k + \sum_{i=1}^{I} \sum_{k=1}^{K} \beta_{3ik} D_i M_k + \epsilon, \tag{1}$$

where β_0 is the intercept, β_{1i} and β_{2k} respectively represent the coefficients of the ith ⁵¹⁹ drug feature D_i and the kth residue-level cancer cell line feature M_k, β_{3ik} is the ⁵²⁰ coefficient of the interaction term formed by D_i and M_k, and ϵ is the error term. ⁵²¹

Considering that not only residue-level features but also higher-level features could ⁵²² independently affect drug response, we expanded the model by incorporating all cancer ⁵²³ cell line features: ⁵²⁴

$$IC_{50} = \beta_0 + \sum_{i=1}^{I} \beta_{1i} D_i + \sum_{j=1}^{J} \beta_{2j} C_j + \sum_{i=1}^{I} \sum_{k=1}^{K} \beta_{3ik} D_i M_k + \epsilon, \tag{2}$$

where β_{2j} is the coefficients of the jth all-level cancer cell line feature C_j. Since all-level ⁵²⁵ features include residue-level features,$\{C_1, ..., C_J\}$ is a superset of $\{M_1, ..., M_K\}$. ⁵²⁶ Considering that the interaction terms formed by the substructures of drug and ⁵²⁷ high-level cancer cell line features have no biological relevance, we did not incorporate ⁵²⁸ all the cancer cell line features as part of interaction terms. For example, we did not ⁵²⁹ consider the interaction between a substructure "Fingerprint_1" and a biological process ⁵³⁰ "lung development" because it is unexplainable. ⁵³¹

In addition to using all-level features to describe a cancer cell line, we further ⁵³² introduced more types of interaction effects into the full QSMART model to capture the ⁵³³

environment of a cancer cell line:

$$IC_{50} = \beta_0 + \sum_{i=1}^{I} \beta_{1i} D_i + \sum_{j=1}^{J} \beta_{2j} C_j + \sum_{i=1}^{I} \sum_{k=1}^{K} \beta_{3ik} D_i M_k + \tag{3}$$

$$\sum_{p=1}^{P} \beta_{4p} PPI_p + \sum_{q=1}^{Q} \beta_{5q} RECx_q + \sum_{r=1}^{R} \beta_{6r} PWYx_r + \sum_{s=1}^{S} \beta_{7s} GOx_s + \epsilon, \tag{4}$$

where β_{4p}, β_{5q}, β_{6r}, and β_{7s} are the coefficients of the pth protein-protein interaction PPI_p, the qth reaction-reaction interaction $RECx_q$, the rth pathway-pathway interaction $PWYx_r$, and the sth biological processes interaction GOx_s, respectively. These four types of interaction effects are formed by all pairs of protein, reaction, pathway, and biological process features, respectively. More details about interaction effects are described below.

Interaction effect

Five types of interaction effects were introduced into the QSMART model: drug-mutation interaction, protein-protein interaction, reaction-reaction interaction, pathway-pathway interaction, and biological processes interaction. These interactions were not necessarily physical interactions; instead, they were predictors that show statistically significant contribution to explaining the variation of IC_{50} values. For drug-mutation interaction, only residues mapping to the reference PKA structure were considered for forming interactions with drugs. To reduce the search space, prior biological knowledge was used to filter interactions with less biological relevance. For protein-protein interaction (PPI), we retain PPIs with scores higher than 700 in the STRING database [57]; gene expression level was used as a weight for PPIs to roughly represent the protein abundance in cancer cell lines. For reaction, pathway, and biological processes interactions, we removed the interactions formed by two entities from the same branch of a tree. For instance, the interaction between the biological process "lung cell differentiation" (GO:0060479) and its parent "lung development" (GO:0030324) was removed since it is unexplainable. Each interaction effect was tested individually by F-test using R (version 3.4.4) [75]. Significant interaction effects (FDR < 0.05) with no less than 30 non-zero values were taken for further feature selection.

Datasets

To reduce more potential sources of noise and bias, we further filter cancer cell lines from the PKI response dataset if (1) their mutation profiles were not detected by whole-genome sequences (2) they have less than 30 drug response entries (3) their gene expression is not available, or (4) their mutation site does not map to a residue in the PKA reference alignment. The dataset was then split into 29 groups, stratified by cancer primary site. Groups with less than 1,000 responses (adrenal gland, biliary tract, placenta, prostate, salivary gland, small intestine, testis, and vulva) were excluded due to low statistical power. "Haematopoietic and lymphoid tissue", the largest group, was further divided into two subsets by primary histology: "haematopoietic neoplasm" and "lymphoid neoplasm". For the case study, we collected cancer cell lines for the non-small cell lung cancer (NSCLC) dataset from the lung cancer dataset if their histology subtype was adenocarcinoma, non-small cell carcinoma, squamous cell carcinoma, large cell carcinoma, giant cell carcinoma, or mixed adenosquamous carcinoma. Remaining samples were classified as "lung (others)". We created cancer type-centric training sets by expanding the drug response dataset with drug features, cancer cell lines features, and significant interaction effects. Categorical data in the training sets were coded into

dummy variables. As a result, we prepared 23 cancer type-centric training sets. The numbers of PKI response, PKIs, and cancer cell lines for each cancer type are shown in Table 1.

Feature screening

Observing high multicollinearity within the features in the first component of our prediction framework (Fig 1), we implemented the variance inflation factor (VIF) criterion [66] to remove highly correlated features. For the multiple regression model with f features, X_i $(i = 1, ..., f)$, the VIF for the ith feature can be expressed by:

$$VIF_i = \frac{1}{1 - R_i^2},\qquad(5)$$

where R_i^2 is the correlation coefficient of the regression between X_i and the remaining $f - 1$ features. $VIF_i > 5$ (i.e. $R_i^2 > 0.8$) was considered to be high collinearity [76] and X_i should be excluded from the model. We first prioritized drug features based on these rules: (1) the later PubChem fingerprint bit positions (complex patterns) have higher priorities than the earlier ones (simple elements), and (2) PubChem fingerprints have higher priorities than calculated chemical descriptors because fingerprints directly represent molecular substructures of the drug. Then, starting from higher priority features moving towards lower priority features, we implemented stepwise selection under VIF control.

Co-expressed genes in the same prediction model also exhibited collinearity. To address this issue, we also used the VIF criterion to filter co-expressed genes in each training set. We prioritize genes based on the Pearson correlation coefficient between their expression and IC_{50} values, then implemented stepwise selection under VIF control.

Feature selection

To combat the problem of p (the number of drug features plus cancer cell line features plus interaction effects) >> n (the number of drug responses) in the training sets, we implemented Lasso [77] with Bayesian information criterion (BIC) [78] by the HDeconometrics package in R [79] (the third component of our prediction framework in Fig 1). After feature selection, the remaining number of selected features for each cancer type are shown in Table 1.

Neural network architecture

For each cancer type, all the selected features provided as input nodes of a neural network, implemented by JMP® [80]. There are three types of neural network architectures in this study: single dense layer (SDL), simple double dense layers (SDDL), and complex double dense layers (CDDL). The numbers of hidden layer nodes follow the geometric pyramid rule [81]. Given N input nodes, there are $\lceil N^{1/2} \rceil$ hidden nodes in the SDL architecture; in the SDDL architecture, there are $\lceil N^{2/3} \rceil$ and $\lceil N^{1/3} \rceil$ hidden nodes in the first and second hidden layers, respectively; in the CDDL architecture, there are N and $\lceil N^{1/2} \rceil$ hidden nodes respectively in the first and second hidden layers. The nodes among the two layers are fully connected. Biases are introduced into the input and hidden layers. The activation function of every node in the neural network is a hyperbolic tangent function (TanH). Newton's method [82] is chosen as an optimizer by JMP.

To avoid overfitting, we implement 10-fold cross-validation, early stopping, and Lasso-style penalty function (absolute value penalty, i.e. L1 regularization [83]). Based

on Occam's razor principle [38], we started from an SDL model for each cancer type. If the performance (average R^2 of the intact validation sets across the 10 folds) is less than a threshold 0.8 in 200 iterations, we increased the iteration to 300; if the performance is still less than the threshold, we implemented an SDDL model for 200 iterations and so on until using a CDDL model for 300 iterations. To increase the reproducibility of this study, fixed random seeds were assigned and all the codes for training and prediction models are available at https://github.com/leon1003/QSMART/.

Comparative prediction models

We compared neural networks with three other prediction algorithms with 10-fold cross-validation: random forests [84], support vector machine (SVM) [85], and Lasso [77]. Random forests were implemented by WEKA (version 3.8.3) [86] with default settings ("maxDepth" = 0, "bagSizePercent" = 100). For each cancer type, the number of iterations was decided based on the iterations used for each of the final neural network models (200 or 300 iterations). SVM was implemented by the SMOreg function (SVM for regression) in WEKA with default kernel ("PolyKernel") and optimizer ("RegSMOImproved") settings. Lasso was implemented by the R package "glmnet" [87] with default parameter setting for Lasso regression (alpha = 1 and family = "gaussian"). Additionally, we also compared our prediction models with two-way ANOVA analyses and MCA [35]. Because the purpose of two-way ANOVA analyses implemented by R was to quantify how much two factors (drug and cancer cell line) can explain the variation of drug response (adjusted R^2 was used), the model used the drug and cancer cell line identifiers as inputs and did not undergo 10-fold cross-validation. The performance of MCA shown in Table 2 is based on its prediction for PKI response (details are available in S1 Data).

Supporting information

S1 Fig. Residual analyses for 23 cancer-centric models and the overall result of using QSMART with neural networks. X-axis: predicted IC_{50}; y-axis: residuals, defined as actual IC_{50} minus predicted IC_{50}. Residuals mean and standard deviation are shown for each cancer type.

S2 Fig. Genome-wide mutational status (genomic fingerprints) across all 23 cancer types. AG: autonomic ganglia; CNS: central nervous system; NSCLC: non-small cell lung cancer; UAT: upper aerodigestive tract.

S3 Fig. Comparison between GDSC1 and GDSC2 in the GDSC release 8.0. GDSC1 (the old drug response dataset) and GDSC2 (the new drug response dataset) were generated based on different types of assays. Cell viability was measured using either Resazurin or Syto60 in GDSC1, while it was measured based on Promega CellTiter-Glo® in GDSC2. In total, there are 22,624 drug-cancer cell line pairs found in both datasets; the experiments of all these pairs were done by Wellcome Sanger Institute. (a) The hexbin plot shows the actual IC_{50} from GDSC1 (x-axis) versus the actual IC_{50} from GDSC2 (y-axis); a fitted regression line and its R^2 are shown. (b) There are 7,283 PKI-cancer cell line pairs found in both GDSC1 and GDSC2. The hexbin plot shows the PKI's actual IC_{50} from GDSC1 (x-axis) versus the PKI's actual IC_{50} from GDSC2 (y-axis). (c) Based on the prediction result of our QSMART with neural network models trained by GDSC1 data, the hexbin plot shows the PKI's predicted IC_{50} (x-axis) versus the PKI's actual IC_{50} from GDSC2 (y-axis).

S4 Fig.   The prediction performances of using QSMART model with neural networks for different PKI target groups. (a) Average actual IC_{50} of different PKI target groups across 23 cancer types. (b) The prediction performances (in R^2) of using QSMART model with neural networks for different PKI target groups. (c) The prediction performances (in RMSE: root-mean-square error) of using QSMART model with neural networks for different PKI target groups. NSCLC: non-small cell lung cancer.

S1 Table.   The number of different-level features and prediction performance of neural networks. AG: autonomic ganglia; AUC: area under the ROC Curve; CNS: central nervous system; DxM: drug-mutation interaction; PPI: protein-protein interaction; GOx: biological process interaction; NSCLC: non-small cell lung cancer; PWYx: pathway-pathway interaction; R^2: coefficient of determination; RECx: reaction-reaction interaction; RMSE: root-mean-square error; UAT: upper aerodigestive tract; #IC_{50}: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks; #Tours: number of times to fit the model.

S2 Table.   Prediction performances of using genomic fingerprints. The best performance for each cancer type is highlighted in bold. AG: autonomic ganglia; CNS: central nervous system; NN: neural networks; NSCLC: non-small cell lung cancer; R^2: coefficient of determination; RF: random forests; SVM: support vector machine; UAT: upper aerodigestive tract; #IC_{50}: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks.

S3 Table.   Prediction performances of using no interaction effects. The best performance for each cancer type is highlighted in bold. AG: autonomic ganglia; CNS: central nervous system; NN: neural networks; NSCLC: non-small cell lung cancer; R^2: coefficient of determination; RF: random forests; SVM: support vector machine; UAT: upper aerodigestive tract; #IC_{50}: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks.

S4 Table.   Prediction performances of using random feature selection. The best performance for each cancer type is highlighted in bold. AG: autonomic ganglia; CNS: central nervous system; DxM: drug-mutation interaction; NN: neural networks; NSCLC: non-small cell lung cancer; R^2: coefficient of determination; RF: random forests; SVM: support vector machine; UAT: upper aerodigestive tract; #IC_{50}: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks.

S5 Table.   Prediction performances of using random 10X feature selection. The best performance for each cancer type is highlighted in bold. AG: autonomic ganglia; CNS: central nervous system; DxM: drug-mutation interaction; NN: neural networks; NSCLC: non-small cell lung cancer; R^2: coefficient of determination; RF: random forests; SVM: support vector machine; UAT: upper aerodigestive tract; #IC_{50}: number of drug responses; #Nodes: number of nodes in the first and second hidden layers of neural networks.

S6 Table.   The result of PANTHER pathway enrichment analysis.

S7 Table. Cancer cell line features. M_{ki}: if the residue corresponding to PKA position i of protein kinase k is mutated (1) or not (0); CSV_{ki}: the conservation score of the residue corresponding to PKA position i of protein kinase k; EXP_k: the gene expression level of protein kinase k; M_{kim}: if the residue corresponding to PKA position i of protein kinase k is mutated to the amino acid type m (1) or not (0); C_{ki}, P_{ki}, H_{ki}, A_{ki}, V_{ki}, or E_{ki}: respectively mean the charge, polarity, hydrophobicity, accessible surface area, side-chain volume, or energy differences caused by the mutation of the residue corresponding to PKA position i of protein kinase k; S_{ki}: the BLOSUM62 substitution score of the mutation occurred at the residue corresponding to PKA position i of protein kinase k; N_k: the length of protein kinase k sequence; M_{kn}: if the nth residue of protein kinase k is mutated (1) or not (0); $L_t(k, n)$, $L_T(k, n)$, $L_d(k, n)$, or $L_D(k, n)$: respectively mean if the nth residue of protein kinase k is located in sequence motif t, structural motif T, subdomain d, or functional domain D (1) or not (0); CSV_{kn}: the conservation score of the nth residue of protein kinase k; CNV_k: the copy number variation status of protein kinase k; $F_f(k)$ or $G_g(k)$: respectively mean if protein kinase k belongs to family f or group p (1) or not (0); $R_r(k)$, $W_w(k)$, or $B_b(k)$: respectively mean if protein kinase k is implicated in reaction r, pathway w, or biological process b (1) or not (0); NCI code: National Cancer Institute (NCI) Thesaurus code.

S1 Data. MCA's performance of PKI response prediction.

S2 Data. Features for Lung (NSCLC) dataset.

S3 Data. Understudied proteins.

S4 Data. PKI target groups and PKI structures.

Acknowledgments

Funding for N.K. (R01GM114409 and U01CA239106) from the National Institutes of Health is acknowledged. Funding for P.M. (R01GM122080 and DMS-1903226) from NIH and NSF is acknowledged.

Author Contributions

Conceptualization: Liang-Chin Huang

Data Curation: Liang-Chin Huang

Formal Analysis: Liang-Chin Huang, Ye Wang, Huimin Cheng, Ping Ma

Funding Acquisition: Natarajan Kannan, Ping Ma

Investigation: Liang-Chin Huang, Wayland Yeung, Natarajan Kannan

Methodology: Liang-Chin Huang, Ping Ma, Khaled Rasheed, Natarajan Kannan

Software: Liang-Chin Huang, Ye Wang, Huimin Cheng, Sheng Li, Khaled Rasheed

Visualization: Liang-Chin Huang, Wayland Yeung

Writing – Original Draft Preparation: Liang-Chin Huang   751

Writing – Review & Editing: Wayland Yeung, Ye Wang, Huimin Cheng, Aarya   753
Venkat, Sheng Li, Ping Ma, Khaled Rasheed, Natarajan Kannan   754

References

1. Arslan MA, Kutuk O, Basaga H. Protein kinases as drug targets in cancer. Curr Cancer Drug Targets. 2006;6(7):623–634.

2. Lehne G, Elonen E, Baekelandt M, Skovsgaard T, Peterson C. Challenging drug resistance in cancer therapy–review of the First Nordic Conference on Chemoresistance in Cancer Treatment, October 9th and 10th, 1997. Acta Oncol. 1998;37(5):431–439.

3. Holohan C, Van Schaeybroeck S, Longley DB, Johnston PG. Cancer drug resistance: an evolving paradigm. Nat Rev Cancer. 2013;13(10):714–726.

4. Sharma SV, Bell DW, Settleman J, Haber DA. Epidermal growth factor receptor mutations in lung cancer. Nat Rev Cancer. 2007;7(3):169–181.

5. Bell DW, Gore I, Okimoto RA, Godin-Heymann N, Sordella R, Mulloy R, et al. Inherited susceptibility to lung cancer may be associated with the T790M drug resistance mutation in EGFR. Nat Genet. 2005;37(12):1315–1316.

6. Tracy S, Mukohara T, Hansen M, Meyerson M, Johnson BE, Janne PA. Gefitinib induces apoptosis in the EGFRL858R non-small-cell lung cancer cell line H3255. Cancer Res. 2004;64(20):7241–7244.

7. Pao W, Miller VA, Politi KA, Riely GJ, Somwar R, Zakowski MF, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. PLoS Med. 2005;2(3):e73.

8. Yang W, Soares J, Greninger P, Edelman EJ, Lightfoot H, Forbes S, et al. Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker discovery in cancer cells. Nucleic Acids Res. 2013;41(Database issue):D955–961.

9. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature. 2012;483(7391):603–607.

10. Nguyen L, Dang CC, Ballester PJ. Systematic assessment of multi-gene predictors of pan-cancer cell line sensitivity to drugs exploiting gene expression data. F1000Res. 2016;5.

11. Geeleher P, Cox NJ, Huang RS. Clinical drug response can be predicted using baseline gene expression levels and in vitro drug sensitivity in cell lines. Genome Biol. 2014;15(3):R47.

12. Jang IS, Neto EC, Guinney J, Friend SH, Margolin AA. Systematic assessment of analytical methods for drug sensitivity prediction from cancer cell line data. Pac Symp Biocomput. 2014; p. 63–74.

13. Ammad-Ud-Din M, Khan SA, Wennerberg K, Aittokallio T. Systematic identification of feature combinations for predicting drug response with Bayesian multi-view multi-task linear regression. Bioinformatics. 2017;33(14):i359–i368.

14. Geeleher P, Zhang Z, Wang F, Gruener RF, Nath A, Morrison G, et al. Discovering novel pharmacogenomic biomarkers by imputing drug response in cancer patients from large genomics studies. Genome Res. 2017;27(10):1743–1751.

15. Ding MQ, Chen L, Cooper GF, Young JD, Lu X. Precision Oncology beyond Targeted Therapy: Combining Omics Data with Machine Learning Matches the Majority of Cancer Cells to Effective Therapeutics. Mol Cancer Res. 2018;16(2):269–278.

16. Wang X, Sun Z, Zimmermann MT, Bugrim A, Kocher JP. Predict drug sensitivity of cancer cells with pathway activity inference. BMC Med Genomics. 2019;12(Suppl 1):15.

17. Li Q, Shi R, Liang F. Drug sensitivity prediction with high-dimensional mixture regression. PLoS ONE. 2019;14(2):e0212108.

18. Zhang N, Wang H, Fang Y, Wang J, Zheng X, Liu XS. Predicting Anticancer Drug Responses Using a Dual-Layer Integrated Cell Line-Drug Network Model. PLoS Comput Biol. 2015;11(9):e1004498.

19. Stanfield Z, Co?kun M, Koyuturk M. Drug Response Prediction as a Link Prediction Problem. Sci Rep. 2017;7:40321.

20. Le DH, Pham VH. Drug Response Prediction by Globally Capturing Drug and Cell Line Information in a Heterogeneous Network. J Mol Biol. 2018;430(18 Pt A):2993–3004.

21. Juan-Blanco T, Duran-Frigola M, Aloy P. Rationalizing Drug Response in Cancer Cell Lines. J Mol Biol. 2018;430(18 Pt A):3016–3027.

22. Yang J, Li A, Li Y, Guo X, Wang M. A novel approach for drug response prediction in cancer cell lines via network representation learning. Bioinformatics. 2019;35(9):1527–1535.

23. Liu H, Zhao Y, Zhang L, Chen X. Anti-cancer Drug Response Prediction Using Neighbor-Based Collaborative Filtering with Global Effect Removal. Mol Ther Nucleic Acids. 2018;13:303–311.

24. Wei D, Liu C, Zheng X, Li Y. Comprehensive anticancer drug response prediction based on a simple cell line-drug complex network model. BMC Bioinformatics. 2019;20(1):44.

25. Rahman R, Matlock K, Ghosh S, Pal R. Heterogeneity Aware Random Forest for Drug Sensitivity Prediction. Sci Rep. 2017;7(1):11347.

26. Lind AP, Anderson PC. Predicting drug activity against cancer cells by random forest models based on minimal genomic information and chemical properties. PLoS ONE. 2019;14(7):e0219774.

27. Dong Z, Zhang N, Li C, Wang H, Fang Y, Wang J, et al. Anticancer drug sensitivity prediction in cell lines from baseline gene expression through recursive feature selection. BMC Cancer. 2015;15:489.

28. Gupta S, Chaudhary K, Kumar R, Gautam A, Nanda JS, Dhanda SK, et al. Prioritization of anticancer drugs against a cancer using genomic features of cancer cells: A step towards personalized medicine. Sci Rep. 2016;6:23857.

29. Ammad-Ud-Din M, Khan SA, Malani D, Murumagi A, Kallioniemi O, Aittokallio T, et al. Drug response prediction by inferring pathway-response associations with kernelized Bayesian matrix factorization. Bioinformatics. 2016;32(17):i455–i463.

30. He X, Folkman L, Borgwardt K. Kernelized rank learning for personalized drug recommendation. Bioinformatics. 2018;34(16):2808–2816.

31. Cichonska A, Pahikkala T, Szedmak S, Julkunen H, Airola A, Heinonen M, et al. Learning with multiple pairwise kernels for drug bioactivity prediction. Bioinformatics. 2018;34(13):i509–i518.

32. Menden MP, Iorio F, Garnett M, McDermott U, Benes CH, Ballester PJ, et al. Machine learning prediction of cancer cell sensitivity to drugs based on genomic and chemical properties. PLoS ONE. 2013;8(4):e61318.

33. Chang Y, Park H, Yang HJ, Lee S, Lee KY, Kim TS, et al. Cancer Drug Response Profile scan (CDRscan): A Deep Learning Model That Predicts Drug Effectiveness from Cancer Genomic Signature. Sci Rep. 2018;8(1):8857.

34. Liu P, Li H, Li S, Leung KS. Improving prediction of phenotypic drug response on cancer cell lines using deep convolutional network. BMC Bioinformatics. 2019;20(1):408.

35. Manica M, Oskooei A, Born J, Subramanian V, Saez-Rodriguez J, Rodriguez Martinez M. Toward Explainable Anticancer Compound Sensitivity Prediction via Multimodal Attention-Based Convolutional Encoders. Mol Pharm. 2019;.

36. Chiu YC, Chen HH, Zhang T, Zhang S, Gorthi A, Wang LJ, et al. Predicting drug response of tumors from integrated genomic profiles by deep neural networks. BMC Med Genomics. 2019;12(Suppl 1):18.

37. Gunning D, Aha DW. DARPA's Explainable Artificial Intelligence Program. AI Magazine. 2019;40(2):44–58.

38. Blumer A, Ehrenfeucht A, Haussler D, Warmuth MK. Occam's Razor. Inf Process Lett. 1987;24(6):377–380. doi:10.1016/0020-0190(87)90114-1.

39. Yap CW. PaDEL-descriptor: an open source software to calculate molecular descriptors and fingerprints. J Comput Chem. 2011;32(7):1466–1474.

40. Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, et al. COSMIC: the Catalogue Of Somatic Mutations In Cancer. Nucleic Acids Res. 2019;47(D1):D941–D947.

41. Chedotal A, Kerjan G, Moreau-Fauvarque C. The brain within the tumor: new roles for axon guidance molecules in cancers. Cell Death Differ. 2005;12(8):1044–1056.

42. Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, et al. PANTHER: a library of protein families and subfamilies indexed by function. Genome Res. 2003;13(9):2129–2141.

43. Gao X, Gao C, Liu G, Hu J. MAP4K4: an emerging therapeutic target in cancer. Cell Biosci. 2016;6:56.

44. Qiu MH, Qian YM, Zhao XL, Wang SM, Feng XJ, Chen XF, et al. Expression and prognostic significance of MAP4K4 in lung adenocarcinoma. Pathol Res Pract. 2012;208(9):541–548.

45. Miled C, Pontoglio M, Garbay S, Yaniv M, Weitzman JB. A genomic map of p53 binding sites identifies novel p53 targets involved in an apoptotic network. Cancer Res. 2005;65(12):5096–5104.

46. the Druggable Genome I. Understudied Proteins; 2019. https://commonfund.nih.gov/idg/understudiedproteins.

47. Gumireddy K, Li A, Chang DH, Liu Q, Kossenkov AV, Yan J, et al. AKAP4 is a circulating biomarker for non-small cell lung cancer. Oncotarget. 2015;6(19):17637–17647.

48. Jagadish N, Parashar D, Gupta N, Agarwal S, Purohit S, Kumar V, et al. A-kinase anchor protein 4 (AKAP4) a promising therapeutic target of colorectal cancer. J Exp Clin Cancer Res. 2015;34:142.

49. Kumar V, Jagadish N, Suri A. Role of A-Kinase anchor protein (AKAP4) in growth and survival of ovarian cancer cells. Oncotarget. 2017;8(32):53124–53136.

50. Duronio RJ, Xiong Y. Signaling pathways that control cell proliferation. Cold Spring Harb Perspect Biol. 2013;5(3):a008904.

51. Gavrin LK, Saiah E. Approaches to discover non-ATP site kinase inhibitors. MedChemComm. 2013;4(1):41–51.

52. Cox KJ, Shomin CD, Ghosh I. Tinkering outside the kinase ATP box: allosteric (type IV) and bivalent (type V) inhibitors of protein kinases. Future Med Chem. 2011;3(1):29–43.

53. Kuan FC, Li SH, Wang CL, Lin MH, Tsai YH, Yang CT. Analysis of progression-free survival of first-line tyrosine kinase inhibitors in patients with non-small cell lung cancer harboring leu858Arg or exon 19 deletions. Oncotarget. 2017;8(1):1343–1353.

54. Kannan S, Pradhan MR, Tiwari G, Tan WC, Chowbay B, Tan EH, et al. Hydration effects on the efficacy of the Epidermal growth factor receptor kinase inhibitor afatinib. Sci Rep. 2017;7(1):1540.

55. Schrödinger, LLC. The PyMOL Molecular Graphics System, Version 1.8; 2015.

56. Yun CH, Boggon TJ, Li Y, Woo MS, Greulich H, Meyerson M, et al. Structures of lung cancer-derived EGFR mutants and inhibitor complexes: mechanism of activation and insights into differential inhibitor sensitivity. Cancer Cell. 2007;11(3):217–227.

57. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic Acids Res. 2015;43(Database issue):D447–452.

58. Kooistra AJ, Kanev GK, van Linden OP, Leurs R, de Esch IJ, de Graaf C. KLIFS: a structural kinase-ligand interaction database. Nucleic Acids Res. 2016;44(D1):D365–371.

59. McSkimming DI, Dastgheib S, Talevich E, Narayanan A, Katiyar S, Taylor SS, et al. ProKinO: a unified resource for mining the cancer kinome. Hum Mutat. 2015;36(2):175–186.

60. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major update to the DrugBank database for 2018. Nucleic Acids Res. 2018;46(D1):D1074–D1082.

61. Li YH, Yu CY, Li XX, Zhang P, Tang J, Yang Q, et al. Therapeutic target database update 2018: enriched resource for facilitating bench-to-clinic research of targeted therapeutics. Nucleic Acids Res. 2018;46(D1):D1121–D1127.

62. Nguyen DT, Mathias S, Bologa C, Brunak S, Fernandez N, Gaulton A, et al. Pharos: Collating protein information to shed light on the druggable genome. Nucleic Acids Res. 2017;45(D1):D995–D1002.

63. Koleti A, Terryn R, Stathias V, Chung C, Cooper DJ, Turner JP, et al. Data Portal for the Library of Integrated Network-based Cellular Signatures (LINCS) program: integrated access to diverse large-scale cellular perturbation response data. Nucleic Acids Res. 2018;46(D1):D558–D566.

64. Jones DC, Hallyburton I, Stojanovski L, Read KD, Frearson JA, Fairlamb AH. Identification of a Î⁰-opioid agonist as a potent and selective lead for drug development against human African trypanosomiasis. Biochem Pharmacol. 2010;80(10):1478–1486.

65. Steinbeck C, Hoppe C, Kuhn S, Floris M, Guha R, Willighagen EL. Recent developments of the chemistry development kit (CDK) - an open-source java library for chemo- and bioinformatics. Curr Pharm Des. 2006;12(17):2111–2120.

66. James G, Witten D, Hastie T, Tibshirani R. An introduction to statistical learning. vol. 112. Springer; 2013.

67. Kawashima S, Ogata H, Kanehisa M. AAindex: Amino Acid Index Database. Nucleic Acids Res. 1999;27(1):368–369.

68. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. Proc Natl Acad Sci USA. 1992;89(22):10915–10919.

69. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, et al. The Pfam protein families database in 2019. Nucleic Acids Res. 2019;47(D1):D427–D432.

70. Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The Reactome Pathway Knowledgebase. Nucleic Acids Res. 2018;46(D1):D649–D655.

71. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, et al. AmiGO: online access to ontology and annotation data. Bioinformatics. 2009;25(2):288–289.

72. Bairoch A. The Cellosaurus, a Cell-Line Knowledge Resource. J Biomol Tech. 2018;29(2):25–38.

73. McSkimming DI, Dastgheib S, Baffi TR, Byrne DP, Ferries S, Scott ST, et al. KinView: a visual comparative sequence analysis tool for integrated kinome research. Mol Biosyst. 2016;12(12):3651–3665.

74. Capra JA, Singh M. Predicting functionally important residues from sequence conservation. Bioinformatics. 2007;23(15):1875–1882.

75. Team RC. type [; 2014].

76. Sheather S. A modern approach to regression with R. Springer Science & Business Media; 2009.

77. Tibshirani R. Regression Shrinkage and Selection Via the Lasso. JOURNAL OF THE ROYAL STATISTICAL SOCIETY, SERIES B. 1994;58:267–288.

78. Schwarz G. Estimating the Dimension of a Model. The Annals of Statistics. 1978;6(2):461–464.

79. R GF. HDeconometrics: Implementation of several econometric models in high-dimension; 2016.

80. Sall J, Stephens ML, Lehman A, Loring S. JMP start statistics: a guide to statistics and data analysis using JMP. Sas Institute; 2017.

81. Masters T. Practical Neural Network Recipes in C++. San Diego, CA, USA: Academic Press Professional, Inc.; 1993.

82. Deuflhard P. Newton methods for nonlinear problems: affine invariance and adaptive algorithms. vol. 35. Springer Science & Business Media; 2011.

83. Ng AY. Feature Selection, L1 vs. L2 Regularization, and Rotational Invariance. In: Proceedings of the Twenty-first International Conference on Machine Learning. ICML '04. New York, NY, USA: ACM; 2004. p. 78–. Available from: http://doi.acm.org/10.1145/1015330.1015435.

84. Breiman L. Random Forests. Mach Learn. 2001;45(1):5–32. doi:10.1023/A:1010933404324.

85. Cortes C, Vapnik V. Support-Vector Networks. Mach Learn. 1995;20(3):273–297. doi:10.1023/A:1022627411411.

86. Witten IH, Frank E, Hall MA, Pal CJ. Data Mining, Fourth Edition: Practical Machine Learning Tools and Techniques. 4th ed. San Francisco, CA, USA: Morgan Kaufmann Publishers Inc.; 2016.

87. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. J Stat Softw. 2010;33(1):1–22.