Title: Allopatric speciation is more prevalent than parapatric ecological

divergence in a recent high-Andean diversification (Asteraceae: *Linochilus*)

Short running title: Allopatric speciation in a high-Andean diversification

Authors: Oscar M. Vargas[1,2*], Santiago Madriñán[3], Beryl B. Simpson[2]

Address:

[1] Department of Biological Sciences, California State Polytechnic University,

Humboldt, Arcata, California, USA

[2] Integrative Biology and Billie Turner Plant Resources Center, The University of

Texas at Austin, Austin, Texas, USA

[3] Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, DC,

Colombia

[*] **Corresponding Author:**

Oscar M. Vargas. Department of Biological Sciences, California State

Polytechnic University, Humboldt, Arcata, CA 95521, USA. Tel 512 300-9236; E-mail:

oscarvargash@gmail.com

ABSTRACT

Elucidating how species accumulate in diversity hotspots is an ongoing debate in evolutionary biology. The páramo, in the Northern Andes, has remarkably high indices of plant diversity, endemicity, and diversification rates. A hypothesis for explaining such indices is that allopatric speciation is high in the páramo given its island-like distribution; an alternative hypothesis is that the altitudinal gradient of the Andean topography provides a variety of niches that drive vertical parapatric ecological speciation. A formal test for evaluating the relative roles of allopatric speciation and parapatric ecological divergence is lacking. The main aim of our study is to test which kind of speciation is more common in an endemic páramo genus. We developed a framework incorporating phylogenetics, species' distributions, and a morpho-ecological trait (leaf area) to compare sister species and infer whether allopatric or parapatric ecological divergence caused their speciation. We applied our framework to the species-rich genus *Linochilus* (63 spp.) and found that the majority of recent speciation events in it (12, 80%) have been driven by allopatric speciation, while a smaller fraction (1, 6.6%) is attributed to parapatric ecological divergence; two pairs produced inconclusive results (13.3%). We conclude that páramo autochthonous diversification is primarily driven by allopatric speciation.

INTRODUCTION

Alexander von Humboldt's and Aimé Bonpland's *Tableau Physique des Andes et Pays Voisins* illustrated how plant species are assembled from lowlands to high altitudes in the tropical Andes (von Humboldt and Bonpland 1805). This representation visualized for the first time how diversity changes locally with elevation, setting a seminal starting point for biodiversity studies. Today, 217 years after the first publication of the *Essay on the Geography of Plants* (von Humboldt and Bonpland 1805), scientists have mapped species richness around the globe and have identified numerous hot-spots of biodiversity. Understanding how species accumulation occurs in such hotspots is pivotal for the fields of evolutionary biology and biogeography.

Because biodiversity hotspots often coincide with areas of topographic complexity (Barthlott et al. 1996; Myers et al. 2000; Mutke and Barthlott 2005; Jenkins et al. 2013) geographical isolation and ecological opportunity are typically cited to explain species richness. In mountain systems, valleys and canyons act as landscape barriers for organisms (Janzen 1967; van der Hammen and Cleef 1986; Muños-Ortiz et al. 2015) promoting allopatric speciation (vicariant or peripatric) via geographical isolation, while slopes and ecological gradients can drive parapatric speciation via ecological divergence (Hughes and Atchison 2015; Pyron et al. 2015). Vicariant speciation occurs when a mother species that is distributed broadly is divided into two daughter populations, which speciate via subsequent independent evolution. Glacial and interglacial cycles are often assumed to have caused these vicariant events (van der Hammen and Cleef 1986; Carstens and Knowles 2007, Flantua et al. 2019). Peripatric speciation occurs when a dispersal event from a source lineage to a new, previously uncolonized area takes place (founder speciation event). If the newly colonized area is geographically isolated enough, then lineages become reproductively isolated with one another, resulting in progenitor and derivative

lineage as different species (Coyne and Orr 2004). Parapatric speciation takes place when a

continuous population that is distributed along an ecological gradient (e.g. an altitudinal

gradient) is subdivided in two or more subpopulations that locally adapt to different niches (e.g.

lower vs. higher elevations); the initial subpopulations become independent species by means of

ecological differentiation, non-random mating, and the formation of reproductive barriers

(Givnish 1997; Schluter 2000; Simpson 1953).

Under a scenario of allopatric speciation, recently diverged species should be similar to

each other thanks to phylogenetic conservatism (Fig. 1A). In the context of sky-islands, sister

species that arose because of geographic speciation should occupy similar niches in different

islands, presenting similar ecological characteristics (Pyron et al. 2015). Alternatively, under a

scenario of parapatric ecological speciation, sister species are expected to occupy different

niches within an island, presenting a signature of ecological divergence (Fig. 1D) (Pyron et al.

2015). Therefore, it is possible to test the predictions of allopatric speciation and parapatric

ecological divergence in sky-islands by studying the distribution and ecological traits of sister

taxa.

94
FIGURE 1. Hypothetical scenarios for speciation in sister species. a) Allopatric speciation event in which
geographical isolation resulted in two species living in separate páramo islands (indicated by island shape
and color) and occupying similar niches (as indicated by similar leaf shapes). b) Allopatric speciation

event in which geographical isolation resulted in two species living in separate páramo islands occupying
different niches as indicated by different leaf types. c) A speciation event in which the reason for
divergence is inconclusive, sister species inhabit the same island and have similar leaf shapes. d) A
parapatric ecological speciation event in which sister species evolved different leaf areas in response to
selection to different niches on the same páramo island.

The Páramo, a high-altitude ecosystem found above the timberline in the Northern Andes

(the Andes of Ecuador, Colombia, and Venezuela) (Cuatrecasas 1968; Luebert and Weigend

2014; Weigend 2002), provides an ideal scenario to test how speciation happens in high

elevation and island-like biodiversity hotspots (Fig. 2). With ca. ~3,400 species of vascular

plants (Luteyn 1999), of which 60% to 100% are estimated to be endemic (Luteyn 1992;

Madriñán et al. 2013), and particularly high diversification rates (Madriñán et al. 2013), the

páramo is considered the most species rich ecosystem of the world's tropical montane regions

(Sklenář et al. 2014). Unlike other South American high-altitude areas farther south (i.e. Perú

and Chile), the páramo is characterized by abundant precipitation throughout the year, and is

therefore defined by both high altitude and humidity. It has been estimated that the páramo

originated when the high altitudes (>3,000 m) of the Northern Andes emerged as a result of rapid

uplift 2–4 Ma (van der Hammen and Cleef 1986; Gregory-Wodzicki 2000). The fragmented

coverage of the paramo (Fig. 2) and its altitudinal gradient (~3000–4500 m) are hypothesized to

have acted as drivers of diversification by promoting both allopatric speciation (vicariant or

peripatric) via geographical isolation (van der Hammen and Cleef 1986) and parapatric

speciation via ecological divergence (Hughes and Atchison 2015, Nevado et al. 2018). Despite

efforts documenting Andean diversification (e.g. Nürk et al. 2015; Uribe-Convers and Tank

2015; Lagomarsino et al. 2016; Pérez-Escobar et al. 2017) a formal test to quantify the relative

prevalence of allopatric versus parapatric ecological speciation of taxa in the region is lacking.

To quantify the relative contributions of allopatric and parapatric speciation in the

páramo, we used a comparative framework that combines phylogenetic, geographical, and

ecological information. We applied this approach to *Linochilus*, a genus restricted to the páramo

and the upper boundary of the cloud forest (Blake 1928; Cuatrecasas 1969; Vargas 2011, 2018;

Vargas et al. 2017; Saldivia et al. 2019). *Lichochilus'* phylogeny was recently inferred using

high-throughput sequencing (Vargas et al. 2017), and a taxonomic monograph is almost

complete for the genus by OMV.

The main aim of our study is to ascertain which kind of speciation, allopatric versus

parapatric ecological divergence, is more common in this this plant genus that inhabits an island-

like system and its lower boundary. We specifically aim to 1) quantify the relative contribution

of allopatric and parapatric ecological speciation in the divergence of sister taxa in *Linochilus*,

and 2) to pinpoint the geographical origin of the *Linochilus* genus, since its presence in the

páramo and the upper boundary of the cloud forest suggest and ecological transition from

paramo to the forest (Vargas & Madriñán 2012).

136
137   **FIGURE 2** Approximate geographical extent of the Páramo ecosystem (left) on the Northern Andes.
138   Páramo la Rusia, Boyacá, Colombia (right); notice how high the Andean forest interdigitates with the
139   Páramo along the creek.

MATERIALS AND METHODS

R Code, input and control files, and bioregionalization maps are available at

https://bitbucket.org/XXXXXXXX/linochilus

FOCAL CLADE

We used *Linochilus* (Asteraceae) because of our recent comprehensive knowledge about

its phylogeny (Vargas et al. 2017), distribution, and taxonomy of its species (Cuatrecasas 1969;

Vargas 2011; Vargas 2018). Additionally, it is almost entirely restricted to the páramo, although

some species dwell in the upper boundary of the cloud forest due to a downslope colonization

event (Vargas and Madriñán 2012). The genus contains 63 species distributed in the disjunct

mountains of the Talamanca Cordillera (Costa Rica), the Sierra Nevada de Santa Marta

(Colombia), and the Northern Andes (Colombia, Venezuela, and Ecuador), exhibiting a variety

of woody habits from decumbent subshrubs only 10 cm tall to small trees 6 m tall (Cuatrecasas

1969; Vargas 2018). Growth form and leaf area of *Linochilus* species are associated with the

habitat they occupy—shrubs and decumbent subshrubs with microphyllous leaves inhabit the

open páramo, while small trees with broad leaves reside at lower altitudes in the upper, more

humid, edge of Andean forest (Vargas and Madriñán 2012). *Linochilus* was recently segregated

from *Diplostephium*, a genus with similar morphology and ecology that primarily inhabits the

Central Andes (Vargas 2018, Saldivia et al. 2019).

SISTER SPECIES COMPARISONS

To measure the relative contribution of allopatric speciation and parapatric ecological

divergence on recent speciation events, we compared the geographical distribution (Appendix

S1) and the leaf areas (Appendix S2) of species shown to be sister in the phylogeny of *Linochilus*

(Vargas et al. 2017). We used leaf area as a proxy to evaluate ecological divergence between

sister species. Leaf area is a functional character that varies with the eco-physiological pressures

of a species' niche, reflecting its adaptation to water availably, irradiance, and elevation (Givnish

1987), therefore providing a proxy for ecological niche. For example, *L. antioquensis* inhabits

the upper Andean forest and has an average leaf area of 848.4 mm^2. In contrast, *L. phylicoides*,

dwells in the physiologically dry open páramo with plenty of access to sunlight, having an

average leaf area of 4.0 mm^2. When possible, we measured the area of 30 leaves from six

different individuals for each species (Appendix S3). We scanned the leaves at 600 dpi from

herbarium material belonging to ANDES, TEX, and US herbaria. Each leaf was outlined using

PHOTOSHOP CS4 (Adobe Systems, San Jose, California) as a single white and black image.

We then used the R package MOMOCS (Bonhomme et al. 2014) to calculate the area of each

leaf from the images created in the previous step. We performed a Wilcoxon signed-rank tests of

the log-transformed leaf areas between sister species using R (R Core Team 2016). We also used

the Wilcoxon signed-rank test to compare the leaf area between clades. It is important to note

that this test considers the distribution of values for leaf areas in each species (instead of just the

average) and it is designed to compare non-independent samples.

We scored a pair as allopatric when they inhabited non-overlapping páramo islands

(mountaintops) and when they inhabited different slopes of the same mountain (this pattern is

possible for species that inhabit the upper boundaries of the Andean forest). We scored a pair as

sympatric when their distribution overlapped at least one páramo island— in other words, the

presence of the two sisters in a single páramo island will make them sympatric. We followed the

paramo delineation of Londoño et al. 2014. In addition to using paramo islands for scoring

species as allopatric or sympatric, we calculated range overlap and range asymmetry between

sisters by overlaying occurrences in a grid following Vargas et al. (2020). We used two grid

sizes, 0.05 and 0.1 decimal degrees, corresponding to 33 and 131 Km2 respectively. With the

grid-ranges of sister species, we calculated the sisters' overlap as the area occupied by both

sisters divided by the summed area of the smaller ranged sister; an overlap of 0 indicates full

allopatry while an overlap of 1 indicates that the smaller-ranged sister is found solely within the

range of its sister (Barraclough and Vogler 2000; Fitzpatrick and Turelli 2006). Range

asymmetry was calculated as the area occupied by the larger-ranged sister divided by the area of

its sister (Fitzpatrick and Turelli 2006). The aforementioned calculations were made on R using

the "raster" package v. 3.5-21 (Hijmans 2022). Distributional data were extracted from curated

COL and US herbaria specimens.

We interpreted the results in the following fashion (Fig. 1):

•   If a sister-species pair is allopatric and there is no significant difference between their

leaf areas, we interpreted this scenario as an event of allopatric speciation driven by

geographical isolation, with the reasoning that leaf area had remained similar because

either relatively little time had passed since divergence and/or because of niche

conservatism. (Fig. 1a; Wiens 2004; Pyron et al. 2015).

•   If a sister-species pair is allopatric and their leaf areas are significantly different, we

interpreted this scenario as an event of allopatric speciation following geographical

isolation (Fig. 1b) in which there was subsequent ecological divergence driven by local

adaption (Rundell and Price 2009; Pyron et al. 2015)

•   If a sister-species pair is scored as sympatric and their leaf areas are different, we

interpreted this scenario as an event of parapatric speciation with ecological divergence

(Rundle and Nosil 2005; Rundell and Price 2009) (Fig. 1d).

• If a sister-species pair has overlapping geographical distributions and there is no

significant difference between their leaf areas, we interpreted this scenario as

inconclusive (Fig. 1c). This pattern could be the result of different processes, e.g.,

allopatric speciation with no ecological divergence followed by secondary contact

(Rundell and Price 2009; Hopkins 2013) or a parapatric speciation event driven by

ecological divergence in a trait other than leaf area (e.g., Snaydon and Davies 1976;

Silvertown et al. 2005).

We acknowledge that the role of parapatric ecological divergence may be underestimated

in this study because we only measured leaf area as an ecological proxy. Ecological traits

independent from leaf area can confer the ability to colonize different páramo niches (e.g. Cortés

et al. 2018), such as underground eco-physiological adaptations to soils with different water

saturation, or physiological adaptations at the anatomical and cellular level. Studying alternative

physio-ecological variables in sister species comparisons could shed light on other types of

ecological divergence (e.g. adaptations to different moisture in soils, microclimatic preferences).

Our framework assumes that:

1) Leaf area represents a good proxy for the organism's niche, which is likely true in our

focal genus. *Linochilus* leaf size is associated with their habitat, broad and large leaves are

present in species that dwell in the upper limit of the Andean forest, medium to small leaves are

characteristic of taxa inhabiting the páramo, and microphyllous leaves are found in lineages that

inhabit the upper boundary of the páramo (a.k.a. superpáramo) (Cuatrecasas 1969, Vargas &

Madriñán 2012).

2) The phylogeny employed represents the true species tree and contains all taxa. The

best hypothesis to date for the species tree in *Linochilus* (Vargas et al. 2017, based on double

digest restriction associated DNA data) contains only a subset of documented *Linochilus* species

(36 of 63). To account for the potential effects of missing species in our phylogeny, we searched

the taxonomic literature for the most morphologically similar species (assuming this is most

likely its sister species) to those species not sampled in the phylogeny and compared their

distributional ranges (Appendix S4). Specifically, we searched for new descriptions of species

where authors indicate the most similar taxon to the new species based on key traits. Diagnostic

characters in *Linochilus* include, but are not limited to, habit, the size and shape of the leaf, the

number of capitula per inflorescence, and several measurements in the corolla of both disk and

ray flowers. Alternative sisters based on morphology were added only to species included in the

first sister species analysis, aiming to maintain a similar number of pairs and make the results

comparable. We provide two sets of results: a) considering only the pairs derived from the

phylogeny and b) pairing non-sampled species to their most likely sister species present on the

phylogeny (we provide a discussion explaining why we believe this second set of results is more

reliable).

3) The páramo developed only in the last 2–4 Ma (van der Hammen and Cleef 1986;

Gregory-Wodzicki 2000) and it is therefore reasonable to assume that its carrying capacity for

the number of species has not been reached, and therefore extinction rates are low. Extremely

high diversification rates support the previous statement (Madriñán et al. 2013)

4) The modern species ranges are representative of past ranges, a reasonable assumption

given the recent divergence of sister taxa pairs.

EVALUATING NICHE CONSERVATISM

We used leaf area as a proxy to infer the ecological niche of *Linochilus* species. To test

for phylogenetic signal reflected in the evolution of the leaf area, we averaged log-transformed

leaf data and calculated Pagel's (1999) lambda (λ) using the R package PHYTOOLS (Revell

2012). To calculate λ, the observed phylogeny was compared to modified trees in which the

internal branches are compressed to various degrees. When λ = 0 the observed data follow a

model in which internal branches are completely collapsed (star phylogeny), meaning that the

trait evolves independently from the phylogeny. When λ = 1 the observed data follow a model in

which the internal branches are not modified, meaning that the evolution of the trait is

phylogenetically dependent (Harmon 2018; but see Revell et al. 2008).

BIOGEOGRAPHIC ANALYSIS

To elucidate the biogeographic history of *Linochilus* species, we performed a historical

biogeographic analysis. We defined the biogeographic areas based on the páramo complexes

defined by Londoño et al. (2014) which was restricted to Colombian páramos, and added three

complexes to completely cover the distribution of *Linochilus* (and the páramo). Contours of

páramo areas were edited with QGIS 2.8Wien (QGIS Development Team 2005). Our areas were

outlined as follows (areas not included by Londoño et al. (2014) are indicated with a star):

Northern Páramos (N). Páramos of the "Sierra Nevada de Santa Marta" and the "Serranía

del Perijá."

Talamanca* (T). Páramos located in the Talamanca Cordillera of Central America.

Mérida* (T). Páramos located in the Mérida Cordillera of Venezuela.

Eastern Cordillera (E). Páramos located in Eastern Cordillera of Colombia.

Antioquia (A). Cluster of páramos comprised by areas in the Western and Central

Cordilleras of Colombia mostly located in the department of Antioquia, Colombia.

Western Cordillera (W). Páramos located in the Western Cordillera of Colombia with the

exception of those located in the department of Antioquia, Colombia.

Central Cordillera (C). Páramos located in the Central Cordillera of Colombia with the

exception of those located in the department of Antioquia, Colombia.

Southern Páramos (S). Páramos located in the Colombian Massif and the Ecuadorian

Andes.

We pruned the chronogram of Vargas et al. (2017) to include only *Linochilus* species and

used BioGeoBEARS (Matzke 2013) to infer the biogeographic history of the genus.

BioGeoBEARS implements different models of ancestral range inference, DEC (Dispersal-

Extinction-Cladogenesis; Ree and Smith 2008), DIVALIKE (a likelihood version of Dispersal–

Vicariance Analysis; Ronquist 1997), and BAYAREAALIKE (a likelihood implementation of

the BAYAREA model; Landis et al. 2013), and it evaluates the addition of the J parameter

(Matzke 2014) to each one of the models to account for founder-event speciation (DEC+J,

DIVALIKE+J, BAYAREALIKE+J but see Ree and Sanmartín 2018). We opted not to use a

constrained model (e.g. limiting the presence areas to time windows based on their inferred

history) because the paleoaltitudes of the Northern and Central Andes are still debated (Luebert

and Weigend 2014). We assigned areas to tips based on the same occurrence dataset used for the

sister species comparisons (Appendix S1).

RESULTS

SISTER SPECIES COMPARISONS

TABLE 1 Relative signal of geographical isolation and parapatric ecological divergence in *Linochilus* based on sister species comparisons.

Sister sp. pairs based on:	Geographical isolation	Parapatric ecological divergence	Inconclusive
Phylogeny of Vargas et al. (2017)	9 (64.3%)	3 (21.4%)	2 (14.3%)
Phylogeny supplemented with missing sampling	12 (80%)	1 (6.6%)	2 (13.3%)

Independently of the framework used to infer allopatry/sympatry and the method

employed to infer sister species (phylogeny vs. phylogeny supplemented with missing sampling),

our results unequivocally reveal a strong signal of geographic isolation and little ecological

divergence *Linochilus* sister species (Tables 1–2, Fig 3, Appendix S5–7). When only species

included in the Vargas et al. (2017) phylogeny are considered (row 1, Table 1; Appendix S5–7),

nine out of 14 sister species pairs (64.3%) are allopatric, with only one of these pairs presenting

different leaf areas, suggesting divergence by geographical isolation with little ecological

divergence after speciation. A further three species pairs (21.4%) occur sympatrically and have

evidence of ecological divergence in the leaf areas, while two cases (14.3%) present inconclusive

evidence. When species not sampled in the phylogeny of Vargas et al. (2017) are considered (see

Appendix S4) by pairing them to their most likely sampled sister species (row 2, Table 1–2, Fig

3) the contribution of allopatric speciation cases increase to 12 out of 15 (80%), while the

number of cases for parapatric ecological divergence decreases to 1 (6.6%); the number of

inconclusive cases remains about the same, 2 (13.3%). The aforementioned results are virtually

the same to using grids of 0.1 and 0.05 decimal degrees to quantify range overlap (Table 3,

Appendix S6). For example, range overlap calculated with 0.1 grid produces the same results as

our island framework in the analysis supplemented with missing sampling (Tables 2–3). The

most fine-scale analysis, using a grid of 0.05 decimal degrees, flips one of the pairs (*L. romeroi –*

L. saxatilis) from sympatry to allopatry (this pair is likely to be parapatric with species

occupying different vegetational belts), with this pair being the only conflicting result among the

approaches.

TABLE 2 Sister species comparisons based on the phylogeny of Vargas et al. (2017) and supplemented
with non-sampled species (underlined). Allopatry and sympatry is determined based on overlapping
occurrences on paramo islands as defined by Londoño et al. 2014.

Sister 1	Sister 2	Wilcoxon test leaf	Distribution	Inferred divergence	Age (Ma)
L. phylicoides	*L. lacunosus*	0.4722	Allopatric	Geog. isolation	0.16 (0.01–0.58)
L. obtusus	*L. venezuelensis*	1.0000	Sympatric	Inconclusive	0.22 (0.01–0.81)
L. violaceus	*L. cinerascens*	0.2897	Allopatric	Geog. isolation	0.28 (0.01–0.86)
L. rosmarinifolius	*L. cyparissias*	–	Allopatric	Geog. isolation	–
L. floribundus	*L. farallonensis / perijaensis*	–	Allopatric	Geog. isolation	–
L. rhomboidalis	*L. apiculatus*	2.50E-09*	Allopatric	Geog. isolation	0.59 (0.01–1.54)
L. alveolatus	*L. costaricensis*	1.0000	Allopatric	Geog. isolation	0.56 (0.01–1.46)
L. rhododendroides	*L. schultzii*	1.0000	Allopatric	Geog. isolation	0.92 (0.02–2.14)
L. tenuifolius	*L. ellipticus*	–	Allopatric	Geog. isolation	–
L. oblongifolius	*L. mutiscuanus*	0.4140	Allopatric	Geog. isolation	0.20 (0.01–0.76)
L. sp. nov. ANT	*L. antioquensis*	0.1663	Allopatric	Geog. isolation	0.52 (0.01–1.47)
L. huertasii	*L. julianii*	–	Allopatric	Geog. isolation	1.41 (0.05–3.18)
L. eriophorus	*L. chrysotrichus*	–	Allopatric	Geog. isolation	–
L. colombianus	*L. glutinosus*	1.20E-11*	Sympatric	Ecological	2.97 (0.11–6.17)
L. romeroi	*L. saxatilis*	–	Sympatric	Inconclusive	–

EVALUATING NICHE CONSERVATISM

Our results suggest that there is a strong signal of niche conservatism in the leaf area of

Linochilus. Graphing the boxplots of leaf areas per species in front of the phylogeny reveals a

pattern in which closely related species tend to have similar leaf areas, likely occupying similar

niches (Fig. 4). These observations are supported by the Pagel's (1999) lambda (λ) of 0.98

calculated with leaf area data on the phylogeny, P= 2.8e-09 against the null hypothesis of λ=0.

This Pagel's lambda of almost one suggests that the evolution of the leaf area is highly

dependent on the phylogeny, possessing a strong phylogenetic signal. The phylogenetic signal

found, builds on the sister species comparisons, where only a small fraction of pairs (13.3%)

presents statistically significantly different leaf areas (Table 2).

TABLE 3 Range overlap calculated using 0.1 and 0.05 decimal degrees grids. *Linochilus* sister species
comparisons based on the phylogeny of Vargas et al. (2017) supplemented with non-sampled species
(underlined). *L. floribundus* is found twice in the table because there are two hypothesized sisters for it in
the taxonomic literature.

		Range overlap		Range asymmetry	
Sister 1	**Sister 2**	**0.1**	**0.05**	**0.1**	**0.05**
L. phylicoides	*L. lacunosus*	0	0	10.7	16.4
L. obtusus	*L. venezuelensis*	0.4	0.2	1.0	1.1
L. violaceus	*L. cinerascens*	0	0	5.7	6.3
L. rosmarinifolius	*L. cyparissias*	0	0	21.8	25.3
L. floribundus	*L. farallonensis*	0	0	42.0	46.0
L. floribundus	*L. perijaensis*	0	0	42.5	46.6
L. rhomboidalis	*L. apiculatus*	0	0	1.7	1.5
L. alveolatus	*L. costaricensis*	0	0	15.2	8.1
L. rhododendroides	*L. schultzii*	0	0	4.0	5.2
L. tenuifolius	*L. ellipticus*	0	0	17.1	20.1
L. oblongifolius	*L. mutiscuanus*	0	0	3.0	4.0
L. sp. nov. ANT	*L. antioquensis*	0	0	1.0	1.0
L. huertasii	*L. julianii*	0	0	7.0	7.0
L. eriophorus	*L. chrysotrichus*	0	0	7.0	9.0
L. colombianus	*L. glutinosus*	0.2	0.2	2.6	2.6
L. romeroi	*L. saxatilis*	1.0	0	3.0	1.5

FIGURE 3 Distribution of sister species based on the phylogeny Vargas et al. (2017) supplemented with missing sampling. Darker pixels indicate higher elevations, orange polygons delineate paramo complexes. ∗ : allopatric based on páramo islands. + : allopatric based on 0.1 decimal degree grid analysis. ‡ : allopatric based on 0.05 decimal degree grid analysis. Pairs *L. rosmarinifolius–L. cyparissias* and *L. oblongifolius–L. mutiscuanus,* all inhabiting upper boundary of the cloud forest, are codified as allopatric in the island framework as they inhabit different slopes in the mountains they co-occur.

BIOGEOGRAPHIC ANALYSES

The best scoring model in the biogeographic reconstruction for *Linochilus* was the

BAYAREALIKE+J with an AICc of 226.78 followed by DEC+J with 232.62 (Table 4). The

BAYAREALIKE+J reconstruction (Fig. 5) shows that the Colombian Eastern Cordillera played

a major role in the diversification of *Linochilus.* This area, which contains the most species of the

genus, was shown to be the ancestral range for most *Linochilus* ancestors (61%). The ancestral

range for the node representing the ancestor for all *Linochilus* species is inconclusive, with two

areas sharing aproximately two thirds of the relative probability: the Eastern Cordillera and the

Northern Colombian Páramos (Sierra Nevada de Santa Marta + the Serranía del Perijá). The

reconstruction shows that the *Denticulata* clade (clade D, Fig. 5), which downslope colonized the

upper limit of the cloud forest from the páramo, originated in the Eastern Colombian Cordillera.

The biogeographic analysis also shows that approximately two thirds of the species sampled are

restricted to a single paramo complex (27 spp., 71%), while the other third is found in two or

more complexes (11 spp. 29%). These biogeographic results evidence high endemism in the

genus where most species are restricted to a few paramo islands, reinforcing the importance of

geographic isolation in the diversification of the genus.

360
FIGURE 4. *Linochilus* phylogeny (Vargas et al. 2017) with boxplots for leaf area. Blue bars indicate
confidence intervals for node ages. Photos on the left correspond to the superscript numeration on the tips
of the phylogeny; photos 1 and 2 correspond to sympatric sister species with different leaf areas, while
photos 3 and 4 correspond to allopatric sister species with similar leaf areas. Photos on the right
correspond to the species indicated by the gray dashed line. The *Denticulata* clade is indicated by the
letter D. Pairs are indicated by gray boxes, * : allopatric, ♣ : different leaf areas.

TABLE 4 Comparison of the different biogeographic models for ancestral range inference evaluated by
BioGeoBEARS in the phylogeny of *Linochilus* (Vargas et al. 2017).

Model	LnL	Num. params	AIC
DEC	-115.09205	2	234.184098
DEC+J	-113.31082	3	232.621645
DIVALIKE	-115.63941	2	235.278824
DIVALIKE+J	-115.64036	3	237.280724
BAYAREALIKE	-121.933	2	247.865999
BAYAREALIKE+J	-110.39081	3	226.78162

369
370
FIGURE 5 BAYAREALIKE+J biogeographical ancestral reconstruction based on the *Linochilus*
phylogeny of Vargas et al. (2017) with percent probabilities of the different ancestral areas as pie charts.
Letters indicate biogeographic areas considered in the analysis. N: northern páramos. T: Talamanca. M:
Mérida. E: Eastern Cordillera. A: Antioquia's páramos. W: Western Cordillera. C: Central Cordillera. S:
southern páramos. Letters above each pie charts indicate the most probable area or area combination for
that node.

DISCUSSION

In this study, we developed a framework to quantify the relative contributions of

geographic isolation and parapatric ecological divergence in recent speciation events. We

incorporated phylogenetics, geographic distributions, and a morpho-ecological trait. Our

framework was applied to *Linochilus*, a genus of plants restricted to the páramo, which is the

most species-rich tropical montane ecosystem. Because of the island-like distribution of the

páramo and the within-páramo elevation gradient (Fig 2.), it has been suggested that allopatric

speciation and parapatric ecological divergence are the main drivers of speciation in the high

Andes (van der Hammen and Cleef 1986, Hughes and Atchison 2015). The application of our

approach to *Linochilus* revealed that most recent speciation events are allopatric (Table 1), with

most sister species presenting allopatry and similar leaf areas. The signal of allopatric speciation

between sister species is robust independently of the method used to estimate allopatry/sympatry.

64.3% sisters are allopatric using presence/absence in páramo islands and the sister species

inferred from Vargas et al. (2017). 80% sisters are allopatric when missing sampling is

accounted for. Allopatry is inferred similarly (relative to presence/absence in paramo islands) for

most sisters when using grids of 0.1 and 0.05 decimal degrees (Tables 1–3). Likewise, the

contribution of parapatric ecological divergence to recent speciation events is low ($< 25\%$) in

both sets of results, phylogeny: 21.4% and phylogeny supplemented with missing sampling:

6.6%.

We believe that the most reliable estimate for calculating the relative contribution of

allopatric speciation vs. parapatric ecological divergence is the phylogeny supplemented with

missing sampling on the island framework results (second row, Table 1; Table 2). These results

agree with the 0.1 grid results, are corrected for missing sampling in the phylogeny, and are more

conservative at estimating allopatric pairs (relative to using a 0.05 degree grid). Therefore, *we*

will focus our discussion on the species pairs derived from the phylogeny and supplemented for

missing sampling (Tables 2–3).

ALLOPATRIC VS. PARAPATRIC ECOLOGICAL SPECIATION IN THE PÁRAMO

Our results suggest that most recent speciation events (12, 80%) in *Linochilus* are driven by allopatric speciation, while only a few are driven by parapatric ecological divergence events (1, 6.6%) (Table 2, Fig. 3). Out of the 12 allopatric pairs, only one shows a signal of subsequent ecological divergence. Inconclusive divergence in pairs (2, 13.3%) was represented by a sympatric pair with similar leaf areas, and a sympatric pair with no leaf data. Independently of range overlap, only two pairs haven significantly different leaf areas. This lack of ecological divergence suggests a strong signal of niche conservatism in *Linochilus*, as does a Pagel's λ of 0.98 (a value closer to one indicates strong phylogenetic signal in the evolution of leaf area). In the context of the páramo flora, our results are consistent with the hypothesis that páramos are island-like in promoting allopatric speciation. Our results suggest that allopatric speciation alone can explain most of the recent and rapid speciation found in other páramo genera where most species are restricted to one or few páramo islands: *Bartsia*, *Espeletia, Escallonia*, *Hypericum*, *Jamesonia-Eriosorus, Lachemilla,* and *Lupinus,* (Drummond et al. 2012; Zapata 2013; Sánchez-Baracaldo and Thomas 2014; Nürk et al. 2015; Uribe-Convers and Tank 2015; Diazgranados and Barber 2017; Contreras-Ortiz et al. 2018; Morales-Briones et al. 2018ab).

Although our results suggest a minor role of parapatric ecological divergence (6.6%) (Table 1), our analysis focused on sister species that diverged recently and does not consider more ancient divergent events. However, when we look at the distribution of leaf area on the phylogeny, we observe that taxa in clade D (*Linochilus* series *Denticulata*) have significantly larger leaves suited for dwelling in the upper zone of Andean forest (Fig. 3): Wilcoxon P < 2.2e-16 for both *L.* ser. *Denticulata* vs. its sister clade, and *L.* ser. *Denticulata* vs. *Linochilus*'s most species-rich clade (the clade originating with the most common ancestor of *L. heterophyllus* and

L. phylicoides). The significantly larger leaves in *L.* ser. *Denticulata* suggests that an event of

ecological shift, from microphyllous to macrophyllous leaves, took place ca. 3 Ma near the

ancestor of the *Denticulata* clade leading to the evolution of at least 20 species (32% out of all

species in *Linochilus*). Based on the larger leaf area found in the *Denticulata* clade, which allows

Linochilus species to be competitive in the upper zone of the cloud forest, we hypothesize that

this downslope colonization event by the *Denticulata* clade is a case ecological divergence.

Regardless of the reason for ecological divergence (after allopatric speciation or during

parapatric ecological divergence), ecological divergence may *boost* allopatric speciation by

allowing a lineage to colonize a new niche and rapidly speciate in it by means of allopatric

speciation in an island-like system. In the specific case of *Linochilus*, the evolution of larger

leaves, which happened in one single event ca. 3 Ma, allowed a lineage to colonize a lower

vegetational belt and speciate. A similar pattern is found in Andean *Senecio,* where

monophyletic forest and páramo clades have been documented to speciate in parallel (Dušková

et al. 2017). In the context of adaptive radiations, we propose that ecological speciation happens

fewer times when compared with allopatric speciation, but its effect could be significant at

geological time scales; in other words, while allopatric speciation can explain most of the recent

speciation, ecological divergence explains the origin of a single (or few) key adaptation(s)

facilitating the colonization of new niches. *Espeletia*, whose crown origin is estimated at 2.5 Ma,

shows a peak in morphological differentiation relatively deep in its phylogeny at 1.5 Ma

(Pouchon et al. 2018) in addition to have a strong signal of allopatric speciation (Diazgranados

and Barber 2017). Other potential diversification *boosters* are genome duplication events

(Morales-Briones et al. 2018a) and pollination shifts (Lagomarsino et al. 2016).

ALTERNATIVE HYPOTHESES AND FURTHER CONSIDERATIONS

Our framework is unable to distinguish between vicariant and peripatric speciation. Testing which kind of allopatric speciation occurs is challenging in island systems—typically peripatric speciation predicts that the founder species will be distributed in a significantly smaller area than the source lineage (Coyne and Orr 2004; Anacker and Strauss 2014; Grossenbancher et al. 2014; Skeels and Cardillo 2019), but geographically restricted patches (e.g. páramos) can confound testing for significant differences in the distributions. Additionally, páramo islands have shifted their elevational distribution due to Pleistocene climate fluctuations, connecting páramo islands during glaciation periods and disconnecting them during interglacial periods (Simpson 1974, 1975; Flantua et al. 2019). This "flickering connectivity" could be a major driver of geographic isolation in páramo plants, especially for taxa with low seed dispersal ability (i.g. *Espeletia* complex with ca. 100 spp.). In *Linochilus*, which has a fruit that easily disperses with wind, we find that 10 sisters (66.6%) present range asymmetry in which the widespread-sister range is > 3 times larger than the small-ranged sister, suggesting peripatric speciation as a lead player in the diversification of *Linochilus;* we advise to take this result with caution for the aforementioned reasons. Finally, the best fit of the BAYAREALIKE+J as the best model for the biogeographic analysis in *Linochilus* supports the idea that founder speciation events (peripatric speciation), modeled by the J parameter, are important for the speciation of *Linochilus* in the páramos.

Our sister-species framework assumes that speciation is a bifurcating process in which every speciation event produces two reciprocal monophyletic species. In the context of vicariant speciation in the páramo, a glacial-interglacial event could result in the fragmentation of one previously continually distributed population, into more than two daughter proto-species; the

complex topography of the Colombian Eastern Cordillera provides a probable location for this

process to occur (Fig 2, Fig 7-10 in van der Hammen and Cleef 1986). Hypothetically, parapatric

and peripatric speciation can also violate a bifurcating speciation model because a widely

distributed population could be the source for multiple independent parapatric and or peripatric

divergent events (e.g. upslope colonization and adaptation in different mountains, multiple

dispersal events, Dexter et al. 2017).

How does the result of allopatric speciation driving diversification align with recent finds

of pervasive hybridization-introgression in *Linochilus* (Vargas et al. 2017) and other páramo

plants (Morales-Briones et al. 2018a; Nevado et al. 2018; Pouchon et al. 2018)? High gene flow

indices can be explained by three non-exclusive scenarios. First, under allopatric speciation,

recent geographically isolated populations (proto-species) should be reproductively compatible

for the time that it takes to reproductive barriers to develop. During that time window, gene

transfer between these proto-species is likely to occur leaving a signal of introgression among

sister species. Second, it is possible that, given geographic isolation, the biological pressure to

enforce reproductive barriers is relaxed, making geographically isolated proto-species

reproductively compatible for a long period of time; this would allow hybridization post-

morphological differentiation—that is, hybridization after the time point when they are

morphologically different and therefore considered by taxonomists as different species. Third,

considering the small geographical ranges and likely low population numbers of many

Linochilus species, it is possible that introgression has a positive effect by increasing genetic

variability and counteracting genetic drift —making gene flow adaptive. Testing these

hypotheses with phylogeographic approaches would help to elucidate the role of hybridization in

the evolution of páramo plants.

Finally, it has been suggested that hybridization can cause speciation because it has the

potential of producing genotypes pre-adapted to un-exploited niches (Anderson and Stebbins

1954; Seehausen 2004); this hypothesis predicts a signal of ecological divergence between the

hybrid and its parental species. Taking into account that we only observed two cases of

parapatric ecological divergence in sister species, our results suggest that hybrid speciation might

not play a major role in the recent diversification history of *Linochilus*.

SPATIOTEMPORAL HISTORY OF *LINOCHILUS* IN THE CONTEXT OF THE

PÁRAMO

Our biogeographic reconstruction of *Linochilus* shows that the genus originated in the

Northern Andes 6.46 Ma, predating the estimated origin of the páramo 2–4 Ma (van der

Hammen and Cleef 1986; Gregory-Wodzicki 2000). Despite the fact that *Linochilus*'s inferred

age 95% confidence interval of 1.71–11.37 can accommodate such incongruence, other primarily

páramo genera like *Arcytophyllum, Brunfelsia, Jamesonia+Eriosorus, Lysipomia, Valeriana,* and

Vasconcellea also show ages older than 4 Ma (Luebert and Weigend 2014). An explanation for

this early origin could be that ancestors of these lineages inhabited the summits of middle

elevation mountains (<2000 m) extant at that time. Mid-elevation tropical mountains often have

open and semi-dry areas at upper elevations, which are somewhat physiologically similar to the

páramo. These physiologically dry patches are caused by well-drained soils and strong winds

similar to the contemporary *campos de altitude* and *campus rupestres* in Brazil (Safford 1999;

Alves et al. 2014). It is possible that middle elevation mountaintops provided an early habitat for

Linochilus ancestors before higher elevations were available at 4 Ma. A second alternative is that

páramos were available before 2–4 Ma as suggested by Ehlers and Poulsen (2009). A third

scenario is that *Linochilus* originated in the Sierra Nevada de Santa Marta (SNSM), a mountain

range located in northern Colombia which is now separated from the main Andes Cordilleras.

SNSM's paleoelevation remains largely unstudied (Villagómez et al. 2011), but our

biogeographic analysis provides some evidence for the last hypothesis because the *Linochilus*

species endemic to the SNSM (*L. coriaceus, L. inesianus,* and *P. romeroi*) comprise a clade that

that is sister to the rest of *Linochilus*, making the SNSM the second most likely area of origin for

the genus after the Eastern Cordillera (Fig. 5).

Our biogeographic analysis also suggests that the Eastern Cordillera of Colombia played

a major role in the diversification of the genus given the many extant and ancestral species

whose distributional range include this area. The Eastern Cordillera contains the most páramo

land area with discrete patches (Londoño et al. 2014) making it ideal for autochthonous

allopatric speciation. Our reconstruction also indicates that the Eastern Cordillera was the source

for the colonization of three other mountain ranges: the Colombian Central Cordillera, the

Colombian Western Cordillera, and the Talamanca Cordillera in Central America.

Achenes of *Linochilus* are small and a have a pappus that allows for long-distance

dispersal of their seeds by wind (Cuatrecasas 1969). Collectively, *Linochilus* is found on almost

every páramo island, with the exception of the southernmost páramos, south of the Girón-Paute

valley in Ecuador, which is a biogeographic barrier for numerous Andean genera (Jørgensen et

al. 1995). Specific examples of long-distance dispersal in *Linochilus* are shown by the two

species reported at the westernmost páramos in Costa Rica. *L. costaricensis* is endemic to Costa

Rica and is probably a direct descendant of *L. alveolatus*, which inhabits the Colombian Eastern

Cordillera. The second Costa Rican species, *L. floribundus,* is also reported for Colombia and

Ecuador (Vargas 2011; 2018). The aforementioned examples of long-distance dispersal build on

previous results about the potential dominant role of peripatric speciation in the genus and in

other high montane taxa with seeds or fruits adapted to wind dispersal.

CONCLUSIONS

Our comparative framework that incorporates phylogenetics, geographical distributions,

and morpho-ecological characters unveiled a high signal of allopatric speciation, supporting it as

the process driving most of the recent speciation events in *Linochilus*. The island-like

distribution of the páramo is likely a primary factor of autochthonous allopatric speciation via

geographic isolation, explaining the particularly high accumulation of plant species in the

páramo (Simpson and Todzia 1990) and their high speciation rates (Madriñán et al. 2013).

Despite the comparatively small role of parapatric ecological speciation identified in recent

Linochilus sister taxa, we propose that ecological divergence has a role that is infrequent but

potentially powerful in island-like systems. When ecological divergence does occur, it allows a

lineage to colonize a new niche and then rapidly speciate by means of allopatric speciation

among islands. Ecological divergence events that *boost* diversification are thus expected to be

detectable at deeper geological time scales (> 1 Ma). We conclude that geographic isolation and

parapatric ecological divergence are positively synergistic processes in the history of the

diversification of the paramo flora, contributing significantly to the global latitudinal species

gradient.

ACKNOWLEDGEMENTS

We thank Stefani Brandt, Julia Harencar, Shelley Sianta, Kathleen Kay, and Dena

Grossenbacher for helpful comments and discussion during the writing of this manuscript. This

study would not have been possible without the scientific collections housed at Herbario

Nacional Colombiano (COL), the Billie Turner Plant Resources Center (TEX), and the United

States National Herbarium at the Smithsonian Institution (US). Financial support was provided

by The University of Texas at Austin (Plant Biology Program Awards, the C. L. Lundell Chair of

Systematic Botany, The Linda Escobar Award), the Garden Club of America (2012 Award in

Tropical Botany), and the Smithsonian Institution (Cuatrecasas Award 2006).

566 CONFLICT OF INTEREST

567 We declare no competing interest.

568 REFERENCES

Alves, R. J. V, N. G. Silva, J. A. Oliveira, and D. Medeiros. 2014. Circumscribing campo
rupestre - megadiverse Brazilian rocky montane savanas. Brazilian J. Biol. 74:355–362.

Anderson, E., and G. L. Stebbins. 1954. Hybridization as an evolutionary stimulus. Evolution
8:378–388.

Barthlott, W., W. Lauer, and A. Placke. 1996. Global distribution of species diversity in vascular
plants: towards a world map of phytodiversity. Erdkunde 50:317–327.

Barraclough, T. G., and A. P. Vogler. 2000. Detecting the geographical pattern of speciation
from species-level phylogenies. Am. Nat. 155:419–434.

Blake, S. F. 1928. Review of the genus *Diplostephium*. Am. J. Bot. 15:43–64.

Bonhomme, V., S. Picq, C. Gaucherel, and J. Claude. 2014. Momocs: Outline Analysis Using R.
J. Stat. Softw. 56:1–24.

Carstens, B. C., and L. L. Knowles. 2007. Shifting distributions and speciation: Species
divergence during rapid climate change. Mol. Ecol. 16:619–627.

Contreras-Ortiz, N., G. Atchison, C. Hughes, and S. Madriñán. 2018. Convergent evolution of
high elevation plant growth forms and geographically structured variation in Andean
Lupinus (Leguminosae). Bot. J. Linn. Soc. 187:118–136.

Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinauer.

Cortés, A. J., L. N. Garzón, J. B. Valencia, and S. Madriñán. 2018. On the Causes of Rapid
Diversification in the Páramos: Isolation by Ecology and Genomic Divergence in
Espeletia. Front. Plant Sci. 9.

Cuatrecasas, J. 1968. Páramo vegetation and its life forms. Colloq. Geogr. 9:163–186.

Cuatrecasas, J. 1969. Prima Flora Colombiana. 3. Compositae-Astereae. Webbia 24:1–335.

Diazgranados, M., and J. Barber. 2017. Geography shapes the phylogeny of frailejones
(Espeletiinae Cuatrec., Asteraceae): a remarkable example of recent rapid radiation in sky
islands. PeerJ 5:e2968.

Dexter, K. G., M. Lavin, B. M. Torke, A. D. Twyford, T. A. Kursar, P. D. Coley, C. Drake, R.
Hollands, and R. T. Pennington. 2017. Dispersal assembly of rain forest tree communities
across the Amazon basin. Proc. Natl. Acad. Sci. U. S. A. 114:2645–2650.

Drummond, C. S., R. J. Eastwood, S. T. S. Miotto, and C. E. Hughes. 2012. Multiple Continental
Radiations and Correlates of Diversification in *Lupinus* (Leguminosae): Testing for Key
Innovation with Incomplete Taxon Sampling. Syst. Biol. 61:443–460.

Dušková, E., P. Sklenář, F. Kolář, D. L. A. Vásquez, K. Romoleroux, T. Fér, and K. Marhold.
2017. Growth form evolution and hybridization in *Senecio* (Asteraceae) from the high
equatorial Andes. Ecol. Evol. 7:6455–6468.

Flantua, S. G. A., A. O'Dea, R. E. Onstein, C. Giraldo, and H. Hooghiemstra. 2019. The
flickering connectivity system of the north Andean páramos. J. Biogeogr. 46:1808–1825.

Ehlers, T. A., and C. J. Poulsen. 2009. Influence of Andean uplift on climate and paleoaltimetry
estimates. Earth Planet. Sci. Lett. 281:238–248. Elsevier B.V.

Fitzpatrick, B. M., and M. Turelli. 2006. The geography of mammalian speciation: mixed signals
from phylogenies and range maps. Evolution 60:601–615.

Givnish, T. J. 1987. Comparative studies of leaf form: assessing the relative roles of selective
pressures and phylogenetic constraints. New Phytol. 106:131–160.

Givnish, T. J. 1997. Adaptive radiation and molecular systematics: aims and conceptual issues.
Pp. 1–54 *in* Molecular evolution and adaptive radiation. Cambridge University Press,
New York, USA.

Gregory-Wodzicki, K. M. 2000. Uplift history of the Central and Northern Andes: a review.
Geol. Soc. Am. Bull. 112:1091–1105.

Grossenbacher, D. L., S. D. Veloz, and J. P. Sexton. 2014. Niche and range size patterns suggest
that speciation begins in small, ecologically diverged populations in North American
monkeyflowers (*Mimulus* spp.). Evolution 68:1270–1280.

Harmon, L. J. 2018. Phylogenetic comparative methods: Learning from trees. Available via
https://doi.org/10.32942/osf.io/e3xnr

Hijmans, R. J. 2022. Raster: geographic data and modelling. R package version 3.5-21. Available
via https://CRAN.R-project.org/packageraster.

Hopkins, R. 2013. Reinforcement in plants. New Phytol. 197:1095–1103.

Hughes, C. E., and G. W. Atchison. 2015. The ubiquity of alpine plant radiations: from the
Andes to the Hengduan Mountains. New Phytol. 207:275–282.
Janzen, D. H. 1967. Why mountain passes are higher in the tropics. Am. Nat. 101:233–249.

Jenkins, C. N., S. L. Pimm, and L. N. Joppa. 2013. Global patterns of terrestrial vertebrate
diversity and conservation. Proc. Natl. Acad. Sci. 110:E2602–E2610.

Jørgensen, P. M., C. Ulloa-Ulloa, and J. Madsen. 1995. A floristic analysis of the high Andes of
Ecuador. Pp. 221–237 *in* S. P. Churchill, ed. Biodiversity and Conservation of
Neotropical Montane Forest. The New York Botanical Garden, New York.
Lagomarsino, L. P., F. L. Condamine, A. Antonelli, A. Mulch, and C. C. Davis. 2016. The
abiotic and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae).
New Phytol. 210:1430–1442.

Landis, M. J., N. J. Matzke, B. R. Moore, and J. P. Huelsenbeck. 2013. Bayesian analysis of
biogeography when the number of areas is large. Syst. Biol. 62:789–804.

Londoño, C., A. Cleef, and S. Madriñán. 2014. Angiosperm flora and biogeography of the
páramo region of Colombia, Northern Andes. Flora Morphol. Distrib. Funct. Ecol. Plants
209:81–87.

Löwenberg-Neto, P. 2014. Neotropical region: A shapefile of Morrone's (2014) biogeographical
regionalisation. Zootaxa 3802:300.

Luebert, F., and M. Weigend. 2014. Phylogenetic insights into Andean plant diversification.
Front. Ecol. Evol. 2:1–17.

Luteyn, J. L. 1992. Páramos: why study them. Pp. 1–14 *in* J. L. Luteyn and H. Baslev, eds.
Paramo: an Andean ecosystem under human influence. London, UK.

Luteyn, J. L. 1999. Páramos: a checklist of plant diversity, geographical distribution, and
botanical literature. The New York Botanical Garden, New York, USA.

Madriñán, S., A. J. Cortés, and J. E. Richardson. 2013. Páramo is the world's fastest evolving
and coolest biodiversity hotspot. Front. Genet. 4:192.

Matzke, N. J. 2013. BioGeoBEARS: Biogeography with Bayesian (and likelihood) evolutionary
analysis in R Scripts, CRAN: The Comprehensive R Archive Network, Vienna, Austria.

Matzke, N. J. 2014. Model selection in historical biogeography reveals that founder-event
speciation is a crucial process in island clades. Syst. Biol. 63:951–970.

Morales-Briones, D. F., A. Liston, and D. C. Tank. 2018a. Phylogenomic analyses reveal a deep
history of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae).
New Phytol., doi: 10.1111/nph.15099.

Morales-Briones, D. F., K. Romoleroux, F. Kolář, and D. C. Tank. 2018b. Phylogeny and
evolution of the neotropical radiation of *Lachemilla* (Rosaceae): uncovering a history of
reticulate evolution and implications for infrageneric classification. Syst. Bot. 43:17–34.

Muños-Ortiz, A., Á. A. Velásquez-Alvarez, C. E. Guarnizo, and A. J. Crawford. 2015. Of peaks
and valleys: testing the roles of orogeny and habitat heterogeneity in driving allopatry in
mid-elevation frogs (Aromobatidae: *Rheobates*) of the northern Andes. J. Biogeogr.
42:193–205.

Mutke, J., and W. Barthlott. 2005. Patterns of vascular plant diversity at continental to global
scales. Biol. Skr. 55:521–537.

Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. da Fonseca, and J. Kent. 2000.
Biodiversity hotspots for conservation priorities. Nature 403:853–858.

Nevado, B., N. Contreras-Ortiz, C. Hughes, and D. A. Filatov. 2018. Pleistocene glacial cycles
drive isolation, gene flow and speciation in the high-elevation Andes. New Phytol.
219:779–793.

Nürk, N. M., S. Uribe-Convers, B. Gehrke, D. C. Tank, and F. R. Blattner. 2015. Oligocene
niche shift, Miocene diversification – cold tolerance and accelerated speciation rates in
the St. John's Worts (*Hypericum*, Hypericaceae). BMC Evol. Biol. 15:1–13.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Pérez-Escobar, O. A., G. Chomicki, F. L. Condamine, A. P. Karremans, D. Bogarín, N. J.
Matzke, D. Silvestro, and A. Antonelli. 2017. Recent origin and rapid speciation of
Neotropical orchids in the world's richest plant biodiversity hotspot. New Phytol.
215:891–905.

Pouchon, C., A. Fernández, J. M. Nassar, F. Boyer, S. Aubert, S. Lavergne, and J. Mavárez.
2018. Phylogenomic Analysis of the Explosive Adaptive Radiation of the *Espeletia*
Complex (Asteraceae) in the Tropical Andes. Syst. Biol. 67:1041–1060.

Pyron, R. A., G. C. Costa, M. A. Patten, and F. T. Burbrink. 2015. Phylogenetic niche
conservatism and the evolutionary basis of ecological speciation. Biol. Rev. 90:1248–
1262.

QGIS Development Team. 2005. QGIS Geographic Information System. Open source geospatial
foundation project. Open source geospatial foundation project.

R Core Team. 2016. R: A Language and Environment for Statistical Computing. Vienna, Austria.

Ree, R. H., and I. Sanmartín. 2018. Conceptual and statistical problems with the DEC+J model
of founder-event speciation and its comparison with DEC via model selection. J.
Biogeogr. 45:741–749.

Ree, R. H., and S. A. Smith. 2008. Maximum Likelihood Inference of Geographic Range
Evolution by Dispersal, Local Extinction, and Cladogenesis. Syst. Biol. 57:4–14.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other
things). Methods Ecol. Evol. 3:217–223.

Revell, L. J., L. J. Harmon, and D. C. Collar. 2008. Phylogenetic signal, evolutionary process,
and rate. Syst. Biol. 57:591–601.

Ronquist, F. 1997. Dispersal-Vicariance Analysis: A New Approach to the Quantification of
Historical Biogeography. Syst. Biol. 46:195–203.

Rundell, R. J., and T. D. Price. 2009. Adaptive radiation, nonadaptive radiation, ecological
speciation and nonecological speciation. Trends Ecol. Evol. 24:394–399.

Rundle, H. D., and P. Nosil. 2005. Ecological speciation. Ecol. Lett. 8:336–352.

Saldivia, P., O. M. Vargas, D. A. Orlovich, and J. M. Lord. 2019. Nomenclatural priority of the
genus *Linochilus* over *Piofontia* (Asteraceae: Astereae). Phytotaxa 424:158–166.

Safford, H. D. 1999. Brazilian Páramos II. Macro- and mesoclimate of the *campos de altitude*
and affinities with high mountain climates of the tropical Andes and Costa Rica. J.
Biogeogr. 26:713–737.

Sánchez-Baracaldo, P., and G. H. Thomas. 2014. Adaptation and convergent evolution within
the *Jamesonia-Eriosorus* complex in high-elevation biodiverse Andean hotspots. PLoS
One 9:e110618.

Schluter, D. 2000. The ecology of adaptive radiation. Oxford University Press, Oxford, UK.

Seehausen, O. 2004. Hybridization and adaptive radiation. Trends Ecol. Evol. 19:198–207.

Silvertown, J., C. Servaes, P. Biss, and D. Macleod. 2005. Reinforcement of reproductive
isolation between adjacent populations in the Park Grass Experiment. Heredity (Edinb).
95:198–205.

Simpson, B. B. 1974. Glacial migrations of plants: island biogeographical evidence. Science
185:698–700.

Simpson, B. B. 1975. Pleistocene changes in the flora of the high tropical Andes. Paleobiology
1:273–294.

Simpson, B. B., and C. A. Todzia. 1990. Patterns and processes in the development of the high
Andean flora. Am. J. Bot. 77:1419–1432.

Simpson, G. G. 1953. The Major Features of Evolution. Columbia University Press, New York,
USA.

Skeels, A., and M. Cardillo. 2019. Reconstructing the geography of speciation from
contemporary biodiversity data. Am. Nat. 193.

Sklenář, P., I. Hedberg, and A. M. Cleef. 2014. Island biogeography of tropical alpine floras. J.
Biogeogr. 41:287–297.

Snaydon, R. W., and M. S. Davies. 1976. Rapid population differentiation in a mosaic
environment IV. Populations of *Anthoxanthum odoratum* at sharp boundaries. Heredity
(Edinb). 37:9–25.

Uribe-Convers, S., and D. C. Tank. 2015. Shifts in diversification rates linked to biogeographic
movement into new areas: an example of a recent radiation in the Andes. Am. J. Bot.
102:1854–1869.

van der Hammen, T., and A. M. Cleef. 1986. Development of the high Andean páramo flora and
vegetation. Pp. 153–201 *in* F. Vuilleumier and M. Monasterio, eds. High Altitude
Tropical Biogeography. Oxford University Press, New York, USA.

Vargas, O. M. 2011. A nomenclator of *Diplostephium* (Asteraceae: Astereae): a list of species
with their synonyms and distribution. Lundellia 14:32–51.

Vargas, O. M. 2018. Reinstatement of the Genus *Piofontia*: A Phylogenomic-based Study
Reveals the Biphyletic Nature of *Diplostephium* (Asteraceae: Astereae). Syst. Bot.
43:485–496.

Vargas, O. M., and S. Madriñán. 2012. Preliminary phylogeny of *Diplostephium* (Asteraceae):
speciation rate and character evolution. Lundellia 15:1–15.

Vargas, O. M., E. M. Ortiz, and B. B. Simpson. 2017. Conflicting phylogenomic signals reveal a
pattern of reticulate evolution in a recent high-Andean diversification (Asteraceae:
Astereae: *Diplostephium*). New Phytol. 214:1736–1750.

Vargas, O. M., B. Goldston, D. L. Grossenbacher, and K. M. Kay. 2020. Patterns of speciation
are similar across mountainous and lowland regions for a Neotropical plant radiation
(Costaceae: *Costus*). Evolution 74:2644–2661.

Villagómez, D., R. Spikings, A. Mora, G. Guzmán, G. Ojeda, E. Cortés, and R. Van Der Lelij.
2011. Vertical tectonics at a continental crust-oceanic plateau plate boundary zone:
Fission track thermochronology of the Sierra Nevada de Santa Marta, Colombia.
Tectonics 30: TC4004.

von Humboldt, A., and A. Bonpland. 1805. Essai sur la géographie des plantes: accompagné
d'un tableau physique des régions équinoxiales, fondé sur des mesures exécutées, depuis
le dixième degré de latitude boréale jusqu'au dixième degré de latitude australe, pendant
les années 1799. A Paris:Chez Levrault, Schoell et compagnie, libraires.

Weigend, M. 2002. Observations on the biogeography of the Amotape-Huancabamba zone in
northern Peru. Bot. Rev. 68:38–54.

Wiens, J. J. 2004. Speciation and ecology revisited: phylogenetic niche conservatism and the
origin of species. Evolution 58:193–197.

Zapata, F. 2013. A multilocus phylogenetic analysis of *Escallonia* (Escalloniaceae):
Diversification in montane South America. Am. J. Bot. 100:526–545.