Energy Crisis in Human and Mouse Models of *SLC4A11*-associated Corneal Endothelial Dystrophies

Running Title: Energy Crisis in SLC4A11-deficient Corneal Endothelium

Wenlin Zhang[1], Ricardo Frausto[1], Doug Chung[1], Christopher G. Griffis[1], Liyo Kao[2], Angela Chen[1], Rustam Azimov[2], Alapakkam Sampath[1], Ira Kurtz[2, 3], Anthony Aldave[1]*

[1]Stein Eye Institute, UCLA, Los Angeles, CA 90095

[2]Division of Nephrology, David Geffen School of Medicine at UCLA, Los Angeles, CA 90095

[3]Brain Research Institute, UCLA, Los Angeles, CA 90095

*Corresponding Author: Anthony Aldave, aldave@jsei.ucla.edu

Conflict of Interests: There is no conflict of interests.

Abstract

Mutations in the solute-linked carrier family 4 member 11 (*SLC4A11)* gene are associated with congenital hereditary endothelial dystrophy (CHED), Fuchs endothelial corneal dystrophy and Harboyan syndrome, in all of which visually significant cornea edema may require corneal transplantation. However, the pathogenesis of SLC4A11-associated corneal endothelial dystrophies remains to be elucidated. Recent evidence suggested cellular respiration reprogramming and mitochondrial oxidative stress in SLC4A11-deficient corneal endothelium. Given the complexity of cellular metabolic regulation and its cell type specific impact on cellular physiology, we systemically analyzed the transcriptome of *SLC4A11* knock-down primary human corneal endothelium (*SLC4A11* KD pHCEnC) and corneal endothelial cells derived from *Slc4a11*$^{-/-}$ mice (*Slc4a11*$^{-/-}$ MCEnC) to provide a comprehensive characterization of the transcriptome profile changes resulting from loss of SLC4A11. To identify the conserved molecular mechanisms that lead to cornea endothelial dysfunction in both the human and murine models, we performed comparative transcriptomic analysis. Our analysis identified inhibition of cell metabolism and ion transport function as well as mitochondria dysfunction as shared between *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC. Functional analysis confirmed the absence of SLC4A11-mediated NH_4Cl-induced membrane depolarization in *Slc4a11*$^{-/-}$ MCEnC. Transcriptome of *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC identified downregulation of Na^+-HCO_3^- transporter (NBCe1, *SLC4A4*), a key player in corneal endothelial 'pump' function, and upregulation of Syntaxin 17 (*STX17*), an initiator of mitophagy. NBCe1 and STX17 were further analyzed in *Slc4a11*$^{-/-}$ MCEnC for functional impact and in *SLC4A11* KD pHCEnC and corneal endothelium from individuals with CHED for protein expression, all

showed consistent changes with transcriptome. CHED corneal endothelium also showed decreased immunostaining intensity for mitochondria markers suggesting decreased mitochondira density . In *SLC4A11* KD pHCEnC and *Slc4a11*[-/-] MCEnC, steady state ATP depletion and ATP sensing AMPK-p53 pathway activation were observed, consistent with the prediction using transcriptome data that transcriptional factor p53 were responsible for the transcriptomic changes. These findings suggest that insufficient energy fueling the corneal endothelial 'pump', as a result of metabolic inhibition and failing mitochondria, is the direct cause of clinical phenotype of corneal edema in *SLC4A11*-associated corneal endothelial dystrophies.

INTRODUCTION

Solute-linked carrier family 4 member 11 (SLC4A11) is one of the highly expressed

differentiation markers for corneal endothelium [1-3]. Mutations in *SLC4A11* are

associated with congenital hereditary endothelial dystrophy (CHED), Harboyan

syndrome (CHED with perceptive deafness) and Fuchs endothelial corneal

dystrophy (FECD) [4-7]. Additional evidence of a role for *SLC4A11* in FECD comes

from reports of the hypermethylation of the *SLC4A11* promoter and transcriptional

downregulation of *SLC4A11* in FECD corneal endothelium [8,9]. Children with

CHED often present with bilateral corneal edema at or shortly after birth with

significant vision impairment. Corneal transplantation is the only means of restoring

vision and is associated with a guarded prognosis in terms of long term recovery of

vision and graft survival [10]. In addition, these children are at risk of developing

perceptive deafness later in life (Harboyan Syndrome) [11]. FECD affects as much

as 5% of the U.S. population over 40 years of age [12], and visually significant

corneal edema secondary to FECD is the most common indication for keratoplasty in

US and worldwide [13,14]. Together, CHED and FECD constitute common

indications for corneal transplantation in published series from around the world

[15,16].

SLC4A11 is functionally characterized as an NH_3 and alkaline pH stimulated

H^+ transporter, while permeability to Na^+, OH^- and water has also been reported [17-

21]. SLC4A11 is essential in facilitating energy producing glutaminolysis, maintaining

antioxidant signaling and preventing apoptosis in corneal endothelial cell [22-25].

During development and in the event of oxidative DNA damage, SLC4A11 gene

expression is directly upregulated by activated p53 during development and in the

response to DNA damage [26,27]. In SLC4A11-associated corneal endothelial

dystrophies, the cornea edema that develops as a result of pathologic *SLC4A11*

mutations is evidence of corneal endothelial dysfunction, either from direct cell

loss/death or from disturbances in the corneal endothelial 'pump-leak' system [28-30].

Corneal endothelial 'pump-leak' system achieves corneal transparency by

maintaining the dynamic balance between the passive fluid leak from anterior

chamber into corneal stroma and the active pump moving fluid from corneal stroma

to anterior chamber. The fluid leak in corneal endothelium is particularly significant

because hydrophilic glycosaminoglycans in the stroma are only 13~23% hydrated in

physiological condition, producing a significant swelling pressure [31-33]. In addition,

corneal endothelial tight junctions are focally incomplete, leading to a

nonhomogeneous seal of the lateral intercellular space and offering little resistance

to the paracellular passage of water and solutes [34-36]. The fluid pump activity is

associated with a host of ion channels/exchangers/pumps positioned strategically at

the apical and basolateral membranes of the corneal endothelial cells [29,37] and

driven by an ionic electrical-chemical gradient set up by the highly expressed Na^+/K^+-

ATPase [38]. As such, corneal endothelium has the second highest density of

mitochondria among any cell types in the body (second to photoreceptors) to

generate sufficient ATP to fuel the Na^+/K^+-ATPase driven endothelial 'pump'. As

mentioned above, SLC4A11 plays a significant role in facilitating ATP-generating

glutaminolysis in corneal endothelium [24]. Thus, it is not surprising that

glutaminolysis inhibition, mitochondria membrane potential (MMP) depolarization,

enriched mitochondrial reactive oxidative species (ROS) and increased mitochondria

turnover have been observed in the corneal endothelium of the $Slc4a11^{-/-}$ mouse

[24,25,39], which recapitulates the CHED corneal phenotype of significant cornea

edema, Descemet membrane thickening and progressive corneal endothelial cell

loss [40]. Thus, the association between SLC4A11 and corneal endothelial

mitochondrial function suggests SLC4A11 is involved not only in moving ions across

the plasma membrane but also in the supply of energy to the endothelial 'pump'.

Approximately 94 mutations, including nonsense, missense, frameshift and

intronic mutations, have been identified in *SLC4A11* in individuals with CHED

[4,6,11,41-58]. Coding region mutations have been identified in 17 of the 19 exons of

SLC4A11, with no spatial clustering within ~~to any~~ ~~of the xx functional domains of~~

SLC4A11 protein. While a large number of these mutations result in SLC4A11

protein misfolding and failure to mature to the plasma membrane [5,6,59-61], some

mutations affect SLC4A11 transporter function without impacting membrane

trafficking [19,62,63] or cause aberrant *SLC4A11* pre-mRNA splicing and

subsequent reduced SLC4A11 expression [57]. Collectively, these observations

suggest that loss of SLC4A11 function is the primary pathogenetic mechanism in

CHED rather than mutant SLC4A11 protein misfolding/mislocalization in ER and

secondary unfolded protein response. Therefore we investigated the impact of

SLC4A11 reduction on the corneal endothelial transcriptome in human and murine

corneal endothelial cells, with validation in corneal endothelium from individuals with

CHED, and elucidated the upstream molecular mechanism leading to the observed

transcriptomic changes.

MATERIALS AND METHODS

Isolation and Culturing of Primary Human Corneal Endothelial Cells

Primary cultures of human corneal endothelial cells (pHCEnC) were isolated from

donor corneas as previously described [64]. Cells were plated in laminin coated cell

culture plastic and cultured in a 1:1 mixture of F12-Ham's and M199 (F99) medium.

Experiments were performed when the cells achieved an intact and confluent

monolayer.

Knock-down of SLC4A11 in pHCEnC

Confluent pHCEnC were transfected with 10nM anti-SLC4A11 siRNA (siRNA-

CrCrGrArArArGrUrArCrCrUrGrArArGrUrUrArArArGrArACT) or scrambled siRNA

(OriGene Technologies) using Lipofectamine LTX (Life Technologies). At 72 hrs

post-transfection, the cells were lysed in either Tri Reagent (Sigma-Aldrich) or

radioimmunoprecipitation assay (RIPA) buffer for RNA and protein isolation,

respectively.

Immortalized Mouse Corneal Endothelial Cell (MCEnC) Culture

Immortalized *Slc4a11*$^{+/+}$ and *Slc4a11*$^{-/-}$ MCEnC lines were derived from *Slc4a11*$^{+/+}$

and *Slc4a11*$^{-/-}$ mice corneal endothelium respectively as previously described [25].

All MCEnC lines were cultured on flasks coated with a mixture of 40 μg/cm^2

chondroitin sulfate (Sigma-Aldrich) and 40 ng/ cm^2 laminin (Sigma-Aldrich). Details of

culture medium used can be found in supplemental material.

Total RNA Isolation from pHCEnC and MCEnC

Total RNA from cultured pHCEnC was isolated in TRI Reagent and purified with the

RNeasy Clean-Up Kit (Qiagen). Total RNA from cultured MCEnC was isolated and

purified using the RNeasy Plus Mini Kit (Qiagen). The integrity of the purified RNA

was analyzed by the Agilent 2100 Electrophoresis Bioanalyzer System (Agilent

Technologies).

RNA Sequencing (RNA-Seq) of total RNA from pHCEnC and MCEnC

Purified total RNA from pHCEnC was prepared for RNA-seq libraries using KAPA

mRNA HyperPrep Kit. Libraries were sequenced on the Illumina Hi-Seq 4000 and

paired-end 150-bp reads were generated. Purified total RNA from MCEnC was

submitted to the UCLA Technology Center for Genomics & Bioinformatics for library

preparation and sequencing. Single-end 50-bp reads were generated using the

Illumina Hi-seq 3000. The generated FASTQ files and quantitative results are

available from the GEO DataSets database (accession number XXXXX and XXXXX;

NCBI).

RNA-sequencing Data Analyses

The FASTQ files containing the raw reads from the pHCEnC and MCEnC were

aligned to the human (GRCh38/hg38) and mouse (GRCm38/mm10) genomes

respectively using Hisat2. The raw counts of aligned reads were converted to counts

per million (CPM) mapped reads with consideration of library size and normalized by

the method of trimmed mean of M values (TMM). Then linear models for microarray

analysis (LIMMA) coupled with variance modeling at the observation-level (VOOM)

were used for differential gene expression analysis in R software. The following

thresholds defined differential expression: CPM > 1, fold change (fc) > 1, and

adjusted p-value < 0.05. Differential gene expression (DGE) lists were obtained from

the following sample sets respectively: pHCEnC sample set: passage 1 of pHCEnC

with siRNA targeting *SLC4A11* versus scrambled RNA control (*SLC4A11* KD

pHCEnC and scRNA pHCEnC; n=3 each); MCEnC early sample set: early passages

of *Slc4a11*$^{-/-}$ MCEnC versus wild-type control (p6 *Slc4a11*$^{-/-}$ and p7 *Slc4a11*$^{+/+}$

MCEnC; n=4 each); and MCEnC late sample set: late passages of $Slc4a11^{-/-}$

MCEnC versus wild-type control (p39 $Slc4a11^{-/-}$ and p40 $Slc4a11^{+/+}$ MCEnC; n=4

each).

Ingenuity Pathway Analysis

Ingenuity Pathway Analysis (IPA®, QIAGEN Bioinformatics) was used to perform

comparative transcriptome analysis among three sample sets (pHCEnC, MCEnC

early and MCEnC late), canonical/biological function pathway enrichment and

upstream regulator prediction analyses on each sample set. In the comparative

transcriptome analysis, canonical and biological function pathways with a predicted

activation z-score in each of the three sample sets were sorted by the sign and value

of the z-score to identify the most enriched pathways. Enriched canonical and

biological function pathways without an assigned z-score and predicted upstream

regulators were ranked by mean enrichment p-values among sample sets.

Quantitative PCR

Quantitative PCR was performed to validate the differential expression of selected

genes from transcriptome DGE list on separate batches of RNA samples from

SLC4A11 KD pHCEnC and $Slc4a11^{-/-}$ MCEnC. Total RNA of 1 μg from each sample

was reversed transcribed using SuperScript III First-Strand synthesis kit (Sigma).

Quantitative PCR was performed on the LightCycler 480 System (Roche) using the

KAPA SYBR FAST Universal Kit (Kapa Biosystems) with an annealing temperature

of 60 °C. Primers used are listed in Supplemental Table 1. Relative gene expression

was obtained by comparison to the housekeeping gene *PPIA/Ppia or ACTB/Actb*

and was calculated by the comparative Ct ($2^{-\Delta Ct}$) method.

Immunofluorescence

Five-micrometer sections of paraffin-embedded corneas from healthy donors and

individuals with CHED were de-paraffinized and re-hydrated in a graded ethanol

series (100%, 95% and 70% and 50%) for 5 min and subject to antigen retrieval in

10 mM Na-citrate. After blocking with 2% bovine serum albumin (BSA) in PBS,

sections were incubated overnight at 4 °C with primary antibodies (listed in

Supplemental Table 1) in 2% BSA. Slides were then incubated with 1:200 Alexa

Fluor® 488 F(ab')2-Goat anti-Rabbit IgG (Invitrogen) and 1:200 Alexa Fluor® 568

F(ab')2-Goat Anti-Mouse IgG (Invitrogen) with DAPI in PBS with 2% BSA for one

hour. Sections were mounted with prolong anti-fade mounting reagent (Invitrogen)

and imaged with Olympus FV-1000 inverted confocal fluorescence microscope.

Florescence intensity of the images were quantified using FLUOVIEW 4.2 and

ImageJ software.

Single Cell Patch-clamp Recording

Single cell recordings were made from MCEnC cultured on 25-mm glass coverslips

to ~10–20% confluence to isolate single adhered cells. Membrane voltages were

measured using whole-cell patch electrodes in current clamp mode ($I_{hold} = 0$) as

previously described [65]. Ammonium evoked responses were sampled at 1kHz and

digitally low-pass filtered at 50Hz using a 7-pole Butterworth filter. Data were

reported as mean (95%CI). Details of data acquisition and statistical analysis can be

found in supplemental material.

Intracellular pH Measurement

Intracellular pH (pH_i) measurements were performed by monitoring intracellular free

H^+ concentration using pH-sensitive fluorescent dye BCECF-AM (2'7'-

bis(carboxyethyl)-5(6)-carboxyfluorescein-acetoxymethyl ester, B1170; Thermo

Fisher Scientific) as described previously [21,22]. Details of experimental settings

can be found in supplemental material.

Western Blotting

Whole-cell lysates from pHCEnC and MCEnC were prepared with RIPA buffer with

proteinase and phosphatase inhibitors. Total protein was quantified by bicinchoninic

acid (BCA) assay, separated and detected on Simple Western system Wes[TM]

(ProteinSimple). Quantification and data analysis were performed in Compass for

SW software (ProteinSimple). Antibodies used are listed in Supplemental Table 1.

Intracellular ATP Assay

MCEnC and pHCEnC were seeded on 1×10^5 /mL in 12- and 24-well plates,

respectively, and cultured to sub-confluence. ATP was extracted by a boiling water

method [66], and measured using a luciferin-luciferase based ATP assay kit

(Molecular Probes) following the manufacturer's instructions.

Statistical Analysis

Statistical analysis was performed in GraphPad Prism 7.0 software, unless otherwise

indicated, with appropriate statistical tests based on the data structure. Specific

statistical tests used for each comparison are indicated in figure legends. Data are

presented as mean ± SEM unless otherwise indicated. Statistical significance is

denoted as follows in the figures: $p < 0.05$, *; $p < 0.01$, **; $p<0.001$, ***; $p<0.0001$,

****.

RESULTS

SLC4A11 Mutations Do Not Result in Decreased Mutant SLC4A11 Expression

Corneal specimens two individuals with CHED were examined with light microscopy

and immunocytochemistry. One of the two individuals (*SLC4A11*^Mu/Mu) demonstrated a

compound heterozygous mutations in *SLC4A11* (NM_001174090.1)

c.[473_480delGCTTCGCCinsC];[2623C>T], p.[(R158PfsX3)];[(Arg875*)] and the

other (*SLC4A11*^Mu/WT) demonstrated a single heterozygous *SLC4A11* coding region

mutation [c.2146C>G (p.Pro716Ala)]. Both corneal specimens demonstrated a

significantly thickened Descemet's membrane and an attenuated corneal endothelial

layer with cytoplasmic inclusions present in some cells (Fig. 1A).

Immunofluorescence staining for SLC4A11 protein in the corneal endothelium was

performed in conjunction with the use of an antibody against an endoplasmic

reticulum marker, protein disulfide isomerase (PDI), to investigate whether *SLC4A11*

mutations lead to disease via protein misfolding and retention in the endoplasmic

reticulum (ER), as previously reported [5,6,60]. There was no difference in the

corneal endothelial SLC4A11 protein staining intensity between the two specimens

from the individuals with CHED and seven healthy controls. In addition, the staining

localized to the cellular membrane and did not co-localize with PDI (Fig. 1B). Thus,

in the corneal endothelium of these two individuals with CHED, mutant SLC4A11

expression is not reduced compared to wild type SLC4A11 expression in controls,

and remains localized to the cellular membrane instead of being retained in the ER.

SLC4A11/Slc4a11 Reduction Induces Corneal Endothelial Transcriptome Changes

Next, we performed transcriptome analysis in pHCEnC and MCEnC with reduced

SLC4A11/Slc4a11, mimicking the loss of SLC4A11 function in CHED corneal

endothelium. Principle component analysis (PCA) was performed to ensure tight

clustering within biological triplicates, and to ensure that the number of dimensions

considered are sufficient in explaining the variances between samples of different

genotypes (Supplemental Figure 1). Based on the shared corneal endothelial

phenotypes between human CHED and the *Slc4a11*[-/-] mouse, we compared the

transcriptomes from *SLC4A11* KD pHCEnC and early and late passage MCEnC

derived from *Slc4a11*[-/-] mice (denoted as pHCEnC, MCEnC early and MCEnC late

sample set, respectively) [25]. A comparison of the genes identified from each

sample set that were differentially expressed in comparison to appropriate controls

revealed 3171 genes that were consensually differentially expressed across three

sample sets (Fig.1C), of which 1041 genes were differentially expressed in the same

direction (Fig.1D). Over a third of the 30 most highly differentially expressed genes

(Fig.1E) have been previously demonstrated to play important functional roles in the

cornea and/or have been associated with corneal diseases, including *SEMA3E* [1],

MST1R [67], *RNF43* [68], SLC2A3 [69], *WNK4* [70], SLC4A4 [71], *CYGB* [72],

SLC9A7 [1,73], *ZNF469* [74-77], *BNC1* [78] and *TTC22* [79].

Loss of SLC4A11 Leads to Generalized Inhibition of Cellular Metabolism

Comparison of enriched canonical pathway from *Slc4a11* KD pHCEnC, *Slc4a11*[-/-]

MCEnC early and *Slc4a11*[-/-] MCEnC late sample sets identified a shared generalized

inhibition (defined by negative activation z-score) of multiple metabolic pathways that

were interconnected via intermediate metabolites (Fig.2A, B). A generalized

decrease in the expression of enzyme-encoding genes in these pathways was

observed (Fig.2C-I), a finding that was confirmed for selected genes from each

pathway using qPCR (Fig.2J).

Altered Expression of Channels and Transporters Impairs Transport Function of

Corneal Endothelium

IPA biological function enrichment analysis identified "transport of molecule" as

the top inhibited function (negative z-score) shared between the transcriptomes of

the pHCEnC, MCEnC early and late samples sets (Fig. 3A). Since SLC4A11 is an

electrogenic $NH_3:H^+$ co-transporter [17], we performed single cell recording of

membrane potential of *Slc4a11*$^{+/+}$ and *Slc4a11*$^{-/-}$ MCEnC in response to

extracellularly perfused 10 mM NH_4Cl . While exposure to NH_4Cl induced a +12.7

mV (95% CI +5 mV, +20 mV) depolarization in *Slc4a11*$^{+/+}$ MCEnC, likely due to the

$NH_3:H^+$ permeability provided by Slc4a11, exposure to NH_4Cl induced a -9.50 mV

(95% CI -19.3 mV, -0.5 mV) hyperpolarization in *Slc4a11*$^{-/-}$ MCEnC (Fig. 3B, C).

To further understand the nature of this depolarization to hyperpolarization shift

resulting from loss of Slc4a11, we examined the list of 1041 genes differentially

expressed in the same direction in pHCEnC, MCEnC early and MCEnC late sample

sets to identify channels and transporters with a $log_2FC > 1$ (Fig. 3D). Twelve genes

encoding ion channels were identified (Fig. 3D) including: Cl^- channels (*CLCN3*,

CLCN2, *ANO5*) and Ca^{2+} channels (*TRPC1*, *TRPV4*, *CACNB1*, *CACNB4*, *TRPV2*),

which were both up- and down-regulated; and K^+ channels (*KCNAB2*, *KCNMA1*),

which were exclusively down-regulated. Twenty six genes encoding transporters

were identified (Fig. 3D), 12 of which were upregulated and 14 of which were down

regulated. including: Na^+-HCO_3^- co-transporter (*SLC4A4*), Ca^{2+}-ATPase (*ATP2B1*),

glucose transporter (*SLC2A1, SLC2A13, SLC2A3, SLC2A12, SLC15A2*), glutamine

transporter(*SLC38A1*), Cl^-/HCO_3^- exchanger (*SLC4A3*), ATP-binding cassette (ABC)

transporters (*ABCA8, ABCC4*), K^+-Cl^- cotransporter (*SLC12A7*), Na^+/H^+ exchanger

regulator (*SLC9A3R2*), Na^+-HPO_3^- co-transporter (*SLC20A1*), lactate transporter

(*SLC16A4*), Na^+-carnitine co-transporter (SLC22A4, *SLC22A18*), SO_4^{2-} transporter

(*SLC26A2*). In particular, electrogenic Na^+-HCO_3^- co-transporter (NBCe1, *SLC4A4*),

which plays an essential role in the corneal endothelial 'pump' function [29], was

down-regulated 4 fold in *SLC4A11* KD pHCEnC and > 100 fold in *Slc4a11*[-/-] MCEnC

(early and late passage) in transcriptomes of three sample sets (Fig.3D). The

downregulation of SLC4A4/Slc4a4 was verified by qPCR in *SLC4A11* KD pHCEnC

and *Slc4a11*[-/-] MCEnC (Fig. 3E) and by Western blot in *SLC4A11* KD pHCEnC (Fig.

3F). Immunofluorescence staining for NBCe1 (SLC4A4) in the corneal endothelium

of two individuals with CHED showed decreased expression of NBCe1 compared to

control cornea endothelium (Fig. 3G, H).

Next, direct functional measurement of Na^+ dependent HCO_3^- transport in

Slc4a11[-/-] MCEnC showed reduced Na^+-HCO_3^- co-transporter activity when

compared to *Slc4a11*[+/+] MCEnC (Fig. 3I). In Fig.3I, intracellar pH (pH_i) was

maintained at a low value when *Slc4a11*[-/-] and *Slc4a11*[+/+] MCEnC were perfused

with Na^+-free bicarbonate containing ([HCO_3^-] 28.3 mM) solution. When switched to

a Na^+ containing bicarbonate solution, pH_i increased as the Na^+-HCO_3^- cotransporter

(NBCe1) started to move the weak base HCO_3^- inward using Na^+ inward

transmembrane electrochemical gradient. We determined the initial slope of this

pH_i rise (pH_i recovery) to serve as an indirect measure of Na^+-HCO_3^- cotransport

activity, and observed reduced Na^+ dependent pH_i recovery in *Slc4a11*[-/-] MCEC

than *Slc4A11*[+/+] MCEnC (Fig. 3I, J).

Mitochondria Dysfunction Leads to Reduced ATP Production

Dilated mitochondria is a characteristic electron microscopy finding in CHED corneal

endothelium, suggestive of mitochondrial involvement in the pathogenesis [80]. We

identified "Mitochondria Dysfunction" is shared between pHCEnC and MCEnC early

and late sample sets as the top enriched IPA canonical pathway (ranked by p-value,

Fig. 4A). Correspondingly, in the list of 1041 genes differentially expressed in the

same direction in pHCEnC, MCEnC early and MCEnC late sample sets, genes

encoding proteins involved in the mitochondria electron transport chain, mediating

mitochondrial ATP flux, import machinery and translation machinery showed a

generalized decreased expression (Fig. 4B, C). The transcriptomes also suggested

'autophagy' was activated (Fig.3A) in *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC.

Together with the indication of "mitochondria dysfunction", we specifically examined

the differential expression of the *STX17*/*Stx17* gene, which encodes SNARE protein

Syntaxin 17 (STX17) and is an mitophagy initiator facilitating the removal of

dysfunctional mitochondria [81]. *STX17/Stx17* was upregulated in the *SLC4A11* KD

pHCEnC and *Slc4a11*$^{-/-}$ MCEnC transcriptomes (Fig. 4D), a finding that was

validated by qPCR in *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC (Fig. 4E), by

Western blot in *SLC4A11* KD pHCEnC (Fig. 4F) as well as by immunofluorescence,

which demonstrated increased STX17 expression in CHED corneal endothelium

compared with control (Fig. 4G). As STX17 can only mediate mitophagy where

mitochondria are present, we co-stained for mitochondrial marker COX4 in CHED

corneal endothelium and control (Fig. 4G), which demonstrated significantly

decreased COX4 indicating reduced mitochondria density in CHED corneal

endothelium (Fig.4G). Quantification of the relative STX17 abundance in reference to

mitochondria density (STX17/COX4 ratio) showed increased STX17 level in CHED

corneal endothelium (Fig. 4H). Quantification of COX4 staining intensity showed

significantly reduced COX4 signal in CHED corneal endothelium than in healthy

controls (Fig4 I). Staining CHED endothelium with another mitochondria marker,

cytochrome C, also showed consistently decreased staining intensity (Fig. 4J, K).

Given the evidence indicating mitochondrial dysfunction and reduced

mitochondria density in the setting of reduced SLC4A11 expression, we

hypothesized that this would result in insufficient ATP energy supply to maintain the

Na^+-K^+-ATPase driven corneal endothelial 'pump', and the cornea edema that

characterizes CHED. Thus, we performed a direct measurement of the steady state

ATP levels in *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC, which revealed

reduced ATP concentrations compared to control scRNA pHCEnC and *Slc4a11*$^{+/+}$

MCEnC, respectively (Fig. 4L, M).

Transcriptomic Changes Associated with Loss of SLC4A11 Are Mediated by

Activation of AMPK-p53 Pathway

To identify the upstream signaling pathway leading to the observed transcriptomic

changes in *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC, we performed upstream

regulator prediction in IPA, which identified P53 (encoded by *TP53* gene) as the top

candidate transcription factor (Fig. 5A). Western blot analysis in *SLC4A11* KD

pHCEnC demonstrated increased Ser15 phosphorylation of p53 and an increased

phosphorylated p53 / total p53 ratio compared to scRNA pHCEnC control (Fig. 5B),

indicative of activation of p53 transcriptional activity. Similarly, in *Slc4a11*$^{-/-}$ MCEnC,

we observed increased Ser18 phosphorylation of p53 (corresponding to Ser15

of human p53) in *Slc4a11*$^{-/-}$ MCEnC late passage, although this was not observed in

early passage (Fig.5C). However, there was increased total p53 level in both

Slc4a11$^{-/-}$ MCEnC early and late passages (Fig.5C, D), indicative of transcriptional

activation of p53.

To identify the kinase responsible for the Ser15 (Ser18 in mouse)

phosphorylation and transcriptional activation of p53, we investigated the role of the

cellular ATP sensor AMP-activated protein kinase (AMPK), given the observed ATP

depletion in *SLC4A11* KD pHEnC and *Slc4a11*$^{-/-}$ MCEnC as well as the reported

capacity of AMPK to mediate Ser15 (Ser18 in mouse) phosphorylation and

transcriptional activation of p53 [82-84]. In the setting of a decreased ATP-to-ADP

(or ATP-to-AMP) ratio, AMPK catalytic α subunit will be phosphorylated at Thr172

whereas phosphorylation of regulatory β1 subunit at Ser182 is not dependent upon

cellular ATP level [85]. In *SLC4A11* KD pHCEnC and *Slc4a11*$^{-/-}$ MCEnC, we

observed an increased ratio of Thr172 phosphorylated AMPKα / total AMPKα, and

no change in the ratio of Ser182 phosphorylated AMPKβ1 / total AMPKβ1 (Fig. 5C).

DISCUSSION

In this manuscript, we utilized human and mouse corneal endothelial cell lines with

reduced expression of SLC4A11/Slc4a11 to investigate the cause of corneal

endothelial dysfunction that characterizes each of the *SLC4A11*-associated corneal

endothelial dystrophies (CHED, FECD and Harboyan syndrome. While several

previous reports have elucidated possible roles for mitochondrial uncoupling, ER

unfolded protein response, oxidative stress and apoptosis in SLC4A11-deficient cell

lines [11,22,23,39,60,86], only one report examined CHED patient corneal

endothelium and demonstrated increased oxidative stress [22]. We identified ATP

depletion in corneal endothelial lacking SLC4A11/Slc4a11, in both transient (72

hours) *SLC4A11* knock-down pHCEnC and permanent *Slc4a11* knockout MCEnC

through prolonged passage (early vs late passage). The reduced corneal endothelial

ATP levels provide a proposed pathogenesis for the corneal endothelial dysfunction,

which clinically manifested as corneal edema. The fact that ATP depletion and ATP-

sensor AMPKα activation were detected within 72 hours of *SLC4A11* knock-down in

pHCEnC suggests this energy shortage from the loss of SLC4A11 function is likely

the initial step in the pathogenesis of *SLC4A11*-associated corneal endothelial

dystrophies that leads to the downstream cascade of oxidative stress and apoptotic

cell loss.

362   When examining the expression and localization of SLC4A11 protein in CHED

corneal endothelium, we provide the initial evidence that mutant SLC4A11 protein is

not retained in the ER of the corneal endothelium. While the prevailing hypothesis is

that majority of *SLC4A11* mutations result in protein misfolding and retention in the

endoplasmic reticulum [5,6,60,61], supporting evidence is from overexpression of

SLC4A11 mutants in a HEK293 cell line model. Mutant SLC4A11 localization was

neither investigated in corneal endothelial cells overexpressing SLC4A11 mutants

nor in corneal endothelium from individuals with CHED. In addition, our

immunostaining of CHED corneal endothelium did not show reduced expression

level of the SLC4A11 mutant protein, consistent with the previous report that CHED

corneal endothelium does not demonstrate reduced SLC4A11 expression at the

mRNA level [22]. Instead, our data supports the alternative hypothesis that

identified *SLC4A11* mutations affect the protein transport function of the SLC4A11

protein in the corneal endothelium [19,62,63].

We used a comparative transcriptomics approach based on the observation of

phenotypic similarities between CHED and the *Slc4a11*$^{-/-}$ mouse corneal phenotype

[40,87]. This approach is based on the high degree of gene orthology between

mouse and human and the organ-dominated hierarchical clustering observed across

mammals on real genes expression data [88]. Comparative transcriptomic analysis

of human and mouse sample sets together is advantageous over independent

transcriptomic analysis of each sample set as it enables us to identify transcriptome

determinants attributable to the loss of SLC4A11/Slc4a11 from that resulting from

other biological factors or technical effects such as primary cell passage numbers,

cell line immortalization, siRNA treatment, differences in culture medium and etc [89].

With this approach, we identified generalized inhibition of multiple metabolic

pathways, as well as mitochondria dysfunction in both *SLC4A11* KD pHCEnC and

Slc4a11$^{-/-}$ MCEnC, mediated via the activation of the AMPK-p53 pathway. The data

presented here support the use of the *Slc4a11*$^{-/-}$ mice as a model for the *SLC4A11*

associated corneal endothelial dystrophies and indicate a favorable translational

potential for therapeutic approaches shown to be efficacious in the *Slc4a11*$^{-/-}$ mice.

When investigating the impact of reduced SLC4A11 on molecule transport

function in corneal endothelial cells, we demonstrated opposite NH_4Cl induced

membrane potential change in $Slc4a11^{-/-}$ versus $Slc4a11^{+/+}$ MCEnC. Membrane

depolarization in reaction to extracellular NH_4Cl exposure were commonly observed

in membranes with permeability to NH_4^+/NH_3 (membranes expressing ammonia or

ammonium transporters/channels such as AQP8, AQP4, AMT1 and Rh-associated

glycoproteins) [90,91]. In $Slc4a11^{+/+}$ MCEnC, extracellular NH_4Cl perfusion induced

similar membrane depolarization. However, in $Slc4a11^{-/-}$ MCEnC, hyperpolarization

was observed. Transcriptome data showed differential expression of 38 ion channels

and transporters, suggesting SLC4A11 not only facilitates NH_4^+/NH_3 transmembrane

movement but may also functions in maintaining the homeostasis of the endothelial

transmembrane ion gradient by regulating the expression and function of other

channels/transporters. This hypothesis is supported by the observation of reduced

expression and activity of the $Na^+-HCO_3^-$ co-transporter (NBCe1, $SLC4A4$) in

$Slc4a11^{-/-}$ MCEnC. Future investigation analyzing the activity of other differentially

expressed ion channels and transporters is needed to facilitate our detailed

understanding of the impact of SLC4A11 reduction on corneal endothelial "pump

function".

Our attempt to identify upstream determinants of the observed transcriptomic

changes revealed activation of AMPK-p53 pathway in $SLC4A11$ KD pHCEnC and

$Slc4a11^{-/-}$ MCEnC. Our data suggests that AMPK activation is likely to be the initial

cellular event in response to SLC4A11 deficiency induced ATP depletion. While

there was consistent AMPK activation (increased Thr172 phosphorylation) in both

acute SLC4A11 deficiency in pHCEnC and chronic SLC4A11 deficiency in MCEnC,

and increased p53 phosphorylation at Ser15 (or Ser18 in mouse) was only observed

in *SLC4A11* KD pHCEnC and *Slc4a11*[-/-] MCEnC late passage, but not in *Slc4a11*[-/-]

MCEnC early passage. However, total p53 level was elevated in both early and late

passages of MCEnC with permeant knockout of *Slc4a11*. Although p53 is generally

stablely expressed at the protein level, recent evidence suggests activation of

AMPKα can induce the transcriptional activation of *p53* [83]. In the context of our

data, it suggests that short-term AMPKα activation in SLC4A11 deficiency induced

post-translational activation of p53, whereas prolonged AMPKα activation in

SLC4A11 deficiency induced transcriptional activation of p53.

ACKNOWLEDGEMENTS

We genuinely thank Dr. Huong Duong in Ho Chi Minh City Eye Hospital for

facilitating the acquisition of postsurgical corneal endothelium specimens from

individuals with CHED.

Support is provided by National Eye Institute Grants 1R01EY022082 (A.J.A.),

P30EY000331 (core grant), the Walton Li Chair in Cornea and Uveitis (A.J.A.), the

Stotter Revocable Trust (SEI Cornea Division), an unrestricted grant from Research

to Prevent Blindness (A.J.A.), National Institute of Diabetes and Digestive and

Kidney Diseases 1R01DK077162 (I.K.), the Allan Smidt Charitable Fund (I.K.), Ralph

Block Family Foundation (I.K.) and Knights Templar Eye Foundation Career Starter

Grant (WZ).

STATEMENT OF AUTHOR CONTRIBUTIONS

W.Z., R.F., C.G.G., L.K.and A.J.A. conceived the experiments, W.Z. and A.J.A.

wrote the manuscript, W.Z., R.F., D.C, C.G.G., L.K., A.C. and R.A. performed the

experiments. W.Z. A.S., I.K., and A.J.A. secured funding. A.S. and I.K. provided

expertise and feedback.

REFERENCE LIST

1.   Frausto RF, Wang C, Aldave AJ. Transcriptome Analysis of the Human Corneal Endothelium.
445   *Investigative Ophthalmology & Visual Science* 2014; **55**: 7821-7830.
446   2.   Chng Z, Peh GS, Herath WB, *et al.* High throughput gene expression analysis identifies
447   reliable expression markers of human corneal endothelial cells. *PLoS One* 2013; **8**: e67546.
448   3.   Chen Y, Huang K, Nakatsu MN, *et al.* Identification of novel molecular markers through
449   transcriptomic analysis in human fetal and adult corneal endothelial cells. *Human Molecular*
450   *Genetics* 2012; **22**: 1271-1279.
451   4.   Aldahmesh MA, Khan AO, Meyer BF, *et al.* Mutational spectrum of SLC4A11 in autosomal
452   recessive CHED in Saudi Arabia. *Invest Ophthalmol Vis Sci* 2009; **50**: 4142-4145.
453   5.   Vithana EN, Morgan PE, Ramprasad V, *et al.* SLC4A11 mutations in Fuchs endothelial corneal
454   dystrophy. *Hum Mol Genet* 2008; **17**: 656-666.
455   6.   Vithana EN, Morgan P, Sundaresan P, *et al.* Mutations in sodium-borate cotransporter
456   SLC4A11 cause recessive congenital hereditary endothelial dystrophy (CHED2). *Nat Genet*
457   2006; **38**: 755-757.
458   7.   Desir J, Moya G, Reish O, *et al.* Borate transporter SLC4A11 mutations cause both Harboyan
459   syndrome and non-syndromic corneal endothelial dystrophy. *J Med Genet* 2007; **44**: 322-326.
460   8.   Khuc E, Bainer R, Wolf M, *et al.* Comprehensive characterization of DNA methylation
461   changes in Fuchs endothelial corneal dystrophy. *PLoS One* 2017; **12**: e0175112.
462   9.   Gottsch JD, Bowers AL, Margulies EH, *et al.* Serial Analysis of Gene Expression in the Corneal
463   Endothelium of Fuchs' Dystrophy. *Investigative Ophthalmology & Visual Science* 2003; **44**:
464   594-599.
465   10.   Dana M-R, Moyes AL, Gomes JAP, *et al.* The Indications for and Outcome in Pediatric
466   Keratoplasty: A Multicenter Study. *Ophthalmology* 1995; **102**: 1129-1138.
467   11.   Siddiqui S, Zenteno JC, Rice A, *et al.* Congenital hereditary endothelial dystrophy caused by
468   SLC4A11 mutations progresses to Harboyan syndrome. *Cornea* 2014; **33**: 247-251.
469   12.   Lorenzetti DW, Uotila MH, Parikh N, *et al.* Central cornea guttata. Incidence in the general
470   population. *Am J Ophthalmol* 1967; **64**: 1155-1158.
471   13.   2018 Eye Banking Statistical Report. *Eye Bank Association of America* 2019.
472   14.   Gain P, Jullienne R, He Z, *et al.* Global Survey of Corneal Transplantation and Eye Banking.
473   *JAMA Ophthalmology* 2016; **134**: 167-173.
474   15.   Dorrepaal SJ, Cao KY, Slomovic AR. Indications for penetrating keratoplasty in a tertiary
475   referral centre in Canada, 1996-2004. *Can J Ophthalmol* 2007; **42**: 244-250.
476   16.   Ghosheh FR, Cremona FA, Rapuano CJ, *et al.* Trends in penetrating keratoplasty in the
477   United States 1980-2005. *Int Ophthalmol* 2008; **28**: 147-153.
478   17.   Zhang W, Ogando DG, Bonanno JA, *et al.* Human SLC4A11 Is a Novel NH3/H+ Co-transporter.
479   *J Biol Chem* 2015; **290**: 16894-16905.
480   18.   Myers EJ, Marshall A, Jennings ML, *et al.* Mouse Slc4a11 expressed in Xenopus oocytes is an
481   ideally selective H+/OH- conductance pathway that is stimulated by rises in intracellular and
482   extracellular pH. *Am J Physiol Cell Physiol* 2016; **311**: C945-C959.
483   19.   Vilas GL, Loganathan SK, Liu J, *et al.* Transmembrane water-flux through SLC4A11: a route
484   defective in genetic corneal diseases. *Hum Mol Genet* 2013; **22**: 4579-4590.
485   20.   Ogando DG, Jalimarada SS, Zhang W, *et al.* SLC4A11 is an EIPA-sensitive Na(+) permeable pHi
486   regulator. *Am J Physiol Cell Physiol* 2013; **305**: C716-727.
487   21.   Kao L, Azimov R, Shao XM, *et al.* Multifunctional ion transport properties of human SLC4A11:
488   comparison of the SLC4A11-B and SLC4A11-C variants. *Am J Physiol Cell Physiol* 2016; **311**:
489   C820-C830.

22.   Guha S, Chaurasia S, Ramachandran C, *et al.* SLC4A11 depletion impairs NRF2 mediated
antioxidant signaling and increases reactive oxygen species in human corneal endothelial
cells during oxidative stress. *Sci Rep* 2017; **7**: 4074.
23.   Liu J, Seet LF, Koh LW, *et al.* Depletion of SLC4A11 causes cell death by apoptosis in an
immortalized human corneal endothelial cell line. *Invest Ophthalmol Vis Sci* 2012; **53**: 3270-
3279.
24.   Zhang W, Li H, Ogando DG, *et al.* Glutaminolysis is Essential for Energy Production and Ion
Transport in Human Corneal Endothelium. *EBioMedicine* 2017; **16**: 292-301.
25.   Zhang W, Ogando DG, Kim ET, *et al.* Conditionally Immortal Slc4a11-/- Mouse Corneal
Endothelial Cell Line Recapitulates Disrupted Glutaminolysis Seen in Slc4a11-/- Mouse Model.
Invest Ophthalmol Vis Sci 2017; **58**: 3723-3731.
26.   Wang Q, Zou Y, Nowotschin S, *et al.* The p53 Family Coordinates Wnt and Nodal Inputs in
Mesendodermal Differentiation of Embryonic Stem Cells. *Cell Stem Cell* 2017; **20**: 70-86.
27.   Younger ST, Kenzelmann-Broz D, Jung H, *et al.* Integrative genomic analysis reveals
widespread enhancer regulation by p53 in response to DNA damage. *Nucleic Acids Res* 2015;
43: 4447-4462.
28.   Edelhauser HF. The balance between corneal transparency and edema: the Proctor Lecture.
Invest Ophthalmol Vis Sci 2006; **47**: 1754-1767.
29.   Bonanno JA. Molecular mechanisms underlying the corneal endothelial pump. *Exp Eye Res*
2012; **95**: 2-7.
30.   Srinivas SP. Dynamic regulation of barrier integrity of the corneal endothelium. *Optom Vis*
Sci 2010; **87**: E239-254.
31.   Hedblom EE. The role of polysaccharides in corneal swelling. *Exp Eye Res* 1961; **1**: 81-91.
32.   Hedbys BO, Mishima S, Maurice DM. The Imbibition Pressure of the Corneal Stroma. *Exp Eye*
Res 1963; **2**: 99-111.
33.   Maurice DM. The transparency of the corneal stroma. *Vision research* 1970; **10**: 107-108.
34.   Noske W, Fromm M, Levarlet B, *et al.* Tight junctions of the human corneal endothelium:
morphological and electrophysiological features. *Ger J Ophthalmol* 1994; **3**: 253-257.
35.   Ma L, Kuang K, Smith RW, *et al.* Modulation of tight junction properties relevant to fluid
transport across rabbit corneal endothelium. *Exp Eye Res* 2007; **84**: 790-798.
36.   Maurice DM, Srinivas SP. Fluorometric measurement of light absorption by the rabbit cornea.
Exp Eye Res 1994; **58**: 409-413.
37.   Bonanno JA. Identity and regulation of ion transport mechanisms in the corneal endothelium.
Prog Retin Eye Res 2003; **22**: 69-94.
38.   Riley MV, Winkler BS, Peters MI, *et al.* Relationship between fluid transport and in situ
inhibition of Na(+)-K+ adenosine triphosphatase in corneal endothelium. *Invest Ophthalmol*
Vis Sci 1994; **35**: 560-567.
39.   Ogando DG, Choi M, Shyam R, *et al.* Ammonia sensitive SLC4A11 mitochondrial uncoupling
reduces glutamine induced oxidative stress. *Redox Biology* 2019; **26**: 101260.
40.   Han SB, Ang HP, Poh R, *et al.* Mice with a targeted disruption of Slc4a11 model the
progressive corneal changes of congenital hereditary endothelial dystrophy. *Invest*
Ophthalmol Vis Sci 2013; **54**: 6179-6189.
41.   Jiao X, Sultana A, Garg P, *et al.* Autosomal recessive corneal endothelial dystrophy (CHED2) is
associated with mutations in SLC4A11. *J Med Genet* 2007; **44**: 64-68.
42.   Hemadevi B, Veitia RA, Srinivasan M, *et al.* Identification of mutations in the SLC4A11 gene
in patients with recessive congenital hereditary endothelial dystrophy. *Archives of*
ophthalmology 2008; **126**: 700-708.
43.   Sultana A, Garg P, Ramamurthy B, *et al.* Mutational spectrum of the SLC4A11 gene in
autosomal recessive congenital hereditary endothelial dystrophy. *Mol Vis* 2007; **13**: 1327-
1332.

44.   Aldave AJ, Yellore VS, Bourla N, *et al.* Autosomal recessive CHED associated with novel
compound heterozygous mutations in SLC4A11. *Cornea* 2007; **26**: 896-900.
45.   Ramprasad VL, Ebenezer ND, Aung T, *et al.* Novel SLC4A11 mutations in patients with
recessive congenital hereditary endothelial dystrophy (CHED2). Mutation in brief #958.
Online. *Human mutation* 2007; **28**: 522-523.
46.   Shah SS, Al-Rajhi A, Brandt JD, *et al.* Mutation in the SLC4A11 gene associated with
autosomal recessive congenital hereditary endothelial dystrophy in a large Saudi family.
Ophthalmic Genet 2008; **29**: 41-45.
47.   Kumar A, Bhattacharjee S, Prakash DR, *et al.* Genetic analysis of two Indian families affected
with congenital hereditary endothelial dystrophy: two novel mutations in SLC4A11. *Mol Vis*
2007; **13**: 39-46.
48.   Paliwal P, Sharma A, Tandon R, *et al.* Congenital hereditary endothelial dystrophy - mutation
analysis of SLC4A11 and genotype-phenotype correlation in a North Indian patient cohort.
Mol Vis 2010; **16**: 2955-2963.
49.   Liskova P, Dudakova L, Tesar V, *et al.* Detailed assessment of renal function in a proband
with Harboyan syndrome caused by a novel homozygous SLC4A11 nonsense mutation.
Ophthalmic research 2015; **53**: 30-35.
50.   Kodaganur SG, Kapoor S, Veerappa AM, *et al.* Mutation analysis of the SLC4A11 gene in
Indian families with congenital hereditary endothelial dystrophy 2 and a review of the
literature. *Mol Vis* 2013; **19**: 1694-1706.
51.   Park SH, Jeong HJ, Kim M, *et al.* A novel nonsense mutation of the SLC4A11 gene in a Korean
patient with autosomal recessive congenital hereditary endothelial dystrophy. *Cornea* 2013;
32: e181-182.
52.   Kumawat BL, Gupta R, Sharma A, *et al.* Delayed onset of congenital hereditary endothelial
dystrophy due to compound heterozygous SLC4A11 mutations. *Indian J Ophthalmol* 2016; **64**:
492-495.
53.   Kim JH, Ko JM, Tchah H. Fuchs Endothelial Corneal Dystrophy in a Heterozygous Carrier of
Congenital Hereditary Endothelial Dystrophy Type 2 with a Novel Mutation in SLC4A11.
Ophthalmic Genet 2015; **36**: 284-286.
54.   Kaul H, Suman M, Khan Z, *et al.* Missense mutation in SLC4A11 in two Pakistani families
affected with congenital hereditary endothelial dystrophy (CHED2). *Clin Exp Optom* 2016; **99**:
73-77.
55.   Cunnusamy K, Bowman CB, Beebe W, *et al.* Congenital Corneal Endothelial Dystrophies
Resulting From Novel De Novo Mutations. *Cornea* 2016; **35**: 281-285.
56.   Hand CK, McGuire M, Parfrey NA, *et al.* Homozygous SLC4A11 mutation in a large Irish
CHED2 pedigree. *Ophthalmic Genet* 2017; **38**: 148-151.
57.   Brejchova K, Dudakova L, Skalicka P, *et al.* IPSC-Derived Corneal Endothelial-like Cells Act as
an Appropriate Model System to Assess the Impact of SLC4A11 Variants on Pre-mRNA
SplicingCorneal Endothelial-like Cells Expressing SLC4A11. *Investigative Ophthalmology &*
Visual Science 2019; **60**: 3084-3090.
58.   Moazzeni H, Javadi MA, Asgari D, *et al.* Observation of nine previously reported and 10 non-
reported SLC4A11 mutations among 20 Iranian CHED probands and
identification of an MPDZ mutation as possible cause of CHED and FECD in one
family. *British Journal of Ophthalmology* 2019: e314377.
59.   Alka K, Casey JR. Ophthalmic Nonsteroidal Anti-Inflammatory Drugs as a Therapy for Corneal
Dystrophies Caused by SLC4A11 Mutation. *Invest Ophthalmol Vis Sci* 2018; **59**: 4258-4267.
60.   Alka K, Casey JR. Molecular phenotype of SLC4A11 missense mutants: Setting the stage for
personalized medicine in corneal dystrophies. *Human mutation* 2018; **39**: 676-690.
61.   Malhotra D, Loganathan SK, Chiu AM, *et al.* Human Corneal Expression of SLC4A11, a Gene
Mutated in Endothelial Corneal Dystrophies. *Scientific Reports* 2019; **9**: 9681.

62.   Li S, Hundal KS, Chen X, *et al.* R125H, W240S, C386R, and V507I SLC4A11 mutations
591   associated with corneal endothelial dystrophy affect the transporter function but not
592   trafficking in PS120 cells. *Exp Eye Res* 2019; **180**: 86-91.
593   63.   Kao L, Azimov R, Abuladze N, *et al.* Human SLC4A11-C functions as a DIDS-stimulatable
594   H(+)(OH(-)) permeation pathway: partial correction of R109H mutant transport. *Am J Physiol*
595   *Cell Physiol* 2015; **308**: C176-188.
596   64.   Frausto RF, Le DJ, Aldave AJ. Transcriptomic Analysis of Cultured Corneal Endothelial Cells as
597   a Validation for Their Use in Cell Replacement Therapy. *Cell Transplant* 2016; **25**: 1159-1176.
598   65.   Wang T, Pahlberg J, Cafaro J, *et al.* Activation of Rod Input in a Model of Retinal
599   Degeneration Reverses Retinal Remodeling and Induces Formation of Functional Synapses
600   and Recovery of Visual Signaling in the Adult Retina. *J Neurosci* 2019; **39**: 6798-6810.
601   66.   Yang NC, Ho WM, Chen YH, *et al.* A convenient one-step extraction of cellular ATP using
602   boiling water for the luciferin-luciferase assay of ATP. *Anal Biochem* 2002; **306**: 323-327.
603   67.   Swamynathan SK, Davis J, Piatigorsky J. Identification of candidate Klf4 target genes reveals
604   the molecular basis of the diverse regulatory roles of Klf4 in the mouse cornea. *Invest*
605   *Ophthalmol Vis Sci* 2008; **49**: 3360-3370.
606   68.   Okumura N, Nakamura T, Kay EP, *et al.* R-spondin1 regulates cell proliferation of corneal
607   endothelial cells via the Wnt3a/beta-catenin pathway. *Invest Ophthalmol Vis Sci* 2014; **55**:
608   6861-6869.
609   69.   Chng Z, Peh GSL, Herath WB, *et al.* High Throughput Gene Expression Analysis Identifies
610   Reliable Expression Markers of Human Corneal Endothelial Cells. *PLoS ONE* 2013; **8**: e67546.
611   70.   Shimizu M, Goto T, Sato A, *et al.* WNK4 is an essential effector of anterior formation in FGF
612   signaling. *Genes Cells* 2013; **18**: 442-449.
613   71.   Bonanno JA. Molecular mechanisms underlying the corneal endothelial pump. *Experimental*
614   *eye research* 2012; **95**: 2-7.
615   72.   Ostojic J, Grozdanic S, Syed NA, *et al.* Neuroglobin and cytoglobin distribution in the anterior
616   eye segment: a comparative immunohistochemical study. *J Histochem Cytochem* 2008; **56**:
617   863-872.
618   73.   Cheong YK, Ngoh ZX, Peh GS, *et al.* Identification of cell surface markers glypican-4 and
619   CD200 that differentiate human corneal endothelium from stromal fibroblasts. *Invest*
620   *Ophthalmol Vis Sci* 2013; **54**: 4538-4547.
621   74.   Yildiz E, Bardak H, Gunay M, *et al.* Novel Zinc Finger Protein Gene 469 (ZNF469) Variants in
622   Advanced Keratoconus. *Curr Eye Res* 2017; **42**: 1396-1400.
623   75.   Vincent AL, Jordan CA, Cadzow MJ, *et al.* Mutations in the zinc finger protein gene, ZNF469,
624   contribute to the pathogenesis of keratoconus. *Invest Ophthalmol Vis Sci* 2014; **55**: 5629-
625   5635.
626   76.   Lechner J, Porter LF, Rice A, *et al.* Enrichment of pathogenic alleles in the brittle cornea gene,
627   ZNF469, in keratoconus. *Hum Mol Genet* 2014; **23**: 5527-5535.
628   77.   Lu Y, Dimasi DP, Hysi PG, *et al.* Common genetic variants near the Brittle Cornea Syndrome
629   locus ZNF469 influence the blinding disease risk factor central corneal thickness. *PLoS Genet*
630   2010; **6**: e1000947.
631   78.   Zhang X, Tseng H. Basonuclin-null mutation impairs homeostasis and wound repair in mouse
632   corneal epithelium. *PLoS One* 2007; **2**: e1087.
633   79.   Chung DD, Frausto RF, Lin BR, *et al.* Transcriptomic Profiling of Posterior Polymorphous
634   Corneal Dystrophy. *Invest Ophthalmol Vis Sci* 2017; **58**: 3202-3214.
635   80.   Ehlers N, Modis L, Moller-Pedersen T. A morphological and functional study of Congenital
636   Hereditary Endothelial Dystrophy. *Acta Ophthalmol Scand* 1998; **76**: 314-318.
637   81.   Xian H, Yang Q, Xiao L, *et al.* STX17 dynamically regulated by Fis1 induces mitophagy via
638   hierarchical macroautophagic mechanism. *Nature Communications* 2019; **10**: 2059.

82.   Nieminen AI, Eskelinen VM, Haikala HM, *et al.* Myc-induced AMPK-phospho p53 pathway
activates Bak to sensitize mitochondrial apoptosis. *Proceedings of the National Academy of*
Sciences 2013; **110**: E1839-E1848.
83.   Okoshi R, Ozaki T, Yamamoto H, *et al.* Activation of AMP-activated Protein Kinase Induces
p53-dependent Apoptotic Cell Death in Response to Energetic Stress. *Journal of Biological*
Chemistry 2008; **283**: 3979-3987.
84.   Jones RG, Plas DR, Kubek S, *et al.* AMP-Activated Protein Kinase Induces a p53-Dependent
Metabolic Checkpoint. *Molecular Cell* 2005; **18**: 283-293.
85.   Herzig S, Shaw RJ. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nature*
Reviews Molecular Cell Biology 2017; **19**: 121.
86.   Loganathan SK, Casey JR. Corneal dystrophy-causing SLC4A11 mutants: suitability for folding-
correction therapy. *Human mutation* 2014; **35**: 1082-1091.
87.   Kirkness CM, McCartney A, Rice NS, *et al.* Congenital hereditary corneal oedema of
Maumenee: its clinical features, management, and pathology. *Br J Ophthalmol* 1987; **71**:
130-144.
88.   Breschi A, Djebali S, Gillis J, *et al.* Gene-specific patterns of expression variation across
organs and species. *Genome Biology* 2016; **17**: 151.
89.   Breschi A, Gingeras TR, Guigo R. Comparative transcriptomics in human and mouse. *Nat Rev*
Genet 2017; **18**: 425-440.
90.   Assentoft M, Kaptan S, Schneider H-P, *et al.* Aquaporin 4 as a NH3 Channel. *The Journal of*
biological chemistry 2016; **291**: 19184-19195.
91.   Mak D-OD, Dang B, Weiner ID, *et al.* Characterization of ammonia transport by the kidney Rh
glycoproteins RhBG and RhCG. *American Journal of Physiology-Renal Physiology* 2006; **290**:
F297-F305.

FIGURE LEGEND

Figure 1. SLC4A11 deficiency leads to transcriptome alteration in corneal

endothelium. (A) Histopathologic examination of control and CHED corneas

demonstrating Descemet membrane thickening and endothelial cell attenuation in

corneas from individuals with CHED (H&E stain, scale bar, 25 µm). (B) Left:

representative images of immunofluorescence staining for SLC4A11 (green signal)

and ER marker PDI (red signal) in corneal endothelium of two individuals with CHED

and healthy control. Nucleus were stained with DAPI (blue signal) (Scale bar, 100

µm); Right: scatterplot of mean fluorescence intensity (MFI) ratio of SLC4A11 over

PDI in corneal endothelium of two individuals with CHED and of 7 healthy controls.

(two-tailed unpaired t-test with Welch's correction, p = 0.5127). (C) Venn diagram of

the endothelial transcriptome in *SLC4A11* KD pHCEnC, *Slc4a11*[-/-] MCEnC early and

late passages indicating the number of differentially expressed genes (DEG) in each

of the sample sets. (D) Venn diagram of the endothelial transcriptome in *SLC4A11*

KD pHCEnC, *Slc4a11*[-/-] MCEnC early and late passages indicating the number of

genes differentially expressed in the same direction in each of the sample sets. (E)

Heatmap of top 30 most highly differentially expressed genes shared between

SLC4A11 KD pHCEnC, *Slc4a11*[-/-] MCEnC early and late passages. Genes are

clustered based on cellular localization of the gene product.

Figure 2. Inhibition of multiple metabolic pathways in SLC4A11-deficient

corneal endothelial cells. (A) Heatmap showing consensually enriched IPA

canonical pathways from comparison of transcriptomes of *SLC4A11* KD pHCEnC,

Slc4a11[-/-] MCEnC early and late passages (sorted by mean "activation z-score") (B)

Schematic illustration of the crosstalk between identified inhibited metabolic

pathways. Differentially expressed genes encoding key enzymes are color coded

using the same color as used in C-I. Transcript levels of differentially expressed

enzyme-coding genes involved in: (C) citric acid (TCA) cycle (*IDH3G,* isocitrate

dehydrogenase 3 (NAD$^+$) gamma; *SUCLA2,* succinate-CoA ligase ADP-forming beta

subunit; *SDHD,* succinate dehydrogenase complex subunit D; and *FH,* fumarate

hydratas); (D) glycolysis (*GLUT1/3,* glucose transporter type 1 and 3; *GPI,* glucose-

6-phosphate isomerase; *PFKL,* phosphofructokinase, liver type; *PFKP,*

phosphofructokinase, platelet; *ALDOA,* aldolase, fructose-bisphosphate A; *TPI1,*

triosephosphate isomerase 1; *PGK1,* phosphoglycerate kinase 1; *ENO1,* enolase 1;

and *LDHD,* lactate dehydrogenase); (E) acetyl-CoA biosynthesis/pyruvate

dehydrogenase (PDH) complex (*PDK3*, pyruvate dehydrogenase kinase 3; *PDHB*,

pyruvate dehydrogenase E1 beta subunit; and *DLAT*, dihydrolipoamide S-

acetyltransferase); (F) glutaminolysis (*SNAT1*, system N amino acid transporter 1;

and *GOT2*, glutamic-oxaloacetic transaminase 2); (G) pentose phosphate shunt

(*G6PD*, glucose-6-phosphate dehydrogenase; and *PGD*, phosphogluconate

dehydrogenase); (H) GDP-Mannose synthesis (*MPI*, mannose phosphate isomerase;

GMPPB, GDP-mannose pyrophosphorylase B in UDP-acetyl-galactosamine

synthesis; *GNPDA1*, glucosamine-6-phosphate deaminase 1; and *PGM2L1*,

phosphoglucomutase 2 like 1); (I) pyrimidine ribonucleotides (UTP/CTP) and

pyrimidine deoxyribonucleotides (dTTP/dCTP) *de novo* synthesis (*CAD*, carbamoyl-

phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase; *CMPK1*,

cytidine/uridine monophosphate kinase 1; *RRM1*, ribonucleotide reductase catalytic

subunit M1; *RRM2*, ribonucleotide reductase regulatory subunit M2; *NME1*,

NME/NM23 nucleoside diphosphate kinase 1 and *NME2*, NME/NM23 nucleoside

diphosphate kinase 2). (J) Selected differentially expressed genes from each

pathway listed above were validated by qPCR in separate RNA isolations from

SLC4A11 KD pHCEnC and *Slc4a11$^{-/-}$* MCEnC early and late passages. Genes from

different pathways are clustered and separated by dashed lines.

Figure 3. SLC4A11 deficiency impact*s* corneal endotheli*al* ion and solute

transport *function.* (A) Heatmap showing consensually enriched IPA biological

function pathways from comparison of transcriptomes of SLC4A11 KD pHCEnC,

Slc4a11$^{-/-}$ MCEnC early and late passages (sorted by mean "activation z-score"). (B)

Representative trace of current-clamped single cell recording during 10 mM NH$_4$Cl

superfusion of *Slc4a11$^{+/+}$* and *Slc4a11$^{-/-}$* MCEnC. (C) Boxplot of membrane potential

changes (*d* Vm) in *Slc4a11$^{+/+}$* (n = 6) and *Slc4a11$^{-/-}$* (n =7) MCEnC in response to

NH$_4$Cl superfusion. Each individual data point is plotted as a dot. [Monte Carlo

resampling two-tailed paired-sampled t-test, difference between genotype *ΔVm* = -

22.2 (-34.8, -10) mV*, p* = 0.0069]. (D) Heatmap showing common differentially

expressed genes encoding ion channel and transporter proteins in *SLC4A11* KD

pHCEnC and early and late *Slc4a11$^{-/-}$* MCEnC. (E) The differential expression of

Na$^+$-HCO$_3^-$ transporter (NBCe1, encoded by *SLC4A4*) mRNA was validated by

qPCR in separate RNA isolations from *SLC4A11* KD pHCEnC and *Slc4a11$^{-/-}$*

MCEnC. (F) Western Blot for NBCe1 in *SLC4A11* KD pHCEnC and scRNA pHCEnC

control showing decreased NBCe1 protein level in *SLC4A11* KD pHCEnC. (G)

Representative images of immunofluorescence staining for NBCe1 (green signal)

and tight junction ZO-1 (red signal) in corneal endothelium of two individuals with

CHED and healthy control. Nuclei were stained with DAPI (blue signal) (scale bar, 10

μm). (H) Scatterplot of mean fluorescence intensity (MFI) ratio of NBCe1 over ZO-1

in corneal endothelium of two individuals with CHED and of seven healthy controls.

(two-tailed unpaired t-test with Welch's correction, *p* = 0.0031). (I) Representative

trace of intracellular pH (pH$_i$) response in *Slc4a11$^{-/-}$* and *Slc4a11$^{+/+}$* MCEnC to

addition of extracellular Na$^+$ in HCO$_3^-$ containing solution. (J) Bar graph of the rate of

intracellular $[H^+]$ change ($d[H^+]/ds$) in *Slc4a11$^{+/+}$* (n = 6) and *Slc4a11$^{-/-}$* (n =8) MCEnC

(two tailed unpaired t-test, *p* < 0.0001).

Figure 4. Mitochondria dysfunction in SLC4A11 deficient corneal endothelium.

(A) Heatmap of commonly enriched IPA canonical and biological function pathways

in transcriptomes of *SLC4A11* KD pHCEnC, early and late Slc4a11[-/-] MCEnC without

assigned activation z-scores (sorted by mean "adjusted p-value"). (B) Transcript

levels of differentially expressed enzyme coding genes involved in mitochondria

electron transport chain (Complex I: *NDUFA1, NDUFA9, NDUFA10*,

NADH:Ubiquinone Oxidoreductase subunit A1, 9 and 10; *NDUFAB1,*

NADH:Ubiquinone Oxidoreductase Subunit AB1; Complex II: SDHD, Succinate

Dehydrogenase Complex Subunit D; Complex III: *UQCRB*, Ubiquinol-Cytochrome C

Reductase Binding Protein; *UQCRC2*, Ubiquinol-Cytochrome C Reductase Core

Protein 2; *CYC1*, Cytochrome C1; Complex IV: *COX5A*, COX5B and *COX7B*,

Cytochrome C Oxidase Subunit 5A, 5B and 7B; Complex V: *ATP5F1*, ATP Synthase

Peripheral Stalk-Membrane Subunit B; *ATP5G1*, ATP Synthase Membrane Subunit

C Locus 1). (C) Transcript levels of differentially expressed protein coding genes that

serve as mitochondrial functional markers (ATP flux: *VDAC2/3,* Voltage-dependent

Anion-selective Channel; import machinery: *TIMM7A, TIMM8A, TIMM50*,

Translocase of Inner Mitochondrial Membrane 7A, 8A, 50; translation machinery:

GFM2, G Elongation Factor Mitochondrial 2*; MRPL10, MRPL18, MRPL20, MRPL33,*

MRPL48, MRPL52, MRPL58, MRPL16, MRPS2, MRPS18A, Mitochondrial

Ribosomal Protein L10*,* L18, L20, L33, L48, L52, L58, L16, S2, S18A;). (D)

Transcript levels of differentially expressed *STX17* in *SLC4A11* KD pHCEnC and

Slc4a11[-/-] MCEnC transcriptomes. (E) Differential expression of *STX17* mRNA in

SLC4A11 KD pHCEnC and *Slc4a11[-/-]* MCEnC validated by qPCR in separate RNA

isolations. (F) Western Blot for STX17 and NBCe1in scRNA pHCEnC control and

SLC4A11 KD pHCEnC showing decreased protein level in *SLC4A11* KD pHCEnC*.*

(G) Representative images of immunofluorescence staining for STX17 (green signal)

and mitochondria marker COX4 (red signal) in corneal endothelium of two individuals

with CHED and healthy control. Nuclei were stained with DAPI (blue signal)(scale

bar, 10 µm). (H) Scatterplot of mean fluorescence intensity (MFI) ratio of STX17 over

COX4 in corneal endothelium of two individuals with CHED and of seven healthy

controls (two-tailed unpaired t-test with Welch's correction, p = 0.0483). (I)

Scatterplot of MFI of COX4 in corneal endothelium of two individuals with CHED and

of seven healthy controls (two-tailed unpaired t-test with Welch's correction, p =

0.0018). (J) Representative images of immunofluorescence staining for cytochrome

C (green signal) in corneal endothelium of two individuals with CHED and of seven

healthy controls. Nucleus were stained with DAPI (blue signal) (scale bar, 5 µm). (K)

Scatterplot of MFI of cytochrome C in corneal endothelium of two individuals with

CHED and of seven healthy controls (two-tailed unpaired t-test with Welch's

correction, p = 0.0304). (L) Bar graph summary of intracellar ATP levels measured in

SLC4A11 KD pHCEnC and scRNA pHCEnC controls (n = 6 each, one-tailed

unpaired t-test, p = 0.0281). (M) Bar graph summary of intracellar ATP levels

measured in early passage (n=6 each, two-tailed unpaired t-test, p = 0.034) and late

passage (n=6 each, p = 0.039) *Slc4a11$^{+/+}$* and *Slc4a11$^{-/-}$* MCEnC.

Figure 5. Activation of AMPK-p53 pathway in SLC4A11 deficient corneal

endothelium. (A) Heatmap of shared predicted upstream regulators of

transcriptomes of SLC4A11 KD pHCEnC, early and late Slc4a11$^{-/-}$ MCEnC (sorted

by mean "adjusted p-value"). (B) Western blot analysis of *SLC4A11* KD pHCEnC

and scRNA controls for SLC4A11, Ser15 phosphorylated p53 (phos-p53 (S15)), total

p53, Thr172 phosphorylated AMPKα (phos-AMPKα (T172)), total AMPKα, Ser182

phosphorylated AMPKβ (phos-AMPKβ1 (S182)) and total AMPKβ1 and β2. (C)

Western blot analysis of early and late *Slc4a11$^{+/+}$* and *Slc4a11$^{-/-}$* MCEnC for Ser18

phosphorylated p53 (phos-p53 (S18)), total p53, Thr172 phosphorylated AMPKα

(phos-AMPKα (T172)), total AMPKα, Ser182 phosphorylated AMPKβ (phos-

AMPKβ1 (S182)) and total AMPKβ1 and β2. (D) Bar graph summary of Western blot

densitometry results for normalized total p53 level in *SLC4A11* KD pHCEnC versus

scRNA pHCEnC (CTRL) (two-tailed unpaired-sample t-test, $p > 0.99$), in early

passage *Slc4a11$^{-/-}$* versus *Slc4a11$^{+/+}$* MCEnC ($p = 0.00027$) and late passage

Slc4a11$^{-/-}$ versus *Slc4a11$^{+/+}$* MCEnC ($p = 0.00027$).

A

Healthy control
▷ Corneal Stroma
▷ Descemet's membrane
▶ Corneal Endothelium

CHED

SLC4A11^{Mu/WT}
▷ Descemet's membrane
▶ Corneal Endothelium

SLC4A11^{Mu/Mu}
▷ Descemet's membrane
▶ Corneal Endothelium

B

DAPI SLC4A11 PDI

Healthy control

CHED

SLC4A11^{Mu/WT}

SLC4A11^{Mu/Mu}

SLC4A11

Mean Fluorescence Intensity (AU / area)

CTRL   CHED

C

743   771   323

3171

1265

335   159

SLC4A11 KD pHCEnC
Slc4a11^{-/-} MCEnC early
Slc4a11^{-/-} MCEnC late

D

2462   737   1662

1041

780   2090

1019

E

log₂FC
-10   0   10

TNFSF10 PTN PAPPA SEMA3E IL12B MST1R RNF43 SLC2A3 PTGIR WNK4 ADGRG2 MEGF10 PCDHB6 LRRC4C TSPAN7 PKP1 SPTA1 SYNPO2 PPL CYGB CYP2S1 SLC9A7 NFE2L3 ATOH8 RNF469 BNC1 TTC22 PCDHGB2 VAT1L

SLC4A11 KD pHCEnC
Slc4a11^{-/-} MCEnC early
Slc4a11^{-/-} MCEnC late

Extracellular Space   Plasma Membrane   Cytoplasm   Nucleus   Other