Phagocytic glia are obligatory intermediates in transmission of mutant huntingtin aggregates across neuronal synapses

Kirby M. Donnelly[1], Olivia R. DeLorenzo[2], Wint M. Thu[1], Gabrielle E. Pisano[1], Liqun Luo[3,4], Ron R. Kopito[3], Margaret M. Panning Pearce[1,2,5]

[1]Department of Biological Sciences and [2]Program in Neuroscience, University of the Sciences, Philadelphia, PA 19063, USA

[3]Department of Biology, Stanford University, Stanford, CA 94305, USA

[4]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA

[5]Correspondence: m.pearce@usciences.edu

ABSTRACT

Emerging evidence supports the hypothesis that pathogenic protein aggregates associated with many neurodegenerative diseases spread from cell to cell through the brain in a manner akin to infectious prions. Here, we show that mutant huntingtin (mHtt) aggregates associated with Huntington disease transfer anterogradely from presynaptic to postsynaptic neurons in the adult *Drosophila* olfactory system. Trans-synaptic transmission of mHtt aggregates is inversely correlated with neuronal activity and blocked by caspase inhibition in presynaptic neurons, implicating synaptic dysfunction and cell death in aggregate spreading. Remarkably, mHtt aggregate transmission across synapses requires the scavenger receptor Draper and involves a transient visit to the glial cytoplasm, indicating that phagocytic glia act as obligatory intermediates in aggregate spreading between synaptically-connected neurons. These findings expand our understanding of phagocytic glia as double-edged players in neurodegeneration—they clear neurotoxic protein aggregates, but also provide opportunities for prion-like seeds to evade phagolysosomal degradation and propagate further in the brain.

INTRODUCTION

Neurodegenerative diseases have emerged as one of the greatest healthcare challenges in our

aging society, and thus a better understanding the underlying pathological mechanisms is critical to

development of more effective therapies or cures for these fatal disorders. A common molecular feature

of many neurodegenerative diseases [e.g., Alzheimer disease (AD), frontotemporal dementias (FTD),

Parkinson disease (PD), amyotrophic lateral sclerosis (ALS), and Huntington disease (HD)] is the

misfolding of certain proteins, driving their accumulation into insoluble, amyloid aggregates (Knowles et

al., 2014). Appearance of proteinaceous deposits in patient brains correlates closely with neuronal loss

and clinical progression, and strategies to lower production or enhance clearance of pathological

proteins in the degenerating brain have shown therapeutic promise in animal models and clinical trials

(Boland et al., 2018; Li et al., 2019; Tabrizi et al., 2019).

Post-mortem histopathological analyses (Braak and Braak, 1991; Braak et al., 2003;

Brettschneider et al., 2013) and *in vivo* imaging studies (Deng et al., 2004; Poudel et al., 2019) indicate

that proteopathic lesions associated with neurodegeneration appear in highly-reproducible and disease-

specific spatiotemporal patterns through the brain. Interestingly, these patterns largely follow

neuroanatomical tracts (Ahmed et al., 2014; Mezias et al., 2017), suggesting a central role for synaptic

connectivity in pathological aggregate spreading. Accumulating evidence supports the idea that

intracellular aggregates formed by tau, α-synuclein, TDP-43, SOD1, and mutant huntingtin (mHtt)

transfer from cell-to-cell and self-replicate by recruiting natively-folded versions of the same protein,

analogous to how infectious prion protein (PrP^{Sc}) templates the conformational change of soluble PrP^C

in prion diseases (Vaquer-Alicea and Diamond, 2019). Numerous studies have pointed to roles for

endocytosis and exocytosis (Asai et al., 2015; Babcock and Ganetzky, 2015; Chen et al., 2019; Holmes

et al., 2013; Lee et al., 2010; Zeineddine et al., 2015), membrane permeabilization (Chen et al., 2019;

Falcon et al., 2018; Flavin et al., 2017; Zeineddine et al., 2015), tunneling nanotubes (Costanzo et al.,

2013; Sharma and Subramaniam, 2019), and neuronal activity (Wu et al., 2016) in entry and/or exit of

pathogenic protein assemblies from cells, but the exact mechanisms by which amyloid aggregates or

on-pathway intermediates could cross one or more biological membranes in the highly complex central

nervous system (CNS) remain an enigma.

Glia are resident immune cells of the CNS, constantly surveying the brain to maintain

homeostasis and to respond rapidly to tissue damage or trauma. Reactive astrocytes and microglia

provide a first line of defense in neurodegeneration by infiltrating sites of neuronal injury, upregulating

immune-responsive genes, and phagocytosing dying neurons and other debris, including protein

aggregates (Asai et al., 2015; Grathwohl et al., 2009; Wyss-Coray et al., 2003). Prolonged activation of

these glial responses results in chronic inflammation, exacerbating synaptic dysfunction and neuronal

loss (Hammond et al., 2018). We and others have previously shown that Draper, a *Drosophila*

scavenger receptor responsible for recognizing and phagocytosing cellular debris (Freeman, 2015),

regulates the load of mHtt (Pearce et al., 2015) and $A\beta_{1-42}$ (Ray et al., 2017) aggregate pathology in the

fly CNS. Remarkably, we also found that a portion of phagocytosed neuronal mHtt aggregates gain

entry into the glial cytoplasm and there nucleate the aggregation of normally-soluble wild-type Htt

(wtHtt) proteins, suggesting that glial phagocytosis provides a route for spreading of prion-like

aggregates in intact brains. Consistent with our results, microglial ablation suppresses pathological tau

transmission between synaptically-connected regions of the mouse brain (Asai et al., 2015), and PrP^{Sc}

transfers from infected astrocytes to co-cultured neurons (Victoria et al., 2016). Thus, these findings

suggest that phagocytic glia could play double-edged roles in neurodegeneration, with normally

neuroprotective clearance mechanisms contributing to dissemination of prion-like aggregate species

through the CNS.

A plethora of studies from the last decade have strengthened the prion-like hypothesis for

neurodegenerative diseases, but we still lack a clear understanding of how pathogenic protein

aggregates spread in an intact CNS. In this study, we adapted our previously-described *Drosophila* HD

model to examine roles for synaptic connectivity and phagocytic glia in prion-like mHtt aggregate

transmission in the fly CNS. HD is an autosomal dominant disorder caused by expansion of a CAG

repeat region in exon 1 of the Htt gene, resulting in production of highly aggregation-prone mHtt

proteins containing abnormally expanded polyglutamine (polyQ$\geq$37) tracts (Bates et al., 2015;

MacDonald et al., 1993). By contrast, wtHtt proteins containing polyQ≤36 tracts only aggregate upon

nucleation by pre-formed Htt aggregate "seeds" (Chen et al., 2001; Preisinger et al., 1999). A growing

body of evidence from cell culture (Chen et al., 2001; Costanzo et al., 2013; Holmes et al., 2013; Ren et

al., 2009; Sharma and Subramaniam, 2019; Trevino et al., 2012) and *in vivo* (Ast et al., 2018; Babcock

and Ganetzky, 2015; Jeon et al., 2016; Masnata et al., 2019; Pearce et al., 2015; Pecho-Vrieseling et

al., 2014) models of HD supports the idea that pathogenic mHtt aggregates have prion-like properties—

they transfer from cell to cell and template the conformational change of soluble wtHtt proteins. Here,

we report that mHtt aggregates formed in presynaptic olfactory receptor neuron (ORN) axons effect

prion-like conversion of wtHtt proteins expressed in the cytoplasm of their postsynaptic partner

projection neurons (PNs) in the adult fly olfactory system. Remarkably, we find that transfer of mHtt

aggregates from presynaptic ORNs to postsynaptic PNs was abolished in Draper-deficient animals and

requires passage of the aggregate seeds through the cytoplasm of Draper-dependent glial cells.

Together, these findings support the conclusion that phagocytic glia are obligatory intermediates in

prion-like transmission of mHtt aggregates between synaptically-connected neurons *in vivo*.

RESULTS

Anterograde transfer of prion-like mHtt aggregates across synapses in the adult fly olfactory system

Aggregates formed by N-terminal fragments of mHtt generated by aberrant splicing (e.g., exon 1; Htt_{ex1}) (Sathasivam et al., 2013) or caspase cleavage (e.g., exon 1-12; Htt_{ex1-12}) (Graham et al., 2006) (Fig. 1A) accumulate in HD patient brains, are highly cytotoxic, and spread between cells in culture and *in vivo* (Babcock and Ganetzky, 2015; Costanzo et al., 2013; Pearce et al., 2015; Pecho-Vrieseling et al., 2014; Ren et al., 2009). We have previously established transgenic *Drosophila* that employ binary expression systems [e.g., Gal4-UAS, QF-QUAS, or LexA-LexAop (Riabinina and Potter, 2016)] to express fluorescent protein (FP) fusions of Htt_{ex1} in non-overlapping cell populations in order to monitor cell-to-cell transfer of $mHtt_{ex1}$ aggregates in intact brains (Donnelly and Pearce, 2018; Pearce et al., 2015). Our experimental approach (Fig. 1B) exploits the fact that $wtHtt_{ex1}$ proteins only aggregate upon encountering $mHtt_{ex1}$ aggregate seeds (Chen et al., 2001; Preisinger et al., 1999), such that transfer of $mHtt_{ex1}$ aggregates from "donor" cells is reported by conversion of cytoplasmic $wtHtt_{ex1}$ from its normally soluble, diffuse state to a punctate, aggregated state in "acceptor" cells. To confirm that $mHtt_{ex1}$ nucleates the aggregation of $wtHtt_{ex1}$ in fly neurons, we co-expressed FP-fusions of these two proteins using pan-neuronal *elav[C155]-Gal4*. In flies expressing only mCherry-tagged $mHtt_{ex1}$ ($Htt_{ex1}Q91$-mCherry) pan-neuronally, aggregates were visible as discrete mCherry+ puncta throughout neuronal cell bodies and neuropil regions of adult fly brains (Fig. 1-figure supplement 1A). By contrast, GFP-tagged $wtHtt_{ex1}$ ($Htt_{ex1}Q25$-GFP) was expressed diffusely throughout neuronal cell bodies and processes in age-matched adults brains (Fig. 1-figure supplement 1B). Co-expression of these two Htt proteins converted $Htt_{ex1}Q25$-GFP to a punctate expression pattern that overlapped with $Htt_{ex1}Q91$-mCherry signal (Fig. 1-figure supplement 1C), whereas the solubility of GFP lacking a polyQ sequence was not affected by $Htt_{ex1}Q91$-mCherry co-expression (Fig. 1-figure supplement 1D). Thus, prion-like conversion of $Htt_{ex1}Q25$-GFP occurs only in the presence of $Htt_{ex1}Q91$-mCherry aggregates and results from a homotypic nucleation reaction rather than $mHtt_{ex1}$-induced global changes to proteostasis in the neuronal cytoplasm.

To examine trans-synaptic transfer of mHtt$_{ex1}$ aggregates, we coupled QF-driven expression of Htt$_{ex1}$Q91-mCherry with Gal4-driven expression of Htt$_{ex1}$Q25-GFP in neuronal cell populations that make well-defined synaptic connections in the adult fly olfactory system (Fig. 1B). Htt$_{ex1}$Q91-mCherry was expressed using *Or67d-QF* in presynaptic ORNs (hereafter "DA1 ORNs") that project axons from the antenna into the central brain, where they form synaptic connections with dendrites of partner PNs (hereafter "DA1 PNs") in the DA1 glomerulus of the antennal lobe (Jefferis et al., 2001). In these same animals, Htt$_{ex1}$Q25-GFP was expressed in ~60% of PNs using *GH146-Gal4*, which labels lateral and ventral DA1 PNs in addition to other PN types (Marin et al., 2002). We did not detect expression of Htt$_{ex1}$Q91-mCherry in DA1 ORNs until ~24 hr before eclosion, consistent with activation of adult olfactory receptor gene expression during late pupal development (Clyne et al., 1999), whereas Htt$_{ex1}$Q25-GFP was expressed in PNs much earlier in development (Stocker et al., 1997). Thus, this genetic approach enables us to monitor prion-like transfer of Htt$_{ex1}$Q91-mCherry aggregates between post-mitotic, synaptically-connected DA1 ORNs and PNs in the adult fly brain.

Prion-like transfer of Htt$_{ex1}$Q91-mCherry aggregates across DA1 ORN-PN synapses was examined by monitoring the solubility of Htt$_{ex1}$Q91-mCherry and Htt$_{ex1}$Q25-GFP in or near the DA1 glomerulus as the flies aged (Fig. 1B, inset). Puncta representing Htt$_{ex1}$Q91-mCherry aggregates were detected in the DA1 glomerulus of 1 day-old adults and increased in number over time (Fig. 1C and D). Numbers of Htt$_{ex1}$Q91-mCherry aggregates in males exceeded those in females at each time point tested, consistent with known sexual dimorphism in DA1 glomerular volume (Stockinger et al., 2005). In these same brains, Htt$_{ex1}$Q25-GFP was expressed diffusely in GH146+ PN cell bodies and dendrites in 1 day-old adults, but induced Htt$_{ex1}$Q25-GFP aggregates appeared in low quantities in the DA1 glomerulus of young adults and accumulated as the flies aged (Fig. 1C and E). Though rare, Htt$_{ex1}$Q25-GFP aggregates also appeared in PN soma in older adults (Fig. 1F and G), suggesting that these aggregates have the capacity to propagate beyond the DA1 glomerulus. In control experiments, Htt$_{ex1}$Q25-mCherry in DA1 ORNs did not alter Htt$_{ex1}$Q25-GFP solubility in PNs (Fig. 1-figure supplement 2A,E,F), and the solubility of GFP expressed in PNs was unaffected by Htt$_{ex1}$Q91-mCherry expression in DA1 ORNs (Fig. 1-figure supplement 2B,E,F). In addition, seeded Htt$_{ex1}$Q25-GFP aggregates still

appeared when the Gal4 inhibitor Gal80 was expressed in ORNs (Fig. 1-figure supplement 2C,E,F) or

when the QF repressor QS was expressed in PNs (Fig. 1-figure supplement 2D,E,F), arguing strongly

against the possibility that $Htt_{ex1}Q91$-mCherry and $Htt_{ex1}Q25$-GFP were co-expressed in any cells in

these transgenic animals. Because $Htt_{ex1}Q25$-GFP did not aggregate in the absence of $Htt_{ex1}Q91$-

mCherry expression (Fig. 1-figure supplement 1B), these findings indicate that the solubility of

$Htt_{ex1}Q25$-GFP in postsynaptic PNs was non-cell autonomously altered by $Htt_{ex1}Q91$-mCherry

aggregates formed in presynaptic ORNs.

Importantly, $Htt_{ex1}Q25$-GFP aggregates were not detected in any other glomeruli labeled by

GH146-Gal4 in these flies (Fig. 1C), suggesting that prion-like conversion of $Htt_{ex1}Q25$-GFP in PNs by

$Htt_{ex1}Q91$-mCherry aggregates in ORNs requires synaptic connectivity. To examine whether spreading

of $Htt_{ex1}Q91$-mCherry can also occur retrogradely across ORN-PN synapses, we expressed $Htt_{ex1}Q91$-

mCherry in GH146+ PNs and $Htt_{ex1}Q25$-GFP in all ORNs using *pebbled-Gal4*. While many $Htt_{ex1}Q91$-

mCherry aggregates were formed in PN soma, dendrites, and axons, we saw no evidence of seeded

$Htt_{ex1}Q25$-GFP aggregation in ORN axons in the antennal lobe (Fig. 1-figure supplement 3), indicating

that $Htt_{ex1}Q91$-mCherry spreading across ORN-PN synapses is restricted to or much more efficient in

the anterograde direction. Interestingly, we found that aggregates formed by $mHtt_{ex1-12}$ caspase-6

cleavage products (RFP-$Htt_{ex1-12}Q138$) (Fig. 1A) (Graham et al., 2006), which have previously been

shown to spread from ORN axons to more distant (non-PN) neurons in the fly CNS (Babcock and

Ganetzky, 2015), also transfer across DA1 ORN-PN synapses in our model (Fig. 1H). Thus, the ability

to spread trans-synaptically is shared by multiple pathogenic N-terminal mHtt fragments, and the

sequences required for prion-like conversion of wtHtt reside within Htt_{ex1}.

wtHtt$_{ex1}$ aggregates in PNs are seeded by smaller mHtt$_{ex1}$ aggregates originating in ORNs

Further examination of confocal stacks of the DA1 glomerulus from flies expressing $Htt_{ex1}Q91$-

mCherry in DA1 ORNs and $Htt_{ex1}Q25$-GFP in GH146+ PNs (Fig. 2A-D) indicated that the vast majority

(>95%) of induced $Htt_{ex1}Q25$-GFP aggregates co-localized with $Htt_{ex1}Q91$-mCherry puncta (arrows in

Fig. 2B and D). This suggests that the solubility of $Htt_{ex1}Q25$-GFP in PNs was altered upon direct

physical interaction with Htt$_{ex1}$Q91-mCherry seeds from DA1 ORNs. In support of this, fluorescence

resonance energy transfer (FRET) analysis showed that the FP tags on Htt$_{ex1}$Q91-mCherry and

Htt$_{ex1}$Q25-GFP were within ~100Å in all co-localized aggregates tested (Fig. 2E). While nearly all

Htt$_{ex1}$Q25-GFP aggregates in the DA1 glomerulus co-localized with Htt$_{ex1}$Q91-mCherry seeds, this was

true of only a small proportion of Htt$_{ex1}$Q25-GFP aggregates in PN soma (Fig. 1F and G), suggesting

that aggregate fragmentation may drive propagation outside of the antennal lobe neuropil. Interestingly,

volumetric analysis of all mCherry+ aggregates in the DA1 glomerulus revealed that seeding-competent

Htt$_{ex1}$Q91-mCherry aggregates belonged to a smaller-sized subpopulation (range = 0.1–2.2 μm^3; mean

= 0.859 $\pm$ 0.052 μm^3), whereas the entire population of Htt$_{ex1}$Q91-mCherry aggregates was more

heterogeneously-sized (range = 0.1–9 μm^3; mean = 2.676 $\pm$ 0.046 μm^3) (Fig. 2F). We previously

reported that Htt$_{ex1}$Q25-YFP aggregates formed in the glial cytoplasm co-localized with a similarly

smaller-sized subpopulation of Htt$_{ex1}$Q91-mCherry aggregates from DA1 ORN axons (Pearce et al.,

2015), and smaller mHtt$_{ex1}$ aggregates are associated with increased seeding-propensity and

neurotoxicity in other HD models (Ast et al., 2018; Chen et al., 2001). Taken together, these findings

strongly suggest that Htt$_{ex1}$Q91-mCherry aggregates formed in presynaptic ORNs gain access to the

cytoplasm of postsynaptic PNs to effect prion-like conversion of Htt$_{ex1}$Q25-GFP, and that Htt$_{ex1}$Q91-

mCherry aggregate transmissibility is inversely correlated with aggregate size.

mHtt$_{ex1}$ aggregate transfer is enhanced across silenced DA1 ORN-PN synapses

Endocytosis, exocytosis, and neuronal activity have been previously implicated in neuron-to-

neuron spreading of mHtt and other pathogenic aggregates (Babcock and Ganetzky, 2015; Pecho-

Vrieseling et al., 2014; Wu et al., 2016), but it is not known how these processes could contribute to

aggregate transfer across endogenous synapses *in vivo*. To examine a role for synaptic activity in DA1

ORN-to-PN transfer of mHtt$_{ex1}$ aggregates, we used well-established genetic tools that block fission or

fusion of synaptic vesicles at the presynaptic membrane, thereby impairing neurotransmission. First, we

used *shibire^{ts1}* (*shi^{ts1}*) (Kosaka and Ikeda, 1983), a temperature-sensitive mutant of the GTPase

Shibire/dynamin that blocks endocytic recycling of synaptic vesicles in flies raised at the restrictive

temperature. Co-expression of shi^{ts1} with Htt$_{ex1}$Q91-mCherry in DA1 ORNs in adult flies raised at 31°C

did not affect numbers of Htt$_{ex1}$Q91-mCherry aggregates in the DA1 glomerulus (Fig. 3A and B), but,

surprisingly, strongly enhanced formation of Htt$_{ex1}$Q25-GFP aggregates in DA1 PN dendrites (Fig. 3A

and C) compared to control flies expressing LacZ. Likewise, co-expression of Htt$_{ex1}$Q91-mCherry with

tetanus toxin light chain (TeTxLC), which inhibits SNARE-mediated fusion of synaptic vesicles with

presynaptic membranes (Sweeney et al., 1995), had no consistent effect on Htt$_{ex1}$Q91-mCherry

aggregate numbers in ORN axons (Fig. 3D and E), but strongly increased numbers of Htt$_{ex1}$Q25-GFP

aggregates in the DA1 glomerulus (Fig. 3D and F). To manipulate neuronal activity by an alternative

approach, we co-expressed the heat-activated *Drosophila* transient-receptor potential A (dTrpA)

channel (Hamada et al., 2008) with Htt$_{ex1}$Q91-mCherry to thermogenetically stimulate DA1 ORNs. In

adult flies shifted from 25°C to 31°C upon eclosion, dTrpA-mediated activation of Htt$_{ex1}$Q91-mCherry-

expressing DA1 ORNs slightly increased Htt$_{ex1}$Q91-mCherry aggregate numbers (Fig. 3G and H), but

substantially decreased formation of Htt$_{ex1}$Q25-GFP aggregates in the DA1 glomerulus (Fig. 3G and I)

compared to control flies expressing LacZ. Together, these results suggest that the ability of

presynaptic ORNs to fire is inversely correlated with prion-like transfer of Htt$_{ex1}$Q91-mCherry

aggregates from ORNs to PNs, perhaps reflecting enhanced aggregate transfer across dysfunctional

synapses characteristic of HD and other neurodegenerative diseases.

Our results using shi^{ts1} and TeTxLC to block DA1 ORN activity contrast with previous reports

showing that spreading of RFP-Htt$_{ex1-12}$Q138 aggregates from ORN axons to non-synaptically

connected neurons was slowed when endocytosis or exocytosis was impaired in ORNs (Babcock and

Ganetzky, 2015) or that botulinum toxin inhibited spreading of mHtt$_{ex1}$ from R6/2 mouse brain slices to

functionally-connected human neurons (Pecho-Vrieseling et al., 2014). We therefore wondered whether

blocking Shibire-mediated endocytosis in ORNs might create a more favorable environment for

aggregate transfer across endogenous synapses in our HD model. To test this, we first quantified

Htt$_{ex1}$Q91-mCherry-expressing DA1 ORN axonal surfaces using membrane-targeted GFP (mCD8-GFP)

(Fig. 4A), a tool widely used to label neuronal membranes (Lee and Luo, 1999; Mosca and Luo, 2014)

and quantify neuron abundance (Burr et al., 2014; MacDonald et al., 2006) in fly brains. Three-

dimensional image analysis (Fig. 4B) revealed that the intensity and volume occupied by ORN axonal

membranes in the DA1 glomerulus were increased ~2-fold in shi[ts1]-expressing flies raised at 31°C as

compared to controls (Fig. 4A-C). Because numbers of Htt$_{ex1}$Q91-mCherry aggregates were unaffected

by shi[ts] expression in DA1 ORNs (Fig. 3B), this finding suggests that blocking endocytosis increases

the surface area of DA1 ORN axons, perhaps reflecting a homeostatic compensatory mechanism

(Davis, 2013) that could create additional exit sites for Htt$_{ex1}$Q91-mCherry aggregates.

We have previously reported that phagocytic glia engulf Htt$_{ex1}$Q91-mCherry aggregates from

DA1 ORNs, resulting in prion-like conversion of Htt$_{ex1}$Q25-YFP aggregates in the glial cytoplasm

(Pearce et al., 2015). To test whether blocking Shibire-mediated endocytosis in ORNs could also

enhance ORN-to-glia transfer of Htt$_{ex1}$Q91-mCherry aggregates, we co-expressed shi[ts1] with Htt$_{ex1}$Q91-

mCherry in DA1 ORNs and monitored formation of Htt$_{ex1}$Q25-YFP aggregates in the glial cytoplasm.

Similar to effects of shi[ts1] on trans-synaptic Htt$_{ex1}$Q91-mCherry aggregate transfer, blocking Shibire-

mediate endocytosis in DA1 ORNs increased Htt$_{ex1}$Q25-YFP aggregate formation in glia without

affecting Htt$_{ex1}$Q91-mCherry aggregate numbers (Fig. 4D, E, and F). These findings suggest that

blocking endocytosis in Htt$_{ex1}$Q91-mCherry-expressing ORNs alters axonal membrane turnover and

stimulates prion-like transfer of Htt$_{ex1}$Q91-mCherry aggregates from ORNs to both PNs and phagocytic

glia, the latter of which we previously showed is mediated by Draper-dependent glial phagocytosis

(Pearce et al., 2015).

Glial Draper mediates ORN-to-PN transfer and alters morphology of neuronal mHtt$_{ex1}$ aggregates

The parallel effects of shi[ts1]-mediated endocytic inhibition on transfer of Htt$_{ex1}$Q91-mCherry

aggregates from DA1 ORNs to DA1 PNs and to glia suggest that mHtt$_{ex1}$ aggregate spreading between

these cell types is coordinated. We previously reported that ORN-to-glia transfer of Htt$_{ex1}$Q91-mCherry

aggregates is strictly dependent on Draper (Pearce et al., 2015), a scavenger receptor responsible for

phagocytic engulfment and clearance of neuronal debris in the fly CNS and other tissues (Etchegaray

et al.; Han et al., 2014; Hoopfer et al., 2006; MacDonald et al., 2006). Therefore, we sought to

determine whether Draper-dependent phagocytic glia might play a role in transferring mHtt$_{ex1}$

255 aggregates from ORNs to PNs. To test this, we quantified numbers of $Htt_{ex1}Q91$-mCherry and

256 $Htt_{ex1}Q25$-GFP aggregates in DA1 ORN axons and PN dendrites, respectively, in animals heterozygous

257 or homozygous for the *draper(drpr)$^{\Delta 5}$* null mutation (Freeman et al., 2003). As we have previously

258 reported (Pearce et al., 2015), *drpr* knockout increased steady-state numbers of $Htt_{ex1}Q91$-mCherry

259 aggregates in DA1 ORN axons (Fig. 5A and B), but strikingly, abolished formation of $Htt_{ex1}Q25$-GFP

260 aggregates in PNs (Fig. 5A and C). *Drpr* knockout also completely blocked formation of $Htt_{ex1}Q25$-GFP

261 aggregates in PNs when shi^{ts1} was co-expressed with $Htt_{ex1}Q91$-mCherry in ORNs (Fig. 5-figure

262 supplement 1), confirming that enhanced transfer of $Htt_{ex1}Q91$-mCherry aggregates from shi^{ts1}-

263 expressing ORNs occurs via the same Draper-dependent mechanism. To rule out the possibility that

264 *GH146-Gal4* drives expression of $Htt_{ex1}Q25$-GFP in some Draper+ glia, we showed that repo-Gal80-

265 mediated inhibition of Gal4 in all glia had no effect on $Htt_{ex1}Q91$-mCherry or $Htt_{ex1}Q25$-GFP aggregate

266 numbers (Fig. 5-figure supplement 2A, C, and D). Moreover, expression of *drpr*-specific siRNAs in PNs

267 had no effect on $Htt_{ex1}Q25$-GFP aggregate formation, consistent with glia being the sole source of *drpr*

268 expression in the CNS (Fig. 5-figure supplement 2B, C, and D). We also did not observe co-localization

269 between ORN-derived $Htt_{ex1}Q91$-mCherry aggregates and GFP-tagged versions of Lamp1

270 (Pulipparacharuvil et al., 2005) or Atg8a (Juhász et al., 2008) in glia (Fig. 5-figure supplement 3),

271 suggesting that neuronal $Htt_{ex1}Q91$-mCherry aggregates do not seed $Htt_{ex1}Q25$-GFP in glial lysosomes

272 or autophagosomes.

273  These results point to a central role for glial Draper in $Htt_{ex1}Q91$-mCherry aggregate transfer

274 between multiple cell types in the fly CNS, but how phagocytic glia could mediate spreading of

275 $Htt_{ex1}Q91$-mCherry aggregates across neuronal synapses was not clear. Intriguingly, the average

276 volume of $Htt_{ex1}Q91$-mCherry aggregates in the DA1 glomerulus significantly increased from 2.201 $\pm$

277 0.044 μm^3 in wild-type animals to 2.431 $\pm$ 0.038 μm^3 in *drpr*-knockout animals (p < 0.0001 by Student's

278 *t*-test), and the relative frequency of two aggregate subpopulations shifted between these groups: in the

279 absence of *drpr*, the proportion of a smaller-sized subpopulation ("A", ~0.1-0.5 μm^3) decreased while a

280 larger-sized subpopulation ("B", ~2.5-4.0 μm^3) increased in abundance (Fig. 5D). The decreased

281 frequency of subpopulation A aggregates in *drpr$^{\Delta 5/\Delta 5}$* flies was especially interesting because these

aggregates closely correlate in volume with those of seeding-competent Htt$_{ex1}$Q91-mCherry aggregates

in PNs (Fig. 2F) or in glia (Pearce et al., 2015). Thus, these results suggest that phagocytic glia

mediate transfer of Htt$_{ex1}$Q91-mCherry aggregates across neuronal synapses, at least in part by

promoting the formation of a subpopulation of smaller, transmission-competent neuronal mHtt$_{ex1}$

aggregates.

Caspase activation in ORNs regulates trans-synaptic transfer of mHtt$_{ex1}$ aggregates

289   Phagocytic glia engulf injured or degenerating neuronal processes and apoptotic cell corpses by

recognizing "eat me" signals exposed on these debris (Wilton et al., 2019). Thus, we asked whether

Htt$_{ex1}$Q91-mCherry aggregate-expressing DA1 ORNs might expose cell surface signals that could drive

Draper-dependent transfer of aggregates to DA1 PNs. mHtt expression induces caspase-dependent

cell death pathways in fly and mammalian models of HD (Ahmed et al., 2014), and we observed that in

drpr-deficient flies, *Drosophila* caspase-1 (Dcp-1) immunoreactivity was increased in ORN axons

expressing Htt$_{ex1}$Q91-GFP as compared to controls expressing Htt$_{ex1}$Q25-GFP (Fig. 6A and B). In

addition, co-expression of the viral effector caspase inhibitor p35 (Hay et al., 1994) with Htt$_{ex1}$Q91-

mCherry in DA1 ORNs inhibited formation of Htt$_{ex1}$Q25-GFP aggregates in DA1 PNs (Fig. 6C and E,

squares), indicating that caspase activation in presynaptic ORNs is required for mHtt$_{ex1}$ aggregate

transfer to postsynaptic PNs. Caspase inhibition also increased numbers of Htt$_{ex1}$Q91-mCherry

aggregates in the DA1 glomerulus (Fig. 6C and D, *squares*), phenocopying effects seen in *drpr*$^{Δ5/Δ5}$

brains (Fig. 5A and B). Conversely, co-expression of p35 with Htt$_{ex1}$Q25-GFP in PNs did not affect

numbers of Htt$_{ex1}$Q91-mCherry and Htt$_{ex1}$Q25-GFP aggregates in the DA1 glomerulus (Fig. 6D and E,

circles), indicating that caspase-dependent signaling in PNs does not regulate mHtt$_{ex1}$ aggregate

spreading across ORN-PN synapses. Together, these results suggest that Htt$_{ex1}$Q91-mCherry-induced

caspase activation in ORN axons signals for engulfment and trans-synaptic transfer of Htt$_{ex1}$Q91-

mCherry aggregates via Draper-dependent phagocytic glia.

mHtt$_{ex1}$ aggregates transfer from ORNs to PNs via the glial cytoplasm

The strict requirement for Draper in prion-like transfer of Htt$_{ex1}$Q91-mCherry aggregates across

ORN-PN synapses can be explained by two non-mutually exclusive models. In one model (Fig. 7A,

route 1), Htt$_{ex1}$Q91-mCherry aggregates spread from presynaptic ORNs to postsynaptic PNs via a glial

cytoplasmic intermediate. This model is consistent with our previous finding that phagocytosed

neuronal Htt$_{ex1}$Q91-mCherry aggregates enter the glial cytoplasm and effect prion-like conversion of

Htt$_{ex1}$Q25-YFP (Pearce et al., 2015). Alternatively (Fig. 7A, *route 2*), Draper-dependent phagocytic glia

could sculpt the synaptic environment in a way that promotes transfer of mHtt$_{ex1}$ aggregates directly

from ORN axons to PN dendrites. To distinguish between these models, we generated transgenic flies

that use three binary expression systems (QF-QUAS, Gal4-UAS, and LexA-LexAop) to independently

express a uniquely-tagged Htt$_{ex1}$ transgene in each of three cell populations: Htt$_{ex1}$Q91-mCherry in DA1

ORNs, Htt$_{ex1}$Q25-3xHA in repo+ glia, and Htt$_{ex1}$Q25-YFP in GH146+ PNs (Fig. 7A). If Htt$_{ex1}$Q91-

mCherry aggregates formed in presynaptic ORNs travel through the glial cytoplasm to postsynaptic

PNs, the same Htt$_{ex1}$Q91-mCherry aggregate seeds should template the aggregation of Htt$_{ex1}$Q25-

3xHA in glia and Htt$_{ex1}$Q25-YFP in PNs, resulting in appearance of triple-labeled

mCherry+/3xHA+/YFP+ puncta in the DA1 glomerulus (Fig. 7A, *route 1*). A transient subpopulation of

double-labeled mCherry+/3xHA+ intermediates might also be observed, but these would be converted

to mCherry+/3xHA+/YFP+ aggregates over time. Alternatively, if Htt$_{ex1}$Q91-mCherry aggregates do not

access the glial cytoplasm *en route* to PNs, only double-labeled mCherry+/3xHA+ and mCherry+/YFP+

aggregates would appear in the DA1 glomerulus and persist over time (Fig. 7A, *route 2*).

Transgenic flies expressing these differentially-tagged Htt$_{ex1}$ transgenes in three distinct CNS

cell types were generated, and no overlapping expression of any two transgenes was detected by

confocal microscopy (Fig. 7B and C). Htt$_{ex1}$Q91-mCherry expression was low and largely diffuse in

young (0-1 day-old) adult flies, but became more punctate as the flies aged (Fig. 7B-D).

mCherry+/3xHA+ puncta representing Htt$_{ex1}$Q91-mCherry aggregates that had transferred from ORNs

to glia began to appear in 2 day-old adults, increased in number during the next 24 hr, but thereafter

stabilized as the flies aged (Fig. 7C and E, *white bars*). Strikingly though, small numbers of

mCherry+/3xHA+/YFP+ puncta appeared in 2 day-old adults and steadily increased in abundance as

the flies aged (Fig. 7C and E, *black bars*), suggesting that these triple-labeled aggregates arose from

$Htt_{ex1}Q91$-mCherry seeds that originated in DA1 ORN axons and transited through the glial cytoplasm

before reaching the cytoplasm of DA1 PN dendrites. Notably, mCherry+/YFP+ puncta representing

aggregates that directly transferred from DA1 ORNs to DA1 PNs were absent in these brains, arguing

against direct transfer of $Htt_{ex1}Q91$-mCherry aggregates across ORN-PN synapses. Importantly, when

drpr was depleted specifically in glia by RNAi in these flies, only mCherry+ puncta were visible in the

DA1 glomerulus of 7 day-old adults (Fig. 7C, *far right*), confirming that glial Draper mediates aggregate

transfer unidirectionally from ORNs to glia to PNs. Taken together, these findings suggest that Draper-

dependent glia act as obligatory intermediates in prion-like spreading of $mHtt_{ex1}$ aggregates from

presynaptic ORNs to postsynaptic PNs in *Drosophila* brains.

DISCUSSION

      The hypothesis that prion-like mechanisms underlie the progression of protein aggregate pathology in HD and other neurodegenerative diseases has gained considerable support, but the mechanisms underlying cell-to-cell aggregate transmission *in vivo*, the influences of cell- and region-specific vulnerability, and the relevance of aggregate spreading to disease progression remain unclear. In this study, we report that prion-like transfer of mHtt$_{ex1}$ aggregates from presynaptic ORNs to postsynaptic PNs occurs via Draper-dependent phagocytic glia in transgenic *Drosophila* brains. ORN-to-PN transmission of mHtt$_{ex1}$ aggregates was enhanced by blocking presynaptic endocytosis and exocytosis and slowed by thermogenetically stimulating ORNs, suggesting an inverse relationship between presynaptic neuronal activity and mHtt$_{ex1}$ aggregate transmission across synapses. Endocytic inhibition in ORNs also enhanced ORN-to-glia mHtt$_{ex1}$ transfer in the antennal lobe, suggesting that neuronal silencing stimulates Draper-dependent phagocytosis of aggregate-containing ORN axons and presumably subsequent transfer to postsynaptic PNs. We also find that trans-synaptic mHtt$_{ex1}$ aggregate transfer requires caspase activity in ORNs and is completely blocked in *drpr*-knockout animals, suggesting that Draper-dependent glia recognize caspase-dependent "eat me" signals (e.g. exposed phosphatidylserine) on the surface of mHtt$_{ex1}$-expressing axons to initiate phagocytic engulfment. Remarkably, a portion of mHtt$_{ex1}$ aggregates originating in ORN axons effect prion-like conversion of cytoplasmic wtHtt$_{ex1}$ proteins first in glia and then in PN dendrites, indicating that aggregates transfer from presynaptic ORNs to postsynaptic PNs through a Draper-dependent glial cytoplasmic intermediate. These findings expand our understanding of the beneficial and harmful effects of phagocytic glia in neurodegeneration—while phagocytic engulfment and lysosomal degradation clear neurotoxic aggregates and cell corpses from the brain, escape of even a few engulfed aggregates from the phagolysosomal system generates a glial cytoplasmic reservoir of prion-like aggregate species that can propagate further in the CNS.

      Though HD is a monogenic disorder and the Htt gene is expressed ubiquitously, it is plausible that prion-like spreading of mHtt aggregates could drive progression of HD pathology from more vulnerable regions of the brain to those less susceptible to mHtt aggregation. We find that mHtt$_{ex1}$

aggregate transmission across ORN-PN synapses occurs selectively in the anterograde direction and

is inversely correlated with synaptic activity, suggesting that aggregate spreading is enhanced across

dysfunctional synapses, an early molecular feature of HD and other neurodegenerative diseases.

Intriguingly, the earliest pathological changes in HD patient brains involve loss of cortical inputs to the

striatum, where more prominent pathological findings (i.e., mHtt aggregate accumulation and massive

loss of medium spiny neurons) appear in later stages of disease (Reiner and Deng, 2018). Moreover,

selective silencing of mHtt expression in both the cortex and striatum produced the most beneficial

effects on striatal degeneration and motor phenotypes in BACHD mice (Wang et al., 2014), suggesting

that mHtt-induced toxicity in presynaptic cortical regions may play a major role in early HD

neuropathogenesis. Our results demonstrating increased aggregate transfer from activity-impaired

ORN axons conflict with reports from other groups showing that neuronal silencing slows cell-to-cell

spreading of mHtt in fly brains (Babcock and Ganetzky, 2015) or mixed-species *ex vivo* co-cultures

(Pecho-Vrieseling et al., 2014). This discrepancy could be due to construct- or cell type-specific effects,

or could possibly reflect different mechanisms regulating synaptic or non-synaptic aggregate

transmission in the brain. Our HD model differs from these others in that we monitored mHtt aggregate

spreading across endogenous synapses formed under physiological conditions. Thus, our findings

uncovering unexpected roles for presynaptic silencing and phagocytic glia in aggregate transfer likely

reflect intimate relationships that glia make with neuronal synapses in intact brains.

Central roles for phagocytic glia in neurodegeneration are becoming increasingly recognized as

genome-wide association studies and transcriptomic analyses identify glial genes associated with

increased risk or development of disease. For example, rare variants in *TREM2*, a gene that encodes a

microglial phagocytic receptor, are associated with increased risk of AD, FTD, and PD, and loss of

TREM2 function exacerbates Aβ-, tau-, and α-synuclein-associated neurotoxicity (Griciuc et al.; Guo et

al., 2019; Leyns et al., 2019; Zhao et al., 2018). Reactive glia alter their morphology and gene

expression profiles in response to diverse types of injury to promote neuronal survival by releasing

trophic factors and clearing debris (Hammond et al., 2018). Phagocytic glia likely play some beneficial

roles in neurodegeneration, since pathogenic aggregates accumulate when glial phagocytosis is

disrupted (Asai et al., 2015; Hong et al., 2016; Pearce et al., 2015; Ray et al., 2017; Wilton et al., 2019),

and administration of antibodies targeting pathological Aβ, tau, and α-synuclein proteins reduces

aggregate accumulation and spreading in mouse brains (Funk et al., 2015; Masliah et al., 2005; Tran et

al., 2014; Yanamandra et al., 2013). However, chronically active and/or dysfunctional glia constitutively

release neuroinflammatory signals and acquire behaviors that exacerbate neuronal damage. For

example, hyperactivate microglia excessively phagocytose synapses in pre-plaque AD mouse brains

(Hong et al., 2016) and induce the formation of neurotoxic A1 astrocytes that persist in many

neurodegenerative diseases (Liddelow et al., 2017; Yun et al., 2018) and during aging (Clarke et al.,

2018). Our findings thus add new insights to recognizing glia as double-edged players in

neurodegeneration and suggest that rebalancing the protective and harmful effects of phagocytic glia

could be a promising new therapeutic strategy.

Our data support a model in which Draper-dependent phagocytic glia engulf mHtt$_{ex1}$ aggregates

or portions of mHtt$_{ex1}$-expressing ORN axons exposing caspase-dependent signals to promote neuron

survival; however, this same process leads to prion-like aggregate spreading to glial and PN

cytoplasms. We provide several lines of evidence to support our conclusion that phagocytosed mHtt$_{ex1}$

aggregates access the cytoplasm of glia and then PNs and to argue against seeding of wtHtt$_{ex1}$ in the

interstitial space. The strict requirement for Draper in ORN-to-glia and ORN-to-glia-to-PN aggregate

transfer indicates that neuronal mHtt$_{ex1}$ and glial wtHtt$_{ex1}$ proteins physically interact during or following

phagocytic engulfment, requiring breaching of one or more membrane barriers that normally exclude

phagocytosed material from the glial cytoplasm. In support of this, prion-like aggregates formed by α-

synuclein, tau, and mHtt$_{ex1}$ have the capacity to rupture cell surface or endolysosomal membranes to

directly access the cytoplasmic compartment (Chen et al., 2019; Falcon et al., 2018; Flavin et al., 2017;

Zeineddine et al., 2015). It is also possible that phagocytosis becomes dysregulated in diseased or

aged brains. Indeed, extracellular Aβ or α-synuclein fibrils accumulate inside phagocytic microglia over

time (Chung et al., 1999; Frackowiak et al., 1992), and the ability of microglia to degrade phagocytosed

material declines with age (Bliederhaeuser et al., 2016; Tremblay et al., 2012). Defects in lysosomal

clearance could also mediate aggregate escape, supported by a recent finding that proteins unable to

be efficiently digested by autophagolysosomes are excreted by cells in a model of the polyQ disease

dentatorubral-pallidoluysian atrophy (Baron et al., 2017). Thus, impairments in phagosome maturation

or degradation of engulfed material by the lysosome could increase opportunities for aggregate escape

and cell-to-cell spreading.

Interestingly, we found that Draper knockout is associated with loss of smaller-sized, seeding-

competent mHtt$_{ex1}$ aggregates in ORN axons, suggesting that the Draper phagocytic pathway regulates

neuronal mHtt$_{ex1}$ aggregate size. How phagocytic glia could alter the morphology of neuronal mHtt$_{ex1}$

aggregates is not immediately clear, but it is possible that dynamin-mediated fission or SNARE-

mediated fusion events during phagosome maturation could generate aggregate fragments that fall

below an upper size limit for entry to the cytoplasm and/or can directly penetrate intracellular

membranes. Draper could also selectively deplete a subpopulation of larger neuronal aggregates from

which smaller transmissible species are produced. Interestingly, seeding propensity is correlated with

smaller mHtt$_{ex1}$ aggregate size in other HD models (Ast et al., 2018; Chen et al., 2001), and secondary

nucleation processes initiated by aggregate fragmentation are thought to drive amplification of prion-like

particles in many neurodegenerative diseases (Knowles et al., 2014).

One of the most striking results we report here is that all induced wtHtt$_{ex1}$ aggregates in PNs

arise from ORN-derived mHtt$_{ex1}$ aggregates that escaped from Draper-dependent phagocytosis into the

glial cytoplasm. How could phagocytosed mHtt$_{ex1}$ aggregates transfer from the glial cytoplasm to the

cytoplasm of postsynaptic PNs? Several mechanisms already proposed for prion-like aggregate

spreading could support inter-cytoplasmic transfer between these cells, including transport through

membrane-protected compartments such as extracellular vesicles or tunneling nanotubes, secretion or

passive release of aggregates from dying cells, and/or direct penetration of lipid bilayers (Brundin et al.,

2010; Davis et al., 2018; Vaquer-Alicea and Diamond, 2019). However, none of these mechanisms

discriminate between transfer of aggregates directly from ORNs to PNs or from glia to PNs, suggesting

that escape of aggregates from phagocytic glia must involve a mechanism that is unique to these cells

in the disease state. In support of this, pathological tau transfers between cultured cells or anatomically-

connected regions of mouse brains via exosomes secreted by microglia (Asai et al., 2015), suggesting

that phagocytic glia could facilitate prion-like spreading via exosomal release after engulfed neuronal

aggregates have reached the glial cytoplasm. Further studies are needed to elucidate the role of

phagocytic glia in aggregate propagation and the mechanistic details for intracellular aggregate

propagation between diverse cell types in the CNS.

Our findings support a mechanism in which glia engulf $mHtt_{ex1}$ aggregates from ORN axons

without phagocytosing elements of PNs because (i) $mHtt_{ex1}$ aggregate transmission only occurred in

the anterograde direction across ORN-PN synapses, (ii) $wtHtt_{ex1}$ aggregates were observed in PN cell

bodies, (iii) $mHtt_{ex1}$ aggregates did not co-localize with markers of lysosomes or autophagosomes in

glia, and (iv) caspase inhibition in ORNs and not in PNs slowed aggregate transfer. In developing

mouse brains, pruned presynaptic structures are observed within microglia largely in the absence of

postsynaptic markers (Schafer et al., 2012; Weinhard et al., 2018), suggesting that glia selectively

"nibble" presynaptic components through a process known as trogocytosis. Interestingly, mammalian

astrocytes expressing the Draper homolog MEGF10 and complement-dependent microglia

preferentially engulf less active, "weaker" synapses to refine neural circuits in developing and adult

mouse brains (Chung et al., 2013; Schafer et al., 2012), and aberrant activation of these pathways

could contribute to early synaptic loss in neurodegenerative diseases (Hong et al., 2016). Our finding

that transfer of $mHtt_{ex1}$ aggregates to glia and PNs was enhanced when ORNs were silenced suggests

that phagocytic glia become activated to clear dysfunctional, aggregate-containing presynaptic

structures that accumulate early in HD and other in neurodegenerative diseases, ultimately leading to

propagation of prion-like aggregate species along synaptic connections in the brain.

In summary, our data demonstrate that $mHtt_{ex1}$ aggregates undergo prion-like transfer between

synaptically-connected neurons via Draper-dependent glia, raising the intriguing possibility that

phagocytic glia could act as a mobile vehicle to spread neuronal aggregate pathology through the CNS.

These findings have important implications for understanding complex relationships between

aggregate-induced cytotoxicity and communication between neuronal and glial cell types in health and

disease. Deciphering the mechanisms that underlie helpful and harmful phagocytic glial cell functions in

the degenerating brain will help to reveal new ways in which glia can be targeted therapeutically in

neurodegenerative diseases.

MATERIALS AND METHODS

Fly husbandry

All fly stocks and crosses were raised on standard cornmeal/molasses media at 25°C, ~50% relative humidity, and on a 12 hr light/12 hr dark cycle, unless otherwise noted. The following drivers were used to genetically access different cell populations in the fly CNS: *elav[C155]-Gal4* (Lin and Goodman, 1994), *Or67d-QF* (Liang et al., 2013), *GH146-Gal4* (Stocker et al., 1997), *GH146-QF* (Potter et al., 2010), and *GH146-LexA::GAD* (Lai et al., 2008) (a kind gift from Tzumin Lee, Janelia Farms), *repo-Gal4* (Sepp et al., 2001), and *Or83b-Gal4* (Kreher et al., 2005). Transgenic flies previously described by our lab include *QUAS-Htt$_{ex1}$Q91-mCherry*, *QUAS-Htt$_{ex1}$Q25-mCherry*, *UAS-Htt$_{ex1}$Q91-mCherry*, *UAS-Htt$_{ex1}$Q25-GFP*, *UAS-Htt$_{ex1}$Q25-YFP*, and *UAS-GFP* transgenes inserted at the attP3 (1st chromosome) and/or attP24 (2nd chromosome) ϕC31 integration sites (Pearce et al., 2015). Other transgenic flies not generated in this study include: *QUAS-nucLacZ* lines #7 and 44 (BDSC #30006 and 30007)*, QUAS-shi^{ts1}* lines #2, 5, and 7 (BDSC #30010 and 30012; kind gift from C. Potter, Johns Hopkins School of Medicine), *QUAS-TeTxLC* lines #4c and 9c and *QUAS-dTrpA* lines #5, 6, and 7 (kind gifts from O. Riabinina and C. Potter, Johns Hopkins), *QUAS-mCD8-GFP* line #5J (BDSC #30002), *QUAS-p35* (a kind gift from H. Steller, Rockefeller University), *UAS-LacZ* (BDSC #8529), *UAS-p35* (BDSC #5072)*, QUAS-Gal80* (BDSC #51948), *UAS-QS* (BDSC #30033), *repo-Gal80* (a kind gift from T. Clandinin, Stanford University), *UAS-FFLuc.VALIUM1* (BDSC #35789), *UAS-GFP-Lamp1* (BDSC #42714), *UAS-Atg8a-GFP* (BDSC #52005), and *UAS-Draper^{RNAi}* and *drpr^{Δ5}* mutant flies (kind gifts from M. Freeman, Vollum Institute). Genotypes for all flies used in this study are listed in Table 1.

Cloning and transgenesis

pUASTattB(Htt$_{ex1}$Q25-GFP) and pUASTattB(Htt$_{ex1}$Q25-3xHA) plasmids were generated by PCR amplification of tagged Htt$_{ex1}$ cDNAs from the pcDNA3 vector backbone and cloned into pUASTattB via XhoI and XbaI restriction sites. cDNA for mRFP-Htt$_{ex1-12}$Q138 (a kind gift from T. Littleton, MIT) (Weiss et al., 2012) was subcloned into the pQUASTattB plasmid (Riabinina et al., 2015) using EcoRI and XbaI restriction sites. Htt$_{ex1}$Q25-YFP cDNA was amplified by PCR from the pUASTattB(Htt$_{ex1}$Q25-YFP)

plasmid (Pearce et al., 2015) and subcloned into the EcoRI site in pLOT downstream of LexA-

responsive LexAop DNA sequences. Plasmids were microinjected into embryos containing the attP3 X

chromosome (for UAS-Htt$_{ex1}$Q25-GFP), attP3B or attP8 X chromosome (for QUAS-mRFP-Htt$_{ex1}$-

$_{12}$Q138), or attP24 2nd chromosome (for UAS-Htt$_{ex1}$Q25-3xHA or LexAop-Htt$_{ex1}$Q25-YFP) ϕC31

integration sites either in-house or at BestGene, Inc (Chino Hills, CA).

Drosophila brain dissection, immunostaining, and imaging

Adult fly brains were dissected, fixed, and stained and/or imaged as previously described

(Pearce et al., 2015). Briefly, brains were dissected in ice-cold phosphate-buffered saline containing

either 0.03% Triton X-100 (PBS/0.03T, when intrinsic fluorescence of expressed FP-fusions was to be

imaged) or 0.3% Triton X-100 (PBS/0.3T, when using indirect immunofluorescence to detect protein

expression). We typically imaged using intrinsic GFP/YFP/mCherry fluorescence when Htt-FP fusions

were bright enough to detect on the confocal to reduce sample preparation time needed prior to

imaging. Dissected brains were transferred to microfuge tubes containing PBS/T + 4%

paraformaldehyde fixative solution on ice and then fixed in the dark at room temperature (RT) for 5 min

(when imaging intrinsic fluorescence) or 20 min (when using immunofluorescence). For direct

fluorescence imaging, brains were washed in PBS/0.03T buffer several times before incubation in

Slowfade Gold Antifade Mountant (Invitrogen, Carlsbad, CA) for at least 1 hr at 4°C in the dark. For

immunostaining, brains were washed several times in PBS/0.3T, and then blocked in PBS/0.3T

containing 5% normal goat serum (Lampire Biological Laboratories, Pipersville, PA) for 30min at RT,

followed by incubation in primary antibodies diluted in blocking solution and incubation for 24-72 hr at

4°C in the dark. Brains were then washed in PBS/0.3T several times at RT and incubated in secondary

antibodies diluted in blocking solution for 20-24 hr at 4°C in the dark. Following another set of washes

in PBS/0.3T at RT, the brains were incubated in Slowfade mountant for at least 16 hr at 4°C in the dark.

Primary antibodies used in this study include rabbit anti-DsRed (1:2000; #632496; Takara Bio

USA, Inc., Mountain View, CA), rabbit anti-mCherry (1:500; #PA5-34974; Invitrogen, Carlsbad, CA),

chicken anti-GFP (1:500; #GFP-1020; Aves Labs, Tigard, OR), chicken anti-GFP (1:1000; #ab13970;

Abcam, Cambridge, UK), rat anti-HA (1:100; clone 3F10; #11867423001; Roche, Basel, Switzerland),

rabbit anti-cleaved Dcp-1 (1:100; #9578S; Cell Signaling Technology, Danvers, MA), and mouse anti-

Bruchpilot (1:100; clone nc82; Developmental Studies Hybridoma Bank, Iowa City, IA). Secondary

antibodies used include FITC-conjugated donkey anti-chicken (1:200; #703-095-155; Jackson Immuno

Research Labs, West Grove, PA) and AlexaFluor 488, 568, or 647-conjugated goat anti-chicken, anti-

mouse, anti-rabbit, or anti-rat IgGs (1:250 each; Invitrogen, Carlsbad, CA).

Confocal microscopy and image analysis and quantification

Brains were bridge-mounted in Slowfade mountant on glass microscopy slides overlayed with

#1.5 coverglass (22 x 22 mm), and edges were sealed using clear nail polish. All images were collected

on Leica SP8 or Zeiss LSM510 laser-scanning confocal systems equipped with 405 nm, 488 nm, 561

nm, and 633 nm lasers and 40X 1.3NA or 63X 1.4NA oil objective lenses. Leica LAS X or Zeiss ZEN

software programs were used to establish optimal settings during each microscopy session and to

collect optical z-slices of whole-mounted brain samples with Nyquist sampling criteria.

Raw confocal data were analyzed in 2D using ImageJ/FIJI software (NIH, Bethesda, MD) or in

3D using Imaris image analysis software (Bitplane, Zürich, Switzerland). Most images were collected at

optical zoom $\geq$3 to capture the DA1 glomerulus, and the image bounding box served as a region of

interest. In Imaris, individual Htt aggregates were identified using the "Surfaces" algorithm (surface

detail set to 0.18 μm and background subtraction at 0.45 μm). "Split touching objects" option was

selected, and seed point diameter was set to 0.8 μm. Thresholding was adjusted manually for each

image, typically within ~5% among samples in the same experiment, to maximally quantify punctate

(i.e., aggregate) objects while minimizing background. While these settings permitted accurate counting

of most heterogeneously-sized Htt_{ex1}Q91-mCherry aggregates in the DA1 glomerulus, some larger

aggregates were aberrantly double-counted or smaller aggregates in close proximity were

inappropriately merged. In these rare cases, individual objects were manually unified or split using the

Imaris software prior to recording total aggregate counts for each image. Because Htt_{ex1}Q25-

G/YFP/3xHA aggregates present in DA1 PNs or glia were harder to discern from surrounding diffuse

Htt$_{ex1}$Q25 signal, these puncta were manually counted by scanning individual z-slices to identify co-

localized aggregates. To measure DA1 ORN axon volume in the DA1 glomerulus, mCD8-GFP signal

was detected using the "Surfaces" function in Imaris (surface detail = 0.2 μm, and background

subtraction = 3 μm). Individual aggregate or surface detail measurements (e.g. volume, intensity) were

calculated using Imaris, and the data were exported to Excel (Microsoft Corporation, Redmond, WA) or

Prism (GraphPad Software, San Diego, CA) software programs for further analysis.

FRET analysis

Htt$_{ex1}$Q25-GFP and Htt$_{ex1}$Q91-mCherry co-localized aggregates were analyzed for FRET using

the acceptor photobleaching method as previously described (Pearce et al., 2015). Briefly, 28 individual

aggregates were analyzed by photobleaching the mCherry acceptor using a 561nm laser set at 100%

intensity and scanning until the mCherry signal was no longer detectable. Donor fluorescence

dequenching was measured by exciting GFP using a 488 nm laser set at 1% intensity before and after

acceptor photobleaching. mCherry fluorescence was also excited before and after photobleaching with

a 561 nm laser set at 1% intensity. Fluorescence emission was collected between 500 and 550 nm for

GFP and 610 and 700 nm for mCherry to generate before and after images. FRET efficiencies

(FRET$_{eff}$) were calculated after background correction using the equation $(GFP_{after}-GFP_{before})/(GFP_{after})$ x

100 and represented by a pixel-by-pixel FRET$_{eff}$ image generated using the FRETcalc plugin

(Stepensky, 2007) in FIJI/ImageJ software.

Statistical analyses

All quantified data were organized and analyzed in Excel or Prism 8 software programs.

Quantifications in graphical form are shown as mean $\pm$ SEM, except for frequency analyses, which are

displayed in histograms. Results of all statistical analyses are described in each Figure Legend.

Sample size is indicated by *n* either directly on graphical panels of figures or in the figure legend and

was selected to yield sufficient statistical power ($\geq$ 5 biological replicates for each condition; each

individual brain represents a single biological replicate). Multiple statistical comparisons were

performed using the following tests and post-hoc corrections where appropriate: Student's *t*-tests for

pairwise comparisons, or one-way or two-way ANOVA followed by Sidak's or Tukey's multiple

comparison tests, respectively, for experiments involving $\geq$3 genotypes. Results of these statistical tests

are shown on each graph and described in the Figure Legends.

ACKNOWLEDGMENTS

The authors would like to thank K. Chao, D. Luginbuhl, V. Reed, B. Temsamrit, and M. Warkala for technical assistance, G. Panning for help with analyzing volume measurements, T.R. Clandinin, M.R. Freeman, H. Krämer, T. Lee, J.T. Littleton, C.J. Potter, and O. Riabinina for kindly sharing reagents, and members of the Kopito, Luo, and Pearce labs for many valuable discussions. This work was supported by grants from the NIH (R01-DC005982 to L.L., R01-NS042842 to R.R.K, and R03-AG063295 to M.M.P.P.), the W.W. Smith Charitable Trusts (to M.M.P.P.), the Pittsburgh Foundation (UN2018-98318 to M.M.P.P.), and start-up funds from University of the Sciences (to M.M.P.P.).

AUTHOR CONTRIBUTIONS

K.M.D., O.R.D., W.M.T., and G.E.P. – Investigation, Validation, Visualization, Writing – review & editing. L.L. and R.R.K. – Conceptualization, Resources, Supervision, Writing – review & editing, Funding Acquisition. M.M.P.P. – Conceptualization, Methodology, Investigation, Validation, Formal Analysis, Visualization, Writing – original draft preparation, Writing – review & editing, Resources, Supervision, Project Administration, Funding Acquisition.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Ahmed, Z., Cooper, J., Murray, T.K., Garn, K., McNaughton, E., Clarke, H., Parhizkar, S., Ward, M.A.,

Cavallini, A., Jackson, S*., et al.* (2014). A novel in vivo model of tau propagation with rapid and

progressive neurofibrillary tangle pathology: the pattern of spread is determined by connectivity,

not proximity. Acta neuropathologica *127*, 667-683.

Asai, H., Ikezu, S., Tsunoda, S., Medalla, M., Luebke, J., Haydar, T., Wolozin, B., Butovsky, O., Kügler,

S., and Ikezu, T. (2015). Depletion of microglia and inhibition of exosome synthesis halt tau

propagation. Nature neuroscience *18*, 1584.

Ast, A., Buntru, A., Schindler, F., Hasenkopf, R., Schulz, A., Brusendorf, L., Klockmeier, K., Grelle, G.,

McMahon, B., Niederlechner, H*., et al.* (2018). mHTT Seeding Activity: A Marker of Disease

Progression and Neurotoxicity in Models of Huntington's Disease. Molecular cell *71*, 675-

688.e676.

Babcock, D.T., and Ganetzky, B. (2015). Transcellular spreading of huntingtin aggregates in the

Drosophila brain. Proceedings of the National Academy of Sciences of the United States of

America *112*, E5427-5433.

Baron, O., Boudi, A., Dias, C., Schilling, M., Nolle, A., Vizcay-Barrena, G., Rattray, I., Jungbluth, H.,

Scheper, W., Fleck, R.A*., et al.* (2017). Stall in Canonical Autophagy-Lysosome Pathways

Prompts Nucleophagy-Based Nuclear Breakdown in Neurodegeneration. Current biology *27*,

3626-3642.e3626.

Bates, G.P., Dorsey, R., Gusella, J.F., Hayden, M.R., Kay, C., Leavitt, B.R., Nance, M., Ross, C.A.,

Scahill, R.I., Wetzel, R*., et al.* (2015). Huntington disease. Nature reviews Disease primers *1*,

15005.

Bliederhaeuser, C., Grozdanov, V., Speidel, A., Zondler, L., Ruf, W.P., Bayer, H., Kiechle, M., Feiler,

M.S., Freischmidt, A., Brenner, D*., et al.* (2016). Age-dependent defects of alpha-synuclein

oligomer uptake in microglia and monocytes. Acta neuropathologica *131*, 379-391.

Boland, B., Yu, W.H., Corti, O., Mollereau, B., Henriques, A., Bezard, E., Pastores, G.M., Rubinsztein,

D.C., Nixon, R.A., Duchen, M.R*., et al.* (2018). Promoting the clearance of neurotoxic proteins in

neurodegenerative disorders of ageing. Nature reviews Drug discovery *17*, 660-688.

Braak, H., and Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes. Acta

neuropathologica *82*, 239-259.

Braak, H., Del Tredici, K., Rub, U., de Vos, R.A., Jansen Steur, E.N., and Braak, E. (2003). Staging of

brain pathology related to sporadic Parkinson's disease. Neurobiology of aging *24*, 197-211.

Brettschneider, J., Del Tredici, K., Toledo, J.B., Robinson, J.L., Irwin, D.J., Grossman, M., Suh, E., Van

Deerlin, V.M., Wood, E.M., Baek, Y*., et al.* (2013). Stages of pTDP-43 pathology in amyotrophic

lateral sclerosis. Annals of neurology *74*, 20-38.

Brundin, P., Melki, R., and Kopito, R. (2010). Prion-like transmission of protein aggregates in

neurodegenerative diseases. Nature reviews Molecular cell biology *11*, 301-307.

Burr, A.A., Tsou, W.-L., Ristic, G., and Todi, S.V. (2014). Using Membrane-Targeted Green Fluorescent

Protein To Monitor Neurotoxic Protein-Dependent Degeneration of Drosophila Eyes. Journal of

neuroscience research *92*, 1100-1109.

Chen, J.J., Nathaniel, D.L., Raghavan, P., Nelson, M., Tian, R., Tse, E., Hong, J.Y., See, S.K., Mok,

S.A., Hein, M.Y*., et al.* (2019). Compromised function of the ESCRT pathway promotes

endolysosomal escape of tau seeds and propagation of tau aggregation. The Journal of

biological chemistry. Pii: jbc.RA119.009432.

Chen, S., Berthelier, V., Yang, W., and Wetzel, R. (2001). Polyglutamine aggregation behavior in vitro

supports a recruitment mechanism of cytotoxicity. Journal of molecular biology *311*, 173-182.

Chung, H., Brazil, M.I., Soe, T.T., and Maxfield, F.R. (1999). Uptake, degradation, and release of

fibrillar and soluble forms of Alzheimer's amyloid beta-peptide by microglial cells. The Journal of

biological chemistry *274*, 32301-32308.

Chung, W.-S., Clarke, L.E., Wang, G.X., Stafford, B.K., Sher, A., Chakraborty, C., Joung, J., Foo, L.C.,

Thompson, A., Chen, C*., et al.* (2013). Astrocytes mediate synapse elimination through

MEGF10 and MERTK pathways. Nature *504*, 394-400.

Clarke, L.E., Liddelow, S.A., Chakraborty, C., Munch, A.E., Heiman, M., and Barres, B.A. (2018).

Normal aging induces A1-like astrocyte reactivity. Proceedings of the National Academy of

Sciences of the United States of America *115*, E1896-e1905.

Clyne, P.J., Warr, C.G., Freeman, M.R., Lessing, D., Kim, J., and Carlson, J.R. (1999). A novel family

of divergent seven-transmembrane proteins: candidate odorant receptors in Drosophila. Neuron

22, 327-338.

Costanzo, M., Abounit, S., Marzo, L., Danckaert, A., Chamoun, Z., Roux, P., and Zurzolo, C. (2013).

Transfer of polyglutamine aggregates in neuronal cells occurs in tunneling nanotubes. Journal

of cell science *126*, 3678-3685.

Davis, G.W. (2013) Homeostatic signaling and the stabilization of neural function. Neuron *80*, 718-728.

Davis, A.A., Leyns, C.E.G., and Holtzman, D.M. (2018). Intercellular Spread of Protein Aggregates in

Neurodegenerative Disease. Annual review of cell and developmental biology *34*, 545-568.

Deng, Y.P., Albin, R.L., Penney, J.B., Young, A.B., Anderson, K.D., and Reiner, A. (2004). Differential

loss of striatal projection systems in Huntington's disease: a quantitative immunohistochemical

study. Journal of chemical neuroanatomy *27*, 143-164.

Donnelly, K.M., and Pearce, M.M.P. (2018). Monitoring Cell-to-cell Transmission of Prion-like Protein

Aggregates in Drosophila Melanogaster. Journal of visualized experiments *12*, 133.

Etchegaray, J.I., Elguero, E.J., Tran, J.A., Sinatra, V., Feany, M.B., and McCall, K. (2016). Defective

Phagocytic Corpse Processing Results in Neurodegeneration and Can Be Rescued by TORC1

Activation. The Journal of neuroscience : the official journal of the Society for Neuroscience *36*,

3170-3183.

Falcon, B., Noad, J., McMahon, H., Randow, F., and Goedert, M. (2018). Galectin-8-mediated selective

autophagy protects against seeded tau aggregation. The Journal of biological chemistry *293*,

2438-2451.

Flavin, W.P., Bousset, L., Green, Z.C., Chu, Y., Skarpathiotis, S., Chaney, M.J., Kordower, J.H., Melki,

R., and Campbell, E.M. (2017). Endocytic vesicle rupture is a conserved mechanism of cellular

invasion by amyloid proteins. Acta neuropathologica *134*, 629-653.

Frackowiak, J., Wisniewski, H.M., Wegiel, J., Merz, G.S., Iqbal, K., and Wang, K.C. (1992).

Ultrastructure of the microglia that phagocytose amyloid and the microglia that produce beta-

amyloid fibrils. Acta neuropathologica *84*, 225-233.

Freeman, M.R. (2015). Drosophila Central Nervous System Glia. Cold Spring Harbor perspectives in

biology *7*, a020552.

Freeman, M.R., Delrow, J., Kim, J., Johnson, E., and Doe, C.Q. (2003). Unwrapping glial biology: Gcm

target genes regulating glial development, diversification, and function. Neuron *38*, 567-580.

Funk, K.E., Mirbaha, H., Jiang, H., Holtzman, D.M., and Diamond, M.I. (2015). Distinct Therapeutic

Mechanisms of Tau Antibodies: Promoting Microglial Clearance Versus Blocking Neuronal

Uptake. The Journal of biological chemistry *290*, 21652-21662.

Graham, R.K., Deng, Y., Slow, E.J., Haigh, B., Bissada, N., Lu, G., Pearson, J., Shehadeh, J., Bertram,

L., Murphy, Z.*, et al.* (2006). Cleavage at the caspase-6 site is required for neuronal dysfunction

and degeneration due to mutant huntingtin. Cell *125*, 1179-1191.

Grathwohl, S.A., Kalin, R.E., Bolmont, T., Prokop, S., Winkelmann, G., Kaeser, S.A., Odenthal, J.,

Radde, R., Eldh, T., Gandy, S.*, et al.* (2009). Formation and maintenance of Alzheimer's

disease beta-amyloid plaques in the absence of microglia. Nature neuroscience *12*, 1361-1363.

Griciuc, A., Patel, S., Federico, A.N., Choi, S.H., Innes, B.J., Oram, M.K., Cereghetti, G., McGinty, D.,

Anselmo, A., Sadreyev, R.I.*, et al.* (2019). TREM2 Acts Downstream of CD33 in Modulating

Microglial Pathology in Alzheimer's Disease. Neuron *103*, 820-835.e827.

Guo, Y., Wei, X., Yan, H., Qin, Y., Yan, S., Liu, J., Zhao, Y., Jiang, F., and Lou, H. (2019). TREM2

deficiency aggravates alpha-synuclein-induced neurodegeneration and neuroinflammation in

Parkinson's disease models. FASEB journal *33*, 12164-12174.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and Garrity, P.A.

(2008). An internal thermal sensor controlling temperature preference in Drosophila. Nature

454, 217-220.

Hammond, T.R., Robinton, D., and Stevens, B. (2018). Microglia and the Brain: Complementary

Partners in Development and Disease. Annual review of cell and developmental biology *34*,

523-544.

Han, C., Song, Y., Xiao, H., Wang, D., Franc, N.C., Jan, L.Y., and Jan, Y.N. (2014). Epidermal cells are

the primary phagocytes in the fragmentation and clearance of degenerating dendrites in

Drosophila. Neuron *81*, 544-560.

Hay, B.A., Wolff, T., and Rubin, G.M. (1994). Expression of baculovirus P35 prevents cell death in

Drosophila. Development *120*, 2121-2129.

Holmes, B.B., DeVos, S.L., Kfoury, N., Li, M., Jacks, R., Yanamandra, K., Ouidja, M.O., Brodsky, F.M.,

Marasa, J., Bagchi, D.P.*, et al.* (2013). Heparan sulfate proteoglycans mediate internalization

and propagation of specific proteopathic seeds. Proceedings of the National Academy of

Sciences of the United States of America *110*, E3138-3147.

Hong, S., Beja-Glasser, V.F., Nfonoyim, B.M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K.M., Shi,

Q., Rosenthal, A., Barres, B.A.*, et al.* (2016). Complement and microglia mediate early synapse

loss in Alzheimer mouse models. Science *352*, 712-716.

Hoopfer, E.D., McLaughlin, T., Watts, R.J., Schuldiner, O., O'Leary, D.D., and Luo, L. (2006). Wlds

protection distinguishes axon degeneration following injury from naturally occurring

developmental pruning. Neuron *50*, 883-895.

Jefferis, G.S., Marin, E.C., Stocker, R.F., and Luo, L. (2001). Target neuron prespecification in the

olfactory map of Drosophila. Nature *414*, 204-208.

Jeon, I., Cicchetti, F., Cisbani, G., Lee, S., Li, E., Bae, J., Lee, N., Li, L., Im, W., Kim, M.*, et al.* (2016).

Human-to-mouse prion-like propagation of mutant huntingtin protein. Acta neuropathologica

132, 577-592.

Juhász, G., Hill, J.H., Yan, Y., Sass, M., Baehrecke, E.H., Backer, J.M., and Neufeld, T.P. (2008). The

class III PI(3)K Vps34 promotes autophagy and endocytosis but not TOR signaling in

Drosophila. The Journal of cell biology *181*, 655-666.

Knowles, T.P., Vendruscolo, M., and Dobson, C.M. (2014). The amyloid state and its association with protein misfolding diseases. Nature reviews Molecular cell biology *15*, 384-396.

Kosaka, T., and Ikeda, K. (1983). Possible temperature-dependent blockage of synaptic vesicle recycling induced by a single gene mutation in Drosophila. Journal of neurobiology *14*, 207-225.

Kreher, S.A., Kwon, J.Y., and Carlson, J.R. (2005). The molecular basis of odor coding in the Drosophila larva. Neuron *46*, 445-456.

Lai, S.-L., Awasaki, T., Ito, K., and Lee, T. (2008). Clonal analysis of Drosophila antennal lobe neurons: diverse neuronal architectures in the lateral neuroblast lineage. Development *135*, 2883-2893.

Lee, H.-J., Suk, J.-E., Patrick, C., Bae, E.-J., Cho, J.-H., Rho, S., Hwang, D., Masliah, E., and Lee, S.-J. (2010). Direct Transfer of α-Synuclein from Neuron to Astroglia Causes Inflammatory Responses in Synucleinopathies. Journal of Biological Chemistry *285*, 9262-9272.

Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. Neuron *22*, 451-461.

Leyns, C.E.G., Gratuze, M., Narasimhan, S., Jain, N., Koscal, L.J., Jiang, H., Manis, M., Colonna, M., Lee, V.M.Y., Ulrich, J.D.*, et al.* (2019). TREM2 function impedes tau seeding in neuritic plaques. Nature neuroscience *22*, 1217-1222.

Li, Z., Wang, C., Wang, Z., Zhu, C., Li, J., Sha, T., Ma, L., Gao, C., Yang, Y., Sun, Y.*, et al.* (2019). Allele-selective lowering of mutant HTT protein by HTT–LC3 linker compounds. Nature *575*, 203-209.

Liang, L., Li, Y., Potter, C.J., Yizhar, O., Deisseroth, K., Tsien, R.W., and Luo, L. (2013). GABAergic projection neurons route selective olfactory inputs to specific higher-order neurons. Neuron *79*, 917-931.

Liddelow, S.A., Guttenplan, K.A., Clarke, L.E., Bennett, F.C., Bohlen, C.J., Schirmer, L., Bennett, M.L., Munch, A.E., Chung, W.S., Peterson, T.C.*, et al.* (2017). Neurotoxic reactive astrocytes are induced by activated microglia. Nature *541*, 481-487.

Lin, D.M., and Goodman, C.S. (1994). Ectopic and increased expression of Fasciclin II alters

motoneuron growth cone guidance. Neuron *13*, 507-523.

MacDonald, J.M., Beach, M.G., Porpiglia, E., Sheehan, A.E., Watts, R.J., and Freeman, M.R. (2006).

The Drosophila cell corpse engulfment receptor Draper mediates glial clearance of severed

axons. Neuron *50*, 869-881.

MacDonald, M.E., Ambrose, C.M., Duyao, M.P., Myers, R.H., Lin, C., Srinidhi, L., Barnes, G., Taylor,

S.A., James, M., Groot, N.*, et al.* (1993). A novel gene containing a trinucleotide repeat that is

expanded and unstable on Huntington's disease chromosomes. Cell *72*, 971-983.

Marin, E.C., Jefferis, G.S., Komiyama, T., Zhu, H., and Luo, L. (2002). Representation of the glomerular

olfactory map in the Drosophila brain. Cell *109*, 243-255.

Masliah, E., Rockenstein, E., Adame, A., Alford, M., Crews, L., Hashimoto, M., Seubert, P., Lee, M.,

Goldstein, J., Chilcote, T.*, et al.* (2005). Effects of alpha-synuclein immunization in a mouse

model of Parkinson's disease. Neuron *46*, 857-868.

Masnata, M., Sciacca, G., Maxan, A., Bousset, L., Denis, H.L., Lauruol, F., David, L., Saint-Pierre, M.,

Kordower, J.H., Melki, R.*, et al.* (2019). Demonstration of prion-like properties of mutant

huntingtin fibrils in both in vitro and in vivo paradigms. Acta neuropathologica *137*, 981-1001.

Mezias, C., LoCastro, E., Xia, C., and Raj, A. (2017). Connectivity, not region-intrinsic properties,

predicts regional vulnerability to progressive tau pathology in mouse models of disease. Acta

neuropathologica communications *5*, 61.

Mosca, T.J., and Luo, L. (2014). Synaptic organization of the Drosophila antennal lobe and its

regulation by the Teneurins. Elife *3*, e03726.

Pearce, M.M., Spartz, E.J., Hong, W., Luo, L., and Kopito, R.R. (2015). Prion-like transmission of

neuronal huntingtin aggregates to phagocytic glia in the Drosophila brain. Nature

communications *6*, 6768.

Pecho-Vrieseling, E., Rieker, C., Fuchs, S., Bleckmann, D., Esposito, M.S., Botta, P., Goldstein, C.,

Bernhard, M., Galimberti, I., Muller, M.*, et al.* (2014). Transneuronal propagation of mutant

huntingtin contributes to non-cell autonomous pathology in neurons. Nature neuroscience *17*,

1064-1072.

Potter, C.J., Tasic, B., Russler, E.V., Liang, L., and Luo, L. (2010). The Q system: a repressible binary

system for transgene expression, lineage tracing, and mosaic analysis. Cell *141*, 536-548.

Poudel, G.R., Harding, I.H., Egan, G.F., and Georgiou-Karistianis, N. (2019). Network spread

determines severity of degeneration and disconnection in Huntington's disease. Human brain

mapping *40*, 4192-4201.

Preisinger, E., Jordan, B.M., Kazantsev, A., and Housman, D. (1999). Evidence for a recruitment and

sequestration mechanism in Huntington's disease. Philosophical transactions of the Royal

Society of London Series B, Biological sciences *354*, 1029-1034.

Pulipparacharuvil, S., Akbar, M.A., Ray, S., Sevrioukov, E.A., Haberman, A.S., Rohrer, J., and Krämer,

H. (2005). Drosophila Vps16A is required for trafficking to lysosomes and

biogenesis of pigment granules. Journal of cell science *118*, 3663-3673.

Ray, A., Speese, S.D., and Logan, M.A. (2017). Glial Draper Rescues Abeta Toxicity in a Drosophila

Model of Alzheimer's Disease. The Journal of neuroscience : the official journal of the Society

for Neuroscience *37*, 11881-11893.

Reiner, A., and Deng, Y.P. (2018). Disrupted striatal neuron inputs and outputs in Huntington's disease.

CNS neuroscience & therapeutics *24*, 250-280.

Ren, P.H., Lauckner, J.E., Kachirskaia, I., Heuser, J.E., Melki, R., and Kopito, R.R. (2009). Cytoplasmic

penetration and persistent infection of mammalian cells by polyglutamine aggregates. Nature

cell biology *11*, 219-225.

Riabinina, O., Luginbuhl, D., Marr, E., Liu, S., Wu, M.N., Luo, L., and Potter, C.J. (2015). Improved and

expanded Q-system reagents for genetic manipulations. Nature methods *12*, 219-222.

Riabinina, O., and Potter, C.J. (2016). The Q-System: A Versatile Expression System for Drosophila.

Methods in molecular biology *1478*, 53-78.

Sathasivam, K., Neueder, A., Gipson, T.A., Landles, C., Benjamin, A.C., Bondulich, M.K., Smith, D.L.,

Faull, R.L., Roos, R.A., Howland, D.*, et al.* (2013). Aberrant splicing of HTT generates the

pathogenic exon 1 protein in Huntington disease. Proceedings of the National Academy of

Sciences of the United States of America *110*, 2366-2370.

Schafer, D.P., Lehrman, E.K., Kautzman, A.G., Koyama, R., Mardinly, A.R., Yamasaki, R., Ransohoff,

R.M., Greenberg, M.E., Barres, B.A., and Stevens, B. (2012). Microglia sculpt postnatal neural

circuits in an activity and complement-dependent manner. Neuron *74*, 691-705.

Sepp, K.J., Schulte, J., and Auld, V.J. (2001). Peripheral glia direct axon guidance across the

CNS/PNS transition zone. Developmental biology *238*, 47-63.

Sharma, M., and Subramaniam, S. (2019). Rhes travels from cell to cell and transports Huntington

disease protein via TNT-like protrusion. The Journal of cell biology *218*, 1972-1993.

Stepensky, D. (2007). FRETcalc plugin for calculation of FRET in non-continuous intracellular

compartments. Biochemical and biophysical research communications *359*, 752-758.

Stocker, R.F., Heimbeck, G., Gendre, N., and de Belle, J.S. (1997). Neuroblast ablation in Drosophila

P[GAL4] lines reveals origins of olfactory interneurons. Journal of neurobiology *32*, 443-456.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirian, L., and Dickson, B.J. (2005). Neural circuitry that

governs Drosophila male courtship behavior. Cell *121*, 795-807.

Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., and O'Kane, C.J. (1995). Targeted expression of

tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and causes

behavioral defects. Neuron *14*, 341-351.

Tabrizi, S.J., Leavitt, B.R., Landwehrmeyer, G.B., Wild, E.J., Saft, C., Barker, R.A., Blair, N.F.,

Craufurd, D., Priller, J., Rickards, H.*, et al.* (2019). Targeting Huntingtin Expression in Patients

with Huntington's Disease. The New England journal of medicine *380*, 2307-2316.

Tran, H.T., Chung, C.H., Iba, M., Zhang, B., Trojanowski, J.Q., Luk, K.C., and Lee, V.M. (2014). Alpha-

synuclein immunotherapy blocks uptake and templated propagation of misfolded alpha-

synuclein and neurodegeneration. Cell reports *7*, 2054-2065.

Tremblay, M.E., Zettel, M.L., Ison, J.R., Allen, P.D., and Majewska, A.K. (2012). Effects of aging and

sensory loss on glial cells in mouse visual and auditory cortices. Glia *60*, 541-558.

Trevino, R.S., Lauckner, J.E., Sourigues, Y., Pearce, M.M., Bousset, L., Melki, R., and Kopito, R.R.

(2012). Fibrillar structure and charge determine the interaction of polyglutamine protein

aggregates with the cell surface. The Journal of biological chemistry *287*, 29722-29728.

Vaquer-Alicea, J., and Diamond, M.I. (2019). Propagation of Protein Aggregation in Neurodegenerative

Diseases. Annual review of biochemistry *88*, 785-810.

Victoria, G.S., Arkhipenko, A., Zhu, S., Syan, S., and Zurzolo, C. (2016). Astrocyte-to-neuron

intercellular prion transfer is mediated by cell-cell contact. Scientific reports *6*, 20762.

Wang, N., Gray, M., Lu, X.-H., Cantle, J.P., Holley, S.M., Greiner, E., Gu, X., Shirasaki, D., Cepeda, C.,

Li, Y*., et al.* (2014). Neuronal targets for reducing mutant huntingtin expression to ameliorate

disease in a mouse model of Huntington's disease. Nature medicine *20*, 536-541.

Weinhard, L., di Bartolomei, G., Bolasco, G., Machado, P., Schieber, N.L., Neniskyte, U., Exiga, M.,

Vadisiute, A., Raggioli, A., Schertel, A*., et al.* (2018). Microglia remodel synapses by

presynaptic trogocytosis and spine head filopodia induction. Nature communications *9*, 1228.

Weiss, K.R., Kimura, Y., Lee, W.C., and Littleton, J.T. (2012). Huntingtin aggregation kinetics and their

pathological role in a Drosophila Huntington's disease model. Genetics *190*, 581-600.

Wilton, D.K., Dissing-Olesen, L., and Stevens, B. (2019). Neuron-Glia Signaling in Synapse Elimination.

Annual review of neuroscience *42*, 107-127.

Wu, J.W., Hussaini, S.A., Bastille, I.M., Rodriguez, G.A., Mrejeru, A., Rilett, K., Sanders, D.W., Cook,

C., Fu, H., Boonen, R.A*., et al.* (2016). Neuronal activity enhances tau propagation and tau

pathology in vivo. Nature neuroscience *19*, 1085-1092.

Wyss-Coray, T., Loike, J.D., Brionne, T.C., Lu, E., Anankov, R., Yan, F., Silverstein, S.C., and

Husemann, J. (2003). Adult mouse astrocytes degrade amyloid-beta in vitro and in situ. Nature

medicine *9*, 453-457.

Yanamandra, K., Kfoury, N., Jiang, H., Mahan, T.E., Ma, S., Maloney, S.E., Wozniak, D.F., Diamond,

M.I., and Holtzman, D.M. (2013). Anti-tau antibodies that block tau aggregate seeding in vitro

markedly decrease pathology and improve cognition in vivo. Neuron *80*, 402-414.

Yun, S.P., Kam, T.I., Panicker, N., Kim, S., Oh, Y., Park, J.S., Kwon, S.H., Park, Y.J., Karuppagounder,

877   S.S., Park, H.*, et al.* (2018). Block of A1 astrocyte conversion by microglia is neuroprotective in

878   models of Parkinson's disease. Nature medicine *24*, 931-938.

879   Zeineddine, R., Pundavela, J.F., Corcoran, L., Stewart, E.M., Do-Ha, D., Bax, M., Guillemin, G., Vine,

880   K.L., Hatters, D.M., Ecroyd, H.*, et al.* (2015). SOD1 protein aggregates stimulate

881   macropinocytosis in neurons to facilitate their propagation. Molecular neurodegeneration *10*, 57.

882   Zhao, Y., Wu, X., Li, X., Jiang, L.L., Gui, X., Liu, Y., Sun, Y., Zhu, B., Pina-Crespo, J.C., Zhang, M.*, et*

883   *al.* (2018). TREM2 Is a Receptor for beta-Amyloid that Mediates Microglial Function. Neuron *97*,

884   1023-1031.

A Htt: ex1, ex1-12, HR 1-7, HR 8-15, HR 17-22, HR 23-28, HR 29-36, polyQ

B projection neurons (PNs) (wtHtt$_{ex1}$), mushroom body and lateral horn, glial cell, olfactory receptor neurons (ORNs) (mHtt$_{ex1\ or\ ex1-12}$), glomeruli, antenna, antennal lobe

glial cell, DA1 ORN axon (mHtt$_{ex1\ or\ ex1-12}$), DA1 glomerulus, DA1 PN dendrite (wtHtt$_{ex1}$)

C Htt$_{ex1}$Q91-mCherry, Htt$_{ex1}$Q25-GFP, merge, 1d, 7d, 14d, DA1

D Htt$_{ex1}$Q91-mCherry puncta, age (days)

E Htt$_{ex1}$Q25-GFP puncta, age (days)

F Htt$_{ex1}$Q91-mCherry, Htt$_{ex1}$Q25-GFP, merge

G Htt$_{ex1}$Q25-GFP puncta, age (days)

H RFP-Htt$_{ex1-12}$Q138, Htt$_{ex1}$Q25-GFP, merge, DA1

Figure 1. mHtt_{ex1} or mHtt_{ex1-12} aggregates generated in presynaptic ORNs nucleate the aggregation of wtHtt_{ex1} in postsynaptic PNs. (A) Cartoon depiction of full-length human Htt (3144 amino acids). The majority of the Htt protein is composed of 36 HEAT repeat domains (HR, gray regions). The variable-length polyQ region (green/magenta) is located near the N-terminus, with the pathogenic threshold (~Q37) indicated by the white dotted line. The C-termini of two N-terminal pathogenic fragments used in this study (ex1 and ex1-12) are indicated by arrows. (B) Diagram illustrating our overall experimental approach. In the fly olfactory system, primary ORNs synapse with secondary PNs in discrete regions of the antennal lobe known as glomeruli (gray circles). PNs send their axons into higher brain centers (mushroom body and/or lateral horn). Several glial cell types also reside in or near the antennal lobe, including Draper-expressing glia whose processes ensheath individual glomeruli in the antennal lobe. To examine spreading of mHtt aggregates between synaptically-connected ORNs and PNs, we generated transgenic flies that express mHtt fragments (*QUAS-Htt$_{ex1}$Q91-mCherry* or *QUAS-RFP-Htt$_{ex1-12}$Q138*) in DA1 ORNs using the *Or67d-QF* driver and *UAS-Htt$_{ex1}$Q25-GFP* in DA1 PNs using *GH146-Gal4*, which labels ~60% of PNs. Inset: Transfer of mHtt aggregates between ORNs and PNs was assessed by monitoring the solubility of Htt$_{ex1}$Q25-GFP in PN dendrites and cell bodies and co-localization of GFP+ puncta with mCherry/RFP+ mHtt aggregates. (C) Maximum intensity z-projections of the DA1 glomerulus from adult males at the indicated ages expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs (*magenta*) and Htt$_{ex1}$Q25-GFP in GH146+ PNs (*green*). Approximate boundaries of the DA1 glomerulus are indicated by the white dotted line. (D) and (E) Quantification of Htt$_{ex1}$Q91-mCherry (D) and Htt$_{ex1}$Q25-GFP (E) puncta in the DA1 glomerulus of adult female (*solid bars*) or male (*open bars*) flies at the indicated ages. (F) Single z-slice images showing Htt$_{ex1}$Q25-GFP puncta (*arrowheads*) in lateral PN soma from 21 day-old males with the same genotype as in (C-E). (G) Quantification of Htt$_{ex1}$Q25-GFP puncta detected in PN soma in female (*solid bars*) and male (*open bars*) flies at the indicated ages. Puncta that were GFP+ only and those that were both GFP+ and mCherry+ at each age are indicated by stacked green and magenta bars, respectively. (H) Maximum intensity z-projections from a 7 day-old adult female expressing RFP-Htt$_{ex1-12}$Q138 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. In (D), (E), and (G), error bars indicate ± SEM; *n* is indicated below each data set. Statistical comparisons made for females or males at different ages were calculated using two-way ANOVA with Tukey's multiple comparisons tests, and Student's *t*-test was used for pairwise comparisons between age-matched females and males (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, or ****$p < 0.0001$). Scale bars = 10 μm in (C) and (H), and 5 μm in (F).

Figure 1 – figure supplement 1. mHtt$_{ex1}$ aggregates induce prion-like conversion of wtHtt$_{ex1}$ in neurons. (A and B) Maximum intensity z-projections of 7 day-old adult female brains expressing either Htt$_{ex1}$Q91-mCherry (A) or Htt$_{ex1}$Q25-GFP (B) using pan-neuronal *elav[C155]-Gal4*. (C) and (D) Co-expression of Htt$_{ex1}$Q91-mCherry and Htt$_{ex1}$Q25-GFP (C) or GFP (D) in all neurons with *elav[C155]-Gal4*. Merged images include DAPI signal (blue) to indicate nuclei. Scale bars = 10 μm in all images.

Figure 1 – figure supplement 2. Controls for spreading of mHtt$_{ex1}$ aggregates between synaptically-connected ORNs and PNs. (A-D) Maximum-intensity z-projections of the DA1 glomerulus from 10 day-old adults expressing (A) Htt$_{ex1}$Q25-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, (B) Htt$_{ex1}$Q91-mCherry in DA1 ORNs and GFP in PNs, (C) Htt$_{ex1}$Q91-mCherry together with Gal80 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, and (D) Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP together with QS in GH146+ PNs. Scale bars = 10 μm. (E and F) Quantification of mCherry+ (E) and GFP+ (F) aggregates for all genotypes shown in (A-D). Horizontal bars indicate mean, and error bars indicate ± SEM; n ≥ 8. (G) Table summarizing genotypes and graph symbols displayed in (A-F).

7d

Htt$_{ex1}$Q91-mCherry (PNs)   Htt$_{ex1}$Q25-GFP (ORNs)   Brp   merge

z-16

z-27

z-39

14d

Htt$_{ex1}$Q91-mCherry (PNs)   Htt$_{ex1}$Q25-GFP (ORNs)   Brp   merge

z-10

z-20

z-54

Figure 1 – figure supplement 3. mHtt$_{ex1}$ aggregates do not transfer retrogradely from PNs to ORNs. Single confocal z-slices of the antennal lobe from 7 (*top 3 rows*) and 14 (*bottom 3 rows*) day-old adults expressing Htt$_{ex1}$Q91-mCherry in ~60% of PNs using *GH146-QF* and Htt$_{ex1}$Q25-GFP in all ORNs using *pebbled-Gal4*. The brains were immunostained with antibodies against mCherry (*magenta*), GFP (*green*), and the neuropil marker Bruchpilot (Brp; *blue*). Individual mCherry+ puncta are indicated by solid white arrows and punctate GFP+ structures by open arrowheads. Scale bars = 10 μm, relative z-positions in each confocal stack are indicated in upper right of merged images.

Figure 2. Features of induced Htt$_{ex1}$Q25-GFP aggregates in PNs. (A) Three-dimensional rendering of the DA1 glomerulus from a 14 day-old adult male expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. (B) A single confocal slice from an adult male with the same genotype as in (A) showing four induced Htt$_{ex1}$Q25-GFP aggregates and associated Htt$_{ex1}$Q91-mCherry signals (arrows). (C and D) Volumetric surface renderings of the raw confocal data shown in (A). Thresholding was adjusted in each channel to ensure that the majority of Htt$_{ex1}$Q91-mCherry puncta were identified as individual objects (mCherry surfaces) and induced Htt$_{ex1}$Q25-GFP aggregates were identified as objects distinguishable from surrounding diffuse signal (GFP surfaces). In the higher magnification view shown in (D), GFP surfaces were set at 50% transparency in the merged image so that co-localized mCherry surfaces are visible. Five individual Htt$_{ex1}$Q91-mCherry + Htt$_{ex1}$Q25-GFP co-localized aggregates are indicated by arrows. (E) A single mCherry+ and GFP+ aggregate before ("Pre-AccPb") and after ("Post-AccPb") photobleaching of the mCherry acceptor fluorophore. A larger mCherry-only aggregate is also shown in the image as a control. Pixel-by-pixel FRET efficiency (FRET$_{eff}$) is shown as a heat map image (*bottom right*; colorbar scale to right of image), calculated using the equation (GFP$_{final}$ - GFP$_{initial}$)/GFP$_{final}$. Average FRET$_{eff}$ for this aggregate was 11.5%. Average FRET$_{eff}$ values for all induced Htt$_{ex1}$Q25-GFP aggregates tested by this method are shown in the histogram above the FRET$_{eff}$ image (*n* = 28). (F) Distribution of volumes for all mCherry+ aggregates (*gray bars*; *n* = 1792) or mCherry+ aggregate seeds that co-localized with induced Htt$_{ex1}$Q25-GFP aggregates (*black bars*; *n* = 100) detected in the DA1 glomerulus of 7 and 14 day-old adult flies with the same genotype as in (A-E). Scale bars = 10 μm (A and C), 5 μm (B), 2 μm (D), and 1 μm (E).

Figure 3. mHtt$_{ex1}$ aggregate transfer from ORNs to synaptically-connected PNs is inversely correlated with presynaptic activity. (A) Maximum-intensity z-projections of the DA1 glomerulus from 10 day-old adult males co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or shi^{ts1} in DA1 ORNs and also expressing Htt$_{ex1}$Q25-GFP in GH146+ PNs. Flies were shifted from the permissive temperature (18°C) to the restrictive temperature (31°C) upon eclosion. (B) and (C) Quantification of Htt$_{ex1}$Q91-mCherry (B) and Htt$_{ex1}$Q25-GFP (C) puncta in the DA1 glomerulus of 4 or 10 day-old adult males expressing LacZ or shi^{ts1} in ORNs using two independent *QUAS-shi^{ts1}* lines. (D) Maximum-intensity z-projections of the DA1 glomerulus from 10 day-old males co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or TeTxLC in DA1 ORNs and expressing Htt$_{ex1}$Q25-GFP in GH146+ PNs. (E) and (F) Quantification of Htt$_{ex1}$Q91-mCherry (E) and Htt$_{ex1}$Q25-GFP (F) puncta in the DA1 glomerulus of 4 or 10 day-old adult males expressing LacZ or TeTxLC in ORNs using two independent *QUAS-TeTxLC* lines. (G) Maximum-intensity z-projections of the DA1 glomerulus from 7 day-old adult females co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or dTrpA in DA1 ORNs and expressing Htt$_{ex1}$Q25-GFP in GH146+ PNs. dTrpA-expressing neurons were activated by shifting the adults from 25°C to 31°C upon eclosion. (H) and (I) Quantification of Htt$_{ex1}$Q91-mCherry (H) and Htt$_{ex1}$Q25-GFP (I) puncta in the DA1 glomerulus of 7 day-old females expressing LacZ or dTrpA in ORNs using three independent *QUAS-dTrpA* lines. For quantifications shown in (B), (C), (D), (E), (H), and (I), error bars indicate ± SEM; *n* is indicated for each data set. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, or n.s. = not significant by two-way ANOVA with Sidak's multiple comparisons tests. Scale bars = 10 μm in all images.

Figure 4. Inhibition of endocytosis increases mHtt$_{ex1}$-expressing ORN axon volume and enhances transfer of neuronal mHtt$_{ex1}$ aggregates to glia. (A) Maximum-intensity z-projections of the DA1 glomerulus from 10 day-old females co-expressing Htt$_{ex1}$Q91-mCherry, mCD8-GFP, and either LacZ or shi^{ts1} in DA1 ORNs. Adult flies were shifted from 18°C to 31°C upon eclosion. (B) Volumetric surface renderings generated for Htt$_{ex1}$Q91-mCherry and mCD8-GFP signals from the same confocal stacks shown in (A). (C) Quantification of mCD8-GFP intensity (*left y-axis, green*) and DA1 ORN axon volume (*right y-axis, black*) in the DA1 glomerulus of 10 day-old adult females expressing LacZ or shi^{ts1} together with Htt$_{ex1}$Q91-mCherry and mCD8-GFP in DA1 ORNs. a.u. = arbitrary units. (D) Maximum-intensity z-projections of the DA1 glomerulus from 5-6 day-old males co-expressing Htt$_{ex1}$Q91-mCherry with either LacZ or shi^{ts1} in DA1 ORNs and expressing Htt$_{ex1}$Q25-YFP in all glia using *repo-Gal4*. Adult flies of each genotype were shifted from 18°C to the restrictive temperature (31°C) upon eclosion. (E and F) Quantification of Htt$_{ex1}$Q91-mCherry (E) and Htt$_{ex1}$Q25-YFP (F) puncta in the DA1 glomerulus of 5-6 day-old males expressing LacZ or shi^{ts1} in ORNs using two independent *QUAS-shi^{ts1}* lines. For quantifications shown in (C), (E), and (F), error bars indicate ± SEM; *n* is indicated below each data set. **p < 0.01, ****p < 0.0001, or n.s. = not significant by two-way ANOVA with Sidak's multiple comparisons tests in (C) and by one-way ANOVA with Tukey's multiple comparisons tests in (E) and (F). Scale bars = 10 μm in all images.

Figure 5. Draper is required for mHtt$_{ex1}$ aggregate transfer from presynaptic ORNs to postsynpatic PNs and regulates neuronal mHtt$_{ex1}$ aggregate size. (A) Maximum-intensity z-projections of the DA1 glomerulus from 13 day-old adult females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs and either heterozygous (*drpr$^{\Delta5/+}$*) or homozygous (*drpr$^{\Delta5/\Delta5}$*) for the *drpr$^{\Delta5}$* null allele. Scale bars = 10 μm. (B and C) Numbers of Htt$_{ex1}$Q91-mCherry (B) or Htt$_{ex1}$Q25-GFP (C) puncta in the DA1 glomerulus of female or male *drpr$^{\Delta5/+}$* or *drpr$^{\Delta5/\Delta5}$* flies. Each symbol represents quantification for an individual DA1 glomerulus. Horizontal bars indicate means, and error bars indicate ± SEM; *n* is shown above each data set. *p < 0.05, **p < 0.01, or ****p < 0.0001 by two-way ANOVA with Tukey's multiple comparisons tests. "n.d." = not determined, as *drpr$^{\Delta5/\Delta5}$* males did not survive to 13 days old. (D) Three-dimensional rendering of mCherry+ surfaces identified in the DA1 glomerulus of 7 day-old *drpr$^{+/+}$* or *drpr$^{\Delta5/\Delta5}$* females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs. Colors indicate aggregate volume according to the colorbar scale below the images. Scale bars = 5 μm. (E) Distribution of volumes calculated for all mCherry+ aggregates in 7 day-old *drpr$^{+/+}$* (*gray line and fill*; *n* = 1556) or *drpr$^{\Delta5/\Delta5}$* (*black line and striped fill*; *n* = 1749) male and female flies. Purple and green shading indicate smaller- ("A") and larger-sized ("B") aggregate subpopulations, respectively, that experienced a shift in abundance between wild-type and *drpr*-null animals.

Figure 5 – figure supplement 1. Draper is required for enhanced transfer of mHtt$_{ex1}$ from shi^{ts}-expressing ORNs to synaptically-connected PNs. (A) Maximum-intensity z-projections of the DA1 glomerulus from 7 day-old *drpr$^{+/+}$* or *drpr$^{\Delta5/\Delta5}$* adult females co-expressing Htt$_{ex1}$Q91-mCherry with either LacZ or shi^{ts1} in ORNs and Htt$_{ex1}$Q25-GFP in PNs. Adult flies were shifted from 18°C to the restrictive temperature (31°C) upon eclosion. Scale bars = 10 μm. (B and C) Quantification of Htt$_{ex1}$Q91-mCherry (B) and Htt$_{ex1}$Q25-GFP (C) aggregates for controls expressing LacZ in ORNs (*squares*), and flies expressing shi^{ts1} in ORNs in *drpr$^{+/+}$* (*circles*) or *drpr$^{\Delta5/\Delta5}$* backgrounds (*triangles*). Horizontal bars indicate means, and error bars indicate ± SEM; *n* is shown below each data set. ****p < 0.0001, or n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons tests.

Figure 5 – figure supplement 2. Gal80-mediated repression of Gal4 in glia and RNAi knockdown of *drpr* in PNs do not affect ORN-to-PN transfer of mHtt$_{ex1}$ aggregates. (A and B) Maximum-intensity z-projections of the DA1 glomerulus from 2 week-old adult males showing that (A) inhibiting possible *GH146-Gal4*-mediated expression in glia with *repo-Gal80* or (B) expression of siRNAs targeting *drpr* in PNs do not affect Htt$_{ex1}$Q25-GFP aggregation in PNs when Htt$_{ex1}$Q91-mCherry is expressed in ORNs. Control flies did not express repo-Gal80 (A) or co-expressed firefly luciferase (FFLuc) with Htt$_{ex1}$Q25-GFP in PNs (B). Scale bars = 10 μm. (C and D) Quantification of Htt$_{ex1}$Q91-mCherry (C) and Htt$_{ex1}$Q25-GFP (D) aggregates for genotypes shown in (A) and (B). Horizontal bars indicate mean, and error bars indicate ± SEM; *n* is shown below each data set. n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons tests.

Figure 5 – figure supplement 3. mHtt$_{ex1}$ aggregates generated in ORNs do not co-localize with markers of lysosomes or autophagosomes in glia. (A and B) Single confocal slices from the DA1 glomerulus of 1 week-old adult females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs using *Or67d-QF* and the lysosomal marker GFP-Lamp1 (A) or the autophagosome marker Atg8a-GFP (B) in glia using *repo-Gal4*. Line scans in (A) and (B) show intensity of mCherry (*magenta*) and GFP (*green*) fluorescence along dotted lines indicated in images. Scale bars = 5 μm; a.u. = arbitrary units.

Figure 6. Caspase activity in ORNs mediates mHtt$_{ex1}$ aggregate transfer from ORNs to PNs. (A) Maximum-intensity z-projections of an antennal lobe from 8 day-old *drpr*-null adult males expressing Htt$_{ex1}$Q25-GFP (*top*) or Htt$_{ex1}$Q91-GFP (*bottom*) in most ORNs using *Or83b-Gal4* driver. Brains were immunostained with antibodies against GFP (*green*) or cleaved Dcp-1 (*grayscale*). (B) Quantification of cDcp-1 immunoreactivity in the antennal lobe of flies expressing Htt$_{ex1}$Q25-GFP (*black bar*) or Htt$_{ex1}$Q91-GFP (*white bar*) in Or83b+ ORNs. (C) Maximum-intensity z-projections of the DA1 glomerulus from 12 day-old males co-expressing Htt$_{ex1}$Q91-mCherry with LacZ (control) or p35 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. (D) and (E) Quantification of Htt$_{ex1}$Q91-mCherry (D) or Htt$_{ex1}$Q25-GFP (E) puncta in the DA1 glomerulus of 12 day-old adult males co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or p35 in ORNs (*squares*) or co-expressing Htt$_{ex1}$Q25-GFP with LacZ or p35 in PNs (*circles*). Numbers of aggregates in flies expressing LacZ or p35 are indicated by solid or open shapes, respectively. For quantifications shown in (B), (D), and (E), horizontal bars indicate means and error bars indicate ± SEM; *n* is shown below each data set; **p < 0.01, ***p < 0.001, or ****p < 0.0001 by Student's t-test in (B) or one-way ANOVA with Tukey's multiple comparisons tests in (D) and (E). Scale bars = 10 μm in all images.

Figure 7. mHtt$_{ex1}$ aggregates transfer from presynaptic ORNs to postsynaptic PNs via a Draper-dependent phagocytic glial cytoplasmic intermediate. (A) Diagram illustrating our experimental approach for testing a role for Draper-dependent glia in Htt$_{ex1}$Q91-mCherry aggregate transfer from DA1 ORNs to DA1 PNs. Flies that combined *Or67d-QF-*, *repo-Gal4-*, and *GH146-LexA:GAD*-driven expression of Htt$_{ex1}$Q91-mCherry in DA1 ORNs, Htt$_{ex1}$Q25-3xHA in all glia, and Htt$_{ex1}$Q25-YFP in ~60% of PNs, respectively, were generated and analyzed for immunoreactivity of each fluorescent protein or epitope tag. If Htt$_{ex1}$Q91-mCherry aggregates travel to PNs via the glial cytoplasm (route 1), triple-labeled (mCherry+/3xHA+/YFP+) aggregates would be expected to appear. By contrast, if Htt$_{ex1}$Q91-mCherry aggregates transfer directly to PNs without accessing the glial cytoplasm (route 2), then only double-labeled (mCherry+/3xHA+ and mCherry+/YFP+) aggregates would be detected. (B) Maximum-intensity z-projections of the DA1 glomerulus from adult females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs (*magenta*), Htt$_{ex1}$Q25-3xHA in glia (*blue*), and Htt$_{ex1}$Q25-YFP in PNs (*green*) at the indicated ages. Brains were immunostained with antibodies against the unique mCherry, 3xHA, and GFP tags on each Htt protein. Scale bars = 10 μm. (C) Single confocal z-slices from females with the same genotype as in (B) at the indicated ages. mCherry+/3xHA+/YFP+ aggregates, mCherry+/3xHA+ aggregates, and 3xHA+-only aggregates are indicated by arrows, open arrowheads, and asterisks, respectively. The hashtag in the 7d image indicates an mCherry+/3xHA+ aggregate for which the associated mCherry+ seed is in a different focal plane. Annotations are shown in yellow on each image for increased visibility. Scale bars = 5 μm. (D) and (E) Quantification of aggregates with only one label (D) or co-labeled aggregates (E) that were detected in adult female brains at the indicated ages. Aggregate species that were not detected in our samples (i.e., mCherry+/YFP+ or 3xHA+/YFP+) are not shown. Error bars indicate ± SEM; *n* is shown below each data set. Statistical comparisons were made between different time points for each aggregate species using two-way ANOVA with Tukey's multiple comparisons tests (*$p < 0.05$, *$p < 0.01$, or ****$p < 0.0001$).

Table 1. Genotypes displayed in each figure

Figure panel(s)	Genotype
Figure 1	
C,D,E,F,G	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/(+/Y)$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$
H	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/QUAS\text{-}RFP\text{-}Htt_{ex1\text{-}12}Q138^{attP3B}$; $GH146\text{-}Gal4/+$
Figure 1, figure supplement 1	
A	$elav[C155]\text{-}Gal4/UAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP3}$
B	$elav[C155]\text{-}Gal4/+$; $UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP24}/+$
C	$elav[C155]\text{-}Gal4/UAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP3}$; $UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP24}/+$
D	$elav[C155]\text{-}Gal4/UAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP3}/+$; $UAS\text{-}GFP^{attP24}/+$
Figure 1, figure supplement 2	
A	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/+$; $GH146\text{-}Gal4/QUAS\text{-}Htt_{ex1}Q25\text{-}mCherry^{attP24}$
B	$Or67d\text{-}QF/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/UAS\text{-}GFP^{attP24}$
C	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/QUAS\text{-}Gal80$; $GH146\text{-}Gal4/QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}$
D	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/+$; $GH146\text{-}Gal4/QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}$; $UAS\text{-}QS/+$
Figure 1, figure supplement 3	
	$pebbled\text{-}Gal4/QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP3}$; $UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP24}/+$; $GH146\text{-}QF/+$
Figure 2	
A,B,C,D,E,F	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/(+/Y)$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$
Figure 3	
A: + LacZ	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}nucLacZ/+$
A: + shi^{ts1}	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}shibire^{ts1}\#7/+$
B,C: LacZ	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}nucLacZ/+$
B,C: shi^{ts1}#1	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}shibire^{ts1}\#5/+$
B,C: shi^{ts1}#2	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}shibire^{ts1}\#7/+$
D: + LacZ	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}nucLacZ/+$
D: + TeTxLC	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/Pin$; $QUAS\text{-}TeTxLC\#9c/+$
E,F: LacZ	$Or67d\text{-}QF,UAS\text{-}Htt_{ex1}Q25\text{-}GFP^{attP3}/Y$; $GH146\text{-}Gal4,QUAS\text{-}Htt_{ex1}Q91\text{-}mCherry^{attP24}/+$; $QUAS\text{-}nucLacZ/+$

E,F: TeTxLC#1	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/QUAS-TeTxLC#4c ; +/(Dh/Tm6b)*
E,F: TeTxLC#2	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/Pin; QUAS-TeTxLC#9c/+*
G: + LacZ	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*
G: + dTrpA	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/QUAS-dTrpA#7*
H,I: LacZ	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*
H,I: dTrpA#1	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/QUAS-dTrpA#7*
H,I: dTrpA#2	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-dTrpA#5 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*
H,I: dTrpA#3	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-dTrpA#6/+*

Figure 4

A,B,C: + LacZ	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; QUAS-mCD8-GFP/+ ; QUAS-nucLacZ/+*
A,B,C: + shi^{ts1}	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; QUAS-mCD8-GFP/+ ; QUAS-shibire^{ts1}#7/+*
D: + LacZ	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-nucLacZ*
D: + shi^{ts1}	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-shibire^{ts1}#7*
E,F: LacZ	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-nucLacZ*
E,F: shi^{ts1}#1	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-shibire^{ts1}#5*
E,F: shi^{ts1}#2	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-shibire^{ts1}#7*

Figure 5

A,B,C: drpr$^{Δ5/+}$	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/+*
A,B,C: drpr$^{Δ5/Δ5}$	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*
D,E: drpr$^{+/+}$	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/(+/Y)*
D,E: drpr$^{Δ5/Δ5}$	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/(+/Y) ; ; drpr^{Δ5}/drpr^{Δ5}*

Figure 5, figure supplement 1

A,B,C: + shi^{ts1}, drpr$^{+/+}$	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-shibire^{ts1}#2 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*
A,B,C: + shi^{ts1}, drpr$^{Δ5/Δ5}$	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-shibire^{ts1}#2 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*
B,C: + LacZ, drpr$^{+/+}$	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*

Figure 5, figure supplement 2	
A,C,D: - repo-Gal80	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*
A,C,D: + repo-Gal80	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/repo-Gal80*
B,C,D: + FFLuc in PNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; UAS-FFLuc ^{attP2}/+*
B,C,D: + Draper^{RNAi} in PNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; UAS-Draper$^{RNAi\#5}$/+*
Figure 5, figure supplement 3	
A	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; UAS-GFP-Lamp1/+ ; repo-Gal4/+*
B	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; UAS-Atg8a-GFP/+ ; repo-Gal4/+*
Figure 6	
A(top),B	*Or83b-Gal4,UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*
A(bottom),B	*Or83b-Gal4,UAS-Htt$_{ex1}$Q91-GFP^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*
C: + LacZ	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*
C: + p35	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+; QUAS-p35/+*
D,E: LacZ in ORNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*
D,E: p35 in ORNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+; QUAS-p35/+*
D,E: LacZ in PNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/UAS-LacZ*
D,E: p35 in PNs	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/UAS-p35*
Figure 7	
B,C,D,E	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/+*
C: + Drpr^{RNAi}	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/UAS-Draper$^{RNAi\#5}$*