1

Phagocytic glia are obligatory intermediates in transmission of mutant huntingtin aggregates across neuronal synapses

Kirby M. Donnelly[1], Olivia R. DeLorenzo[2], Aprem D.A. Zaya[1], Gabrielle E. Pisano[1], Wint M. Thu[1], Liqun Luo[3,4], Ron R. Kopito[3], Margaret M. Panning Pearce[1,2,5]

[1]Department of Biological Sciences and [2]Program in Neuroscience, University of the Sciences, Philadelphia, PA 19104, USA

[3]Department of Biology, Stanford University, Stanford, CA 94305, USA

[4]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA

[5]Correspondence: m.pearce@usciences.edu

ABSTRACT

Emerging evidence supports the hypothesis that pathogenic protein aggregates associated with neurodegenerative diseases spread from cell to cell through the brain in a manner akin to infectious prions. Here, we show that mutant huntingtin (mHtt) aggregates associated with Huntington disease transfer anterogradely from presynaptic to postsynaptic neurons in the adult *Drosophila* olfactory system. Trans-synaptic transmission of mHtt aggregates is inversely correlated with neuronal activity and blocked by inhibiting caspases in presynaptic neurons, implicating synaptic dysfunction and cell death in aggregate spreading. Remarkably, mHtt aggregate transmission across synapses requires the glial scavenger receptor Draper and involves a transient visit to the glial cytoplasm, indicating that phagocytic glia act as obligatory intermediates in aggregate spreading between synaptically-connected neurons. These findings expand our understanding of phagocytic glia as double-edged players in neurodegeneration—by clearing neurotoxic protein aggregates, but also providing an opportunity for prion-like seeds to evade phagolysosomal degradation and propagate further in the brain.

INTRODUCTION

Neurodegenerative diseases have emerged as one of the greatest healthcare challenges in our aging society, and thus a better understanding of the underlying pathological mechanisms is critical for development of more effective treatments or cures for these fatal disorders. A common molecular feature of many neurodegenerative diseases [e.g., Alzheimer disease (AD), frontotemporal dementias (FTD), Parkinson disease (PD), amyotrophic lateral sclerosis (ALS), and Huntington disease (HD)] is the misfolding of certain proteins, driving their accumulation into insoluble, amyloid aggregates (Knowles et al., 2014). Appearance of proteinaceous deposits in patient brains correlates closely with neuronal loss and clinical progression, and strategies to lower production or enhance clearance of pathological proteins in the degenerating brain have shown therapeutic promise in animal models and clinical trials (Boland et al., 2018; Li et al., 2019; Tabrizi et al., 2019).

Post-mortem histopathological analyses (Braak and Braak, 1991; Braak et al., 2003; Brettschneider et al., 2013) and *in vivo* imaging studies (Deng et al., 2004; Poudel et al., 2019) indicate that proteopathic lesions associated with neurodegeneration appear in highly-reproducible and disease-specific spatiotemporal patterns through the brain. Interestingly, these patterns largely follow neuroanatomical tracts (Ahmed et al., 2014; Mezias et al., 2017), suggesting a central role for synaptic connectivity in pathological aggregate spreading. Accumulating evidence supports the idea that intracellular aggregates formed by tau, α-synuclein, TDP-43, SOD1, and mutant huntingtin (mHtt) transfer from cell to cell and self-replicate by recruiting natively-folded versions of the same protein, analogous to how infectious prion protein (PrP^{Sc}) templates the conformational change of soluble PrP^C in prion diseases (Vaquer-Alicea and Diamond, 2019). Numerous studies have pointed to roles for endocytosis and exocytosis (Asai et al., 2015; Babcock and Ganetzky, 2015; Chen et al., 2019; Holmes et al., 2013; Lee et al., 2010; Zeineddine et al., 2015), membrane permeabilization (Chen et al., 2019; Falcon et al., 2018; Flavin et al., 2017; Zeineddine et al., 2015), tunneling nanotubes (Costanzo et al., 2013; Sharma and Subramaniam, 2019), and neuronal activity (Wu et al., 2016) in entry and/or exit of pathogenic protein assemblies from cells, but the exact mechanisms by which amyloid aggregates or

on-pathway intermediates cross one or more biological membranes in the highly complex central

nervous system (CNS) remain an enigma.

Glia are resident immune cells of the CNS and constantly survey the brain to maintain

homeostasis and respond rapidly to tissue damage or trauma. Reactive astrocytes and microglia

provide a first line of defense in neurodegeneration by infiltrating sites of neuronal injury, upregulating

immune-responsive genes, and phagocytosing dying neurons and other debris, including protein

aggregates (Asai et al., 2015; Grathwohl et al., 2009; Wyss-Coray et al., 2003). Prolonged activation of

these glial responses results in chronic inflammation, exacerbating synaptic dysfunction and neuronal

loss (Hammond et al., 2018). We and others have previously shown that Draper, a *Drosophila*

scavenger receptor that recognizes and phagocytoses cellular debris (Freeman, 2015), regulates the

load of mHtt (Pearce et al., 2015) and $A\beta_{1-42}$ (Ray et al., 2017) aggregate pathology in the fly CNS.

Remarkably, we also found that a portion of phagocytosed neuronal mHtt aggregates gain entry into

the glial cytoplasm and once there, nucleate the aggregation of normally-soluble wild-type Htt (wtHtt)

proteins, suggesting that glial phagocytosis provides a path for spreading of prion-like aggregates in

intact brains. Consistent with these findings, microglial ablation suppresses pathological tau

transmission between synaptically-connected regions of the mouse brain (Asai et al., 2015), and PrP^{Sc}

transfers from infected astrocytes to co-cultured neurons (Victoria et al., 2016). Thus, phagocytic glia

may play double-edged roles in neurodegeneration, with normally neuroprotective clearance

mechanisms also driving dissemination of prion-like aggregates through the brain.

A plethora of studies from the last decade have strengthened the prion-like hypothesis for

neurodegenerative diseases, but we still lack a clear understanding of how pathogenic protein

aggregates spread between cells in an intact CNS. In this study, we adapted our previously-described

Drosophila HD model to investigate roles for synaptic connectivity and phagocytic glia in prion-like mHtt

aggregate transmission in adult fly brains. HD is an autosomal dominant disorder caused by expansion

of a CAG repeat region in exon 1 of the Htt gene, resulting in production of highly aggregation-prone

mHtt proteins containing abnormally expanded polyglutamine (polyQ$\geq$37) tracts (Bates et al., 2015;

MacDonald et al., 1993). By contrast, wtHtt proteins containing polyQ$\leq$36 tracts only aggregate upon

nucleation by pre-formed Htt aggregate "seeds" (Chen et al., 2001; Preisinger et al., 1999). A growing

body of evidence from cell culture (Chen et al., 2001; Costanzo et al., 2013; Holmes et al., 2013; Ren et

al., 2009; Sharma and Subramaniam, 2019; Trevino et al., 2012) and *in vivo* (Ast et al., 2018; Babcock

and Ganetzky, 2015; Jeon et al., 2016; Masnata et al., 2019; Pearce et al., 2015; Pecho-Vrieseling et

al., 2014) models of HD supports the idea that pathogenic mHtt aggregates have prion-like properties—

they transfer from cell to cell and self-replicate by nucleating the aggregation of soluble wtHtt proteins.

Here, we report that mHtt aggregates formed in presynaptic olfactory receptor neuron (ORN) axons

effect prion-like conversion of wtHtt proteins expressed in the cytoplasm of postsynaptic partner

projection neurons (PNs) in the adult fly olfactory system. Remarkably, transfer of mHtt aggregates

from presynaptic ORNs to postsynaptic PNs was abolished in Draper-deficient animals and required

passage of the prion-like aggregate seeds through the cytoplasm of phagocytic glial cells. Together,

these findings support the conclusion that phagocytic glia are obligatory intermediates in prion-like

transmission of mHtt aggregates between synaptically-connected neurons *in vivo*, providing new insight

into key roles for glia in HD pathogenesis.

RESULTS

Prion-like transfer of mHtt aggregates between synaptically-connected neurons in the adult fly olfactory system

Aggregates formed by N-terminal fragments of mHtt generated by aberrant splicing (e.g., exon 1; Htt_{ex1}) (Sathasivam et al., 2013) or caspase cleavage (e.g., exon 1-12; Htt_{ex1-12}) (Graham et al., 2006) (Fig. 1A) accumulate in HD patient brains, are highly cytotoxic, and spread between cells in culture and *in vivo* (Babcock and Ganetzky, 2015; Costanzo et al., 2013; Pearce et al., 2015; Pecho-Vrieseling et al., 2014; Ren et al., 2009). We have previously established transgenic *Drosophila* that employ binary expression systems [e.g., Gal4-UAS, QF-QUAS, or LexA-LexAop (Riabinina and Potter, 2016)] to express fluorescent protein (FP) fusions of Htt_{ex1} in non-overlapping cell populations to monitor cell-to-cell transfer of $mHtt_{ex1}$ aggregates in intact brains (Donnelly and Pearce, 2018; Pearce et al., 2015). Our experimental approach (Fig. 1B) exploits the previously-reported finding that $wtHtt_{ex1}$ proteins aggregate upon physically encountering $mHtt_{ex1}$ aggregate seeds (Chen et al., 2001; Preisinger et al., 1999), such that transfer of $mHtt_{ex1}$ aggregates from "donor" cells is reported by conversion of cytoplasmic $wtHtt_{ex1}$ from its normally soluble, diffuse state to a punctate, aggregated state in "acceptor" cells (Fig. 1B-inset). To confirm that $mHtt_{ex1}$ nucleates the aggregation of $wtHtt_{ex1}$ in fly neurons, we co-expressed FP-fusions of these two proteins using pan-neuronal *elav[C155]-Gal4*. In flies expressing only mCherry-tagged $mHtt_{ex1}$ ($Htt_{ex1}Q91$-mCherry) pan-neuronally, aggregates were visible as discrete mCherry+ puncta throughout adult fly brains, with enrichment in neuropil regions (Fig. 1-figure supplement 1A). By contrast, GFP-tagged $wtHtt_{ex1}$ ($Htt_{ex1}Q25$-GFP) was expressed diffusely in the same regions of age-matched adult brains (Fig. 1-figure supplement 1B). Upon co-expression with $Htt_{ex1}Q91$-mCherry, $Htt_{ex1}Q25$-GFP was converted to a punctate expression pattern that almost entirely overlapped with $Htt_{ex1}Q91$-mCherry signal (Fig. 1-figure supplement 1C and F), whereas expression patterns of neither membrane-targeted GFP (mCD8-GFP) nor soluble GFP lacking a polyQ sequence were affected by the presence of $Htt_{ex1}Q91$-mCherry aggregates in neurons (Fig. 1-figure supplement 1D-F). Thus, in the fly CNS, $Htt_{ex1}Q91$ aggregates induce prion-like conversion of normally-soluble $Htt_{ex1}Q25$ via a homotypic nucleation reaction that requires the Htt_{ex1} sequence.

To examine trans-synaptic prion-like transfer of mHtt$_{ex1}$ aggregates, we coupled QF-driven

expression of Htt$_{ex1}$Q91-mCherry with Gal4-driven expression of Htt$_{ex1}$Q25-GFP in neuronal cell

populations that make well-defined synaptic connections in the adult fly olfactory system (Fig. 1B-D).

Htt$_{ex1}$Q91-mCherry was expressed using *Or67d-QF* in ~40 presynaptic ORNs ("DA1 ORNs") that

project axons from the antenna into the central brain, where they form synaptic connections with

dendrites of ~7 partner PNs ("DA1 PNs") in the DA1 glomerulus of the antennal lobe (Fig. 1B-inset, C,

and D) (Jefferis et al., 2001). In these same animals, Htt$_{ex1}$Q25-GFP was expressed in ~60% of PNs

using *GH146-Gal4*, which labels lateral and ventral DA1 PNs in addition to other PN types (Fig. 1C and

D) (Marin et al., 2002). We did not detect expression of Htt$_{ex1}$Q91-mCherry in DA1 ORNs until ~24 hr

before eclosion, consistent with activation of adult olfactory receptor gene expression during late pupal

development (Clyne et al., 1999), whereas Htt$_{ex1}$Q25-GFP was expressed in PNs via *GH146-Gal4*

earlier in development (Stocker et al., 1997). This genetic approach therefore enables us to monitor

prion-like transfer of mHtt aggregates between post-mitotic, synaptically-connected DA1 ORNs and

PNs in the adult fly brain.

Formation and prion-like transfer of mHtt$_{ex1}$ aggregates across DA1 ORN-PN synapses was

examined by monitoring the solubility of Htt$_{ex1}$Q91-mCherry and Htt$_{ex1}$Q25-GFP proteins in or near the

DA1 glomerulus (Fig. 1B-inset and E-G). Whereas Htt$_{ex1}$Q25-mCherry was expressed diffusely in DA1

ORN axons (Fig. 1C), aggregated Htt$_{ex1}$Q91-mCherry was visible as discrete puncta almost entirely

restricted to DA1 axons and axon termini in the DA1 region of the antennal lobe (Fig. 1D and E1-G1).

We used semi-automated 3D segmentation and reconstruction of high-magnification confocal z-stacks

(Fig. 1-figure supplement 2A1 and Video 1) to quantify Htt$_{ex1}$Q91 aggregate formation in the DA1

glomerulus over time. Htt$_{ex1}$Q91 aggregates first appeared in pharate adults and increased in number

as the flies aged (Fig. 1E1'-G1' and 1J). Numbers of Htt$_{ex1}$Q91 aggregates in males exceeded those in

females at each time point (Fig. 1J), consistent with known sexual dimorphism in DA1 glomerular

volume (Stockinger et al., 2005). In these same brains, Htt$_{ex1}$Q25-GFP was expressed diffusely

throughout GH146+ PN cell bodies and processes in young adults (Fig. 1D and E2), but bright

Htt$_{ex1}$Q25-GFP puncta began to appear and accumulate in the DA1 glomerulus as the flies aged (Fig.

1F2 and G2, *arrows*). Because these puncta could be difficult to distinguish from surrounding non-

aggregated $Htt_{ex1}Q25$-GFP signal, and GFP+ puncta representing normal dendritic architecture and/or

intracellular vesicles in the secretory pathway were visible in GH146+ PNs expressing mCD8-GFP (Fig.

1-figure supplement 2B2 and Fig. 1-figure supplement 3B), we defined $Htt_{ex1}Q25$ aggregates as GFP+

puncta that colocalized with $Htt_{ex1}Q91$ aggregates (Fig. 1-figure supplement 2A1 and C1-7, and Video

1). This approach reported similar results to manual quantification of the same data in 2D confocal

slices (Fig. 1-figure supplement 2C1-7 and D), and identical data were obtained when $Htt_{ex1}Q25$-GFP+

segmented surfaces were filtered for colocalization with $Htt_{ex1}Q91$-mCherry+ puncta (Fig. 1-figure

supplement 2A2 and D). By contrast, $Htt_{ex1}Q91$ aggregates in DA1 ORNs did not colocalize with

mCD8-GFP expressed in GH146+ PNs in control animals regardless of whether mCherry+ or GFP+

surfaces were initially segmented (Fig. 1-figure supplement 2B1-2). Thus, our semi-automated

approach to identify "$Htt_{ex1}Q91$+$Htt_{ex1}Q25$" aggregates specifically reports non-cell autonomous

conversion of postsynaptic $wtHtt_{ex1}$ by presynaptic $mHtt_{ex1}$ seeds (Fig. 1B-inset).

Numbers of $Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregates increased as flies expressing $Htt_{ex1}Q91$-mCherry in

DA1 ORNs and $Htt_{ex1}Q25$-GFP in GH146+ PNs aged from 1 to 21 days old (Fig. 1E2'-G2' and 1J).

$Htt_{ex1}Q91$ aggregates outnumbered $Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregates at each time point tested (Fig. 1J

and K), likely reflecting a higher rate of aggregate formation in "donor" ORNs expressing polyQ-

expanded $mHtt_{ex1}$ than in "acceptor" PNs, where $wtHtt_{ex1}$ proteins must be nucleated by $mHtt_{ex1}$ seeds

originating in other cells. In control experiments, soluble $Htt_{ex1}Q25$-mCherry expressed in DA1 ORNs

did not colocalize with $Htt_{ex1}Q25$-GFP expressed in PNs (Fig. 1-figure supplement 3A, E, and F), and

$Htt_{ex1}Q91$ aggregates in DA1 ORNs did not nucleate membrane-bound mCD8-GFP in PNs (Fig. 1-

figure supplement 3B, E, and F). $Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregates still formed when the Gal4 inhibitor

Gal80 was expressed in ORNs (Fig. 1-figure supplement 3C, E, and F) or when the QF repressor QS

was expressed in PNs (Fig. 1-figure supplement 3D-F), arguing strongly against co-expression of

$Htt_{ex1}Q91$-mCherry and $Htt_{ex1}Q25$-GFP using the highly-specific *Or67d-QF* and *GH146-Gal4* drivers.

Together, these findings indicate that $mHtt_{ex1}$ aggregates formed in presynaptic ORNs induce non-cell

autonomous, homotypic aggregation of $wtHtt_{ex1}$ expressed in the cytoplasm of postsynaptic PNs.

Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates were not detected in non-synaptically-connected GH146+ PNs,

including in glomeruli directly adjacent to where DA1 ORNs terminate, suggesting that prion-like

conversion of Htt$_{ex1}$Q25 in PNs by Htt$_{ex1}$Q91 aggregates in ORNs requires synaptic connectivity. To

examine whether Htt$_{ex1}$Q91 aggregates can also spread retrogradely across ORN-PN synapses, we

expressed Htt$_{ex1}$Q91-mCherry in PNs using *GH146-QF* and Htt$_{ex1}$Q25-GFP in all ORNs using *pebbled-*

Gal4. While many Htt$_{ex1}$Q91 aggregates were visible in PN soma, dendrites, and axons in adult brains

(Fig. 1-figure supplement 4A-C), we did not observe colocalization of Htt$_{ex1}$Q25-GFP and Htt$_{ex1}$Q91-

mCherry puncta within the antennal lobe neuropil in 1, 7, and 14 day-old adults (Fig. 1-figure

supplement 4A, B, and D), suggesting that Htt$_{ex1}$Q91 aggregate spreading across ORN-PN synapses is

restricted to or much more efficient in the anterograde direction. We also found that aggregates formed

by mHtt$_{ex1-12}$ caspase-6 cleavage products (RFP-Htt$_{ex1-12}$Q138) (Fig. 1A) (Graham et al., 2006), which

have previously been shown to spread from ORN axons to more distant (non-PN) neurons in the fly

CNS (Babcock and Ganetzky, 2015), also transfer anterogradely across DA1 ORN-PN synapses (Fig.

1H and I). Together, these findings indicate that multiple pathogenic N-terminal mHtt fragments share

the ability to spread trans-synaptically in *Drosophila* brains, and the sequences required for prion-like

conversion of wtHtt reside within Htt$_{ex1}$.

wtHtt$_{ex1}$ aggregates in PNs are seeded by smaller mHtt$_{ex1}$ aggregates originating in ORNs

Higher magnification examination of DA1 glomeruli in flies expressing Htt$_{ex1}$Q91-mCherry (Fig.

2A and B) or RFP-Htt$_{ex1-12}$Q138 (Fig. 2C and D) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs

revealed that most (>85%) Htt$_{ex1}$Q25-GFP puncta in DA1 PNs colocalized with Htt$_{ex1}$Q91-mCherry

aggregates. Segmentation in both the red (Fig 2A1'-D1') and green (Fig. 2A2'-D2') channels

demonstrated that GFP+ surfaces entirely surrounded associated mCherry+ or RFP+ surfaces in these

colocalized aggregates (Fig 2A3'-D3'). These data suggest that aggregated Htt$_{ex1}$Q91 or Htt$_{ex1-12}$Q138

proteins form the "core" of induced Htt$_{ex1}$Q25 aggregates and supports our hypothesis that wtHtt$_{ex1}$

solubility in PNs is altered upon direct physical interaction with pre-formed mHtt seeds. To further

examine molecular interactions between Htt$_{ex1}$Q91 and Htt$_{ex1}$Q25 proteins, we measured fluorescence

resonance energy transfer (FRET) in colocalized aggregates. Remarkably, positive FRET signal was

detected for all $Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregates analyzed by this method (Fig. 2E), indicating that the

FP tags fused to each of these proteins were in close molecular proximity (<10 nm apart), consistent

with direct physical contact.

Comparison of volumes for all segmented $Htt_{ex1}Q91$, $Htt_{ex1-12}Q138$, $Htt_{ex1}Q91$+$Htt_{ex1}Q25$, and

$Htt_{ex1-12}Q138$+$Htt_{ex1}Q25$ aggregates revealed that $Htt_{ex1}Q91$ and $Htt_{ex1-12}Q138$ aggregates increased in

size as the flies aged, most substantially during the first week of adulthood (Fig. 2F). Interestingly, the

mean volume of $Htt_{ex1}Q91$ or $Htt_{ex1-12}Q138$ surfaces that colocalized with $Htt_{ex1}Q25$ aggregates (herein

referred to as "seeding-competent" mHtt aggregates) was less than the mean volume of all $Htt_{ex1}Q91$ or

$Htt_{ex1-12}Q138$ surfaces, and these differences were statistically significant in older animals. When

aggregate volumes were analyzed across all time points, it became apparent that seeding-competent

$Htt_{ex1}Q91$ or $Htt_{ex1-12}Q138$ aggregates clustered in a significantly smaller-sized subpopulation (range =

0.1–3.5 μm^3; mean = 1.845 $\pm$ 0.068 μm^3 for $Htt_{ex1}Q91$ aggregates, mean = 1.293 $\pm$ 0.071 μm^3 for Htt_{ex1-}

$_{12}Q138$ aggregates), whereas the entire population of $Htt_{ex1}Q91$ or $Htt_{ex1-12}Q138$ aggregates was more

heterogeneously-sized (range = 0.1–15 μm^3; mean = 2.19 $\pm$ 0.024 μm^3 for $Htt_{ex1}Q91$ aggregates, mean

= 2.40 $\pm$ 0.048 μm^3 for $Htt_{ex1-12}Q138$ aggregates) (Fig. 2G and H). We previously reported that

$Htt_{ex1}Q25$ aggregates seeded in the cytoplasm of glial cells colocalized with a similarly smaller-sized

subpopulation of seeding-competent $Htt_{ex1}Q91$ aggregates from DA1 ORNs (Pearce et al., 2015), and

smaller $mHtt_{ex1}$ aggregates are associated with increased seeding-propensity and neurotoxicity in other

HD models (Ast et al., 2018; Chen et al., 2001). Taken together, these findings strongly suggest that

$mHtt_{ex1}$ or $mHtt_{ex1-12}$ aggregates formed in presynaptic ORNs effect prion-like conversion of $wtHtt_{ex1}$ in

the cytoplasm of postsynaptic PNs, and that mHtt aggregate transmissibility in the fly CNS is correlated

with smaller aggregate size.

$mHtt_{ex1}$ aggregate transfer is enhanced across silenced DA1 ORN-PN synapses

Endocytosis, exocytosis, and neuronal activity have been previously implicated in neuron-to-

neuron spreading of mHtt and other pathogenic aggregates (Babcock and Ganetzky, 2015; Pecho-

Vrieseling et al., 2014; Wu et al., 2016), but it is not known how these processes contribute to

aggregate transfer across endogenous synapses *in vivo*. To examine a role for synaptic activity in DA1

ORN-to-PN transfer of mHtt$_{ex1}$ aggregates, we used well-established fly genetic tools that block fission

or fusion of synaptic vesicles at the presynaptic membrane to impair neurotransmission. First, we used

shibire^{ts1} (*shi^{ts1}*) (Kosaka and Ikeda, 1983), a temperature-sensitive mutant of the GTPase

Shibire/dynamin that blocks endocytic recycling of synaptic vesicles in flies raised at the restrictive

temperature. Co-expression of shi^{ts1} with Htt$_{ex1}$Q91-mCherry in DA1 ORNs in flies shifted from the

permissive temperature (18°C) to the restrictive temperature (31°C) in adulthood had no effect or

slightly decreased numbers of Htt$_{ex1}$Q91 aggregates in the DA1 glomerulus (Fig. 3A-C), but,

surprisingly, strongly enhanced formation of seeded Htt$_{ex1}$Q25 aggregates in DA1 PN dendrites

compared with control flies expressing LacZ (Fig. 3A, B, and D). Likewise, co-expression of Htt$_{ex1}$Q91-

mCherry with tetanus toxin light chain (TeTxLC), which inhibits SNARE-mediated fusion of synaptic

vesicles with presynaptic membranes (Sweeney et al., 1995), strongly increased numbers of seeded

Htt$_{ex1}$Q25 aggregates in the DA1 glomerulus (Fig. 3E, F, and H). At some time points, TeTxLC co-

expression led to increased numbers of Htt$_{ex1}$Q91 aggregates compared with control animals (Fig. 3E-

G); however, there appeared to be no correlation between numbers of Htt$_{ex1}$Q91 and

Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates in the DA1 glomerulus over time, so the increased numbers of seeded

Htt$_{ex1}$Q25 aggregates were unlikely to be simply due to abundance of presynaptic Htt$_{ex1}$Q91 seeds. To

manipulate neuronal activity by an alternative approach, we co-expressed the heat-activated

Drosophila transient-receptor potential A (dTrpA) channel (Hamada et al., 2008) with Htt$_{ex1}$Q91-

mCherry to thermogenetically stimulate DA1 ORNs. In adult flies shifted from ~21°C to 31°C upon

eclosion, dTrpA-mediated activation of Htt$_{ex1}$Q91-mCherry-expressing DA1 ORNs slightly increased

Htt$_{ex1}$Q91 aggregate numbers, but decreased formation of seeded Htt$_{ex1}$Q25 aggregates in the DA1

glomerulus compared with control flies expressing LacZ (Fig. 3I-L). Together, these results indicate that

prion-like transmission of Htt$_{ex1}$Q91 aggregates from ORNs to PNs is inversely correlated with

presynaptic ORN activity. These findings suggest that aggregate transfer could be enhanced across

dysfunctional synapses, which are an early pathological finding in HD and other neurodegenerative

diseases.

Our results using shi[ts1] and TeTxLC to block DA1 ORN activity contrast with previous reports

showing that spreading of Htt$_{ex1-12}$Q138 aggregates was inhibited from endocytosis- or exocytosis-

impaired ORNs to non-synaptically-connected neurons (Babcock and Ganetzky, 2015) or that

botulinum toxin inhibited spreading of mHtt$_{ex1}$ from R6/2 mouse brain slices to functionally-connected

human neurons (Pecho-Vrieseling et al., 2014). This discrepancy could be due to construct- or cell

type-specific effects or possibly different mechanisms regulating synaptic or non-synaptic aggregate

transmission in the brain. Thus, we wondered whether blocking Shibire-mediated endocytosis in ORNs

might create a more favorable environment for aggregate transfer across endogenous synapses in our

HD model. To test this, we first quantified Htt$_{ex1}$Q91-mCherry-expressing DA1 ORN axonal surfaces

using mCD8-GFP, a tool widely used to label neuronal cell bodies and processes (Lee and Luo, 1999;

Mosca and Luo, 2014) and to quantify neuron or neurite abundance (Burr et al., 2014; MacDonald et

al., 2006) in fly brains. Segmentation and 3D reconstruction of mCD8-GFP+ DA1 ORN surfaces

revealed that fluorescence intensity of and volume occupied by DA1 ORN axons were increased ~2-

fold in shi[ts1]-expressing flies at the restrictive temperature compared with controls expressing LacZ (Fig.

4A-C). This effect appeared to be specific since Htt$_{ex1}$Q91 aggregate abundance in DA1 ORNs was not

increased by shi[ts] co-expression (Fig. 3C), and may reflect a homeostatic compensatory response to

endocytic blockade (Davis, 2013; Dickman et al., 2006) that could create additional exit sites for

Htt$_{ex1}$Q91 aggregates.

To test whether inhibition of Shibire-mediated endocytosis in ORNs has similar effects on ORN-

to-glia transfer of Htt$_{ex1}$Q91 aggregates (Pearce et al., 2015), we co-expressed shi[ts1] with Htt$_{ex1}$Q91-

mCherry in DA1 ORNs and monitored formation of seeded Htt$_{ex1}$Q25 aggregates in the glial cytoplasm.

Similar to effects of shi[ts1] on trans-synaptic Htt$_{ex1}$Q91 aggregate transfer, blocking Shibire-mediate

endocytosis in DA1 ORNs increased seeded Htt$_{ex1}$Q25 aggregate formation in glia without affecting

Htt$_{ex1}$Q91 aggregate numbers (Fig. 4D-G). Together, these findings suggest that shi[ts]-mediated

silencing of Htt$_{ex1}$Q91-mCherry-expressing DA1 ORNs alters axonal surface area and enhances prion-

like transfer of Htt$_{ex1}$Q91 aggregates from presynaptic ORN axons to the cytoplasm of both glia and

postsynaptic PNs.

Glial Draper is required for ORN-to-PN transfer and alters morphology of neuronal mHtt$_{ex1}$ aggregates

The parallel effects of shi^{ts1}-mediated endocytic blockade on transfer of Htt$_{ex1}$Q91 aggregates

from DA1 ORNs to DA1 PNs and to glia suggest that mHtt$_{ex1}$ aggregate spreading between these

different cell types is coordinated. Our prior work showed that ORN-to-glia transfer of mHtt$_{ex1}$

aggregates is strictly dependent on Draper (Pearce et al., 2015), a scavenger receptor responsible for

phagocytic engulfment and clearance of neuronal debris in the fly CNS and other tissues (Etchegaray

et al.; Han et al., 2014; Hoopfer et al., 2006; MacDonald et al., 2006). Therefore, we sought to

determine whether Draper-expressing phagocytic glia might play a role in transferring mHtt$_{ex1}$

aggregates from ORNs to PNs. To test this, we quantified numbers of Htt$_{ex1}$Q91 and seeded Htt$_{ex1}$Q25

aggregates in DA1 ORN axons and PN dendrites, respectively, in animals heterozygous or

homozygous for the *draper(drpr)^{Δ5}* null mutation (Freeman et al., 2003). We previously reported that

drpr knockout (KO) increased steady-state numbers of Htt$_{ex1}$Q91 aggregates in DA1 ORN axons

(Pearce et al., 2015); however, this effect was found to not be statistically significant between *drpr^{Δ5}*

heterozygotes and homozygotes in this study (Fig. 5A-C). We suspect this is the case for two reasons:

(a) our image segmentation parameters improve identification of very small aggregates, which we show

are more abundant when *drpr* is expressed at normal levels (Fig. 5E-G), and (b) the Htt$_{ex1}$Q91-mCherry

transgene used here expresses at lower levels than the transgene used in our previous study. Thus,

Htt$_{ex1}$Q91 aggregates initially form more slowly in DA1 ORN axons (compare numbers of Htt$_{ex1}$Q91

aggregates in young females in Fig. 1J vs. Fig. 7I; the latter experiment used the same higher-

expressing Htt$_{ex1}$Q91-mCherry transgene as in our prior study) and may be less affected by Draper

depletion. Strikingly though, *drpr* KO completely blocked formation of Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates

in postsynaptic PNs (Fig. 5A, B, and D), suggesting that Draper mediates trans-synaptic transfer of

mHtt$_{ex1}$ aggregates. This surprising effect of *drpr* KO on seeded Htt$_{ex1}$Q25 aggregate formation was

also observed when shi^{ts1} was co-expressed with Htt$_{ex1}$Q91-mCherry in ORNs (Fig. 5-figure supplement

1), confirming that enhanced transfer of $Htt_{ex1}Q91$ aggregates from shi^{ts1}-expressing ORNs occurs via

the same Draper-dependent mechanism. To rule out the possibility that *GH146-Gal4* drives expression

of $Htt_{ex1}Q25$-GFP in Draper+ glia, we used repo-Gal80 to inhibit Gal4 in all glia and saw no effect on

$Htt_{ex1}Q91$ or $Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregate numbers (Fig. 5-figure supplement 2A, B, E, and F).

Moreover, expression of *drpr*-specific siRNAs in PNs did not affect formation of either $Htt_{ex1}Q91$ or

$Htt_{ex1}Q91$+$Htt_{ex1}Q25$ aggregates (Fig. 5-figure supplement 2C-F), consistent with glia as the sole

source of *drpr* expression in the fly CNS. We also did not observe significant colocalization between

ORN-derived $Htt_{ex1}Q91$ aggregates and GFP-fusions of Atg8a (Juhász et al., 2008) or Lamp1

(Pulipparacharuvil et al., 2005) in glia (Fig. 5-figure supplement 3), suggesting that neuronal $Htt_{ex1}Q91$

aggregates do not seed $Htt_{ex1}Q25$ in glial lysosomes or autophagosomes.

These results point to an unexpected but central role for glial Draper in $Htt_{ex1}Q91$ aggregate

transfer between multiple cell types in the fly CNS, but how phagocytic glia could mediate spreading of

$Htt_{ex1}Q91$ aggregates across neuronal synapses was not immediately clear. Intriguingly, mean volume

of $Htt_{ex1}Q91$ aggregates in DA1 ORN axons was significantly increased in *drpr* KO animals compared

to wild-type controls (Fig. 5E, F, and G-inset), and the relative frequency of two aggregate

subpopulations shifted between these genotypes: in the absence of *drpr*, the abundance of a smaller-

sized subpopulation (~0.1-1 μm^3) decreased while a larger-sized subpopulation (~2.5-4.0 μm^3)

increased in abundance (Fig. 5E-G). Intriguingly, the smaller aggregate subpopulation correlated well

with the size of $Htt_{ex1}Q91$ aggregates associated with converted $Htt_{ex1}Q25$ in PNs (Fig. 2G) or glia

(Pearce et al., 2015), suggesting that phagocytic glia could at least in part mediate formation of smaller

seeding-competent Htt_{ex1} aggregates. However, molecular features other than size must regulate the

seeding capacity of $mHtt_{ex1}$ aggregates, since smaller-sized aggregates did not completely disappear in

drpr KO animals (Fig. 5G). Taken together, these results indicate that Draper-expressing phagocytic

glia mediate transfer of $Htt_{ex1}Q91$ aggregates across DA1 ORN-PN synapses, perhaps in part by

altering morphological features of neuronal $mHtt_{ex1}$ aggregates.

Caspase activation in ORNs is required for trans-synaptic transfer of $mHtt_{ex1}$ aggregates

Phagocytic glia engulf injured or degenerating neuronal processes and apoptotic cell corpses by

recognizing "eat me" signals exposed on these debris (Wilton et al., 2019), and mHtt expression

induces caspase-dependent apoptosis in fly and mammalian models of HD (Ahmed et al., 2014). Thus,

we asked whether $Htt_{ex1}Q91$-mCherry-expressing DA1 ORNs activate pathways that could stimulate

Draper-dependent transfer of aggregates to DA1 PNs. In $drpr^{\Delta 5}$ heterozygotes, expression of $Htt_{ex1}Q91$-

GFP in all ORNs did not significantly increase cleavage of *Drosophila* caspase-1 (Dcp-1) compared to

flies expressing $Htt_{ex1}Q25$-GFP (Fig. 6A, B, and E); however, Dcp-1 cleavage was significantly

increased in $Htt_{ex1}Q91$- vs. $Htt_{ex1}Q25$-expressing ORNs when *drpr* was knocked out (Fig. 6C-E). These

data suggest that glia efficiently clear $Htt_{ex1}Q91$-expressing ORN axons displaying Dcp-1-dependent

"eat me" signals via Draper-dependent phagocytosis. In addition, co-expression of the viral effector

caspase inhibitor p35 (Hay et al., 1994) with $Htt_{ex1}Q91$-mCherry in DA1 ORNs inhibited formation of

$Htt_{ex1}Q91+Htt_{ex1}Q25$ aggregates in DA1 PNs (Fig. 6F, G, J, and K, *squares*), indicating that inhibition of

apoptotic caspases in presynaptic ORNs phenocopies effects of *drpr* KO on $Htt_{ex1}Q91$ aggregate

transfer from ORNs to PNs. By contrast, co-expression of p35 with $Htt_{ex1}Q25$-GFP in PNs did not affect

numbers of $Htt_{ex1}Q91+Htt_{ex1}Q25$ aggregates in the DA1 glomerulus (Fig. 6H-K, *circles*), suggesting that

caspase-dependent signaling in PNs are not required for $Htt_{ex1}Q91$ aggregate spreading across ORN-

PN synapses. Together, these results suggest that signals mediated by apoptotic caspase activation in

DA1 ORNs, and not DA1 PNs, promote engulfment and trans-synaptic transfer of $Htt_{ex1}Q91$ aggregates

via phagocytic glia.

$mHtt_{ex1}$ aggregates transfer from ORNs to PNs via the glial cytoplasm

The strict requirement for glial Draper in prion-like transfer of $Htt_{ex1}Q91$ aggregates across ORN-

PN synapses can be explained by two non-mutually exclusive models. In one model (Fig. 7A, *route 1*),

$mHtt_{ex1}$ aggregates spread from presynaptic ORNs to postsynaptic PNs via a glial cytoplasmic

intermediate. This model is consistent with our previous finding that phagocytosed neuronal $mHtt_{ex1}$

aggregates gain entry to the glial cytoplasm to effect prion-like conversion of $Htt_{ex1}Q25$ (Pearce et al.,

2015). Alternatively (Fig. 7A, *route 2*), phagocytic glia could sculpt the synaptic environment in a way

that promotes transfer of mHtt$_{ex1}$ aggregates directly from ORN axons to PN dendrites. To distinguish

between these models, we generated transgenic flies that use three binary expression systems (i.e.,

QF-QUAS, Gal4-UAS, and LexA-LexAop) to independently express a uniquely-tagged Htt$_{ex1}$ transgene

in each of three cell populations: Htt$_{ex1}$Q91-mCherry in DA1 ORNs, Htt$_{ex1}$Q25-3xHA in repo+ glia, and

Htt$_{ex1}$Q25-YFP in GH146+ PNs (Fig. 7A). If Htt$_{ex1}$Q91-mCherry aggregates formed in presynaptic ORNs

transfer to postsynaptic PNs via the glial cytoplasm, Htt$_{ex1}$Q91-mCherry aggregate seeds should

template the aggregation first of Htt$_{ex1}$Q25-3xHA in glia and then Htt$_{ex1}$Q25-YFP in PNs, resulting in

appearance of triple-labeled mCherry+/3xHA+/YFP+ puncta in the DA1 glomerulus (Fig. 7A, *route 1*). A

transient double-labeled mCherry+/3xHA+ aggregate subpopulation that have accessed the glial

cytoplasm but not yet reached PNs might also be observed in this scenario. Alternatively, if Htt$_{ex1}$Q91-

mCherry aggregates do not access the glial cytoplasm *en route* to PNs, only double-labeled

mCherry+/3xHA+ and mCherry+/YFP+ aggregates would appear in the DA1 glomerulus (Fig. 7A, *route

2*).

Brains from transgenic flies expressing these differentially-tagged Htt$_{ex1}$ transgenes in ORNs,

glia, and PNs were analyzed, and expression patterns consistent with known morphologies of these cell

types in the antennal lobe were observed by confocal microscopy (Fig. 7B and C, and Figure 7-figure

supplement 1). In contrast to data in other figures, these samples required immunostaining to detect

expression of Htt$_{ex1}$Q25-3xHA and Htt$_{ex1}$Q25-YFP expression in glia and PNs, respectively. We found

that immunolabeled aggregate subtypes were not amenable to volumetric segmentation, and so we

instead manually quantified single-, double-, and triple-labeled Htt$_{ex1}$ aggregates in confocal slices and

used line scan intensity profiling to confirm colocalization. Htt$_{ex1}$Q91-mCherry expression in DA1 ORNs

was partially diffuse and partially punctate in young (0-1 day-old) adult flies (Fig. 7B, D, E, and I), but

became more punctate as the flies aged (Fig. 7C, F, G, and I). mCherry+/3xHA+ puncta representing

Htt$_{ex1}$Q91 aggregates that transferred from ORNs to glia first appeared in 2 day-old adults and

increased in number over the next ~24 hours (Fig. 7E, F, and J, *blue bars*). Remarkably,

mCherry+/3xHA+/YFP+ puncta also began to appear in 2 day-old adults and increased in abundance

until the flies were ~4 days old (Fig. 7F, G, and J, *striped bars*). The timing of appearance of these

different aggregate subtypes supports our hypothesis that $Htt_{ex1}Q91$ seeds originating in DA1 ORN

axons transit through the glial cytoplasm before reaching the cytoplasm of DA1 PN dendrites (Fig. 7A,

route 1). mCherry+/YFP+ puncta representing aggregates that had transferred from DA1 ORNs to DA1

PNs without accessing the glial cytoplasm were not detected in these brains, arguing against direct

transfer of $Htt_{ex1}Q91$-mCherry aggregates from ORNs to PNs (Fig. 7A, *route 2*). Control animals

expressing $Htt_{ex1}Q25$ proteins in paired combinations of ORNs, glia, or PNs (Figure 7-figure

supplement 1) confirmed that $Htt_{ex1}Q25$ aggregates in glia or PNs were only detected when $Htt_{ex1}Q91$

was expressed in ORNs. Remarkably, when we used RNAi to specifically knockdown *drpr* in glia,

numbers of $Htt_{ex1}Q91$-mCherry aggregates increased (Fig. 7H and I), while mCherry+/3xHA+ and

mCherry+/3xHA+/YFP+ aggregates significantly decreased or were absent in 7 day-old adults (Fig. 7H

and J). These data confirm that glial Draper mediates $Htt_{ex1}Q91$ aggregate transmission from ORN

axons to glia and across ORN-PN synapses. Taken together, these findings suggest that phagocytic

glia act as obligatory intermediates in unidirectional prion-like spreading of $mHtt_{ex1}$ aggregates from

presynaptic ORNs to postsynaptic PNs in *Drosophila* brains.

DISCUSSION

The hypothesis that prion-like spreading contributes to progression of protein aggregate pathology in HD and other neurodegenerative diseases is gaining considerable support, and yet we still understand very little about the mechanisms that underlie cell-to-cell spreading *in vivo*, the influences of cell- and tissue-specific vulnerability, and the relevance of aggregate transmission to disease progression. In this study, we report that mHtt$_{ex1}$ aggregates transfer anterogradely from presynaptic ORNs to postsynaptic PNs via an obligatory path through phagocytic glia in *Drosophila* brains. ORN-to-PN transmission of mHtt$_{ex1}$ aggregates was enhanced by blocking endocytosis and exocytosis and slowed by thermogenetically stimulating presynaptic ORNs, suggesting an inverse relationship between presynaptic activity and mHtt$_{ex1}$ aggregate spreading. mHtt$_{ex1}$ aggregate transfer across synapses was inhibited by blocking apoptosis in presynaptic neurons and required the Draper scavenger receptor, which we have previously reported to mediate phagocytic engulfment of mHtt$_{ex1}$ aggregates from ORN axons and prion-like conversion of wtHtt$_{ex1}$ in the glial cytoplasm (Pearce et al., 2015). Here, we expand our understanding of the role that glia play in prion-like diseases by showing that phagocytosed neuronal mHtt$_{ex1}$ aggregates transit through the cytoplasm of glia before reaching postsynaptic PNs. To the best of our knowledge, these findings are the first to uncover a role for a well-conserved phagocytic pathway in prion-like spreading of pathogenic aggregates between neurons *in vivo*.

The increased propensity of mHtt to aggregate as a result of polyQ expansion is held in check by the proteostasis network, and intrinsic differences in proteostatic capacity of neuronal subpopulations could underlie regional selectivity to inclusion body formation in HD brains (Margulis and Finkbeiner, 2014). However, accumulating evidence that mHtt aggregates have prion-like properties suggests that aggregate spreading could also contribute to this by propagating mHtt pathology through networks of synaptically-connected but selectively-vulnerable neurons. We find that mHtt$_{ex1}$ aggregate transmission across ORN-PN synapses occurs selectively in the anterograde direction and is inversely correlated with synaptic activity. Some of the earliest changes seen in HD patient brains involve loss of presynaptic cortical inputs to the striatum, where more prominent pathological findings (e.g., mHtt aggregate accumulation and massive loss of medium spiny neurons)

appear in later stages of disease (Reiner and Deng, 2018). In addition, selective silencing of mHtt in

both the cortex and striatum of BACHD mice inhibited striatal degeneration and motor phenotypes to a

greater extent that silencing in just the striatum (Wang et al., 2014), suggesting that mHtt-induced

toxicity in presynaptic cortical regions could play an important role in early HD development. Thus,

presynaptic dysfunction may be a driving force for mHtt aggregate spreading between synaptically-

connected regions of the brain. While our findings do not directly address secondary consequences of

mHtt$_{ex1}$ aggregate spreading across synapses, we identify a novel mechanism whereby glial responses

to pathological changes in synapses mediate spreading of toxic aggregates between neurons. How

aggregate spreading via glia impacts neuronal viability and functioning at the synaptic and/or circuit

level are important questions for future studies.

Glia survey the brain to maintain homeostasis and can rapidly switch between supportive and

reactive states in response to perturbations in CNS microenvironments. Upon sensing neuronal insult

or injury, reactive microglia and astrocytes undergo dramatic morphological, metabolic, and

transcriptional changes and promote neuronal survival by releasing trophic factors and phagocytosing

debris (Hammond et al., 2018). Central roles for phagocytic glia in neurodegeneration are becoming

increasingly recognized as genome-wide association studies and transcriptomic analyses identify glial

genes associated with increased disease risk. For example, rare variants in the microglial phagocytic

receptor gene *TREM2* are associated with increased risk of AD, FTD, and PD, and loss of TREM2

function exacerbates Aβ-, tau-, and α-synuclein-associated neurotoxicity (Griciuc et al.; Guo et al.,

2019; Leyns et al., 2019; Zhao et al., 2018). Disruptions in key glial phagocytic functions leads to

accumulation of potentially toxic aggregates in the brain (Asai et al., 2015; Hong et al., 2016; Pearce et

al., 2015; Ray et al., 2017; Wilton et al., 2019), and administration of antibodies targeting pathological

Aβ, tau, or α-synuclein proteins inhibits aggregate accumulation and spreading *in vivo* (Funk et al.,

2015; Masliah et al., 2005; Tran et al., 2014; Yanamandra et al., 2013). However, chronically-active or

otherwise dysfunctional glia induce neuroinflammation and exacerbate neuronal damage. For example,

hyperactivate microglia excessively engulf synapses in pre-plaque AD mouse brains (Hong et al., 2016)

and signal for formation of neurotoxic A1 astrocytes, which accumulate in several neurodegenerative

diseases (Liddelow et al., 2017; Yun et al., 2018) and during aging (Clarke et al., 2018). Our findings

thus add new insights to recognizing glia as double-edged players in neurodegeneration and suggest

that rebalancing the protective and harmful effects of glial phagocytosis could be an effective new

therapeutic strategy.

We provide several lines of evidence to support a model in which glia engulf $mHtt_{ex1}$

aggregates, or perhaps portions of $mHtt_{ex1}$ aggregate-containing axons, from ORNs without

internalizing elements of PNs: (i) $mHtt_{ex1}$ aggregate transmission occurred exclusively in the

anterograde direction across ORN-PN synapses, (ii) neuronal $mHtt_{ex1}$ aggregates did not colocalize

with markers of lysosomes or autophagosomes in glia, and (iii) caspase inhibition in ORNs and not in

PNs inhibited aggregate transfer. Our data suggest that glia selectively target presynaptic ORNs by

recognizing apoptotic "eat me" signals induced by $mHtt_{ex1}$ aggregate accumulation, and this process

drives aggregate transfer to postsynaptic PNs. In developing mouse brains, pruned presynaptic

structures are engulfed by microglia largely in the absence of postsynaptic markers (Schafer et al.,

2012; Weinhard et al., 2018), suggesting that mammalian glia have the ability to selectively "nibble"

presynaptic components (a fine-tuned phagocytic process known as trogocytosis). In addition,

astrocytes internalize dystrophic presynaptic terminals that accumulate near amyloid plaques in

transgenic mouse and human AD brains (Gomez-Arboledas et al., 2018). Intriguingly, astrocytes

expressing the mammalian Draper homolog MEGF10 and complement-dependent microglia

preferentially engulf "weaker" synapses to refine neural circuits in developing and adult mouse brains

(Chung et al., 2013; Schafer et al., 2012), and aberrant activation of these pathways could contribute to

early synaptic loss in neurodegenerative disease (Hong et al., 2016). We find that cell-to-cell $mHtt_{ex1}$

aggregate transfer is accelerated from silenced ORN axons, suggesting that activity-impaired synapses

are engulfed by nearby glia, enhancing spread along the ORN-to-glia-to-PN track. Thus, while the

primary objective of phagocytic glia may be to eliminate toxic neuronal debris from the CNS, aberrant

and/or excessive activation of phagocytic pathways could paradoxically promote disease. Premature

elimination of live synapses could also drive network dysfunction, suggesting that attenuating glial

responses could be an effective early intervention to preserve neurological functions in

neurodegenerative disease patients (Carpanini et al., 2019).

The most exciting yet unexpected finding we report here is that prion-like mHtt$_{ex1}$ aggregates do

not directly transfer from ORNs to PNs, but instead make an obligatory detour through the phagocytic

glial cytoplasm. This circuitous route could explain why relatively small numbers of seeded wtHtt$_{ex1}$

aggregates form in glia or PNs, since seeding-competent mHtt$_{ex1}$ must escape from multiple

degradative systems and from the membrane-bound phagolysosomal compartment in order to access

cytoplasmic wtHtt$_{ex1}$. Though transfer through glia is compulsory across ORN-PN synapses, our data

cannot exclude the possibility that mHtt$_{ex1}$ aggregates transfer directly between neurons in other

regions of the brain or at later stages of disease, when neuron-glia communication and cell integrity

could be severely compromised. However, our findings suggest that phagocytic glia drive aggregate

transfer across ORN-PN synapses by altering the transmissibility of mHtt$_{ex1}$ aggregates formed in

presynaptic ORN axons. We previously proposed that Draper-dependent phagocytosis could provide a

temporary conduit for engulfed mHtt$_{ex1}$ aggregates to escape into the glial cytoplasm (Pearce et al.,

2015). This idea is supported by work from others showing that α-synuclein, tau, and mHtt$_{ex1}$

aggregates rupture cell surface or endolysosomal membranes to access the cytoplasmic compartment

(Chen et al., 2019; Falcon et al., 2018; Flavin et al., 2017; Ren et al., 2009; Zeineddine et al., 2015). It

is possible that Draper-dependent phagocytosis could modify mHtt$_{ex1}$ aggregates to increase their

capacity to cross biological membranes, for example by altering molecular features such as rigidity,

frangibility, or size. Indeed, we report here that seeding-competent mHtt$_{ex1}$ aggregates belong to a

smaller-sized aggregate subpopulation whose abundance is regulated by Draper, and other groups

have observed that the seeding propensy of mHtt$_{ex1}$ (Ast et al., 2018; Chen et al., 2001) or tau (Wu et

al., 2013) aggregates is strongly associated with smaller size. This raises the intriguing possibility that

membrane fission or fusion events that occur during dynamin-mediated engulfment or phagosome

maturation could directly fragment or allow for escape of partially-digested neuronal mHtt$_{ex1}$ aggregates

that fall below an upper size limit for entry into the cytoplasm. Indeed, aggregate fragmentation and

secondary nucleation events are thought to be key components of prion-like propagation in many

neurodegenerative diseases (Knowles et al., 2014).

An outstanding question raised by our study is how ORN-derived mHtt$_{ex1}$ aggregates physically

transfer from glia to PN cytoplasms. This could be accomplished by a number of mechanisms already

proposed for cell-to-cell spreading of prion-like aggregates, such as transport through extracellular

vesicles or tunneling nanotubes (Costanzo et al., 2013; Sharma and Subramaniam, 2019),

endocytosis/exocytosis (Asai et al., 2015; Babcock and Ganetzky, 2015; Chen et al., 2019; Holmes et

al., 2013; Lee et al., 2010; Zeineddine et al., 2015), secretion or passive release of aggregates from

dying cells, and direct penetration of lipid bilayers (Brundin et al., 2010; Davis et al., 2018; Vaquer-

Alicea and Diamond, 2019). In a mouse model of tauopathy, pathological tau is transported between

anatomically-connected regions of the brain via exosomes secreted by microglia (Asai et al., 2015),

suggesting that phagocytosed neuronal aggregates may never encounter the extracellular space during

transfer. It is also possible that dysfunction caused by continuous aggregate internalization, genetic

mutations, and/or normal aging could decrease the efficiency by which glia clear aggregates, promoting

their spread. In support of this, extracellular Aβ or α-synuclein fibrils accumulate inside microglia

(Chung et al., 1999; Frackowiak et al., 1992), and the ability of microglia to effectively degrade

phagocytosed material declines with age (Bliederhaeuser et al., 2016; Tremblay et al., 2012). We

therefore favor a mechanism by which phagocytosed aggregates resistant to degradation overwhelm

the glial phagolysosomal system and promote aggregate release, either through active secretion (e.g.,

in an act of self-preservation) or during glial cell death. While we did not observe mHtt$_{ex1}$ aggregate

transmission to non-partner PNs in flies up to 3 weeks old, suggesting that aggregate transfer at these

ages is restricted to synaptic regions ensheathed by only 1-2 glial cells (MacDonald et al., 2006; Wu et

al., 2017), important remaining questions are whether aggregates that have invaded the glial cytoplasm

can transfer to other cells in or near the DA1 glomerulus (e.g., other glia or local interneurons) or to

downstream neurons in the olfactory circuit.

In summary, our data demonstrate that phagocytic glia are active participants in the spread of

prion-like protein aggregates between synaptically-connected neurons *in vivo*. Since microglial

processes are highly motile (Hammond et al., 2018), these findings raise the intriguing possibility that

phagocytic glia could mediate not only trans-synaptic, but also long-range spreading of neuronal

aggregate pathology in the mammalian CNS. Our findings have important implications for

understanding complex relationships between aggregate-induced cytotoxicity and neuron-glia

communication in health and disease. Deciphering the mechanisms that regulate helpful vs. harmful

effects of phagocytic glia in the brain will help to reveal the therapeutic potential of targeting key glial

functions in HD and other neurodegenerative diseases.

MATERIALS AND METHODS

Fly husbandry

All fly stocks and crosses were raised on standard cornmeal/molasses media at 25°C, ~50% relative humidity, and on a 12 hr light/12 hr dark cycle, unless otherwise noted. The following drivers were used to genetically access different cell populations in the fly CNS: *elav[C155]-Gal4* (Lin and Goodman, 1994), *Or67d-QF* (Liang et al., 2013), *GH146-Gal4* (Stocker et al., 1997), *GH146-QF* (Potter et al., 2010), and *GH146-LexA::GAD* (Lai et al., 2008) (a kind gift from Tzumin Lee, Janelia Farms), *repo-Gal4* (Sepp et al., 2001), and *Or83b-Gal4* (Kreher et al., 2005). Transgenic flies previously described by our lab include *QUAS-Htt$_{ex1}$Q91-mCherry*, *QUAS-Htt$_{ex1}$Q25-mCherry*, *UAS-Htt$_{ex1}$Q91-mCherry*, *UAS-Htt$_{ex1}$Q25-GFP*, *UAS-Htt$_{ex1}$Q25-YFP*, and *UAS-GFP* transgenes inserted at the attP3 (1st chromosome) and/or attP24 (2nd chromosome) ϕC31 integration sites (Pearce et al., 2015). Other transgenic flies not generated in this study include: *UAS-mCD8-GFP* (BDSC #5137), *QUAS-nucLacZ* lines #7 and 44 (BDSC #30006 and 30007)*, QUAS-shi^{ts1}* lines #2, 5, and 7 (BDSC #30010 and 30012; kind gift from C. Potter, Johns Hopkins School of Medicine), *QUAS-TeTxLC* lines #4c and 9c and *QUAS-dTrpA* lines #5, 6, and 7 (kind gifts from O. Riabinina and C. Potter, Johns Hopkins), *QUAS-mCD8-GFP* line #5J (BDSC #30002), *QUAS-p35* (a kind gift from H. Steller, Rockefeller University), *UAS-LacZ* (BDSC #8529), *UAS-p35* (BDSC #5072)*, QUAS-Gal80* (BDSC #51948), *UAS-QS* (BDSC #30033), *repo-Gal80* (a kind gift from T. Clandinin, Stanford University), *UAS-FFLuc.VALIUM1* (BDSC #35789), *UAS-GFP-Lamp1* (BDSC #42714), *UAS-Atg8a-GFP* (BDSC #52005), and *UAS-Draper^{RNAi}* and *drpr^{Δ5}* mutant flies (kind gifts from M. Freeman, Vollum Institute). Genotypes for all flies used in this study are listed in Table 1.

Cloning and transgenesis

pUASTattB(Htt$_{ex1}$Q25-GFP) and pUASTattB(Htt$_{ex1}$Q25-3xHA) plasmids were generated by PCR amplification of tagged Htt$_{ex1}$ cDNAs from the pcDNA3 vector backbone and cloned into pUASTattB via XhoI and XbaI restriction sites. cDNA for mRFP-Htt$_{ex1-12}$Q138 (a kind gift from T. Littleton, MIT) (Weiss et al., 2012) was subcloned into the pQUASTattB plasmid (Riabinina et al., 2015) using EcoRI and XbaI

restriction sites. Htt$_{ex1}$Q25-YFP cDNA was amplified by PCR from the pUASTattB(Htt$_{ex1}$Q25-YFP)

plasmid (Pearce et al., 2015) and subcloned into the EcoRI site in pLOT downstream of LexA-

responsive LexAop DNA sequences. Plasmids were microinjected into embryos containing the attP3 X

chromosome (for UAS-Htt$_{ex1}$Q25-GFP), attP3B or attP8 X chromosome (for QUAS-mRFP-Htt$_{ex1-}$

$_{12}$Q138), or attP24 2nd chromosome (for UAS-Htt$_{ex1}$Q25-3xHA or LexAop-Htt$_{ex1}$Q25-YFP) ϕC31

integration sites either in-house or at BestGene, Inc (Chino Hills, CA).

Drosophila brain dissection and sample preparation

Adult fly brains were dissected, fixed, and stained and/or imaged as previously described

(Pearce et al., 2015). Briefly, brains were dissected in ice-cold phosphate-buffered saline containing

either 0.03% Triton X-100 (PBS/0.03T, when intrinsic fluorescence of FP-fusions was imaged) or 0.3%

Triton X-100 (PBS/0.3T, when indirect immunofluorescence was used to detect protein expression).

Where possible, we imaged using intrinsic GFP/YFP/mCherry fluorescence when Htt-FP fusions

expression levels were high enough to detect on the confocal. Dissected brains were transferred to

microfuge tubes containing PBS/T + 4% paraformaldehyde fixative solution on ice and then fixed in the

dark at room temperature (RT) for 5 min (when imaging intrinsic fluorescence) or 20 min (when using

immunofluorescence). For direct fluorescence imaging, brains were washed in PBS/0.03T buffer

several times before incubation in Slowfade Gold Antifade Mountant (Invitrogen, Carlsbad, CA) for at

least 1 hr at 4°C in the dark. For immunostaining, brains were washed several times in PBS/0.3T, and

then blocked in PBS/0.3T containing 5% normal goat serum (Lampire Biological Laboratories,

Pipersville, PA) for 30min at RT, followed by incubation in primary antibodies diluted in blocking solution

and incubation for 24-72 hr at 4°C in the dark. Brains were then washed in PBS/0.3T several times at

RT and incubated in secondary antibodies diluted in blocking solution for 20-24 hr at 4°C in the dark.

Following another set of washes in PBS/0.3T at RT, the brains were incubated in Slowfade mountant

for at least 16 hr at 4°C in the dark. Brains were then bridge-mounted in Slowfade mountant on glass

microscopy slides overlayed with #1.5 coverglass (22 x 22 mm), and edges were sealed using clear

nail polish.

Primary antibodies used in this study include rabbit anti-DsRed (1:2000; #632496; Takara Bio

USA, Inc., Mountain View, CA), rabbit anti-mCherry (1:500; #PA5-34974; Invitrogen, Carlsbad, CA),

chicken anti-GFP (1:500; #GFP-1020; Aves Labs, Tigard, OR), chicken anti-GFP (1:1000; #ab13970;

Abcam, Cambridge, UK), chicken anti-GFP (1:500; #A10262; Invitrogen, Carlsbad, CA), rat anti-HA

(1:100; clone 3F10; #11867423001; Roche, Basel, Switzerland), rabbit anti-cleaved Dcp-1 (1:100;

#9578S; Cell Signaling Technology, Danvers, MA), and mouse anti-Bruchpilot (1:100; clone nc82;

Developmental Studies Hybridoma Bank, Iowa City, IA). Secondary antibodies used include FITC-

conjugated donkey anti-chicken (1:200; #703-095-155; Jackson Immuno Research Labs, West Grove,

PA) and AlexaFluor 488, 568, or 647-conjugated goat anti-chicken, anti-mouse, anti-rabbit, or anti-rat

IgGs (1:250 each; Invitrogen, Carlsbad, CA).

Image acquisition

All data were collected on a Leica SP8 laser-scanning confocal system equipped with 405 nm,

488 nm, 561 nm, and 633 nm lasers and 40X 1.3NA or 63X 1.4NA oil objective lenses. Leica LAS X

software was used to establish optimal settings during each microscopy session and to collect optical z-

slices of whole-mounted brain samples with Nyquist sampling criteria. Optical zoom was used to further

magnify and establish regions of interest in each sample. For most images, confocal data were

collected from ~60 x 60 x 25 μm (*xyz*) stacks centered on a single DA1 glomerulus, which was located

using fluorescent signal from Htt$_{ex1}$ protein expressed in DA1 ORN terminals. Exceptions to this are

shown in Fig. 1-figure supplement 1A-E [250 x 250 x ~60 μm (*xyz*) stacks capturing fluorescence signal

in the anterior central brain] and Fig. 1C-D, Fig. 1-figure supplement 4A-B, and Fig. 6A-D [~150 x 150 x

30 μm (*xyz*) stacks of the anterior portion of a single antennal lobe].

Post-imaging analysis

Raw confocal data were analyzed in 2D using ImageJ/FIJI (NIH, Bethesda, MD) or in 3D using

Imaris (Bitplane, Zürich, Switzerland), and all quantitative data were analyzed independently by two

researchers blinded to the experimental conditions. For semi-automated quantification of aggregates in

DA1 glomeruli, raw confocal data were deconvolved to reduce blur, rendered in 3D, and cropped if

necessary to establish the region of interest for further analysis. mHtt fluorescence was segmented in

3D stacks using the "Surfaces" algorithm (surface detail set to 0.25 μm and background subtraction at

0.75 μm), with the "split touching objects" option selected, and seed point diameter was set to 0.85 μm.

Background thresholding and seed point classification were adjusted manually for each image to

optimize segmentation of heterogeneously-sized $Htt_{ex1}Q91$-mCherry puncta ("aggregates") and

minimize capturing of diffuse signal. These settings differed <5-10% among individual samples in the

same experiment. In rare cases (<5% of aggregates in each image), some larger aggregates were

aberrantly split or smaller aggregates in close proximity were incorrectly merged; in these cases, the

objects were manually unified or split using the software program. To quantify seeded $wtHtt_{ex1}$

aggregates, red mHtt surfaces that colocalized with $Htt_{ex1}Q25$-GFP were identified by applying a filter

for mean intensity in the green channel. Colocalizing aggregates were selected by adjusting the

threshold to capture discrete $Htt_{ex1}Q25$ puncta with high contrast compared to surrounding diffuse

signal. Segmentation of $Htt_{ex1}Q25$ fluorescence in Fig. 1-figure supplement 2, Fig. 2A-D, and Video 1

was carried out using the same settings described above in the green channel. To measure DA1 ORN

axon volume and intensity in Fig. 4A-C, mCD8-GFP fluorescence was segmented using the "Surfaces"

function in Imaris, with surface detail set to 0.2 μm and background subtraction at 3 μm. Detailed

surface measurements (e.g. volume or intensity) were calculated in Imaris, and the data were exported

to Excel (Microsoft Corporation, Redmond, WA) or Prism (GraphPad Software, San Diego, CA) for

further analysis.

For data shown in Fig. 7 and Fig. 7-figure supplement 1, indirect immunofluorescence was used

to detect $Htt_{ex1}Q25$-3xHA expression in glia and amplify low $Htt_{ex1}Q25$-YFP signal in PNs. We found

that semi-automated image segmentation as described above reported fewer than half of

immunolabeled colocalized aggregates identified by manual counting, likely because of poor antibody

penetration into the aggregate core and $Htt_{ex1}Q25$-YFP aggregates with low contrast that made these

data less amenable to 3D segmentation. Instead, we manually counted aggregates in these samples by

scanning individual z-slices and scoring discrete puncta with increased signal relative to adjacent

diffuse signal in one, two, or all three channels. Line scans confirmed colocalization of signals from the

different Htt_{ex1} proteins, as shown in Fig. 7D-H.

FRET analysis

$Htt_{ex1}Q25+Htt_{ex1}Q91$ colocalized aggregates were analyzed for FRET using the acceptor

photobleaching method as previously described (Pearce et al., 2015). Briefly, 28 individual aggregates

were analyzed by photobleaching the mCherry acceptor appended to $Htt_{ex1}Q91$ using a 561nm laser

set at 100% intensity and scanning until fluorescence was no longer detectable. Donor fluorescence

dequenching was measured by exciting GFP fused to $Htt_{ex1}Q25$ using a 488 nm laser set at 1%

intensity before and after acceptor photobleaching. mCherry fluorescence was also excited before and

after photobleaching with a 561 nm laser set at 1% intensity. Fluorescence emission was collected

between 500 and 550 nm for GFP and 610 and 700 nm for mCherry to generate before and after

images as shown in Fig. 2E. FRET efficiencies ($FRET_{eff}$) were calculated after background correction

using the equation $(GFP_{after}-GFP_{before})/(GFP_{after})$ x 100 and represented by a pixel-by-pixel $FRET_{eff}$

image generated using the FRETcalc plugin (Stepensky, 2007) in FIJI/ImageJ.

Statistical analyses

All quantified data were organized and analyzed in Excel or Prism 8. Quantifications in graphical

form are shown as mean $\pm$ SEM, except for frequency analyses, which are displayed in histograms.

Results of all statistical analyses are described in each Figure Legend. Sample size (*n*) for each figure

is indicated in Table 1 and was selected to yield sufficient statistical power ($\geq$ 5 biological replicates

from $\geq$ 3 brains for each condition; a single DA1 glomerulus represents one biological replicate).

Multiple statistical comparisons were performed using the following tests and post-hoc corrections

where appropriate: Student's *t*-tests for pairwise comparisons, or one-way or two-way ANOVA followed

by Sidak's or Tukey's multiple comparison tests for experiments involving $\geq$3 genotypes. Results of

these statistical tests are shown on each graph, and symbols used to indicate statistical significance

are defined in the Figure Legends.

ACKNOWLEDGMENTS

The authors would like to thank K. Chao, D. Luginbuhl, V. Reed, B. Temsamrit, and M. Warkala for technical assistance, G. Panning for help with analyzing detailed aggregate measurements, T.R. Clandinin, M.R. Freeman, H. Krämer, T. Lee, J.T. Littleton, C.J. Potter, and O. Riabinina for kindly sharing reagents, and members of the Kopito, Luo, and Pearce labs for many valuable discussions. This work was supported by grants from the NIH (R01-DC005982 to L.L., R01-NS042842 to R.R.K, and R03-AG063295 to M.M.P.P.), the W.W. Smith Charitable Trusts (to M.M.P.P.), the Pittsburgh Foundation (UN2018-98318 to M.M.P.P.), and start-up funds from University of the Sciences (to M.M.P.P.).

AUTHOR CONTRIBUTIONS

K.M.D., O.R.D., A.D.A.Z., G.E.P., and W.M.T. – Investigation, Validation, Visualization, Writing – review & editing. L.L. and R.R.K. – Conceptualization, Resources, Supervision, Writing – review & editing, Funding Acquisition. M.M.P.P. – Conceptualization, Methodology, Investigation, Validation, Formal Analysis, Visualization, Writing – original draft preparation, Writing – review & editing, Resources, Supervision, Project Administration, Funding Acquisition.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Ahmed, Z., Cooper, J., Murray, T.K., Garn, K., McNaughton, E., Clarke, H., Parhizkar, S., Ward, M.A., Cavallini, A., Jackson, S.*, et al.* (2014). A novel in vivo model of tau propagation with rapid and progressive neurofibrillary tangle pathology: the pattern of spread is determined by connectivity, not proximity. Acta neuropathologica *127*, 667-683.

Asai, H., Ikezu, S., Tsunoda, S., Medalla, M., Luebke, J., Haydar, T., Wolozin, B., Butovsky, O., Kügler, S., and Ikezu, T. (2015). Depletion of microglia and inhibition of exosome synthesis halt tau propagation. Nature neuroscience *18*, 1584.

Ast, A., Buntru, A., Schindler, F., Hasenkopf, R., Schulz, A., Brusendorf, L., Klockmeier, K., Grelle, G., McMahon, B., Niederlechner, H.*, et al.* (2018). mHTT Seeding Activity: A Marker of Disease Progression and Neurotoxicity in Models of Huntington's Disease. Molecular cell *71*, 675-688.e676.

Babcock, D.T., and Ganetzky, B. (2015). Transcellular spreading of huntingtin aggregates in the Drosophila brain. Proceedings of the National Academy of Sciences of the United States of America *112*, E5427-5433.

Baron, O., Boudi, A., Dias, C., Schilling, M., Nolle, A., Vizcay-Barrena, G., Rattray, I., Jungbluth, H., Scheper, W., Fleck, R.A.*, et al.* (2017). Stall in Canonical Autophagy-Lysosome Pathways Prompts Nucleophagy-Based Nuclear Breakdown in Neurodegeneration. Current biology *27*, 3626-3642.e3626.

Bates, G.P., Dorsey, R., Gusella, J.F., Hayden, M.R., Kay, C., Leavitt, B.R., Nance, M., Ross, C.A., Scahill, R.I., Wetzel, R.*, et al.* (2015). Huntington disease. Nature reviews Disease primers *1*, 15005.

Bliederhaeuser, C., Grozdanov, V., Speidel, A., Zondler, L., Ruf, W.P., Bayer, H., Kiechle, M., Feiler, M.S., Freischmidt, A., Brenner, D.*, et al.* (2016). Age-dependent defects of alpha-synuclein oligomer uptake in microglia and monocytes. Acta neuropathologica *131*, 379-391.

Boland, B., Yu, W.H., Corti, O., Mollereau, B., Henriques, A., Bezard, E., Pastores, G.M., Rubinsztein,

D.C., Nixon, R.A., Duchen, M.R.*, et al.* (2018). Promoting the clearance of neurotoxic proteins in

neurodegenerative disorders of ageing. Nature reviews Drug discovery *17*, 660-688.

Braak, H., and Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes. Acta

neuropathologica *82*, 239-259.

Braak, H., Del Tredici, K., Rub, U., de Vos, R.A., Jansen Steur, E.N., and Braak, E. (2003). Staging of

brain pathology related to sporadic Parkinson's disease. Neurobiology of aging *24*, 197-211.

Brettschneider, J., Del Tredici, K., Toledo, J.B., Robinson, J.L., Irwin, D.J., Grossman, M., Suh, E., Van

Deerlin, V.M., Wood, E.M., Baek, Y.*, et al.* (2013). Stages of pTDP-43 pathology in amyotrophic

lateral sclerosis. Annals of neurology *74*, 20-38.

Brundin, P., Melki, R., and Kopito, R. (2010). Prion-like transmission of protein aggregates in

neurodegenerative diseases. Nature reviews Molecular cell biology *11*, 301-307.

Burr, A.A., Tsou, W.-L., Ristic, G., and Todi, S.V. (2014). Using Membrane-Targeted Green Fluorescent

Protein To Monitor Neurotoxic Protein-Dependent Degeneration of Drosophila Eyes. Journal of

neuroscience research *92*, 1100-1109.

Carpanini, S.M., Torvell, M., Morgan, B.P. (2019). Therapeutic inhibition of the complement system in

diseases of the central nervous system. Frontiers in Immunology *10*, 362.

Chen, J.J., Nathaniel, D.L., Raghavan, P., Nelson, M., Tian, R., Tse, E., Hong, J.Y., See, S.K., Mok,

S.A., Hein, M.Y.*, et al.* (2019). Compromised function of the ESCRT pathway promotes

endolysosomal escape of tau seeds and propagation of tau aggregation. The Journal of

biological chemistry. Pii: jbc.RA119.009432.

Chen, S., Berthelier, V., Yang, W., and Wetzel, R. (2001). Polyglutamine aggregation behavior in vitro

supports a recruitment mechanism of cytotoxicity. Journal of molecular biology *311*, 173-182.

Chung, H., Brazil, M.I., Soe, T.T., and Maxfield, F.R. (1999). Uptake, degradation, and release of

fibrillar and soluble forms of Alzheimer's amyloid beta-peptide by microglial cells. The Journal of

biological chemistry *274*, 32301-32308.

Chung, W.-S., Clarke, L.E., Wang, G.X., Stafford, B.K., Sher, A., Chakraborty, C., Joung, J., Foo, L.C.,
Thompson, A., Chen, C.*, et al.* (2013). Astrocytes mediate synapse elimination through
MEGF10 and MERTK pathways. Nature *504*, 394-400.

Clarke, L.E., Liddelow, S.A., Chakraborty, C., Munch, A.E., Heiman, M., and Barres, B.A. (2018).
Normal aging induces A1-like astrocyte reactivity. Proceedings of the National Academy of
Sciences of the United States of America *115*, E1896-e1905.

Clyne, P.J., Warr, C.G., Freeman, M.R., Lessing, D., Kim, J., and Carlson, J.R. (1999). A novel family
of divergent seven-transmembrane proteins: candidate odorant receptors in Drosophila. Neuron
22, 327-338.

Costanzo, M., Abounit, S., Marzo, L., Danckaert, A., Chamoun, Z., Roux, P., and Zurzolo, C. (2013).
Transfer of polyglutamine aggregates in neuronal cells occurs in tunneling nanotubes. Journal
of cell science *126*, 3678-3685.

Davis, G.W. (2013) Homeostatic signaling and the stabilization of neural function. Neuron *80*, 718-728.

Davis, A.A., Leyns, C.E.G., and Holtzman, D.M. (2018). Intercellular Spread of Protein Aggregates in
Neurodegenerative Disease. Annual review of cell and developmental biology *34*, 545-568.

Deng, Y.P., Albin, R.L., Penney, J.B., Young, A.B., Anderson, K.D., and Reiner, A. (2004). Differential
loss of striatal projection systems in Huntington's disease: a quantitative immunohistochemical
study. Journal of chemical neuroanatomy *27*, 143-164.

Dickman D.K., Lu, Z., Meinertzhagen, I.A., and Schwarz, T.L. (2006). Altered synaptic development
and active zone spacing in endocytosis mutants. Current biology *16*, 591-598.

Donnelly, K.M., and Pearce, M.M.P. (2018). Monitoring Cell-to-cell Transmission of Prion-like Protein
Aggregates in Drosophila Melanogaster. Journal of visualized experiments *12*, 133.

Etchegaray, J.I., Elguero, E.J., Tran, J.A., Sinatra, V., Feany, M.B., and McCall, K. (2016). Defective
Phagocytic Corpse Processing Results in Neurodegeneration and Can Be Rescued by TORC1
Activation. The Journal of neuroscience : the official journal of the Society for Neuroscience *36*,
3170-3183.

Falcon, B., Noad, J., McMahon, H., Randow, F., and Goedert, M. (2018). Galectin-8-mediated selective

autophagy protects against seeded tau aggregation. The Journal of biological chemistry *293*,

2438-2451.

Flavin, W.P., Bousset, L., Green, Z.C., Chu, Y., Skarpathiotis, S., Chaney, M.J., Kordower, J.H., Melki,

R., and Campbell, E.M. (2017). Endocytic vesicle rupture is a conserved mechanism of cellular

invasion by amyloid proteins. Acta neuropathologica *134*, 629-653.

Frackowiak, J., Wisniewski, H.M., Wegiel, J., Merz, G.S., Iqbal, K., and Wang, K.C. (1992).

Ultrastructure of the microglia that phagocytose amyloid and the microglia that produce beta-

amyloid fibrils. Acta neuropathologica *84*, 225-233.

Freeman, M.R. (2015). Drosophila Central Nervous System Glia. Cold Spring Harbor perspectives in

biology *7*, a020552.

Freeman, M.R., Delrow, J., Kim, J., Johnson, E., and Doe, C.Q. (2003). Unwrapping glial biology: Gcm

target genes regulating glial development, diversification, and function. Neuron *38*, 567-580.

Funk, K.E., Mirbaha, H., Jiang, H., Holtzman, D.M., and Diamond, M.I. (2015). Distinct Therapeutic

Mechanisms of Tau Antibodies: Promoting Microglial Clearance Versus Blocking Neuronal

Uptake. The Journal of biological chemistry *290*, 21652-21662.

Gomez-Arboledas, A., Davila, J.C., Sanchez-Mejias, E., Navarro, V., Nuñez-Diaz, C., Sanchez-Varo,

R., Sanchez-Mico, M.V., Trujillo-Estrada, L., Fernandez-Valenzuela, J.J., Vizuete, M., *et al.*

(2018). Phagocytic clearance of presynaptic dystrophies by reactive astrocytes in Alzheimer's

disease. Glia *66*, 637-653.

Graham, R.K., Deng, Y., Slow, E.J., Haigh, B., Bissada, N., Lu, G., Pearson, J., Shehadeh, J., Bertram,

L., Murphy, Z.*, et al.* (2006). Cleavage at the caspase-6 site is required for neuronal dysfunction

and degeneration due to mutant huntingtin. Cell *125*, 1179-1191.

Grathwohl, S.A., Kalin, R.E., Bolmont, T., Prokop, S., Winkelmann, G., Kaeser, S.A., Odenthal, J.,

Radde, R., Eldh, T., Gandy, S.*, et al.* (2009). Formation and maintenance of Alzheimer's

disease beta-amyloid plaques in the absence of microglia. Nature neuroscience *12*, 1361-1363.

Griciuc, A., Patel, S., Federico, A.N., Choi, S.H., Innes, B.J., Oram, M.K., Cereghetti, G., McGinty, D., Anselmo, A., Sadreyev, R.I., *et al.* (2019). TREM2 Acts Downstream of CD33 in Modulating Microglial Pathology in Alzheimer's Disease. Neuron *103*, 820-835.e827.

Guo, Y., Wei, X., Yan, H., Qin, Y., Yan, S., Liu, J., Zhao, Y., Jiang, F., and Lou, H. (2019). TREM2 deficiency aggravates alpha-synuclein-induced neurodegeneration and neuroinflammation in Parkinson's disease models. FASEB journal *33*, 12164-12174.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and Garrity, P.A. (2008). An internal thermal sensor controlling temperature preference in Drosophila. Nature *454*, 217-220.

Hammond, T.R., Robinton, D., and Stevens, B. (2018). Microglia and the Brain: Complementary Partners in Development and Disease. Annual review of cell and developmental biology *34*, 523-544.

Han, C., Song, Y., Xiao, H., Wang, D., Franc, N.C., Jan, L.Y., and Jan, Y.N. (2014). Epidermal cells are the primary phagocytes in the fragmentation and clearance of degenerating dendrites in Drosophila. Neuron *81*, 544-560.

Hay, B.A., Wolff, T., and Rubin, G.M. (1994). Expression of baculovirus P35 prevents cell death in Drosophila. Development *120*, 2121-2129.

Holmes, B.B., DeVos, S.L., Kfoury, N., Li, M., Jacks, R., Yanamandra, K., Ouidja, M.O., Brodsky, F.M., Marasa, J., Bagchi, D.P., *et al.* (2013). Heparan sulfate proteoglycans mediate internalization and propagation of specific proteopathic seeds. Proceedings of the National Academy of Sciences of the United States of America *110*, E3138-3147.

Hong, S., Beja-Glasser, V.F., Nfonoyim, B.M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K.M., Shi, Q., Rosenthal, A., Barres, B.A., *et al.* (2016). Complement and microglia mediate early synapse loss in Alzheimer mouse models. Science *352*, 712-716.

Hoopfer, E.D., McLaughlin, T., Watts, R.J., Schuldiner, O., O'Leary, D.D., and Luo, L. (2006). Wlds protection distinguishes axon degeneration following injury from naturally occurring developmental pruning. Neuron *50*, 883-895.

831   Jefferis, G.S., Marin, E.C., Stocker, R.F., and Luo, L. (2001). Target neuron prespecification in the

832   olfactory map of Drosophila. Nature *414*, 204-208.

833   Jeon, I., Cicchetti, F., Cisbani, G., Lee, S., Li, E., Bae, J., Lee, N., Li, L., Im, W., Kim, M.*, et al.* (2016).

834   Human-to-mouse prion-like propagation of mutant huntingtin protein. Acta neuropathologica

835   *132*, 577-592.

836   Juhász, G., Hill, J.H., Yan, Y., Sass, M., Baehrecke, E.H., Backer, J.M., and Neufeld, T.P. (2008). The

837   class III PI(3)K Vps34 promotes autophagy and endocytosis but not TOR signaling in

838   Drosophila. The Journal of cell biology *181*, 655-666.

839   Knowles, T.P., Vendruscolo, M., and Dobson, C.M. (2014). The amyloid state and its association with

840   protein misfolding diseases. Nature reviews Molecular cell biology *15*, 384-396.

841   Kosaka, T., and Ikeda, K. (1983). Possible temperature-dependent blockage of synaptic vesicle

842   recycling induced by a single gene mutation in Drosophila. Journal of neurobiology *14*, 207-225.

843   Kreher, S.A., Kwon, J.Y., and Carlson, J.R. (2005). The molecular basis of odor coding in the

844   Drosophila larva. Neuron *46*, 445-456.

845   Lai, S.-L., Awasaki, T., Ito, K., and Lee, T. (2008). Clonal analysis of Drosophila antennal

846   lobe neurons: diverse neuronal architectures in the lateral neuroblast lineage. Development

847   *135*, 2883-2893.

848   Lee, H.-J., Suk, J.-E., Patrick, C., Bae, E.-J., Cho, J.-H., Rho, S., Hwang, D., Masliah, E., and Lee, S.-J.

849   (2010). Direct Transfer of α-Synuclein from Neuron to Astroglia Causes Inflammatory

850   Responses in Synucleinopathies. Journal of Biological Chemistry *285*, 9262-9272.

851   Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function

852   in neuronal morphogenesis. Neuron *22*, 451-461.

853   Leyns, C.E.G., Gratuze, M., Narasimhan, S., Jain, N., Koscal, L.J., Jiang, H., Manis, M., Colonna, M.,

854   Lee, V.M.Y., Ulrich, J.D.*, et al.* (2019). TREM2 function impedes tau seeding in neuritic plaques.

855   Nature neuroscience *22*, 1217-1222.

Li, Z., Wang, C., Wang, Z., Zhu, C., Li, J., Sha, T., Ma, L., Gao, C., Yang, Y., Sun, Y.*, et al.* (2019).

Allele-selective lowering of mutant HTT protein by HTT–LC3 linker compounds. Nature *575*,

203-209.

Liang, L., Li, Y., Potter, C.J., Yizhar, O., Deisseroth, K., Tsien, R.W., and Luo, L. (2013). GABAergic

projection neurons route selective olfactory inputs to specific higher-order neurons. Neuron *79*,

917-931.

Liddelow, S.A., Guttenplan, K.A., Clarke, L.E., Bennett, F.C., Bohlen, C.J., Schirmer, L., Bennett, M.L.,

Munch, A.E., Chung, W.S., Peterson, T.C.*, et al.* (2017). Neurotoxic reactive astrocytes are

induced by activated microglia. Nature *541*, 481-487.

Lin, D.M., and Goodman, C.S. (1994). Ectopic and increased expression of Fasciclin II alters

motoneuron growth cone guidance. Neuron *13*, 507-523.

MacDonald, J.M., Beach, M.G., Porpiglia, E., Sheehan, A.E., Watts, R.J., and Freeman, M.R. (2006).

The Drosophila cell corpse engulfment receptor Draper mediates glial clearance of severed

axons. Neuron *50*, 869-881.

MacDonald, M.E., Ambrose, C.M., Duyao, M.P., Myers, R.H., Lin, C., Srinidhi, L., Barnes, G., Taylor,

S.A., James, M., Groot, N.*, et al.* (1993). A novel gene containing a trinucleotide repeat that is

expanded and unstable on Huntington's disease chromosomes. Cell *72*, 971-983.

Margulis, J. and Finkbeiner, S. (2014). Proteostasis in striatal cells and selective neurodegeneration in

Huntington's disease. Frontiers in Cellular Neuroscience *8*, 218.

Marin, E.C., Jefferis, G.S., Komiyama, T., Zhu, H., and Luo, L. (2002). Representation of the glomerular

olfactory map in the Drosophila brain. Cell *109*, 243-255.

Masliah, E., Rockenstein, E., Adame, A., Alford, M., Crews, L., Hashimoto, M., Seubert, P., Lee, M.,

Goldstein, J., Chilcote, T.*, et al.* (2005). Effects of alpha-synuclein immunization in a mouse

model of Parkinson's disease. Neuron *46*, 857-868.

Masnata, M., Sciacca, G., Maxan, A., Bousset, L., Denis, H.L., Lauruol, F., David, L., Saint-Pierre, M.,

Kordower, J.H., Melki, R.*, et al.* (2019). Demonstration of prion-like properties of mutant

huntingtin fibrils in both in vitro and in vivo paradigms. Acta neuropathologica *137*, 981-1001.

Mezias, C., LoCastro, E., Xia, C., and Raj, A. (2017). Connectivity, not region-intrinsic properties,

predicts regional vulnerability to progressive tau pathology in mouse models of disease. Acta

neuropathologica communications *5*, 61.

Mosca, T.J., and Luo, L. (2014). Synaptic organization of the Drosophila antennal lobe and its

regulation by the Teneurins. Elife *3*, e03726.

Pearce, M.M., Spartz, E.J., Hong, W., Luo, L., and Kopito, R.R. (2015). Prion-like transmission of

neuronal huntingtin aggregates to phagocytic glia in the Drosophila brain. Nature

communications *6*, 6768.

Pecho-Vrieseling, E., Rieker, C., Fuchs, S., Bleckmann, D., Esposito, M.S., Botta, P., Goldstein, C.,

Bernhard, M., Galimberti, I., Muller, M*., et al.* (2014). Transneuronal propagation of mutant

huntingtin contributes to non-cell autonomous pathology in neurons. Nature neuroscience *17*,

1064-1072.

Potter, C.J., Tasic, B., Russler, E.V., Liang, L., and Luo, L. (2010). The Q system: a repressible binary

system for transgene expression, lineage tracing, and mosaic analysis. Cell *141*, 536-548.

Poudel, G.R., Harding, I.H., Egan, G.F., and Georgiou-Karistianis, N. (2019). Network spread

determines severity of degeneration and disconnection in Huntington's disease. Human brain

mapping *40*, 4192-4201.

Preisinger, E., Jordan, B.M., Kazantsev, A., and Housman, D. (1999). Evidence for a recruitment and

sequestration mechanism in Huntington's disease. Philosophical transactions of the Royal

Society of London Series B, Biological sciences *354*, 1029-1034.

Pulipparacharuvil, S., Akbar, M.A., Ray, S., Sevrioukov, E.A., Haberman, A.S., Rohrer, J., and Krämer,

H. (2005). Drosophila Vps16A is required for trafficking to lysosomes and

biogenesis of pigment granules. Journal of cell science *118*, 3663-3673.

Ray, A., Speese, S.D., and Logan, M.A. (2017). Glial Draper Rescues Abeta Toxicity in a Drosophila

Model of Alzheimer's Disease. The Journal of neuroscience : the official journal of the Society

for Neuroscience *37*, 11881-11893.

Reiner, A., and Deng, Y.P. (2018). Disrupted striatal neuron inputs and outputs in Huntington's disease.

CNS neuroscience & therapeutics *24*, 250-280.

Ren, P.H., Lauckner, J.E., Kachirskaia, I., Heuser, J.E., Melki, R., and Kopito, R.R. (2009). Cytoplasmic

penetration and persistent infection of mammalian cells by polyglutamine aggregates. Nature

cell biology *11*, 219-225.

Riabinina, O., Luginbuhl, D., Marr, E., Liu, S., Wu, M.N., Luo, L., and Potter, C.J. (2015). Improved and

expanded Q-system reagents for genetic manipulations. Nature methods *12*, 219-222.

Riabinina, O., and Potter, C.J. (2016). The Q-System: A Versatile Expression System for Drosophila.

Methods in molecular biology *1478*, 53-78.

Sathasivam, K., Neueder, A., Gipson, T.A., Landles, C., Benjamin, A.C., Bondulich, M.K., Smith, D.L.,

Faull, R.L., Roos, R.A., Howland, D.*, et al.* (2013). Aberrant splicing of HTT generates the

pathogenic exon 1 protein in Huntington disease. Proceedings of the National Academy of

Sciences of the United States of America *110*, 2366-2370.

Schafer, D.P., Lehrman, E.K., Kautzman, A.G., Koyama, R., Mardinly, A.R., Yamasaki, R., Ransohoff,

R.M., Greenberg, M.E., Barres, B.A., and Stevens, B. (2012). Microglia sculpt postnatal neural

circuits in an activity and complement-dependent manner. Neuron *74*, 691-705.

Sepp, K.J., Schulte, J., and Auld, V.J. (2001). Peripheral glia direct axon guidance across the

CNS/PNS transition zone. Developmental biology *238*, 47-63.

Sharma, M., and Subramaniam, S. (2019). Rhes travels from cell to cell and transports Huntington

disease protein via TNT-like protrusion. The Journal of cell biology *218*, 1972-1993.

Stepensky, D. (2007). FRETcalc plugin for calculation of FRET in non-continuous intracellular

compartments. Biochemical and biophysical research communications *359*, 752-758.

Stocker, R.F., Heimbeck, G., Gendre, N., and de Belle, J.S. (1997). Neuroblast ablation in Drosophila

P[GAL4] lines reveals origins of olfactory interneurons. Journal of neurobiology *32*, 443-456.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirian, L., and Dickson, B.J. (2005). Neural circuitry that

governs Drosophila male courtship behavior. Cell *121*, 795-807.

Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., and O'Kane, C.J. (1995). Targeted expression of

tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and causes

behavioral defects. Neuron *14*, 341-351.

Tabrizi, S.J., Leavitt, B.R., Landwehrmeyer, G.B., Wild, E.J., Saft, C., Barker, R.A., Blair, N.F.,

Craufurd, D., Priller, J., Rickards, H*., et al.* (2019). Targeting Huntingtin Expression in Patients

with Huntington's Disease. The New England journal of medicine *380*, 2307-2316.

Tran, H.T., Chung, C.H., Iba, M., Zhang, B., Trojanowski, J.Q., Luk, K.C., and Lee, V.M. (2014). Alpha-

synuclein immunotherapy blocks uptake and templated propagation of misfolded alpha-

synuclein and neurodegeneration. Cell reports *7*, 2054-2065.

Tremblay, M.E., Zettel, M.L., Ison, J.R., Allen, P.D., and Majewska, A.K. (2012). Effects of aging and

sensory loss on glial cells in mouse visual and auditory cortices. Glia *60*, 541-558.

Trevino, R.S., Lauckner, J.E., Sourigues, Y., Pearce, M.M., Bousset, L., Melki, R., and Kopito, R.R.

(2012). Fibrillar structure and charge determine the interaction of polyglutamine protein

aggregates with the cell surface. The Journal of biological chemistry *287*, 29722-29728.

Vaquer-Alicea, J., and Diamond, M.I. (2019). Propagation of Protein Aggregation in Neurodegenerative

Diseases. Annual review of biochemistry *88*, 785-810.

Victoria, G.S., Arkhipenko, A., Zhu, S., Syan, S., and Zurzolo, C. (2016). Astrocyte-to-neuron

intercellular prion transfer is mediated by cell-cell contact. Scientific reports *6*, 20762.

Wang, N., Gray, M., Lu, X.-H., Cantle, J.P., Holley, S.M., Greiner, E., Gu, X., Shirasaki, D., Cepeda, C.,

Li, Y*., et al.* (2014). Neuronal targets for reducing mutant huntingtin expression to ameliorate

disease in a mouse model of Huntington's disease. Nature medicine *20*, 536-541.

Weinhard, L., di Bartolomei, G., Bolasco, G., Machado, P., Schieber, N.L., Neniskyte, U., Exiga, M.,

Vadisiute, A., Raggioli, A., Schertel, A*., et al.* (2018). Microglia remodel synapses by

presynaptic trogocytosis and spine head filopodia induction. Nature communications *9*, 1228.

Weiss, K.R., Kimura, Y., Lee, W.C., and Littleton, J.T. (2012). Huntingtin aggregation kinetics and their

pathological role in a Drosophila Huntington's disease model. Genetics *190*, 581-600.

Wilton, D.K., Dissing-Olesen, L., and Stevens, B. (2019). Neuron-Glia Signaling in Synapse Elimination. Annual review of neuroscience *42*, 107-127.

Wu, J.W., Herman, M., Liu, L., Simoes, S., Acker, C.M., Figueroa, H., Steinberg, J.I., Margittai, M., Kayed, R., Zurzolo, C., Di Paolo, G., and Duff, K.E. (2013). Small misfolded tau species are internalized via bulk endocytosis and anterogradely and retrogradely transported in neurons. Journal of biological chemistry *288*, 1856-1870.

Wu, J.W., Hussaini, S.A., Bastille, I.M., Rodriguez, G.A., Mrejeru, A., Rilett, K., Sanders, D.W., Cook, C., Fu, H., Boonen, R.A.*, et al.* (2016). Neuronal activity enhances tau propagation and tau pathology in vivo. Nature neuroscience *19*, 1085-1092.

Wu, B., Li, J., Chou, Y-H., Luginbuhl, D., and Luo, L. (2017) Fibroblast growth factor signaling instructs ensheathing glia wrapping of *Drosophila* olfactory glomeruli. Proceedings of the National Academy of Sciences of the United States of America *114*, 7505-7512.

Wyss-Coray, T., Loike, J.D., Brionne, T.C., Lu, E., Anankov, R., Yan, F., Silverstein, S.C., and Husemann, J. (2003). Adult mouse astrocytes degrade amyloid-beta in vitro and in situ. Nature medicine *9*, 453-457.

Yanamandra, K., Kfoury, N., Jiang, H., Mahan, T.E., Ma, S., Maloney, S.E., Wozniak, D.F., Diamond, M.I., and Holtzman, D.M. (2013). Anti-tau antibodies that block tau aggregate seeding in vitro markedly decrease pathology and improve cognition in vivo. Neuron *80*, 402-414.

Yun, S.P., Kam, T.-I., Panicker, N., Kim, S., Oh, Y., Park, J.S., Kwon, S.H., Park, Y.J., Karuppagounder, S.S., Park, H.*, et al.* (2018). Block of A1 astrocyte conversion by microglia is neuroprotective in models of Parkinson's disease. Nature medicine *24*, 931-938.

Zeineddine, R., Pundavela, J.F., Corcoran, L., Stewart, E.M., Do-Ha, D., Bax, M., Guillemin, G., Vine, K.L., Hatters, D.M., Ecroyd, H.*, et al.* (2015). SOD1 protein aggregates stimulate macropinocytosis in neurons to facilitate their propagation. Molecular neurodegeneration *10*, 57.

Zhao, Y., Wu, X., Li, X., Jiang, L.L., Gui, X., Liu, Y., Sun, Y., Zhu, B., Pina-Crespo, J.C., Zhang, M.*, et al.* (2018). TREM2 Is a Receptor for beta-Amyloid that Mediates Microglial Function. Neuron *97*, 1023-1031.

FIGURE LEGENDS

Figure 1. mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates formed in presynaptic ORNs induce the aggregation of

wtHtt$_{ex1}$ expressed in postsynaptic PNs. (A) Primary structure of full-length human Htt (3144 amino

acids), including HEAT repeats (HR, *gray regions*) and the N-terminal variable-length polyQ region

(*green/red box*), with the pathogenic threshold (~Q37) indicated by a white dotted line. C-termini of two

N-terminal mHtt fragments used in this study (Htt$_{ex1}$ and Htt$_{ex1-12}$) are indicated. (B) Overall experimental

approach. In the fly olfactory system, ORNs synapse with PNs in discrete regions of the antennal lobe

known as glomeruli (*gray circles*). PNs send axons into higher brain centers (i.e., mushroom body

and/or lateral horn). Draper-expressing glial cells project processes in the antennal lobe, where they

ensheath individual glomeruli. To monitor spreading of mHtt aggregates between synaptically-

connected ORNs and PNs, we generated transgenic flies that express mHtt$_{ex1}$ or mHtt$_{ex1-12}$ fragments in

DA1 ORNs and wtHtt$_{ex1}$ in DA1 PNs. Inset: Transfer of mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates between ORNs

and PNs was assessed by monitoring the solubility and colocalization of mHtt and wtHtt fluorescent

signals. (C and D) Maximum intensity z-projections of antennal lobes from 7 day-old adult males

expressing either Htt$_{ex1}$Q25-mCherry (C) or Htt$_{ex1}$Q91-mCherry (D) in DA1 ORNs using *Or67d-QF* and

Htt$_{ex1}$Q25-GFP in GH146+ PNs using *GH146-Gal4*. Raw data are shown in grayscale for individual

channels and pseudocolored in merged images. Merged images include Bruchpilot

immunofluorescence in blue to mark neuropil, which was used to approximate the boundaries of the

DA1 glomerulus (white dotted lines). Scale bars = 20 μm. (E-G) High-magnification confocal z-stacks of

DA1 glomeruli from 1 day-old (E), 14 day-old (F), and 21 day-old (G) adult males expressing Htt$_{ex1}$Q91-

mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Boxed regions in (F and G) are shown at

higher magnification in insets. Raw data are shown in grayscale in individual channels (Htt$_{ex1}$Q91: E1,

F1, G1; Htt$_{ex1}$Q25: E2, F2, G2) and pseudocolored in merged images (E3, F3, G3). mCherry+

"Htt$_{ex1}$Q91 surfaces" (E1', F1', G1') and "Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces" (E2', F2', G2') identified by

semi-automated image segmentation are shown adjacent to raw data and pseudocolored red and

yellow, respectively, in the "merged surfaces" images (E3', F3', G3'). Arrows (*yellow* on grayscale

images, *white* on merged images) indicate Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces. Scale bars = 10 μm. (H and I)

Confocal z-stacks from 1 day-old (H) and 21 day-old (I) adult females expressing RFP-Htt$_{ex1-12}$Q138 in

DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Boxed region in (I) is shown at higher magnification in

insets. RFP+ surfaces identified by semi-automated image segmentation are shown in the last column,

with Htt$_{ex1-12}$Q138-only surfaces in red and Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10

μm. (J) and (K) Numbers of Htt$_{ex1}$Q91 or Htt$_{ex1-12}$Q138 ("mHtt") surfaces (J) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 or

Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 ("mHtt+wtHtt") surfaces (K) identified in adult males (*open bars*) or females

(*solid bars*) expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs or adult females expressing RFP-Htt$_{ex1-12}$Q138

in DA1 ORNs (*striped bars*) at the indicated ages. Data are shown as mean ± SEM; *p < 0.05, **p <

0.01, ***p < 0.001, or ****p < 0.0001 by two-way ANOVA followed by Tukey's multiple comparisons

tests. *s indicate statistical significance comparing flies of the same genotype and sex at different ages

(black *s compare males or females expressing Htt$_{ex1}$Q91, and gray *s compare females expressing

Htt$_{ex1-12}$Q138 over time). #s indicate statistical significance comparing different genotypes at the same

age (black #s compare males vs females expressing Htt$_{ex1}$Q91, and gray #s compare females

expressing Htt$_{ex1}$Q91 vs. females expressing Htt$_{ex1-12}$Q138).

1030

Figure 1 – figure supplement 1. mHtt$_{ex1}$ nucleates prion-like conversion of wtHtt$_{ex1}$ in fly

neurons. (A-E) Confocal z-stacks of 14 day-old adult female brains expressing (A) Htt$_{ex1}$Q91-mCherry

or (B) Htt$_{ex1}$Q25-GFP alone, or co-expressing Htt$_{ex1}$Q91-mCherry with (C) Htt$_{ex1}$Q25-GFP, (D)

membrane-targeted mCD8-GFP, or (E) soluble GFP in all neurons using *elav[C155]-Gal4*. Dimensions

of each confocal stack are 250 x 250 x ~60 (*xyz*) μm. Merged images include DAPI signal (*blue*) to

label nuclei. Scale bars = 20 μm. (F) Colocalization of mCherry and GFP fluorescent signals calculated

as Pearson's correlation coefficients for genotypes shown in (A-E). Data are shown as mean ± SEM;

****p < 0.0001 by one-way ANOVA with Tukey's multiple comparisons test.

1039

Figure 1 – figure supplement 2. Semi-automatic quantification of seeded wtHtt$_{ex1}$ aggregates. (A

and B) Semi-automatic image processing workflow for high-magnification confocal z-stacks of DA1

glomeruli from 14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either (A1,2)

Htt$_{ex1}$Q25-GFP or (B1,2) mCD8-GFP in GH146+ PNs. Raw data were preprocessed by deconvolution

to reduce noise (*top panels*), segmented in the mCherry (A1 and B1) or GFP (A2 and B2) channels

(*middle panels*), and filtered for co-localizing fluorescence signal in the other channel (*bottom panels*).

Arrows in (A1 and A2) indicate seven Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta identified by this method. Scale bars

= 10 μm. (C1-7) Selected single 0.35 μm confocal z-slices from the same confocal stack shown in (A1

and A2). Slice number is indicated at the top right of each image. Individual Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta

identified by semi-automated image segmentation in (A) are indicated with arrows (*yellow* in individual

channels, *white* on merged images) in each slice. Two additional co-localized Htt$_{ex1}$Q91+Htt$_{ex1}$Q25

puncta identified by manual counting are indicated with asterisks in (A2). Scale bars = 10 μm. Insets

show Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta at higher zoom (inset dimensions = 9.12 μm x 9.12 μm). Htt$_{ex1}$Q91-

mCherry (*red*) and Htt$_{ex1}$Q25-GFP (*green*) fluorescence intensity profiles for lines indicated in merged

insets are shown below images. Lines were scanned from leftmost to rightmost point. (D) Comparison

of manual quantification (C) vs semi-automated segmentation approaches (A1 and A2) for 14 day-old

males with the same genotype in (A and C). Data are shown as mean ± SEM; n.s. = not significant by

one-way ANOVA followed by Tukey's multiple comparisons test.

1058

Figure 1 – figure supplement 3. Controls for prion-like transmission of mHtt$_{ex1}$ aggregates from

presynaptic DA1 ORNs to postsynaptic PNs. (A-D) Confocal z-stacks of DA1 glomeruli from 10 day-

old adults expressing (A) Htt$_{ex1}$Q25-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, (B)

Htt$_{ex1}$Q91-mCherry in DA1 ORNs and mCD8-GFP in GH146+ PNs, (C) Htt$_{ex1}$Q91-mCherry together

with Gal80 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, and (D) Htt$_{ex1}$Q91-mCherry in DA1

ORNs and Htt$_{ex1}$Q25-GFP together with QS in GH146+ PNs. mCherry+ surfaces identified by semi-

automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and

Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (E and F) Quantification of (E) Httex1Q91

and (F) Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates in the DA1 glomeruli of flies with genotypes shown in (A-D).

Data are shown as mean ± SEM. *p < 0.05, ****p < 0.0001 by one-way ANOVA with Tukey's multiple

comparisons test.

1070

Figure 1 – figure supplement 4. mHtt$_{ex1}$ aggregates do not transfer retrogradely from PN dendrites to ORN axons. Single confocal slices (A1-3 and B1-3) and confocal z-stacks (A4 and B4) of the antennal lobe from 7 (A) and 14 (B) day-old adult females expressing Htt$_{ex1}$Q91-mCherry in ~60% of PNs using *GH146-QF* and Htt$_{ex1}$Q25-GFP in all ORNs using *pebbled-Gal4*. Dissected brains were immunostained with antibodies against mCherry (*red*), GFP (*green*), and the neuropil marker Bruchpilot (*blue*, shown in merged images). GFP+ puncta identified in single slices are indicated by open arrowheads; none of these were found to be mCherry+. Semi-automated segmentation of the Htt$_{ex1}$Q91-mCherry fluorescent signal ("merge surfaces" in A4 and B4) identified numerous Htt$_{ex1}$Q91 aggregates (graphed in C) throughout the antennal lobe neuropil and surrounding region. A small number of Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces (graphed in D) were identified in these brains (arrows in A4 and B4); however, none of these were located within the boundaries of the antennal lobe. Scale bars = 10 μm; slice numbers are indicated at the top right in (A1-3 and B1-3). Quantified data in (C and D) are shown as mean ± SEM; n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test.

Figure 2. wtHtt$_{ex1}$ aggregates in postsynaptic PNs are nucleated by mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates from presynaptic ORNs. (A-D) High-magnification confocal z-stacks of DA1 glomeruli from adult flies expressing Htt$_{ex1}$Q91-mCherry (A and B) or RFP-Htt$_{ex1-12}$Q138 (C and D) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Raw data (A1-3, B1-3, C1-3, D1-3) are shown adjacent to surfaces identified by 3D segmentation of the red (A1', B1', C1', D1') or green (A2', B2', C2', D2') channels. Htt$_{ex1}$Q25-GFP surfaces are shown at 50% transparency in "merged surfaces" images (A3', B3', C3', D3') for visibility of co-localized Htt$_{ex1}$Q91-mCherry or RFP-Htt$_{ex1-12}$Q138 surfaces. Scale bars = 1 μm. (E1-4) A single confocal slice through the center of a Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregate before (E1, E2) and after (E1', E2') mCherry acceptor photobleaching. Data are shown as a heat map to highlight changes in fluorescence intensities after photobleaching. Scale bar = 1 μm. FRET efficiency (FRET$_{eff}$) for this aggregate is shown in (E4), and average FRET$_{eff}$ values for all Htt$_{ex1}$Q91+Htt$_{ex1}$Q25

aggregates tested are shown in (E3). (F) Volumes of $Htt_{ex1}Q91$ (*solid red boxes*), $Htt_{ex1}Q91+Htt_{ex1}Q25$

(*solid green boxes*), $Htt_{ex1-12}Q138$ (*striped red boxes*), and $Htt_{ex1-12}Q138+Htt_{ex1}Q25$ (*striped green*

boxes) aggregates identified in the DA1 glomerulus at the indicated ages. Box widths indicate

interquartile ranges, vertical lines inside each box indicate medians, whiskers indicate

minimums/maximums, and "+"s indicate means for each data set. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

****$p < 0.0001$ by one-way ANOVA followed by Tukey's multiple comparisons test. Statistical

significance is indicated by "*"s when comparing the same aggregate sub-population at different ages

($Htt_{ex1}Q91$ surfaces in *black* and $Htt_{ex1-12}Q138$ surfaces in *gray*), by "#"s when comparing $Htt_{ex1}Q91$ vs

$Htt_{ex1}Q91+Htt_{ex1}Q25$ (*black*) or $Htt_{ex1-12}Q138$ vs $Htt_{ex1-12}Q138+Htt_{ex1}Q25$ (*gray*) aggregates at the same

ages, and by "^"s when comparing $Htt_{ex1}Q91$ vs $Htt_{ex1-12}Q138$ aggregates at the same ages. (G and H)

Distribution of volumes for (G) $Htt_{ex1}Q91$ (*light gray bars*) and $Htt_{ex1}Q91+Htt_{ex1}Q25$ (*dark gray bars*) or

(H) $Htt_{ex1-12}Q138$ (*light gray bars*) and $Htt_{ex1-12}Q138+Htt_{ex1}Q25$ (*dark gray bars*) aggregates, combined

from 7, 14, and 21 day-old flies. Mean volume of $Htt_{ex1}Q91$ or $Htt_{ex1-12}Q138$ aggregates and

$Htt_{ex1}Q91+Htt_{ex1}Q25$ or $Htt_{ex1-12}Q138+Htt_{ex1}Q25$ aggregates are indicated by black and white dotted

lines, respectively, on each histogram.

1112

Figure 3. $mHtt_{ex1}$ aggregate transfer from ORNs to synaptically-connected PNs is inversely

correlated with presynaptic activity. (A, B, E, F, I, and J) Confocal z-stacks of DA1 glomeruli from 10

day-old males (A and B, and E and F) or 7 day-old females (I and J) co-expressing $Htt_{ex1}Q91$-mCherry

with either LacZ (A, E, and I), shi^{ts1} (B), TeTxLC (F), or dTrpA (J) in DA1 ORNs and $Htt_{ex1}Q25$-GFP in

GH146+ PNs. In (A-B), flies were raised at the permissive temperature (18°C) and shifted to the

restrictive temperature (31°C) upon eclosion, and in (I-J), flies were raised at room temperature (~21°C)

and shifted to 31°C upon eclosion. mCherry+ surfaces identified by semi-automated image

segmentation are shown in the last panels, with $Htt_{ex1}Q91$-only surfaces in red and $Htt_{ex1}Q91+Htt_{ex1}Q25$

surfaces in yellow. Scale bars = 10 μm. (C and D, G and H, and K and L) Quantification of $Htt_{ex1}Q91$ (C,

G, and K) and $Htt_{ex1}Q91+Htt_{ex1}Q25$ (D, H, and L) aggregates identified in DA1 glomeruli from adult

males of the indicated ages co-expressing $Htt_{ex1}Q91$-mCherry with LacZ or shi^{ts1} using two independent

QUAS-shi^{ts1} lines in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (C and D), adult males of the

indicated ages co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or TeTxLC using two independent *QUAS-*

TeTxLC lines in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (G and H), and 7 day-old females

expressing Htt$_{ex1}$Q91-mCherry with either LacZ or dTrpA using three independent *QUAS-dTrpA* lines in

DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (K and L). Data are shown as mean ± SEM; *p < 0.05,

**p < 0.01, ***p < 0.001, ****p < 0.0001, n.s. = not significant by one- or two-way ANOVA with Tukey's

multiple comparisons test comparing shi^{ts1}-, TeTxLC-, or dTrpA-expressing flies to their respective

controls expressing LacZ.

1132

Figure 4. Inhibiting Shibire-mediated endocytosis increases mHttex1-expressing ORN axon

volume and enhances transfer of mHtt$_{ex1}$ aggregates from DA1 ORN axons to glia. (A and B)

Confocal z-stacks of DA1 glomeruli from 10 day-old females co-expressing Htt$_{ex1}$Q91-mCherry, mCD8-

GFP, and either LacZ (A) or shi^{ts1} (B) in DA1 ORNs. Flies were shifted from the permissive temperature

(18°C) to the restrictive temperature (31°C) upon eclosion. Raw data are shown in grayscale, and 3D

segmented surfaces are shown in red for Htt$_{ex1}$Q91 and as a heat map for mCD8-GFP to highlight

differences in intensity between the genotypes. Scale bars = 10 μm. (C) Quantification of mCD8-GFP

intensity (left y-axis, *green*) and volume (right y-axis, *black*) of DA1 glomeruli from 10 day-old adult

females co-expressing LacZ or shi^{ts1} with Htt$_{ex1}$Q91-mCherry and mCD8-GFP in DA1 ORNs. a.u. =

arbitrary units. Data are shown as mean ± SEM; ****p < 0.0001 by Student's t-test. (D and E) Confocal

z-stacks of DA1 glomeruli from 5-6 day-old males expressing Htt$_{ex1}$Q91-mCherry with either LacZ (D) or

shi^{ts1} (E) in DA1 ORNs and Htt$_{ex1}$Q25-YFP in repo+ glia. Adult flies were shifted from 18°C to 31°C

upon eclosion. mCherry+ surfaces identified by semi-automated image segmentation are shown in the

last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars

= 10 μm. (F and G) Quantification of Htt$_{ex1}$Q91 (F) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (G) aggregates in DA1

glomeruli of 5-6 day-old males expressing LacZ or shi^{ts1} using two independent *QUAS-shi^{ts1}* lines. Data

are shown as mean ± SEM; *p < 0.05, ****p < 0.0001 by one-way ANOVA with Tukey's multiple

comparisons test comparing shi^{ts1}-expressing flies to control flies expressing LacZ.

1151

Figure 5. Draper mediates mHtt$_{ex1}$ aggregate transfer from presynaptic DA1 ORNs to

postsynaptic PNs and regulates neuronal mHtt$_{ex1}$ aggregate size. (A and B) Confocal z-stacks of

DA1 glomeruli from 13 day-old adult females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and

Htt$_{ex1}$Q25-GFP in GH146+ PNs, either heterozygous (A; *drpr*$^{+/-}$) or homozygous (B; *drpr*$^{-/-}$) for the *drpr*$^{\Delta 5}$

null allele. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last

row, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10

µm. (C and D) Quantification of Htt$_{ex1}$Q91 (C) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (D) aggregates in DA1 glomeruli

from female or male *drpr*$^{+/-}$ or *drpr*$^{-/-}$ flies at the indicated ages. Data are shown as mean ± SEM; *p <

0.05, **p < 0.01, ****p < 0.0001 by two-way ANOVA with Sidak's multiple comparisons test for *drpr*$^{+/-}$

vs. *drpr*$^{-/-}$ flies at the same ages. "n.d." = not determined; 13 day-old *drpr*$^{-/-}$ flies were not viable. (E and

F) Htt$_{ex1}$Q91 surfaces identified in DA1 glomeruli from 7 day-old *drpr*$^{+/-}$ or *drpr*$^{-/-}$ females expressing

Htt$_{ex1}$Q91-mCherry in DA1 ORNs. mCherry+ surfaces are color-coded according to the following

volume ranges: cyan = 0-1.0 µm^3; gray = 1.01-2.49 µm^3; magenta = 2.5-9.0 µm^3. Gray and magenta

surfaces were set to 70% transparency to improve visibility of smaller cyan surfaces. Scale bars = 10

µm. (G) Relative frequency of volumes for all Htt$_{ex1}$Q91 aggregates identified in 7 day-old *drpr*$^{+/+}$ (*solid*

line) or *drpr*$^{-/-}$ (*dotted line; gray shading*) males and females. The inset graph shows mean Htt$_{ex1}$Q91

aggregate volume ± SEM for the two genotypes. ****p < 0.0001 by unpaired Student's t-test.

1169

Figure 5 – figure supplement 1. Draper is required for enhanced transfer of mHtt$_{ex1}$ from shi^{ts1}-

expressing DA1 ORNs to GH146+ PNs. (A and B) Confocal stacks of DA1 glomeruli from 7 day-old

drpr$^{+/+}$ (A) or *drpr*$^{-/-}$ (B) adult females co-expressing Htt$_{ex1}$Q91-mCherry with shi^{ts1} in DA1 ORNs and

Htt$_{ex1}$Q25-GFP in GH146+ PNs. Adult flies were shifted from 18°C to 31°C upon eclosion. mCherry+

surfaces identified by semi-automated image segmentation are shown in the last panels, with

Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 µm. (C and

D) Quantification of Htt$_{ex1}$Q91 (C) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (D) aggregates for the same genotypes

shown in (A and B). Data are shown as mean ± SEM; ****p < 0.0001 by Student's t-test comparing *drpr*

$^{+/+}$ vs *drpr* $^{-/-}$ flies.

1179

Figure 5 – figure supplement 2. Gal80-mediated repression of Gal4 in glia or RNAi knockdown of

drpr in PNs do not alter ORN-to-PN prion-like transfer of mHtt$_{ex1}$ aggregates. (A and B) Confocal

stacks of DA1 glomeruli from 10-14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs

and Htt$_{ex1}$Q25-GFP in GH146+ PNs with (A) no additional transgenes ("control") or (B) repo-Gal80 to

inhibit Gal4-mediated expression of Htt$_{ex1}$Q25-GFP in glia. (C and D) Confocal z-stacks of DA1

glomeruli from 14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and co-expressing

Htt$_{ex1}$Q25-GFP together with firefly luciferase (FFLuc; C) or dsRNA targeting draper (Drpr^{RNAi}; D) in

GH146+ PNs. In (A-D), mCherry+ surfaces identified by semi-automated image segmentation are

shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in

yellow. Scale bars = 10 μm. (E and F) Quantification of Htt$_{ex1}$Q91 (E) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (F)

aggregates for genotypes shown in (A-D). Data are shown as mean ± SEM; n.s. = not significant by

one-way ANOVA with Tukey's multiple comparisons tests.

1192

Figure 5 – figure supplement 3. mHtt$_{ex1}$ aggregates generated in ORNs do not co-localize with

markers of lysosomes or autophagosomes in glia. (A1-2 and B1-2) Single confocal slices from DA1

glomeruli of 4-5 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either the

autophagosomal marker Atg8a-GFP (A1 and A2) or the lysosomal marker GFP-Lamp1 (B1 and B2) in

repo+ glia. Scale bars = 10 μm. Htt$_{ex1}$Q91-mCherry (red) and Atg8a- or Lamp1-GFP (green)

fluorescence intensity profiles for lines indicated in (A1-2 and B1-2) are shown below images; a.u. =

arbitrary units. Lines were scanned from leftmost to rightmost point. (C and D) Confocal stacks of DA1

glomeruli from 4-5 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either Atg8a-

GFP (C) or GFP-Lamp1 (D) in repo+ glia. mCherry+ surfaces identified by semi-automated image

segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+GFP

surfaces in yellow. Scale bars = 10 μm. (E and F) Quantification of Htt$_{ex1}$Q91 (E) and Htt$_{ex1}$Q91+GFP

(F) surfaces for the same genotypes as in (C and D) and for control animals expressing mCD8-GFP in

glia. Data are shown as mean ± SEM; n.s. = not significant by one-way ANOVA with Tukey's multiple

comparisons test.

1207

Figure 6. Caspase activation in ORNs mediates mHtt$_{ex1}$ aggregate transfer from ORNs to PNs.

(A-D) Maximum-intensity projections of antennal lobes from 8 day-old adult males expressing

Htt$_{ex1}$Q25-GFP (A and C) or Htt$_{ex1}$Q91-GFP (B and D) in most ORNs using *Or83b-Gal4* in *drpr^{Δ5}*

heterozygotes (*drpr$^{+/-}$*; A and B) or homozygotes (*drpr$^{-/-}$*; C and D). Brains were immunostained for

GFP (left panels) or cleaved Dcp-1 (right panels). Scale bars = 20 μm. (E) Quantification of cDcp-1

immunofluorescence from 8 day-old adult males with the same genotypes as in (A-D). Data are shown

as mean ± SEM; **p < 0.01, ***p < 0.001, n.s. = not significant by one-way ANOVA with Tukey's

multiple comparisons test. (F-I) Confocal z-stacks of DA1 glomeruli from 14 day-old males expressing

Htt$_{ex1}$Q91-mCherry with LacZ (F) or p35 (G) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, or

Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP with LacZ (H) or p35 (I) in GH146+ PNs.

mCherry+ surfaces identified by 3D segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only

surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (J and K)

Quantification of Htt$_{ex1}$Q91 (J) or Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (K) aggregates in DA1 glomeruli of flies with the

same genotypes in (F and G) (*squares*) or (H and I) (*circles*). Numbers of aggregates in flies expressing

LacZ or p35 are indicated by solid or open shapes, respectively. Data are shown as mean ± SEM; *p <

0.05, n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test.

1224

Figure 7. mHtt$_{ex1}$ aggregates transfer from presynaptic ORNs to postsynaptic PNs via the

cytoplasm of phagocytic glia. (A) Diagram illustrating our experimental approach for examining a role

for Draper-expressing glia in mHtt$_{ex1}$ aggregate transfer from DA1 ORNs to DA1 PNs. Flies that

combined *Or67d-QF-*, *repo-Gal4-*, and *GH146-LexA:GAD*-driven expression of Htt$_{ex1}$Q91-mCherry in

DA1 ORNs, Htt$_{ex1}$Q25-3xHA in all glia, and Htt$_{ex1}$Q25-YFP in ~60% of PNs, respectively, were

generated and analyzed by confocal microscopy of immunostained brains. If Htt$_{ex1}$Q91-mCherry

aggregates travel to PNs via the glial cytoplasm (route 1), triple-labeled (mCherry+/3xHA+/YFP+)

aggregates should be observed. By contrast, if $Htt_{ex1}Q91$-mCherry aggregates transfer directly to PNs

without accessing the glial cytoplasm (route 2), then only double-labeled (mCherry+/3xHA+ and

mCherry+/YFP+) aggregates would be detected. (B-C) Confocal z-stacks of DA1 glomeruli from adult

females of the indicated ages expressing $Htt_{ex1}Q91$-mCherry in DA1 ORNs (red), $Htt_{ex1}Q25$-3xHA in

glia (blue), and $Htt_{ex1}Q25$-YFP in PNs (green) at the indicated ages. Brains were immunostained with

antibodies against the mCherry, 3xHA, and YFP tags unique to each Htt protein. Scale bars = 10 µm.

(D-H) Single 0.35 µm confocal z-slices from females of the indicated ages with (D-G) the same

genotype as in (B and C) or (H) also expressing dsRNAs targeting *drpr* (+Drpr^{RNAi}). Colocalizing

mCherry+/3xHA+ or mCherry+/3xHA+/YFP+ aggregates are indicated by open arrowheads or arrows,

respectively, shown in yellow on grayscale and white on merged images for increased visibility. Scale

bars = 5 µm. $Htt_{ex1}Q91$-mCherry (red), $Htt_{ex1}Q25$-3xHA (blue), and $Htt_{ex1}Q25$-YFP (green) fluorescence

intensity profiles for lines "a" and "b" are shown to the right of each merged image. Lines were scanned

from leftmost to rightmost point. Arrowheads and arrows on graphs indicate peak mCherry fluorescence

in colocalized mCherry+/3xHA+ and mCherry+/3xHA+/YFP+ aggregates, respectively. (I and J)

Quantification of (I) mCherry-only or (J) mCherry+/3xHA+ and mCherry+/3xHA+/YFP+ aggregates

identified in control (*solid bars*) or Drpr^{RNAi}-expressing (*open bars*) animals over time. +Drpr^{RNAi} animals

were only analyzed at 7 days-old. Data are shown as mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001,

or ****p < 0.0001 by one- or two-way ANOVA followed by Tukey's multiple comparisons test. *s indicate

statistical significance comparing control flies at different ages [black *s compare mCherry-only

aggregates in (I) and mCherry+/3xHA+/YFP+ aggregates in (J), and blue *s compare mCherry+/3xHA+

aggregates in (J)], and #s indicate statistical significance comparing mCherry+/3xHA+ and

mCherry+/3xHA+/YFP+ aggregates, respectively, in control vs Drpr^{RNAi}-expressing flies at the same

age.

Figure 7 - figure supplement 1. Controls for monitoring transmission of mHtt$_{ex1}$ aggregates from

presynaptic DA1 ORNs to postsynaptic PNs via a glial intermediate. (A-C) Confocal z-stacks (A1,

B1, and C1) and single 0.35 µm z-slices (A2, B2, and C2) of DA1 glomeruli from 4-6 day-old adult

females expressing (A1-2) Htt_{ex1}Q25-mCherry in DA1 ORNs and Htt_{ex1}Q25-3xHA in repo+ glia, (B1-2)

Htt_{ex1}Q25-mCherry in DA1 ORNs and Htt_{ex1}Q25-YFP in GH146+ PNs, or (C1-2) Htt_{ex1}Q25-3xHA in

repo+ glia and Htt_{ex1}Q25-YFP in GH146+ PNs. All brains were immunostained with antibodies against

all three epitope tags (i.e., mCherry, YFP, and 3xHA) to show specific staining. Scale bars = 10 µm in

(A1, B1, and C1) and 5 µm in (A2, B2, and C2). Htt_{ex1}Q25-mCherry (red), Htt_{ex1}Q25-3xHA (blue), and

Htt_{ex1}Q25-YFP (green) fluorescence intensity profiles for lines "a" and "b" shown in merged slices (A2,

B2, and C2) are shown below images. Lines were scanned from leftmost to rightmost point.

1266

Video 1. Semi-automated quantification of $mHtt_{ex1}$ and seeded $wtHtt_{ex1}$ aggregates in the DA1

glomerulus. Animation illustrating semi-automated approach for quantifying $mHtt_{ex1}$ and $wtHtt_{ex1}$

aggregates in brains expressing Htt_{ex1}Q91-mCherry in DA1 ORNs and Htt_{ex1}Q25-GFP in GH146+ PNs.

Data shown in video is the same as in Fig. 1-figure supplement 2A1-2 and C1-7. Segmentation of raw

high-magnification 3D confocal data (0:00) in the red channel (0:07) identified distinct Htt_{ex1}Q91-

mCherry surfaces (0:09), which were filtered for those that co-localize with high-intensity GFP signal to

isolate the subpopulation associated with Htt_{ex1}Q25-GFP puncta (0:13). Volumetric surfaces

representing Htt_{ex1}Q91 (*red*) and Htt_{ex1}Q91+Htt_{ex1}Q25 (*yellow*) aggregates (0:18) are shown for each

data set analyzed by this method. The animation also illustrates segmentation of raw data in the green

channel (0:28) to identify the brightest Htt_{ex1}Q25-GFP objects in each data set (0:29). Overlap of

mCherry+ and GFP+ surfaces, with GFP+ surfaces set at 50% transparency (0:30), highlights co-

localization of smaller Htt_{ex1}Q91 "seeds" surrounded by Htt_{ex1}Q25 signal in these aggregates.

Figure 1. mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates formed in presynaptic ORNs induce the aggregation of wtHtt$_{ex1}$ expressed in postsynaptic PNs. (A) Primary structure of full-length human Htt (3144 amino acids), including HEAT repeats (HR, *gray regions*) and the N-terminal variable-length polyQ region (*green/red box*), with the pathogenic threshold (~Q37) indicated by a white dotted line. The C-termini of two N-terminal mHtt fragments used in this study (ex1 and ex1-12) are indicated. (B) Overall experimental approach. In the fly olfactory system, ORNs synapse with PNs in discrete regions of the antennal lobe known as glomeruli (*gray circles*). PNs send axons into higher brain centers (i.e., mushroom body and/or lateral horn). Draper-expressing glial cells project processes in the antennal lobe, where they ensheath individual glomeruli. To monitor spreading of mHtt aggregates between synaptically-connected ORNs and PNs, we generated transgenic flies that express mHtt$_{ex1}$ or mHtt$_{ex1-12}$ fragments in DA1 ORNs and wtHtt$_{ex1}$ in DA1 PNs. Inset: Transfer of mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates between ORNs and PNs was assessed by monitoring the solubility and co-localization of mHtt and wtHtt fluorescent signals. (C and D) Maximum intensity z-projections of antennal lobes from 7 day-old adult males expressing either Htt$_{ex1}$Q25-mCherry (C) or Htt$_{ex1}$Q91-mCherry (D) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Raw data are shown in grayscale for individual channels and pseudocolored in merged images. Merged images include Bruchpilot immunofluorescence in blue to mark neuropil, which was used to approximate the boundaries of the DA1 glomerulus (*white dotted lines*). Scale bars = 20 μm. (E-G) High-magnification confocal z-stacks of DA1 glomeruli from 1 day-old (E), 14 day-old (F), and 21 day-old (G) adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Boxed regions in (F and G) are shown at higher magnification in insets. Raw data are shown in grayscale in individual channels (Htt$_{ex1}$Q91: E1, F1, G1; Htt$_{ex1}$Q25: E2, F2, G2) and pseudocolored in merged images (E3, F3, G3). mCherry+ "Htt$_{ex1}$Q91 surfaces" (E1', F1', G1') and "Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces" (E2', F2', G2') identified by semi-automated image segmentation are shown adjacent to raw data and pseudocolored red and yellow, respectively, in the "merged surfaces" images (E3', F3', G3'). Arrows (yellow on grayscale images, white on merged images) indicate Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces. Scale bars = 10 μm. (H and I) Confocal z-stacks from 1 day-old (H) and 21 day-old (I) adult females expressing RFP-Htt$_{ex1-12}$Q138 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Boxed region in (I) is shown at higher magnification in insets. RFP+ surfaces identified by semi-automated image segmentation are shown in the last column, with Htt$_{ex1-12}$Q138-only surfaces in red and Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (J) and (K) Numbers of Htt$_{ex1}$Q91 or Htt$_{ex1-12}$Q138 ("mHtt") surfaces (J) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 or Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 ("mHtt+wtHtt") surfaces (K) identified in adult males (*open bars*) or females (*solid bars*) expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs or adult females expressing RFP-Htt$_{ex1-12}$Q138 in DA1 ORNs (*striped bars*) at the indicated ages. Data are shown as mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001, or ****p < 0.0001 by two-way ANOVA followed by Tukey's multiple comparisons tests. *s indicate statistical significance comparing flies of the same genotype and sex at different ages (black *s compare males or females expressing Htt$_{ex1}$Q91, and gray *s compare females expressing Htt$_{ex1-12}$Q138 over time). #s indicate statistical significance comparing different genotypes at the same age (black #s compare males vs females expressing Htt$_{ex1}$Q91, and gray #s compare females expressing Htt$_{ex1}$Q91 vs. females expressing Htt$_{ex1-12}$Q138).

Figure 1 – figure supplement 1. mHtt$_{ex1}$ nucleates prion-like conversion of wtHtt$_{ex1}$ in fly neurons. (A-E) Confocal z-stacks of 14 day-old adult female brains expressing (A) Htt$_{ex1}$Q91-mCherry or (B) Htt$_{ex1}$Q25-GFP alone, or co-expressing Htt$_{ex1}$Q91-mCherry with (C) Htt$_{ex1}$Q25-GFP, (D) membrane-targeted mCD8-GFP, or (E) soluble GFP in all neurons using *elav[C155]-Gal4*. Dimensions of each confocal stack are 250 x 250 x ~60 (*xyz*) μm. Merged images include DAPI signal (*blue*) to label nuclei. Scale bars = 20 μm. (F) Co-localization of mCherry and GFP fluorescent signals calculated as Pearson's correlation coefficients for genotypes shown in (A-E). Data are shown as mean ± SEM; ****p < 0.0001 by one-way ANOVA with Tukey's multiple comparisons test.

Figure 1 – figure supplement 2. Semi-automatic quantification of seeded wtHtt$_{ex1}$ aggregates. (A and B) Semi-automatic image processing workflow for high-magnification confocal z-stacks of DA1 glomeruli from 14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either (A1,2) Htt$_{ex1}$Q25-GFP or (B1,2) mCD8-GFP in GH146+ PNs. Raw data were preprocessed by deconvolution to reduce noise (*top panels*), segmented in the mCherry (A1 and B1) or GFP (A2 and B2) channels (*middle panels*), and filtered for co-localizing fluorescence signal in the other channel (*bottom panels*). Arrows in (A1 and A2) indicate seven Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta identified by this method. Scale bars = 10 μm. (C1-7) Selected single 0.35 μm confocal z-slices from the same confocal stack shown in (A1 and A2). Slice number is indicated at the top right of each image. Individual Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta identified by semi-automated image segmentation in (A) are indicated with arrows (yellow in individual channels, white on merged images) in each slice. Two additional co-localized Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta identified by manual counting are indicated with asterisks in (A2). Scale bars = 10 μm. Insets show Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 puncta at higher zoom (inset dimensions = 9.12 μm x 9.12 μm). Htt$_{ex1}$Q91-mCherry (*red*) and Htt$_{ex1}$Q25-GFP (*green*) fluorescence intensity profiles for lines indicated in merged insets are shown below images. Lines were scanned from leftmost to rightmost point. (D) Comparison of manual quantification (C) vs semi-automated segmentation approaches (A1 and A2) for 14 day-old males with the same genotype in (A and C). Data are shown as mean ± SEM; n.s. = not significant by one-way ANOVA followed by Tukey's multiple comparisons test.

Figure 1 – figure supplement 3. Controls for prion-like transmission of mHtt$_{ex1}$ aggregates from presynaptic DA1 ORNs to postsynaptic PNs. (A-D) Confocal z-stacks of DA1 glomeruli from 10 day-old adults expressing (A) Htt$_{ex1}$Q25-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, (B) Htt$_{ex1}$Q91-mCherry in DA1 ORNs and mCD8-GFP in GH146+ PNs, (C) Htt$_{ex1}$Q91-mCherry together with Gal80 in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, and (D) Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP together with QS in GH146+ PNs. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (E and F) Quantification of (E) Htt$_{ex1}$Q91 and (F) Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates in the DA1 glomeruli of flies with genotypes shown in (A-D). Data are shown as mean ± SEM. *p < 0.05, ****p < 0.0001 by one-way ANOVA with Tukey's multiple comparisons test.

Figure 1 – figure supplement 4. mHtt$_{ex1}$ aggregates do not transfer retrogradely from PN dendrites to ORN axons. Single confocal slices (A1-3 and B1-3) and confocal z-stacks (A4 and B4) of the antennal lobe from 7 (A) and 14 (B) day-old adult females expressing Htt$_{ex1}$Q91-mCherry in ~60% of PNs using *GH146-QF* and Htt$_{ex1}$Q25-GFP in all ORNs using *pebbled-Gal4*. Dissected brains were immunostained with antibodies against mCherry (*red*), GFP (*green*), and the neuropil marker Bruchpilot (*blue*, shown in merged images). GFP+ puncta identified in single slices are indicated by open arrowheads; none of these were found to be mCherry+. Semi-automated segmentation of the Htt$_{ex1}$Q91-mCherry fluorescent signal ("merge surfaces" in A4 and B4) identified numerous Htt$_{ex1}$Q91 aggregates (graphed in C) throughout the antennal lobe neuropil and surrounding region. A small number of Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces (graphed in D) were identified in these brains (arrows in A4 and B4); however, none of these were located within the boundaries of the antennal lobe. Scale bars = 10 μm; slice numbers are indicated at the top right in (A1-3 and B1-3). Quantified data in (C and D) are shown as mean ± SEM; n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test.

Figure 2. wtHtt$_{ex1}$ aggregates in postsynaptic PNs are nucleated by mHtt$_{ex1}$ or mHtt$_{ex1-12}$ aggregates from presynaptic ORNs. (A-D) High-magnification confocal z-stacks of DA1 glomeruli from adult flies expressing Htt$_{ex1}$Q91-mCherry (A and B) or RFP-Htt$_{ex1-12}$Q138 (C and D) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Raw data (A1-3, B1-3, C1-3, D1-3) are shown adjacent to surfaces identified by 3D segmentation of the red (A1', B1', C1', D1') or green (A2', B2', C2', D2') channels. Htt$_{ex1}$Q25-GFP surfaces are shown at 50% transparency in "merged surfaces" images (A3', B3', C3', D3') for visibility of co-localized Htt$_{ex1}$Q91-mCherry or RFP-Htt$_{ex1-12}$Q138 surfaces. Scale bars = 1 μm. (E1-4) A single confocal slice through the center of a Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregate before (E1, E2) and after (E1', E2') mCherry acceptor photobleaching. Data are shown as a heat map to highlight changes in fluorescence intensities after photobleaching. Scale bar = 1 μm. FRET efficiency (FRET$_{eff}$) for this aggregate is shown in (E4), and average FRET$_{eff}$ values for all Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 aggregates tested are shown in (E3). (F) Volumes of Htt$_{ex1}$Q91 (*solid red boxes*), Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (*solid green boxes*), Htt$_{ex1-12}$Q138 (*striped red boxes*), and Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 (*striped green boxes*) aggregates identified in the DA1 glomerulus at the indicated ages. Box widths indicate interquartile ranges, vertical lines inside each box indicate medians, whiskers indicate minimums/maximums, and "+"s indicate means for each data set. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by one-way ANOVA followed by Tukey's multiple comparisons test. Statistical significance is indicated by "*"s when comparing the same aggregate sub-population at different ages (Htt$_{ex1}$Q91 surfaces in black and Htt$_{ex1-12}$Q138 surfaces in gray), by "#"s when comparing Htt$_{ex1}$Q91 vs Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (black) or Htt$_{ex1-12}$Q138 vs Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 (gray) aggregates at the same ages, and by "^"s when comparing Htt$_{ex1}$Q91 vs Htt$_{ex1-12}$Q138 aggregates at the same ages. (G and H) Distribution of volumes for (G) Htt$_{ex1}$Q91 (*light gray bars*) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (*dark gray bars*) or (H) Htt$_{ex1-12}$Q138 (*light gray bars*) and Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 (*dark gray bars*) aggregates, combined from 7, 14, and 21 day-old flies. Mean volume of Htt$_{ex1}$Q91 or Htt$_{ex1-12}$Q138 aggregates and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 or Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 aggregates are indicated by black and white dotted lines, respectively, on each histogram.

Figure 3. mHtt$_{ex1}$ aggregate transfer from ORNs to synaptically-connected PNs is inversely correlated with presynaptic activity. (A-B, E-F, and I-J) Confocal z-stacks of DA1 glomeruli from 10 day-old males (A-B and E-F) or 7 day-old females (I-J) co-expressing Htt$_{ex1}$Q91-mCherry with either LacZ (A, E, and I), shi^{ts1} (B), TeTxLC (F), or dTrpA (J) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. In (A-B), flies were raised at the permissive temperature (18°C) and shifted to the restrictive temperature (31°C) upon eclosion, and in (I-J), flies were raised at room temperature (~21°C) and shifted to 31°C upon eclosion. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (C-D, G-H, and K-L) Quantification of Htt$_{ex1}$Q91 (C, G, and K) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (D, H, and L) aggregates identified in DA1 glomeruli from adult males of the indicated ages co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or shi^{ts1} using two independent *QUAS-shi^{ts1}* lines in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (C-D), adult males of the indicated ages co-expressing Htt$_{ex1}$Q91-mCherry with LacZ or TeTxLC using two independent *QUAS-TeTxLC* lines in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (G-H), and 7 day-old females expressing Htt$_{ex1}$Q91-mCherry with either LacZ or dTrpA using three independent *QUAS-dTrpA* lines in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs (K-L). Data are shown as mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, n.s. = not significant by one- or two-way ANOVA with Tukey's multiple comparisons test comparing shi^{ts1}-, TeTxLC-, or dTrpA-expressing flies to their respective controls expressing LacZ.

Figure 4. Inhibiting Shibire-mediated endocytosis increases mHtt$_{ex1}$-expressing ORN axon volume and enhances transfer of mHtt$_{ex1}$ aggregates from DA1 ORN axons to glia. (A and B) Confocal z-stacks of DA1 glomeruli from 10 day-old females co-expressing Htt$_{ex1}$Q91-mCherry, mCD8-GFP, and either LacZ (A) or shi^{ts1} (B) in DA1 ORNs. Flies were shifted from the permissive temperature (18°C) to the restrictive temperature (31°C) upon eclosion. Raw data are shown in grayscale, and 3D segmented surfaces are shown in red for Htt$_{ex1}$Q91 and as a heat map for mCD8-GFP to highlight differences in intensity between the genotypes. Scale bars = 10 μm. (C) Quantification of mCD8-GFP intensity (*left y-axis, green*) and volume (*right y-axis, black*) of DA1 glomeruli from 10 day-old adult females co-expressing LacZ or shi^{ts1} with Htt$_{ex1}$Q91-mCherry and mCD8-GFP in DA1 ORNs. a.u. = arbitrary units. Data are shown as mean ± SEM; ****$p < 0.0001$ by Student's *t*-test. (D and E) Confocal z-stacks of DA1 glomeruli from 5-6 day-old males expressing Htt$_{ex1}$Q91-mCherry with either LacZ (D) or shi^{ts1} (E) in DA1 ORNs and Htt$_{ex1}$Q25-YFP in repo+ glia. Adult flies were shifted from 18°C to 31°C upon eclosion. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (F and G) Quantification of Htt$_{ex1}$Q91 (F) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (G) aggregates in DA1 glomeruli of 5-6 day-old males expressing LacZ or shi^{ts1} using two independent *QUAS-shi^{ts1}* lines. Data are shown as mean ± SEM; *$p < 0.05$, ****$p < 0.0001$ by one-way ANOVA with Tukey's multiple comparisons test comparing shi^{ts1}-expressing flies to control flies expressing LacZ.

Figure 5. Draper mediates mHtt$_{ex1}$ aggregate transfer from presynaptic DA1 ORNs to postsynaptic PNs and regulates neuronal mHtt$_{ex1}$ aggregate size. (A and B) Confocal z-stacks of DA1 glomeruli from 13 day-old adult females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, either heterozygous (A; *drpr$^{+/-}$*) or homozygous (B; *drpr$^{-/-}$*) for the *drpr^{Δ5}* null allele. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last row, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (C and D) Quantification of Htt$_{ex1}$Q91 (C) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (D) aggregates in DA1 glomeruli from female or male *drpr$^{+/-}$* or *drpr$^{-/-}$* flies at the indicated ages. Data are shown as mean ± SEM; *p < 0.05, **p < 0.01, ****p < 0.0001 by two-way ANOVA with Sidak's multiple comparisons test for *drpr$^{+/-}$* vs. *drpr$^{-/-}$* flies at the same ages. "n.d." = not determined; 13 day-old *drpr$^{-/-}$* males were not viable. (E and F) Htt$_{ex1}$Q91 surfaces identified in DA1 glomeruli from 7 day-old *drpr$^{+/+}$* or *drpr$^{-/-}$* females expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs. Htt$_{ex1}$Q91 aggregates are color-coded according to the following volume ranges: cyan = 0-1.0 μm³; gray = 1.01-2.49 μm³; magenta = 2.5-9.0 μm³. Gray and magenta surfaces were set to 70% transparency to improve visibility of smaller cyan surfaces. Scale bars = 10 μm. (G) Relative frequency of volumes for all Htt$_{ex1}$Q91 aggregates identified in 7 day-old *drpr$^{+/+}$* (*solid line*) or *drpr$^{-/-}$* (*dotted line; gray shading*) males and females. The inset graph shows mean Htt$_{ex1}$Q91 aggregate volume ± SEM for the two genotypes. ****p < 0.0001 by unpaired Student's *t*-test.

Figure 5 – figure supplement 1. Draper is required for enhanced transfer of mHtt$_{ex1}$ from shi^{ts1}-expressing DA1 ORNs to GH146+ PNs. (A and B) Confocal z-stacks of DA1 glomeruli from 7 day-old *drpr*$^{+/+}$ (A) or *drpr*$^{-/-}$ (B) adult females co-expressing Htt$_{ex1}$Q91-mCherry with shi^{ts1} in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs. Adult flies were shifted from 18°C to 31°C upon eclosion. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (C and D) Quantification of Htt$_{ex1}$Q91 (C) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (D) aggregates for the same genotypes shown in (A and B). Data are shown as mean ± SEM; ****p < 0.0001 by Student's *t*-test comparing *drpr*$^{+/+}$ vs *drpr*$^{-/-}$ flies.

Figure 5 – figure supplement 2. Gal80-mediated repression of Gal4 in glia or RNAi knockdown of *drpr* in PNs do not alter ORN-to-PN prion-like transfer of mHtt$_{ex1}$ aggregates. (A and B) Confocal z-stacks of DA1 glomeruli from 10-14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs with (A) no additional transgenes ("control") or (B) *repo-Gal80* to inhibit *Gal4*-mediated expression of Htt$_{ex1}$Q25-GFP in glia. (C and D) Confocal z-stacks of DA1 glomeruli from 14 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and co-expressing Htt$_{ex1}$Q25-GFP together with firefly luciferase (FFLuc; C) or dsRNA targeting *draper* (Drpr[RNAi]; D) in GH146+ PNs. In (A-D), mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (E and F) Quantification of Htt$_{ex1}$Q91 (E) and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (F) aggregates for genotypes shown in (A-D). Data are shown as mean ± SEM; n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons tests.

Figure 5 – figure supplement 3. mHtt$_{ex1}$ aggregates generated in ORNs do not co-localize with markers of lysosomes or autophagosomes in glia. (A1-2 and B1-2) Single confocal slices from DA1 glomeruli of 4-5 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either the autophagosomal marker Atg8a-GFP (A1 and A2) or the lysosomal marker GFP-Lamp1 (B1 and B2) in repo+ glia. Scale bars = 10 μm. Htt$_{ex1}$Q91-mCherry (*red*) and Atg8a- or Lamp1-GFP (*green*) fluorescence intensity profiles for lines indicated in (A1-2 and B1-2) are shown below images; a.u. = arbitrary units. Lines were scanned from leftmost to rightmost point. (C and D) Confocal z-stacks of DA1 glomeruli from 4-5 day-old adult males expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs and either Atg8a-GFP (C) or GFP-Lamp1 (D) in repo+ glia. mCherry+ surfaces identified by semi-automated image segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+GFP surfaces in yellow. Scale bars = 10 μm. (E and F) Quantification of Htt$_{ex1}$Q91 (E) and Htt$_{ex1}$Q91+GFP (F) surfaces for the same genotypes as in (C and D) and for control animals expressing mCD8-GFP in glia. Data are shown as mean ± SEM; n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test.

Figure 6. Caspase activation in ORNs mediates mHtt$_{ex1}$ aggregate transfer from ORNs to PNs. (A-D) Maximum-intensity projections of antennal lobes from 8 day-old adult males expressing Htt$_{ex1}$Q25-GFP (A and C) or Htt$_{ex1}$Q91-GFP (B and D) in most ORNs using *Or83b-Gal4* in *drpr$^{\Delta5}$* heterozygotes (*drpr$^{+/-}$*; A and B) or homozygotes (*drpr$^{-/-}$*; C and D). Brains were immunostained for GFP (*left panels*) or cleaved Dcp-1 (*right panels*). Scale bars = 20 μm. (E) Quantification of cDcp-1 immunofluorescence from 8 day-old adult males with the same genotypes as in (A-D). Data are shown as mean ± SEM; **p < 0.01, ***p < 0.001, n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test. (F-I) Confocal z-stacks of DA1 glomeruli from 14 day-old males expressing Htt$_{ex1}$Q91-mCherry with LacZ (F) or p35 (G) in DA1 ORNs and Htt$_{ex1}$Q25-GFP in GH146+ PNs, or Htt$_{ex1}$Q91-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-GFP with LacZ (H) or p35 (I) in GH146+ PNs. mCherry+ surfaces identified by 3D segmentation are shown in the last panels, with Htt$_{ex1}$Q91-only surfaces in red and Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 surfaces in yellow. Scale bars = 10 μm. (J and K) Quantification of Htt$_{ex1}$Q91 (J) or Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 (K) aggregates in DA1 glomeruli of flies with the same genotypes in (F and G) (*squares*) or (H and I) (*circles*). Numbers of aggregates in flies expressing LacZ or p35 are indicated by solid or open shapes, respectively. Data are shown as mean ± SEM; *p < 0.05, n.s. = not significant by one-way ANOVA with Tukey's multiple comparisons test.

Figure 7. mHtt$_{ex1}$ aggregates transfer from presynaptic ORNs to postsynaptic PNs via the cytoplasm of phagocytic glia. (A) Diagram illustrating our experimental approach for examining a role for Draper-expressing glia in mHtt$_{ex1}$ aggregate transfer from DA1 ORNs to DA1 PNs. Flies that combined *Or67d-QF-*, *repo-Gal4-*, and *GH146-LexA:GAD*-driven expression of Htt$_{ex1}$Q91-mCherry in DA1 ORNs, Htt$_{ex1}$Q25-3xHA in all glia, and Htt$_{ex1}$Q25-YFP in ~60% of PNs, respectively, were generated and analyzed by confocal microscopy of immunostained brains. If Htt$_{ex1}$Q91-mCherry aggregates travel to PNs via the glial cytoplasm (route 1), triple-labeled (mCherry+/3xHA+/YFP+) aggregates should be observed. By contrast, if Htt$_{ex1}$Q91-mCherry aggregates transfer directly to PNs without accessing the glial cytoplasm (route 2), then only double-labeled (mCherry+/3xHA+ and mCherry+/YFP+) aggregates would be detected. (B-C) Confocal z-stacks of DA1 glomeruli from adult females of the indicated ages expressing Htt$_{ex1}$Q91-mCherry in DA1 ORNs (*red*), Htt$_{ex1}$Q25-3xHA in glia (*blue*), and Htt$_{ex1}$Q25-YFP in PNs (*green*) at the indicated ages. Brains were immunostained with antibodies against the mCherry, 3xHA, and YFP tags unique to each Htt protein. Scale bars = 10 μm. (D-H) Single 0.35 μm confocal z-slices from females of the indicated ages with (D-G) the same genotype as in (B and C) or (H) also expressing dsRNA targeting *drpr* (+Drpr^{RNAi}). Colocalizing mCherry+/3xHA+ or mCherry+/3xHA+/YFP+ aggregates are indicated by open arrowheads or arrows, respectively, shown in yellow on grayscale and white on merged images for increased visibility. Scale bars = 5 μm. Htt$_{ex1}$Q91-mCherry (*red*), Htt$_{ex1}$Q25-3xHA (*blue*), and Htt$_{ex1}$Q25-YFP (*green*) fluorescence intensity profiles for lines "a" and "b" are shown to the right of each merged image. Lines were scanned from leftmost to rightmost point. Arrowheads and arrows on plots indicate peak mCherry fluorescence in colocalized mCherry+/3xHA+ and mCherry+/3xHA+/YFP+ aggregates, respectively. (I and J) Quantification of (I) mCherry-only or (J) mCherry+/3xHA+ and mCherry+/3xHA+/YFP+ aggregates identified in control (*solid bars*) or Drpr^{RNAi}-expressing (*open bars*) animals over time. +Drpr^{RNAi} animals were only analyzed at 7 days-old. Data are shown as mean ± SEM; *p < 0.05, **p < 0.01, ***p < 0.001, or ****p < 0.0001 by one- or two-way ANOVA followed by Tukey's multiple comparisons test. *s indicate statistical significance comparing control flies at different ages [black *s compare mCherry-only aggregates in (I) and mCherry+/3xHA+/YFP+ aggregates in (J), and blue *s compare mCherry+/3xHA+ aggregates in (J)], and blue and black #s indicate statistical significance comparing mCherry+/3xHA+ or mCherry+/3xHA+/YFP+ aggregates, respectively, in control vs Drpr^{RNAi}-expressing flies at the same age.

Figure 7 - figure supplement 1. Controls for monitoring transmission of mHtt$_{ex1}$ aggregates from presynaptic DA1 ORNs to postsynaptic PNs via a glial intermediate. (A-C) Confocal z-stacks (A1, B1, and C1) and single 0.35 µm z-slices (A2, B2, and C2) of DA1 glomeruli from 4-6 day-old adult females expressing (A1-2) Htt$_{ex1}$Q25-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-3xHA in repo+ glia, (B1-2) Htt$_{ex1}$Q25-mCherry in DA1 ORNs and Htt$_{ex1}$Q25-YFP in GH146+ PNs, or (C1-2) Htt$_{ex1}$Q25-3xHA in repo+ glia and Htt$_{ex1}$Q25-YFP in GH146+ PNs. All brains were immunostained with antibodies against all three epitope tags (i.e., mCherry, YFP, and 3xHA) to show specific staining. Scale bars = 10 µm in (A1, B1, and C1) and 5 µm in (A2, B2, and C2). Htt$_{ex1}$Q25-mCherry (*red*), Htt$_{ex1}$Q25-3xHA (*blue*), and Htt$_{ex1}$Q25-YFP (*green*) fluorescence intensity profiles for lines "a" and "b" shown in merged slices (A2, B2, and C2) are shown below images. Lines were scanned from leftmost to rightmost point.

Table. Genotypes and sample sizes displayed in each figure

Figure	Genotype	Sample size (*n*) on graphs
Figure 1		
C	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4/QUAS-Htt$_{ex1}$Q25-mCherry^{attP24}*	
D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4/QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}*	
E,F,G,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	J,K: Htt$_{ex1}$ males - 1d (*14*), 7d (*15*), 14d (*20*), 21d (*14*) J,K: Htt$_{ex1}$ females - 1d (*10*), 7d (*10*), 14d (*17*), 21d (*18*)
H,I,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-RFP-Htt$_{ex1-12}$Q138^{attP3B} ; GH146-Gal4/+*	J,K: Htt$_{ex1-12}$ females - 1d (*10*), 7d (*12*), 14d (*14*), 21d (*20*)
Figure 1, figure supplement 1		
A,F	*elav[C155]-Gal4/UAS-Htt$_{ex1}$Q91-mCherry^{attP3}*	F: Htt$_{ex1}$Q91 - *8*
B,F	*elav[C155]-Gal4/+ ; UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+*	F: Htt$_{ex1}$Q25 - *6*
C,F	*elav[C155]-Gal4/UAS-Htt$_{ex1}$Q91-mCherry^{attP3} ; UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+*	F: Htt$_{ex1}$Q91 + Htt$_{ex1}$Q25 - *6*
D,F	*elav[C155]-Gal4/UAS-Htt$_{ex1}$Q91-mCherry^{attP3} ; UAS-mCD8-GFP/+*	F: Htt$_{ex1}$Q91 + mCD8 - *6*
E,F	*elav[C155]-Gal4/UAS-Htt$_{ex1}$Q91-mCherry^{attP3} ; UAS-GFP^{attP24}/+*	F: Htt$_{ex1}$Q91 + GFP - *7*
Figure 1, figure supplement 2		
A,B,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	D - *14*
Figure 1, figure supplement 3		
A,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4/QUAS-Htt$_{ex1}$Q25-mCherry^{attP24}*	E,F - *6*
B,E,F	*Or67d-QF/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/UAS-mCD8-GFP*	E,F - *8*
C,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-Gal80 ; GH146-Gal4/QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}*	E,F: *6*
D,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4/QUAS-Htt$_{ex1}$Q91-mCherry^{attP24} ; UAS-QS/+*	E,F: *6*
Figure 1, figure supplement 4		
A1-4,B1-4,C,D	*pebbled-Gal4/QUAS-Htt$_{ex1}$Q91-mCherry^{attP3} ; UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+ ; GH146-QF/+*	C,D - 7d (*5*), 14d (*5*)
Figure 2		
A,B,E,F,G	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	E3 - *28* F: Htt$_{ex1}$Q91 - 1d (*1138*), 7d (*2235*), 14d (*4309*), 21d (*5435*) F: Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 - 1d (*8*), 7d (*36*), 14d (*319*), 21d (*392*) G: Htt$_{ex1}$Q91 - *13,132* G: Htt$_{ex1}$Q91+Htt$_{ex1}$Q25 - *754*
C,D,F,H	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-RFP-Htt$_{ex1-12}$Q138^{attP3B} ; GH146-Gal4/+*	F: Htt$_{ex1-12}$Q138 - 1d (*271*), 7d (*469*), 14d (*1196*), 21d (*1944*)

		F: Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 - 1d (*1*), 7d (*16*), 14d (*73*), 21d (*175*) H: Htt$_{ex1-12}$Q138 - *3887* H: Htt$_{ex1-12}$Q138+Htt$_{ex1}$Q25 - *265*
Figure 3		
A,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*	C,D: +LacZ - 4d (*12*), 10d (*8*)
B,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+; QUAS-shibire^{ts1}#5/+*	C,D: +shi^{ts1}#1 - 4d (*10*), 10d (*8*)
C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+; QUAS-shibire^{ts1}#7/+*	C,D: +shi^{ts1}#2 - 4d (*10*), 10d (*11*)
E,G,H	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*	G,H: +LacZ - 4d (*8*), 10d (*10*)
F,G,H	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/QUAS-TeTxLC#4c ; +/(Dh/Tm6b)*	G,H: +TeTxLC#1 - 4d (*10*), 10d (*9*)
G,H	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/Pin; QUAS-TeTxLC#9c/+*	G,H: +TeTxLC#2 - 4d (*10*), 10d (*6*)
I,K,L	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*	K,L: +LacZ - *10*
K,L	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-dTrpA#5 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	K,L: +TrpA#1 - *6*
J,K,L	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-dTrpA#6/+*	K,L: +TrpA#2 - *11*
K,L	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/+ ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/QUAS-dTrpA#7*	K,L: +TrpA#3 - *10*
Figure 4		
A,C	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; QUAS-mCD8-GFP/+ ; QUAS-nucLacZ/+*	C: +LacZ - *10*
B,C	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; QUAS-mCD8-GFP/+ ; QUAS-shibire^{ts1}#7/+*	C: +shi^{ts1} - *8*
D,F,G	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-nucLacZ*	F,G: +LacZ - *9*
E,F,G	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-shibire^{ts1}#7*	F,G: +shi^{ts1}#1 - *8*
F,G	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/Y ; UAS-Htt$_{ex1}$Q25-YFP^{attP24}/+ ; repo-Gal4/QUAS-shibire^{ts1}#5*	F,G: +shi^{ts1}#2 - *10*
Figure 5		
A,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/+*	C,D: *drpr$^{+/-}$*, females - 7d (*12*), 13d (*20*) C,D: *drpr$^{+/-}$* males - 7d (*18*)
B,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*	C,D: *drpr$^{-/-}$* females - 7d (*10*), 13d (*13*) C,D: *drpr$^{-/-}$* males - 7d (*6*)
E,G	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/(+/Y)*	G: *drpr$^{+/+}$* - *1556*
F,G	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/(+/Y) ; ; drpr^{Δ5}/drpr^{Δ5}*	G: *drpr$^{-/-}$* - *1749*
Figure 5, figure supplement 1		
A,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-shibire^{ts1}#2 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	C,D: +shi^{ts1}, *drpr$^{+/+}$* - *11*
B,C,D	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/QUAS-shibire^{ts1}#2 ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*	C,D: +shi^{ts1}, *drpr$^{-/-}$* - *8*

Figure 5, figure supplement 2		
A,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+*	E,F: control - *5*
B,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/repo-Gal80*	E,F: +repo-Gal80 - *6*
C,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; UAS-FFLuc ^{attP2}/+*	E,F: +FFLuc - *11*
D,E,F	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; UAS-Draper$^{RNAi#5}$/+*	E,F: +Drpr^{RNAi} - *6*
Figure 5, figure supplement 3		
A1-2,C,E,F	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; UAS-Atg8a-GFP/+ ; repo-Gal4/+*	E,F: Atg8a - *12*
B1-2,D,E,F	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; UAS-GFP-Lamp1/+ ; repo-Gal4/+*	E,F: Lamp1 - *12*
E,F	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; UAS-mCD8-GFP/+ ; repo-Gal4/+*	E,F: mCD8 - *10*
Figure 6		
A,E	*Or83b-Gal4,UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+ ; drpr^{Δ5}/+*	E: Htt$_{ex1}$Q25, *drpr $^{+/-}$* - *6*
B,E	*Or83b-Gal4,UAS-Htt$_{ex1}$Q91-GFP^{attP24}/+ ; drpr^{Δ5}/+*	E: Htt$_{ex1}$Q91, *drpr $^{+/-}$* - *7*
C,E	*Or83b-Gal4,UAS-Htt$_{ex1}$Q25-GFP^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*	E: Htt$_{ex1}$Q25, *drpr $^{-/-}$* - *6*
D,E	*Or83b-Gal4,UAS-Htt$_{ex1}$Q91-GFP^{attP24}/+ ; drpr^{Δ5}/drpr^{Δ5}*	E: Htt$_{ex1}$Q91, *drpr $^{-/-}$* - *8*
F,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+ ; QUAS-nucLacZ/+*	J,K: +LacZ in ORNs - *8*
G,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+; QUAS-p35/+*	J,K: +p35 in ORNs - *8*
H,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/UAS-LacZ*	J,K: +LacZ in PNs - *7*
I,J,K	*Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/Y ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/UAS-p35*	J,K: +p35 in PNs - *7*
Figure 7		
B,C,D,E,F, G,I,J	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/+*	I,J: control - 0d (*12*), 1d (*10*), 2d (*12*), 3d (*10*), 4d (*11*), 5d (*10*), 7d (*14*), 14d (*11*)
H,I,J	*Or67d-QF,QUAS-Htt$_{ex1}$Q91-mCherry^{attP3}/+ ; GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/UAS-Draper$^{RNAi#5}$*	I,J: +Drpr^{RNAi} - 7d (*8*), 14d (*7*)
Figure 7-figure supplement 1		
A1-2	*GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/+*	
B1-2	*Or67d-QF/+ ; GH146-LexA::GAD,LexAop-Htt$_{ex1}$Q25-YFP^{attP24}/QUAS-Htt$_{ex1}$Q25-mCherry^{attP24}*	
C1-2	*Or67d-QF/+ ; QUAS-Htt$_{ex1}$Q25-mCherry^{attP24}/UAS-Htt$_{ex1}$Q25-3xHA^{attP24} ; repo-Gal4/+*	
Video 1		
	Or67d-QF,UAS-Htt$_{ex1}$Q25-GFP^{attP3}/(+/Y) ; GH146-Gal4,QUAS-Htt$_{ex1}$Q91-mCherry^{attP24}/+	