# Genetic profiling of Vietnamese population from large-scale genomic analysis of non-invasive prenatal testing data

Ngoc Hieu Tran[1,2,*], Thanh Binh Vo[1,3], Van Thong Nguyen[4], Nhat Thang Tran[5], Thu-Huong Nhat Trinh[6], Hong-Anh Thi Pham[1,3], Thi Hong Thuy Dao[1,3], Ngoc Mai Nguyen[1,3], Yen-Linh Thi Van[1,3], Vu Uyen Tran[1,3], Hoang Giang Vu[1,3], Quynh-Tram Nguyen Bui[1,3], Phuong-Anh Ngoc Vo[1,3], Huu Nguyen Nguyen[1,3], Quynh-Tho Thi Nguyen[3], Thanh-Thuy Thi Do[3,7], Phuong Cao Thi Ngoc[3,7], Dinh Kiet Truong[3], Hoai-Nghia Nguyen[8,*], Hoa Giang[1,3,*], Minh-Duy Phan[1,3,9,*]

[1]Gene Solutions, Vietnam

[2]David R. Cheriton School of Computer Science, University of Waterloo, Canada

[3]Medical Genetics Institute, Vietnam

[4]Hung Vuong hospital, Vietnam

[5]University Medical Center, Ho Chi Minh city, Vietnam

[6]Tu Du hospital, Vietnam

[7]Division of Molecular Hematology, Lund Stem Cell Center, Lund University, Lund, Sweden

[8]University of Medicine and Pharmacy at Ho Chi Minh city, Vietnam

[9]School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Australia

*Corresponding authors.

Ngoc Hieu Tran (nh2tran@uwaterloo.ca); ORCID: 0000-0001-5186-8852

Hoa Giang (gianghoa@gmail.com)

Hoai-Nghia Nguyen (nhnghia81@gmail.com)

Minh-Duy Phan (m.phan1@uq.edu.au); ORCID: 0000-0002-3426-1044

**Abstract**

The under-representation of several ethnic groups in existing genetic databases and studies

have undermined our understanding of the genetic variations and associated traits or diseases

in many populations. Cost and technology limitations remain the challenges in performing large-

scale genome sequencing projects in many developing countries, including Vietnam. As one of

the most rapidly adopted genetic tests, non-invasive prenatal testing (NIPT) data offers an

alternative untapped resource for genetic studies. Here we performed a large-scale genomic

analysis of 2,683 pregnant Vietnamese women using their NIPT data and identified a

comprehensive set of 8,054,515 single-nucleotide polymorphisms, among which 8.2% were

new to the Vietnamese population. Our study also revealed 24,487 disease-associated genetic

variants and their allele frequency distribution, especially 5 pathogenic variants for prevalent

genetic disorders in Vietnam. We also observed major discrepancies in the allele frequency

distribution of disease-associated genetic variants between the Vietnamese and other

populations, thus highlighting a need for genome-wide association studies dedicated to the

Vietnamese population. The resulted database of Vietnamese genetic variants, their allele

frequency distribution, and their associated diseases presents a valuable resource for future

genetic studies.

**Keywords:** genetics; genomics; non-invasive prenatal testing; genetic diseases

## Background

Following the successful initiative of the 1000 genomes project [1], several large-scale genome and exome sequencing projects have been conducted, either as international collaboration efforts such as ExACT [2], gnomAD [3], or for a specific country or population [4-8]. Those projects have provided comprehensive profiles of human genetic variation in some populations, paving the way for unprecedented advance in treatment of common genetic diseases. However, the lack of diversity and the under-representation of several populations in genome sequencing projects and genome-wide association studies (GWAS) have increasingly become a critical problem [9,10]. For instance, Gurdasani *et al.* found that the representation of ethnic groups in GWAS was significantly biased, with nearly 78% of the participants having European ancestries, whereas the two major populations, Asian and African, only accounted for 11% and 2.4%, respectively [10]. Vietnam has a population of 96.5 million, the 15th highest in the world and the 9th highest in Asia. Yet there was merely one dataset of 99 Vietnamese individuals that had been studied as part of the 1000 genomes project (population code KHV, the Kinh ethnic group in Ho Chi Minh City, Vietnam). A recent study has sequenced genomes and exomes of another 305 individuals to further expand the Vietnamese genetic database [11]. However, costs and technologies to perform large-scale genome sequencing projects still remain a challenge for most developing countries, including Vietnam.

An alternative approach has been proposed recently to re-use the low-coverage genome sequencing data from non-invasive prenatal testing (NIPT) for large-scale population genetics studies [12,13]. NIPT is a method that sequences cell-free DNA from maternal plasma at an ultra-low depth of 0.1-0.2x to detect fetal aneuploidy [14]. By combining a sufficiently large number of NIPT samples, one could obtain a good representation of the population genetic variation. The benefits of re-using NIPT data for population genetics are manifold. First, the data can be re-used at no extra cost given the approval and consent of the participants. As one of

the most rapidly adopted genetic tests, NIPT has been successfully established and become a

standard screening procedure with thousands to millions of tests performed world-wide,

including many developing countries such as Vietnam [14]. Using NIPT data for population

genetic studies may also reduce privacy concerns since the genetic variants can only be

analyzed by aggregating a large number of samples and the results can only be interpreted at

the population level. A single sample tells little about the genetic information of an individual due

to low sequencing depth. Last but not least, previous studies have suggested that sequencing a

large number of individuals at a low depth might provide more accurate inferences of the

population genetic structure than the traditional approach of sequencing a limited number of

individuals at a higher depth, especially when the budget is limited [15,16].

In this paper, we presented the first study of Vietnamese genetic variations from non-invasive

prenatal testing data, and to the best of our knowledge, the third of such kinds in the world

[12,13]. We analyzed the NIPT data of 2,683 pregnant Vietnamese women to identify genetic

variants and their allele frequency distribution in the Vietnamese population. We also studied

the relationships between the Vietnamese genetic profile and common genetic disorders, and

discovered pathogenic variants related to prevalent diseases in Vietnam. Finally, we highlighted

the differences in the distribution of disease-associated genetic variants between the

Vietnamese and other populations, thus highlighting a need for genome-wide association

studies dedicated to the Vietnamese population. The resulted database of Vietnamese genetic

variants from NIPT data is made available to facilitate future research studies in population

genetics and associated traits or diseases.

## Results

**Data collection**

A total of 2,683 pregnant Vietnamese women who performed non-invasive prenatal testing

during the period from 2018-2019 at the Medical Genetics Institute, Vietnam, were recruited to

the study. The participants have approved and given written consent to the anonymous re-use

of their genomic data for the study. All information of the participants is confidential and not

available to the authors, except the records that their NIPT and pregnancy results are normal.

The study was approved by the institutional ethics committee of the University of Medicine and

Pharmacy, Ho Chi Minh city, Vietnam. The whole genome of each participant was sequenced to

an average of 3.6 million paired-end reads of 2x75 bp, which corresponds to a sequencing

depth of 0.17x per sample.

**Genome coverage and sequencing depth of the NIPT dataset**

Data pre-processing was first performed on each of 2,683 samples and the results were stored

in binary alignment map (BAM) format, one BAM file per sample. The data pre-processing steps

include: quality control of raw data using FastQC [17], trimmomatic [18]; alignment of paired-end

reads to the human reference genome (build GRCh38) using bwa [19], samtools [20],

MarkDuplicates [21]; and summary of alignment results using Qualimap [22], bedtools [23], IGV

[24]. The quality of raw data and alignment results are presented in Supplementary Figures S1

and S2. Raw data showed high sequencing quality, no error or bias was observed. The

mapping quality and insert size distributions followed closely what expected across the

reference genome. The overall sequencing error rate was about 0.3%. More details of the data

pre-processing steps can be found in the Methods section.

The average genome coverage and depth were 14.59% and 0.17x per sample, respectively,

and aggregated to 95.09% and 462x across 2,683 samples (Figure 1a). Although the

sequencing depth per sample was low, there might be more than one read from the same

sample overlapping at a genome position. This problem may affect the estimation of allele

frequency because the estimation is based on the assumption that a sample may contribute

only 0 or 1 allele (read) at any given genome position [12,13]. For instance, we found that the

average percentage of genome positions with depth 2x (i.e. covered by two overlapping reads)

in a sample was 1.75% (Figure 1a). These overlapping reads occurred randomly across the

reference genome and the samples. At any genome position, there were on average 47 out of

2,683 samples that each contributed two reads (Supplementary Figure S3). To address this

problem, we followed a filtering strategy from previous studies [12,13] to keep only one read if

there were overlapping reads in a sample. Thus, for every genome position, each sample could

only contribute up to one read, and when the samples were aggregated, all reads at any

position were obtained from different samples. In addition, we also removed alignments with low

mapping quality scores (MAPQ < 30).

After the filtering step, the sequencing depth was reduced from 0.17x to 0.12x per sample. The

aggregated sequencing depth of 2,683 samples was 364x and the genome coverage was

91.56% (Figure 1a). The distributions of genome coverage and sequencing depth are presented

in Figures 1b-d. The sequencing depth was approximately uniform across the reference

genome, except for low-mappability regions and chromosome Y. The distributions of

sequencing depth and MAPQ score also closely followed the mappability of the human

reference genome obtained from Umap [25]. The average sequencing depth of chromosome Y

was about 2.1% that of the whole genome, consistent with the proportion of fetal DNA in NIPT

samples (8-10%) [14].

**Variant calling and validation**

We aggregated 2,683 samples into one and used Mutect2 from GATK [26,27] for variant calling

and allele frequency estimation. In addition to its main function of somatic calling, Mutect2 can

also be used on data that represents a pool of individuals, such as our NIPT dataset, to call

multiple variants at a genome site [12,13,28]. The called variants were further checked against

strand bias, weak evidence, or contamination using FilterMutectCalls. The allele frequencies

were estimated based on the numbers of reads aligned to the reference and the alternate

alleles. For validation, we compared our NIPT call set to the KHV (Kinh in Ho Chi Minh City,

Vietnam) and EAS (East Asian) populations from the 1000 genomes project [1], as well as the

dbSNP database (version 151, [29]).

We identified a total of 8,054,515 SNPs from the NIPT dataset. The transition to transversion

ratio was 2.0 over the whole genome and 2.8 over protein coding regions, which was similar to

the observed ratios from previous genome or exome sequencing projects. As expected, a

majority of these SNPs, 7,390,020 or 91.8%, had been reported earlier in the KHV call set

(Figure 2a). Since the KHV population only had 99 individuals, we further looked into its

common SNPs that were shared by at least two individuals. We found that the NIPT call set

recovered 90.5% of the KHV common SNPs (Supplementary Figure S4; 6,889,016 / 7,609,526

= 90.5%). This sensitivity is in line with the genome coverage reported earlier in Figure 1a. An

important advantage of NIPT data is the ability of sampling a large number of individuals to

better represent a population and to accurately estimate the allele frequency. We found a strong

Pearson correlation of 98.8% between the allele frequency of the NIPT call set and that of the

KHV call set (Figure 2b). Furthermore, thanks to its larger sample size, the NIPT allele

frequency indeed showed better resolution than the KHV one, as evidenced by vertical trails in

Figure 2b or a zoomed-in view in Supplementary Figure S5.

Our NIPT call set included 664,495 (8.2%) SNPs that had not been reported in the KHV call set.

Among them, 67,153 (0.8%) were found in the EAS call set, another 517,020 (6.4%) were found

in the dbSNP database, and the remaining 80,322 (1.0%) were novel SNPs (Figure 2a).

Majority of those SNPs had allele frequencies less than 10%. The overall allele frequency

distribution of our NIPT call set is presented in Figure 2c.

We used VEP (Variant Effect Predictor [30]) to analyze the effects of 8,054,515 variants in the

NIPT call set (Figure 2d). About 1.5% of the SNPs were located in the coding and UTR regions,

12% in the upstream and downstream regions, and 4% in the TF regulatory regions. More than

80% of the SNPs were located in the intron and intergenic regions. We also noted that the new

SNPs and those in KHV had similar proportions of coding, UTR, upstream and downstream,

and TF regulatory regions (Figure 2d).

**Analysis of pathogenic SNPs and their allele frequencies in Vietnamese population**

We searched the NIPT call set against the ClinVar database (version 20191105, [31]) to explore

the associations between Vietnamese genomic variants and common genetic diseases. We

identified 24,487 SNPs with ClinVar annotations that have been reviewed by at least one

research group (Table 1). Among them, five SNPs were classified as "Pathogenic" or "Likely

pathogenic", 117 SNPs were found to affect a "drug response", and a majority of the remaining

were classified as "Benign" or "Likely benign". We also noted that 391 ClinVar-annotated SNPs

(1.6%), including 1 pathogenic SNP, had not been reported in the KHV call set.

Table 2 and Supplementary Table S1 present the details of five pathogenic SNPs identified in

our NIPT call set. Their associated genetic diseases include: erythropoietic protoporphyria, non-

syndromic genetic deafness, Joubert syndrome, hemochromatosis type 1, and 5-alpha

reductase deficiency. The SNP rs9332964 C>T in *SRD5A2*, which is associated with 5-alpha

reductase deficiency, had not been reported in the KHV call set. 5-alpha reductase deficiency is

an autosomal recessive disorder that affects male sexual development. This SNP is rare in the

world and East Asia populations, but was found to be more common in the Vietnamese

population (allele frequency of 0.05%, 0.67%, and 2.90%, respectively). This SNP had also

been reported in a recent study [11] at a very low allele frequency of 1.36%.

We noticed that the allele frequencies of the five pathogenic SNPs varied considerably between

the Vietnamese, the East Asia, and the world populations (Table 2). For instance, the SNP

rs72474224 C>T in *GJB2* is commonly liked to non-syndromic hearing loss and deafness, which

is also the most prevalent genetic disorder in the Vietnamese population. We found that its

allele frequency in the Vietnamese population was 60% higher than in the East Asia population,

which in turn was an order of magnitude higher than in the world population (13.40%, 8.35%,

and 0.76%, respectively). The allele frequency was consistent with the estimated carrier

frequency of 1 in 5 in the Vietnamese population. Similarly, the allele frequency of rs2272783

A>G in *FECH*, which is associated with erythropoietic protoporphyria, was nearly three times

higher in the Vietnamese and East Asia populations than in the world population (28.10%,

32.57%, and 11.23%, respectively). The prevalence of this pair of SNP and disease in East and

Southeast Asia has been reported previously in [32]. On the other hand, the allele frequency of

rs1799945 C>G in *HFE*, which is associated with hemochromatosis type 1, was about two and

three times lower in the Vietnamese and East Asia populations than in the world population

(5.10%, 3.41%, and 10.82%, respectively). Such discrepancies were also observed for "Benign"

variants, e.g., those related to autosomal recessive non-syndromic hearing loss (Supplementary

Table S2). The variations strongly suggest that population-specific genome-wide association

studies are required to provide a more accurate understanding of the clinical significance of

genetic variants and the true disease prevalence in the Vietnamese population.

## Discussion

In this study, we analyzed the genomes of 2,683 pregnant Vietnamese women from their non-

invasive prenatal testing data. The genomes were originally sequenced at a low depth of

approximately 0.17x per sample for the purpose of fetal aneuploidy testing [14]. Here we

combined the 2,683 samples to a total sequencing depth of 364x and performed variant calling

and analysis for the Vietnamese population. We identified a comprehensive set of 8,054,515

SNPs at a high level of sensitivity and accuracy: 90.5% of Vietnamese common SNPs were

recovered; 99% of identified SNPs were confirmed in existing databases; and a strong

correlation of 98.8% to known allele frequencies. The results were exciting given that the total

sequencing depth of our dataset, 364x, was merely equivalent to sequencing 20 individuals at a

moderate depth of 20x. It also suggests that there is still plenty of room for improvement by

increasing the number of NIPT samples. For instance, Liu *et al.* have demonstrated a large-

scale population genetic analysis based on hundreds of thousands of NIPT samples for the

Chinese population [12].

Another benefit of using NIPT data is cost-effective. In our study, the dataset was re-used at no

cost with written consent from the participants. The whole analysis pipeline was done within a

week on a cloud computing platform for a few hundred dollars. Thus, the overall cost was

negligible compared to that of a typical genome sequencing project. The cost advantage of this

approach may play a major role in large-scale genome sequencing projects, especially in

developing countries where technologies and resources are still limited.

Our study revealed 24,487 disease-associated genetic variants, especially five pathogenic

variants for prevalent genetic disorders in the Vietnamese population. We also found major

discrepancies in the allele frequency distribution of genetic variants between the Vietnamese,

the East Asia, and the world populations. Thus, a comprehensive genetic profile and genome-

wide association studies dedicated to the Vietnamese population are highly desired. Knowing

the distribution of genetic disorders in the population will be useful for public health policy and

planning, preventive medicine, early genetic screening strategies, etc.

Some technical and design limitations in our study could be addressed in future research to

improve the application of NIPT data in population genetics studies. First, currently there is no

variant calling tool that is designed specifically for NIPT data. Here we used Mutect2 and

previous studies also used similar somatic calling tools with the purpose of identifying all

possible variants at a genome site [12,13]. Thus, we took a conservative approach to consider

only SNPs but not indels to ensure a reliable call set. Another limitation was the exclusion of

chromosome Y due to its low coverage as a result of limited amount of fetal DNA in NIPT

samples. This problem could be addressed by increasing the number of samples to obtain

enough sequencing coverage for reliable variant calling. NIPT data is also biased by sex, with

only ~5% of the data coming from male population (assuming a 10% cell-free fetal DNA fraction

with 50% male fetuses). In general, sufficiently large sample size is essential to use NIPT data

for population genetics research. Thus, privacy policy, code of ethics, and standards of practice

need to be established to protect the confidential information and data of the participants.

## Conclusions

We showed that non-invasive prenatal testing data could be reliably used to reconstruct the

genetic profile of the Vietnamese population. Our study identified pathogenic variants for

prevalent genetic diseases in the Vietnamese population and called for a need for population-

specific genome-wide association studies. The resulted database presents a valuable resource

for future studies of genetic variations and associated traits or diseases, not only for the

Vietnamese population but also for other Southeast-Asia and Asia populations. Our results also

demonstrated that non-invasive prenatal testing data provides a valuable and cost-effective

resource for large-scale population genetic studies.

## Methods

**Sample preparation**

Cell-free DNA (cfDNA) in maternal plasma was extracted using MagMAX Cell-Free DNA

Isolation Kit from Thermo Fisher Scientific (Waltham, MA, USA). Library preparation was done

using NEBNext Ultra II DNA Library Prep Kit from New England BioLabs (Ipswich, MA, USA).

The samples were sequenced on the NextSeq 550 platform using paired-end 2x75 bp Reagent

Kit from Illumina (San Diego, CA, USA).

**Bioinformatics analysis pipeline**

Quality check of raw sequencing data was performed using FastQC (version 0.11.8, [17]).

Paired-end reads were trimmed to 75 bp, adapters ("TruSeq3-PE-2.fa") and low-quality bases

were removed using trimmomatic (version 0.39, [18]). We only kept pairs with both reads

surviving the trimming (about 95.7% of the dataset). The reads were then aligned to the human

reference genome, build GRCh38 (hg38), using bwa mem (version 0.7.17-r1188, [19]).

Supplementary hits were marked as secondary for Picard compatibility. Alignment results were

sorted and indexed using samtools (version 1.9, [20]). Potential PCR duplicates were marked

using MarkDuplicates from GATK (version 4.1.1.0, [26]).

Alignments with mapping quality scores less than 30 were discarded. In-house Python scripts

were developed to mark overlapping alignments and to keep only one of them. Qualimap

(version 2.2.1, [22]), bedtools (version 2.25.0 [23]), and IGV (version 2.4.19, [24]) were used to

summarize the alignment results and to calculate genome coverage and sequencing depth.

Mutect2 from GATK (version 4.1.1.0 [26,27]) was used in tumor-only mode for variant calling. All

samples were assigned the same sample name to combine them before variant calling.

FilterMutectCalls was used to exclude variants with weak evidence, strand bias, or

contamination. bcftools (version 1.9, [33]) was used to filter, summarize, and compare VCF

(Variant Call Format) files. VEP (version 98, [30]) was used to predict the effects of variants and

to annotate them against dbSNP (version 151, [29]) and ClinVar (version 20191105, [31])

databases.

The whole analysis pipeline and parameter settings can be found in the attached Python scripts.

Declarations

**Ethics approval and consent to participate**

The study was approved by the institutional ethics committee of the University of Medicine and

Pharmacy, Ho Chi Minh city, Vietnam. The participants who performed NIPT triSure at Medical

Genetics Institute, Vietnam, have approved and given written consent to the anonymous re-use

of their genomic data for this study.

**Consent for publication**

All authors have read and approved the manuscript for publication.

**Availability of data and materials**

292     The database of genetic variants identified from NIPT data and the Python scripts for

293     bioinformatics analysis pipeline is available on GitHub: https://github.com/nh2tran/NIPT_WGS

**Competing interests**

295     NHT, TBV, HATP, HTD, NMN, YLTV, VUT, HGV, QTNB, PANV, HNN, HG and MDP are current

296     employees of Gene Solutions, Vietnam. The other authors declare no competing interests.

**Funding**

298     This study was funded by Gene Solutions, Vietnam. The funder did not have any additional role

299     in the study design, data collection and analysis, decision to publish, or preparation of the

300     manuscript.

**Authors' contributions**

TBV, HATP, HTD, NMN, YLTV, VUT, HGV, QTNB, PANV performed experiments.

VTN, NTTM, THNT, TTTD recruited patients and performed clinical analysis.

HNN, QTTN, PCTN, DKT designed experiments and analyzed data.

HNN supervised the project.

NHT, HG, MDP analyzed the data and wrote the manuscript, designed experiments and

analyzed sequencing data.

**Acknowledgements**

309     The authors thank Dr. Kwok Pui Choi for critical reading of the manuscript.

## References

1. The 1000 Genomes Project Consortium (2015). A global reference for human genetic variation. Nature 526:68-74.

2. Lek M et al (2016). Analysis of protein-coding genetic variation in 60,706 humans. Nature 536:285-91.

3. Karczewski KJ et al (2019). Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes. bioRxiv doi: 10.1101/531210.

4. The UK10K Consortium (2015). The UK10K project identifies rare variants in health and disease. Nature 526:82-90.

5. Gudbjartsson DF et al (2015). Large-scale whole-genome sequencing of the Icelandic population. Nat Genet. 47:435-44.

6. Maretty L et al (2017). Sequencing and de novo assembly of 150 genomes from Denmark as a population reference. Nature 548:87-91.

7. The Genome of the Netherlands Consortium (2014). Whole-genome sequence variation, population structure and demographic history of the Dutch population. Nat Genet. 46:814-25.

8. Wu D et al (2019). Large-Scale Whole-Genome Sequencing of Three Diverse Asian Populations in Singapore. Cell 179:736-749.

9. Editorial (2019). Diversity matters. Nat Rev. Genet. 20:495.

10. Gurdasani D, Barroso I, Zeggini E, Sandhu MS (2019). Genomics of disease risk in globally diverse populations. Nat Rev Genet. 20:520-535.

11. Le VS et al. A Vietnamese human genetic variation database (2019). Hum Mutat. 40:1664-1675.

12. Liu S et al (2018). Genomic Analyses from Non-invasive Prenatal Testing Reveal Genetic

Associations, Patterns of Viral Infections, and Chinese Population History. Cell 175:347-

359.

13. Budis J et al (2019). Non-invasive prenatal testing as a valuable source of population

specific allelic frequencies. J Biotechnol.299:72-78.

14. Phan MD et al (2019). Establishing and validating noninvasive prenatal testing procedure

for fetal aneuploidies in Vietnam. J Matern Fetal Neonatal Med. 32:4009-4015.

15. Li Y, Sidore C, Kang HM, Boehnke M, Abecasis GR (2011). Low-coverage sequencing:

implications for design of complex trait association studies. Genome Res. 21:940-51.

16. Fumagalli M (2013). Assessing the effect of sequencing depth and sample size in

population genetics inferences. PLoS One 8:e79667.

17. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

18. Bolger AM, Lohse M, Usadel B (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114-20.

19. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997v2 [q-bio.GN].

20. Li H et al. The Sequence Alignment/Map format and SAMtools (2009). Bioinformatics

25:2078-9.

21. http://broadinstitute.github.io/picard/

22. Okonechnikov K, Conesa A, Garcia-Alcalde F (2016). Qualimap 2: advanced multi-sample

quality control for high-throughput sequencing data. Bioinformatics 32:292-4.

23. Quinlan AR, Hall IM (2010). BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26:841-2.

24. Robinson JT et al. Integrative Genomics Viewer (2011). Nat Biotechnol. 29:24-6.

25. Karimzadeh M, Ernst C, Kundaje A, Hoffman MM (2018). Umap and Bismap: quantifying

genome and methylome mappability. Nucleic Acids Res. 46:e120.

26. Van der Auwera GA et al (2013). From FastQ data to high confidence variant calls: the

Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 43:11.10.1-

33.

27. Cibulskis K et al (2013). Sensitive detection of somatic point mutations in impure and

heterogeneous cancer samples. Nat Biotechnol. 31:213-9.

28. https://software.broadinstitute.org/gatk/documentation/article?id=11136#2.1

29. Sherry ST et al (2001). dbSNP: the NCBI database of genetic variation. Nucleic Acids

Res. 29:308-11.

30. McLaren W (2016). The Ensembl Variant Effect Predictor. Genome Biol. 17:122.

31. Landrum MJ et al (2014). ClinVar: public archive of relationships among sequence

variation and human phenotype. Nucleic Acids Res. 42:D980-5.

32. Gouya L et al (2006). Contribution of a common single-nucleotide polymorphism to the

genetic predisposition for erythropoietic protoporphyria. Am J Hum Genet. 78:2-14.

33. Li H (2011). A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics 27:2987-2993.

## Figure Legends and Tables

**Figure 1.** Distributions of genome coverage and sequencing depth of the NIPT dataset. (a)

Average genome coverage and sequencing depth per sample and from all samples combined.

(b) Summary histogram of sequencing depth over all genome positions. (c) Distribution of

sequencing depth per chromosome. (d)  IGV tracks of sequencing depth, bwa MAPQ score, and

Umap k50 mappability across the whole genome.

**Figure 2.** Summary of the NIPT call set. (a) Venn diagram comparison between the NIPT call

set, the KHV and EAS call sets from the 1000 genomes project, and the dbSNP database. The

percentages were calculated with respect to the NIPT call set. (b) Scatter plot comparison of

allele frequency estimated from the NIPT and the KHV call sets. (c) Allele frequency distribution

of the NIPT call set. (d) Distribution of locations and effects of variants in the NIPT call set.

**Table 1. Summary of ClinVar annotations for the NIPT call set.**

| ClinVar clinical significance | in KHV | not in KHV |
|---|---|---|
| Benign or Likely benign | 23,414 | 300 |
| Uncertain significance | 353 | 74 |
| Pathogenic or Likely pathogenic | 4 | 1 |
| Drug response | 114 | 3 |
| Others | 211 | 13 |
| **Total number of annotations** | **24,096** | **391** |

**Table 2. Pathogenic variants identified from the NIPT call set.**

| Variant information | | | | | | ClinVar annotations | | | Allele frequency | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| chr | position | dbSNP | Ref | Alt | ID | Gene | Conditions | NIPT | gnomAD EAS | gnomAD |
| chr18 | 57571588 | rs2272783 | A | G | 562 | FECH | Erythropoietic protoporphyria | 28.10% | 32.57% | 11.23% |
| chr13 | 20189473 | rs72474224 | C | T | 17023 | GJB2 | Nonsyndromic hearing loss and deafness | 13.40% | 8.35% | 0.76% |
| chr13 | 72835359 | rs17089782 | G | A | 217689 | PIBF1 | Joubert syndrome | 6.80% | 5.13% | 1.36% |
| chr6 | 26090951 | rs1799945 | C | G | 10 | HFE | Hemochromatosis type 1 | 5.10% | 3.41% | 10.82% |
| chr2 | 31529325 | rs9332964 | C | T | 3351 | SRD5A2 | 5-alpha reductase deficiency | 2.90% | 0.67% | 0.05% |

**(a)**

| | Raw data | After filter |
|---|---|---|
| **Coverage and depth per bam** | | |
| Genome coverage | 14.59% | 12.17% |
| Positions with depth 1x | 12.60% | 12.17% |
| Positions with depth 2x | 1.75% | 0.00% |
| Positions with depth 3x | 0.20% | 0.00% |
| Positions with depth >= 4x | 0.05% | 0.00% |
| Average depth | 0.17 | 0.12 |
| **Aggregating 2,683 samples** | | |
| Genome coverage | 95.09% | 91.56% |
| Average depth | 462 | 364 |

**(b)**

average=364, error=67

**(c)**

**(d)**

**(a)**

NIPT: 8,054,515 SNPs

KHV: 12,710,234 SNPs

597,342
(7.4%)

7,390,020
(91.8%)

67,153
(0.8%)

novel
80,322
(1.0%)

dbSNP
517,020
(6.4%)

EAS: 22,724,409 SNPs

**(b)**

Pearson correlation = 98.8%

NIPT allele frequency

KHV allele frequency

**(c)**

Number of SNPs

- novel (80,322)
- in EAS, dbSNP (584,173)
- in KHV (7,390,020)

Allele frequency

**(d)**

novel (80,322)

| 0.6 | 30.2 | 1.0 | 13.4 | 3.2 | 50.7 | 0.9 |

in EAS, dbSNP (584,173)

| 0.6 | 38.7 | 0.7 | 15.3 | 2.9 | 40.6 | 1.2 |

in KHV (7,390,020)

| 0.6 | 52.6 | 0.9 | 11.7 | 4.0 | 29.4 | 0.8 |

Coding region   Intron   5' & 3' UTR   Up & Downstream   TF & regulatory   Intergenic   Others