The ClpX and ClpP2 Orthologs of *Chlamydia trachomatis* Perform Discrete and Essential Functions in Organism Growth and Development

Nicholas A. Wood[1], Amanda M. Blocker[2], Mohamed A. Seleem[3], Martin Conda-Sheridan[3], Derek J. Fisher[2], Scot P. Ouellette[1#]

[1]Department of Pathology and Microbiology, College of Medicine, University of Nebraska Medical Center, Omaha, NE

[2]School of Biological Sciences, Southern Illinois University Carbondale, Carbondale, IL

[3]Department of Pharmaceutical Sciences, College of Pharmacy, University of Nebraska Medical Center, Omaha, NE

Keywords: *Chlamydia*, differentiation, division, protein turnover, protein quality control, Clp protease, ClpX, ClpP

Running Title: ClpX ATPase of *Chlamydia*

#Corresponding Author:

Department of Pathology and Microbiology, College of Medicine, University of Nebraska Medical Center, 985900 Nebraska Medical Center (DRC2 5022), Omaha, NE

Tel: +1-402-559-0763 Fax: +1-402-559-5900 Email: scot.ouellette@unmc.edu

Abstract:

Chlamydia trachomatis (Ctr) is an obligate intracellular bacterium that undergoes a complex developmental cycle in which the bacterium differentiates between two functionally and morphologically distinct forms, the EB and RB, each of which expresses its own specialized repertoire of proteins. Both primary (EB to RB) and secondary (RB to EB) differentiation require protein turnover, and we hypothesize that proteases are critical for mediating differentiation. The Clp protease system is well conserved in bacteria and important for protein turnover. Minimally, the system relies on a serine protease subunit, ClpP, and a AAA+ ATPase, such as ClpX, that recognizes and unfolds substrates for ClpP degradation. In *Chlamydia*, ClpX is encoded within an operon 3' to *clpP2*. We present evidence that the chlamydial ClpX and ClpP2 orthologs are essential to organism viability and development. We demonstrate here that chlamydial ClpX is a functional ATPase and forms the expected homohexamer *in vitro*. Overexpression of a ClpX mutant lacking ATPase activity had a limited impact on DNA replication or secondary differentiation but, nonetheless, reduced EB viability with observable defects in EB morphology noted. Conversely, the overexpression of a catalytically inactive ClpP2 mutant significantly impacted developmental cycle progression by reducing the overall number of organisms. Blocking *clpP2X* transcription using CRISPR interference led to a decrease in bacterial growth, and this effect was complemented *in trans* by a plasmid copy of *clpP2*. Taken together, our data indicate that ClpX and the associated ClpP2 serve distinct functions in chlamydial developmental cycle progression and differentiation.

Words: 246/250

Importance

Chlamydia trachomatis is the leading cause of infectious blindness globally and the most

reported bacterial sexually transmitted infection both domestically and internationally. Given the

economic burden, the lack of an approved vaccine, and the use of broad-spectrum antibiotics for

treatment of infections, an understanding of chlamydial growth and development is critical for

the advancement of novel, targeted antibiotics. The Clp proteins comprise an important and

conserved protease system in bacteria. Our work highlights the importance of the chlamydial Clp

proteins to this clinically important bacterium. Additionally, our study implicates the Clp system

playing an integral role in chlamydial developmental cycle progression, which may help

establish models of how *Chlamydia* spp. and other bacteria progress through their respective

developmental cycles. Our work also contributes to a growing body of Clp-specific research that

underscores the importance and versatility of this system throughout bacterial evolution and

further validates Clp proteins as drug targets.

Words: 150/150

Introduction

        Chlamydia trachomatis (Ctr) is the leading cause of both bacterial sexually transmitted infections (STIs) and infectious blindness worldwide (1, 2). When left untreated in women, STIs can result in chronic sequelae, including pelvic inflammatory disease, ectopic pregnancy, and tubal infertility. A better understanding of Ctr molecular processes may help reveal essential systems that can be leveraged for more targeted intervention strategies.

        C. trachomatis is an obligate intracellular bacterial pathogen that differentiates between distinct functional and morphological forms during the course of its developmental cycle (see (3) for review). The elementary body (EB) is small (~0.3 μm in diameter), infectious, but non-dividing (4, 5). EBs attach to host cells and are internalized into host membrane-derived vacuoles that are rapidly modified into the inclusion (6-9). Within this inclusion, EBs undergo primary differentiation into larger (~1.0 μm in diameter) reticulate bodies (RBs). RBs are non-infectious but divide using a polarized budding mechanism (10) until secondary differentiation from RBs to EBs occurs. Many studies have detailed the transcriptional and proteomic differences between EBs and RBs (11-14). Given that chlamydial differentiation is not preceded by an unequal division and redistribution of intracellular proteins, as occurs in other bacteria like *Bacillus subtilis* (see (15) for review) or *Caulobacter crescentus* (16), and that EBs and RBs have distinct proteomes, we hypothesize that proteomic turnover plays an integral role in chlamydial differentiation.

        Previously, our groups characterized the two ClpP paralogs of *C. trachomatis*. We established that the *clp* protease-associated genes are expressed mid-developmental cycle and that ClpP1 and ClpP2 perform unique functions in chlamydial physiology based on differential effects related to overexpression of catalytically inactive mutants (17). In addition to the two

ClpP paralogs, Ctr encodes ClpX and ClpC orthologs (18). ClpX and ClpC are Type I AAA+

(**A**TPase **A**ssociated with diverse cellular **A**ctivities) unfoldases that utilize ATP hydrolysis to

linearize protein substrates for either degradation by the ClpP protease or refolding (19, 20).

Type I AAA+ ATPases encode Walker A and Walker B motifs, which are responsible for ATP

binding and hydrolysis, respectively (21, 22). These AAA+ unfoldases oligomerize to form

homo-hexamers that then recognize substrates through multiple different mechanisms (see (23)

for review).

Here, we characterized the role of ClpX in chlamydial growth and development. Because

ClpX is encoded within an operon with ClpP2, we also investigated effects of overexpression

and knockdown of both components. Ctr ClpX is highly conserved, possesses ATPase activity,

and formed the expected homohexamer *in vitro*. Interestingly, overexpression of wild-type ClpX,

ClpP2, and ClpP2X constructs in Ctr had little effect on bacterial growth, but overexpression of

the inactive mutants (alone or in tandem) had a substantial negative effect on recoverable

inclusion forming units (IFUs). However, the reduction in IFUs upon inactive ClpX

overexpression resulted from non-functional EB generation while the IFU reduction upon

inactive ClpP2 overexpression was the result of a block in developmental cycle progression. Our

results indicate that chlamydial ClpX is a true ortholog of bacterial ClpX and that the ClpP2X

gene products are integral to chlamydial growth and development.

Words: **500/500**

Results

The chlamydial ClpX retains conserved motifs of, and exhibits predicted structural homology

to, ClpX orthologs. To initiate our study, we first performed bioinformatic and *ab initio*

structural modeling analyses to determine whether the chlamydial ClpX (ClpX$_{Ctr}$) possesses the

expected conserved regions and motifs consistent with its proposed function as an AAA+

ATPase. Using multiple sequence alignment, we aligned ClpX$_{Ctr}$ to ClpX orthologs and

annotated conserved motifs identified in other studies (Fig. 1a). ClpX$_{Ctr}$ retains the N-terminal

metal binding domain (24, 25), the Walker A and B motifs for ATP binding and hydrolysis,

respectively (21, 23), the sensor motifs for recognition of nucleotide bound state (26), the RKH

motif and pore loops for substrate recognition (27-29) and unfolding (30, 31), the arginine finger

for inter-subunit sensing of nucleotide state in the ClpX hexamer (22, 32), and the IGF Loop for

interaction with ClpP (33, 34). Interestingly, the predicted secondary structure of ClpX$_{Ctr}$ shows

few notable aberrations (see Discussion) from other prototypical bacterial ClpX orthologs and is

predicted to form the expected homohexamer by structural modeling (Fig. 1b, two subunits

removed for clarity). The spatial conservation of AAA+ and ClpX-specific motifs (colored in

Fig. 1b as in the multiple sequence alignment) indicates that the chlamydial ClpX likely

functions using a mechanism similar or identical to other ClpX orthologs. Taken together, these

in silico studies suggest that ClpX$_{Ctr}$ functions as a canonical AAA+ ATPase.

Chlamydial ClpX forms the expected homohexamer and possesses ATPase activity. To

determine the oligomeric state of ClpX$_{Ctr}$ *in vitro*, we purified recombinant protein and analyzed

its migration by native PAGE. At the same time, we also constructed a Walker B ATPase mutant

(E187A) ClpX$_{Ctr}$ as a control for biochemical studies. Following the incubation of 10 µg of wild-

type or mutant ClpX$_{Ctr}$ for 20 minutes in a HEPES based buffer, we loaded the entire volume

into a 4-20% gradient gel. We observed the ClpX$_{Ctr}$ proteins migrating above the 242 kDa band

of the molecular weight ladder, which is close to the expected hexameric size of 283 kDa (Fig.

2a). We then sought to assess ATPase activity of recombinant wild-type and ATPase mutant

ClpX$_{Ctr}$ using the Kinase-Glo® endpoint assay, which measures ATP remaining in a sample at

the end of the reaction period. Indeed, ClpX$_{Ctr}$ hydrolyzed ATP, while the ATPase mutant

isoform showed a significant defect in ATP hydrolysis (Fig. 2b). These data indicate that ClpX$_{Ctr}$

(i) forms a homohexamer of the predicted size and (ii) possesses ATPase activity that is

abrogated by a mutation in the Walker B motif.

We next tested whether wild-type and ATPase mutant ClpX$_{Ctr}$ interact with each other

using the Bacterial Adenylate Cyclase Two-Hybrid (BACTH) assay. This system is predicated

on the reconstitution of adenylate cyclase activity by bringing two complementary fragments of

the enzyme (T25 and T18) into close proximity by interacting proteins. Generation of cAMP by

the reconstituted adenylate cyclase drives ß-galactosidase production that can be measured

qualitatively by the presence of blue colonies and growth on minimal medium (Fig. 2c) or

quantitatively by measuring enzyme activity directly (Fig. 2d). We performed a series of

pairwise interaction tests between the wild-type and mutant ClpX$_{Ctr}$. In each instance, we

observed a positive interaction that was quantifiable and on par with the positive control (T25-

Zip vs T18-Zip). We conclude from these data that the mutant isoform can interact with the wild-

type isoform.

Overexpression of ATPase mutant ClpX or catalytically inactive ClpP2 has both distinct and

overlapping effects. We previously measured the effects of overexpression of both wild-type and

catalytically inactive ClpP2$_{Ctr}$ on chlamydial growth and observed a modest reduction in growth

at 24 hours post-infection (hpi) (17). We wanted to more carefully assess growth differences

during the chlamydial developmental cycle in the presence of overexpressed wild-type and

mutant ClpX$_{Ctr}$ and ClpX$_{E187A}$, ClpP2$_{Ctr}$ and ClpP2$_{S98A}$, or both together (ClpP2X$_{Ctr}$ and

ClpP2$_{S98A}$ClpX$_{E187A}$). To do this, we performed growth curves where we induced expression, or

not, at 10 hpi and quantified growth at various timepoints after induction. Immunofluorescence

analysis (IFA) of replicate treatments and quantification of recoverable inclusion forming units

(IFUs; a proxy for EBs) revealed distinct effects upon overexpression of the individual

components (Fig. 3a-c) as well as with the entire operon (Fig. 3d&e). We noted that

overexpression of wild-type ClpP2$_{Ctr}$ showed no appreciable effect at either 24 or 48 hpi (14 and

38 h pulses of induction, respectively), whereas overexpression of ClpP2$_{S98A}$ appeared to reduce

the number of organisms present within the inclusion at 48 hpi but not at 24 hpi (Fig. 3a). These

observations correlated with a marked decrease in EB production in the later time points of

mutant ClpP2$_{S98A}$ but not wild-type ClpP2$_{Ctr}$ overexpression (Fig. 3b). Conversely, ATPase

mutant ClpX$_{E198A}$ overexpression resulted in smaller inclusions and a decrease in IFUs (Fig.

3a&c). Effects on IFU recovery suggest that Ctr is more sensitive to ClpX$_{Ctr}$ rather than ClpP2$_{Ctr}$

disruption earlier in the developmental cycle, as the IFU reduction is exacerbated sooner with

ClpX$_{E198A}$ overexpression (observe the differences at 24hpi in Fig. 3b&c). As noted for the

overexpression of individual wild-type isoforms, there was no significant impact on IFU

recovery of overexpressing both wild-type ClpP2$_{Ctr}$ and ClpX$_{Ctr}$ in tandem. Consistent with the

effects of overexpressing individual mutant isoforms, overexpression of the mutant ClpP2$_{S98A}$

and ClpX$_{E187A}$ isoforms in tandem showed an exacerbated phenotype throughout the

developmental cycle as noted by both IFA and IFU assays (Fig. 3d&e). Importantly, the wild-

type chromosomal copies of $ClpP2_{Ctr}$ and $ClpX_{Ctr}$ continue to be expressed during these

overexpression assays. Therefore, the true impact of overexpression of the mutant isoforms is

likely underrepresented.

Functional disruption of ClpP2 blocks developmental cycle progression while ClpX disruption

reduces EB viability. Given that the IFU assay only measures EB viability from a population and

not total bacterial numbers or differentiation status, we next wanted to address these nuances of

the chlamydial developmental cycle. We first measured genomic DNA as a proxy for total

number of bacteria (i.e. both RBs and EBs). Overexpression of any wild-type protein had no

significant impact on DNA accumulation. From 24 hpi to 48 hpi, we observed a significant drop

in the rate of increase of gDNA levels when $ClpP2_{S98A}$ was overexpressed alone or in the mutant

operon configuration (Fig. 4a). Overexpression of the $ClpX_{E187A}$ showed a trend (two-fold)

towards decreased DNA levels that was not statistically significant. This was surprising given the

roughly 20-fold reduction in IFUs (Fig. 3), suggesting total bacterial numbers during $ClpX_{E187A}$

overexpression are unaffected.

To determine if there was an impact on differentiation status during overexpression of

wild-type and mutant isoforms, we next assessed HctB levels, an EB-specific gene product (35-

37), by western blot as an indicator of secondary differentiation. We normalized the integrated

density of HctB to the integrated density of MOMP (major outer membrane protein; present in

both EBs and RBs) to ensure that we were comparing HctB levels to the total number of bacteria

to give a more robust readout of differentiated bacteria within the population. The relative HctB

levels in samples where $ClpP2_{S98A}$ was overexpressed were significantly reduced as expected,

indicating the generation of fewer EBs and consistent with IFU and genomic DNA data, whereas

the other experimental conditions showed no changes in relative HctB levels (Fig. 4b&c). These

data suggest that overexpression of $ClpX_{E187A}$ does not impact total number of bacteria, as

measured by gDNA levels, or RB-to-EB differentiation, as measured by HctB levels.

Nevertheless, the recoverable IFUs were decreased when $ClpX_{E187A}$ is overexpressed (Fig.

3C&E), indicating a defect in the EBs being produced.

To further explore this possibility, we prepared samples for transmission electron

microscopy to examine at higher resolution the morphology of EBs and RBs from $ClpP2_{S98A}$ and

$ClpX_{E187A}$ overexpressing strains. As RBs condense to EBs, an intermediate form can be

observed where the chlamydial nucleic acids are visible as an electron dense spot. Consistent

with other measured effects (Fig. 4d&e; Suppl. Fig. S1a), $ClpP2_{S98A}$ overexpression resulted in

smaller inclusions with fewer organisms whereas $ClpX_{E187A}$ appeared to have no effect on

bacterial numbers (Fig. 4d&e). In both induced and uninduced samples for each strain, we noted

normal RBs (Fig. 4d&e, white triangles) and normal EBs (Fig. 4d&e, black triangles). However,

we observed numerous abnormal developmental forms following $ClpX_{E187A}$ overexpression that

appeared to be multi-nucleated or abnormally condensing intermediate forms (Fig. 4e, gray

triangles). These data are consistent with the limited effect of $ClpX_{E187A}$ overexpression on the

production of HctB (Fig. 4b&c) but the decreased recoverable IFUs (Fig. 3 a&c), further

supporting that disruption of normal ClpX function may affect EB viability.

Knockdown of the* clpP2X *operon reduces recoverable progeny and results in reduced plasmid

retention while complementation with* clpP2 *partially restores infectivity. Overexpression of

mutant isoforms of $ClpP2_{Ctr}$ and/or $ClpX_{Ctr}$ was sufficient to disrupt chlamydial development in

the presence of endogenous $ClpP2X_{Ctr}$. However, we wanted to directly block the chromosomal

copies by employing an improved version of the chlamydial CRISPR interference (CRISPRi)

strategy previously described by us ((38) and Ouellette et al, *in prep*). CRISPRi relies on the

inducible expression of a catalytically inactive Cas9 (dCas9) in combination with a guide RNA

(gRNA) to block transcription at specific chromosomal sites (39). We transformed Ctr L2 with

vectors encoding inducible dCas9 and constitutive gRNAs targeting either the *clpP2X* or *incA*

intergenic regions (Fig. 5A and Suppl. Figs. S2&3). Of note, *pcnB2*, a predicted poly-A

polymerase, is encoded at the 3' end of the *clpP2X* operon and was included in our analyses. To

determine whether we could also complement the effects of knockdown, we constructed vectors

encoding either *clpP2_FLAG* or *pcnB2_FLAG* in a transcriptional fusion 3' to dCas9. These

vectors were transformed into *C. trachomatis* L2. Under conditions when dCas9 is expressed,

ClpP2_FLAG or PcnB2_FLAG is also expressed (Fig. 5b and Suppl. Fig. S4). IncA knockdown

served as a control since *incA* is a non-essential gene (40). These CRISPRi transformants were

used to infect HEp2 cells.

232   When dCas9 expression was induced at 10hpi in the various transformants, we observed

a marked and rapid decrease in *clpP2*, *clpX*, and *pcnB2* transcript levels compared to the

uninduced controls at 14hpi (Fig. 5c). Importantly, we did not observe a decrease in transcript

levels for unrelated genes associated with different stages of the developmental cycle (*clpP1,*

euo, and *omcB*; Suppl. Fig. S2; (12, 17, 41, 42)). We also observed abnormal inclusion

morphology with individual organisms aggregated to one side of the inclusion (Fig. 5a).

Complementation with ClpP2_FLAG resulted in decreased *clpX* transcripts as compared to the

uncomplemented strain but an increased level of *clpP2* transcripts, as expected (Fig. 5f).

However, ClpP2 complementation did not fully restore inclusion morphology as organisms still

displayed   some   aggregation   within   the   inclusion,   albeit   intermediate   between   the

uncomplemented knockdown and control conditions (Fig. 5b). Complementation with

PcnB2_FLAG did not impact *clpP2* and *clpX* transcript levels during knockdown but did

increase *pcnB2* transcripts approximately 10-fold relative to the uninduced samples (Suppl. Fig.

S4b). Abnormal inclusion morphology was also evident (Suppl. Fig. S4a). As previously

observed, IncA expression was uniformly blocked after dCas9 induction (Suppl. Fig. S3; (38)).

We also attempted a similar *clpX* complementation strategy but saw no effect on *clpP2* or *pcnB2*

transcript levels during knockdown (unpublished observation). Given that we tagged our dCas9

variant with an SsrA recognition motif and that ClpX recognizes the SsrA tag (see below), the

dCas9 was likely targeted for degradation by the plasmid-encoded ClpX, subsequently

eliminating any knockdown.

We next assayed chlamydial growth as measured by IFU recovery and chlamydial

morphology by immunofluorescence after inducible knockdown of the target genes. Expression

of dCas9 was induced at 4hpi, and IFUs were harvested at 24 and 48hpi and titred on fresh cell

monolayers in the presence of penicillin, the selection agent. When *clpP2X* expression was

blocked at 4hpi, we noted a 5-fold decrease in penicillin-resistant (i.e. transformants containing

the CRISPRi plasmid) IFUs at 24hpi but a more than 200-fold decrease at 48hpi (Fig. 5d).

Immunofluorescence assay images supported this drop in viable organisms, revealing a severely

diminished number of organisms within the inclusion (Fig. 5a). In performing these assays in the

presence of penicillin, we also observed numerous penicillin-sensitive organisms (i.e. aberrant

RBs (43)) during the titration step, suggesting that the plasmid conferring resistance and

encoding the CRISPRi system was being lost after induction of dCas9 expression. To test this,

we quantified plasmid retention in the *clpP2X* knocked down samples and observed that

blocking *clpP2X* expression resulted in ~75% plasmid loss at 24hpi and greater than 90% loss at

48hpi (Fig. 5e), which was closely recapitulated with *pcnB2* complementation (Suppl. Fig.

S4c&d). These effects on IFUs and plasmid retention were not observed for *incA* knockdown

(Suppl. Fig. S3b&c). We note that *incA* knockdown did result in a reproducible, but transient,

increase in IFUs at 24hpi that returned to "normal" levels at 48hpi (Fig. S3b). The reasons for

this are not clear. Nonetheless, we conclude from these data that blocking *clpP2X* expression is

deleterious to *Chlamydia*, further highlighting its essentiality to this pathogen. Interestingly,

complementation with *clpP2* largely rescued the drop in recoverable IFUs while maintaining

plasmid loss as compared to the uncomplemented condition (Fig. 5g&h). This suggests that

much of the loss in bacterial viability during *clpP2X* knockdown is attributable to reduced ClpP2

levels.

ClpX function is required for degradation of an SsrA-tagged substrate in Ctr. Recently, ClpX-

specific inhibitors were synthesized by the Sieber group and were shown to interfere with ClpX

ATPase activity (44). One compound, identified as 334, was shown to have potent anti-ClpX

activity whereas a derivative, 365, was inactive. We performed *ab initio* modelling and

molecular dynamics simulations (45) to determine if these compounds could interact with an

ADP-bound hexameric ClpX$_{Ctr}$. For 334, a high scoring model (-9.1 kcal/mol binding affinity,

RMSD ~ 0) was predicted with the drug binding near to the ATP binding pocket, suggesting a

mechanism of action where 334 likely occludes the ATPase site (Suppl. Fig. S5). Whether the

effect stems from the blocking of ATP binding and subsequent destabilization of the complex,

attenuation of ATPase function by preventing a conformational change of the complex, or steric

hindrance of complex formation remains to be elucidated. Conversely, compound 365 bound

outside of the ATP pocket with a much lower score (Suppl. Fig. S6).

Given the predicted binding of the ClpX inhibitors on the structure of ClpX$_{Ctr}$, we sought

to determine whether these drugs effectively blocked ClpX function *in vivo*. We utilized a

chlamydial SsrA-tagged GFP variant [GFP(VAA)], which has been previously shown to be a

substrate of ClpX$_{Ctr}$ *in vitro* (46), or GFP tagged with a mutated SsrA tag [GFP(VDD)]. In

mutating VAA to VDD, GFP is rendered resistant to ClpX recognition (47). Using an

anhydrotetracycline-inducible pBOMB4 derivative encoding constitutive mCherry as a metric of

chlamydial growth (herein referred to as pBOMBmC), we cloned each GFP variant into

pBOMBmC and transformed these constructs into Ctr L2. Samples were infected, and GFP

expression was induced at 8 hpi. All induced samples were then allowed to develop until 16 hpi,

when media was replaced by either DMEM only or DMEM containing the indicated antibiotic

(Fig. 6a&b, Suppl. Fig. S7). Immediately following drug addition, induction of GFP expression

was maintained or not, where removal of aTc is denoted by Pulse/Chase. GFP signal was

normalized to the constitutive mCherry signal expressed on the plasmid backbone. We observed

that removal of inducer results in rapid loss of GFP(VAA) but not GFP(VDD) signal compared

to continuous induction, indicating that GFP(VAA) is degraded as expected (Fig. 6a&b, Suppl.

Fig. S7). Treatment with compound 334 resulted in sustained GFP(VAA) signal, consistent with

its documented effects on ClpX (44). Importantly, compound 365, the inactive 334 derivative,

and ACP1b, a ClpXP uncoupling drug with a mechanism of action similar to the ADEP

compounds (17, 48), failed to protect GFP(VAA) from degradation (Fig. 6a), which further

supports the specificity of compound 334 in mediating this effect. We then tested the effect of

these drugs on chlamydial viability and noted a multiple log$_{10}$ decrease in recoverable IFUs

(Supp. Fig. S8), supporting that blocking ClpX function is highly detrimental to *Chlamydia*.

However, in attempting to assess the effect of these compounds on *Chlamydia*, we noted reduced

host cell viability with prolonged treatment (Supp. Fig. S8d), making interpretation of those data

unclear.

Discussion

Given the unique roles and protein repertoires of the chlamydial developmental forms

(EB/RB), we hypothesize that protein degradation is a critical factor in the differentiation process

from one form to the other. The Clp system is highly conserved in both prokaryotic and

eukaryotic systems where it has been described to perform important functions in both

proteostasis and pathogenesis (49). The Clp system is nominally composed of a proteolytic

subunit, ClpP, and a AAA+ ATPase that functions as an unfoldase to recognize substrates and

feed them into the ClpP barrel for degradation (23). The work presented here expands our

understanding of the Clp protease system of an obligate intracellular bacterial pathogen,

Chlamydia trachomatis. Focusing on characterization of $ClpX_{Ctr}$ and the function of the *clpP2X*

operon, we demonstrated the importance of the ClpXP protease system during chlamydial

growth and development.

Multiple lines of evidence support that the chlamydial ClpX is a *bona fide* AAA+

ATPase. Firstly, multiple sequence alignment of $ClpX_{Ctr}$ to orthologs of other bacteria revealed a

perfect conservation of the motifs involved in nucleotide binding, ATP hydrolysis, and

nucleotide-state sensing (Fig. 1) (50, 51). Secondly, homology-directed and *ab initio* modelling

of $ClpX_{Ctr}$ revealed that the spatial orientation of these domains is conserved as well (Fig. 1),

though we acknowledge that structural studies are critical to drawing conclusions about $ClpX_{Ctr}$

conformational states. Thirdly, $ClpX_{Ctr}$ interacts with itself to form a homohexamer that

possesses ATPase activity (Fig. 2). Importantly, this ATPase activity could be disrupted by a

targeted mutation in the Walker B motif while having no effect on the oligomerization properties

of the protein. Fourthly, a characterized ClpX inhibitor that disrupts its ATPase activity also

blocked SsrA-mediated degradation *in vivo* and disrupted the growth of *C. trachomatis* serovar

L2 (Fig. 6; Suppl. Fig S7). Finally, overexpression of a ClpX$_{Ctr}$ ATPase mutant *in vivo*

negatively impacted chlamydial growth and development (Figs. 3&4).

While we have characterized the ATPase function of ClpX$_{Ctr}$ and its role in chlamydial

growth, further work remains to determine whether this ClpX ortholog functions as an unfoldase.

Nevertheless, our bioinformatics analysis supports this as ClpX$_{Ctr}$ retains substrate recognition

motifs, including both pore loops and the RKH motif for gripping and translocation of substrates

(27-31). *Chlamydia* spp. also encode the tmRNA/ssrA tagging system for ribosomal rescue (18,

52-55), which fits a model where ClpX$_{Ctr}$ may play an integral role in turnover of tagged,

partially translated peptides. A recent article, using an *in vitro* assay for SsrA-tagged GFP

degradation, suggests this function of chlamydial ClpX may be conserved (46). Our *in vivo* data

exhibiting SsrA-tagged GFP protection with the addition of a ClpX inhibitor also suggests

ClpX$_{Ctr}$ can target SsrA-tagged substrates, but whether this tagging is for ribosomal rescue or

more specific purposes (56, 57) remains to be determined and is currently under investigation by

our research group.

In *Chlamydia*, *clpX* is encoded in an operon with *clpP2*. Our data indicate that, not

surprisingly, the ClpP2X$_{Ctr}$ system is highly regulated and essential. We previously demonstrated

that treatment with ClpP-targeting antibiotics severely impacts growth of *Chlamydia* (17). Here,

we performed a systematic analysis of the effects of overexpression of wild-type or inactivated

ClpP2X$_{Ctr}$ components. The overexpression of wild-type ClpP2$_{Ctr}$ and/or ClpX$_{Ctr}$ had no

biologically or statistically significant effect on chlamydial growth that we could measure.

However, extended overexpression of mutant $ClpP2_{Ctr}(S98A)$ and/or $ClpX_{Ctr}(E187A)$ resulted in

abrogation of chlamydial growth as measured by recovery of infectious progeny. Three

observations should be noted. Firstly, the effect of inducibly-expressed proteins is measured in

the presence of the endogenous chromosomally-expressed copies of these proteins. Therefore, it

is likely that the mutants would have even more dramatic effects on chlamydial growth and that

disruption of these components results in lethality. For $ClpX_{Ctr}$, this is supported by the effects of

the ClpX inhibitor on *Chlamydia* (Supp. Fig. 8), which effectively stopped chlamydial growth.

Secondly, we demonstrated that the mutant proteins could interact *in vitro* with wild-type

isoforms (Fig. 2 and (17)). Therefore, we can infer that overexpression of the mutant proteins

leads to their incorporation into the endogenous Clp machinery to disrupt or impair its function.

Thirdly, to our knowledge, ours is the first study to ectopically express two different tagged

proteins in *Chlamydia*, showing both the feasibility of this approach and its potential utility to

dissect chlamydial biology.

The overexpression of the catalytically inactive mutant Clp proteins in *Chlamydia*

revealed potentially subtle differences in the role of each component in chlamydial growth and

development. Surprisingly, we noted a roughly 50% reduction in detectable genomes (Fig. 4A)

when $ClpX_{Ctr}(E187A)$ was expressed whereas IFUs were reduced roughly 20-fold (Fig. 3). The

production of EBs as measured by HctB levels did not appreciably change (Fig. 4b&c). This

suggests that, while development is hindered, the drop in IFUs may be due to defective EB

viability, infectivity, or inclusion establishment and not a reduction in secondary differentiation

per se. Support for this comes from electron microscopy images, which revealed unusual

morphologies after overexpression of the mutant $ClpX_{Ctr}$ isoform (Fig. 4d; Suppl. Fig. S1). As

noted above, the SsrA-mediated degradation of partially translated products may be critical

during the differentiation process. As RBs condense to EBs, DNA replication, transcription, and

translation are significantly abrogated, and we hypothesize that those partially translated proteins

trapped during this differentiation step require degradation. This may explain the effects we

observed on decreased EB viability during overexpression of the ClpX mutant isoform. Whether

the degradation of such partial peptides occurs during RB-to-EB differentiation or during the

next infection when an EB differentiates to an RB remains to be investigated.

386   Conversely, for $ClpP2_{Ctr}$(S98A) overexpression, the substantial IFU decrease coupled

with a sharp drop in gDNA levels indicate that $ClpP2_{Ctr}$ plays a role in developmental cycle

progression. HctB levels are also significantly reduced, which is consistent with the lack of EB

generation. Taken together, these data may indicate that $ClpP2_{Ctr}$ is integral to developmental

cycle progression or differentiation and that its function is tightly regulated. We cannot,

however, conclude that secondary differentiation is directly affected due to the fact that total

organism numbers are severely reduced. Rather, our proposed model suggests that $ClpP2_{Ctr}$

disruption may affect both factors by a mechanism that we are currently working to identify.

Conversely, $ClpX_{Ctr}$ may serve a more prominent $ClpP2_{Ctr}$-independent function in

differentiation of the organism (Figure 7). Interestingly, several studies on the periplasmic

protease, HtrA, in *Chlamydia* have implicated this protease in membrane reorganization (58-60).

Therefore, one model may be that the ClpXP system may be required for cytosolic turnover

whereas HtrA may be required for membrane reorganization during differentiation.

399   We successfully generated chlamydial transformants with an inducible knockdown

system to repress $ClpP2X_{Ctr}$ expression. To date, this study is the first of its kind in *Chlamydia* to

knock down genes that are essential, highlighting the utility of CRISPRi in studies of chlamydial

biology while providing insight into possible $ClpP2X_{Ctr}$ function. We also demonstrated that

these knockdowns may be complemented by insertion of the gene of interest under the inducible

promoter as a transcriptional fusion with the dCas9 (Fig. 5 and Ouellette et al, *in prep*). Notably,

we observed a large decrease in IFU production coupled with an increase in plasmid loss after

inhibition of *clpP2X* expression (Fig. 5). These effects were not observed when targeting a non-

essential gene. Additionally, we demonstrated that the impact on IFUs, but not the effect on

plasmid retention, was reversed by complementation. Of note, penicillin does not kill chlamydiae

but blocks cell division (61, 62), which keeps the organism transcriptionally in an RB-like state

(63). This suggests that knocking down an essential gene(s) puts selective pressure on the

chlamydiae to lose the plasmid encoding the CRISPRi system. This has important ramifications

for long-term experiments and functional analyses. Nevertheless, the CRISPRi system represents

a significant advance for our ability to study essential systems in this obligate intracellular

bacterium.

A recent *in vitro* study suggested that homoheptamers of ClpP1 form a complex with

homoheptamers of ClpP2 that interact with ClpX (46), yet we observed distinct differences when

overexpressing mutant isoforms of these proteins in *Chlamydia* (17). If these ClpP proteins also

interact with the other chlamydial AAA+ ATPase, ClpC, then it is not surprising that we

observed differences between ClpP2 and ClpX in our assays since disrupting ClpP2 could also

be impacting ClpC function. This would also beg the question why we did not observe similar

impacts on chlamydial growth when overexpressing each mutant isoform (17). One possibility

may be that ClpX interacts with ClpP2 in a ClpP1P2 complex whereas ClpC interacts with

ClpP1 in a ClpP1P2 complex. In this scenario, disrupting ClpP2 would act epistatically to ClpX.

This is not inconsistent with our data and proposed model. Clearly, the composition of the Clp

complexes remains to be determined *in vivo*, and this is an area of ongoing investigation.

In conclusion, we have demonstrated the importance of the ClpP2X$_{Ctr}$ system to

chlamydial development, but many questions remain unanswered. These include why ClpP2$_{Ctr}$

and ClpX$_{Ctr}$ may serve independent purposes and what substrates this system may be targeting.

Additionally, we need to identify any cofactors, chaperones, adaptor proteins, or a lack thereof

that may be pertinent to this system. We plan to dissect the structural motifs of ClpP2$_{Ctr}$ and

ClpX$_{Ctr}$ to determine if any of the noted differences from other bacterial Clp proteins may alter

activity, which may aid in our goal of further functional assessment. Finally, continued

experimentation to address our overarching hypothesis that protein turnover is critical to

chlamydial differentiation, and that the Clp system is a key mediator of this process, is required.

Overall, we conclude that the chlamydial ClpP2X$_{Ctr}$ system is critical to the development of

these obligate intracellular bacteria.

Materials and Methods:

Strains and Cell Culture: The human epithelial cell line HEp2 was used for the overexpression assays, gDNA and protein extractions, and antibiotic studies. McCoy mouse fibroblasts were used for chlamydial transformation, and human epithelial HeLa cells were used for plaque purification. All of these cell lines were passaged routinely in Dulbecco's Modified Eagle's Medium (DMEM, Gibco/ThermoFisher) and 10% FBS (Sigma; St. Louis, MO) and verified to be mycoplasma-free using LookOut® Mycoplasma PCR Detection Kit (Sigma). Density gradient purified *Chlamydia trachomatis* L2/434/Bu (ATCC VR902B) EBs were used for the antibiotic studies. *C. trachomatis* serovar L2 EBs (25667R) naturally lacking the endogenous plasmid were prepared and used for transformation [see (64)].

Bioinformatics Analysis: Gene sequences of *Chlamydia trachomatis* were obtained from STDGen database (http://stdgen.northwestern.edu) or KEGG Genome Browser (65-67). RefSeq protein sequences from *Escherichia coli, Bacillus subtilis, Mycobacterium tuberculosis, Staphylococcus aureus,* and *Pseudomonas aeruginosa* were acquired from the NCBI protein database (https://www.ncbi.nlm.nih.gov/guide/proteins/). ClpX pairwise protein alignments to find sequence identity were performed using NCBI Protein BLAST function (https://blast.ncbi.nlm.nih.gov/Blast.cgi) (68). Multiple sequence alignments were performed using Clustal Omega (69) with default settings and were presented using Jalview Version 2 (70). PDB files for predicted monomeric 3D structures were acquired from the Phyre2 website (http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index) (71). Complexes were modelled using SWISS-MODEL available on the ExPASy server (72-75). Protein models and model alignments were rendered using the UCSF Chimera package from the Computer Graphics

Laboratory, University of California, San Francisco (supported by NIH P41 RR-01081) (76).

Docking analyses were performed using AutoDock Vina (45). Molecules were prepped using

Dunbrack rotamer libraries (77, 78) to replace incomplete side chains and ANTECHAMBER for

charge assignment and topology generation (79).

Plasmid Construction: A full list of the primers and plasmids used is included in the

supplementary material. The Gateway® recombination system of cloning was used for plasmids

for the Bacterial Adenylate Cyclase Two-Hybrid (BACTH) system (80). The genes were

amplified from *Chlamydia trachomatis* L2 genomic DNA with added *attB* recombination sites.

The PCR products were then incubated with a pDONR^{TM}221 entry vector (containing *attP*

recombination sites) in the presence of BP Clonase II (Invitrogen) to insert the gene via flanking

attP recombination sites and remove the *ccdB* insert, resulting in an entry vector containing the

gene of interest flanked by *attL* sites. These constructs were transformed into DH5α chemically

competent *E. coli* and plated onto kanamycin-containing LB agar. Plasmid was isolated and used

for the LR reaction into one of three destination vectors (pST25-DEST, pSNT25-DEST, or

pUT18C-DEST). The same entry vector for any given gene was used for all three LR reactions

to insert into the destination vector. Entry vector and destination were incubated in a 1:1 ratio.

DH5α *E. coli* were transformed with 2 μL of the reaction mix. Purified plasmid from an

individual colony was sequence verified prior to use in the BACTH assay (see below).

Constructs for chlamydial transformation were created using the HiFi Cloning (New

England Biolabs) protocol. Primers were designed to add a poly-Histidine (6xHis) tag to the

gene of interest with the overlap to insert into the shuttle vector. Primers were generated using

the NEBuilder® assembly tool available from New England BioLabs (http://nebuilder.neb.com).

The backbone used was the pTLR2 derivative of the pASK plasmid (81). For the CRISPRi plasmid, the *S. aureus* dCas9 was PCR amplified from pX603-AAV-CMV::NLS-dSaCas9(D10A,N580A)-NLS-3xHA-bGHpA (a gift from Dr. F. Zhang; Addgene plasmid # 61594 (39)) and inserted into a derivative of pBOMB4-Tet::L2 (kind gift of Dr. T. Hackstadt, NIH; (82)) modified to weaken its ribosome binding site (Ouellette *in prep*). The gRNA cassettes were designed as previously described (38)), ordered as gBlock fragments from IDTDNA (Coralville, IA), and inserted into the BamHI site of the pBOMB4-Tet derivative encoding *Sa_dCas9* to produce, for example, the plasmid pBOMBLCRia::L2 (*clpP2X*). HiFi reactions were assembled according to the manufacturer's protocol. The reaction was transformed into DH10ß *E. coli*, and isolated plasmid was verified by restriction enzyme digest and sequencing by Eurofins Genomics. Sequence verified plasmids were transformed into *dam-/dcm- E. coli* (New England BioLabs) to produce demethylated plasmid, which was verified as described earlier prior to transformation into *C. trachomatis* (see below).

For mutation of ClpX Walker B motif, Q5 mutagenesis (New England BioLabs) was used. Primers were designed encoding the E187A mutation for PCR linearization of the plasmid. ClpX BACTH constructs were used as a template for the PCR amplification, and plasmids were re-circularized by KLD reaction. The resulting reactions were transformed into DH5α *E. coli* for plasmid production. Plasmids were isolated, and mutations were verified by Sanger sequencing (Eurofins Genomics) prior to use in the BACTH system. These plasmids also served as template for the PCR reactions to produce PCR products for insertion of the mutant *clpX* gene into the pTLR2 plasmid.

Strains created or used in this study are listed in the supplementary material. Transformed *E. coli* strains were maintained on LB agar plates, with antibiotics as necessary. To extract

chlamydial genomic DNA, EBs were subjected to heat and proteinase K treatment prior to

phenol:chloroform extraction (83). Sodium hydroxide lysis was utilized for the extraction of *E.*

coli genomic DNA. For cloning into the pLATE31 plasmid, the aLICator LIC Cloning and

Expression Kit 3 (Thermo Scientific) was used according to the manufacturer's specifications.

Plasmids were first cloned into DH5α *E. coli* for plasmid propagation. Transformants were

screened for inserts using colony PCR with Fermentas Master Mix (Thermo Scientific) and

positive clones were grown for plasmid isolation (GeneJet Plasmid Miniprep Kit, Thermo

Scientific). Sequence verified plasmids were then transformed into BL21(DE3) Δ*clpPAX E. coli*

(46) for subsequent protein purification.

Purification of Recombinant ClpX: His-tagged Ctr ClpX and Ctr ClpX$_{(E187A)}$ were purified from

500 mL cultures of BL21(DE3) Δ*clpPAX E. coli* transformed with the respective plasmid based

on the protocol described in (17). Samples were induced with 0.5 mM IPTG and incubated with

shaking for 20 hours at 18°C. Cultures were pelleted and frozen at -80°C prior to purifications.

Samples were suspended in buffer A (25 mM Tris Base [pH 7.5], 300 mM NaCl, and 10 mM

Imidazole), sonicated, bound to HisPur Cobalt Resin (Thermo Scientific), and washed in buffer

A Proteins were eluted from the resin using buffer B (25 mM Tris Base [pH 7.5], 300 mM NaCl,

and 300 mM Imidazole). Buffer exchange for ATPase assay buffer (25 mM HEPES [pH 7.2],

200 mM KCl, 20 mM MgCl$_2$, and 10% glycerol) was performed using a Millipore Amicon Ultra

15 filtration units (3 kDa cut-off). ClpX proteins were quantified using the Bio-Rad Protein

assay, assessed for purity on 10% SDS-PAGE gels with Coomassie staining (Suppl. Fig. S9), and

identified using anti-His-tag western blot. Blotting was performed using a mouse monoclonal

anti-6x His antibody (1:1000; Millipore HIS.H8) and a goat anti-mouse IgG HRP conjugated

secondary antibody (1:2000). Protein samples were aliquoted and stored at -80°C.

In Vitro *Analysis of ClpX Homo-Oligomerization:* 10 µg of purified protein was incubated at

for 20 minutes at 37°C in oligomerization buffer (25 mM Tris Base [pH 7.5], 5 mM KCl, 5 mM

$MgCl_2$, 1 mM DTT, and 1% glycerol) prior to mixing with a 5x native sample buffer (5 mM Tris

[pH 6.8], 38 mM glycine, 0.06% bromophenol blue). Assays were analyzed on a BioRad

MiniProtean 4-20% gradient gel for Native-PAGE. Gels were assessed using Coomassie

staining.

Assessment of ClpX ATPase activity **in vitro.**

A 49.5 µL reaction containing 1.5 µg of recombinant wild-type ClpX or $ClpX_{(E187A)}$ in ATPase

assay buffer (see above) supplemented with 1.7% DMSO was preincubated for 10 minutes at

room temperature without ATP. Next, ATP dissolved in ATPase assay buffer was added to 1 µM

giving a final volume of 50 µL, and the reaction was incubated at 30□ C for 1.5 hours. After the

1.5 hours, the reactions were incubated for an additional 30 minutes at room temperature. 50 µL

of Kinase-Glo® reagent (Promega) was then added and incubated at room temperature for 10

minutes. Luminescence of the reaction, reflecting ATP not consumed by ClpX or $ClpX_{(E187A)}$,

was then measured using a BioTek Synergy H1 plate reader. Reactions were performed in

duplicate at least three times with at least two independent protein preparations.

Determining Protein-Protein Interactions with the BACTH System: The Bacterial Adenylate

Cyclase Two-Hybrid (BACTH) assay was utilized to test interaction between ClpX wild-type

and mutant (84). The genes of interest are translationally fused to one of either subunit, denoted

as T18 and T25, of the *B. pertussis* adenylate cyclase toxin, which can complement adenylate

cyclase deficient (Δ*cya*) DHT1 *E. coli*. Wild-type and mutant *clpX* genes cloned into one of the

pST25, pSNT25, or pUT18C Gateway® vectors was tested for both homotypic and heterotypic

interactions (9, 80). Plasmids from each background were co-transformed into chemically

competent DHT1 *E. coli*, which were plated on a double antibiotic minimal M63 medium

selection plate supplemented with 0.5 mM IPTG for induction of the protein, 40 μg/mL Xgal,

0.04% casein hydrolysate, and 0.2% maltose. Leucine zipper motifs were used for controls in

pKT25 and pUT18C backgrounds on the appropriate antibiotic selection plates because these

have been previously shown to interact (85). Blue colonies, indicative of positive interaction,

were screened using the β-galactosidase assay. Random positive colonies were selected and

grown in M63 minimal media with the appropriate antibiotics. 0.1% SDS and chloroform were

used to permeabilize the bacteria prior to addition of 0.1% o-nitrophenol-β-galactoside (ONPG).

1 M $NaHCO_3$ was used to stop the reaction after precisely 20 minutes of incubation at room

temperature. Absorbance at the 405 nm wavelength was recorded and normalized to bacterial

growth (OD_{600}), dilution factor, and time (in minutes) of incubation prior to stopping the

reaction. Totals were reported in relative units (RU) of β-galactosidase activity.

Chlamydial Transformation: The protocol followed was a modification of the method

developed by Mueller and Fields (86) and as previously described (17). For transformation, 10^6

C. trachomatis serovar L2 EBs (25667R) naturally lacking the endogenous plasmid were

incubated with 2 μg of unmethylated plasmid in a volume of 50 μL $CaCl_2$ at room temperature

for 30 minutes. Reaction volume was sufficient for one well of a six well plate of McCoy mouse

fibroblasts. Transformants were mixed with 1 mL of HBSS and added to 1 mL of HBSS in a six

well plate. The plates were centrifuged at room temperature for 15 minutes, 400 xg. The plate

was then incubated at 37° C for 15 minutes. After incubation, the HBSS was aspirated and

replaced with antibiotic-free DMEM+10% FBS. 8 hours post-infection, the media was replaced

with DMEM containing 1 μg/mL cycloheximide and 1 U/mL penicillin. Cells infected with

transformants were passaged every 48 hours until a population of penicillin resistant bacteria was

established. EBs were harvested and frozen in sucrose/phosphate (2SP; (64)) solution at -80° C.

Determining the Effect of Overexpression of Wild-Type and Mutant Clp Proteins via

Immunofluorescence and Inclusion Forming Unit Analysis: *C. trachomatis* transformants

containing plasmids encoding the 6xHis-tagged protein of interest were used to infect a confluent

monolayer of HEp2 cells. Penicillin treatment was maintained throughout the duration of the

infection. At 10 hpi, samples were induced or not with 10 nM anhydrotetracycline (aTc). At the

given timepoints, three wells of a 24 well plate were scraped in 2SP, vortexed with three 1 mm

glass beads, and frozen at -80° C. At the same timepoint, a coverslip was fixed in 3.25%

formaldehyde and 0.025% glutaraldehyde for two minutes, followed by permeabilization with

cold 90% methanol for one minute. Coverslips were labeled with primary goat anti-major outer

membrane protein (MOMP; Meridian, Cincinnati, OH), rabbit anti-6xHis (Abcam, Cambridge,

MA), and DAPI. Appropriate donkey secondary antibodies were used (Invitrogen, Carlsbad,

CA). Images were acquired on an Axio ImagerZ.2 equipped with Apotome.2 optical sectioning

hardware and X-Cite Series 120PC illumination lamp. Frozen IFU samples were titrated onto a

fresh monolayer of HEp2s without antibiotics. At 24 hpi, samples were fixed with methanol for

10 minutes, stained for MOMP, and enumerated.

Genomic DNA Isolation and qPCR Enumeration of Genomic Equivalents: At 24 or 48 hpi,

one well of a six well plate was scraped into the media overlay and pelleted at 17000 xg, 4° C for

15 minutes. Each sample was resuspended in 500 µL of cold PBS, frozen three times at -80° C,

and processed using the Qiagen DNeasy Blood and Tissue Kit according to the manufacturer's

specifications. DNA concentrations were assessed using a spectrophotometer prior to dilution

down to 5 ng/µL. 5 µL of the resulting dilution was used for a 25 µL qPCR reaction volume

using SYBR® Green PCR Master Mix (Applied Biosystems). Each reaction was performed in

triplicate. A standard curve using Ctr L2 genomic DNA was generated for interpolation of

sample Ct values. This experiment was performed three times for three biological replicates.

Analysis of HctB Levels Upon Clp Overexpression: At 24 or 48 hpi, one well of a six well plate

per test condition was rinsed twice with HBSS. To lyse the cells, 500 µL of denaturing lysis

buffer (8 M Urea, 10 mM Tris, 2.5% 2-mercaptoethanol, 1% SDS) was added to each well and

incubated for 15 minutes at room temperature. 300 units of Universal Nuclease (Pierce) per mL

of lysis buffer was added immediately prior to addition to the wells. Following incubation,

samples were centrifuged at 17000 xg, 4° C for 15 minutes to remove any insoluble material.

Samples were quantitated using the EZQ Protein Quantitation Kit (Pierce). 50 µg of each sample

was run in a 4-20% gradient SDS-PAGE gel (BioRad) and transferred to a PVDF 0.45 µm pore

size membrane for 1 h at 300 mA. The membrane was probed using goat anti-MOMP (Meridian)

and rabbit anti-HctB (generously provided by Dr. T. Hackstadt, NIH) primary antibodies

followed by staining with donkey anti-goat 680 and donkey anti-rabbit 800 (LI-COR) secondary

antibodies. The membrane was imaged on an Azure c600 imaging system. The channels were

gray-scaled and equally contrast corrected, and the resulting images were used for integrated

density measurement with FIJI software (87). To assess relative HctB levels, the HctB integrated

density of each sample was normalized to its respective MOMP integrated density to avoid bias

due to lower overall organism numbers. The ratios were then used to compare induced versus

uninduced relative HctB levels. These experiments were performed three times for a total of

three biological replicates.

Transmission electron microscopy (TEM) assessment of the effect of overexpression of mutant

Clp isoforms. Samples were infected and induced as previously discussed (see above). At 48 hpi,

samples were fixed using 2% Glutaraldehyde, 2% Formaldehyde in 0.1M Sorensen's phosphate

buffer, pH 7.2. Samples were then stained post-fixation in 1% Osmium Tetroxide in water for 1

hour. Samples were dehydrated in an Ethanol series 50%, 70%, 90%, 95%, 100% 3 changes of

100%, all steps 15 minutes each and were then soaked in Propylene Oxide 100% 3 changes for

15 minutes each. Samples were left overnight in a fume hood in a 1:1 mixture of Propylene

Oxide and Embed 812. The following day the samples were placed in molds with fresh Embed

812 and polymerized overnight in an oven set at 65° C. Blocks were thin sectioned 90

nanometers thick on a Leica UC6 Ultramicrotome using a Diatome diamond knife. Sections were

placed on uncoated 200 mesh copper grids and stained with 2% Uranyl Acetate and Reynold's

Lead Citrate. Sections were examined on a FEI Tecnai G2 TEM operated at 80Kv.

Confirmation of* clpP2X *and* incA *knockdown. Briefly, two wells of a six-well plate per

condition were infected with pBOMBLCRia-*clpP2X* transformed CtrL2 at an MOI of 0.8. At

either four or ten hpi, samples were or were not induced with 10 nM aTc. At each given

timepoint, total RNA was collected using Trizol reagent (Invitrogen) and was extracted with

chloroform as described previously (17, 83, 88-90). The aqueous layer was precipitated using

isopropanol, as per the manufacturer's instructions. Samples were DNase treated using the

TURBO DNA-free kit (Ambion), and 1 µg of the resulting RNA was reverse transcribed using

SuperScript III reverse transcriptase (Invitrogen). Equal volumes of cDNA were loaded for each

qPCR reaction. To extract genomic DNA, one well per condition was harvested and processed

using the DNeasy blood and tissue kit (Qiagen) according to the manufacturer's instructions as

noted above. Samples were diluted to 5 ng/µL, and 5 µL of the resulting dilution was used per

qPCR reaction. cDNA and gDNA samples were quantified using 25 µL reactions with 2x SYBR

PowerUP Green Master Mix (Invitrogen) analyzed on a QuantStudio 3 (Applied Biosystems)

thermal cycler using the standard cycling conditions. A standard curve using purified wild-type

CtrL2 genomic DNA was generated for sample quantification. Data are displayed as the ratio of

cDNA to gDNA normalized to the 10h uninduced sample. For *incA* knockdown, HEp2 cells

were infected with the pBOMBLCRia-*incA* transformant, induced with 10 nM aTc as above, and

fixed at 24hpi with methanol. Cells were labeled with primary guinea pig anti-major outer

membrane protein (MOMP; kind gift of Dr. E. Rucks, UNMC), rabbit anti-Sa_dCas9 (Abcam,

Cambridge, MA), sheep anti-IncA (Dr. E. Rucks), and DAPI. Appropriate donkey secondary

antibodies were used (Invitrogen). Coverslips were mounted on glass slides using ProLong Glass

Antifade mounting media (ThermoFisher). Images were acquired on a Zeiss Axio ImagerZ.2

equipped with Apotome.2 optical sectioning hardware and X-Cite Series 120PC illumination

lamp.

Determination of the effect of* clpP2X *knockdown on Ctr. 24-well plates of HEp2 cells were infected at an MOI of 0.8 with either pBOMBLCRia-*clpP2X* or pBOMBLCRia-*incA* transformed into CtrL2. Samples were induced or not at 4 hpi and were harvested, fixed, and titered as previously described. To monitor the expression of dCas9 and, if present, the complemented gene of interest, immunofluorescence samples on glass coverslips were fixed and permeabilized using ice cold 90% methanol at 24 or 48 hpi. Each sample was then stained using goat anti-MOMP (Meridian BioScience), rabbit anti-Sa_dCas9, and mouse anti-FLAG (Sigma-Aldrich) primary antibodies. Samples were then stained with the appropriate donkey secondary antibodies and were imaged as discussed above. Each titration was fixed using 4% formaldehyde and 0.025% glutaraldehyde to preserve GFP fluorescence. IFU counts of GFP positive inclusions are displayed as a percentage of the uninduced sample at the given timepoint. Plasmid retention for each condition is displayed as the percent of GFP positive to total number of inclusions for each condition.

GFP_ssrA stability assay. Chlamydial transformants harboring either pBOMBmC-GFP(VAA) or pBOMBmC-GFP(VDD) plasmids were used to infect a fresh monolayer of McCoy mouse fibroblasts on coverslips in a 24 well plate. Samples were induced or not at 8 hpi. At 16 hpi, all media was replaced with fresh DMEM with or without the indicated compound. GFP expression was either continued through 20 hpi or was removed from 16 to 20 hpi for pulse/chase samples. At 20 hpi, all samples were formaldehyde/glutaraldehyde fixed as previously discussed. Coverslips were then mounted and imaged as previously discussed using a 40x objective. GFP and mCherry integrated densities were measured for approximately 60 fields of view on two separate coverslips per condition. These experiments were repeated for two biological replicates.

To determine the ratio of GFP to mCherry, GFP integrated density measurements were divided

by those of mCherry.

Effect of ClpX-targeting compounds on chlamydial growth and host cell viability. Stocks of

ClpX-specific inhibitor 334 and its derivative, 365, were synthesized as previously reported (44),

resuspended at 25 mg/mL in DMSO, and frozen at -20°C. Methods for the synthesis,

purification, and analysis of these compounds is available in Supplementary Information. A dose

curve of treatment was performed to determine an inhibitory concentration of the compounds on

Ctr, and 25 µg/mL was chosen (data not shown). For the 24 and 48 h samples, 500 µL of DMEM

containing 25 µg/mL of the compounds were added at 8 hpi, and samples were harvested at the

indicated timepoint. For the time of infection samples, compounds were added at 15 minutes

post-infection and removed at 8 hpi. For the reactivation samples, DMEM containing the

respective compound was added at 8 hpi, washed out three times with HBSS at 24 hpi, and then

replaced with DMEM only for 24 additional hours prior to harvest. To determine the effect on

preformed EBs, compound was added at 24 hpi, and samples were harvested at 48 hpi. To

harvest, three wells of a 24-well plate were scraped into 2SP, vortexed with 3 mm glass beads,

and frozen at -80^0 C. Samples were titrated onto a fresh monolayer of HEp2 cells with no

treatment for enumeration. For assessment of cell viability, HEp2 cells were plate in 96 well dark

wall plate and were treated as described for IFU experiments. All wells were then mixed with

PrestoBlue HS reagent (Invitrogen), incubated for 10 minutes, and measured for fluorescence

using a Tecan plate reader. Values are given as a percentage of the untreated sample.

Acknowledgements

We thank Dr. H. Caldwell (NIH/NIAID) for eukaryotic cell lines, Dr. T. Hackstadt

(RML/NIAID) for providing antibodies and the pBOMB4-Tet::L2 plasmid, Dr. P. Scott Hefty

(KU) for the pTLR2-gfp::L2 plasmid, and Dr. Peter Sass (University of Tuebingen) for the

BL21(DE3) Δ*clpPAX E. coli* strain used in this research. We thank Dr. Lisa Rucks antibodies

used in this study and for critical review of the manuscript as well as members of the Ouellette

and Rucks labs for helpful discussions of the work. This project was supported by a National

Science Foundation CAREER award (1810599) and an NIAID/National Institutes of Health

award (R21AI141933-01) to SPO and by University funds to DJF. This project was also funded

by NIGMS/National Institutes of Health award to the Nebraska Center for Molecular Target

Discovery and Development (1P20GM121316□01A1, PI: Robert Lewis, Project Leader, MCS).

References:

1. CDC. 2017. 2017 Sexually Transmitted Diseases Surveillance. https://www.cdc.gov/std/stats16/chlamydia.htm#foot-1. Accessed
2. WHO. 2018. Trachoma. http://www.who.int/mediacentre/factsheets/fs382/en/. Accessed
3. AbdelRahman YM, Belland RJ. 2006. The chlamydial developmental cycle. FEMS Microbiology Reviews 29:949-959.
4. Hatch TP, Allan I, Pearce JH. 1984. Structural and polypeptide differences between envelopes of infective and reproductive life cycle forms of *Chlamydia* spp. Journal of Bacteriology 157:13-20.
5. Hackstadt T, Baehr W, Ying Y. 1991. *Chlamydia trachomatis* developmentally regulated protein is homologous to eukaryotic histone H1. Proceedings of the National Academy of Sciences of the United States of America 88:3937-3941.
6. Clifton DR, Fields KA, Grieshaber SS, Dooley CA, Fischer ER, Mead DJ, Carabeo RA, Hackstadt T. 2004. A chlamydial type III translocated protein is tyrosine-phosphorylated at the site of entry and associated with recruitment of actin. Proceedings of the National Academy of Sciences of the United States of America 101:10166.
7. Lane BJ, Mutchler C, Al Khodor S, Grieshaber SS, Carabeo RA. 2008. Chlamydial Entry Involves TARP Binding of Guanine Nucleotide Exchange Factors. PLoS Pathogens 4:e1000014.
8. Hodinka RL, Davis CH, Choong J, Wyrick PB. 1988. Ultrastructural study of endocytosis of *Chlamydia trachomatis* by McCoy cells. Infection and Immunity 56:1456-1463.
9. Moore ER, Ouellette SP. 2014. Reconceptualizing the chlamydial inclusion as a pathogen-specified parasitic organelle: an expanded role for Inc proteins. Frontiers in Cellular and Infection Microbiology 4:157.
10. Abdelrahman Y, Ouellette SP, Belland RJ, Cox JV. 2016. Polarized Cell Division of *Chlamydia trachomatis*. PLoS pathogens 12:e1005822-e1005822.
11. Saka HA, Thompson JW, Chen Y-S, Kumar Y, Dubois LG, Moseley MA, Valdivia RH. 2011. Quantitative proteomics reveals metabolic and pathogenic properties of Chlamydia trachomatis developmental forms. Molecular Microbiology 82:1185-1203.
12. Belland RJ, Zhong G, Crane DD, Hogan D, Sturdevant D, Sharma J, Beatty WL, Caldwell HD. 2003. Genomic transcriptional profiling of the developmental cycle of *Chlamydia trachomatis*. Proceedings of the National Academy of Sciences of the United States of America 100:8478-8483.
13. Vandahl BBS, Birkelund S, Christiansen G. 2004. Genome and proteome analysis of Chlamydia. PROTEOMICS 4:2831-2842.
14. Skipp PJS, Hughes C, McKenna T, Edwards R, Langridge J, Thomson NR, Clarke IN. 2016. Quantitative Proteomics of the Infectious and Replicative Forms of Chlamydia trachomatis. PLoS ONE 11:e0149011.
15. Higgins D, Dworkin J. 2012. Recent progress in Bacillus subtilis sporulation. FEMS microbiology reviews 36:131-148.
16. Hughes V, Jiang C, Brun Y. 2012. *Caulobacter crescentus*. Current Biology 22:R507-R509.

17. Wood NA, Chung K, Blocker A, Rodrigues de Almeida N, Conda-Sheridan M, Fisher
DJ, Ouellette SP. 2018. Initial Characterization of the Two ClpP Paralogs of *Chlamydia*
trachomatis Suggests Unique Functionality for Each. Journal of Bacteriology.
18. Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger
L, Tatusov RL, Zhao Q, Koonin EV, Davis RW. 1998. Genome Sequence of an Obligate
Intracellular Pathogen of Humans: *Chlamydia trachomatis*. Science 282:754.
19. Singh SK, Grimaud R, Hoskins JR, Wickner S, Maurizi MR. 2000. Unfolding and
internalization of proteins by the ATP-dependent proteases ClpXP and ClpAP.
Proceedings of the National Academy of Sciences of the United States of America
97:8898-8903.
20. Hwang BJ, Park WJ, Chung CH, Goldberg AL. 1987. *Escherichia coli* contains a soluble
ATP-dependent protease (Ti) distinct from protease La. Proceedings of the National
Academy of Sciences of the United States of America 84:5550-5554.
21. Walker JE, Saraste M, Runswick MJ, Gay NJ. 1982. Distantly related sequences in the
alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring
enzymes and a common nucleotide binding fold. The EMBO Journal 1:945-951.
22. Wendler P, Ciniawsky S, Kock M, Kube S. 2012. Structure and function of the AAA+
nucleotide binding pocket. Biochimica et Biophysica Acta (BBA) - Molecular Cell
Research 1823:2-14.
23. Baker TA, Sauer RT. 2012. ClpXP, an ATP-powered unfolding and protein-degradation
machine. Biochimica et biophysica acta 1823:15-28.
24. Wojtyra UA, Thibault G, Tuite A, Houry WA. 2003. The N-terminal Zinc Binding
Domain of ClpX Is a Dimerization Domain That Modulates the Chaperone Function.
Journal of Biological Chemistry 278:48981-48990.
25. Donaldson LW, Wojtyra U, Houry WA. 2003. Solution Structure of the Dimeric Zinc
Binding Domain of the Chaperone ClpX. Journal of Biological Chemistry 278:48991-
48996.
26. Erzberger JP, Berger JM. 2006. Evolutionary relationships and structural mechanisms of
AAA+ proteins. Annu Rev Biophys Biomol Struct 35:93-114.
27. Siddiqui SM, Sauer RT, Baker TA. 2004. Role of the processing pore of the ClpX AAA+
ATPase in the recognition and engagement of specific protein substrates. Genes &
development 18:369-374.
28. Martin A, Baker TA, Sauer RT. 2008. Diverse pore loops of the AAA+ ClpX machine
mediate unassisted and adaptor-dependent recognition of ssrA-tagged substrates.
Molecular cell 29:441-450.
29. Martin A, Baker TA, Sauer RT. 2008. Pore loops of the AAA+ ClpX machine grip
substrates to drive translocation and unfolding. Nature structural & molecular biology
15:1147-1151.
30. Iosefson O, Olivares AO, Baker TA, Sauer RT. 2015. Dissection of Axial-Pore Loop
Function during Unfolding and Translocation by a AAA+ Proteolytic Machine. Cell
reports 12:1032-1041.
31. Iosefson O, Nager AR, Baker TA, Sauer RT. 2015. Coordinated gripping of substrate by
subunits of a AAA+ proteolytic machine. Nature chemical biology 11:201-206.
32. Ogura T, Whiteheart SW, Wilkinson AJ. 2004. Conserved arginine residues implicated in
ATP hydrolysis, nucleotide-sensing, and inter-subunit interactions in AAA and AAA+
ATPases. Journal of Structural Biology 146:106-112.

33. Kim Y-I, Levchenko I, Fraczkowska K, Woodruff RV, Sauer RT, Baker TA. 2001.
Molecular determinants of complex formation between Clp/Hsp100 ATPases and the
ClpP peptidase. Nature Structural Biology 8:230.
34. Amor AJ, Schmitz KR, Baker TA, Sauer RT. 2019. Roles of the ClpX IGF loops in ClpP
association, dissociation, and protein degradation. Protein Science 28:756-765.
35. Brickman TJ, Barry CE, 3rd, Hackstadt T. 1993. Molecular cloning and expression of
hctB encoding a strain-variant chlamydial histone-like protein with DNA-binding
activity. Journal of bacteriology 175:4274-4281.
36. Barry Iii CE, Brickman TJ, Hackstadt T. 1993. Hc1-mediated effects on DNA structure: a
potential regulator of chlamydial development. Molecular Microbiology 9:273-283.
37. Fahr MJ, Douglas AL, Xia W, Hatch TP. 1995. Characterization of late gene promoters
of Chlamydia trachomatis. Journal of bacteriology 177:4252-4260.
38. Ouellette SP. 2018. Feasibility of a Conditional Knockout System for *Chlamydia* Based
on CRISPR Interference. Frontiers in Cellular and Infection Microbiology 8:59.
39. Ran FA, Cong L, Yan WX, Scott DA, Gootenberg JS, Kriz AJ, Zetsche B, Shalem O, Wu
X, Makarova KS, Koonin EV, Sharp PA, Zhang F. 2015. *In vivo* genome editing using
Staphylococcus aureus Cas9. Nature 520:186-191.
40. Suchland RJ, Rockey DD, Bannantine JP, Stamm WE. 2000. Isolates of *Chlamydia*
trachomatis That Occupy Nonfusogenic Inclusions Lack IncA, a Protein Localized to the
Inclusion Membrane. Infection and Immunity 68:360.
41. Wichlan DG, Hatch TP. 1993. Identification of an early-stage gene of *Chlamydia psittaci*
6BC. Journal of Bacteriology 175:2936-2942.
42. Shaw EI, Dooley CA, Fischer ER, Scidmore MA, Fields KA, Hackstadt T. 2002. Three
temporal classes of gene expression during the *Chlamydia trachomatis* developmental
cycle. Molecular Microbiology 37:913-925.
43. Johnson FWA, Hobson D. 1977. The effect of penicillin on genital strains of Chlamydia
trachomatis in tissue culture. Journal of Antimicrobial Chemotherapy 3:49-56.
44. Fetzer C, Korotkov VS, Thänert R, Lee KM, Neuenschwander M, von Kries JP, Medina
E, Sieber SA. 2017. A Chemical Disruptor of the ClpX Chaperone Complex Attenuates
the Virulence of Multidrug-Resistant Staphylococcus aureus. Angewandte Chemie
International Edition 56:15746-15750.
45. Trott O, Olson AJ. 2010. AutoDock Vina: improving the speed and accuracy of docking
with a new scoring function, efficient optimization, and multithreading. Journal of
computational chemistry 31:455-461.
46. Pan S, Malik IT, Thomy D, Henrichfreise B, Sass P. 2019. The functional ClpXP
protease of Chlamydia trachomatis requires distinct clpP genes from separate genetic
loci. Scientific Reports 9:14129.
47. Flynn JM, Levchenko I, Seidel M, Wickner SH, Sauer RT, Baker TA. 2001. Overlapping
recognition determinants within the ssrA degradation tag allow modulation of
proteolysis. Proceedings of the National Academy of Sciences of the United States of
America 98:10584-10589.
48. Leung E, Datti A, Cossette M, Goodreid J, McCaw Shannon E, Mah M, Nakhamchik A,
Ogata K, El Bakkouri M, Cheng Y-Q, Wodak Shoshana J, Eger Bryan T, Pai Emil F, Liu
J, Gray-Owen S, Batey Robert A, Houry Walid A. 2011. Activators of Cylindrical
Proteases as Antimicrobials: Identification and Development of Small Molecule
Activators of ClpP Protease. Chemistry & Biology 18:1167-1178.

49. Olivares AO, Baker TA, Sauer RT. 2016. Mechanistic insights into bacterial AAA+
proteases and protein-remodelling machines. Nature reviews Microbiology 14:33-44.
50. Martin A, Baker TA, Sauer RT. 2007. Distinct static and dynamic interactions control
ATPase-peptidase communication in a AAA+ protease. Molecular cell 27:41-52.
51. Singh SK, Rozycki J, Ortega J, Ishikawa T, Lo J, Steven AC, Maurizi MR. 2001.
Functional Domains of the ClpA and ClpX Molecular Chaperones Identified by Limited
Proteolysis and Deletion Analysis. Journal of Biological Chemistry 276:29420-29429.
52. Thomson NR, Holden MTG, Carder C, Lennard N, Lockey SJ, Marsh P, Skipp P,
O'Connor CD, Goodhead I, Norbertzcak H, Harris B, Ormond D, Rance R, Quail MA,
Parkhill J, Stephens RS, Clarke IN. 2008. Chlamydia trachomatis: genome sequence
analysis of lymphogranuloma venereum isolates. Genome research 18:161-171.
53. Moore SD, Sauer RT. 2007. The tmRNA System for Translational Surveillance and
Ribosome Rescue. Annual Review of Biochemistry 76:101-124.
54. Moore SD, Sauer RT. 2005. Ribosome rescue: tmRNA tagging activity and capacity in
Escherichia coli. Molecular Microbiology 58:456-466.
55. Hudson CM, Williams KP. 2014. The tmRNA website. Nucleic Acids Research
43:D138-D140.
56. Abo T, Inada T, Ogawa K, Aiba H. 2000. SsrA☐mediated tagging and proteolysis of
LacI and its role in the regulation of *lac* operon. The EMBO Journal 19:3762.
57. Dulebohn D, Choy J, Sundermeier T, Okan N, Karzai AW. 2007. Trans-Translation:☐
The tmRNA-Mediated Surveillance Mechanism for Ribosome Rescue, Directed Protein
Degradation, and Nonstop mRNA Decay. Biochemistry 46:4681-4693.
58. Marsh JW, Wee BA, Tyndall JDA, Lott WB, Bastidas RJ, Caldwell HD, Valdivia RH,
Kari L, Huston WM. 2015. A Chlamydia trachomatis strain with a chemically generated
amino acid substitution (P370L) in the cthtrA gene shows reduced elementary body
production. BMC microbiology 15:194-194.
59. Marsh JW, Lott WB, Tyndall JDA, Huston WWM. 2013. Proteolytic activation of
Chlamydia trachomatis HTRA is mediated by PDZ1 domain interactions with protease
domain loops L3 and LC and beta strand β5. Cellular & molecular biology letters 18:522-
537.
60. Huston WM, Tyndall JDA, Lott WB, Stansfield SH, Timms P. 2011. Unique residues
involved in activation of the multitasking protease/chaperone HtrA from Chlamydia
trachomatis. PloS one 6:e24547-e24547.
61. Matsumoto A, Manire GP. 1970. Electron Microscopic Observations on the Effects of
Penicillin on the Morphology of Chlamydia psittaci. Journal of Bacteriology 101:278-
285.
62. Skilton RJ, Cutcliffen LT, Barlow D, Wang Y, Salim O, Lambden PR, Clarke IN. 2009.
Penicillin induced persistence in *Chlamydia trachomatis*: high quality time lapse video
analysis of the developmental cycle. PloS one 4:e7723-e7723.
63. Ouellette SP, Karimova G, Subtil A, Ladant D. 2012. *Chlamydia* co-opts the rod shape-
determining proteins MreB and Pbp2 for cell division. Molecular Microbiology 85:164-
178.
64. Wang Y, Kahane S, Cutcliffe LT, Skilton RJ, Lambden PR, Clarke IN. 2011.
Development of a Transformation System for *Chlamydia trachomatis*: Restoration of
Glycogen Biosynthesis by Acquisition of a Plasmid Shuttle Vector. PLoS Pathogens
7:e1002258.

65.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a
reference resource for gene and protein annotation. Nucleic Acids Research 44:D457-
D462.
66.   Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic
Acids Research 28:27-30.
67.   Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. 2017. KEGG: new
perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Research
45:D353-D361.
68.   Altschul SF, Wootton JC, Gertz EM, Agarwala R, Morgulis A, Schäffer AA, Yu Y-K.
2005. Protein Database Searches Using Compositionally Adjusted Substitution Matrices.
The FEBS journal 272:5101-5109.
69.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Molecular
Systems Biology 7:539-539.
70.   Troshin PV, Procter JB, Barton GJ. 2011. Java bioinformatics analysis web services for
multiple sequence alignment—JABAWS:MSA. Bioinformatics 27:2001-2002.
71.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web
portal for protein modelling, prediction and analysis. Nature protocols 10:845-858.
72.   Waterhouse A, Rempfer C, Heer FT, Studer G, Tauriello G, Bordoli L, Bertoni M,
Gumienny R, Lepore R, Bienert S, de Beer TA P, Schwede T. 2018. SWISS-MODEL:
homology modelling of protein structures and complexes. Nucleic Acids Research
46:W296-W303.
73.   Waterhouse A, Studer G, Tauriello G, Bordoli L, Bienert S, de Beer Tjaart AP, Schwede
T. 2016. The SWISS-MODEL Repository—new features and functionality. Nucleic
Acids Research 45:D313-D319.
74.   Guex N, Peitsch MC, Schwede T. 2009. Automated comparative protein structure
modeling with SWISS-MODEL and Swiss-PdbViewer: A historical perspective.
ELECTROPHORESIS 30:S162-S173.
75.   Bertoni M, Kiefer F, Biasini M, Bordoli L, Schwede T. 2017. Modeling protein
quaternary structure of homo- and hetero-oligomers beyond binary interactions by
homology. Scientific Reports 7:10480.
76.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.
2004. UCSF Chimera—A visualization system for exploratory research and analysis.
Journal of Computational Chemistry 25:1605-1612.
77.   Dunbrack RL. 2002. Rotamer Libraries in the 21st Century. Current Opinion in Structural
Biology 12:431-440.
78.   Dunbrack Jr RL, Cohen FE. 1997. Bayesian statistical analysis of protein side-chain
rotamer preferences. Protein Science 6:1661-1681.
79.   Wang J, Wang W, Kollman PA, Case DA. 2006. Automatic atom type and bond type
perception in molecular mechanical calculations. Journal of Molecular Graphics and
Modelling 25:247-260.
80.   Ouellette SP, Gauliard E, Antosová Z, Ladant D. 2014. A Gateway®☐compatible
bacterial adenylate cyclase☐based two☐hybrid system. Environmental Microbiology
Reports 6:259-267.

81. Wickstrum J, Sammons LR, Restivo KN, Hefty PS. 2013. Conditional Gene Expression
in *Chlamydia trachomatis* Using the Tet System. PLoS ONE 8:e76743.
82. Bauler LD, Hackstadt T. 2014. Expression and targeting of secreted proteins from
Chlamydia trachomatis. Journal of bacteriology 196:1325-1334.
83. Ouellette SP, Hatch TP, AbdelRahman YM, Rose LA, Belland RJ, Byrne GI. 2006.
Global transcriptional upregulation in the absence of increased translation in Chlamydia
during IFNγ☐mediated host cell tryptophan starvation. Molecular Microbiology 62:1387-
1401.
84. Karimova G, Pidoux J, Ullmann A, Ladant D. 1998. A bacterial two-hybrid system based
on a reconstituted signal transduction pathway. Proceedings of the National Academy of
Sciences of the United States of America 95:5752-5756.
85. Robichon C, Karimova G, Beckwith J, Ladant D. 2011. Role of Leucine Zipper Motifs in
Association of the *Escherichia coli* Cell Division Proteins FtsL and FtsB. Journal of
Bacteriology 193:4988-4992.
86. Mueller KE, Wolf K, Fields KA. 2017. Chlamydia trachomatis Transformation and
Allelic Exchange Mutagenesis. Current Protocols in Microbiology 45:11A.3.1-11A.3.15.
87. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. Nature methods 9:676-682.
88. Ouellette SP, Abdelrahman YM, Belland RJ, Byrne GI. 2005. The Chlamydia
pneumoniae type III secretion-related lcrH gene clusters are developmentally expressed
operons. Journal of bacteriology 187:7853-7856.
89. Ouellette SP, Rueden KJ, Gauliard E, Persons L, de Boer PA, Ladant D. 2014. Analysis
of MreB interactors in Chlamydia reveals a RodZ homolog but fails to detect an
interaction with MraY. Frontiers in Microbiology 5:279.
90. Ouellette SP, Rueden KJ, Rucks EA. 2016. Tryptophan Codon-Dependent Transcription
in Chlamydia pneumoniae during Gamma Interferon-Mediated Tryptophan Limitation.
Infection and Immunity 84:2703-2713.
978
979

Figure Legends

Figure 1: Bioinformatic analysis of chlamydial ClpX supports its role as a AAA+ ATPase.

(A) Multiple sequence alignment of chlamydial ClpX with the ClpX orthologs of various other

bacteria. Ec = *Escherichia coli*, Ctr = *Chlamydia trachomatis*, Bs = *Bacillus subtilis*, Mtb =

Mycobacterium tuberculosis, Pa = *Pseudomonas aeruginosa*, Sa = *Staphylococcus aureus*.

Alignment was performed using Clustal Omega with default settings and presented using Jalview

version 2. Alignment was colored by % identity in shades of blue or as indicated below the

alignment. (**B) 3D model** of ClpX was generated using SwissModel and presented in UCSF

Chimera. Conserved motifs colored as above in (A) except the IGF loops, which are colored lime

green. Two subunits of the hexamer were hidden for easier visualization into the complex. Top

and bottom are representations following a 90° rotation either clockwise or counterclockwise

around the X axis. Of note, this model was generated using an ADP-bound form of ClpX as a

template.

Figure 2: Ctr ClpX is a functional ATPase that forms the expected hexamer. (A) Native-

PAGE assay of recombinant ClpX and ClpX$_{(E187A)}$. Expected hexameric size is approximately

283 kDa. **(B) ATP hydrolysis** end point assay using Kinase-Glo®. Relative luminescence units

(RLU) are displayed on the Y axis. Error bars represent standard deviation, and differences

between samples are significant (**** = $p < 0.0001$, one-way ANOVA). **(C) Bacterial**

Adenylate Cyclase Two Hybrid (BACTH) assays showing pairwise, homotypic interaction of

ClpX and ClpX$_{(E187A)}$ as well as heterotypic interaction of ClpX and ClpX$_{(E187A)}$. **(D) β-**

Galactosidase activity of the BACTH interactions from (C), displayed in arbitrary units on the

Y axis. All conditions $p < 0.05$ by unpaired t test compared to negative control.

1003

Figure 3: Overexpression of Clp mutant isoforms negatively impacts Ctr at later

timepoints. (A) IFA of ClpP2 and ClpX wild-type and mutant overexpression at 24 and 48 hpi.

Samples were induced with 10 nM aTc at 10 hpi and stained for MOMP (green), 6xHis (red),

and DNA (blue). Scale bar = 10 μm. Images were acquired on a Zeiss Apotome at 100x

magnification. **(B-C) One step growth curves** of wild-type and mutant ClpP2 (B) or ClpX (C).

Samples induced with 10 nM aTc at 10 hpi. IFUs recovered displayed as Log_{10}. Values represent

the average of two independent experiments. **(D) IFA of ClpP2X** wild-type or mutant operons at

24 and 48 hpi. Samples stained for MOMP (pink), FLAG (ClpP2, red), 6xHis (ClpX, green), and

DNA (blue). Parameters as described in (B). **(E) One step growth curves** of ClpP2X wild-type

and mutant overexpression. IFUs recovered displayed as Log_{10}.

1014

Figure 4: Functional disruption of ClpX and ClpP2 perturbs chlamydial development but

in different manners. (A) Fold changes of detectable gDNA from 24 to 48 hpi for each strain.

Samples were induced at 10 hpi with 10 nM aTc. * = $p \leq 0.05$, ** = $p \leq 0.001$ by paired t-test.

Values displayed as the average of three independent experiments with error bars representing

standard deviation. **(B) Western blot** analysis of HctB levels at 48 hpi with and without

overexpression. One well of a six well plate was lysed into 500 μL of denaturing lysis buffer. 50

μg of protein from the clarified lysate for each sample was assessed. Blots were probed for

MOMP (IR680) and HctB (IR800). Grayscale blot shown is representative of three independent

experiments. **(C) Quantified integrated** density of the staining from (B). HctB levels were

normalized to MOMP levels in each sample to account for differences in bacteria in each sample.

Values displayed as levels of induced to uninduced HctB/MOMP ratios for each strain. * = $p \leq$

0.05, ** = p ≤ 0.001, ns = not significant by multiple comparisons t-test. **(D-E) Representative**

images of electron micrographs of the indicated strains and conditions. White triangles indicate

normal RBs, black triangles indicate normal EBs, and gray triangles indicate abnormal

developmental forms.

1030

Figure 5: Knockdown of ClpP2X negatively impacts *Chlamydia*, while complementation

with ClpP2 partially restores normal development. (A-B) Representative

immunofluorescence images of *clpP2X* knockdown with either (A) no complementation or (B)

clpP2 complementation. **(C) Transcript levels** upon knockdown of *clpP2X*. Data displayed are

the values of three independent biological replicates of triplicate RT-qPCR reactions. Values are

normalized to the 10h uninduced ClpP2 value for each experiment. Samples were induced using

10 nM aTc. **(D) IFU titer** following induction of knockdown at 4 hpi. Values are presented on a

Log_{10} scale percent of the respective uninduced titer. **(E) Plasmid retention** based on the ratio of

GFP positive to total number of inclusions is displayed in percent for each condition. **(F-H) IFU**

titer and plasmid retention of *clpP2X* knockdown with *clpP2* complementation. Conditions

used are the same as in (C-E).

1042

Figure 6: Chemical disruption of ClpX function protects SsrA-tagged GFP. (A) The ratio of

integrated density measurements of GFP to mCherry for SsrA(VAA)-tagged GFP. All samples

were induced or not at 8 hpi with 20 nM aTc. Media was replaced with or without drug at 16 hpi,

and aTc induction was maintained or removed (for pulse/chase samples). Samples were fixed at

20 hpi, and GFP and mCherry intensity were quantified using Fiji. **** = p ≤ 0.0001, ns = not

significant b**y** ordinary one-way ANOVA. **(B) Same conditions** as in (A) but with GFP(VDD).

1049

Figure 7: Proposed model for ClpP2X function in *Chlamydia*. An RB ultimately has two

fates: differentiation or replication. Based on our data showing the impact of ClpP2 disruption on

the developmental cycle, we hypothesize that ClpP2 may function in triggering either event

based on its degradative target at either timepoint. Furthermore, we posit that ClpX serves an

important function in EB organization, given that ClpX affects recoverable IFUs without

reducing the amount of HctB produced.

1056

Supplemental Figure Legends

Figure S1: Transmission electron microscopy of the effects of overexpression of mutant Clp

isoforms on chlamydial morphology. (A) Representative ClpP2$_{S98A}$ uninduced or induced

samples at 48 hpi. Samples were induced or not with 10 nM aTc at 10 hpi and were fixed and

processed at 48 hpi. Arrows indicate abnormal forms in the induced samples. Scale bar = 2 μm.

(B) Representative ClpX$_{E187A}$ uninduced or induced samples at 48 hpi. Samples treated, fixed,

and processed as previously discussed. Arrows indicate abnormal forms with intrabacterial

aggregates. Scale bar = 2 μm.

1065

1066 **Figure S2: (A) RT-qPCR of *clpP1*, *euo*, and *omcB* upon *clpP2X knockdown*.** Data shown are

1067 the average of three biological replicates, each with three technical replicates. Values are

1068 normalized to the 10h timepoint for each respective gene. Error bars represent standard

1069 deviation. Samples induced as indicated with 10 nM aTc. **(B) RT-qPCR of** *clpP2X* knockdown

1070 with *pcnB2* complementation. Values normalized to the ClpP2 10h samples.

1071

1072 **Figure S3: (A) Immunofluorescence staining** to confirm knockdown of IncA upon induction of

1073 dCas9 expression. Sa_dCas9 was induced or not at 4 hpi. Samples were harvested at 24 hpi and

1074 were stained for chlamydial MOMP (green), IncA (magenta), and DNA (Blue). Scale bar = 10

1075 μm. **(B-C) Recoverable IFUs** and plasmid retention upon *incA* knockdown. **(D-E) Recoverable**

1076 **IFUs** and plasmid retention of *clpP2X* knockdown with *pcnB2* complementation.

1077

Figure S4: Representative immunofluorescence images of **(A)** *clpP2X* knockdown with *pcnB2*

complementation. Samples were induced as in IFU experiments. Chlamydial MOMP in green,

FLAG (complemented gene) in red, dCas9 in magenta, and DNA in blue. **(B) Transcript levels**

upon knockdown of *clpP2X* with *pcnB2* complementation. Data displayed are the values of three

independent biological replicates of triplicate RT-qPCR reactions. Values are normalized to the

10h uninduced ClpP2 value for each experiment. Samples were induced using 10 nM aTc. **(C)**

IFU titer following induction of knockdown and complementation at 4 hpi. Values are presented

on a Log_{10} scale percent of the respective uninduced titer. **(D) Plasmid retention** of *clpP2X*

knockdown with *pcnB2* complementation.

1087

Figure S5: Docking simulation of the ClpX inhibitor 334 on a ClpX model. (A) PDB

structure of 334. Inset is the 2D structure of the drug. **(B) Ribbon model** of docked 334 within

the ClpX hexamer. The best scoring model is shown. Only the two ClpX subunits making

contact with the model are shown (A in gray, B in seafoam green). The Walker A motif (red),

Walker B motif (purple), sensor 1 motif (dark green), and sensor 2 motif (orange) of subunit A

are colored for visualization. The arginine finger is labeled light green of subunit B. The center

picture is oriented as outside of the complex looking inward, and the other images are rotations

as indicated. **(C) Surface rendering** of the ClpX subunits with docked 334 are shown with

coloration as in (B).

1097

Figure S6: Docking simulation of the ClpX inhibitor 365 on a ClpX model. (A) PDB

structure of 365. Inset is the 2D structure of the drug. **(B) Ribbon model** of docked 365 within

the ClpX hexamer. The best scoring model is shown. Only the two ClpX subunits making

contact with the model are shown (A in gray, B in seafoam green). The Walker A motif (red),

Walker B motif (purple), sensor 1 motif (dark green), and sensor 2 motif (orange) of subunit A

are colored for visualization. The arginine finger is labeled light green of subunit B. The center

picture is oriented as outside of the complex looking inward, and the other images are rotations

as indicated. **(C) Surface rendering** of the ClpX subunits with docked 365 are shown with

coloration as in (B).

1107

Figure S7: Chemical disruption of ClpX function blocks degradation of an SsrA-tagged

substrate in *C. trachomatis* **L2.** Representative immunofluorescence images corresponding to

data presented in Figure 6 of the main text. Representative images are of one field of view of one

biological replicate. All samples were induced or not at 12 hpi with 20 nM aTc. Media was

replaced with or without drug at 16 hpi, and aTc induction was maintained or removed (for

pulse/chase samples). Samples were fixed at 20 hpi and were imaged as previously described

using the same exposure time for all samples. Scale bars = 20 μm. Individual color channels

were grayscaled and inverted to give clear distinction of fluorescent protein intensity.

1116

Figure S8: Chemical disruption of ClpX function is highly detrimental to *Chlamydia*. (A)

Immunofluorescence assay (IFA) of 24 hours post-infection (hpi). All drugs added at a final

concentration of 25 µg/mL. Drugs were added at 8 hpi for the 24 h treatment samples. Drugs

were added 15 minutes post-infection and removed at 8 hpi for the TOI: 8 h pulse samples.

MOMP is stained in green, and DNA is stained with DAPI. Images were acquired on a Zeiss

LSM800 microscope at 63x magnification. Scale bar = 10 µm. TOI = Time of Infection. **(B) IFA**

of 48 hpi samples. Samples stained for MOMP (green) and DNA (blue). Scale bar = 10 µm.

Drug was added at 8 hpi for 48 h and reactivation samples. Media removed, drug washed out,

and media with drug added back to +Drug at 24 h and 48 h samples. Reactivation samples media

replaced with DMEM, no drug. **(C) Recoverable inclusion forming units (IFUs)** from the

indicated conditions. Totals present as Log_{10} IFUs recovered. Standard deviation displayed on

graphs as error bars. **(D) PrestoBlue cell viability** assay upon drug treatment from 8 to 24 hpi.

Values displayed as a % of the untreated samples.

1130

1131　**Figure S9: Example ClpX protein purifications.** Recombinant, 6x His-tagged

1132　$ClpX/ClpX_{(E187A)}$ were purified using cobalt-based immobilized metal affinity chromatography.

1133　Samples were quantified and 1 and 5 µg aliquots were run on 10% SDS-PAGE followed by

1134　staining with Coomassie brilliant blue. Three ClpX and two $ClpX_{(E187A)}$ purification were

1135　performed using BL21(DE3) Δ*clpPAX E. coli*.

1136

1137　**Supplemental Table 1.** List of primers, plasmids, and strains used in this study.

Fig. 1

Fig. 2

Fig. 3

A. gDNA Fold Changes

B. HctB Levels

C.

| ClpP2 | ClpP2mut | ClpX | ClpXmut | ClpP2X | ClpP2Xmut |

MOMP (all Ctr)

HctB (EBs Only)

D. ClpP2mut Uninduced | ClpP2mut Induced

E. ClpXmut Uninduced | ClpXmut Induced

Fig. 4

Fig. 5

A.

B.

Fig. 6

Fig. S9