Evaluating the consequences of common assumptions in run

reconstructions on Pacific-salmon biological status assessments

Stephanie J. Peacock[1,*], Eric Hertz[2], Carrie A. Holt[3],

Brendan Connors[4], Cameron Freshwater[5], and Katrina Connors[6]

1. 8625 33 Ave NW, Calgary, AB T3B 1M2. Email: stephanie.j.peacock@gmail.com

2. Pacific Salmon Foundation, 300 – 1682 West 7th Avenue Vancouver, BC V6J 4S6. Email:

ehertz@psf.ca

3. Fisheries and Oceans Canada, Pacific Biological Station, 3190 Hammond Bay Road,

Nanaimo, BC V9T 6N7. Email: Carrie.Holt@dfo-mpo.gc.ca

4. Fisheries and Oceans Canada, Institute of Ocean Sciences, 9860 West Saanich Road,

Sidney, BC V8L 4B2. Email: Brendan.Connors@dfo-mpo.gc.ca

5. Fisheries and Oceans Canada, Pacific Biological Station, 3190 Hammond Bay Road,

Nanaimo, BC V9T 6N7. Email: Cameron.Freshwater@dfo-mpo.gc.ca

6. Pacific Salmon Foundation, 300 – 1682 West 7th Avenue Vancouver, BC V6J 4S6. Email:

kconnors@psf.ca

* Corresponding author. Phone: (778) 266-0575

Abstract

Information on biological status is essential for designing, implementing, and evaluating

management strategies and recovery plans for threatened or exploited species. However, the

data required to quantify status are often limited, and it is important to understand how

assessments of status may be biased by assumptions in data analysis. For Pacific salmon,

biological status assessments based on spawner abundances and spawner-recruitment (SR)

analyses often involve "run reconstructions" that impute missing spawner data, expand

observed spawner abundance to account for unmonitored streams, assign catch to individual

stocks, and quantify age-at-return. Using a stochastic simulation approach, we quantified how

common assumptions in run reconstructions biased assessments of biological status based on

spawner abundance. We found that status assessments were robust to most common

assumptions in run reconstructions, even in the face of declining monitoring coverage, but that

overestimating catch tended to increase rates of status misclassification. Our results lend

confidence to biological status assessments based on spawner abundances and SR analyses,

even in the face of incomplete data.

Introduction

Timely and effective management of fish and wildlife relies on accurate information

about the current biological status of populations. Status assessments form the basis of many

conservation decision-support tools, though they have limited influence without clear and

binding linkages to policy and decision making. Methods to determine the allocation of

resources that maximize biodiversity conservation, such as Priority Threat Management (Martin

et al. 2018; Carwardine et al. 2019; Walsh et al. 2020), are most effective when baseline

information on status can help identify which threatened populations are at highest risk of

extirpation without management intervention, and which are most likely to benefit from

recovery planning given limited resources. Metrics of biological status may also be useful for

defining recovery goals and as performance measures when evaluating the effectiveness of

different management strategies (e.g., Walsh et al. 2020). Knowledge of current biological

status is therefore needed for both implementing and evaluating management strategies and

recovery plans.

Depsite the recognized need for information on biological status, complete data on

abundance, trends, and demographic parameters are rarely available for populations or species

that need to be assessed (IUCN 2017). Improving the analytical approaches used to quantify

status is one way to reduce assumptions and limit the bias in status outcomes from low quality

or quantity data (e.g., Fleischman et al. 2013; Staton et al. 2020), but the methods widely

applied in assessing fish and wildlife can be slow to change (Peterman 2018) and analytical

resources are sometimes not available to implement new advanced approaches on all

populations or species of interest. Thus, evaluating the impact of assumptions in the

approaches that are widely applied is critical to understanding the potential biases, uncertainty,

and limitations of status assessments (Chen et al. 2003; Wetzel and Punt 2011).

Closed-loop simulation models are useful tools for evaluating the potential biases in

status assessments due to various ecological and management uncertainties (Walters 1986).

The major advantages of simulation approaches (over, for example, retrospective analyses) are

that the model can be tailored to different biological and management scenarios, allowing

management strategy evaluation (e.g., Punt 1992; Holt and Peterman 2006, 2008), and the true

status of the system is known perfectly, allowing quantification of bias in estimated parameters

(e.g., Pyper and Peterman 1998; Dorner et al. 2009) and estimated status (e.g., Peacock and

Holt 2012; Holt and Folkes 2015; Holt et al. 2018). Simulation approaches are extremely

flexible and enable testing common assumptions that underpin status assessments at broad

spatial scales in the face imperfect information, and have been applied extensively in fisheries

research. In this study, we apply a simulation approach to understanding potential biases in

biological status assessments of Pacific salmon introduced by assumptions in the analysis of

spawner and recruitment data.

Assessing biological status of Pacific salmon

Pacific salmon (*Oncorhynchus* spp.) are a highly exploited group of species and many

populations have experienced declines in recent decades due to overfishing, changing ocean

conditions, and freshwater habitat degradation (e.g., COSEWIC 2016, 2017; Brown et al. 2019).

Pacific salmon are anadromous and semelparous, returning from ocean rearing grounds to

spawn in freshwater before dying, and are typically vulnerable to fisheries upon their return to

coastal waters. The data required to assess biological status of Pacific salmon can include

annual estimates of the number of returning adult salmon to individual rivers, fisheries catch or

harvest rates, and the age composition of returning salmon. Often, these data are incomplete

and require imputation. "Run reconstructions" have been undertaken for many salmon stocks

to expand spawner abundances to account for unmonitored streams and estimate the

abundance of recruits to coastal fisheries prior to spawning (i.e., recruitment; Cave and Gazey

1994; English et al. 2007, 2016, 2018). The exact procedure undertaken depends on the life-

history traits and available data for each population, and can include complexities such as

spatial and temporal variability in returns among spawning populations. Status asssessments

that rely on run reconstructions have been adopted by, for example, local management

organisations, the Marine Stewardship Council (www.msc.org), the Pacific Salmon Treaty (PSC

2019), COSEWIC (e.g., COSEWIC 2016, 2017), and the Pacific Salmon Foundation (Connors et al.

2013, 2018, 2019).

Fisheries and Oceans Canada (DFO) also relies on run reconstructions when assessing

the biological status of Conservation Units (CUs) – groups of wild salmon that, if lost, are

unlikely to recolonize within an acceptable timeframe – under Canada's Wild Salmon Policy

(WSP; Fisheries and Oceans Canada 2005). In their "integrated status assessments", DFO

engages experts to consider the unique context of each CU and the quality and quantity of data

that may be used to estimate status (e.g., DFO 2015, 2016, 2018a). However, the required

resources and time (typically 1-3 years) have meant that integrated status asssssments have

been completed for only 9% of CUs since the WSP was released nearly 15 years ago, and

reports are often 2-4 years out of date when they are released (DFO 2019). Data-driven

approaches focused on calculating quantitative metrics without expert elicitation can provide

timely status updates that are standardized and comparable across CUs using transparent and

repeatable methodology. For example, the Pacific Salmon Foundation (PSF) has undertaken a

widespread effort to apply a data-driven approach to assessing spawner abundance under the

WSP framework, with results for the north and central coast openly available through their

Pacific Salmon Explorer (PSE) – www.salmonexplorer.ca (Connors et al. 2013, 2018, 2019), and

are currently expanding their assessments to all remaining salmon CUs in British Columbia (BC).

Despite the benefits of this approach, the impact of compound assumptions in run

reconstructions (Table 1) on status outcomes needs to be quantified to understand potential

biases and lend confidence and credibility to data-driven assessments.

In this study, we quantifed bias in biological status outcomes for Pacific salmon from

data-driven assessments due to assumptions and uncertainty in run reconstructions (Table 1).

We used a simulation approach that allowed us to explore how bias changed under different

biological and management conditions. As a case study, we tailored our simulation model to

represent a generic chum salmon CU from the central coast of BC because these CUs have

recently been assessed using a data-driven approach (Connors et al. 2018) and have a relatively

simple run-reconstruction model that does not include run timing. Furthermore, there are

conservation concerns for both north and central coast chum salmon, which have not

recovered despite significant reductions in harvest rates over the past two decades (DFO

2018b). Thus, central coast chum salmon offer a useful case study for an initial investigation of

basic assumptions underlying biological status assessments. However, our simulation model is

flexible enough to accommodate different species and life-history traits of Pacific salmon. We

further explore a broad range of biological (e.g., trends in capacity) and management (e.g.,

monitoring coverage) scenarios to yield more general insight into the circumstances under

which assumptions may bias status assessments.

Methods

Benchmark calculations and assumptions

Under the WSP's biological status assessment framework, quantifiable metrics are

calculated from available data and compared against biological reference points, or

"benchmarks", to arrive at a status outcome of red, amber, or green (Fisheries and Oceans

Canada 2005; Figure 1). A red status indicates that a CU has low spawner abundance and/or

reduced spatial distribution and management intervention is required to avoid extirpation. A

green status indicates that the CU is able to sustain maximum annual catch under existing

environmental conditions. The specific benchmarks delineating these status zones must

consider uncertainties in metrics and the unique biological characteristics (e.g., age

composition) of the CU being assessed.

Multiple metrics of biological status have been proposed under the WSP that cover four

broad classes of indicators: current spawner abundance, trends in abundance over time, spatial

distribution of spawners, and fishing mortality (Holt et al. 2009). We focused our analysis only

on the current spawner abundance metric and considered two types of benchmarks (Figure 1,2)

that have been frequently applied to determine biological status of Pacific salmon CUs,

including in the Pacific Salmon Explorer (PSE; Connors et al. 2013, 2018, 2019). The first type of

benchmark is associated with maximum sustainable yield, derived from a SR relationship

(Figure 2b). An upper SR benchmark of 80% of the spawner abundance that is projected to

maintain long-term maximum yield, or 80% S_{MSY}, has been recommended by Holt et al. (2009,

2018) and will be applied to future assessments in the PSE (previous PSE assessments have

applied S_{MSY}; Connors et al. 2018, 2019). S_{MSY} can be calculated explicitly from the productivity

and density-dependence parameters of the Ricker SR relationship (Scheuerell 2016). Multiple

lower benchmarks have been suggested (Holt et al. 2009, 2018), and we used a lower

benchmark of the spawner abundance that leads to S_{MSY} in one generation in the absence of

fishing mortality, called S_{GEN} (Korman and English 2013; DFO 2015), as applied in the PSE. These

benchmarks delinate the green, amber and red status zones: If the geometric mean spawner

abundance over the most recent generation (S_{AVG}) is above the upper benchmark of 80% S_{MSY},

then the CU is assigned a green status, if S_{AVG} is between 80% S_{MSY} and S_{GEN} , then the CU is

assigned an amber status, and if S_{AVG} is less than S_{GEN}, then the CU is assigned a red status.

The second type of benchmarks we considered were percentile benchmarks (Clark et al.

2014; Holt and Folkes 2015; Holt et al. 2018), also called historic spawners benchmarks (e.g.,

Connors et al. 2018). The upper and lower benchmarks are the 50[th] and 25[th] percentiles of

historical spawner abundance, respectively (Figure 2a). The calculation of these benchmarks

has fewer data requirements, as the SR relationship need not be estimated (i.e. there is no

need for age-at-return or harvest data). As such, the percentile benchmarks can be applied to

data-limited CUs for which spawner data are patchy or age and harvest data are not available.

We focused on a basic run-reconstruction model and associated assumptions that are

commonly made when assessing spawner abundance against the benchmarks above (Figure 2c;

Table 1). At a minimum, in order to apply percentile benchmarks, a time series of total spawner

abundance at the CU scale is required. Conservation Units are typically comprised of multiple

spawning populations that may or may not be monitored in any given year. Spawning

populations in individual streams (henceforth "populations") may exhibit unique dynamics as

their productivity is (in part) limited by density-dependent processes in freshwater. A simple

sum of the observed spawner abundance within a CU may be misleading if the same

populations are not monitored consistently. On BC's north and central coast, monitored

populations have been designated as either "indicator streams" or "non-indicator streams",

with indicator streams being prioritized for monitoring and thus having more continuous and

reliable spawner estimates (English 2016). In addition, there may be populations that have

never been monitored and for which spawner abundance is unknown. To reconstruct spawner

abundance to the CU, three "Expansion Factors" have been applied to account for (1) spawners

returning to indicator streams that are not monitored in a given year, (2) spawners returning to

non-indicator streams, and (3) observation efficiency and populations that are never monitored

(Table 1).

The application of SR benchmarks also requires time series of the total number of

salmon returning to the CU to reconstruct recruitment, including those caught in fisheries and

those that make it to spawn but are not counted. The number of returning salmon in a CU that

are caught in fisheries is estimated based on the catch statistics for Pacific Fisheries

Management Areas (PFMAs) adjacent to the geographic location of the CU (Figure 3). It is

assumed that salmon caught in a PFMA were destined to spawn in streams that empty into that

PFMA, although there is the potential for bias in that fish may be caught while migrating

through the PFMA or fish destined for streams in the focal PFMA may be caught in other

PFMAs. Furthermore, in most cases, there is not a perfect spatial correspondence between

PFMAs and CUs (Figure 3). Streams in multiple CUs may flow into a single PFMA, which is

common for small CUs, such as with sockeye salmon which spawn in individual lakes, each of

which is associated with a unique CU (Holtby and Ciruna 2007). In the simplest case, the catch

from that PFMA may be assigned to CUs based on the relative spawner abundance to each CU.

However, differences in run-timing among CUs may complicate the assignment of catch and

necessitate more complex run-reconstruction models. A single CU may also be comprised of

populations that are caught in multiple PFMAs, particularly for species with large CUs such as

pink and chum salmon whose geographic boundary often spans hundreds of kilometers and

includes dozens of spawning populations (Figure 3). In these cases, an average harvest rate

across PFMAs may be applied which may not reflect the variable harvest rates experienced by

the different populations within these large CUs. The impact of observation bias in the catch

assigned to each CU on status assessments is unknown and is a focal aspect of this study (see

Sensitivity analyses, below).

Figure 1. Illustration of the WSP status assessment framework (adapted from Holt et al. 2009). We focused on the

geometric mean spawner abundance (metric, blue) under the spawner abundance indicator. This metric was

assessed against two types of benchmarks: spawner-recruitment (SR) and percentile (see Figure 2). Faded boxes

represent other types of metrics and indicators that may be included in integrated status assessments but were

beyond the scope of what we considered.

Figure 2. a) benchmarks are the 50th (horizontal green line) and 25th (horizontal red line) percentiles of historical

spawner abundance (points). The current spawner abundance is calculated as the geometric mean spawner

abundance over the most recent generation (4 years, blue points and line). b) -recruitment benchmarks are based

on the shape of the Ricker relationship (solid line) fit to data on spawner abundance (x-axis) and corresponding

recruitment (catch + escapement, y-axis). The upper and lower benchmarks are 80% S_{MSY} (green) and S_{GEN} (red),

respectively. S_{GEN} is defined as the spawner abundance that leads to S_{MSY} (grey) in one generation in the absence of

fishing mortality. Under both types of benchmarks, the current spawner abundance in the example shown is above

the upper benchmark, and this CU would be assessed as 'green'. c) The calculation of benchmarks and

benchmarks requires run reconstruction to expand observed spawners abundances, assign catch to CUs, and

calculate recruitment (Table 1).

Figure 3. Our study focused on central coast chum salmon Conservation Units (CUs) as a case study for how

common run-reconstruction assumptions affect the assessment of biological status. salmon CUs (green; different

CUs shaded differently in central coast inset) relatively large do not correspond to the Pacific Fishery Management

Areas (PFMAs; white or light blue shaded regions) for which catch is reported. Therefore, assumptions have to be

made when assigning catch to CUs that may result in under- or over-estimation of catch. Map produced using

PBSmapping (Schnute et al. 2015) with data from (Martin Huang, pers. comm. <Martin.Huang@dfo-mpo.gc.ca>).

Finally, to calculate recruitment for a given cohort of spawners, assumptions about the

age-at-return of spawners in any given year are required (except in the case of pink salmon,

which have a fixed 2-year generation time). The total return in a given year is assigned to brood

years based on the proportion of fish that return at a certain age, but these proportions are

often not estimated every year. For chum salmon on the central coast, the distribution of age-

at-return is assumed to be constant over time and is based on the average of available data

(English et al. 2018). In this case, interannual variability in age-at-return may introduce

uncertainty into the calculation of brood-year recruitment and bias resulting assessments of

status (Zabel and Levin 2002).

Simulation model

We developed and applied a stochastic simulation model of salmon population

dynamics that allows control over various biological and management factors that may

influence the accuracy of status assessments. This approach built on previous studies that

evaluated uncertainties in fisheries management (e.g., Holt and Peterman 2008) and other

factors influencing the performance of metrics and benchmarks under the Wild Salmon Policy

(e.g., Peacock and Holt 2012; Holt and Folkes 2015; Holt et al. 2016, 2018). The simulation

model is comprised of submodels for salmon population dynamics, observation of spawners,

assessment, harvest, and performance (Figure 4). In this subsection, we describe the general

equations for the simulation model, with details of parameterization in the following

subsection.

Figure 4. Schematic of the simulation model comprised of submodels for population dynamics (including harvest),

observation, assessment, and performance. Adapted from Holt et al. (2016).

Population dynamics

We simulated the dynamics of multiple spawning populations returning to indicator or

non-indicator streams within a single hypothetical CU. Although some CUs consist of just a

single spawning population (e.g., lake-type sockeye salmon), many CUs (especially pink and

chum salmon) span thousands of square kilometers (Figure 3) and can include multiple

spawning populations whose dynamics may differ due to local adaptation and finite rearing and

spawning habitats.

We based our simulations on the life history of chum salmon, which generally return to

spawn as 3-, 4-, or 5-year-olds. The number of salmon returning in return year t and population

j, $R_{t,j}$, was calculated as:

$$R_{t,j} = R'_{t-3,j}\, p_{t-3,3} + R'_{t-4,j}\, p_{t-4,4} + R'_{t-5,j}\, p_{t-5,5}, \tag{1}$$

where $p_{y,g}$ is the proportion of recruits from brood year y returning at age g. Throughout this

model description, we use R to denote returns, or catch plus escapement of fish returning in a

year, and R' to denote recruitment, or the total number of offspring from a brood year that

survive to maturity.

We assumed that the annual proportion of recruits returning at a given age was the

same among populations, but incorporated interannual variability in age-at-maturity by

allowing the proportion of recruits that return at age g to vary among brood years, y:

$$p_{y,g} = \frac{\overline{p}_g \exp(\overline{\omega}\, \varepsilon_{y,g})}{\sum_{G=3}^{5} \overline{p}_G \, \exp(\overline{\omega}\, \varepsilon_{y,G})}, \tag{2}$$

262   where $\overline{p}_g$ is the average proportion of individuals maturing as g year-olds, $\overline{\omega}$ is a parameter

263   that controls interannual variability in proportions of fish returning at each age (Figure S4) and

264      $\varepsilon_{y,g}$ are standard normal deviates (Schnute and Richards 1995).

265   The number of salmon that escape the fishery and return to spawn was calculated as:

$$S_{t,j} = \left(1 - h_{t,j}\right) R_{t,j}, \qquad (3)$$

where $R_{t,j}$ is the number of returning salmon from eq. (1) and $h_{t,j}$ is the realized harvest rate

for population j in year t. We incorporated outcome uncertainty (i.e., deviations from the

target harvest rates) by drawing the realized harvest rates for each year and population from a

Beta distribution with mean equal to a target harvest rate, h_t', and variance σ_h^2 (Holt and

Peterman 2008).

We considered two different scenarios for determining the target harvest rate (Figure

5). First, we considered a simple, abundance-based harvest control rule (HCR) where h_t'

increased with the total return to the CU from a minimum of 0.05 (to account for bycatch and

unavoidable mortality and also avoid problems associated with low target HRs when

incorporating Beta-distributed outcome uncertainty) up to an asymptote, h_{MAX} (Holt and

Peterman 2008):

$$h_t' = \max\left(h_{MAX} \left(1 - \exp\left[m \sum_j R_{t,j}\right]\right), 0.05\right), \qquad (4)$$

where m is the shape parameter of the HCR. The low harvest rates at low returns under this

HCR prevented the CU from declining to red status in simulations, and so as to broaden our

results to include CUs with true red status, we also considered a constant high target harvest

rate of $h_t' = 0.60$ regardless of the total return (Figure 5). In the Supplementary Material, we

present an intermediate scenario with a constant moderate target harvest of $h_t' = 0.42$.

Figure 5. The two target harvest-rate cases we simulated were (1) a simple Harvest Control Rule (eq. (4); solid line)

with parameters estimated from historical harvest rates and total return from five central coast chum CUs (grey

points), (2) a constant high target harvest rate of $h'_t = 0.6$ (dotted line).

Each population in our model was harvested in proportion to its abundance, such that

the true total catch of fish that would have returned to streams within the CU was calculated

as:

$$C_t = h_{t,j} \sum_j R_{t,j}. \tag{5}$$

Although realized harvest rates differed among populations, we did not incorporate persistent

biases in realized harvest rates among populations and thus assumed that all populations were

equally vulnerable to the fishery. The extent to which this assumption is violated will depend

on the size of the CU, the number of populations within it, the magnitude of variation in run-

timing and body size among populations, and where fisheries are prosecuted. Any such biases

among populations within a CU would likely be small because run-timing was a consideration

when delineating CUs (Holtby and Ciruna 2007). However, biases among CUs may be significant

and we investigate this by varying the observation bias in the total catch to the CU (see

Sensitivity analyses, below).

Finally, we assumed the spawner-recruitment dynamics followed a Ricker relationship

(Ricker 1954) yielding the number of recruits from brood year y and population j:

$$R'_{y,j} = S_{y,j} \, \exp\!\left(a_j - b_{j,y} \, S_{y,j} + \phi_{y,j}\right), \tag{6}$$

where a_j is the $\log_e$ recruits per spawner at low spawner abundance (i.e., productivity), $b_{j,y}$ is

the density-dependence parameter (which we allowed to change linearly through time; see

below), and $\phi_{y,j}$ are the normally distributed recruitment deviates applied for year y and

population j (eq. (7)).

We allowed productivity to differ among populations, where a_j was drawn from a

truncated normal distribution with mean $\overline{a}$ and variance σ_a^2, with a lower limit of $a_j = 0.4$

because SR benchmarks are not calculable for very low productivity (Holt and Ogden 2013; Holt

et al. 2018). For central coast chum salmon, we found only 1% of spawning populations (2/181)

had $a_j < 0.4$ (Online Supplement). A linear change in the density-dependence parameter was

simulated for some populations as a decline in the capacity of the freshwater habitat (i.e.,

$S_{MAX,j,y} = 1/b_{j,y}$, or the spawner abundance that leads to maximum recruitment). This decline

in capacities captured the potential consequences of cumulative stressors to freshwater habitat

among watersheds on the central coast (Connors et al. 2018). For all populations, the initial

capacity $S_{MAX,j,1}$ was drawn from a lognormal distribution whose mean and standard deviation

differed for populations in indicator versus non-indicator streams, as indicator streams tend to

be larger systems (English 2016; see Parameterization). The productivity and density-

dependence parameters were drawn independently for each Monte Carlo (MC) iteration of the

model.

We incorporated temporal autocorrelation in recruitment deviates:

$$\phi_{y,j} = \tau \, \phi_{y-1,j} + \sigma_v \sqrt{1 - \tau^2} \, v_{y,j}, \tag{7}$$

319   where τ is the temporal autocorrelation coefficient, σ_v is the standard deviation in recruitment

320   deviates without temporal autocorrelation (Ricker 1975, Holt and Bradford 2011), and and $v_{y,j}$

321   is a multivariate normal random variable with mean zero and variance-covariance matrix:

$$\Sigma_{j \times j} = \begin{bmatrix} \sigma_v^2 & \rho \, \sigma_v^2 & \dots & \rho \, \sigma_v^2 \\ \rho \, \sigma_v^2 & \sigma_v^2 & \dots & \rho \, \sigma_v^2 \\ \vdots & \vdots & \ddots & \vdots \\ \rho \, \sigma_v^2 & \rho \, \sigma_v^2 & \dots & \sigma_v^2 \end{bmatrix}_{j \times j}. \tag{8}$$

Here, ρ is the correlation in recruitment deviates among populations.

We simulated the 'true' population dynamics over 50 years, after an initialization period

of seven years to seed eq. (6) given the variable age-at-return of chum salmon. For each year in

this initialization, we assumed that the number of spawners was equal to 20% of $S_{MAX,j,1}$. For

the first year of the initialization, we set $\phi_{y-1,j}$ from eq. (7) to zero.

Observation submodel

In the observation submodel, we incorporated both incomplete monitoring coverage of

streams and imperfect observation of spawners in streams that were monitored. In any given

year, population *j* was observed with probability $\psi_{t,j}$. We included a linear decline in

monitoring coverage (i.e., the probability of a population being observed) over time based on

observations of declining monitoring coverage on the north and central coast of BC (Price et al.

2008, 2017; English 2016). We calculated the annual probability of being monitored separately

for indicator and non-indicator streams based on observations that monitoring coverage of

non-indicator streams is generally lower and has declined more severely than coverage of

indicator streams (English 2016). See Parameterization for further details.

Spawner abundances were 'observed' with log-normal error:

$$\hat{S}_{t,j} = z_{t,j} \left[S_{t,j} \, \exp(\delta_{t,j}) \right], \tag{9}$$

where $z_{t,j} \sim \text{Bernoulli}(\psi_{t,j})$, $\delta_{t,j} \sim N(\overline{\delta}, \ \sigma_\delta)$, $\overline{\delta}$ is the mean observation error, and σ_δ is the

standard deviation in observation error of spawner abundances. Thus, this combines both the

probability of a population being monitored and the distribution of observation errors around

true spawner abundances if monitored. We included a negative bias in the observation of

spawners ($\overline{\delta} \leq 0$) such that the mean observed spawner abundance is on average lower than

the true spawner abundance. In general, it is challenging to enumerate spawners in all reaches

of a stream and in all streams within a CU. The reported spawner abundance is considered an

underestimate of the total spawners in a CU, which motivates the application of Expansion

Factor III for observer (in)efficiency when performing run reconstructions (Table 1). The

calculation of Expansion Factor I required that we impose the constraint that at least one

indicator stream was monitored each year, so if $z_{t,j}$ = 0 for all indicator streams in a year, we

randomly selected one indicator stream to be monitored.

The catch to the entire CU in return year t was observed with log-normal error:

$$\hat{C}_t = C_t \, \exp(\chi_t), \tag{10}$$

where $\chi_t \sim N(\bar{\chi}, \sigma_\chi)$, σ_χ is the standard deviation in catch error, and $\bar{\chi}$ is a bias in catch. We

assumed a default value of $\bar{\chi}$ =0, but in sensitivity analyses we varied this parameter to simulate

scenarios in which fish are caught from other CUs or fish from the focal CU were caught in other

PFMAs.

Previous models (e.g., Holt et al. 2018) have included error in the "estimated age-at-

return" separately for each return year. For central coast chum salmon, annual age-at-return

data are rarely sampled comprehensively so the same average is generally applied across all

years (Peacock et al. 2014; English et al. 2018). Therefore, for each year in a MC trial we

applied the same age-at-return, which was drawn independently for each MC trial using eq. (2)

with observation error, $\overline{\omega_r}$ (Table A1).

Assessment submodel

As described above, the minimum requirement to calculate benchmarks and assess

population status using percentile benchmarks is a time-series of spawner abundance to the

CU. For SR benchmarks, harvest rates and age-at-return must also be estimated in order to

calculate recruitment. The basic procedure of these run reconstructions is outlined in Table 1,

and begins by expanding observations of spawners to indicator streams to the total spawner

abundance to the CU by applying three Expansion Factors. The equations and criteria

governing these Expansion Factors are detailed in the Online Supplement and in English et al.

(2012, 2016, 2018). Briefly, Expansion Factor I ($F'_{t,d}$) imputes missing spawner abundances for

populations in unmonitored indicator streams and is calculated for each year, *t*, within a

decade, *d*, of the spawner time series. It relies on the decadal contribution of each indicator-

stream population to the total escapement to all indicator streams (English et al. 2016).

Expansion Factor II (F''_d) expands observations of spawners from indicator streams to include

populations in non-indicator streams that are less frequently monitored, and is the same for

each year within a decade, *d*. Expansion Factor II is calculated as the average proportion of

total monitored spawners (in indicator and non-indicator streams) that are in non-indicator

streams for that decade. For decades with insufficient information to calculate either of these

Expansion Factors, for example due to declining monitoring coverage, a reference decade may

be used. Expansion Factor III (F''') is determined by the regional DFO staff familiar with the

escapement monitoring techniques used in each statistical area and is assumed to be constant

through time (English et al. 2018). In our model, we assumed that all populations were at least

partially monitored, and that Expansion Factor III accounted for observation (in)efficiency, but

in reality, Expansion Factor III may also account for populations in unmonitored streams.

The observed number of salmon returning in year t is the sum of observed catch and

expanded escapement to the CU:

$$\hat{R}_t = \hat{C}_t + F'_{t,d}\, F''_d\, F''' \sum_{j \in I} \hat{S}_{t,j}, \tag{11}$$

where the summation includes observed spawner abundance to the I indicator streams only,

with the non-indicator streams being accounted for through Expansion Factor II, F''_d.

We do not explicitly account for en route or pre-spawning mortality of fish that escaped

the fishery, and assume that pre-spawning mortality is relatively small and accounted for in the

productivity of the population through the Ricker SR dynamics. Observed recruitment for

brood year y is calculated as the sum of age 3, 4, and 5 fish returning in years $y + 3$, $y + 4$, and

$y + 5$, respectively:

$$\hat{R}'_y = \hat{R}_{y+3}\, \hat{pr}_3 + \hat{R}_{y+4}\, \hat{pr}_4 + \hat{R}_{y+5}\, \hat{pr}_5, \tag{12}$$

yielding the 'reconstructed' spawner-recruit pairs for brood year y.

To calculate estimated SR benchmarks, we fit a linearized Ricker model to the observed

data at the aggregate CU-level. The estimated productivity and density-dependence

parameters were then used to calculate upper and lower SR benchmarks (80% of S_{MSY} and

S_{GEN}) for the CU.

Performance submodel

For each MC simulation, we estimated status using both SR and percentile benchmarks

calculated from the observed spawner-recruit pairs for the CU, including observation biases and

incomplete monitoring coverage. Estimated status under both types of benchmarks was

compared to the true status, which was calculated by comparing the current true spawner

abundance (without observation error) against the upper and lower SR benchmarks (80% of

S_{MSY} and S_{GEN}) from the underlying SR parameters. This meant that we evaluated estimates of

percentile (and SR) benchmarks against the true underlying benchmarks derived from "true" SR

parameters. Because we simulated the true dynamics at the scale of spawning populations and

there was no "true" CU-level value for the Ricker parameters, calculating the true S_{MSY} (and

thus the SR benchmarks) at the CU-level was not straightforward. We chose to calculate the

true SR benchmarks for each population from the underlying Ricker parameters for that

simulation and then summed the benchmarks across all populations to yield the "true" CU-level

benchmarks. Although this CU-level benchmark will underestimate the level required to

maintain all component populations above their individual benchmarks in any given year, the

objective of the WSP is to maintain the overall CU and populations within CUs are assumed be

recolonizable within reasonable time frames (Fisheries and Oceans Canada 2005). When

declines in capacity were included in the simulation, we calculated true status from the initial

capacity parameters before the decline. Although this may have increased the chance that

estimated status was lower than true status (increasing the rate of pessimistic

misclassifications), we took this approach to avoid a shifting baseline in benchmarks.

Performance was evaluated in two ways that capture the difference between estimated

and true status: (1) the proportion of MC simulations for which status was correctly assessed as

green, amber, or red, and the proportion of simulations for which status was either

underestimated (pessimistic) or overestimated (optimistic), and (2) the percent relative bias of

observed average spawner abundance (S_{AVG}) and of the four benchmarks (S_{GEN}, 80% S_{MSY}, S_{25},

S_{50}) compared to their true values for each MC simulation.

For each parameterization investigated (see below) we ran 4000 MC trials, which was sufficient

to ensure the mean percent error in performance measures was < 3% (Figure S3).

Parameterization

Some of the parameters in our simulation model were unknown or unknowable, in

which case we followed assumptions made for southern BC chum salmon by Holt et al. (2018).

Other parameters were available specifically for central coast chum salmon or could be

estimated from available data; details of parameter estimation are given in the Online

Supplement. As mentioned above, in order to understand the assessment biases under

different true statuses we considered two cases: (1) high productivity and a conservative

harvest control rule (HCR), which we refer to as the "base case" because it is most

representative of central coast chum salmon, and (2) low productivity and a constant high

target harvest rate, which represented a CU at high risk of extirpation. In the Online

Supplement, we also present results from a third case intermediate between these two with

low productivity and a constant moderate harvest rate. Unless otherwise indicated in the

sensitivity analysis (below), parameters defaulted to the values described here and listed in the

Appendix, Table A1.

The mean proportion of adults maturing at ages 3, 4, and 5 in eq. (2) was 0.23, 0.64, and

0.13, respectively, based on the average age-at-return applied in run reconstructions of central

coast chum (Challenger et al. 2018; English et al. 2018).

We estimated the parameters in the HCR (Figure 5) from harvest rates and total return

sizes for five central coast chum CUs (English et al. 2018; Salmon Watersheds Program - Pacific

Salmon Foundation 2019).

To estimate the SR parameters for spawning populations (eq. (6)), we fit a linearized

Ricker model to population-level spawner-recruit pairs from nine central coast chum CUs with

individual productivity and density-dependence parameters for each population (Figure S5).

Note that this approach may lead to biased SR parameters as we did not account for temporal

autocorrelation and errors in variables when analysing the data, but they nonetheless provide a

useful starting point for our simulation study. From the model fits, we calculated (1) mean and

variance in productivity among populations, (2) mean and variance in the initial capacity (i.e.,

$1/S_{MAX}$) for indicator and non-indicator streams, (3) the residual variance within populations, (4)

the average correlation in residuals between populations, and (5) the temporal autocorrelation

in residuals within populations. The mean population-level productivity was $\bar{a}$ = 1.40 (Figure

S5), which we applied in our base case. For the low-productivity case, we chose $\bar{a}$ = 0.56, which

was the 2.5[th] percentile of population-level productivity estimates (Figure S5). The residual

variance within populations was used as an estimate of the variance in recruitment deviations

(σ_v^2) in simulations, but was likely an overestimate as we did not account for observation error

separately when fitting the Ricker model. Thus, we explore the results under a range of values

for σ_v in the Supplemental Results (Online Supplement). The correlation in residuals among

populations was estimated at $\rho = 0.46$, but we also investigated lower ($\rho = 0$) and higher ($\rho =$

0.9) levels of synchrony in the Online Supplement.

As the default case, we considered a decline in the capacity of streams (i.e., one over

the density-dependence parameter of the Ricker model) that reflected observed declines in

freshwater habitat (Office of the Auditor General of Canada 2004). Within the central coast

chum CUs, 29% of watersheds are considered to be at moderate risk, and 21% at high risk, of

cumulative habitat pressures over the last 60 years from stressors such as logging, water

licenses for withdrawal of water from streams, and stream crossings (Connors et al. 2018). We

hypothesized that declines in capacity that differ among spawning populations may affect the

accuracy of Expansion Factors I and II, particularly in combination with declining monitoring

coverage (Price et al. 2017). We incorporated a linear decline in capacity (i.e., S_{MAX} = 1/*b*) over

the 50-year time series of between 25% and 50%, representing a moderate decline, for 29% of

populations (the percentage of central coast chum watersheds deemed to be at moderate risk

of cumulative habitat pressures by Connors et al. 2018) and a linear decline between 50% and

75%, representing severe decline, for 21% of populations (the percentage deemed to be at high

risk). The exact percent decline for each population was randomly drawn from a uniform

distribution within the above range for each MC simulation. The remaining 50% of populations

had stable capacity over the 50-year simulation. In a sensitivity analysis, we investigated four

additional scenarios for declining capacity (Table 2).

In the observation submodel, we chose the bias in the observation of spawners to

match Expansion Factor III, which corrects for observer efficiency, with a range of values

explored in a sensitivity analysis (below). Because Expansion Factor III is simply a constant

multiplier applied to spawner abundance, it does not affect estimated status by the percentile

benchmarks. However, Expansion Factor III affects estimated harvest rates and recruitment,

and therefore bias in this Expansion Factor was expected to lead to bias in estimated status

under the SR benchmarks. The value of Expansion Factor III applied in past status assessments

has been constant at F''' = 1.5 for all central coast chum CUs (English et al. 2016), and so we

applied a default value of $\bar{\delta}$ = log$_e$(1/1.5) = -0.4. The majority of central coast chum streams are

surveyed visually by ground (fish counts or other sampling) with some aerial (fish counts) or

boat (fish counts or other sampling) surveys (English 2016), similar to southern BC chum

streams (Holt et al. 2018). We assumed σ_δ = 0.5 following Holt et al. (2018), which is the

maximum estimated uncertainty for visually surveyed spawners (Cousens et al. 1982; Szerlong

and Rundio 2008).

We incorporated a linear decline over the last 27 years of simulations in the proportion

of indicator and non-indicator streams monitored each year from 0.76 and 0.72, respectively, to

0.72 and 0.05 based on English (2016). These declines are representative of overall declines in

monitoring across species, but we also consider the trends specific to chum salmon in the

Online Supplement (Figure S6).

We assumed no bias in the observation of catch ($\bar{\chi}$ = 0) by default, but consider a range

of bias in sensitivity analyses (below). The standard deviation in the observation error of catch

accounts for differences between observed (i.e., reported) and actual catch due to

uncertainties with mixed-stock fisheries and in reporting and estimation of recreational

fisheries and subsistence use. We set this to σ_{χ} = 0.2 (Holt et al. 2018), which is less than the

observation error in spawners.

Sensitivity analyses

We examined the sensitivity of status assessments over a range of values for several

different model parameters that we considered most likely to affect status assessments due to

their influence on the assumptions in run reconstructions (Table 1; Table 2). We investigated

each of the questions below under both the base case and low-productivity high-harvest case

described above, with other parameters at their default values (Table A1) unless otherwise

noted. The specific questions that we addressed were:

1. How does the number of spawning populations and the proportion designated as

indicator streams affect status assessments? The lower the proportion of streams that

are indicators, the greater the magnitude of Expansion Factor II.

2. How does a decline in monitoring coverage affect status assessments? The fewer

indicator streams that are monitored, the greater the magnitude of Expansion Factor I

and the potential uncertainty in expanded spawner abundance. Here, we consider two

scenarios (Table 2; Figure S6): constant monitoring coverage at historical proportions

among all streams and an observed decline in coverage starting in the mid-1980s as has

been observed on the north and central coast (English 2016; Figure S6). In the Online

Supplement we consider two additional scenarios: observed declines in monitoring

specific to chum salmon streams and a sharp, recent decline in monitoring of indicator

streams.

3. How do declines in capacity affect status assessments? The application of Expansion

Factors I and II assumes that the relative contributions of populations to aggregate

abundance in the CU does not change over time, but declines in capacity that differ

among populations may violate this assumption.

4. How does spawner observation bias affect status assessments, given that the value of

Expansion Factor III is fixed over time and often the same among CUs (English et al.

2018)?

5. How does catch observation bias (e.g., over- or under-estimating catch of salmon) affect

status assessments? This represents scenarios where there are errors in estimates of

CU proportions in the aggregate catch in a mixed-stock fishery, or violation in the

assumption of homogenous spatial and temporal distribution of CUs when CU

proportions are not monitored in such fisheries.

6. How does interannual variability in age-at-return affect status assessments?

We investigated the impact of declines in monitoring coverage (question #2 above) in

combination with declines in capacity of spawning populations (question #3) in a bivariate

sensitivity analysis.

Results

The different productivity and harvest rate combinations we considered led to different

true CU statuses. Under high productivity and an abundance-based harvest control rule (HCR) –

the base case corresponding to central coast chum salmon – 86.3% of simulations resulted in

true green status (Figure 6a,b). Conversely, under low productivity and high harvest rates,

67.7% of simulations resulted in true red status (Figure 6c,d).

Under the base case when true status was mostly green, misclassifications resulted in

estimated status lower than the true status meaning assessments were biologically pessimistic

(henceforth referred to as "pessimistic misclassifications"). This was particularly true of the

percentile benchmarks, for which 55.3% of simulations resulted in a pessimistic

misclassification with 11.8% of simulations having misclassified green status as red. Pessimistic

misclassifications were due to positive bias in benchmarks and not bias in the current spawner

abundance (Figure 7), resulting in status being underestimated. For productive populations (as

in the base case), most observed spawner abundances tended to be far above lower

benchmarks and closer to equilibrium values. As a result, lower and upper benchmarks of 25th

and 50th percentile of historical spawner abundance tended to overestimate the "true" SR-

based benchmarks.

When true status was mainly red, under low productivity and high harvest rates,

biologically optimistic misclassifications (henceforth "optimistic misclassifications") were more

common, which may be riskier from a conservation and management standpoint. For example,

46.5% and 44.3% of simulations had an estimated status higher than true status under the

spawner-recruitment (SR) and percentile benchmarks, respectively (Figure 6c,d). These more

frequent optimistic misclassifications were due to a negative bias in benchmarks, in particular

the lower SR benchmark of S_{GEN} (Figure 7a), likely due to a poor ability to estimate S_{MSY} under

low productivity when spawner abundances tend to cluster near the origin.

Under both types of benchmarks, bias did not decrease when monitoring coverage was

held constant at 100% (Figure S9), suggesting that the application of Expansion Factors I and II

were not contributing factors.

Figure 6. Estimated status according to the spawner-recruitment (SR) benchmarks (left) and the percentile

benchmarks (right), over the true status for each of two cases (high productivity and an HCR (a-b) and low

productivity and high harvest rates (c-d)). Grey cells indicate pessimistic misclassifications, which may lead to overly

conservative management actions, and black cells indicate optimistic misclassifications, which may lead to overly

risky management actions. Coloured cells indicate correct classifications for red, amber, and green zones,

respectively.

Figure 7. Relative bias in spawner-recruitment (SR) benchmarks (a), percentile benchmarks (b), and current

spawner abundance (S_{AVG}; black) (median ± interquartile range among 4000 MC simulations) for high productivity

and a harvest control rule (HCR) and the low-productivity high-harvest case.

Sensitivity analyses

The number of spawning populations within the CU and the proportion of those

populations spawning in indicator streams had little impact on status assessments (Figure S10).

Under the base case, the relative bias in estimates of S_{MSY} and S_{GEN} were lower in larger CUs,

with half as many pessimistic misclassifications for larger CUs under the SR benchmarks (40%

for 10 populations versus 20% for 140 populations with 30% indicator streams; Figure S10a).

This trend was not, however, observed in the low-productivity high-harvest case when true

status was predominantly red (Figure S12).

The monitoring-coverage scenarios that we considered, representative of observed

declines in monitoring on the north and central coast, had no effect on status outcomes or the

relative bias in benchmarks. This was true in the base case (Figure 8, Figure S13) and under low

592   productivity (Figures S14 – S15). Even under severe declines in capacity of 50% to 75% for all

593   spawning populations, our results suggest that the observed declines in monitoring coverage on

594   the north and central coast are unlikely to bias status assessments. This result held regardless

595   of whether the recruitment deviates among component populations within the CU were not

596   correlated (ρ = 0; Figure S16) or highly correlated (ρ = 0.9; Figure S17).

Figure 8. The effect of monitoring coverage (no change and decline; Table 2) and the percentage of spawning

populations with severe declines in capacity (x-axis) on performance measures under the base case of high

productivity and HCR. (a-b) The proportion of simulations with correct green, amber, or red status or pessimistic

misclassifications (grey) and optimistic misclassifications (black) under the SR benchmarks (a) and percentile

benchmarks (b). (c-d) The percent relative bias (median $\pm$ interquartile range among 4000 MC simulations) in the

current spawner abundance (S_{AVG}; black) and lower and upper benchmarks (red and green, respectively) under the

SR benchmarks (c) and the percentile benchmarks (d). See Online Supplement for results under the low-productivity

high-harvest case.

Under the base case, declines in capacity of the CU were associated with poorer

estimated status and an increase in misclassification rates (Figure 8a-b). Pessimistic

misclassifications increased because CUs more frequently had a true status of amber but were

609   misclassified as red. Optimistic misclassifications increased, particularly under the SR

610   benchmarks (Figure 8a), because the relative bias in the current spawner abundance (S_{AVG})

611   remained unchanged, but the bias in benchmarks decreased (Figure 8c-d). In the low-

612   productivity high-harvest case, the results were similar but with biologically optimistic

613   misclassifications dominating as status was predominantly amber or red (Figure S15).

Figure 9. The effect of observation bias in the number of spawners (x-axis) on performance measures under the

base case. (a-b) The proportion of simulations with correct green, amber, or red status or pessimistic

misclassifications (grey) and optimistic misclassifications (black) under the SR benchmarks (a) and percentile

benchmarks (b). (c-d) The percent relative bias (median $\pm$ interquartile range among 4000 MC simulations) in the

current spawner abundance (S_{AVG}; black circle) and lower and upper benchmarks (red and green, respectively)

under the SR benchmarks (c) and the percentile benchmarks (d). The asterisk in (a-b) and grey zone in (c-d) indicate

the default parameter value of $\bar{\bar{\delta}} = -0.4$, and the bias that matches the Expansion Factor III of $F''' = 1.5$ applied in all

simulations. See Online Supplement for results under the low-productivity high-harvest case (results were similar).

As the bias in the observation of spawners approached zero ($\bar{\bar{\delta}} \to 0$), misclassifications

under the SR benchmarks declined in all cases, even as the observation bias became less than

the Expansion Factor III applied to correct for it (Figure 9a, Figures S18-S19). The relative bias in

the current spawner abundance was minimized when the observation bias matched the

assumed value of Expansion Factor III ($\bar{\delta}$ = -0.4 corresponding to F''' = 1.5; Figure 9c-d). When

observed spawner abundance was biased low ($\bar{\delta}$ = -1.6), the relative bias in the upper

benchmark of 80% S_{MSY} was higher than the relative bias in current spawner abundance (S_{AVG})

or the lower benchmark (S_{GEN}), and thus CUs with a true green status were more likely to be

misclassified as amber. Status outcomes under the percentile benchmarks were unaffected by

changes in observation bias of spawners (Figure 9b); the bias in both benchmarks and current

abundance showed similar changes as observation bias declined (Figure 9d) such that the

resulting status outcome was unchanged. In the Online Supplement, we also investigated a

change in observation bias halfway through the simulation (Figure S20), but a change from the

base value of $\bar{\delta}$ = -0.4 to $\bar{\delta}$ = -1.6, -0.7, and 0 did not have any impact on status outcomes or

biases in benchmarks (Figure S21).

Underestimation of catch (i.e., negative values of $\bar{\chi}$) resulted in fewer misclassifications

than overestimating catch (Figure 10, Figures S22 – S23). As the catch bias increased from $\bar{\chi}$ = -

1.0 (63% underestimation) to $\bar{\chi}$ = 1.0 (271% overestimation), the relative bias in the lower SR

benchmark of S_{GEN} declined while the relative bias in the upper benchmark of 80% S_{MSY}

increased (Figure 10b). This is due to the errors in variables that occur when catch is

underestimated: productivity and recruitment tend to be underestimated, thus leading to

lower estimates of S_{MSY} and higher estimates of S_{GEN} (Holt and Folkes 2015). Under the base

case, the true status was green in the majority of simulations and so the increasing bias in the

upper benchmark dominated the overall status assessments and led to the increase in

pessimistic misclassifications with increasing $\bar{\chi}$. In the low-productivity high-harvest case, true

status was mostly red and so the increasingly negative bias in S_{GEN} resulted in more optimistic

misclassifications as the observation bias in catch increased (Figure S23). In all cases,

overestimating catch by ~50% (i.e., $\bar{\chi}$ = 0.4) led to a 7 - 10% increase in misclassification rate

(Figure S24). Although these changes in misclassification rates may seem small, there is

potential for large catch errors in run reconstructions, especially when multiple CUs overlap

with a single PFMA. Catch does not factor into the calculation of percentile benchmarks, so

status under the percentile benchmarks was unaffected by changing catch bias.

655   Finally, increasing interannual variability in age-at-maturity resulted in more frequent

status misclassifications, but the effect was relatively small. Under the base case, increasing $\bar{\omega}$

from 0.2 to the default value of 0.8 resulted in an increase in misclassifications from 26.3% to

29.7%, but very little change in the bias in benchmarks (Figure S25). Further increasing the

interannual variability to $\bar{\omega}$ = 1.6 led to 34.2% of simulations being misclassified, but this level of

interannual variability is high (see Figure S4 for example) compared to data on age-at-return for

central coast chum salmon (Challenger et al. 2018; English et al. 2018). The increase in

misclassifications was smaller under the low-productivity high-harvest case (Figure S25).

Figure 10. The effect of observation bias in catch (x-axis) on performance measures under the base case. (a) The

proportion of simulations with correct green, amber, or red status or pessimistic misclassifications (grey) and

optimistic misclassifications (black) under the SR benchmarks. (b) The percent relative bias (median $\pm$ interquartile

range among 4000 MC simulations) in the current spawner abundance (S_{AVG}; black circle) and lower and upper

benchmarks (red and green, respectively) under the SR benchmarks. The asterisk in (a) and grey zone in (b) indicate

the default parameter value of $\bar{\chi}$ = 0. Observation bias in catch ranges from 63% underestimation ($\bar{\chi}$ = -1) to 271%

overestimation ($\bar{\chi}$ = 1). See Online Supplement for results under the low-productivity high-harvest case.

Discussion

In this study, we quantified the impact of common assumptions in run reconstructions

(Figure 2c) on biological status outcomes. In general, the assumptions that we hypothesized

might introduce biases (Table 1) had little effect on status outcomes, suggesting that attempts

to assess status in the face of limited or uncertain data are worthwhile. In particular, there was

almost no effect of declines in monitoring coverage on the accuracy of benchmarks or resulting

status outcomes, even in the face of different trends in capacity and reduced synchrony (i.e.,

zero autocorrelation in recruitment deviates) among spawning populations within the CU.

Extreme declines in monitoring may impact assessments – certainly, if no spawning populations

are monitored then our ability to assess status will inevitably be compromised – but the current

approach to imputing and expanding spawner abundances proved robust to declines in

monitoring coverage in the range documented for the north and central coast (English 2016;

Price et al. 2017).

Perhaps unexpectedly, misclassifications were not minimized when the value of

Expansion Factor III, correcting for observer efficiency, accurately reflected the true observation

bias in the underlying simulation. Rather, misclassifications under the spawner-recruitment

(SR) benchmarks declined as the observation bias in spawners approached zero, due to the

combination of bias in current spawner abundance, which *is* minimized when observation bias

matches Expansion Factor III, and biases in benchmarks, which depend on underlying status.

Status outcomes under the percentile benchmarks were unaffected by observation bias, as this

bias was assumed to be constant over time and affected the current and historical spawner

abundances equally.

Under our base case of high productivity and an abundance-based harvest control rule

(HCR) consistent with historical central coast chum salmon harvest rates, most simulations had

a true green status, but pessimistic misclassifications as amber were common under both SR

and percentile benchmarks. The estimated status from our simulations roughly matched the

status outcomes for seven central coast chum CUs from that period, with the majority of CUs

having green status under the SR benchmarks and amber status under the percentile

benchmarks (Connors et al. 2018). (Note that status of central coast chum salmon CUs has

since declined; see the Pacific Salmon Explorer (www.salmonexplorer.ca) for the most up-to-

date assessments.) Consistent with the real status assessments, status under the percentile

benchmarks tended to be poorer than status under the SR benchmarks. Our simulations

attributed this to a higher relative bias in percentile benchmarks, as found for south-coast chum

salmon under high-productivity scenarios (Holt et al. 2018). Pessimistic misclassifications may

lead to overly conservative management actions, consistent with the Precautionary Principle

(Foster et al. 2000), but may also result in foregone harvest and adverse socio-economic

consequences (Walters et al. 2019).

For simulations with low productivity and high harvest rates (i.e., true red status), the

bias in benchmarks was negative, resulting in a high proportion of optimistic misclassifications.

This represents a high-risk management scenario from a conservation standpoint, particularly

as the true underlying status is red. Further, the absolute biases in benchmarks were more

severe under declines in capacity when status was predominantly red (productivity low)

compared to predominantly green (productivity high). The bias in SR benchmarks was

particularly sensitive to the underlying true status, presumably because productivity and

recruitment, which influence status, also affect the bias in SR parameters (Subbey et al. 2014)

that arises due to recruitment-spawner linkage inherent in the data (Walters 1985; Korman et

al. 1995) and/or due to error in spawner estimates (Walters and Ludwig 1981; Kehler et al.

2002; Kope 2006).

Our results suggest that overestimating catch should be avoided. In particular, under

low productivity and high harvest rates, optimistic misclassifications associated with

oversestimating catch and therefore underestimating the lower benchmark, S_{GEN}, may put

populations at further risk. Under the base case of high productivity and an HCR,

overestimating catch resulted in more frequent pessimistic misclassifications as the upper

benchmark (S_{MSY}) was overestimated, resulting in CUs with a true green status being estimated

as amber. In both cases, the impact of overestimating catch has the potential to significantly

bias assessments: overestimating catch by ~50% led to a 7-10% increase in misclassification

rates. This level of catch overestimation (and higher levels) may occur when fish caught in a

Pacific Fisheries Management Area (PFMA) and assigned to the CU that overlaps with that

PFMA were actually bound for other CUs. This could occur in mixed-stock fisheries if genetic

stock identification is not undertaken to validate assumptions regarding run-timing and

migration patterns. Increased efforts to quantify catch composition, run timing, and spatial

distribution of Pacific salmon CUs are therefore needed to more accurately estimate harvest

rates and minimize misclassifications of biological status.

Limitations, challenges, and future research

Closed-loop simulation models are powerful tools for evaluating management strategies

and quantifying the biases in parameter estimation and status outcomes (Walters 1986;

Peterman 2018), but are not without their weaknesses. As with any model, our simulation

model was an approximation of reality and thus we had to make a number of assumptions.

Both the true population dynamics and assessment submodel assumed a Ricker SR relationship

(Ricker 1954), but we recognize that there is considerable model uncertainty. Distinguishing

among different models (e.g., Ricker versus Beverton Holt) in assessing status would be

challenging, particularly under the declines in capacity that we simulated to capture observed

changes in freshwater habitat. The Ricker model is commonly used to set management targets

and for simulating population dynamics of Pacific salmon (e.g., Peterman et al. 2000; Peacock

and Holt 2012; Holt and Folkes 2015), and a full exploration of other true models was beyond

our scope.

We considered true SR dynamics to operate at the spatial scale of spawning populations

(i.e., individual streams), but there is evidence that the processes influencing productivity and

density-dependence may operate at broader, regional spatial scales (e.g., Malick and Cox 2016).

Other simulation models have incorporated straying among spawning populations within a CU

(e.g., Peacock and Holt 2012; Holt and Folkes 2015). While that approach incorporates density-

dependence that may occur at broader spatial scales, it also requires additional assumptions to

be made about the probabilities of straying among streams, which are largely unknown.

Simulating true dynamics at the scale of individual spawning populations also

complicates the calculation of true status at the CU-level from SR benchmarks. We chose to

calculate SR benchmarks at the spawning-population-level and then sum across spawning

populations to calculate S_{MSY} and S_{GEN} at the CU-level. There are other approaches to calculate

aggregate benchmarks, but each has its own potential biases. For example, SR relationships

could be fit to the "true" data aggregated at the CU-level and SR benchmarks calculated from

the resulting CU-level estimates of productivity and density dependence. The way in which

spawning-population-level benchmarks are aggregated to CU-level benchmarks may affect

performance in our simulations, and a full exploration of how different methods of aggregation

affect our results warrants future consideration.

Despite these limitations, the simulation model that we adapted and applied is flexible

enough that it can accommodate different Pacific salmon species and life-history traits, opening

the door to future work investigating the impact of different assumptions and the impact of the

assumptions that we focused on under additional scenarios. Here, we considered a relatively

simple run-reconstruction model, but further work is needed to quantify how observation

errors and uncertainty in the spatial and temporal distribution of returns affects status

outcomes when more complex run-reconstruction models are used. Temporal shifts in

biological parameters, including age-at-return (e.g., associated with environmental change and

selective fisheries) and productivity (non-stationarity has been widely observed in Pacific

salmon; e.g., Peterman and Dorner 2012; Malick and Cox 2016; Dorner et al. 2017), are also

areas that warrant further exploration. Additional simulations could also help inform methods

in the assessment process, such as the optimal time-series length for detecting changes,

whether benchmarks should be updated with each assessment, and the best analytical

approach to calculate SR benchmarks (e.g., Bayesian hierarchical models vs. single-stock

ordinary least squares).

Conclusions

Pacific salmon are one of the most data-rich groups of fish due to their high economic,

social, and cultural value, but nonetheless our knowledge of their dynamics is uncertain.

Assessing the biological status of Pacific salmon CUs is a conservation and management priority

given the continued declines of many stocks (e.g., COSEWIC 2016, 2017) and escalating threats

to salmon conservation. Current government-led integrated status assessments under the WSP

include expert opinion on the data uncertainties and risks unique to each CU (e.g., DFO 2015,

2016, 2018a), making the process time and resource intensive. This has limited their timely

application to all 460+ Pacific salmon CUs in Canada (DFO 2019), and also means that the

process is not entirely transparent or reproducible. The Pacific Salmon Foundation has

implemented data-driven biological status assessments based on a subset of the indicators and

benchmarks recommended under the WSP, with the results accessible through the Pacific

Salmon Explorer (www.salmonexplorer.ca). These data-driven assessments, similar to those

undertaken by other management and conservation organizations (e.g., Marine Stewardship

Council, COSEWIC, Pacific Salmon Commission), require assumptions to estimate spawner and

recruit time series for CUs. Given the lack of expert scrutiny of the data for each individual CU

under this data-driven approach, it was important to understand how these assumptions may

bias status outcomes under a range of biological and management scenarios.

We found that the data-driven biological status assessments were relatively insensitive

to common assumptions in expanding spawner abundances within the parameter ranges we

explored, but the rate and type of status misclassifications depended on the underlying status

of the CU and may be of greater concern for CUs with poor status. To ensure the accuracy of

data-driven status assessments, increased efforts to collect data on catch composition, age-at-

return, and spawner abundances are needed. Such information will help, for example, to

define plausible ranges of error in catch estimation to lend confidence to estimates of

recruitment and thus assessments under SR benchmarks. Nonetheless, our research suggests

that current efforts to assess status in the face of imperfect and incomplete data are

worthwhile for central coast chum salmon and other similar Pacific salmon populations and can

provide a timely approach to assessing status for CUs that complements more thorough

integrated status assessments.

Acknowledgements

The authors thank the Pacific Salmon Foundation's Population Science Advisory Committee, in

particular David Peacock and Randall Peterman, and Steven Cox-Rogers and Charmaine Carr-

Harris for their valuable feedback throughout this project. Thanks to Doug Stewart and Kate

McGivney for sharing their insights on central coast chum salmon and Brooke Davis and two

anonymous reviewers for feedback on earlier drafts. This research was funded by the Pacific

Salmon Foundation's Salmon Watersheds Program.

Literature cited

Brown, G.S., Baillie, S.J., Thiess, M.E., Bailey, R.E., Candy, J.R., Parken, C.K., and Willis, D.M.

2019. Pre-COSEWIC review of southern British Columbia Chinook Salmon (*Oncorhynchus*

tshawytscha) conservation units, Part I: background. DFO Can. Sci. Advis. Sec. Res. Doc.

2019/11: vii + 67 pp. Available from

http://publications.gc.ca/collections/collection_2019/mpo-dfo/fs70-5/Fs70-5-2019-011-

eng.pdf [accessed 1 December 2019].

Carwardine, J., Martin, T.G., Firn, J., Reyes, R.P., Nicol, S., Reeson, A., Grantham, H.S., Stratford,

D., Kehoe, L., and Chadès, I. 2019. Priority Threat Management for biodiversity

conservation: A handbook. J. Appl. Ecol. 56(2): 481–490. doi:10.1111/1365-2664.13268.

Cave, J.D., and Gazey, W.J. 1994. A preseason simulation model for fisheries on Fraser River

sockeye salmon (*Oncorhynchus nerka*). Can. J. Fish. Aquat. Sci. **51**(7): 1535–1549.

doi:10.1139/f94-153.

Challenger, W., Mochizuki, T., English, K., and Bychkov, Y. 2018. North and Central Coast Salmon

Database and Analysis System User Manual. Sidney, BC. Available from

https://salmonwatersheds.ca/libraryfiles/lib_449.pdf [accessed 1 December 2019].

Chen, Y., Chen, L., and Stergiou, K.I. 2003. Impacts of data quantity on fisheries stock

assessment. Aquat. Sci. 65(1): 92–98. doi:10.1007/s000270300008.

Clark, R., Eggers, D., Munro, A., Fleischman, S., Bue, B., and Hasbrouck, J. 2014. An evaluation of

the percentile approach for establishing sustainable escapement goals in lieu of stock

productivity information. Alaska Dep. Fish Game, Fish. Manuscr. No. 14–06. Anchorage,

AK. doi:10.4027/amdlfs.2016.08.

Connors, B.M., Pickard, D.C., Porter, M., Farrell, G., Bryan, K., Casely, S., and Huang, S. 2013.

Skeena Salmon Conservation Unit Snapshots. Report prepared by ESSA Technologies Ltd.

for the Pacific Salmon Foundation. v + 70p. Available from

https://salmonwatersheds.ca/libraryfiles/lib_420.pdf [accessed 1 December 2019].

Connors, K., Hertz, E., Jones, E., Honka, L., Kellock, K., and Alexander, R. 2019. The Nass Region:

Snapshots of salmon population status. The Pacific Salmon Foundation, Vancouver, BC.

Available from https://salmonwatersheds.ca/libraryfiles/lib_453.pdf [accessed 1

December 2019].

Connors, K., Jones, E., Kellock, K., Hertz, E., Honka, L., and Belzile, J. 2018. BC Central Coast: A

Snapshot of Salmon Populations and Their Habitats. The Pacific Salmon Foundation,

Vancouver, BC. Available from https://salmonwatersheds.ca/library/lib_442/ [accessed 1

December 2019].

COSEWIC. 2016. COSEWIC Assessment and Status Report on the Coho Salmon *Oncorhynchus*

kisutch, Interior Fraser population, in Canada. Committee on the Status of Endangered

Wildlife in Canada Ottawa, ON. Available from https://www.canada.ca/en/environment-

climate-change/services/species-risk-public-registry/cosewic-assessments-status-

reports/coho-salmon-interior-fraser-2016.html [accessed 1 December 2019].

COSEWIC. 2017. COSEWIC assessment and status report on the Sockeye Salmon *Oncorhynchus*

nerka, 24 Designatable Units in the Fraser River Drainage Basin, in Canada. Committee on

the Status of Endangered Wildlife in Canada. Ottawa, On. Available from

https://www.sararegistry.gc.ca/virtual_sara/files/cosewic/srSockeyeSalmon2017e.pdf

[accessed 1 December 2019].

Cousens, N.B.F., Thomas, G.A., Swann, C.G., and Healey, M.C. 1982. A Review of Salmon

Escapement Estimation Techniques. Can. Tech. Rep. Fish. Aquat. Sci. **1108**: 1–128.

DFO. 2015. Wild Salmon Policy Biological Status Assessment for Conservation Units of Interior

Fraser River Coho Salmon (*Oncorhynchus kisutch*). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep.

2015/022. Available from https://waves-vagues.dfo-mpo.gc.ca/Library/364851.pdf

[accessed 1 December 2019].

DFO. 2016. Integrated Biological Status of Southern British Columbia Chinook Salmon

(*Oncorhynchus tshawytscha*) Under the Wild Salmon Policy. DFO Can. Sci. Advis. Sec. Sci.

Advis. Rep. 2016/042. Available from https://waves-vagues.dfo-

mpo.gc.ca/Library/40595419.pdf [accessed 1 December 2019].

DFO. 2018a. The 2017 Fraser sockeye salmon (*Oncorhynchus nerka*) integrated biological status

re-assessment under the Wild Salmon Policy. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep.

2018/017. Available from https://waves-vagues.dfo-mpo.gc.ca/Library/40712163.pdf

[accessed 1 December 2019].

DFO. 2018b. Northern BC Salmon Integrated Fisheries Management Plan. Fisheries and Oceans

Canada, Vancouver, BC. Available from https://waves-vagues.dfo-

mpo.gc.ca/Library/40797168.pdf [accessed 1 December 2019].

DFO. 2019. Wild Salmon Policy 2018-2022 Implementation Plan: Annual Report 2018-2019.

Available from https://www.pac.dfo-mpo.gc.ca/fm-gp/species-especes/salmon-

saumon/wsp-pss/annual-annuel/2018-2019-eng.html [accessed 1 December 2019].

Dorner, B., Catalano, M.J., and Peterman, R.M. 2017. Spatial and temporal patterns of

covariation in productivity of Chinook salmon populations of the northeastern Pacific

Ocean. Can. J. Fish. Aquat. Sci. **75**(7): 1082–1095. doi:10.1139/cjfas-2017-0197.

Dorner, B., Peterman, R.M., and Su, Z. 2009. Evaluation of performance of alternative

management models of Pacific salmon (Oncorhynchus spp.) in the presence of climatic

change and outcome uncertainty using Monte Carlo simulations. Can. J. Fish. Aquat. Sci.

66(12): 2199–2221. doi:10.1139/F09-144

English, K.K. 2016. Review of escapement indicator streams for the North and Central Coast

salmon monitoring program. Report prepared by LGL Limited for the Pacific Salmon

Foundation and Fisheries and Oceans Canada. Available from

https://salmonwatersheds.ca/libraryfiles/lib_440.pdf [accessed 1 December 2019].

English, K.K., Bailey, R.E., and Robichaud, D. 2007. Assessment of Chinook Salmon returns to the

Fraser River watershed using run reconstruction techniques, 1982-04. Can. Sci. Advis. Sec.

Res. Doc. 2007/020: vi + 76. Available from

https://www.psf.ca/sites/default/files/RES2007_020_e.pdf [accessed 1 December 2019].

English, K.K., Peacock, D., Challenger, W., and Mochizuki, T. 2016. North and Central Coast

Salmon Escapement, Catch, Run Size and Exploitation Rate Estimates for each Salmon

Conservation Unit for 1954-2014. Report prepared by LGL Limited for the Pacific Salmon

Foundation and Fisheries and Oceans Canada. Available from

https://salmonwatersheds.ca/libraryfiles/lib_435.pdf [accessed 1 December 2019].

English, K.K., Peacock, D., Challenger, W., Noble, C., Beveridge, I., Robichaud, D., Beach, K.,

Hertz, E., and Connors, K. 2018. North and Central Coast Salmon Escapement, Catch, Run

Size and Exploitation Rate Estimates for each Salmon Conservation Unit for 1954-2017.

Report prepared by LGL Limited for the Pacific Salmon Foundation. Available from

https://salmonwatersheds.ca/library/lib_451/ [accessed 1 December 2019].

Fisheries and Oceans Canada. 2005. Canada's Policy for Conservation of Wild Pacific Salmon.

Vancouver, BC. Available from https://www.pac.dfo-mpo.gc.ca/fm-gp/species-

especes/salmon-saumon/wsp-pss/policy-politique/index-eng.html [accessed 1 December

2019].

Fleischman, S.J., Catalano, M.J., Clark, R.A., and Bernard, D.R. 2013. An age-structured state-

space stock recruit model for Pacific salmon Oncorhynchus spp. Can. J. Fish. Aquat. Sci. **70**:

401–414. doi:10.1139/cjfas-2012-0112.

Foster, K.R., Vecchia, P., and Repacholi, M.H. 2000. Science and the Precautionary Principle.

Science **288**(5468): 979–981. Available from https://www.jstor.org/stable/3075092

[accessed 1 December 2019].

Holt, C., Davis, B., Dobson, D., Godbout, L., Luedke, W., Tadey, J., and Van Will, P. 2018.

Evaluating Benchmarks of Biological Status for Data-limited Populations (Conservation

Units) of Pacific Salmon, Focusing on Chum Salmon in Southern BC. DFO Can. Sci. Advis.

Sec. Res. Doc. 2018/011: ix + 87. Available from http://www.dfo-mpo.gc.ca/csas-

sccs/Publications/ResDocs-DocRech/2018/2018_011-eng.html [accessed 1 December

2019].

Holt, C.A., Brooke, D., and Cox-Rogers, S. 2016. Adapting benchmarks of biological status for

persistent changes in productivity and variability in exploitation history with a focus on

data-limited populations (Conservation Units) of Pacific salmon in north and central BC.

Draft final report for the PSC. Can. Sci. Advis. Secr. Res. Doc. 86. Available from

https://www.psc.org/fund-project/adapting-benchmarks/ [accessed 1 December 2019].

Holt, C.A., Cass, A., Holtby, L.B., Riddell, B.E., and Holtby, B. 2009. Indicators of status and

benchmarks for Conservation Units in Canada's Wild Salmon Policy. Can. Sci. Advis. Sec.

Res. Doc. 2009/058: viii+74. Available from

https://www.psf.ca/sites/default/files/339096.pdf [accessed 1 December 2019].

Holt, C.A., and Folkes, M.J.P. 2015. Cautions on using percentile-based benchmarks of status for

data-limited populations of Pacific salmon under persistent trends in productivity and

uncertain outcomes from harvest management. Fish. Res. **171**: 188–200.

doi:10.1016/j.fishres.2015.01.002.

Holt, C.A., and Ogden, A. 2013. Software for Assessing Status of Conservation Units under

Canada's Wild Salmon Policy: Instructional Manual. Can. Tech. Rep. Fish. Aquat. Sci. *In*

Prep.

Holt, C.A., and Peterman, R.M. 2006. Missing the target: uncertainties in achieving

938   management goals in fisheries on Fraser River, British Columbia, sockeye salmon

939   (*Oncorhynchus nerka*). Can. J. Fish. Aquat. Sci. **63**(12): 2722–2733. doi:10.1139/F06-155

940   Holt, C.A., and Peterman, R.M. 2008. Uncertainties in population dynamics and outcomes of

941   regulations in sockeye salmon (*Oncorhynchus nerka*) fisheries: implications for

942   management. Can. J. Fish. Aquat. Sci. **65**(7): 1459–1474. Available from

943   https://www.nrcresearchpress.com/doi/full/10.1139/F08-053 [accessed 1 December

944   2019].

945   Holtby, L.B., and Ciruna, K.A. 2007. Conservation Units for Pacific salmon under the Wild

946   Salmon Policy. Can. Sci. Advis. Sec. Res. Doc. 2007/070: viii+74. Available from

947   http://www.dfo-mpo.gc.ca/csas-sccs/publications/resdocs-docrech/2007/2007_070-

948   eng.htm [accessed 1 December 2019].

949   IUCN. 2017. Guidelines for Using the IUCN Red List - Categories and Criteria. Version 6.1. July

950   2006. Iucn 11(February): 60. Available from

951   https://www.iucnredlist.org/resources/redlistguidelines [accessed 13 April 2020].

952   Kehler, D.G., Myers, R.A., and Field, C.A. 2002. Measurement error and bias in the maximum

953   reproductive rate for the Ricker model. Can. J. Fish. Aquat. Sci. **59**(5): 854–864.

954   doi:10.1139/f02-060.

Kope, R. 2006. Cumulative effects of multiple sources of bias in estimating spawner-recruit

parameters with application to harvested stocks of Chinook salmon (*Oncorhynchus*

tshawytscha). Fish. Res. **82**(1–3): 101–110. doi:10.1016/j.fishres.2006.08.016.

Korman, J., and English, K. 2013. Benchmark Analysis for Pacific Salmon Conservation Units in

the Skeena Watershed. A report prepared for the Pacific Salmon Foundation, Vancouver,

BC. Available from https://salmonwatersheds.ca/libraryfiles/lib_318.pdf [accessed 1

December 2019].

Korman, J., Peterman, R.M., and Walters, C.J. 1995. Empirical and theoretical analyses of

correction of time-series bias in stock-recruitment relationships of sockeye salmon

(*Oncorhynchus nerka*). Can. J. Fish. Aquat. Sci. **52**(10): 2174–2189. doi:10.1139/f95-810.

Malick, M.J., and Cox, S.P. 2016. Regional-Scale Declines in Productivity of Pink and Chum

Salmon Stocks in Western North America. PLoS One **11**(1): e0146009.

doi:10.1371/journal.pone.0146009.

Martin, T.G., Kehoe, L., Mantyka-Pringle, C., Chades, I., Wilson, S., Bloom, R.G., Davis, S.K.,

Fisher, R., Keith, J., Mehl, K., Diaz, B.P., Wayland, M.E., Wellicome, T.I., Zimmer, K.P., and

Smith, P.A. 2018. Prioritizing recovery funding to maximize conservation of endangered

species. Conserv. Lett. **11**(6): 1–9. doi:10.1111/conl.12604.

Office of the Auditor General of Canada. 2004. 2005 Report of the Commissioner of the

Environment and Sustainable Development. In Fisheries and Oceans Canada — salmon

stocks, habitat, and aquaculture. Chapter 5 [online]. Available from http://www.oag-

bvg.gc.ca/internet/docs/c20041005ce.pdf [accessed 8 April 2019].

Peacock, S.J., Connors, B.M., Krkošek, M., Irvine, J.R., and Lewis, M.A. 2014. Can reduced

predation offset negative effects of sea louse parasites on chum salmon? Proc. R. Soc. B

Biol. Sci. **281**: 20132913. doi:10.1098/rspb.2013.2913.

Peacock, S.J., and Holt, C.A. 2012. Metrics and sampling designs for detecting trends in the

distribution of spawning Pacific salmon (*Oncorhynchus* spp.). Can. J. Fish. Aquat. Sci. **69**(4):

681–694. doi:10.1139/f2012-004.

Peterman, R.M. 2018. Continuous learning, teamwork, and lessons for young scientists. ICES J.

Mar. Sci. **76**(1): 28–40. doi:10.1093/icesjms/fsy141.

Peterman, R.M., and Dorner, B. 2012. A widespread decrease in productivity of sockeye salmon

(*Oncorhynchus nerka*) populations in western North America. Can. J. Fish. Aquat. Sci. **69**(8):

1255–1260. doi:10.1139/F2012-063.

Peterman, R.M., Pyper, B.J., and Grout, J. a. 2000. Comparison of parameter estimation

methods for detecting climate-induced changes in productivity of Pacific salmon

(*Oncorhynchus* spp.). Can. J. Fish. Aquat. Sci. **57**(1): 181–191. doi:10.1139/cjfas-57-1-181.

Price, M.H.H., Darimont, C.T., Temple, N.F., and MacDuffee, S.M. 2008. Ghost runs:

management and status assessment of Pacific salmon (*Oncorhynchus* spp.) returning to

British Columbia's central and north coasts. Can. J. Fish. Aquat. Sci. **65**: 2712–2718. doi:

10.1139/F08-174.

Price, M.H.H., English, K.K., Rosenberger, A.G., MacDuffee, M., and Reynolds, J.D. 2017.

Canada's Wild Salmon Policy: an assessment of conservation progress in British Columbia.

Can. J. Fish. Aquat. Sci. **74**(10): 1507–1518. doi:10.1139/cjfas-2017-0127.

PSC. 2019. Treaty Between the Government of Canada and the Government of the United

States of America Concerning Pacific Salmon. Pacific Salmon Commision, Vancouver, BC.

Available from http://www.psc.org/ [accessed 1 December 2019].

Punt, A.E. 1992. Selecting management methodologies for marine resources, with an

illustration for Southern African hake. South African J. Mar. Sci. **12**(1): 943–958.

doi:10.2989/02577619209504754.

Pyper, B.J., and Peterman, R.M. 1998. Comparison of methods to account for autocorrelation in

correlation analyses of fish data. Can. J. Fish. Aquat. Sci. **55**(9): 2127–2140.

doi:10.1139/cjfas-55-9-2127.

Ricker, W.E. 1954. Stock and recruitment. J. Fish. Res. Board Canada **11**(5): 559–623. Available

from https://www.nrcresearchpress.com/doi/abs/10.1139/f54-039 [accessed 1 December

2019].

Salmon Watersheds Program - Pacific Salmon Foundation. 2019. Salmon Watersheds Data

Library. Available from http://data.salmonwatersheds.ca/data-library/ [accessed 1

December 2019].

Scheuerell, M.D. 2016. An explicit solution for calculating optimum spawning stock size from

Ricker's stock recruitment model. PeerJ **4**(1): e1623. doi:10.7717/peerj.1623.

Schnute, J.T., Boers, N., and Haigh, R. 2015. PBSmapping: Mapping Fisheries Data and Spatial

Analysis Tools. Available from https://cran.r-project.org/package=PBSmapping [accessed 1

December 2019].

Schnute, J.T., and Richards, L.J. 1995. The influence of error on population estimates from

catch-age models. Can. J. Fish. Aquat. Sci. **52**(10): 2063–2077. doi:10.1139/f95-800.

Staton, B.A., Catalano, M.J., Connors, B.M., Coggins, L.G., Jones, M.L., Walters, C.J., Fleischman,

S.J., and Gwinn, D.C. 2020. Evaluation of methods for spawner-recruit analysis in mixed-

stock Pacific salmon fisheries. Can. J. Fish. Aquat. Sci. (*in press*): 1–43. doi:10.1139/cjfas-

2019-0281.

Subbey, S., Devine, J.A., Schaarschmidt, U., and Nash, R.D.M. 2014. Modelling and forecasting

stock–recruitment: current and future perspectives. ICES J. Mar. Sci. **71**(8): 2307–2322.

Szerlong, R.G., and Rundio, D.E. 2008. A statistical modeling method for estimating mortality

and abundance of spawning salmon from a time series of counts. Can. J. Fish. Aquat. Sci.

65: 17–26. doi:10.1139/F07-141.

Walsh, J.C., Connors, K., Hertz, E., Kehoe, L., Martin, T.G., Connors, B., Bradford, M.J.,

Freshwater, C., Frid, A., Halverson, J., Moore, J.W., Price, M.H.H., and Reynolds, J.D. (2020).

Prioritizing conservation actions for Pacific salmon in Canada. J. Appl. Ecol. (*in press*): 1–37.

doi:10.1101/2020.02.03.931691.

Walters, C., English, K., Korman, J., and Hilborn, R. 2019. The managed decline of British

Columbia's commercial salmon fishery. Mar. Policy **101**: 25–32.

doi:10.1016/j.marpol.2018.12.014.

Walters, C.J. 1985. Bias in the Estimation of Functional Relationships from Time Series Data.

Can. J. Fish. Aquat. Sci. **42**(1): 147–149. doi:10.1139/f85-018.

Walters, C.J. 1986. Adaptive Management of Renewable Resources. MacMillan Publishing

Company, New York.

Walters, C.J., and Ludwig, D. 1981. Effects of Measurement Errors on the Assessment of Stock-

Recruitment Relationships. Can. J. Fish. Aquat. Sci. **38**(6): 704–710.

Wetzel, C.R., and Punt, A.E. 2011. Performance of a fisheries catch-at-age model (Stock

Synthesis) in data-limited situations. Mar. Freshw. Res. **62**(8): 927. doi:10.1071/mf11006.

Zabel, R.W., and Levin, P.S. 2002. Simple assumptions on age composition lead to erroneous

conclusions on the nature of density dependence in age-structured populations. Oecologia

133(3): 349–355. doi:10.1007/s00442-002-1051-0.

Tables

Table 1. Summary of common steps in run reconstructions (Figure 2) and associated assumptions and potential biases that we investigated.

Step in run recon-struction	Description	Assumption	Potential biases	Associated factor(s) investigated in this study
(1) Expansion Factor I	Infills observed spawner abundances in indicator streams to account for indicator streams that were not monitored in a given year.	The contribution of each indicator-stream population to total escapement is constant within a decade.	May be biased if contributions are changing due to, for example, changes in capacity or productivity that differ among populations. Under low monitoring coverage, the magnitude of expansion is greater and thus we expect that any potential bias would be larger.	→ Diverging capacities of populations through time → Declining monitoring coverage
(2) Expansion Factor II	Expands spawner abundances from (1) to include spawners in non-indicator streams.	The contribution of non-indicator-stream populations to total escapement is constant within a decade.	May be biased if contributions are changing due to, for example, changes in capacity or productivity that differ between populations in indicator and non-indicator streams.	→ Diverging capacities of populations through time → Declining monitoring coverage

				→ Number of streams and proportion that are indicators
(3) Expansion Factor III	Expands spawner abundances from (2) to account for observer efficiency and for populations that are never monitored, yielding the estimated total spawner abundance for the CU.	The proportion of total spawner abundance estimated in (2) is known, constant over time, and independent of spawner abundance.	May be biased if observer efficiency is not known or poorly estimated, if survey methods change over time (e.g., a weir to overhead counts), or if observation bias varies with total spawner abundance.	→ Bias in the observation of spawners (under the same value of Expansion Factor III) → Declining monitoring coverage
(4) Catch assignment to CUs	Catch from PFMAs is assigned to CUs in proportion to the spawner abundance for that CU.	Fish caught in an PFMA were destined to spawn in streams that flow into that PFMA.	Over- or under-estimation of catch due to different run timing among CUs that flow into the same PFMA (if information on run timing is uncertain, unavailable, or not incorporated into run-reconstruction models).	→ Bias in the observation of catch

| (5) Calculating recruitment using age-at-return | The total return to the CU is assigned to brood years based on the proportion of fish returning at different ages. | Often, annual age-at-return data are not available for each CU, and so age-at-return is assumed to be constant over time, using the average of available data. | Variability in brood year recruitment will be underestimated if there is high inter-annual variability, or temporal changes, in age-at-return that is not accounted for in assessments. | → Inter-annual variability in age-at-return |

1048

1049   *Table 2. Summary of factors that we investigated in sensitivity analyses to determine their impact on bias in status*

1050   *assessments.*

Factor	Scenario	Details
Number of populations and proportion that spawn in indicator streams	Central coast chum*	35 populations, 15 (43%) indicator streams
	Small/Low	10 populations, 3 (30%) indicator streams
	Small/High	10 populations, 8 (80%) indicator streams
	Large/Low	140 populations, 42 (30%) indicator streams
	Large/High	140 populations, 119 (85%) indicator streams
Monitoring coverage	Constant	**Indicator**: historical 76% with no change
		Non-indicator: historical 72% with no change
	Observed decline*	**Indicator**: 76% with a change of -5% over the last 26 years
		Non-indicator: 72% with a change of -67% over the last 26 years
Declines in capacity†	0	All spawning populations have stable capacities
	25	25% of populations severe and 25% moderate declines, and 50% stable capacity
	50	50% of populations severe and 50% moderate declines in capacity
	100	100% of populations display severe declines in capacity
Bias in observation of spawners	Range in bias from $\bar{\delta}$ = -1.6, which would correspond to the maximum value of Expansion Factor III that has been applied (F''' = 5.0,; English et al. 2018) to $\bar{\delta}$ = 0 in increments of 0.2 (*default value $\bar{\delta}$ = -0.4, corresponding to F'''=1.5)	
Bias in observation of catch	Range in bias from $\bar{\chi}$ = -1.0 (63% underestimation) to $\bar{\chi}$ = 1.0 (271% overestimation) in increments of 0.2 (*default value $\bar{\chi}$ = 0).	

Interannual variability in age-at-return	Range in variability from $\bar{\omega}$ = 0.2 to $\bar{\omega}$ = 1.6 in increments of 0.2 (*default value $\bar{\omega}$ = 0.8; Figure S4)

*Default values. †The default values for decline in capacity did not correspond exactly to the

scenarios considered in the sensitivity analyses, but were based on habitat assessments for

central coast (21% of populations having severe declines and 29% of populations having

moderate declines (Connors et al. 2018)).

Figure captions

Figure 1. Illustration of the WSP status assessment framework (adapted from Holt et al. 2009).

We focused on the geometric mean spawner abundance (metric, blue) under the spawner

abundance indicator. This metric was assessed against two types of benchmarks: spawner-

recruitment (SR) and percentile (see Figure 2). Faded boxes represent other types of metrics

and indicators that may be included in integrated status assessments but were beyond the

scope of what we considered.

Figure 2. a) benchmarks are the 50th (horizontal green line) and 25th (horizontal red line)

percentiles of historical spawner abundance (points). The current spawner abundance is

calculated as the geometric mean spawner abundance over the most recent generation (4

years, blue points and line). b) -recruitment benchmarks are based on the shape of the Ricker

relationship (solid line) fit to data on spawner abundance (x-axis) and corresponding

recruitment (catch + escapement, y-axis). The upper and lower benchmarks are 80% S_{MSY}

(green) and S_{GEN} (red), respectively. S_{GEN} is defined as the spawner abundance that leads to S_{MSY}

(grey) in one generation in the absence of fishing mortality. Under both types of benchmarks,

the current spawner abundance in the example shown is above the upper benchmark, and this

CU would be assessed as 'green'. c) The calculation of benchmarks and benchmarks requires

run reconstruction to expand observed spawners abundances, assign catch to CUs, and

calculate recruitment (Table 1).

Figure 3. Our study focused on central coast chum salmon Conservation Units (CUs) as a case

study for how common run-reconstruction assumptions affect the assessment of biological

status. salmon CUs (green; different CUs shaded differently in central coast inset) relatively

large do not correspond to the Pacific Fishery Management Areas (PFMAs; white or light blue

shaded regions) for which catch is reported. Therefore, assumptions have to be made when

assigning catch to CUs that may result in under- or over-estimation of catch. Map produced

using PBSmapping (Schnute et al. 2015) with data from (Martin Huang, pers. comm.

<Martin.Huang@dfo-mpo.gc.ca>).

Figure 4. Schematic of the simulation model comprised of submodels for population dynamics

(including harvest), observation, assessment, and performance. Adapted from Holt et al. (2016).

Figure 5. The two target harvest-rate cases we simulated were (1) a simple Harvest Control Rule

(eq. (4); solid line) with parameters estimated from historical harvest rates and total return

from five central coast chum CUs (grey points), (2) a constant high target harvest rate of $ht' =$

0.6 (dotted line).

Figure 6. Estimated status according to the spawner-recruitment (SR) benchmarks (left) and the

percentile benchmarks (right), over the true status for each of two cases (high productivity and

an HCR (a-b) and low productivity and high harvest rates (c-d)). Grey cells indicate pessimistic

misclassifications, which may lead to overly conservative management actions, and black cells

indicate optimistic misclassifications, which may lead to overly risky management actions.

Coloured cells indicate correct classifications for red, amber, and green zones, respectively.

Figure 7. Relative bias in spawner-recruitment (SR) benchmarks (a), percentile benchmarks (b),

and current spawner abundance (S_{AVG}; black) (median $\pm$ interquartile range among 4000 MC

simulations) for high productivity and a harvest control rule (HCR) and the low-productivity

high-harvest case.

Figure 8. The effect of monitoring coverage (no change and decline; Table 2) and the

percentage of spawning populations with severe declines in capacity (x-axis) on performance

measures under the base case of high productivity and HCR. (a-b) The proportion of simulations

with correct green, amber, or red status or pessimistic misclassifications (grey) and optimistic

misclassifications (black) under the SR benchmarks (a) and percentile benchmarks (b). (c-d) The

percent relative bias (median $\pm$ interquartile range among 4000 MC simulations) in the current

spawner abundance (S_{AVG}; black) and lower and upper benchmarks (red and green,

respectively) under the SR benchmarks (c) and the percentile benchmarks (d). See Online

Supplement for results under the low-productivity high-harvest case.

Figure 9. The effect of observation bias in the number of spawners (x-axis) on performance

measures under the base case. (a-b) The proportion of simulations with correct green, amber,

or red status or pessimistic misclassifications (grey) and optimistic misclassifications (black)

under the SR benchmarks (a) and percentile benchmarks (b). (c-d) The percent relative bias

(median $\pm$ interquartile range among 4000 MC simulations) in the current spawner abundance

(S_{AVG}; black circle) and lower and upper benchmarks (red and green, respectively) under the SR

benchmarks (c) and the percentile benchmarks (d). The asterisk in (a-b) and grey zone in (c-d)

indicate the default parameter value of δ = -0.4, and the bias that matches the Expansion Factor

III of $F''' = 1.5$ applied in all simulations. See Online Supplement for results under the low-

productivity high-harvest case (results were similar).

Figure 10. The effect of observation bias in catch (x-axis) on performance measures under the

base case. (a) The proportion of simulations with correct green, amber, or red status or

pessimistic misclassifications (grey) and optimistic misclassifications (black) under the SR

benchmarks. (b) The percent relative bias (median $\pm$ interquartile range among 4000 MC

simulations) in the current spawner abundance (S_{AVG}; black circle) and lower and upper

benchmarks (red and green, respectively) under the SR benchmarks. The asterisk in (a) and grey

zone in (b) indicate the default parameter value of $\chi = 0$. Observation bias in catch ranges from

63% underestimation ($\chi = -1$) to 271% overestimation ($\chi = 1$). See Online Supplement for

results under the low-productivity high-harvest case.

1126

Appendices

Appendix A: Table of default parameter values

Table A1. Default values for parameters in the simulation model that were used unless otherwise specified (e.g., in

sensitivity analyses). See text for further explanation of the values and the Online Supplement for details of

estimation for those based on raw data. For parameters that were part of sensitivity analyses, the range in

parameter values that was explored is highlighted.

Submodel	Parameter	Symbol	Default value	Range	Ref.
General	Years over which the simulation is run	t : return year y : brood year	50	-	
	Number of indicator streams	I	15	$1 - 41$	1
	Number of non-indicator streams	J	20	1 - 100	1
Population dynamics	Productivity	$\bar{a}$	1.40	-	2
		σ_a^2	0.22	-	2
	$a_i \sim N(\bar{a}, \sigma_a^2)$				
	Density dependence (initial)	$\mu_{b,I}$	7.95	-	2
		$\sigma_{b,I}^2$	2.07	-	2
	$b_{i,t=0} = 1/S_{MAX_i}$				
	$\log S_{MAX_i} \sim N(\mu_{b,I}, \sigma_{b,I}^2)$	$\mu_{b,J}$	6.95	-	2
		$\sigma_{b,J}^2$	1.39	-	2

	Temporal autocorrelation	τ	0.422	-	2
	Variance in recruitment deviates within spawning populations	σ_v^2	1.28	-	2
	Correlation among spawning populations in recruitment deviates	ρ	0.46	-	2
	Average proportions for age-at-maturity	$\overline{p}_g$	$\overline{p}_3 = 0.23$ $\overline{p}_4 = 0.64$ $\overline{p}_5 = 0.13$	-	3
	Interannual variability in age-at-maturity	$\overline{\omega}$	0.8	-	4
	Maximum target harvest rate	h'_{MAX}	0.42	-	5
	Shape parameter for that harvest rule	m	1.13×10^{-5}		5
	Standard deviation in outcome uncertainty around harvest rate	σ_h	0.13	-	5
Observation	Lognormal bias in observed spawners	$\overline{\delta}$	- 0.4	-1.6 to 0.0	6
	Lognormal variance in observed spawners	σ_δ^2	0.25	-	4

Lognormal bias in observed catch	$\bar{\chi}$	0	-1.0 to 1.0	
Lognormal variance in observed catch	σ^2_χ	0.04	-	4
Variability in observed age-at-return	$\overline{\omega}_r$	0.1	-	4

References:

1. Based on the range of indicator and non-indicator streams reported in the Pacific Salmon Explorer

(www.salmonexplorer.ca) for the eight central coast chum CUs.

2. Calculated from river-level stock-recruitment data for central coast chum CUs. See Online Supplement for

details.

3. From the NCCDBV2 (Challenger et al. 2018).

4. Same as assumed in Holt et al. (2018) for south-coast chum salmon.

5. Calculated from CU-level harvest rates and total return size, from the Salmon Watersheds Data Library (Salmon

Watersheds Program - Pacific Salmon Foundation 2019). See Online Supplement for details.

6. Based on expert opinion. Expansion Factor of F''' = 1.5 in Table A3-A4 of English et al. (2016).

1143