1

Clumpy coexistence in phytoplankton: The role of functional similarity in community assembly

Caio Graco-Roza [1, 4]*, Angel M Segura [2], Carla Kruk [3], Patrícia Domingos [1], Janne Soininen [4], Marcelo Manzi Marinho[1]

[1] Laboratory of Phytoplankton of Ecology and Physiology, Department of Plant Biology, University of Rio de Janeiro State, Rua São Francisco Xavier 524—PHLC Sala 511a, 20550-900, Rio de Janeiro, Brazil. Telephone: +552123340822

[2] Modelización Estadística de Datos e Inteligencia Artificial (MEDIA) CURE-Rocha, Universidad de la República, Uruguay.

[3] Sección Limnología, IECA, Facultad de Ciencias, Universidad de la República, Uruguay.

[4] University of Helsinki, Department of Geosciences and Geography, PO Box 64, FI00014, Helsinki, Finland. *correspondence author: caiogracor@gmail.com

Abstract

Emergent neutrality (EN) suggests that species must be sufficiently similar or sufficiently

different in their niches to avoid interspecific competition. Such a scenario results in a transient

pattern with clumps and gaps of species abundance along the niche axis (*e.g.*, represented by

body size). From this perspective, clumps are groups of coexisting species with negligible

fitness differences andstochastic abundance fluctuations. Plankton is an excellent model

system for developing and testing ecological theories, especially those related to size structure

and species coexistence. We tested EN predictions using the phytoplankton community along

the course of a tropical river considering (i) body size structure, (ii) functional clustering of

species in terms of morphology-based functional groups (MBFG), and (iii) the functional

similarity among species concerning their functional traits. Two main clumps in the body size

axis (clump I and II) were conspicuous through time and were detected in different stretches

of the river. Clump I comprised medium-sized species from the MBFGs IV, V, and VI while

clump II included large-bodied species from the MBFGs V and VI. Pairwise differences in

species biovolume correlated with species functional similarity when the whole species pool

was considered, but not among species within the same clump. Although clumps comprised

multiple MBFGs, the dominant species within the clump belonged always to the same MBFG.

Also, within-clump species biovolume increased with functional distinctiveness considering

both seasons and stretches, except the lower course. These results suggest that species within

clumps behave in a quasi-neutral state, but even minor shifts in trait composition may affect

species biovolume.Our findings point that EN belongs to the plausible mechanisms explaining

community assembly in river ecosystems.

Keywords: emergent neutrality; functional distinctiveness; functional similarity; species

coexistence

Introduction

Understanding the mechanisms promoting species coexistence and shaping community structure has been a long-standing goal in community ecology. The former idea that the number of coexisting species is limited by the number of growth-limiting resources or niche dimensions (Gause 1936, Hardin 1960) and its derivate idea, "*the paradox of the plankton*" (Hutchinson 1957), have been widely explained in terms of endogenous and exogenous spatio-temporal mechanisms (Roy and Chattopadhyay 2007). Trait-based approaches are useful to test this matter due to their potential to generalize patterns beyond species' identity, especially because traits influence the species' ability to acquire resources and persist through environmental changes (McGill et al. 2006, Díaz et al. 2013, 2016). Nonetheless, the niche-based theory proposes that the environment filters community composition through species' ecological requirements, which can be perceived through species' traits. Also, intra- and inter-specific interactions potentially drive community assembly, in local communities (Götzenberger et al. 2012). In contrast, the more recent neutral theory suggests that diversity results from random dispersal, speciation, and extinction rates with no role of niche differences in species coexistence (Hubbell 2001). This type of dynamics should then result in a random distribution of functional traits along environmental gradients (Kraft et al. 2008, Cornwell and Ackerly 2009).

More recently, it was shown that community organization is driven by eco-evolutionary processes such as speciation and nutrient uptake kinetics resulting in groups comprising different species with similar ecological requirements (Gravel et al. 2006, Scheffer and van Nes 2006, Hubbell 2006). This finding led to the 'emergent neutrality hypothesis' (EN; HOLT, 2006) that has been supported by observational studies, e.g., for phytoplankton from brackish waters (Segura et al. 2011), birds from the North of Mexico (Thibault et al. 2011)

and beetles at the global scale (Scheffer et al. 2015). EN suggests that species must be

sufficiently similar, and thus, behave neutrally, or different enough in their niches to avoid

competition. Such a scenario would result in species-rich aggregations or clumps along the

niche axis (Scheffer and van Nes 2006, Vergnon et al. 2009, Fort et al. 2010). Modelling

studies have shown that such predictions apply for both steady environmental conditions

(Fort et al. 2010), and also fluctuating resource conditions (Sakavara et al. 2018). Empirical

evidence about EN is still scarce, however (Scheffer et al. 2018).

The clumpy pattern arises from the exceedingly slow displacement rate of species under

intense competition, that is, species within the same clump overlap in their niche such that

the displacement rate of competing species is similar to the competition at the intraspecific

level, leading to stochastic fluctuations in species abundances through time (Scheffer et al.

2018). Thus, the number of clumps corresponds to the number of species to be expected to

stably coexist at equilibrium, but the identity of the dominant species is expected to be

random among the clump residents. However, the assignment of species to clumps is

challenged by the fact that trait differences among species are continuous (Villéger et al.

2008) and the threshold to include a species within a clump varies with the statistical

approach that is applied (Segura et al. 2011, D'Andrea et al. 2019). Therefore, given this

methodological limitation, it is difficult to state empirically whether species behave neutrally

within clumps (i.e., when the strength of interspecific interactions equals the intraspecific

interactions) or if results are an artefact of clump construction.

Zooming in on the uniqueness of trait combinations of species, i.e. functional distinctiveness,

within clumps may advance our comprehension of biotic interactions and move towards a

measurable value of similarity at which species coexistence is driven stochastically.

Functional distinctiveness reflects the non-shared functions among species within a given

species pool (Violle et al. 2017), mirroring the concept of functional similarity (Pavoine et al.

2017). However, functional distinctiveness is not directly linked to functional similarity at the

pairwise level (Coux et al. 2016, Ricotta et al. 2016, Violle et al. 2017). For example, two

species may be equally distinct, i.e. the degree to which a species differs from all the others

within the species pool concerning their functional traits, and still not be similar in their trait

composition at a pairwise level (Coux et al. 2016). This suggests that both pairwise functional

similarity and group-based functional distinctiveness are complementary metrics to assess the

role of trait combination in community assembly. To this end, phytoplankton communities

are useful for biodiversity theory testing due to their species-rich communities, rapid

responses (in human time-scales) and well-characterized relationships between morphology

and physiological and ecological responses (Litchman and Klausmeier 2008, Kruk and

Segura 2012, Litchman et al. 2012).

Body size is considered a master ecological trait and it is often used to characterize species

niche differences (Downing et al. 2014). In phytoplankton, the body size is related to

physiology and life-history (Litchman and Klausmeier 2008), photosynthetic processes

(Marañón 2008), nutrient uptake kinetics (Litchman et al. 2010) and other eco-evolutionary

processes, e.g. the relationship among predation rates, nutrient uptake and organisms body

size (Sauterey et al. 2017). Although body size may relate to different processes, using a

single trait as a proxy for niche differences may not evidence species differences generated

by hidden/unknown niche axes (i.e. ecological dimensions of the niche) and impair the

understanding of clumpy patterns (Barabás et al. 2013, D'Andrea et al. 2018). The use of

multiple traits emerges as a powerful tool to disentangle phytoplankton functional structure

and evaluate competing hypotheses (Reynolds et al. 2014, Chen et al. 2015, Bortolini and

Bueno 2017, Aquino et al. 2018). Morphology-based functional groups (MBFG)

classification of phytoplankton species (Kruk et al. 2010) is a multidimensional combination

of morphological traits that cluster organisms into seven groups with similar physiology and

ecological responses, potentially overcoming the limitations of using a single trait dimension

only. Assessing the functional distinctiveness of species within the same functional cluster

(e.g., clumps, MBFGs) could help to study the existence of functional equivalence (i.e.,

neutrality) among species. Overall, the functional similarity among species is a useful tool to

compare species in a multidimensional space, particularly because the environment may filter

different functional traits across space and time (D'Andrea et al. 2020).

Rivers are highly heterogeneous systems characterized by a continuous water flow that

affects the ecosystem's morphology (e.g., meandering), sedimentation patterns, organisms'

dispersal, and more specifically the phytoplankton abundance and distribution (Reynolds and

Descy 1996, Wetzel 2001). Several theories, e.g., the River Continuum (Vannote et al. 1980)

and Flood Pulse (Junk et al. 1989) concepts explain the longitudinal distribution and

abundance of riverine phytoplankton communities. However, an explicit study of

communities' body size structure and species coexistence under EN in riverine ecosystems is

lacking. For example, phytoplankton species should attain higher biomass at the middle

reaches or in the upper reaches of low-gradient stretches (Descy et al. 2017). Also,

competition rates vary along the river course because water turbulence reduces the likelihood

of biotic interactions (Reynolds et al. 1994), meaning that clumpy coexistence may not be

observed in riverine phytoplankton. Alternatively, if functional trait combinations of species

within the local species pool result from eco-evolutionary processes (Scheffer et al. 2015),

the clumpy pattern should also be apparent in riverine phytoplankton communities. Here, we

push forward three hypotheses to be tested in a tropical river by investigating phytoplankton

community size structure both seasonally and spatially. We expect that:

H$_1$ – There are peak aggregations of species abundance (i.e., clumps) along the body size

axis of phytoplankton in the river that remain constant across space and time as a result of

eco-evolutionary processes.

H$_2$ – Pairwise-differences in species abundances increase with functional dissimilarity at the

community-level but not at the clump level because species within the same clump behave in

a quasi-neutral state. Thus, the dominance within clump varies stochastically between species

as fitness differences are negligible.

H$_3$ – Species abundance increases with functional distinctiveness with respect to other species

within the clumps. Although abundance fluctuates stochastically at the pairwise level, the

number of species bearing similar trait combinations may affect the likelihood of the

interactions within clumps. Therefore, species with the most distinct trait combinations

concerning their clump peers are less likely to share the same ecological requirements, and by

consequence, attain higher abundance.

Methods

Study area

Samples were taken monthly at nine stations along the Piabanha river between May 2012 and

April 2013. Piabanha river is in the tropical region of Brazil and has a drainage basin of

approximately 4500 km² (Figure 1). The headwater is on Petrópolis at 1546m altitude and

drains to the medium valley of Paraíba do Sul river crossing three cities and with agricultural

activities in their watershed. We set three river stretches (lower, medium, and upper courses)

based on the location of steep slopes on the river elevation profile (Figure 1). Data from two

meteorological stations (Bingen and Posse; Figure 1), located in the upper and lower courses

of the river, were used to measure rainfall. We analyzed meteorological data up to three days

before each sampling campaign. We then classified seasons as a dry season (May - October)

and a wet season (November – April) based on the rainfall data.

Figure 1. Map of the study area. The watershed area of the Piabanha river showing the river

course (blue line), the meteorological stations Bingen and Posse, and the sampling sites are

coloured according to river stretches (white circles = upper course, blue circles = medium

course, red circles = lower course). The vertical dotted red line in the elevational profile

figure indicates the locations of steep slopes used to define the boundaries of the river

stretches.

Sampling and sample analysis

In the field, we measured water temperature (°C), dissolved oxygen (DO, mg L^{-1}), and

turbidity by a multiparameter probe sonde (YSI model 600 QS). Water discharge (WD, m^3

s^{-1}) was measured with the SonTek RiverSurveyor – M9. Furthermore, water samples were

taken and kept frozen (one or 2 weeks) until the laboratory analysis for ammonium ($N \cdot NH_4^+$,

mg L^{-1}), nitrate ($N \cdot NO_3^-$, mg L^{-1}), nitrite ($N \cdot NO_2^-$, mg L^{-1}), total phosphorus (TP, mg L^{-1})

and soluble reactive phosphorus (SRP, mg L^{-1})(Figure 2). Ammonium, nitrite, and nitrate

were summed up and are expressed as dissolved inorganic nitrogen (DIN, mg L^{-1}). The water

samples were filtered (except for total phosphorus analysis) using borosilicate filters

(Whatman GF/C), and nutrient concentrations were measured following APHA (2005). A

complete description of the spatial and seasonal patterns of the environmental variables

measured in the Piabanha river can be found in Graco-Roza et al. (2020).

Phytoplankton samples

Subsurface samples of phytoplankton were collected with a bottle of 200 mL and fixed with

Lugol. In the laboratory, phytoplankton species were identified, and population densities

were estimated under an inverted microscope (Olympus CKX41) (Utermöhl 1958). At least

100 individuals of the most abundant species were counted in each sample (Lund et al. 1958,

Uhelingher 1964). Biovolume (mm^3 L^{-1}) of phytoplankton species was estimated by

multiplying the density of each population (ind. L^{-1}) by the average individual volume of the

species (V, $\mu m^3 org^{-1}$). The volume of each species was estimated by measuring geometrical

dimensions and approximating to defined geometrical forms following Hillebrand et al.

(1999). Geometrical dimensions were measured in 20 organisms from each species (when

possible) and the average was used to characterize individual bod size (volume). We recall

that biovolume represents the biomass density and volume is an organism's trait. Species'

surface area (S, μm^2) was estimated, the maximum linear dimension (MLD, μm) was

measured, and the presence or absense of aerotopes, mucilage, flagella, and siliceous

exoskeletal structures was noted in each species (Figure 2). We then used the volume and

surface area of the species to estimate the individual surface volume ratio (SV). Species were

then classified into MBFG according to Kruk et al. (2010), based on the above mentioned

morphological traits. This classification included the following seven groups: (I) small

organisms with high SV, (II) small, flagellated organisms with siliceous exoskeletal

structures, (III) large filaments with aerotopes, (IV) organisms of medium size lacking

specialised traits, (V) flagellates unicells with medium to large size, (VI) non-flagellated

organisms with siliceous exoskeletons and (VII) large mucilaginous colonies. For further

details on MBFG classification, we refer to Kruk et al. (2010) or Segura et al. (2013a). The

information on the traits measured for each species, and the classification into MBFGs can be

found in the Table S1.

Figure 2. Study design. Sketch diagram showing the steps from data acquisition to

hypothesis testing, and the grouping variables used to analyse the data. Water samples were

taken from the river for the estimation of local environmental variables, and phytoplankton

qualitative and quantitative analysis. Phytoplankton species were measured based on

approximate geometrical forms for the estimation of individual volume and surface. Other

traits were also measured during the quantitative and qualitative analysis. Samples were

divided into seasonal and spatial groups prior to the analysis. Species biovolume were used

along with its traits to test the relationship between trait and environment, and highlight the

functional traits. Only body size was used to test the presence of significant clumps (H_1),

while the other functional traits were used to test the effects of functional similarity within

significant clumps (H_2-H_3).

Traits-environment relationship

We tested the relationship between morphological traits and the environmental variables

using a three-table ordination (RLQ) combined with a fourth-corner analysis (Dray et al.

2014) (Figure 2). Both RLQ and fourth-corner methods require the information from three

tables; a) a data frame including the measurements of environmental variables across the

sampling sites (R table), b) a matrix containing species abundances or occurrences across the

sampling sites (L table), and c) a data frame comprising the trait values for each species (Q

table). Also, both methods rely on the analysis of the fourth-corner matrix, crossing the

information between tables R and Q, weighted by table L. The RLQ analysis (Legendre et al.

1997) provides ordination scores to summarize the joint structure among the three tables, but

it does not allow the identification of traits or environmental variables contributing

significantly to the structure. The fourth corner method (Dolédec et al. 1996) tests the

significance of bivariate associations between each trait and environmental variables but

disregards the covariance among traits or environmental variables. Here, we combined the

RLQ analysis with the fourth corner method by applying the fourth corner method to the

output of the RLQ analysis instead of the original raw values (Dray et al. 2014). By doing

this, we summarized the main patterns in the multivariate space and tested the global

significance of the trait–environment relationships using the S_{RLQ} multivariate statistic and

the fourth corner sequential testing procedure (Dray and Legendre 2008). Applying the fourth

corner method in the output of the RLQ determines (a) the relationship between individual

traits and RLQ environmental scores (a.k.a environmental gradients, and (b) the relationship

between environmental variables and RLQ traits scores (a.k.a. trait syndromes)(Dray et al.

2014).

Before applying the RLQ method, we first log-transformed ($\log_{10} x+1$) species biovolume,

species traits (SV, MLD, and V), and environmental variables (except pH and temperature).

A correspondence analysis (Benzécri 1973) was performed on the L table using the function

dudi.ca from 'ade4', and a Hill-Smith analysis (Hill and Smith 1976) on the R and Q tables

separately using the function dudi.hill from 'ade4'. We used Hill-smith analysis because both

R and Q table included categorical or binary variables. The RLQ analysis was conducted in

the output of the ordinations using the function rlq from 'ade4'. We tested the significance of

the joint structure among the RLQ tables using the fourth-corner method with a stepwise

permutation procedure of 999 permutations using the function rlq.fourthcorner from 'ade4'.

The null hypothesis that given fixed traits, species abundances are independent of

environmental conditions was evaluated by permuting sites (rows of tables L or R) while

keeping the species traits (table Q) fixed. The null hypothesis that given fixed environmental

conditions, species abundances are independent of functional traits was evaluated by

permuting species (columns of table L or rows of table Q) while the environmental conditions

(R table) were kept fixed (Dray et al. 2014). Rejecting both null hypotheses imply that tables

R, L, and Q are significantly linked. Because the fourth-corner analysis explores one trait and

one environmental variable at a time, multiple statistical tests are performed simultaneously

increasing the probability of type I error (i.e. false significant associations), thus we adjusted

p-values for multiple testing using the false discovery rate method (Benjamini and Hochberg

1995). We divided the value of the fourth-corner correlation by the square-root of the first

eigenvalue of the correspondence analysis of the L matrix, which is the maximum possible

value (Peres-Neto et al. 2017).

Clumpy patterns

To test for the existence of peak aggregations of species biovolume along the body size axis

of phytoplankton - **H₁**, we analyzed the community structure in each season (dry and wet) and

river stretches (upper, medium, and lower course) (Figure 2). First, the individual volume of

species was log-transformed (log_2) and used as the main niche axis ($X = log_2$ volume)

following Segura et al. 2011. Hence, we divided the niche axis into equally spaced segments

(one segment per unit log_2 volume) and for each segment (j), we estimated the Shannon

entropy (H) using the biovolume of the observed species (Fort et al. 2010, Segura et al.

2011). The entropy index was defined as:

$$H_j = \sum_{i=1}^{n} p_i \, log_2(p_i) \tag{1}$$

here p_i is the fraction of biovolume of species i in the community of n species. Finally, we

tested the significance of the entropy (H) by comparing the observed H against an expected

uniform distribution under the null hypothesis of homogeneous H. For this, we created 1000

communities by sampling the volume of species from a random uniform distribution bounded

by observed individual volumes. Then, each species had a biovolume assigned to it, which

was taken from randomization of the observed biovolume matrix, keeping both the empirical

species rank-biovolume pattern and total biovolume in the sample. For each segment, the

observed H was compared with the distributions of H generated under the null hypothesis,

with significance defined according to standard 5% criterion (Fort et al. 2010, Segura et al.

2011). Finally, we considered a significant segment or two consecutive significant segments

as a clump.

Functional dissimilarity

To test whether differences in species biovolume increases with functional dissimilarity – **H₂**,

we first calculated the functional dissimilarity and the differences in biovolume among pairs

of species using the whole community and using only the species from the significant clumps

separately (Figure 2). The functional dissimilarity was obtained by calculating Gower's

general dissimilarity coefficient on all the species functional traits, that is, all the traits that

showed a significant ($p < 0.05$) relationship with the environmental gradients in the fourth

corner method. The dissimilarity coefficient was estimated using the function gowdis

from'FD'. We used Gower's dissimilarity (*Gd*) because it can handle mixed variable types

(continuous, binary, ordinal, and categorical traits). *Gd* defines a distance value d_{jk} between

two species as

299
$$d_{jk} = \frac{1}{N} \sum_{i=1}^{n} \left| \frac{(x_{ij} - x_{ik})}{\max(x_i) - \min(x_i)} \right| \tag{2}$$

where, N is the number of functional traits considered, x_{ij} the value of trait i for species j, and

x_{ik} the value of the trait i for species k. Therefore, $Gd = d_{jk}$ = functional dissimilarity. We thus

tested **H₂** by conducting Mantel tests with 999 randomizations on the matrices of functional

dissimilarity and differences in biovolume using the function mantel from 'vegan'. We

performed the Mantel test considering: a) all species present in a given season or river

stretch, and b) separately for the species of each significant clump that were present in a

given season or river stretch.

Functional distinctiveness (F_{Dist})

To test whether species biovolume increases with functional distinctiveness at the clump-

level – **H₃**, we estimated the functional distinctiveness (F_{Dist}) as the Euclidean distance of a

species to the average trait position (centroid) in the multidimensional functional space for

the set of species of each of the significant clumps using the equations proposed by Anderson

(2006) (Figure 2). First, we applied a Principal Coordinates Analysis (PCoA) in the species-

by-traits data table using Gower's dissimilarity (*Gd*) and obtained species coordinates in the

functional space using all the axes from the PCoA. Hence, F_{Dist} was calculated as

$$F_{Dist} = \sqrt{\Delta^2\left(u_{ij}^+, c_{i'j'}^+\right) - \Delta^2\left(u_{ij}^-, c_{i'j'}^-\right)} \qquad (3)$$

where Δ^2 is the squared Euclidean distance between u_{ij}, the principal coordinate for the *j*th

species in the *i*th clump, and c_i, the coordinate of the centroid for the *i*th clump. The super-

scripted '+' and '-' indicate the real and imaginary parts respectively (see Anderson 2006, for

details). We did not weight the clump-centroid by species biovolume because it would

artificially give higher distinctiveness for less abundant species and bias our analysis.

Besides, we calculated F_{Dist} using only species from the significant clumps and normalized

the F_{Dist} to range between zero and one by dividing the actual F_{Dist} values by the F_{Dist} of the

most distinct species of the clump. We tested the **H₃**, by modelling the relationship between

species biovolume and F_{Dist} using linear models. We used $\log_{10}$ biovolume as the dependent

variable, with F_{Dist} and Clump (i.e. the clump to which a species belong) as the independent

variables for each season and river stretch separately.

Statistical analyses

Statistical analyses were performed on R v.4.0.4 (R Core Team 2020) using the packages

'ade4' v.1.7.16 (Chessel et al. 2004, Dray and Dufour 2007, Dray et al. 2007, Bougeard and

Dray 2018, Thioulouse et al. 2018), 'FD' v.1.0.12 (Laliberte et al. 2010, Laliberté et al.

2014), the suite of packages 'tidyverse' v.1.3.0 (Wickham et al. 2019), and the package

'vegan' v.2.5.7 (Oksanen et al. 2020).

Results

Our samples included 150 species that were classified in six (MBFG I, III, IV, V, VI, and

VII) from the seven MBFGs based on their functional traits (Table 1). MBFGs IV, V, and VI

included 87% of the total number of species. Species from MBFG IV included filamentous,

colonial, and unicellular species ranging from 21 μm³ to 8181 μm³ lacking specialized

morphological traits (e.g. flagella, siliceous exoskeletal structures). MBFG V comprised

unicellular flagellated species ranging in volume from 31μm³ to 31864 μm³, and MBFG VI

included unicellular and chain-forming species with a siliceous exoskeletal body that ranged

in volume from 48μm³ to 19045μm³.

Table 1. Distribution of species among the morphological-based functional groups.

MBFG	Number of species	Representative taxa
I	9	*Chroococcales* sp., *Chroococcus* sp.
III	3	*Limnothrix* sp.
IV	60	*Pseudanabaena limnetica, Pseudanabaena catenata*
V	13	*Euglena* sp., *Cryptomonas* sp.
VI	57	*Cymbella* sp., *Synedra* sp.
VII	8	*Dictyosphaerium* sp.
Total	150	

Regarding the trait-environment relationship, the first two RLQ axes preserved well the

variance of the ordinations and explained altogether 84.37 % of the variation, with 66.71%

corresponding to the first axis alone. The S_{RLQ} statistic indicated a significant global

relationship between trait syndromes and environmental gradients ($r = 0.18$, p-value < 0.01).

Mainly, the first trait syndrome (RLQ axis 1_{Trait}) correlated significantly with the first

environmental gradient (RLQ axis $1_{Environment}$; $r = 0.17$, p < 0.01) while the second trait

syndrome (RLQ axis 2_{Trait}) correlated significantly with the second environmental gradient

(RLQ axis $2_{Environment}$; $r = 0.12$, $p < 0.01$). Yet, there was no significant relationship between

the first trait syndrome and the second environmental gradient, nor between the second trait

syndrome and the first environmental gradient (Table 2).

Table 2. Combined fourth-corner-RLQ analysis to test the relationship between

functional syndromes (RLQ axis$_{Trait}$) and environmental gradients (RLQ axis$_{Environment}$).

	RLQ axis 1_{Trait}	RLQ axis 2_{Trait}
	Pearson's r (adjusted p-value)	
RLQ axis $1_{Environment}$	**0.17 (< 0.01)**	0.00 (1.00)
RLQ axis $2_{Environment}$	0.00 (1.00)	**0.12 (< 0.01)**

Results of the RLQ showed that most of the biovolume-based variation in trait syndromes

were related to the flow regime of the river Piabanha. The first RLQ axis summarised the

spatial gradient in the environmental conditions, specifically the increase in turbidity,

temperature, pH, and water discharge from the upper to the lower courses while the second

RLQ axis summarised the seasonal gradient with the smaller nutrient concentrations and

higher water turbidity in the wet season (Figure 3A). Noteworthy, the spatial and seasonal

gradients in abiotic conditions coupled with the distribution of species from different

MBFGs. The MBFGs I, III, and IV attained higher abundances in the upper course,

contrasting with MBFG VI that showed the highest abundances at the lower course (Figure

3B). Regarding the seasonal gradient, MBFG V had higher abundances during the dry season

(Figure 3B). Indeed, the fourth-corner method showed that the first trait syndrome correlated

positively with turbidity ($r = 0.16$, $p = 0.01$), temperature ($r = 0.14$, $p = 0.02$), pH ($r = 0.13$, p

= 0.03), water discharge ($r = 0.13$, $p = 0.03$) and total phosphorus ($r = 0.12$, $p = 0.04$), and

negatively with the upper course ($r = -0.21$, $p < 0.01$; Figure 3C). Besides, the second trait

syndrome correlated negatively with dissolved inorganic nitrogen ($r = -0.11$, $p = 0.04$) and

positively with turbidity ($r = 0.13$, $p = 0.01$; Figure 3C). For the spatial environmental

gradient, there was a positive correlation with the species volume ($r = 0.15$, $p = 0.01$) and the

presence of siliceous exoskeletal structures ($r = 0.22$, $p < 0.01$), and a negative correlation

with surface volume ratio ($r = -0.19$, $p < 0.01$; Figure 3D). For the seasonal environmental

gradient, there was a positive significant correlation with species maximum linear dimension

($r = 0.12$, $p = 0.01$), and a negative significant correlation with the presence of flagella ($r = -0.12$, $p = 0.01$; Figure 3D).

0.12, $p = 0.01$; Figure 3D).

376
Figure 3. Results of the (A, B) RLQ ordination and (C, D) hypothesis testing through

fourth-corner analysis. A) The relationships between species traits and environmental

variables. B) The distribution of species in the functional space. Each point in the ordination

plot represents the position of a species modelled according to its traits on RLQ axes 1 and 2.

The black lines connect the species to the centroid of its morphology-based functional groups

- MBFG. Colours represent MBFGs, C) The correlation between species traits and the

environmental gradients (RLQ axis $_{environment}$), and D) the relationship between environmental

variables and the trait syndromes (RLQ axis $_{trait}$). The grey boxes in C and D indicate

significant relationships, and the values within the boxes indicate the Pearson's *r*. Aer,

aerotopes; Muc, mucilage; Si, siliceous exoskeletal structures; Fla, flagella; MLD, maximum

linear dimension; SV, surface volume ratio; V, volume; WD, water discharge; T,

temperature; Turb, turbidity; SRP, soluble reactive phosphorus; DIN, dissolved inorganic

nitrogen; DO, dissolved oxygen; TP, total phosphorus; WS, wet season; DS, dry season; UC,

upper course; MC, medium course; LC, lower course.

Overall, species volume ranged from 4.19 µm³ to 31864 µm³ totalling 14 equally spaced

segments (S) of volume along the niche axis. From the 14 segments, three of them showed

significant ($p < 0.05$) entropy values, specifically S9, S13, and S14. This resulted in a

biovolume aggregation (i.e. clumps) in two regions of the niche axis considering both

seasonal (Figure 4; Column 1) and spatial categories (Figure 5; Column 1). The first clump

included 24 species from the MBFGs IV, V, and VI at the range of S9 (512µm³ - 1024µm³),

particularly eight species from MBFG IV (e.g. *Pseudanabaena catenata* and *P. limnetica*),

four species from the MBFG V (e.g. *Strombomonas* sp., *cf Cryptomonas* sp.), and 12 species

from MBFG VI (e.g*., Fragillaria capuccina* var. gracilis, *Achnantes cf. rupestoides*). The

second clump (hereafter Clump II) included six species at the range of S14, being two species

from MBFG V (i.e. *Euglena* sp.) and four species from MBFG VI (e.g. *Pinnularia* sp.,

Synedra sp.). However, during the wet season, the S13 also had significant entropy values

and six more species from MBFG VI (e.g., *Achnantes inflata*, *Cymbella* sp.) were included

in the clump II.

Species from the same MBFG tended to cluster in the functional space even if they belonged

to different clumps (Figure 4 – 5; Column 2). Yet, species within the same MBFG did not attain

the highest abundance in more than one clump (Figure 4 – 5; Column 3). The mean biovolume

of species within clumps differed between seasons, but the identity of the most abundant

species did not vary (Figure 4 – Column 3). *Pseudanabaena* sp. (spp. 028) and *P. catenata*

(spp. 012) had the highest biovolumes of clump I at both dry (Figure 4 – A2) and wet (Figure

4 – B3) seasons. Within clump II, *Synedra* sp. (spp. 080) attained the highest biovolume during

the dry season (Figure 4 – A3) while *Cymbella* sp. (spp. 051) had the highest biovolume in the

wet season (Figure 4 – B3).

Figure 4. Seasonal distribution of phytoplankton biovolume along the body size axis, the

ordination of species from the significant size segments (S) in the functional space, and

the mean biovolume of the five most abundant species of each significant size segment

during the (A) dry and (B) wet seasons of the Piabanha river, RJ. (1) Stem plots show size

distribution in the sampling sites of the river Piabanha. Each stem represents a species with its

body size (in $\log_2$) plotted on the abscissa and the mean biovolume plotted on the ordinate. The

red dotted line indicates the entropy value of each size segment (i.e., unit of $\log_2$ volume) and

the asterisk highlights the significant entropy values tested through 1000 randomizations. (2)

The species of the corresponding significant size-segment are ordinated in the functional space.

The size of the circles represents the species contribution to the total biovolume of the size

segment, the black line connects species to the centroid (see equation 3), and the number in the

center of the clump indicates the size class it encompasses. (3) Bar plots show the biovolume

of the five most abundant species from each significant size segment. Species are coloured

according to their morphology-based functional groups (MBFG). The code for species can be

found in the supplementary material, Table S1.

Regarding the river stretches, only the clump I had significant entropy values for species from

the S9 (Figure 5 – A1), with *Pseudanabaena* sp. (spp. 028) and *P. catenata* (spp. 012)

contributing most of the biovolume (Figure 4 – A3). At the medium course, both clumps I and

II had significant entropy values with *Pseudanabaena* sp. (spp. 028) attaining the highest

biovolume within clump I, and *Synedra* sp. (spp. 080) attaining the highest biovolume within

clump II (Figure 5 – C3). At the lower course, only clump II had significant entropy values at

the S14 with *Synedra* sp. (spp. 080) as the most representative species (Figure 4 – C3).

Figure 5. Spatial distribution of phytoplankton biovolume along the body size axis, the ordination of species from the significant size segments (S) in the functional space, and the mean biovolume of the five most abundant species of each significant size segment at the (A) upper, (B) medium, and (C) lower courses of the Piabanha river, RJ. (1) Stem plots show size distribution in the sampling sites of the river Piabanha. Each stem represents a species with its body size (in log_2) plotted on the abscissa and the mean biovolume plotted on

the ordinate. The red dotted line indicates the entropy value of each size segment (i.e., unit of

$\log_2$ volume) and the asterisk highlights the significant entropy values tested through 1000

randomizations. (2) The species of the corresponding significant size-segment are ordinated in

the functional space. The size of the circles represents the species contribution to the total

biovolume of the size segment, the black line connects species to the centroid (see equation 3),

and the number in the center of the clump indicates the size class it encompasses. (3) Bar plots

show the biovolume of the five most abundant species from each significant size segment.

Species are coloured according to their morphology-based functional groups (MBFG). The

code for species can be found in the supplementary material, Table S1.

Mantel tests showed that species' pairwise differences in biovolume correlated with functional

dissimilarity irrespectively of the season or river stretch when the whole community was

analyzed (Table 3). The correlation was highest during the wet season (Mantel $r = 0.23$, p $<$

0.01) and at the lower course (Mantel $r = 0.26$, p < 0.01). For the clump-level pairwise

differences, we only found a significant correlation at the upper course (Mantel $r = 0.23$, p $<$

0.02; Table 3). In contrast, functional distinctiveness at clump level presented a significant

positive relationship for both the dry season ($\beta = 16.44$, $R^2 = 0.40$, p < 0.01) and the wet season

($\beta = 18.32$, $R^2 = 0.40$, p < 0.01), and also for the upper ($\beta = 5.68$, $R^2 = 0.40$, p < 0.01) and

medium ($\beta = 14.46$, $R^2 = 0.34$, p $= 0.02$) courses (Table 4), indicating that species with the

most distinct trait combinations within the clumps also attain the highest biovolume.

Essentially, such pattern was observed only for the species within clump I, except during the

wet season where species from clump II also showed a significant positive relationship ($\beta = $ -

17.83, p $= 0.02$; Table 4)

Table 3. Mantel correlation results. Mantel correlation between the differences in species

biovolume and functional dissimilarity for the whole community, and separately for the species

within significant clumps (Figure 3 – 4) along the seasons (dry and wet) and river stretches

(upper, medium, and lower courses). Species number of each stratum (whole community or

clumps) are given. The relationships were tested for significance using 999 permutations,

whenever possible.

Seasons and river stretches	Stratum	Number of species	Mantel r	p-value (permutations)
Dry season				
	Whole community	135	**0.25**	**< 0.01 (999)**
	Clump I	22	0.13	0.09 (999)
	Clump II	4	-0.14	0.71 (23)
Wet Season				
	Whole community	123	**0.29**	**< 0.01 (999)**
	Clump I	20	0.06	0.24 (999)
	Clump II	11	-0.16	0.69 (999)
Upper course				
	Whole community	100	**0.17**	**< 0.01 (999)**
	Clump I	17	**0.23**	**0.02 (999)**
Medium course				
	Whole community	135	**0.21**	**< 0.01 (999)**
	Clump I	23	0.06	0.22 (999)
	Clump II	5	-0.16	0.65 (119)

Lower course

Whole community	122	**0.33**	**<0.01 (999)**	
Clump II	5	-0.23	0.75 (119)	

Note:	Bold values indicate significant correlations (p < 0.05).

Table 4. Linear model results. Regression parameters of the relationship between species biovolume and functional distinctiveness at the clump level. The coefficients are shown along with the p-values of each independent variable.

	Dependent variable: $\log_{10}$ Biovolume				
Independent variables	Dry season	Wet season	Upper course	Medium course	Lower course
F_{Dist}	16.44	18.33	5.68	14.46	9.65
	p < 0.01	p < 0.01	p > 0.01	p = 0.02	p = 0.06
Clump II	12.29	17.73		11.61	5.77
	p = 0.30	p = 0.01		p = 0.29	p = 0.48
$F_{Dist} \times$ Clump II	-11.90	-17.83		-11.12	-4.57
	p = 0.34	p = 0.02		p = 0.34	p = 0.59
Intercept	-18.39	-20.420	-7.26	-16.64	-12.52
	p < 0.01	p < 0.01	p < 0.01	p < 0.01	p = 0.01
Observations	26	31	17	28	26
Adjusted R^2	0.41	0.39	0.40	0.34	0.55
F Statistic	6.75 (df = 3; 22)	7.58 (df = 3; 27)	11.87 (df = 1; 15)	5.70 (df = 3; 24)	11.01 (df = 3; 22)

Discussion

Present results showed that (i) the clumps in body size are a conspicuous feature of

phytoplankton community structure in riverine systems across seasons and river stretches; (ii)

species within clumps showed a random distribution of biomass concerning their pairwise

functional dissimilarity, but not at the whole-community level; and (iii) species biovolume

generally increases for species far apart from the centroid of multivariate trait space (i.e.,

functional distinctiveness) within clumps. Altogether these results support the Emergent

neutrality hypothesis and show studying species beyond pairwise interactions helps to explain

the biomass distribution of functionally similar species, paving the way to analyze intra-

clumps trait distributions.

Multimodal aggregation of species biovolume along body size axis only points to the

integration of niche-based processes and neutrality driving community assembly (Vergnon et

al. 2009), supporting H_1. Alternative hypothesis such as pure neutrality (Hubbell 2001) or

high dimensional hypothesis (HDH, Clark et al. 2007) are not supported by present results

because pure neutrality predicts a uniform distribution of species biovolume and traits along

the niche axis (Hubbell 2001), and the HDH does not predict any particular trait distribution

(Vergnon et al. 2009, Ingram et al. 2018).

One alternative theory that is likely to explain clumpy aggregations is Holling's textural

hypothesis (Holling 1992), which suggests that multimodal species size distribution is the

result of environmental constraints. Our results do not support textural hypothesis, as river

stretches and seasons were markedly different in hydrology, nutrient concentrations, and

other relevant descriptors of riverine landscapes fluxes, but that was not reflected in the stable

clumpy size structure of the phytoplankton registered in the present study (Figure 4). The

stability found in the clumps agree with empirical results registered in Segura et al. (2011,

2013b) and theoretical findings on the location of clumps (Fort et al. 2009).

However,morphological trait composition of species in the different clumps reflected

different environmental templates. The dominant species from the clump I belonged to

MBFG IV (Figure 4–5) and presented highest biovolume under low-flow and high nutrient

conditions (Figure 3), which is in line with previous findings for this MBFG (Chen et al.

2015). Within the second clump (II), the most abundant species belonged to MBFG VI and

had their highest biovolume under high-flow and turbid conditions, in line with the ecology

of silicious organisms (diatoms) able to cope with turbulent environments (Bortolini and

Bueno 2017). The empirical evidence is consistent with studies from coastal and estuarine

environments (Segura et al. 2011, 2013b) and are in line with recent modelling results

suggesting that clumpy patterns arise in environments subjected to resources fluctuation

(Sakavara et al. 2018), such as rivers. The trade-off between resources among competing

species (Tilman 1982), which is a required ingredient for the emergence of clumps should be

further explored.

The analysis of multiple trait dimensions, combining morphology based functional groups

(MBFG) and quantitative distance metrics helped to describe the changes in species traits

within clumps. We found that the species from the same clump are distributed across multiple

MBFGs but only species from the same MBFG attain the highest biovolume within a clump,

reinforcing that body size is a good proxy for niche differences of the species (Blanckenhorn

2000, Gallego et al. 2019). MBFGs helped to detect differences at a finer degree because they

synthesize multiple trait dimensions as suggested previously to understand community

organization (D'Andrea et al. 2018). Given that species within the same MBFG share similar

ecological strategies (Kruk et al. 2010, Kruk and Segura 2012), under EN premises they

should also perform similarly (Scheffer et al. 2018). This is in line with the significant

functional dissimilarity observed at the whole community level but not for each clump

separately, agreeing with H_2. However, the effects of traits in species' fitness are context-

dependent, whereas the use of traits for assigning MBFGs is static, therefore, testing the

significance of traits given the observed environmental conditions might help to unveil

community assembling processes at an even finer degree (Kremer et al. 2017).

We also outlined the role of functional similarity in community assembly by studying the

effects of functional distinctiveness on species biovolume at clump level. Within the clump I,

species biovolume increased with functional distinctiveness, but this pattern was weaker

within clump II (Table 4). It may be that such patterns stem from the fact that phytoplankton

growth-rate decreases with body size while increases with surface volume ratio, providing

that populations of smaller species (such as in clump I) are less sensitive to losses by flushing

rates (Kruk et al. 2010). On the other hand, large-sized species have often an elongated shape

that provides advantage under turbulent conditions with low light availability (Reynolds et al.

1994). Differently, aerotopes and mucilage are useful to reach the surface in deep stratified

lakes but are not key in small rivers or streams were turbulent fluxes dominates and this traits

are not useful to recuperate the position in the water column. Furthermore, in rivers, large-

bodied phytoplanktonic species are often randomly introduced from different habitats (e.g.

periphyton or epiphyton) (Wang et al. 2014, Descy et al. 2017), which has also been found

true for the Piabanha river especially under high flow conditions (Graco-Roza et al. 2020).

This mechanism can help to explain the weak relationship between functional distinctiveness

and biovolume within clump II, which explain the niche overlap found in large-sized species

as the result of inmigration and emigration out of the pelagic zone.

Emergent neutrality results from eco-evolutionary processes that lead species selection

towards a limited number of functional groups (Scheffer and van Nes 2006). This implies that

the clumps observed here are not likely a result of competitive exclusion at the Piabanha

river, but a convergent evolution of competing species over time (Macarthur and Levins

1967). Therefore, even when the competition rates are relaxed due to sufficient nutrient

supply, some other limiting factors that are not consumed by biotic organisms such as heat

energy or turbulence determine species biovolume. Looking at species differences at a high-

order level (instead of pairwise differences) helped to detect the effects of trait composition

on the biovolume disrtribution in quasi-neutral clumps. In fact, our results showed that it is

possible to predict the biovolume of species within clumps, but only when immigration from

different habitats are relaxed and biotic interactions are more likely to occur. Therefore, our

findings partially agree with **H₃** - there is a positive relationship between species abundance

and species functional distinctiveness within clumps, but the environmental conditions seem

to play a key role in the outcome.

There are some possibly influential aspects in our study design that should be discussed.

First, given that phytoplankton communities have short generation time (Reynolds 2006), the

resolution (monthly) of our data makes it difficult to us to capture the turnover in species

abundance rank at the finest possible scale. It may be that studying the daily or even weekly

variation in abundance rank would help us disentangle the so-called stochastic abundance

fluctuations (Caracciolo et al. 2021). However, it has been shown that the study of

phytoplankton community processes on a monthly to yearly time scale helps to understand

the long-term ecological and evolutive dynamics of communities (Segura et al. 2011).

Secondly, considering intraspecific trait variability allows one to disentangle species

responses to environmental variation (Wong and Carmona 2021). Here, we only assessed trait

at species level and it is possible that comparing the overlap in trait values between species

might unveil the quasi-neutral relationship between pairs or clumps of species.

In summary, we provided evidence of both neutral and niche mechanisms driving planktonic

community assembly and support the view that emergent neutrality is a likely mechanism to

explain species coexistence in an open and environmentally heterogeneous ecosystem. The

use of MBFG classification and functional space to describe species within clumps revealed

that under the same size range, species with a greater degree of functional similarity

unpredictably alternate their dominance. The position and dominance of the clumps were

related to the environmental conditions, but the biovolume of species within the clumps was

better predicted by functional distinctiveness than by pairwise functional similarity. This

addresses the difficulty to avoid the ghost of hidden niches (Barabás et al. 2013) and also

provides evidence from multiple angles that point to EN as a plausible mechanism in shaping

species coexistence in riverine landscapes.

Acknowledgements

CGR PhD scholarship was funded by Fudação de Apoio a Pesquisa do Estado do Rio de

Janeiro (FAPERJ), by Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior (CAPES), and by Ella and Georg Ehrnrooth foundation. MMM was partially

supported by CNPq (303572/2017-5).

References

Anderson, M. J. 2006. Distance-based tests for homogeneity of multivariate dispersions.

Biometrics 62:245–253.

APHA, A. 2005. WEF: Standard Methods for the Examination of Water and Wastewater:

Centennial Edition. Washington, DC.

Aquino, C. A. N., J. C. Bortolini, C. C. R. Favaretto, N. Y. Sebastien, and N. C. Bueno. 2018.

Functional phytoplankton distribution predicts the environmental variability between

two subtropical rivers. Revista Brasileira de Botanica 41:835–847.

Barabás, G., R. D'Andrea, R. Rael, G. Meszéna, and A. Ostling. 2013. Emergent neutrality or

hidden niches? Oikos.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological) 57:289–300.

Benzécri, J.-P. 1973. L'analyse des données. Dunod Paris.

Blanckenhorn, W. U. 2000. The Evolution of Body Size: What Keeps Organisms Small? The Quarterly Review of Biology 75:385–407.

Bortolini, J. C., and N. C. Bueno. 2017. Temporal dynamics of phytoplankton using the morphology-based functional approach in a subtropical river. Revista Brasileira de Botanica 40:741–748.

Bougeard, S., and S. Dray. 2018. Supervised Multiblock Analysis in {R} with the {ade4} Package. Journal of Statistical Software 86:1–17.

Caracciolo, M., G. Beaugrand, P. Hélaouët, F. Gevaert, M. Edwards, F. Lizon, L. Kléparski, and E. Goberville. 2021. Annual phytoplankton succession results from niche-environment interaction. Journal of Plankton Research 43:85–102.

Chen, N., L. Liu, Y. Li, D. Qiao, Y. Li, Y. Zhang, and Y. Lv. 2015. Morphology-based classification of functional groups for potamoplankton. Journal of Limnology 74:559–571.

Chessel, D., A.-B. Dufour, and J. Thioulouse. 2004. The {ade4} Package -- {I}: One-Table Methods. R News 4:5–10.

Clark, J. S., M. Dietze, S. Chakraborty, P. K. Agarwal, I. Ibanez, S. LaDeau, and M. Wolosin. 2007. Resolving the biodiversity paradox. Ecology Letters 10:647–659.

Cornwell, W. K., and D. D. Ackerly. 2009. Community assembly and shifts in plant trait distributions across an environmental gradient in coastal California. Ecological Monographs 79:109–126.

Coux, C., R. Rader, I. Bartomeus, and J. M. Tylianakis. 2016. Linking species functional roles to their network roles. Ecology letters 19:762–770.

D'Andrea, R., J. Guittar, J. P. O'Dwyer, H. Figueroa, S. J. Wright, R. Condit, and A. Ostling.

2020. Counting niches: Abundance-by-trait patterns reveal niche partitioning in a

Neotropical forest. Ecology 101.

D'Andrea, R., A. Ostling, and J. P. O'Dwyer. 2018. Translucent windows: how uncertainty in

competitive interactions impacts detection of community pattern. Ecology Letters

21:826–835.

D'Andrea, R., M. Riolo, and A. M. Ostling. 2019. Generalizing clusters of similar species as

a signature of coexistence under competition. PLOS Computational Biology

15:e1006688.

Descy, J. P., F. Darchambeau, T. Lambert, M. P. Stoyneva-Gaertner, S. Bouillon, and A. V.

Borges. 2017. Phytoplankton dynamics in the Congo River. Freshwater Biology 62:87–

101.

Díaz, S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer,

C. Wirth, I. Colin Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B.

Reich, A. T. Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright,

S. N. Sheremet Ev, H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D.

Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk, N. Rüger, M. D. Mahecha,

and L. D. Gorné. 2016. The global spectrum of plant form and function. Nature

529:167–171.

Díaz, S., A. Purvis, J. H. C. Cornelissen, G. M. Mace, M. J. Donoghue, R. M. Ewers, P.

Jordano, and W. D. Pearse. 2013. Functional traits, the phylogeny of function, and

ecosystem service vulnerability. Ecology and Evolution 3:2958–2975.

Dolédec, S., D. Chessel, C. J. F. ter Braak, and S. Champely. 1996. Matching species traits to

environmental variables: a new three-table ordination method. Environmental and

Ecological Statistics 3:143–166.

Downing, A. S., S. Hajdu, O. Hjerne, S. A. Otto, T. Blenckner, U. Larsson, and M. Winder.

2014. Zooming in on size distribution patterns underlying species coexistence in Baltic

Sea phytoplankton. Ecology Letters 17:1219–1227.

Dray, S., P. Choler, S. Dolédec, P. R. Peres-Neto, W. Thuiller, S. Pavoine, and C. J. F. Ter

Braak. 2014. Combining the fourth-corner and the RLQ methods for assessing trait

responses to environmental variation. Ecology 95:14–21.

Dray, S., and A.--B. Dufour. 2007. The {ade4} Package: Implementing the Duality Diagram

for Ecologists. Journal of Statistical Software 22:1–20.

Dray, S., A.-B. Dufour, and D. Chessel. 2007. The {ade4} Package -- {II}: Two-Table and

{K}-Table Methods. R News 7:47–52.

Dray, S., and P. Legendre. 2008. Testing the species traits environment relationships: The

fourth-corner problem revisited. Ecology 89:3400–3412.

Fort, H., M. Scheffer, and E. H. van Nes. 2009. The paradox of the clumps mathematically

explained. Theoretical Ecology 2:171–176.

Fort, H., M. Scheffer, and E. Van Nes. 2010. The clumping transition in niche competition: A

robust critical phenomenon. Journal of Statistical Mechanics: Theory and Experiment

2010.

Gallego, I., P. Venail, and B. W. Ibelings. 2019. Size differences predict niche and relative

fitness differences between phytoplankton species but not their coexistence. ISME

Journal 13:1133–1143.

Gause, G. F. 1936. The Struggle for Existence. Soil Science 41:159.

Götzenberger, L., F. de Bello, K. A. Bråthen, J. Davison, A. Dubuis, A. Guisan, J. Lepš, R.

Lindborg, M. Moora, M. Pärtel, L. Pellissier, J. Pottier, P. Vittoz, K. Zobel, and M.

Zobel. 2012. Ecological assembly rules in plant communities-approaches, patterns and

prospects. Biological Reviews 87:111–127.

Graco-Roza, C., J. B. O. J. B. O. Santos, V. L. M. V. L. M. Huszar, P. Domingos, J.

Soininen, and M. M. M. M. Marinho. 2020. Downstream transport processes modulate

the effects of environmental heterogeneity on riverine phytoplankton. Science of The

Total Environment 703:135519.

Gravel, D., C. D. Canham, M. Beaudet, and C. Messier. 2006. Reconciling niche and

neutrality: The continuum hypothesis. Ecology Letters 9:399–409.

Hardin, G. 1960. The competitive exclusion principle. Science 131:1292–1297.

Hill, M. O., and A. J. E. Smith. 1976. PRINCIPAL COMPONENT ANALYSIS OF

TAXONOMIC DATA WITH MULTI-STATE DISCRETE CHARACTERS. TAXON

25:249–255.

Hillebrand, H., C. D. Dürselen, D. Kirschtel, U. Pollingher, and T. Zohary. 1999. Biovolume

calculation for pelagic and benthic microalgae. Journal of Phycology 35.

Holling, C. S. 1992. Cross-Scale Morphology, Geometry, and Dynamics of Ecosystems.

Ecological Monographs 62:447–502.

Holt, R. D. 2006. Emergent neutrality. Trends in Ecology and Evolution 21:531–533.

Hubbell, S. P. 2001. The Unified Neutral Theory of Biodiversity and Biogeography. Page

Monographs in Population Biology. Princeton University Press, Princeton, Oxford.

Hubbell, S. P. 2006. Neutral theory and the evolution of ecological equivalence. Ecology

87:1387–1398.

Hutchinson, G. E. 1957. Concluding Remarks. Cold Spring Harbor Symposia on Quantitative

Biology 22:415–427.

Ingram, T., R. Costa-Pereira, and M. S. Araújo. 2018. The dimensionality of individual niche

variation. Ecology 99:536–549.

Junk, W. J., P. B. Bayley, and R. E. Sparks. 1989. The flood pulse concept in river-floodplain

systems. Canadian special publication of fisheries and aquatic sciences 106:110–127.

Kraft, N. J. B., R. Valencia, and D. D. Ackerly. 2008. Functional traits and niche-based tree

community assembly in an Amazonian forest. Science 322:580–582.

Kremer, C. T., A. K. Williams, M. Finiguerra, A. A. Fong, A. Kellerman, S. F. Paver, B. B.

Tolar, and B. J. Toscano. 2017. Realizing the potential of trait-based aquatic ecology:

New tools and collaborative approaches. Limnology and Oceanography 62:253–271.

Kruk, C., V. L. M. Huszar, E. T. H. M. Peeters, S. Bonilla, L. Costa, M. LüRling, C. S.

Reynolds, and M. Scheffer. 2010. A morphological classification capturing functional

variation in phytoplankton. Freshwater Biology 55:614–627.

Kruk, C., and A. M. Segura. 2012. The habitat template of phytoplankton morphology-based

functional groups. Hydrobiologia 698:191–202.

Laliberte, E., P. Legendre, E. Laliberté, and P. Legendre. 2010. A distance-based framework

for measuring functional diversity from multiple traits. Ecology 91:299–305.

Laliberté, E., P. Legendre, B. Shipley, and M. E. Laliberté. 2014. Package 'FD.' Measuring

functional diversity from multiple traits, and other tools for functional ecology.

Legendre, P., R. G. Galzin, and M. L. Harmelin-Vivien. 1997. Relating behavior to habitat:

Solutions to the fourth-corner problem. Ecology 78:547–562.

Litchman, E., K. F. Edwards, C. A. Klausmeier, and M. K. Thomas. 2012. Phytoplankton

niches, traits and eco-evolutionary responses to global environmental change. Marine

Ecology Progress Series 470:235–248.

Litchman, E., and C. A. Klausmeier. 2008. Trait-Based Community Ecology of

Phytoplankton. Annual Review of Ecology, Evolution, and Systematics 39:615–639.

Litchman, E., P. de Tezanos Pinto, C. A. Klausmeier, M. K. Thomas, and K. Yoshiyama.

2010. Linking traits to species diversity and community structure in phytoplankton.

Hydrobiologia 653:15–28.

Lund, J. W. G., C. Kipling, and E. D. Le Cren. 1958. The inverted microscope method of

estimating algal numbers and the statistical basis of estimations by counting.

Hydrobiologia 11:143–170.

Macarthur, R., and R. Levins. 1967. The Limiting Similarity, Convergence, and Divergence

of Coexisting Species. The American Naturalist 101:377–385.

Marañón, E. 2008. Inter-specific scaling of phytoplankton production and cell size in the

field. Journal of Plankton Research.

McGill, B. J., B. J. Enquist, E. Weiher, and M. Westoby. 2006. Rebuilding community

ecology from functional traits. Trends in Ecology and Evolution 21:178–185.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin,

R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner.

2020. vegan: Community Ecology Package.

Pavoine, S., M. B. Bonsall, A. Dupaix, U. Jacob, and C. Ricotta. 2017. From phylogenetic to

functional originality: Guide through indices and new developments. Ecological

Indicators 82:196–205.

Peres-Neto, P. R., S. Dray, and C. J. F. ter Braak. 2017. Linking trait variation to the

environment: critical issues with community-weighted mean correlation resolved by the

fourth-corner approach. Ecography 40:806–816.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. Vienna,

Austria.

Reynolds, C., and J. Descy. 1996. The production biomass and structure of phytoplankton in

large rivers. Arch. Hydrobiol. 113:161–187.

Reynolds, C. S. 2006. The Ecology of Phytoplankton. Page The Ecology of Phytoplankton.

Cambridge University Press, Cambridge.

Reynolds, C. S., J. Alex Elliott, and M. A. Frassl. 2014. Predictive utility of trait-separated

phytoplankton groups: A robust approach to modeling population dynamics. Journal of

Great Lakes Research 40:143–150.

Reynolds, C. S., J. P. Descy, and J. Padisák. 1994. Are phytoplankton dynamics in rivers so

different from those in shallow lakes? Hydrobiologia 289:1–7.

Ricotta, C., F. de Bello, M. Moretti, M. Caccianiga, B. E. L. Cerabolini, and S. Pavoine.

2016. Measuring the functional redundancy of biological communities: a quantitative

guide. Methods in Ecology and Evolution 7:1386–1395.

Roy, S., and J. Chattopadhyay. 2007. Towards a resolution of 'the paradox of the plankton':

A brief overview of the proposed mechanisms. Ecological Complexity 4:26–33.

Sakavara, A., G. Tsirtsis, D. L. Roelke, R. Mancy, and S. Spatharis. 2018. Lumpy species

coexistence arises robustly in fluctuating resource environments. Proceedings of the

National Academy of Sciences of the United States of America 115:738–743.

Sauterey, B., B. Ward, J. Rault, C. Bowler, and D. Claessen. 2017. The implications of eco-

evolutionary processes for the emergence of marine plankton community biogeography.

American Naturalist 190:116–130.

Scheffer, M., and E. H. van Nes. 2006. Self-organized similarity, the evolutionary emergence

of groups of similar species. Proceedings of the National Academy of Sciences of the

United States of America 103:6230–6235.

Scheffer, M., E. H. Van Nes, and R. Vergnon. 2018. Toward a unifying theory of

biodiversity. Proceedings of the National Academy of Sciences of the United States of

America 115:639–641.

Scheffer, M., R. Vergnon, E. H. van Nes, J. G. M. Cuppen, E. T. H. M. Peeters, R. Leijs, and

A. N. Nilsson. 2015. The Evolution of Functionally Redundant Species; Evidence from

Beetles. PLOS ONE 10:e0137974.

Segura, A. M., D. Calliari, C. Kruk, D. Conde, S. Bonilla, and H. Fort. 2011. Emergent

neutrality drives phytoplankton species coexistence. Proceedings of the Royal Society

B: Biological Sciences 278:2355–2361.

Segura, A. M., C. Kruk, D. Calliari, and H. Fort. 2013a. Use of a morphology-based

functional approach to model phytoplankton community succession in a shallow

subtropical lake. Freshwater Biology 58:504–512.

Segura, A. M., C. Kruk, D. Calliari, F. García-Rodriguez, D. Conde, C. E. Widdicombe, and

H. Fort. 2013b. Competition drives clumpy species coexistence in estuarine

phytoplankton. Scientific Reports 3:1–6.

Thibault, K. M., E. P. White, A. H. Hurlbert, and S. K. M. Ernest. 2011. Multimodality in the

individual size distributions of bird communities. Global Ecology and Biogeography.

Thioulouse, J., S. Dray, A.--B. Dufour, A. Siberchicot, T. Jombart, and S. Pavoine. 2018.

Multivariate Analysis of Ecological Data with {ade4}. Springer.

Tilman, D. 1982. Resource Competition and Community Structure. Page Monographs in

population biology. Princeton University Press.

Uhelingher, V. 1964. Étude statistique dês méthodes de dénobrement planctonique. Archives

des Sciences 77:121–123.

Utermöhl, H. 1958. Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. SIL

Communications, 1953-1996 9:1–38.

Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The

River Continuum Concept. Canadian Journal of Fisheries and Aquatic Sciences 37:130–

137.

Vergnon, R., N. K. Dulvy, and R. P. Freckleton. 2009. Niches versus neutrality: Uncovering

the drivers of diversity in a species-rich community. Ecology Letters 12:1079–1090.

Villéger, S., N. W. H. H. Mason, and D. Mouillot. 2008. New multidimensional functional

diversity indices for a multifaceted framework in functional ecology. Ecology 89:2290–

2301.

Violle, C., W. Thuiller, N. Mouquet, F. Munoz, N. J. B. Kraft, M. W. Cadotte, S. W.

Livingstone, and D. Mouillot. 2017. Functional Rarity: The Ecology of Outliers. Trends

in Ecology and Evolution 32:356–367.

Wang, C., X. Li, Z. Lai, Y. Li, A. Dauta, and S. Lek. 2014. Patterning and predicting

phytoplankton assemblages in a large subtropical river. Fundamental and Applied

Limnology / Archiv für Hydrobiologie 185:263–279.

Wetzel, R. G. 2001. Limnology: Lake and River Ecosystems. Third edition. Academic Press.

Wickham, H., M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G. Grolemund, A.

Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K. Müller, J.

Ooms, D. Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo,

and H. Yutani. 2019. Welcome to the Tidyverse. Journal of Open Source Software

4:1686.

Wong, M. K. L., and C. P. Carmona. 2021. Including intraspecific trait variability to avoid

distortion of functional diversity and ecological inference: Lessons from natural

assemblages. Methods in Ecology and Evolution 12:946–957.

812