The Genomic Loci of Specific Human tRNA Genes Exhibit Ageing-Related DNA Hypermethylation

Richard J. Acton [1], [2], [3], Wei Yuan [4], [5], Fei Gao [6], Yudong Xia [6], Emma Bourne [7], Eva Wozniak [7], Jordana Bell [4], Karen Lillycrop [3], Jun Wang [6], Elaine Dennison [2], Nicholas Harvey [2], Charles A. Mein [7], Tim D. Spector [4], Pirro G. Hysi [4], Cyrus Cooper [2], Christopher G. Bell [1] *

1 William Harvey Research Institute, Barts & The London School of Medicine and Dentistry, Charterhouse Square, Queen Mary University of London, London, U.K.

2 MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, U.K.

3 Human Development and Health, Institute of Developmental Sciences, University of Southampton, Southampton, U.K.

4 Department of Twin Research & Genetic Epidemiology, St Thomas Hospital, King's College London, London, U.K.

5 Institute of Cancer Research, Sutton, U.K.

6 BGI-Shenzhen, Shenzhen, China

7 Barts & The London Genome Centre, Blizard Institute, Barts & The London School of Medicine and Dentistry, Queen Mary University of London, London, U.K.

* Corresponding author:

Abstract

The epigenome deteriorates with age, potentially impacting on ageing-related disease.

Here, we interrogate the DNA methylation state of the genomic loci of human tRNA. Whilst arising from only ~46kb (<0.002% genome), this information transfer machinery is the second most abundant cellular transcript. tRNAs also control metabolic processes known to affect ageing, through core translational and additional regulatory roles.

We identified a genomic enrichment for age-related DNA hypermethylation at tRNA loci. Analysis in 4,350 MeDIP-seq peripheral-blood DNA methylomes (16-82 years), classified 44 and 21 hypermethylating specific tRNAs at study- and genome-wide significance, respectively, contrasting with 0 hypomethylating. Validation and replication (450k array & independent targeted Bisuphite-sequencing) supported thehypermethylation of this functional unit. The strongest consistent signals, also independent of major cell-type change, occur in tRNA-iMet-CAT-1-4 and tRNA-Ser-AGA-2-6.

This study is this first comprehensive evaluation of the genomic DNA methylation state of human tRNA genes and reveals a discreet hypermethylation with advancing age.

Introduction

Ageing is implicated as a risk factor in multiple chronic diseases [1]. Understanding how the ageing process leads to deteriorating biological function is now a major research focus, with hopes to increase the human 'healthspan' and ameliorate the extensive physical, social and economic costs of these ageing-related disorders [2]. Epigenetic processes, which influence or can inform us about cell-type specific gene expression, are altered with age and are, furthermore, one of the fundamental hallmarks of this progression [3,4].

DNA methylation (DNAm) is the most common epigenetic modification of DNA and age-associated changes in this mark in mammalian tissues have been recognised for decades [5]. In fact, these alterations in DNAm with age are extensive with thousands of loci across the genome affected. Many represent 'drift' arising from the imperfect maintenance of the epigenetic state [6]. However, specific genomic regions show distinct directional changes, with loss of DNA methylation in repetitive or transposable elements [7], as well as gains in certain promoters, including the targets of polycomb repressor complex [8] and bivalent domains [9]. These observations with the advent of high-throughput DNAm arrays also permitted the identification of individual CpG sites that exhibit consistent changes with age, enabling the construction of predictors of chronological age known as epigenetic or DNAm 'clocks' [10–13]. Additionally, it was observed that 'acceleration' of this DNAm-derived measure is a biomarker of 'biological' ageing due to associations with morbidity and mortality (Reviewed in [14] & [15]). In a previous investigation of ageing-related DNAm changes within common disease-associated GWAS regions, we identified hypermethylation of the specific transfer RNA gene, tRNA-iMet-CAT-1-4 [16]. The initiator methionine tRNA possesses certain unique properties, including its capacity to be rate limiting for translation [17], association with the translation initiation factor eIF2 [18], and ability to impact the expression of other tRNA genes [19].

tRNAs are evolutionarily ancient [20] and fundamental in the translation process for all domains of life. This translation machinery and the regulation of protein synthesis are controlled by conserved signalling pathways shown to be modifiable in longevity and ageing interventions [21]. Additionally, beyond their core role in the information flow from DNA to protein sequence, tRNAs can fragment into numerous tRNA-derived small RNAs (tsRNAs) [22] with signalling and regulatory functions [23–26].

When liberally considering the 610 loci annotated as tRNA genes, as well as tRNA pseudo genes, nuclear encoded mitochondrial tRNA genes and possibly some closely related repetitive sequences, these features (gtRNAdb [27]) cover <46 kb (including introns) which represents <0.002% of the human genome [28]. Yet a subset of these genes produce the second most abundant RNA species next to ribosomal RNA [29] and are required for the production of all proteins. tRNA genes are transcribed by RNA polymerase III (polIII) [30] and

have internal type II polIII promoters [31]. DNAm is able to repress the expression of tRNA genes
experimentally [32] but may also represent co-ordination with the local repressive chromatin state [33].
Transcription is also repressed by the highly conserved polIII specific transcription factor Maf1 [34,35], who's
activity is modulated by the Target of Rapamycin Kinase Complex 1 (TORC1) [36]. TORC1 is a highly
conserved hub for signals that modulate ageing [37].

tRNAs as well as tsRNAs are integral to the regulation of protein synthesis and stress response, two processes
known to be major modulators of ageing. Metabolic processes are also recognised to modulate the age estimates
of DNAm clocks [38]. Partial inhibition of translation increases lifespan in multiple model organisms [39] and
PolIII inhibition increases longevity acting downstream of TORC1 [40]. Furthermore, certain tsRNAs circulating
in serum can be modulated by ageing and caloric restriction [41].

We directly investigated ageing-related changes in the epigenetic DNA methylation state of the entire tRNA
gene set, facilitated by the availability of a large-scale MeDIP-seq dataset. Arrays poorly cover this portion of
genome, with even the latest EPIC (850k) arrays only covering <15% of the tRNA genes, with robust probes, and
in total only ~4.7% of all the tRNA gene CpGs [42]. tRNA genes sit at the heart not only of the core biological
process of translation but at a nexus of signalling networks operating in several different paradigms, from small
RNA signalling to large scale chromatin organisation [43]. In summary, tRNA biology, protein synthesis, nutrient
sensing, stress response and ageing are all intimately interlinked. In this study, we have identified tRNA gene
DNA hypermethylation and independently replicated this newly described ageing-related observation.

Results

DNA Methylation of Specific tRNA Gene Loci Changes with Age

Due to tRNAs critical role in translation and evidence of their modulation in ageing and longevity-related
pathways, we interrogated these genes for evidence of ageing-related epigenetic changes. Our discovery set was a
large-scale peripheral blood-derived DNA methylome dataset comprising of 4350 samples. This sequencing-based
dataset had been generated by Methylated DNA Immunoprecipitation (MeDIP-seq) [44], which relies on the
enrichment of methylated fragments of 200-500 bp to give a regional DNAm assessment (500 bp semi-overlapping
windows, see Methods). There are 416 high confidence tRNA genes in the human genome, we initially examined
an expanded set of 610 tRNA-like features identified by tRNAscan including pseudogenes, possible repetitive
elements, and nuclear encoded mitochondrial tRNAs [27,45]. Of these 492 were autosomal and did not reside in
blacklisted regions of the genome [46]. Due to the small size of these tRNAs (60-86bp, median 73bp, excluding
introns which are present in ˜30 tRNAs with sizes from 10-99bp, median 19bp), this fragment-based method
enabled a robust examination of the epigenetic state of these highly similar sequences. This was supported by a
mappability assessment. The median mappability score density for all tRNA genes was 0.90 for 50mers when
considering tRNA genes $\pm 500 bp$ reflecting the regional nature of the MeDIP-seq assay. In contrast the 50mer
mappability density is 0.68 for tRNA genes alone, which would be representative of the mappability of reads
generated using a technique such as whole-genome bisulfite sequencing (see Figures S1 & S2).

We identified 21 genome-wide significant and 44 study-wide significant results (p $< 4.34 \times 10^{-9}$ and
8.36×10^{-5}, respectively, via linear regression (n=4350, see Methods). Study-wide significance was calculated
conservatively with the Bonferroni method for all 598 autosomal tRNAs and closely related features. There was a
strong directional trend with all results at both significance levels being due to increases in DNA methylation.
Age-related changes in cell type proportion are strong in heterogeneous peripheral blood, and include a myeloid
skew, loss of naive T cells and increases in senescent cells [47]. A subset of 3 genome-wide and 16 study-wide
significant hypermethylation results remained significant even after correcting for potential cell-type changes by
including lymphocytes, monocytes, neutrophils and eosinophil cell count data (n=3001, Listed in Figure 1, see
Figure S3). Going forward we refer to these cell-type corrected sets of 3 and 16 tRNA genes as the genome-wide
and study-wide significant tRNA genes respectively.

tRNA	Window	MeDIP		450k Array		Targeted BiS-seq	
		Slope	p-value	Slope	p-value	Slope	p-value
tRNA-Gln-CTG-7-1	Chr1:147,800,750-147,801,250	0.84	2.60e-05				
tRNA-Glu-TTC-1-1	Chr2:131,094,500-131,095,000	1.11	4.64e-09				
	Chr2:131,094,250-131,094,750	1.00	1.12e-07				
	Chr2:131,094,750-131,095,250	1.00	3.28e-07				
tRNA-His-GTG-1-2	Chr1:146,544,500-146,545,000	0.92	1.38e-06				
tRNA-His-GTG-2-1	Chr1:149,155,750-149,156,250	1.05	2.98e-08				
	Chr1:149,155,500-149,156,000	0.83	1.37e-05				
tRNA-Ile-AAT-10-1	Chr6: 27,251,500- 27,252,000	1.07	1.45e-08			1.30	1.22e-03
	Chr6: 27,251,750- 27,252,250	0.90	1.86e-06			1.30	1.22e-03
tRNA-Ile-AAT-4-1	Chr17: 8,130,000- 8,130,500	1.19	2.98e-10	19.63	8.92e-06	-0.74	6.88e-04
	Chr17: 8,130,250- 8,130,750	0.77	3.99e-05	19.63	8.92e-06	-0.74	6.88e-04
tRNA-Ile-TAT-2-2	Chr6: 26,987,750- 26,988,250	0.97	7.25e-07	4.16	1.17e-02	-0.60	3.84e-01
tRNA-iMet-CAT-1-4	Chr6: 26,330,500- 26,331,000	1.28	2.83e-11	13.01	6.07e-06	4.54	9.35e-04
	Chr6: 26,330,250- 26,330,750	1.13	2.89e-09	13.01	6.07e-06	4.54	9.35e-04
tRNA-Leu-TAG-2-1	Chr14: 21,093,250- 21,093,750	1.04	9.38e-08			2.49	8.77e-03
	Chr14: 21,093,500- 21,094,000	0.94	8.50e-07			2.49	8.77e-03
tRNA-Pro-AGG-2-2	Chr6: 26,555,500- 26,556,000	1.04	3.97e-08				
	Chr6: 26,555,250- 26,555,750	1.01	9.58e-08				
tRNA-Ser-ACT-1-1	Chr6: 27,261,250- 27,261,750	0.97	3.53e-07			0.66	1.45e-01
tRNA-Ser-AGA-2-6	Chr17: 8,129,750- 8,130,250	1.21	1.16e-10	20.87	6.72e-05	0.62	4.28e-02
	Chr17: 8,130,000- 8,130,500	1.19	3.03e-10	20.87	6.72e-05	0.62	4.28e-02
tRNA-Ser-TGA-2-1	Chr6: 27,513,000- 27,513,500	0.90	3.58e-06	87.21	1.38e-04	-0.25	5.74e-01
tRNA-Val-AAC-1-2	Chr5:180,590,750-180,591,250	0.91	3.28e-06				
tRNA-Val-AAC-4-1	Chr6: 27,648,500- 27,649,000	1.07	1.25e-08	40.06	9.90e-03		
	Chr6: 27,648,750- 27,649,250	0.95	4.31e-07	40.06	9.90e-03		
tRNA-Val-CAC-2-1	Chr6: 27,247,750- 27,248,250	0.85	2.33e-05	59.16	5.05e-06		

Fig 1. Ageing-related DNA methylation changes in tRNA loci (16 Study-wide significant after blood cell-type correction, $p < 8.36 \times 10^{-5}$). Columns left to right: tRNA gene; Genomic location of MeDIP-seq significant window (hg19); MeDIP results slope (arbitrary units) and p-value; 450k Array results slope (beta) and p-value (Blood-cell corrected and combined for all probes overlapping tRNA); Targeted Bisulphite-sequencing (BiS-seq) slope (beta) and p value (combined all CpGs in targeted region, see Methods) For slope: Orange = hypermethylation and Blue = hypomethylation with age. For p-values: Colour gradient high significance = dark blue to low significance = yellow, scaled relative to column. Blank grey cells indicate unavailable data.

Due to the related nature of these twin samples, we also analysed these data in two subsets of n = 1198 & 1206 by selecting one twin from each pair into the separate sets. This analysis also included correction for Blood Cell counts. Even in these smaller sets, 5 and 7 tRNA genes, respectively, reached study-wide significance. In these sets 5/5 and 6/7 of these were present in 16 study-wide significant tRNA genes.

Furthermore, we examined a subset of samples with longitudinal data (n=658 methylomes from 329 individuals, median age difference 7.6 yrs). At the nominal significance threshold ($p < 0.05$) this yielded a split of 41 hypermethylating tRNA genes and 22 hypomethylating tRNA genes. Of these hypermethylated tRNAs, 2

are in the previously identified genome-wide significant set of 3 (with tRNA-iMet-CAT-1-4 ranked 3rd by p-value) and 9 are in the study-wide significant set of 16. Full results for each age model are provided in Supplementary File S1

tRNA Genes are Enriched for Age Related DNA Hypermethylation

Whilst ageing changes are pervasive throughout the DNA methylome, we identified a strong enrichment for this to occur within the tRNA genes (Fisher's Exact Test p $= 1.05 \times 10^{-27}$) (see Figure 2 b). This is still significant if the 6 of the study-wide significant 16 tRNAs that have any overlap with polycomb or bivalent regions are excluded (p $= 4.66 \times 10^{-15}$)

Fig 2. A) tRNA genes are enriched for age-related hypermethylation compared to the genomic background, (Fisher's Exact Test p $< 1.05 \times 10^{-27}$, n $= 3001$). **B)** tRNA genes show more significant hypermethylated loci than CpG Density matched permutations. Each permutation represented a random set of windows matching the CpG density of the functional unit (tRNA gene loci, bivalent domains, and polycomb group target promoters). These are subsequently assessed for significant age-related DNAm changes (see Methods). The red line is the observed number of significant loci.

CpG density itself is known to have a clear impact on the potential for variability of the DNA methylome as well as ageing-related changes [48,49]. To assess whether this hypermethylation finding was being merely driven by the inherent CpG density of the tRNA genes, we performed a CpG density matched permutation analysis (1,000X, see Methods). This supported the specific nature of these age-related DNAm changes within the functional tRNA genes (Empirical p-value $< 1.0 \times 10^{-3}$, Figure 2 a). As a point of comparison for this genomic functional unit, we also performed the same permutations for the known age-related changes in the promoters of genes that are polycomb group targets [8] and those with a bivalent chromatin state [9]. We were able to reproduce the enrichment of the polycomb group targets and bivalent regions (Empirical p-value $< 1.0 \times 10^{-3}$) in our dataset.

tRNA-iMet-CAT-1-4 (Figure 3a) is located in the largest tRNA gene cluster in the human genome at chr6p22.2-1. This cluster contains 157 tRNA genes spanning the 2.67Mb from tRNA-iMet-CAT-1-2 to tRNA-Leu-AAG-3-1, and also hosts a histone gene microcluster. tRNA-Ile-AAT-4-1 and tRNA-Ser-AGA-2-6 are neighbours and are located on chromosome 17 (Figure 3b). Notably tRNA-Ile-AAT-4-1 and tRNA-Ser-AGA-2-6 have a third close neighbour tRNA-Thr-AGT-1-2 which does not show significant age-related hypermethylation. We observe a similar pattern of sharp peaks of significance closely localised around the other loci in the study-wide significant set. GENCODE 19 places tRNA-Ile-AAT-4-1 in the 3'UTR of a Nonsense-mediated decay transcript of *CTC1* (CST Telomere Replication Complex Component 1, CTC1-002, ENST00000449476.2) and not of its primary transcript.

Fig 3. MeDIP-seq results (blood cell-type corrected model) for **A)** tRNA-iMet-CAT-1-4 as well as **B)** tRNA-Ser-AGA-2-6 and tRNA-Ile-AAT-4-1 exhibiting ageing-related DNA hypermethylation. Top to Bottom: Chromosome ideogram; Gene locations (GENCODE 19); MeDIP-seq DNA methylation ageing $-log_{10}$ p-value results (shown for each 500 bp overlapping window); MeDIP-seq slope of change with age; MeDIP-seq Medium coverage (Reads per Millions, RPM) calculated across all samples; CpG density (%); CpG locations (red lines); 500bp overlapping windows. The sharp peaks suggest that the results are localised to individual tRNA genes not the entire genic locus. One window overlapping tRNA-Ile-AAT-4-1 also partially overlaps tRNA-Ser-AGA-2-6. The 3' UTR of the CTC1 transcript CTC1-002 (ENST00000449476.2, GENCODE 19), which is subject to nonsense mediate decay, overlaps tRNA-Ile-AAT-4-1. tRNA genes with similar CpG density are exhibiting differing age-related DNAm change patterns.

To place these hypermethylating tRNA genes in their genomic context we examined how the extended set of 44 non-blood cell-type corrected study-wide significant tRNAs clustered with other tRNA genes. We clustered the tRNA genes by grouping together all tRNAs within 5Mb of one another and then required that a cluster contain at least 5 tRNA genes with a density of at least 5 tRNA genes per Mb (see Methods). This yielded 12 major tRNA gene clusters containing a total of 353 tRNA genes, 42 of the 44 study-wide significant tRNA genes within these clusters (Figure 4a). The hypermethylating tRNA genes are spread evenly among these clusters proportionately to their size (% ageing tRNA per cluster; non-significant one-way ANOVA).

Fig 4. A) Covered study-wide significant tRNA genes (n = 14/16) are more methylated in a centenarian than in a neonate or an adult (26 year) in Whole Genome Bisulfite Sequencing (WGBS) data from Heyn et al. [50]. Each point represents individual CpG methylation levels within a tRNA gene. **B)** No significant clusters of ageing-related tRNA. Proportion of non-blood cell corrected ageing-related study-wide significant tRNAs (42/44, red) within tRNA clusters (tRNAs within 5Mb with >5 tRNA genes/Mb, see Methods).

Age-related tRNA gene set DNA Hypermethylation is even observed in one Newborn versus one Centenarian

We examined an available Whole Genome Bisulphite sequencing (WGBS) dataset from Heyn *et al.* [50] (see Methods) These data consisted of blood-derived DNA WGBS in one newborn child and one 26 year old, and centenarian (103 years). In their analysis, the centenarian was found to have more hypomethylated CpGs than the neonate across all genomic compartments, including promoters, exonic, intronic, and intergenic regions. However, even in this examination of 3 individuals of 3 different age in the 55% of tRNA that possessed coverage, we observed DNA hypermethylation with age among the study-wide significant hypermethylating tRNA genes. The centenarian was significantly more methylated in this set of tRNAs than the neonate (Wilcoxon rank sum test, 6.14% increase (95% CI -Inf - 4.31), W = 717, p = 6.14×10^{-4}, see Figure 4b).

Age-related Changes Independently Replicated with Targeted Bisulfite Sequencing

In order to further robustly support these-ageing related changes, we attempted to replicate these findings ourselves in an independent ageing dataset. Furthermore, we employed a different technology targeted bisulfite (BiS) sequencing to also further validate the MeDIP-seq-derived results. These data provide individual CpG resolution to identify what may be driving the regional DNAm changes observed, and precise quantitation of the magnitude of change.

We performed this targeted BiS-seq in blood-derived DNA from 8 pools of age-matched individuals at 4 time-points (~4, ~28, ~63, ~78 years) from a total of 190 individuals, as detailed in Table 1. This approach permitted us to assay tRNA gene DNA methylation across a large number of individuals whilst requiring a minimal amount of DNA from each (80-100ng), and costing ~1/24th as much as performing sequencing individually. A total of 79 tRNA loci generated reliable results post-QC (see Methods). These tRNAs covered a total of 458 CpGs with a median of 6 CpGs per tRNA (range 1-9). Median Coverage per site across pools, technical replicates and batches was 679 reads (mean 5902).

Firstly, we ran the 8 Pooled samples on the Illumina EPIC (850k) array to confirm that our pooling approach was applicable for DNAm ageing-related evaluation. This showed an $R^2 = 0.98$ between pool mean chronological age and Horvath clock DNAm predicted age [11](see Figure 5c). Therefore, this confirmed the utility of our pooling approach. We also used these array derived data to estimate the major blood cell proportions for each of these pools with the Houseman algorithm [51].

Table 1. Summary information on participants in each pool. Total of 190 participants in 8 pools, at 4 time points: 4, 28, 63, & 78 years

Pool	Mean Age	Sex	Min Age	Max Age	n
Pool 1	4.07	Male	3.99	4.38	20
Pool 2	4.09	Female	3.99	4.36	20
Pool 3	28.07	Female	25.87	29.80	25
Pool 4	28.23	Female	26.05	30.01	25
Pool 5	63.40	Female	62.80	63.80	25
Pool 6	63.26	Female	62.70	63.70	25
Pool 7	77.96	Female	75.50	80.50	25
Pool 8	77.22	Female	74.40	80.10	25

Table 2. Pairwise Differences in Methylation between Age groups by tRNA. p-values are for pairwise methylation differences (see Methods)[52].

tRNA	num. CpGs	comparison	p-value	delta
tRNA-Ile-AAT-4-1	8	4 vs. 28	1.518e-01	-0.2
		4 vs. 63	1.774e-01	-0.234
		4 vs. 78	3.060e-01	0.0113
		28 vs. 63	7.152e-01	-0.0334
		28 vs. 78	1.553e-01	0.212
		63 vs. 78	2.057e-01	0.245
tRNA-iMet-CAT-1-4	5	4 vs. 28	8.403e-02	0.0116
		4 vs. 63	1.716e-01	0.0125
		4 vs. 78	1.997e-04*	0.0368
		28 vs. 63	3.943e-01	0.000869
		28 vs. 78	1.724e-02*	0.0252
		63 vs. 78	6.224e-02	0.0243
tRNA-Ser-AGA-2-6	9	4 vs. 28	4.222e-01	0.0573
		4 vs. 63	3.968e-01	0.0274
		4 vs. 78	4.651e-01	0.0423
		28 vs. 63	1.095e-01	-0.0299
		28 vs. 78	2.126e-01	-0.015
		63 vs. 78	2.201e-01	0.0149

We noted that individual tRNA loci exhibiting age-related changes in DNAm had duplicate or isodecoder (same anticodon but body sequence variation) sequences in the genome, which despite exact or near sequence identity did not show similar changes. tRNA-iMet-CAT-1-4 for instance is 1 of 8 identical copies in the genome and was the only locus that showed significant changes. The results of pairwise differential methylation tests between age groups for the 6 top tRNAs from the MeDIP-seq models are listed in Table 2. tRNA-iMet-CAT-1-4 shows a pairwise increase of 3.7% from age 4 years to age 78 years.

Of the 3 top hits in MeDIP-seq, tRNA-iMet-CAT-1-4 (Figure 5a(iii)) and tRNA-Ser-AGA-2-6 (Figure 5a(viii)) exceeded nominal significance (p-values = 9.35×10^{-4} & 4.28×10^{-2}, respectively). However, tRNA-Ile-AAT-4-1 (Figure 5a(xi)) showed a nominal decrease in DNAm with age. tRNA tRNA-Leu-TAG-2-1

from the study-wide significant set also showed nominally significant hypermethylation with age (Figure 5a(xviii)). Also, four of the individual CpGs in tRNA-iMet-CAT-1-4 exhibited nominally significant increases in DNAm with Age (Figure 5b).

Fig 5. Comparison of age-related hypermethylating tRNA genes to closely related tRNA genes using targeted bisulfite sequencing (BiS-seq). A) Combined CpGs within tRNA loci results (experiment-wide Bonferroni p = 7.14×10^{-4}); *i-iv*, Comparison of select tRNA-iMet-CAT loci: Hypermethylation is specific to iMet-CAT-1-4 (iii) not other isodecoders (i, ii, & iv); *v-viii*, Comparison of select tRNA-Ser-AGA loci: Hypermethylation is specific to Ser-AGA-2-6 A (viii) and to a lesser extent Ser-AGA-2-1 (v), whilst not other isodecoders (vi, vii); *ix-xiv*, Comparison of select tRNA-Ile loci: Hypermethylation is specific to Ile-AAT-10-1 (ix), Ile-AAT-4-1 (xi) displays hypomethylation contrary to previous MeDIP findings, Ile-TAT-2-2 & 2-3 lack hypermethylation (previously non-significant in blood-corrected MeDIP, although significant in uncorrected), whilst no change in Ile-AAT-11-1 (x) and Ile-TAT-2-1 (xii); *xv-xviii*, Comparison of select tRNA-Leu loci: Hypermethylation in Leu-AAG-3-1 (xv) consistent with 450k and Leu-TAG-2-1 (xviii) consistent with MeDIP, whilst no change in Leu-TAA-1-1 (xvi) & Leu-TAA-3-1 (xvii). **B)** tRNA-iMet-CAT-1-4 Individual CpG analysis: 4 CpGs within this tRNA show consistent methylation increases (all p < 0.05). **C)** Mean chronological age is tightly correlated with DNAm Horvath clock [11] predicted age for the 8 pooled samples (see Table 1 for pool details).

Select Duplicates & Isodecoders of Hypermethylating tRNA loci remain unchanged We targeted a selection of these duplicate and isodecoder loci for bisulfite sequencing in order to confirm that the identified DNAm changes are specific to a given locus and not general to related tRNAs. Examining the tRNA-iMet-CAT-1 family, only the previously identified 1-4 version confirmed significant hypermethylation (not 1-2, 1-3 or 1-5)(Figure 5a(i-iv)). Likewise the tRNA-Ser-AGA-2-6 version was supported compared to 2-1,2-4 and 2-5(Figure 5a(v-viii))). Full Age model results are available in Supplementary File S2.

DNA methylation 450k Array Data Validates the MeDIP-seq Results

Although DNA methylation array poorly covers the tRNA gene set, we wished to attempt to see if this BiS-based but differing and well-established technology was supportive at all of our DNA hypermethylation findings. TwinsUK had available 450k array on 587 individuals, and this platform includes 143 probes, covering 103 tRNAs. All the 3 top tRNAs in the MeDIP-seq results were covered by this data set, and 7 of the 16 study-wide significant set. 9 tRNAs show significant ($p < 4.58 \times 10^{-4}$) increases in DNA methylation with age in models corrected for blood cell counts including all 3 of the 3 tRNAs identified in the MeDIP-seq as genome-wide significant and 5 of the 7 study-wide significant set present on the array (Figure 6). Although it should be noted that 56 of these 143 probes are within the non-robust set of Zhou et al. [42], including 1 of the genome-wide, and 1 of the study-wide results (covering tRNA-Ile-AAT-4-1 & tRNA-Val-AAC-4-1), respectively. Full age model results for all tRNA probes are provided in Supplementary File S3.

Fig 6. Age-related DNA methylation change in tRNA genes Five of seven tRNAs study-wide
significant in MeDIP covered by the 450k array also show significant hypermethylation in this data. tRNAs are
labelled if they are significant here or were in the MeDIP-seq data (Red). Model slope: the model coefficient for
the methylation values. Unfilled circles indicate those probes with potential issues generated by Zhou et al. [42],
this includes tRNA-Ile-AAT-4-1. Significance threshold: $0.05/103 \approx 4.58 \times 10^{-4}$ (the number of tRNA genes
examined).

Ageing-Related tRNA Loci show increased Enhancer-Related Chromatin Signatures   188

We further explored the activity of the tRNA genes in public Chromatin segmentation data in blood (Epilogos   189

Blood & T-cells set) [53,54]. This shows proportionally more Enhancer-related (Enh, EnhBiv & EnhG)   190

chromatin states at tRNA genes hypermethylating with age than the stronger Promoter-related (TSS) in other   191

tRNAs. (Figure 7 b). Whereas these characteristics are less frequently predominant in the rest of the tRNAs   192

(Figure 7 b). Age-hypermethylating tRNA are enriched for enhancer chromatin states compared to the rest of the 193

tRNA genes (Fisher's Exact test p = 0.01).

Fig 7. A) Heatmap of Mean DNA methylation of tRNA sorted Blood Cell Types (data from Reinus et al., probes covering 8 of the 16 study-wide significant tRNA in 7 cell-type fractions from 6 males via GSE35069, [55,56]). This illustrates the ability of tRNA gene methylation to separate the myeloid from the lymphoid lineage and the higher methylation in the lymphoid lineage of the 3 cell-type corrected genome-wide significant tRNAs (iMet-CAT-1-4, etc.) **B)** Chromatin Segmentation data for 'Blood & T-cell' from Epilogos 15 State model [53,54]. Proportion represents the frequency of the predominant chromatin state at a given tRNA (+/- 200bp) for 14/16 study-wide significant tRNAs covered compared to other 371 available tRNAs. The ageing-related hypermethylated tRNAs are enriched for enhancer chromatin state (Fisher's Exact p = 0.01).

Age Hypermethylating tRNAs are more methylated in Lymphoid than Myeloid cells

Three tRNA genes remained genome-wide significant and 16 study-wide significant following correction for major cell-type fraction. This is suggestive of either cell-type independent change or, presumably less likely, a very large effect in a minor cell-type fraction. tRNAs have exhibited tissue-specific expression [57–59] and blood cell-type populations change with age. Specifically, there is shift to favour the production of cells in the myeloid lineage [47]. These points lead us to examine tRNA gene DNAm in sorted cell populations. We used a publically available 450k array dataset [55]) that has been used in the construction of cell-type specific DNAm references for cell-type fraction prediction using the Houseman algorithm [51] (see Methods). This consists of data from 6 individuals (aged 38 ± 13.6/yrs) from seven isolated cell populations (CD4+ T cells, CD8+ T cells, CD56+ NK cells, CD19+ B cells, CD14+ monocytes, neutrophils, and eosinophils). We found that tRNA gene DNAm could separate myeloid from lymphoid lineages (Figures 7 a & S4).

Of the eight study-wide significant tRNAs with array coverage, we identified that collectively these eight are significantly more methylated in the lymphoid than the myeloid lineage (1.1% difference, Wilcoxon rank sum test $p = 1.50 \times 10^{-6}$ 95% CI 0.7%-∞). Thus, any age related increases in myeloid cell proportion would be expected to dampen rather than exaggerate the age-related hypermethylation signal that we observed. In addition

tRNA-Ile-AAT-4-1 and tRNA-Ser-AGA-2-6 have the highest variance in their DNAm of all 129 tRNAs covered in this dataset. This could represent ageing-related changes as these samples range across almost 3 decades. Another possibly may be that these loci as well as hypermethylating are also increasing their variability with age in a similar fashion to those identified by Slieker *et al.* [60]. In that study they identified that those loci accruing methylomic variability were associated with fundamental ageing mechanisms.

tRNA Genes also Hypermethylate with Age in Solid Tissue

Some tRNA gene expression has been shown to be highly tissue specific [57–59]. It follows that our observations of changes in DNAm with age in blood might be specific to that tissue. We used a mix of 450k and 27k array data from 'solid tissue normal' samples made available by TCGA (The Cancer Genome Atlas) and data from foetal tissue [61,62] downloaded from GEO (see Methods). The samples from TCGA range in age from 15-90 (n = 733). Only 43 tRNA genes had adequate data to compare across tissues in this dataset and 115 in the foetal tissue data.

Only 2 of the 3 tRNA genes we identified as genome-wide significant and a further 1 of the study-wide significant tRNA genes are present in the set of 45 tRNA genes in the TCGA data, thus limiting our ability to draw conclusions about the tissue specificity of our results. Solid tissue samples have a strong preponderance for low levels of methylation consistent with the active transcription of many tRNA genes and show slight increasing methylation with age but age accounts for very little of the variance (linear regression slope estimate = 1.52; $R^2 = 0.0002$; p-value 1.34×10^{-3} (Figure S5d). In a pan-tissue analysis we found that 10 tRNA genes showed changes in DNAm with age, 9 of which were hypermethylation (p-value $< 1.1 \times 10^{-3}$). One of these tRNA genes, tRNA-Ser-TGA-2-1 was also present in study-wide significant set of tRNA genes. Figures S6 & S7 illustrate minimal tissue specific differences. Interestingly, however, tRNA-iMet-CAT-1-4 and tRNA-Ser-AGA-2-6 appeared more variable in methylation state than many other tRNAs in the TCGA normal tissue samples (Figure S6) and indeed have the highest variance in DNA methylation across tissues (Figure S5c).

Mice also show age-related tRNA gene DNA hypermethylation

We examined the DNA methylation of the mouse tRNA genes in using data from a reduced representation bisulfite sequencing (RRBS) experiment performed by Petkovich et al. [63]. These data from 152 mice covered 51 tRNA genes and 385 CpGs after QC (see Methods), representing ˜11% of mouse tRNA genes. The mice ranged in age from 0.67-35 months.

Three of the 51 tRNAs showed Bonferroni significant DNA methylation changes with age (p-value <

1.08×10^{-4}) and all were in the hypermethylation direction. These three are tRNA-Asp-GTC-1-12, 239
tRNA-Ile-AAT-1-4, tRNA-Glu-TTC-1-3 (Figure 8). Full Age model Results are available in Supplementary File 240
S4. 241

Fig 8. DNA methylation of CpGs in 3 tRNA which significantly hypermethylate with age in mice, in data from Petkovich et al. [63]. 6 CpGs reach bonferroni significance and 7 show nominally significant increases.

In order to investigate the mouse results further we made use of data from Thornlow et al. [64], which had 242
established tRNA ortholog sets for 29 mammalian species. They identified 197 mouse tRNAs as having direct 243
human orthologs with 44 of these included in the mouse results from Petkovich et al. [63]. However, 244
unfortunately, although 2 of the top 3 tRNAs (tRNA-Ser-AGA-2-6 & tRNA-Ile-AAT-4-1) have mouse orthologs 245
(tRNA-Ser-AGA-2-2 & tRNA-Ile-AAT-1-1), they are not covered in these mouse data. Furthermore, none of the 246
tRNAs showing significant hypermethylation in mice (at nominal p<0.05) had human orthologs in our 247
MeDIP-seq study-wide significant hypermethylating set. Therefore, whilst we cannot directly compare specific 248

tRNA loci due to this lack of coverage, it is interesting that the small number of significant tRNA genes in the
mouse data also hypermethylate with age.

Discussion

Our work has identified a previously unknown enrichment for age-related epigenetic changes within the tRNA genes of the human genome. This observation was strongly directional with increasing DNA methylation with age [65]. The MeDIP-seq dataset employed brought advantages in exploring this undefined terrain of the human tRNA genes. Firstly, being genome-wide it provides much increased access, as these regions are poorly covered by current arrays. Secondly, being a fragment-based regional assessment of DNA methylation, the individual but highly similar small tRNA genes can be surrounded by unique sequence.

We determined by genome-wide permutation that this strong hypermethylation signal was specific to the tRNA genes, and not merely driven by the underlying CpG density of these loci. A targeted BiS-seq experiment validated the defined nature of the tRNA change in an independent dataset, with a successful pooling approach, which may also be useful for other ageing-related targeted DNA methylome evaluations. Additionally, we gained support for our results from limited DNA methylation array data.

Whilst the changes in DNA methylation we have observed are relatively small (i.e. 3.7% between 4-year to 78-year pools in tRNA-iMet-CAT-1-4), this is consistent with the effect size seen in the majority of positions differentially methylatyed with age in many other studies [60,66], except for the noted extreme outliers, such as *EVOLV2* [67]. Furthermore, effect size cut-offs are observed to remove a large fraction of true age-DMPs [68].

We subsequently explored further what was driving this age-related phenomenon and its possible biological implications. As this result was observed in peripheral blood, we were well aware that we were examining DNA derived from a heterogeneous cell type population [69]. Moreover, that there are well known age-related proportional changes in peripheral blood cell composition [47]. The TwinsUK MeDIP-seq and 450k array DNA methylation data included measured haematological values. Therefore, we adjusted for major cell type effects, such as a myeloid skew, and distinct tRNAs were still significant. Although, a caveat to our study is that this can not exclude changes in minor specific sub-cell fractions types. However, that these age-related effects were strong enough to be observed in both a regional MeDIP-seq assessment and a pooled sequencing approach, implies that they not extremely subtle. We examined age-related tRNA gene DNA methylation changes in the limited subset of mouse tRNA genes covered in publicly available RRBS data (˜13%) and were able to identify tRNAs exhibiting DNA hypermethylation with age in this set. This suggests that age-related tRNA gene hypermethylation may not be unique to humans, but at least observed across mammals.

Due to the high number of hypermethylating tRNAs prior to cell-type correction, we were also curious whether the epigenetic state of this small tRNA gene fraction of genome could capture and in fact be a defined fingerprint of cell type. We found that tRNA gene DNA methylation could separate myeloid from lymphoid lineages. There

was also some suggestion of more fine-grained blood cell-type signatures in tRNA DNAm, such as the separation of CD19+ B cells from CD4/8+ T cells. Ageing is also known to lead to an increase in senescent cells (*e.g.* CD8+ CD28- cells). Whether these epigenetic changes in the tRNA genes uniquely represent these cell-types will require technical advances to enable future single cell DNA methylome analysis to accurately assess these regions. If further supported, the epigenetic state of these tRNA loci may aid the taxonomy of cell-type definition.

We observed a predominantly unmethylated state across fetal (Figure S7) and adult tissues (Figure S6) at tRNA gene loci, consistent with the high rate of transcription at many tRNA genes. We suspect that the tRNA genes largely remain unmethylated through development and that the moderate increases in DNAm that we are observing with age at these loci are being driven by changes arising primarily in older individuals. Distinct biological changes have been observed recently in aged individuals [70,71]. Our data are supportive of the tRNA gene set representing a distinct functional unit, which lacks the logarithm change in the young observed strongly in CpGs employed in 'clocks' [13]. This would also be consistent with the lack of significant differences in the tRNA loci detected between the neonate and the 26-year-old adult in the Heyn et al. data. This low baseline DNA methylation of the tRNA genes may also explain why we predominantly observe hypermethylation with age. Whether this is driven by mechanisms, such as aberrant DNA methylation targeting of the tRNA loci or specific sub-celltype effects with age, will require further experimental investigation.

This signal within the tRNA families was observed to occur at specific Isodecoders. Isodecoders expand in number with organismal complexity and the high prevalence in mammals has been suggested due to their additional regulatory functionality [72,73]. They also have distinct translational efficiency [74], which can also have consequences in human disease [75]. After correcting for major cell types, we identified 2 tRNA genes tRNA-iMet-CAT-1-4 and tRNA-Ser-AGA-2-6 which had the most consistent hypermethylation across 3 different assays. Regarding the inconsistent tRNA-Ile-AAT-4-1 result, it is covered by a MeDIP-seq 500bp window which exhibited genome-wide significant hypermethylation, but also partially overlaps tRNA-Ser-AGA-2-6 (Figure 3B). Whilst the 450k array probe overlapping tRNA-Ile-AAT-4-1 (cg06382303) appears to replicate this hypermethylation, it is a borderline case for exclusion flagged by Zhou et al. [42] due to non-unique 30bp 3' subsequence. In the targeted Bisulfite sequencing data, tRNA-Ile-AAT-4-1 exhibited a loss of methylation. Therefore, this may suggest that the hypermethylation signal observed at this locus in the MeDIP-seq data could have been 'pulled up' by the neighbouring tRNA-Ser-AGA-2-6 hypermethylation signal. Of the 16 study-wide significant tRNA genes, only these two of these have a shared significant window, furthermore, in the expanded set of 44 only tRNA-Thr-AGT-1-2 could be similarly affected.

Furthermore, there is great complexity to the fragmentation of tRNA [23], with physiological processes such as stress shown to induce fragment production [76]. These resultant tsRNAs can feedback on protein synthesis by

regulating ribosome biogenesis [77] and others have diverse regulatory functions such as targeting transposable 314
element transcripts [78]. They are also observed to circulate in the blood in a cell-free fashion, and fragment 315
levels can be modulated by ageing and calorie restriction [41]. The isodecoder specific nature of our findings 316
frame a possible hypothesis for regulatory change with age and future work will be required to unravel this 317
potential. tsRNA abundance has been associated with locus specific tRNA gene expression, in some cases 318
independent of mature tRNA levels [79]. This has important implications for the interpretation of our results 319
given the multi-copy nature of genes like tRNA-iMet-CAT-1-4, as even if expression levels of mature iMet tRNAs 320
are unaffected by changes in one copy's DNA methylation, these changes could still influence the levels of 321
particular tsRNAs derived from specific tRNA loci. 322

The location of tRNA-Ser-AGA-2-6 and tRNA-Ile-AAT-4-1 immediately downstream of *CTC1* and of 323
tRNA-Ile-AAT-4-1 within the 3'UTR of an alternate isoform of *CTC1*, which undergoes nonsense-mediated decay, 324
raises the possibility that the gene body epigenetic regulation of *CTC1* may impact on the state of these tRNA 325
genes. *CTC1*'s function in telomere maintenance [80], DNA replication licensing [81], and it's role in a rare 326
progeroid condition [82] indicate that it has ageing-relevant biology. The possible relationship between the the 327
regulation of *CTC1* and that of the tRNA genes downstream of it warrants further study. 328

Whilst, the expression of the tRNA genes has long been simplified as 'constitutive', some observations have 329
indicated that many tRNA genes are expressed in a tissue-specific fashion in diverse organisms [58,59]. Although 330
others have found the majority of isodecoders are transcribed in different cell types [28]. Several transcription 331
factors acting via TFIIIB [83] have a negative (the tumour suppressors p53 [84] and Rb [85]) or positive (the 332
proto-oncogene c-Myc) influence [83]. Regulatory sequence in the flanking or the internal regions of tRNA genes 333
do not explain tRNA expression variation [86]. Whilst DNAm is able to repress the expression of tRNA genes 334
[32], the broader chromatin environment also affects tRNA transcription. Due to the co-ordinated nature of 335
epigenomic modifications, it may also be revealing to evaluate ageing-related histone modification in these tRNA 336
loci. Furthermore, recent data from Gerber et al [87] indicates Pol-II may also have a dynamic regulatory role in 337
tRNA expression. 338

Changes in the epigenetic state of specific tRNA could be modulating transcription efficiency or even codon 339
availability in the ageing cell. tRNA gene dosage is quite closely matched to amino acid usage frequency in the 340
human exome. However, the transcriptome codon usage frequency and tRNA gene expression have been claimed 341
to vary with the replicative state of cells, separating differentiated from replicating cells [88]. Others have argued 342
that these differences are substantially explained by variation in GC content [89] and that codon usage 343
frequencies are observed to be mostly invariant in the transcriptomes of a wide range of tissues, as well as across 344
developmental time [86]. Although, experimental stress-related states have revealed changes with an 345

over-representation of codons that are translated by rare tRNAs [90].   346

tRNA sequences themselves are under strong structural (both secondary and tertiary) [72] as well as   347
functional constraint, which leads to an order of magnitude reduction in variation compared the background   348
genomic mutation rate [28]. Despite strong purifying selection maintaining very low variation in tRNA gene   349
bodies, tRNA genes are subject to high levels of transcription-associated mutagenesis (TAM) leading to elevated   350
mutation rates over evolutionary time in their immediate flanking sequences [91]. The possible effects of genetic   351
variation on DNA methylation mean that polymorphic tRNA could be another potential caveat to our work.   352
Although, there is no significant population variation in, for example, tRNA iMet sequences in 1,000 Genomes   353
data. Indeed, there are only 11 new isodecoder sequences with high confidence (tRNAscan scores ≥ 50) at $>1\%$ 354
population frequency [28]. There is also some evidence for tRNA copy number variation at specific loci, although   355
this remains under-characterised [92,93]. Another potential cause we considered was whether age-related somatic   356
copy number increases could be occurring in these loci. Population or somatic copy number expansions could   357
lead to increased methylated reads in MeDIP-seq without any epigenetic state change. However, this would not   358
be consistent with the targeted and array BiS conversion methodologies, where the proportion of methylated to   359
unmethylated reads would still be constant.   360

It is worth noting the parallels with known cancer and ageing epigenetic changes, and that tRNAs are also   361
dysregulated in cancer [94], with proposed utility as prognostic markers [95]. Furthermore, the early replicating   362
state of tRNA loci, potentially associated with high expression [96], may make them prone to hypermethylate, as   363
is observed in early replicating loci in both cancer [97] and senescent cells [98]. Interestingly, tRNA gene loci may   364
also play a role in local as well as large scale genome organisation [43]. tRNA gene clusters act as insulators [99]   365
and have extensive long-range chromatin interactions with other tRNA gene loci [43]. The coordinated   366
transcription of tRNAs at subnuclear foci and the B-box sequence elements bound by TFIIIC and not PolIII may   367
represent an organising principle for 3D-chromatin by providing spatial constraints [100]. Therefore, these tRNA   368
epigenetic changes could contribute to the structural changes that are also observed in ageing [101].   369

In conclusion, due to the unique challenges that make the tRNA genes difficult to examine it has remained   370
epigenetically under-characterised despite its critical importance for cell function. We directly interrogated the   371
epigenetic DNA methylation state of the functionally important tRNA genes, across the age spectrum in a range   372
of datasets as well as methodologies and identified an enrichment for age-related DNA hypermethylation in the   373
human tRNA genes.   374

Methods

Participants

Participants in the 'EpiTwins' study are adult volunteers from the TwinsUK Register. The participants were aged between 16 and 82 years, with a median of ˜55 years (cohort profile [102]). Ethics for the collection of these data were approved by Guy's & St Thomas' NHS Foundation Trust Ethics Committee (EC04/015—15-Mar-04) and written informed consent was obtained from all participants.

Participants for our targeted bisulfite sequencing of select tRNA loci were drawn from two studies. Samples from participants aged 4 and 28 years are from the MAVIDOS [103] study and participants aged 63 and 78 years are from the Hertfordshire cohort study [104]. Due to a limited number of available samples, the two 4 year old pools contained DNA from 20 individuals each, with all other pools having 25 contributing individuals. Pool 1, the first 4 year old pool used DNA from all male samples, with all other pools using all female samples. Thus, the total number of participants was 190 (see Table 1). Samples from the 28 year old time point are all from pregnant women at ˜11 weeks gestation.

tRNA annotation information

Genomic coordinates of the tRNA genes were downloaded from GtRNAdb [27]. The 2 tRNAs located in chr1_gl000192_random are tRNA-Gly-CCC-8-1 & tRNA-Asn-ATT-1-2 (Supplementary File S5). The 213 probes overlapping tRNA genes were derived from intersecting the tRNA gene annotation data from gtRNAdb with the Illumina 450k array manifest annotation for the hg19 genome build using `bedtools` v2.17.0 [105]. We excluded 107 tRNAs from blacklisted regions of hg19 [46].

tRNA Gene Clustering

We clustered the tRNA genes by grouping together all tRNAs within 5Mb of one another using the `bedtools merge` tool v2.17.0 [105]. We used a command of the form: `bedtools merge -c 4 -o collapse -d <N> -i hg19-tRNAs.bed` where `<N>` is the binsize. We varied the binsize and selected 5Mb as it is at approximately this size that number of clusters with more than one tRNAs exceeds the number of singleton tRNAs (Figure S9). We added the further requirements that these groupings contain at least 5 tRNA genes with a density of at least 5 tRNA genes per Mb to be considered as clusters.

DNA methylome data

TwinsUK MeDIP-seq methylomes

The Methylated DNA Immunoprecipitation sequencing (MeDIP-seq) data was processed as previously described [16,106]. These processed data are available from the European Genome-phenome Archive (EGA) (https://www.ebi.ac.uk/ega) under study number EGAS00001001910 and dataset EGAD00010000983. The dataset used in this work consists of 4350 whole blood methylomes with age data. 4054 are female and 270 male. 3001 have full blood counts. There are 3652 individuals in this data set. These individuals originate from 1933 unique families. There are 1234 monozygotic (MZ) twin pairs (2468 individuals), and 458 dizygotic (DZ) twin pairs (916 individuals).

MeDIP-seq used a monoclonal anti-5mC antibody to bind denatured fragmented genomic DNA at methylated CpG sites. This antibody-bound fraction of DNA was isolated and sequenced [44]. MeDIP-seq 50-bp single-end sequencing reads were aligned to the hg19/GRCh37 assembly of the human genome and duplicates were removed. `MEDIPS` (v1.0) was used for the MeDIP-seq specific analysis [107]. This produced reads per million base pairs (RPM) values binned into 500bp windows with a 250bp slide in the BED format, resulting in ~12.8 million windows on the genome. MeDIP-seq data from regions of interest was extracted using Bedtools v2.17.0 [105].

Analysis of DNA methylome data for Significant Ageing-related changes

All analysis was performed in R/3.5.2. Linear models were fitted to age using the MeDIP-seq DNA methylome data, as quantile normalised RPM scores at each 500bp window. Models were fitted with: 1. No covariates; 2. Batch information as a fixed effect; 3. Blood cell-type counts for neutrophils, monocytes, eosinophils, and lymphocytes as fixed effects; and 4. Batch and Blood Cell counts as fixed effects. Model 1 & 2 were fitted on the full set of 4350 as batch information was available for all samples but blood cell count data was only available for a subset of 3001 methylomes. Models 1 & 2 fitted in the n=3001 subset were similar to those fitted in the complete set of 4350. Models 3 & 4 were fitted in the n=3001 subset with full covariate information and sets of significant tRNAs identified at study-wide and genone wide levels in model 4 were used in subsequent analyses. Models were also fitted for two unrelated subsets created by selecting one twin from each pair (Monozygotic or Dizygotic), yeilding sets with n = 1198 & 1206 DNA methylomes. One additional model was fitted for longitudinal analysis, samples were selected by identifying individuals with a DNA methylome at more than one time point and filtering for only those with a minimum of 5 years between samples. This yielded 658 methylomes from 329 individuals with age differences of 5-16.1 yrs, median 7.6 yrs. Models for this set included participant identifier as a fixed effect in addition to blood cell counts and batch information.

Permutation Analysis for Enrichment with Age-related Changes

We performed a permutation analysis to determine whether the CpG distribution of sets of the tRNA genes was the principle driver of the ageing-related changes observed. Windows overlapping tRNAs have a higher proportion of windows with a greater CpG density than their surrounding sequences (see supplementary Figure S8). CpGs residing within moderate CpG density loci are the most dynamic in the genome [48] and CpG dense CpG island regions include specific ageing-related changes [8,9,16]. For comparison we also performed the permuation in the CGI regions from the Polycomb group protein target promoters in Teschendorff *et al.* [8] and bivalent loci from ENCODE ChromHmm 'Poised Promoter' classification in the GM12878 cell-line [54]. A random set of 500bp windows representing an equivalent CpG density distribution of the feature set in question were selected from the genome-wide data. Above a certain CpG density there are insufficient windows to sample without replacement within a permutation. Furthermore, above $\tilde{} \geq 18\%$ CpG density CpG Islands become consistently hypomethylated [108]. Therefore, all windows with a CpG density of $\geq 18\%$ (45 CpGs per 500bp) were grouped and sampled from the same pool. i.e. a window overlapping a tRNA gene which had a 20% density could be represented in permutation by one with any density $\geq 18\%$. This permutation was performed 1,000 times to determine an Empirical p value by calculating the number of times the permutation result exceeded the observed number of significant windows in the feature set. $Empirical\ p - value = \frac{r+1}{N+1}$, where r is the sum of significant windows in all permutations and N is number of permutations [109].

Neonate and Centenarian Whole Genome Bisulfite Sequencing

DNA methylation calls were downloaded from GEO:GSE31263 and intersected with tRNA genes using bedtools v2.17.0 [105].

Sample pooling and EPIC array

We performed an Illumina Infinium DNA methylation EPIC array ((C) Illumina) and targeted bisulfite sequencing of select tRNA gene loci. Here we used DNA extracted from whole blood and pooled into 8 samples from unrelated individuals at 4 time-points with 2 pools at each time-point. The timepoints were 4, 28, 63, and 78 years. Using the EPIC array we were able to infer the DNAm age using the Horvath DNAm clock [11] and blood cell-type composition of our samples using the Houseman algorithm [51].

Targeted Bisulfite Sequencing

We selected tRNA loci for targeted sequencing in which have had observed changes and DNAm with age and closely related tRNAs in which changes were not observed. Primer design was performed using 'methPrimer' [110] (Supplementary File S6). A total of 84 tRNA loci were targeted and 79 subsequently generated reliable results post-QC. The targeted tRNAs covered a total of 723 CpGs with a median of 8 CpGs per tRNA (range 1-13), data passing QC was generated for 458 CpGs, median 6 (range 1-9) per tRNA.

Quality was assessed before and after read trimming using `fastqc` [111] and `multiqc` [112] to visualise the results. Targeting primers were trimmed with `cutadapt` [113] and a custom `perl5` script. Quality trimming was performed with `trim_galore` [114]. Alignment and methylation calling was performed with `Bismark` (v0.20.0) [115] making use of `bowtie2` [116]. The alignment was performed against both the whole hg19 genome and just the tRNA genes +/- 100bp to assess the possible impact of off-target mapping. Mapping to the whole genome did produce purported methylation calls at a larger number of loci than mapping just to the tRNA genes (683,783 vs 45,861 respectively). Introducing a minimum coverage threshold of 25 reads dramatically reduced this and brought the number of sites into line with that in the tRNA gene set (36,065 vs 33,664 respectively) suggesting a small number of ambiguously mapping reads. All subsequent analysis was performed using the alignment to just the tRNA genes with a minimum coverage of 25 reads.

We performed pairwise differential methylation analysis of the tRNA genes at the different time points using `RnBeads` [52] with `limma` [117] and a minimum coverage of 25 reads. We also performed linear regression predicting age from DNA methylation at the targeted tRNA sites, permitting us to compare rates of increase with age. For the linear regression, we used only CpG sites with more than 25 reads mapped to the regions of the genome targeted for amplification.

TwinsUK Illumina 450k array methylomes

Illumina Infinium DNA methylation 450k arrays ((C) Illumina) were also performed on TwinsUK participants, in 770 Blood-derived DNA samples which had matched MeDIP-seq data. These data were preprocessed in the form of methylation 'beta' values pre-processed as previously described [16,106]. Cell-type correction was performed using cell-count data and the following model: `lm(age ~ beta + eosinophils + lymphocytes + monocytes + neutrophils)`.

Chromatin Segmentation Data

Epilogos chromatin segmentation data [53] was downloaded for the tRNA gene regions +/- 200bp from https://explore.altius.org/tabix/epilogos/hg19.15.Blood_T-cell.KL.gz using the `tabix` utility. The data used was the 'Blood & T-cell' 15 State model based on segmentation of 14 cell-types. This data was manipulated and visualised with `R` and `ggplot2`.

Isolated Blood Cell Type Specific Data

Data from 7 cell-type fractions from 6 Male individuals was downloaded from GSE35069 [55] using `GEOquery` [118]. Five of the 6 top age hypermethylating tRNAs are covered by this array dataset. Heatmaps were created with the `ComplexHeatmap` R package [56].

Cancer and Tissue Specific Methylation Data

Data was downloaded from the TCGA (The Cancer Genome Atlas) via the GDC (genomic data commons) data portal [119] using the `GenomicDataCommons` R package. Data from foetal tissue [61,62] was downloaded from GEO (GSE72867, GSE30654). From the TCGA, we selected samples for which DNAm data was available from both the primary site and normal solid tissue, and for which we could infer an approximate age (within one year). We selected those probes overlapping tRNA genes yielding 73,403 data points across 19 tissues with an age range of 15-90yrs (median 63.4) (Supplementary File S7)

Mouse RRBS Analysis

We downloaded methylation calls and coverage information resulting from RRBS performed by Petkovich *et al.* [63] from GEO using `GEOquery` [118] GSE80672. These data from 152 mice covered 68 tRNA and 436 CpGs after QC requiring >50 reads per CpG and >10 data points per tRNA. We excluded 5 tRNAs from blacklisted regions of mm10 [46]. After QC there were 58 tRNA genes and 385 CpGs. We performed simple linear modeling to predict age from methylation level at each tRNA and each CpG.

Data availability

The MeDIP-seq data supporting the results of this article are available in the EMBL-EBI European Genome-phenome Archive (EGA) under Data set Accession number EGAD00010000983 (`https://www.ebi.ac.uk/ega/datasets/EGAD00010000983`). The targeted BiS-sequencing data is available at: `https://www.ncbi.nlm.nih.gov/bioproject/PRJNA635108`

Code availablity

Available at `https://github.com/RichardJActon/tRNA_paper_code`

Acknowledgements 513

We gratefully acknowlege the individuals from TwinsUK, Mavidos and the Hertfordshrie cohort. TwinsUK $_{514}$ received funding from the Wellcome Trust (Ref: 081878/Z/06/Z), European Community's Seventh Framework $_{515}$ Programme (FP7/2007-2013), the National Institute for Health Research (NIHR)-funded BioResource, Clinical $_{516}$ Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in $_{517}$ partnership with King's College London. Further funding support for the EpiTwin project was obtained from the $_{518}$ European Research Council (project number 250157) and BGI. SNP Genotyping was performed by The $_{519}$ Wellcome Trust Sanger Institute and National Eye Institute via NIH/CIDR. The authors would like to thank $_{520}$ Nikki Graham for her assistance with the identification and pooling of the MAVIDOS and Hertfordshire DNA $_{521}$ samples. The authors also acknowledge the use of the IRIDIS High Performance Computing Facility, and $_{522}$ associated support services at the University of Southampton, in the completion of this work. The MRC-LEU is $_{523}$ supported by the Medical Research Council (MRC). CGB recieved support from Diabetes UK (16/0005454). $_{524}$ RJA was in receipt of a MRC Doctoral fund (1820097). $_{525}$

Author Contributions ⁢526

RJA designed experiments and analysed all the processed and experimental data. CGB conceived and designed $_{527}$ the experiments. TDS, KW and JW conceived and provided TwinsUK MeDIP-seq data. YX, FG and JW $_{528}$ produced raw MeDIP-seq data with WY and JB processing and quality controlling these data. WY contributed $_{529}$ an analysis concept. CC, NH, ED, and KL provided MAVIDOS and Hertfordshire sample data. EB, EW, and $_{530}$ CAM performed the targeted BiS sequencing experiment. PGH contributed additional data and discussion of $_{531}$ results. RJA and CGB wrote the paper. All authors reviewed and approved the final manuscript. $_{532}$

References

1. Partridge L, Deelen J, Slagboom PE. Facing up to the global challenges of ageing. Nature. Springer US; 2018;561: 45–56. doi:10.1038/s41586-018-0457-8

2. Campisi J, Kapahi P, Lithgow GJ, Melov S, Newman JC, Verdin E. From discoveries in ageing research to therapeutics for healthy ageing. Nature. Springer US; 2019;571: 183–192. doi:10.1038/s41586-019-1365-2

3. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. Cell. 2013;153: 1194–217. doi:10.1016/j.cell.2013.05.039

4. Booth LN, Brunet A. The Aging Epigenome. Molecular Cell. Elsevier Inc. 2016;62: 728–744. doi:10.1016/j.molcel.2016.05.013

5. Wilson VL, Jones PA. DNA methylation decreases in aging but not in immortal cells. Science (New York, NY). 1983;220: 1055–7. Available: http://www.ncbi.nlm.nih.gov/pubmed/6844925

6. Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, et al. From The Cover: Epigenetic differences arise during the lifetime of monozygotic twins. Proceedings of the National Academy of Sciences. 2005;102: 10604–10609. doi:10.1073/pnas.0500398102

7. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from conflicts to benefits. Nature Reviews Genetics. Nature Publishing Group; 2017;18: 71–86. doi:10.1038/nrg.2016.139

8. Teschendorff AE, Menon U, Gentry-Maharaj A, Ramus SJ, Weisenberger DJ, Shen H, et al. Age-dependent DNA methylation of genes that are suppressed in stem cells is a hallmark of cancer. Genome research. 2010;20: 440–6. doi:10.1101/gr.103606.109

9. Rakyan VK, Down TA, Maslau S, Andrew T, Yang TP, Beyan H, et al. Human aging-associated DNA hypermethylation occurs preferentially at bivalent chromatin domains. Genome Research. 2010;20: 434–439. doi:10.1101/gr.103101.109

10. Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, et al. Genome-wide methylation profiles reveal quantitative views of human aging rates. Molecular cell. Elsevier Inc. 2013;49: 359–367. doi:10.1016/j.molcel.2012.10.016

11. Horvath S. DNA methylation age of human tissues and cell types. Genome biology. 2013;14: R115. doi:10.1186/gb-2013-14-10-r115

12. Weidner CI, Lin Q, Koch CM, Eisele L, Beier F, Ziegler P, et al. Aging of blood can be tracked by DNA methylation changes at just three CpG sites. Genome biology. 2014;15: R24. doi:10.1186/gb-2014-15-2-r24

13. Bell CG, Lowe R, Adams PD, Baccarelli AA, Beck S, Bell JT, et al. DNA methylation aging clocks: challenges and recommendations. Genome biology. 2019;20: 249. doi:10.1186/s13059-019-1824-y

14. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of ageing. Nature Reviews Genetics. Springer US; 2018;19: 371–384. doi:10.1038/s41576-018-0004-3

15. Field AE, Robertson NA, Wang T, Havas A, Ideker T, Adams PD. DNA Methylation Clocks in Aging: Categories, Causes, and Consequences. Molecular Cell. Elsevier Inc. 2018;71: 882–895. doi:10.1016/j.molcel.2018.08.008

16. Bell CG, Xia Y, Yuan W, Gao F, Ward K, Roos L, et al. Novel regional age-associated DNA methylation changes within human common disease-associated loci. Genome Biology. 2016;17: 193. doi:10.1186/s13059-016-1051-8

17. Rideout EJ, Marshall L, Grewal SS. Drosophila RNA polymerase III repressor Maf1 controls body size and developmental timing by modulating tRNAiMet synthesis and systemic insulin signaling. Proceedings of the National Academy of Sciences. 2012;109: 1139–1144. doi:10.1073/pnas.1113311109

18. Kolitz SE, Lorsch JR. Eukaryotic initiator tRNA: Finely tuned and ready for action. FEBS Letters. 2010;584: 396–404. doi:10.1016/j.febslet.2009.11.047

19. Pavon-Eternod M, Gomes S, Rosner MR, Pan T. Overexpression of initiator methionine tRNA leads to global reprogramming of tRNA expression and increased proliferation in human epithelial cells. RNA. 2013;19: 461–466. doi:10.1261/rna.037507.112

20. Eigen M, Lindemann B, Tietze M, Winkler-Oswatitsch R, Dress A, Haeseler A von. How old is the genetic code? Statistical geometry of tRNA provides an answer. Science. 1989;244: 673–679. doi:10.1126/science.2497522

21. Tavernarakis N. Ageing and the regulation of protein synthesis: a balancing act? Trends in Cell Biology. 2008;18: 228–235. doi:10.1016/j.tcb.2008.02.004

22. Pliatsika V, Loher P, Magee R, Telonis AG, Londin E, Shigematsu M, et al. MINTbase v2.0: a comprehensive database for tRNA-derived fragments that includes nuclear and mitochondrial fragments from all The Cancer Genome Atlas projects. Nucleic Acids Research. Oxford University Press; 2018;46: D152–D159. doi:10.1093/nar/gkx1075

23. Schimmel P. The emerging complexity of the tRNA world: mammalian tRNAs beyond protein synthesis. Nature reviews Molecular cell biology. Nature Publishing Group; 2018;19: 45–58. doi:10.1038/nrm.2017.77

24. Lee YS, Shibata Y, Malhotra A, Dutta A. A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). Genes & development. 2009;23: 2639–49. doi:10.1101/gad.1837609

25. Li S, Xu Z, Sheng J. tRNA-Derived Small RNA: A Novel Regulatory Small Non-Coding RNA. Genes. 2018;9: 246. doi:10.3390/genes9050246

26. Xu W-L, Yang Y, Wang Y-D, Qu L-H, Zheng L-L. Computational Approaches to tRNA-Derived Small RNAs. Non-Coding RNA. 2017;3: 2. doi:10.3390/ncrna3010002

27. Chan PP, Lowe TM. GtRNAdb: a database of transfer RNA genes detected in genomic sequence. Nucleic acids research. 2009;37: D93–7. doi:10.1093/nar/gkn787

28. Parisien M, Wang X, Pan T. Diversity of human tRNA genes from the 1000-genomes project. RNA Biology. 2013;10: 1853–1867. doi:10.4161/rna.27361

29. Lodish H, Berk A, Zipursky SL, Matsudaira P, Baltimore D, Darnell J. Molecular Cell Biology, 4th edition [Internet]. 4th ed. New York: W. H. Freeman; 2000. Available: https://www.ncbi.nlm.nih.gov/books/NBK21475/

30. Schramm L. Recruitment of RNA polymerase III to its target promoters. Genes & Development. 2002;16: 2593–2620. doi:10.1101/gad.1018902

31. Canella D, Praz V, Reina JH, Cousin P, Hernandez N. Defining the RNA polymerase III transcriptome: Genome-wide localization of the RNA polymerase III transcription machinery in human cells. Genome research. 2010;20: 710–21. doi:10.1101/gr.101337.109

32. Besser D, Götz F, Schulze-Forster K, Wagner H, Kröger H, Simon D. DNA methylation inhibits transcription by RNA polymerase III of a tRNA gene, but not of a 5S rRNA gene. FEBS letters. 1990;269: 358–62. doi:10.1016/0014-5793(90)81193-R

33. Varshney D, Vavrova-Anderson J, Oler AJ, Cowling VH, Cairns BR, White RJ. SINE transcription by RNA polymerase III is suppressed by histone methylation but not by DNA methylation. Nature communications. Nature Publishing Group; 2015;6: 6569. doi:10.1038/ncomms7569

34. Murawski M, Szczesniak B, Zoladek T, Hopper AK, Martin NC, Boguta M. maf1 mutation alters the subcellular localization of the Mod5 protein in yeast. Acta biochimica Polonica. 1994;41: 441–8. Available: http://www.ncbi.nlm.nih.gov/pubmed/7732762

35. Pluta K, Lefebvre O, Martin NC, Smagowicz WJ, Stanford DR, Ellis SR, et al. Maf1p, a Negative Effector of RNA Polymerase III in Saccharomyces cerevisiae. Molecular and Cellular Biology. 2001;21: 5031–5040. doi:10.1128/MCB.21.15.5031-5040.2001

36. Mange F, Praz V, Migliavacca E, Willis IM, Schütz F, Hernandez N. Diurnal regulation of RNA polymerase III transcription is under the control of both the feeding–fasting response and the circadian clock. Genome Research. 2017;27: 973–984. doi:10.1101/gr.217521.116

37. Kennedy BK, Lamming DW. The Mechanistic Target of Rapamycin: The Grand ConducTOR of Metabolism and Aging. Cell Metabolism. Elsevier Inc. 2016;23: 990–1003. doi:10.1016/j.cmet.2016.05.009

38. Nwanaji-Enwerem JC, Weisskopf MG, Baccarelli AA. Multi-tissue DNA methylation age: Molecular relationships and perspectives for advancing biomarker utility. Ageing Research Reviews. Elsevier; 2018;45: 15–23. doi:10.1016/j.arr.2018.04.005

39. Hansen M, Taubert S, Crawford D, Libina N, Lee S-J, Kenyon C. Lifespan extension by conditions that inhibit translation in Caenorhabditis elegans. Aging Cell. 2007;6: 95–110. doi:10.1111/j.1474-9726.2006.00267.x

40. Filer D, Thompson MA, Takhaveev V, Dobson AJ, Kotronaki I, Green JWM, et al. RNA polymerase III limits longevity downstream of TORC1. Nature. Nature Publishing Group; 2017;552: 263–267. doi:10.1038/nature25007

41. Dhahbi JM, Spindler SR, Atamna H, Yamakawa A, Boffelli D, Mote P, et al. 5' tRNA halves are present as abundant complexes in serum, concentrated in blood cells, and modulated by aging and calorie restriction. BMC Genomics. 2013;14: 298. doi:10.1186/1471-2164-14-298

42. Zhou W, Laird PW, Shen H. Comprehensive characterization, annotation and innovative use of Infinium DNA methylation BeadChip probes. Nucleic Acids Research. 2016;45: gkw967. doi:10.1093/nar/gkw967

43. Van Bortle K, Phanstiel DH, Snyder MP. Topological organization and dynamic regulation of human tRNA genes during macrophage differentiation. Genome Biology. Genome Biology; 2017;18: 180. doi:10.1186/s13059-017-1310-3

44. Down TA, Rakyan VK, Turner DJ, Flicek P, Li H, Kulesha E, et al. A Bayesian deconvolution strategy for immunoprecipitation-based DNA methylome analysis. Nature Biotechnology. 2008;26: 779–785. doi:10.1038/nbt1414

45. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. Nucleic Acids Research. 2016;44: W54–W57. doi:10.1093/nar/gkw413

46. Amemiya HM, Kundaje A, Boyle AP. The ENCODE Blacklist: Identification of Problematic Regions of the Genome. Scientific Reports. 2019;9: 9354. doi:10.1038/s41598-019-45839-z

47. Geiger H, Haan G de, Florian MC. The ageing haematopoietic stem cell compartment. Nature Reviews Immunology. Nature Publishing Group; 2013;13: 376–389. doi:10.1038/nri3433

48. Ziller MJ, Gu H, Müller F, Donaghey J, Tsai LT-Y, Kohlbacher O, et al. Charting a dynamic DNA methylation landscape of the human genome. Nature. 2013;500: 477–81. doi:10.1038/nature12433

49. Christensen BC, Houseman EA, Marsit CJ, Zheng S, Wrensch MR, Wiemels JL, et al. Aging and Environmental Exposures Alter Tissue-Specific DNA Methylation Dependent upon CpG Island Context. Schübeler D, editor. PLoS Genetics. 2009;5: e1000602. doi:10.1371/journal.pgen.1000602

50. Heyn H, Li N, Ferreira HJ, Moran S, Pisano DG, Gomez A, et al. Distinct DNA methylomes of newborns and centenarians. Proceedings of the National Academy of Sciences. 2012;109: 10522–10527. doi:10.1073/pnas.1120658109

51. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al. DNA methylation arrays as surrogate measures of cell mixture distribution. BMC Bioinformatics. 2012;13: 86.

doi:10.1186/1471-2105-13-86

52. Müller F, Scherer M, Assenov Y, Lutsik P, Walter J, Lengauer T, et al. RnBeads 2.0: comprehensive analysis of DNA methylation data. Genome Biology. Genome Biology; 2019;20: 55. doi:10.1186/s13059-019-1664-9

53. Meuleman W. Epilogos [Internet]. 2019. Available: https://epilogos.altius.org/%20https://github.com/Altius/epilogos

54. Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, et al. Mapping and analysis of chromatin state dynamics in nine human cell types. Nature. 2011;473: 43–49. doi:10.1038/nature09906

55. Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlén S-E, Greco D, et al. Differential DNA Methylation in Purified Human Blood Cells: Implications for Cell Lineage and Studies on Disease Susceptibility. Ting AH, editor. PLoS ONE. 2012;7: e41361. doi:10.1371/journal.pone.0041361

56. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics. 2016;32: 2847–2849. doi:10.1093/bioinformatics/btw313

57. Schmitt BM, Rudolph KLM, Karagianni P, Fonseca NA, White RJ, Talianidis I, et al. High-resolution mapping of transcriptional dynamics across tissue development reveals a stable mRNA–tRNA interface. Genome Research. 2014;24: 1797–1807. doi:10.1101/gr.176784.114

58. Dittmar KA, Goodenbour JM, Pan T. Tissue-specific differences in human transfer RNA expression. PLoS genetics. 2006;2: e221. doi:10.1371/journal.pgen.0020221

59. Sagi D, Rak R, Gingold H, Adir I, Maayan G, Dahan O, et al. Tissue- and Time-Specific Expression of Otherwise Identical tRNA Genes. Chisholm AD, editor. PLOS Genetics. 2016;12: e1006264. doi:10.1371/journal.pgen.1006264

60. Slieker RC, Iterson M van, Luijk R, Beekman M, Zhernakova DV, Moed MH, et al. Age-related accrual of methylomic variability is linked to fundamental ageing mechanisms. Genome biology. Genome Biology; 2016;17: 191. doi:10.1186/s13059-016-1053-6

61. Yang Z, Wong A, Kuh D, Paul DS, Rakyan VK, Leslie RD, et al. Correlation of an epigenetic mitotic clock with cancer risk. Genome Biology. Genome Biology; 2016;17: 205. doi:10.1186/s13059-016-1064-3

62. Nazor KL, Altun G, Lynch C, Tran H, Harness JV, Slavin I, et al. Recurrent Variations in DNA Methylation in Human Pluripotent Stem Cells and Their Differentiated Derivatives. Cell Stem Cell. 2012;10: 620–634. doi:10.1016/j.stem.2012.02.013

63. Petkovich DA, Podolskiy DI, Lobanov AV, Lee S-G, Miller RA, Gladyshev VN. Using DNA Methylation Profiling to Evaluate Biological Age and Longevity Interventions. Cell Metabolism. 2017;25: 954–960.e6. doi:10.1016/j.cmet.2017.03.016

64. Thornlow BP, Armstrong J, Holmes AD, Howard JM, Corbett-Detig RB, Lowe TM. Predicting transfer RNA gene activity from sequence and genome context. Genome Research. 2020;30: 85–94. doi:10.1101/gr.256164.119

65. Ehrlich M. DNA hypermethylation in disease: mechanisms and clinical relevance. Epigenetics. Taylor & Francis; 2019;14: 1141–1163. doi:10.1080/15592294.2019.1638701

66. Xu Z, Taylor JA. Genome-wide age-related DNA methylation changes in blood and other tissues relate to histone modification, expression and cancer. Carcinogenesis. 2014;35: 356–364. doi:10.1093/carcin/bgt391

67. Slieker RC, Relton CL, Gaunt TR, Slagboom PE, Heijmans BT. Age-related DNA methylation changes are tissue-specific with ELOVL2 promoter methylation as exception. Epigenetics and Chromatin. BioMed Central; 2018;11: 1–11. doi:10.1186/s13072-018-0191-3

68. Zhu T, Zheng SC, Paul DS, Horvath S, Teschendorff AE. Cell and tissue type independent age-associated DNA methylation changes are not rare but common. Aging. 2018;10: 3541–3557. doi:10.18632/aging.101666

69. Lappalainen T, Greally JM. Associating cellular epigenetic models with human phenotypes. Nature Reviews Genetics. Nature Publishing Group; 2017;18: 441–451. doi:10.1038/nrg.2017.32

70. Moskowitz DM, Zhang DW, Hu B, Le Saux S, Yanes RE, Ye Z, et al. Epigenomics of human CD8 T cell differentiation and aging. Science Immunology. 2017;2: eaag0192. doi:10.1126/sciimmunol.aag0192

71. Hashimoto K, Kouno T, Ikawa T, Hayatsu N, Miyajima Y, Yabukami H, et al. Single-cell transcriptomics reveals expansion of cytotoxic CD4 T cells in supercentenarians. Proceedings of the National Academy of Sciences of the United States of America. 2019;116: 24242–24251. doi:10.1073/pnas.1907883116

72. Goodenbour JM, Pan T. Diversity of tRNA genes in eukaryotes. Nucleic acids research. 2006;34: 6137–46. doi:10.1093/nar/gkl725

73. Keam SP, Young PE, McCorkindale AL, Dang THY, Clancy JL, Humphreys DT, et al. The human Piwi protein Hiwi2 associates with tRNA-derived piRNAs in somatic cells. Nucleic Acids Research. 2014;42: 8984–8995. doi:10.1093/nar/gku620

74. Geslain R, Pan T. Functional Analysis of Human tRNA Isodecoders. Journal of Molecular Biology. 2010;396: 821–831. doi:10.1016/j.jmb.2009.12.018

75. Kirchner S, Cai Z, Rauscher R, Kastelic N, Anding M, Czech A, et al. Alteration of protein function by a silent polymorphism linked to tRNA abundance. Hurst L, editor. PLOS Biology. 2017;15: e2000779. doi:10.1371/journal.pbio.2000779

76. Li S, Shi X, Chen M, Xu N, Sun D, Bai R, et al. Angiogenin promotes colorectal cancer metastasis via tiRNA production. International Journal of Cancer. 2019; ijc.32245. doi:10.1002/ijc.32245

77. Kim HK, Fuchs G, Wang S, Wei W, Zhang Y, Park H, et al. A transfer-RNA-derived small RNA

regulates ribosome biogenesis. Nature. Nature Publishing Group; 2017;552: 57–62. doi:10.1038/nature25005   724

78. Martinez G, Choudury SG, Slotkin RK. tRNA-derived small RNAs target transposable element   725
transcripts. Nucleic Acids Research. 2017;45: 5142–5152. doi:10.1093/nar/gkx103   726

79. Torres AG, Reina O, Stephan-Otto Attolini C, Ribas de Pouplana L. Differential expression of human   727
tRNA genes drives the abundance of tRNA-derived fragments. Proceedings of the National Academy of Sciences.   728
2019;116: 201821120. doi:10.1073/pnas.1821120116   729

80. Gu P, Jia S, Takasugi T, Smith E, Nandakumar J, Hendrickson E, et al. CTC1-STN1 coordinates G- and   730
C-strand synthesis to regulate telomere length. Aging Cell. 2018;17: e12783. doi:10.1111/acel.12783   731

81. Wang Y, Brady KS, Caiello BP, Ackerson SM, Stewart JA. Human CST suppresses origin licensing and   732
promotes AND-1/Ctf4 chromatin association. Life Science Alliance. 2019;2: e201800270.   733
doi:10.26508/lsa.201800270   734

82. Sargolzaeiaval F, Zhang J, Schleit J, Lessel D, Kubisch C, Precioso DR, et al. CTC1 mutations in a   735
Brazilian family with progeroid features and recurrent bone fractures. Molecular Genetics & Genomic Medicine.   736
2018;6: 1148–1156. doi:10.1002/mgg3.495   737

83. Gomez-Roman N, Grandori C, Eisenman RN, White RJ. Direct activation of RNA polymerase III   738
transcription by c-Myc. Nature. 2003;421: 290–294. doi:10.1038/nature01327   739

84. Crighton D. p53 represses RNA polymerase III transcription by targeting TBP and inhibiting promoter   740
occupancy by TFIIIB. The EMBO Journal. 2003;22: 2810–2820. doi:10.1093/emboj/cdg265   741

85. Sutcliffe JE, Brown TRP, Allison SJ, Scott PH, White RJ. Retinoblastoma Protein Disrupts Interactions   742
Required for RNA Polymerase III Transcription. Molecular and Cellular Biology. 2000;20: 9192–9202.   743
doi:10.1128/MCB.20.24.9192-9202.2000   744

86. Schmitt BM, Rudolph KLM, Karagianni P, Fonseca NA, White RJ, Talianidis I, et al. High-resolution   745
mapping of transcriptional dynamics across tissue development reveals a stable mRNA-tRNA interface. Genome   746
research. 2014;24: 1797–807. doi:10.1101/gr.176784.114   747

87. Gerber A, Ito K, Chu C-S, Roeder RG. Gene-Specific Control of tRNA Expression by RNA Polymerase II.   748
Molecular Cell. Elsevier Inc. 2020;78: 765–778.e7. doi:10.1016/j.molcel.2020.03.023   749

88. Gingold H, Tehler D, Christoffersen NR, Nielsen MM, Asmar F, Kooistra SM, et al. A Dual Program for   750
Translation Regulation in Cellular Proliferation and Differentiation. Cell. Elsevier Inc. 2014;158: 1281–1292.   751
doi:10.1016/j.cell.2014.08.011   752

89. Rudolph KLM, Schmitt BM, Villar D, White RJ, Marioni JC, Kutter C, et al. Codon-Driven   753
Translational Efficiency Is Stable across Diverse Mammalian Cell States. Galtier N, editor. PLoS genetics.   754
2016;12: e1006024. doi:10.1371/journal.pgen.1006024   755

90. Gingold H, Dahan O, Pilpel Y. Dynamic changes in translational efficiency are deduced from codon usage of the transcriptome. Nucleic acids research. 2012;40: 10053–63. doi:10.1093/nar/gks772

91. Thornlow BP, Hough J, Roger JM, Gong H, Lowe TM, Corbett-Detig RB. Transfer RNA genes experience exceptionally elevated mutation rates. Proceedings of the National Academy of Sciences. 2018;115: 8996–9001. doi:10.1073/pnas.1801240115

92. Iben JR, Maraia RJ. tRNA gene copy number variation in humans. Gene. 2014;536: 376–384. doi:10.1016/j.gene.2013.11.049

93. Darrow EM, Chadwick BP. A novel tRNA variable number tandem repeat at human chromosome 1q23.3 is implicated as a boundary element based on conservation of a CTCF motif in mouse. Nucleic Acids Research. 2014;42: 6421–6435. doi:10.1093/nar/gku280

94. Huang S-q, Sun B, Xiong Z-p, Shu Y, Zhou H-h, Zhang W, et al. The dysregulation of tRNAs and tRNA derivatives in cancer. Journal of Experimental & Clinical Cancer Research. Journal of Experimental & Clinical Cancer Research; 2018;37: 101. doi:10.1186/s13046-018-0745-z

95. Krishnan P, Ghosh S, Wang B, Heyns M, Li D, Mackey JR, et al. Genome-wide profiling of transfer RNAs and their role as novel prognostic markers for breast cancer. Scientific Reports. Nature Publishing Group; 2016;6: 32843. doi:10.1038/srep32843

96. Müller CA, Nieduszynski CA. DNA replication timing influences gene expression level. The Journal of Cell Biology. 2017;216: 1907–1914. doi:10.1083/jcb.201701061

97. Du Q, Bert SA, Armstrong NJ, Caldon CE, Song JZ, Nair SS, et al. Replication timing and epigenome remodelling are associated with the nature of chromosomal rearrangements in cancer. Nature Communications. Springer US; 2019;10: 416. doi:10.1038/s41467-019-08302-1

98. Cruickshanks HA, McBryan T, Nelson DM, VanderKraats ND, Shah PP, Tuyn J van, et al. Senescent cells harbour features of the cancer epigenome. Nature Cell Biology. 2013;15: 1495–1506. doi:10.1038/ncb2879

99. Raab JR, Chiu J, Zhu J, Katzman S, Kurukuti S, Wade PA, et al. Human tRNA genes function as chromatin insulators. The EMBO journal. Nature Publishing Group; 2012;31: 330–50. doi:10.1038/emboj.2011.406

100. Noma K-i, Cam HP, Maraia RJ, Grewal SIS. A role for TFIIIC transcription factor complex in genome organization. Cell. 2006;125: 859–72. doi:10.1016/j.cell.2006.04.028

101. Sun L, Yu R, Dang W. Chromatin Architectural Changes during Cellular Senescence and Aging. Genes. 2018;9: 211. doi:10.3390/genes9040211

102. Moayyeri A, Hammond CJ, Valdes AM, Spector TD. Cohort Profile: TwinsUK and Healthy Ageing Twin Study. International Journal of Epidemiology. 2013;42: 76–85. doi:10.1093/ije/dyr207

103. Harvey NC, Javaid K, Bishop N, Kennedy S, Papageorghiou AT, Fraser R, et al. MAVIDOS Maternal Vitamin D Osteoporosis Study: study protocol for a randomized controlled trial. The MAVIDOS Study Group. Trials. 2012;13: 13. doi:10.1186/1745-6215-13-13

104. Syddall H, Aihie Sayer A, Dennison E, Martin H, Barker D, Cooper C. Cohort Profile: The Hertfordshire Cohort Study. International Journal of Epidemiology. 2005;34: 1234–1242. doi:10.1093/ije/dyi127

105. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. University of Utah; 2010;26: 841–842. doi:10.1093/bioinformatics/btq033

106. Bell CG, Gao F, Yuan W, Roos L, Acton RJ, Xia Y, et al. Obligatory and facilitative allelic variation in the DNA methylome within common disease-associated loci. Nature Communications. 2018;9: 8. doi:10.1038/s41467-017-01586-1

107. Lienhard M, Grimm C, Morkel M, Herwig R, Chavez L. MEDIPS: genome-wide differential coverage analysis of sequencing data derived from DNA enrichment experiments. Bioinformatics. 2014;30: 284–286. doi:10.1093/bioinformatics/btt650

108. Bell CG, Wilson GA, Butcher LM, Roos C, Walter L, Beck S. Human-specific CpG "beacons" identify loci associated with human-specific traits and disease. Epigenetics. 2012;7: 1188–99. doi:10.4161/epi.22127

109. North BV, Curtis D, Sham PC. A Note on the Calculation of Empirical P Values from Monte Carlo Procedures. The American Journal of Human Genetics. 2003;72: 498–499. doi:10.1086/346173

110. Li L-C, Dahiya R. MethPrimer: designing primers for methylation PCRs. Bioinformatics (Oxford, England). 2002;18: 1427–31. Available: http://www.ncbi.nlm.nih.gov/pubmed/12424112

111. Andrews S. FastQC [Internet]. Cambridge: Babraham Bioinformatics; 2010. Available: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

112. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics (Oxford, England). 2016;32: 3047–8. doi:10.1093/bioinformatics/btw354

113. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnetjournal. 2011;17: 10. doi:10.14806/ej.17.1.200

114. Krueger F. Trim Galore [Internet]. 2015. Available: https://www.bioinformatics.babraham.ac.uk/projects/trim%7B/_%7Dgalore/

115. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. Bioinformatics. 2011;27: 1571–1572. doi:10.1093/bioinformatics/btr167

116. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature methods. 2012;9: 357–9. doi:10.1038/nmeth.1923

117. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research. 2015;43: e47–e47. doi:10.1093/nar/gkv007

118. Sean D, Meltzer PS. GEOquery: A bridge between the Gene Expression Omnibus (GEO) and BioConductor. Bioinformatics. 2007;23: 1846–1847. doi:10.1093/bioinformatics/btm254

119. Grossman RL, Heath AP, Ferretti V, Varmus HE, Lowy DR, Kibbe WA, et al. Toward a Shared Vision for Cancer Genomic Data. New England Journal of Medicine. 2016;375: 1109–1112. doi:10.1056/NEJMp1607591

120. Derrien T, Estellé J, Marco Sola S, Knowles DG, Raineri E, Guigó R, et al. Fast Computation and Applications of Genome Mappability. Ouzounis CA, editor. PLoS ONE. 2012;7: e30377. doi:10.1371/journal.pone.0030377

Supplementary materials

Supplementary Figures

Fig S1. Example of mappability data from the encode mappability tracks [120] for the initiator methionine tRNA genes.

Fig S2. Mappability score density of the tRNA genes increases with read length and is greater when flanking regions ($\pm 500bp$) are included. Mappability score density is computed as the area under the encode mappability tracks [120] over the length of the region.

Fig S3. Human tRNA genes overview. From the outside in: Chromosome ideograms scaled by the number of tRNA genes (total = 598), as excludes chromosome X (10), Y (0) and contig chr1_gl000192_random (2; see Methods). tRNA genes within 20kbp of one another are grouped with breaks inserted between these clusters. Radial grey lines represent the location of tRNA genes in the genome. $-log_{10}(p-value)$ for the blood cell-type and batch corrected age model are shown for each window overlapping a tRNA gene in green. Mean methylation across all samples (n=3001) in RPM (reads per million base pairs) is shown in blue. Genome-wide significant cell-type & batch corrected ($p < 4.34 \times 10^{-9}$) tRNAs show in red. The 158 Loci covered by 213 probes on the 450k array which directly overlap a tRNA gene are shown with green triangles. The 84 loci targeted for bisulfite sequencing in this study are indicated in magenta. Mappability score density is computed as the area under the encode mappability tracks [120] over the length of the region.

Fig S4. Heatmap Mean Methylation of probes covering each tRNA in 7 cell-type fractions from 6 Male individuals. Showing all 150 tRNAs covered by 213 probes on the Illumina 450k array. Data from GSE35069 [55] downloaded using `GEOquery` [56,118].

Fig S5. Global properties of tRNA methylation data for 45 tRNA genes across 19 tissues with matched normal and tumour samples from 733 cases in TCGA [61,62].

Fig S6. Mean Methylation of 43 tRNAs in 19 tissues. Possible pseudogene (tRNA-Asn-ATT-1-1) is shown in a separate cluster beneath the main heatmap [56].

Fig S7. Mean Methylation of 115 tRNAs in 11 tissues. Possible pseudogenes are shown in a separate cluster beneath the main heatmap [56].

Fig S8. CpG Density in windows overlapping tRNA genes compared to that of non-tRNA overlapping windows in flanking sequences (+/-5kb)

Fig S9. tRNA gene cluster numbers at different bin sizes total: total number of tRNA clusters. singletons: number of tRNAs in clusters alone. moreThan1: number of tRNAs in clusters with more than one tRNA. maxSize: the number of tRNAs in the largest cluster.

Mean methylation by tRNA
In Solid Tissue Normal Samples

Mean methylation by tRNA
Sorted Blood Cell-types (GSE35069)

$-\log_{10}$(p-value) Age Model

Mean Methylation (MeDIP-seq /RPM)

Mappability
* tRNA only (50mer)
+ tRNA ± 500bp (50 mer)

0.0
0.5
1.0

● Targeted BiS-seq
▼ 450k array

tRNA mappability

tRNA region only Vs tRNA +/- 500bp

Mappability Score Density = total score / length of region

Encode Mapability data for iMet isoacceptor tRNA genes (Page 1 of 1)

Age-related DNA methylation change in tRNA genes

Twins UK - Illumina 450k array

A

Age-hyper tRNA gene count proportionate to cluster size

B WGBS for age-hypermethylating tRNA Genes

A Significant Age Related Hypermethylation

Percentage of significant windows by feature type

B CpG Density Permutation Analysis

tRNA	Window	MeDIP		450k Array		Targeted BiS-seq	
		Slope	p-value	Slope	p-value	Slope	p-value
tRNA-Gln-CTG-7-1	Chr1:147,800,750-147,801,250	0.84	2.80e-05				
tRNA-Glu-TTC-1-1	Chr2:131,094,500-131,095,000	1.11	4.84e-09				
	Chr2:131,094,250-131,094,750	1.00	1.12e-07				
	Chr2:131,094,750-131,095,250	1.00	3.28e-07				
tRNA-His-GTG-2-3	Chr1:146,541,500-146,545,000	0.92	1.39e-08				
tRNA-His-GTG-2-1	Chr1:149,155,750-149,156,250	1.05	2.98e-08				
	Chr1:149,155,500-149,156,000	0.83	1.37e-05				
tRNA-Ile-AAT-10-1	Chr6: 27,251,500-27,252,000	1.07	1.45e-08			1.30	1.22e-03
	Chr6: 27,251,750-27,252,250	0.90	1.86e-06			1.30	1.22e-03
tRNA-Ile-AAT-4-1	Chr6: 8,130,000-8,130,500	1.19	2.98e-10	19.63	8.92e-06	-0.74	6.88e-04
	Chr6: 8,130,250-8,130,750	0.77	3.898e-05	19.63	8.92e-06	-0.74	6.88e-04
tRNA-Ile-TAT-2-2	Chr6: 26,987,750-26,988,250	0.97	7.25e-07	4.16	1.17e-02	-0.60	3.84e-01
tRNA-iMet-CAT-1-5	Chr6: 26,330,500-26,331,000	1.24	2.83e-11	13.01	6.07e-06	4.54	9.35e-04
	Chr6: 26,330,750-26,331,250	1.13	2.89e-09	13.01	6.07e-06	4.54	9.35e-04
tRNA-Leu-TAG-2-1	Chr14: 21,093,250-21,093,750	1.04	9.36e-07			2.49	8.77e-03
	Chr14: 21,093,500-21,094,000	0.94	8.58e-07			2.49	8.77e-03
tRNA-Pro-AGG-2-2	Chr6: 26,555,500-26,556,000	1.04	3.97e-08				
	Chr6: 26,555,250-26,555,750	0.94	1.89e-05				
tRNA-Ser-ACT-1-1	Chr6: 27,261,250-27,261,750	0.97	3.55e-07			0.66	1.45e-01
tRNA-Ser-AGA-2-6	Chr7: 8,129,750-8,130,250	1.21	1.16e-10	20.87	6.72e-05	0.62	1.28e-02
	Chr6: 8,130,000-8,130,500	1.19	3.03e-10	20.87	6.72e-05	0.62	1.28e-02
tRNA-Ser-TGA-2-1	Chr2: 27,513,000-27,513,500	0.90	3.58e-06	87.21	1.38e-04	0.45	5.74e-01
tRNA-Val-AAC-1-2	Chr5:180,590,750-180,591,250	0.94	3.28e-06				
tRNA-Val-AAC-4-1	Chr6: 27,618,500-27,619,000	1.07	1.25e-08	40.06	9.90e-03		
	Chr6: 27,618,250-27,618,750	0.95	1.31e-07	40.06	9.90e-03		
tRNA-Val-CAC-2-1	Chr6: 27,247,750-27,248,250	0.85	2.33e-05	59.16	5.05e-06		