Identification of GABAergic neurons

innervating the zebrafish lateral habenula

[1,2]Mahathi Ramaswamy, [1]Ruey-Kuang Cheng and [1,3]Suresh Jesuthasan*

[1]Lee Kong Chian School of Medicine, Nanyang Technological University Singapore,

Singapore.

[2]NUS Graduate School for Integrative Sciences and Engineering, National University of

Singapore, Singapore.

[3]Institute of Molecular and Cell Biology, Singapore.

*Author for correspondence (sureshj@ntu.edu.sg)

Abstract

Habenula neurons are constantly active. The level of activity affects mood and behaviour, with increased activity in the lateral habenula reflecting exposure to punishment and a switch to passive coping and depression. Here, we identify GABAergic neurons that could reduce activity in the lateral habenula of larval zebrafish. GAD65/67 immunohistochemistry and imaging of *gad1b:DsRed* transgenic fish suggest the presence of GABAergic terminals in the neuropil and between cell bodies in the lateral habenula. Retrograde tracing with the lipophilic dye DiD suggests that the former derives from the thalamus, while the latter originates from a group of cells in the posterior hypothalamus that are located between the posterior tuberal nucleus and hypothalamic lobes. Two-photon calcium imaging indicates that blue light causes excitation of thalamic GABAergic neurons and terminals in the neuropil, while a subpopulation of lateral habenula neurons show reduced intracellular calcium levels. Whole-cell electrophysiological recording indicates that blue-light reduces membrane potential of lateral habenula neurons. These observations suggest that GABAergic input from the thalamus may mediate inhibition in the zebrafish lateral habenula. Mechanisms governing release of GABA from the neurons in the posterior hypothalamus, which are likely to be in the tuberomammillary nucleus, remain to be defined.

Introduction

The habenula is a regulator of mood and behaviour in vertebrates (Lecourtier & Kelly, 2007; Namboodiri *et al.*, 2016). It consists of two major subdivisions, the medial and lateral habenula, and receives input from several regions of the forebrain (Mok & Mogenson, 1974; Herkenham & Nauta, 1977; Turner *et al.*, 2016). The lateral habenula has become a focus of substantial interest recently, as activity here controls the release of broadly acting neuromodulators such as dopamine and serotonin (Wang & Aghajanian, 1977; Hikosaka, 2010; Proulx *et al.*, 2014).

In addition to transient activity that is evoked by sensory stimuli (Dreosti *et al.*, 2014; Krishnan

et al., 2014; Cheng *et al.*, 2017; Zhang *et al.*, 2017) and punishment (Matsumoto & Hikosaka,

2009), the habenula displays persistent or spontaneous activity (Gao *et al.*, 1996; Jetti *et al.*,

2014; Sakhi *et al.*, 2014; Baño-Otálora & Piggins, 2017). This activity is dependent on synaptic

input (Kim & Chang, 2005), and changes with time of day (Zhao & Rusak, 2005; Sakhi *et al.*,

2014) or as a result of a recent experience. Persistent elevated activity in the lateral habenula

is implicated in depression (Li *et al.*, 2011; Yang *et al.*, 2018; Andalman *et al.*, 2019).

Given these effects, there is a growing interest in understanding how activity in the lateral

habenula is controlled. In many brain regions, excitation is provided by afferent neurons, with

inhibition coming from local GABAergic interneurons. The habenula largely lacks GABAergic

neurons (Smith *et al.*, 1987; Pandey *et al.*, 2018; Zhang *et al.*, 2018), but receives GABAergic

input (Gottesfeld *et al.*, 1979). Although GABAergic input is normally inhibitory, a subset of

habenula neurons show extended firing following transient hyperpolarization (Chang & Kim,

2004). In the lateral habenula of mice, afferent neurons from the entopeduncular nucleus and

the VTA co-release glutamate and GABA (Root *et al.*, 2014; Shabel *et al.*, 2014). A small

population of GABAergic neurons in the lateral preoptic area (LPO) of rodents projects to the

lateral habenula (Barker *et al.*, 2017). Another input source to the habenula is the antero-dorsal

thalamus (Cheng *et al.*, 2017; Zhang *et al.*, 2017; Fernandez *et al.*, 2018), a region that is rich

in GABAergic neurons (Mueller, 2012). In the mouse, GABAergic neurons from the ventral

lateral geniculate nucleus and intergeniculate leaflet project to the lateral habenula, and enable

light-dependent inhibition (Huang *et al.*, 2019). Here we examine the sources of GABAergic

input to the lateral habenula of zebrafish, an experimental system with an easily accessible

habenula due to eversion of the brain during development (Mueller *et al.*, 2011; Folgueira *et*

al., 2012). Our data suggests that there are two sources - the thalamus and the posterior

hypothalamus. This data on connectivity is expected to shed light on the mechanisms by which

brain states are controlled.

Materials and Methods

Zebrafish lines

Transgenic lines used include *SqKR11Et (Lee et al., 2010; Teh et al., 2010)*, *Et(-*

0.6hsp70l:Gal4-VP16)s1020t (Scott & Baier, 2009), *Tg(UAS:Kaede)s1999t* (Scott *et al.*,

2007), *Tg(elavl3:GCaMP6f)a12200* (Wolf *et al.*, 2017), *TgBAC(gad1b:DsRed)nns26* (Satou

et al., 2013), *Tg(vGlut2:GFP)nns43 (Satou et al., 2012)*, *Tg(dao:GAL4VP16)rw0148a* (Amo

et al., 2014), *Tg(UAS:GCaMP6s)sq205* and *Gt(hspGFFDMC76A)* (Muto *et al.*, 2017) . The

GAL4 enhancer trap line is referred to as *s1020t* in the text for brevity.

All experiments were carried out under guidelines approved by the IACUC of Biopolis

(number 191420).

Neural tracing

A saturated solution of DiD (ThermoFisher Scientific) was made by dissolving a small crystal

in 50µL of ethanol. Larval fish were anesthetized with tricaine, mounted in 2% low-melting

agarose and placed under a compound microscope (Zeiss Examiner). A small amount of dye

was pressure injected into either the left or right habenula by visualizing the fish under a water-

immersion 40x objective. The fish was rested for 30 minutes to allow the dye to diffuse and

then imaged using a Zeiss LSM800 confocal microscope, under a 40x water-immersion

objective. Fish with labelled cells outside the habenula were not used.

Antibody label

Larvae were fixed in 4% paraformaldehyde in phosphate buffered saline (PBS) overnight and then rinsed several times with PBS. After peeling the skin off the brains, samples were incubated in 1%BSA and then overnight at 4 degrees in primary antibody (anti-GAD65/67, Abcam ab11070, 1:500 (Cheng *et al.*, 2017)). The following day the brains were washed several times with PBS and then incubated in secondary antibody (Alexa 488 goat anti-rabbit, 1:1000) overnight at 4˚C. After multiple washes with PBS, brains were mounted in low-melting agarose and imaged on a laser scanning confocal microscope (Zeiss LSM800) using a 40x water dipping objective. A total of 7 fish were imaged.

Electrophysiology

Whole-cell patch clamp and loose-patch recordings were performed from lateral habenula neurons in 5-10 days post-fertilization larvae using procedures previously described in Lupton et al. (2017), except that fish in the current study were not pinned onto a Sylgard dish. Fish were anesthetized by mivacurium before being mounted in a 2% low-melting agarose in a glass-bottom dish that was then immersed with external saline (composition in mM: 134 NaCl, 2.9 KCl, 1.2 MgCl2, 10 Hepes, 10 Glucose, 2.1 CaCl2, pH 7.8). In loose-patch recordings, the pipette was loaded with standard Ringer's solution to record neural signals from extracellular space while in whole-cell patch recordings, the pipette was loaded with a potassium gluconate–based internal solution (composition in mM: 115 K gluconate, 15 KCl, 2 MgCl2,10 Hepes, 10 EGTA, 4 MgATP, pH 7.2) to record neural signals intracellularly. The pipettes were pulled with thick-walled borosilicate capillaries (1.5 mm OD; 0.86 mm ID; Warner Instruments) using a Flaming Brown P-1000 pipette puller (Sutter Instruments) to give a tip diameter of 1–1.5 μm with an initial resistance of 10–20 MΩ for loose-patch recording. Neural signals were acquired using Multiclamp 700B amplifier, Digidata 1440A digitizer, and pCLAMP software v.10 (all

113   from Molecular Devices). In whole-cell patch recordings, the resistance reached gigaohm seal

114   before breaking into the cells. The data were low-pass filtered at 2 kHz using a Bessel filter

115   and sampled at 20 kHz at a gain of 1. Spike events were detected offline using Clampfit v.10.7

116   (downloaded from Molecular Devices). Graphs were plotted using Microsoft Excel for whole-

117   cell patch data and in Clampfit for loose-patch data.

Calcium imaging

119   Larval fish were imaged as described previously (Cheng *et al.*, 2017). Imaging was carried out

120   using a Nikon A1RMP two-photon microscope with a 25x water-dipping objective (NA=1.1),

121   with the Ti-Sapphire tuned to 930 nm and galvano scanning. 5 mm blue LEDs (460-490 nm),

122   controlled by the Nikon Elements software, were used to provide 20 second pulses of light. For

123   imaging the thalamus, multiple planes were captured sequentially.

Image analysis

125   Time-series registration, ROI extraction and rastermap calculation were performed using

126   Suite2p (Pachitariu *et al.*, 2017). Data was normalized into z-scores using Matlab.

Results

The zebrafish habenula is innervated by GABAergic neurons

GAD65/67 immunofluorescence indicates that GABAergic terminals are present in the

neuropil of medial (also called dorsal, based on adult morphology (Amo *et al.*, 2010)) and

lateral (or ventral) habenula of larval zebrafish (Figure 1A, B; arrowheads). In addition,

GABAergic neurites were visible between cell soma in the ventro-lateral habenula. The

GAD65/67 label in the habenula was distinct from the uniform label seen in cell bodies (see

white arrow in Figure 1A), and was seen in the absence of habenula neuron label. As an

independent means of visualizing the cell body and neurites of GABAergic neurons, we imaged

gad1b:DsRed, vGlut2:GFP double transgenic fish, which express DsRed and GFP in

GABAergic and glutamatergic neurons respectively. DsRed could be detected in neurites in

the lateral habenula, surrounding GFP-expressing neurons (Figure 1C). No DsRed labelled cell

was detected in the habenula, in contrast to other brain regions that contained fully labelled cell

bodies. This observation, in conjunction with the GAD65/67 antibody labelling, indicates that

the zebrafish habenula receives GABAergic input.

Neuronal tracing identifies sources of GABAergic afferents targeting

To identify the source of the GABAergic inputs, we examined transgenic lines in which

habenula afferents are labelled. Two major sources of input to the zebrafish habenula are the

entopeduncular nucleus (Turner *et al.*, 2016) and the thalamus (Cheng *et al.*, 2017). The

entopeduncular nucleus is located in the ventral telencephalon, in the vicinity of the lateral

forebrain bundle (Turner *et al.*, 2016). Neurons in this location, which are labelled in the

Et(sqKR11) line (Lee *et al.*, 2010) (Supplementary Movie 1), were seen to innervate the

neuropil of lateral habenula, but no neurites could be detected in between cell soma in lateral

habenula (Figure 1D). In contrast, neurites that are labelled by the *s1020tEt* driver

(Supplementary Movie 2), were visible between cells of the lateral habenula (Figure 1E-G).

These neurites co-expressed DsRed under the control of the *gad1b* promoter (Figure 1H),

indicating that they are GABAergic. GABAergic cell bodies were seen in the thalamus in

neurons expressing GCaMP6s under the control of the *s1020tEt* driver, but not in the

entopeduncular nucleus, which had few 1020tEt positive cells (Supplementary Movie 2). Thus,

GABAergic afferents to the lateral habenula may derive from the thalamus.

To confirm the location of the GABAergic afferents, the lipophilic tracer DiD was injected into

the lateral habenula of *Tg(gad1b:DsRed)* fish (Fig 2A; n = 17 fish; see Movie 3 for an example).

In all cases, terminals between soma were visible in the contralateral lateral habenula,

suggesting that there may be bilateral or contralateral origin of these afferents. Retrogradely

labelled neurons were visible in the ipsilateral thalamus in only 12 of these fish (Fig 2B, D-G),

and out of 42 labelled thalamic neurons, 16 were GABAergic, as judged by *gad1b:DsRed*

expression. The fact that 5 fish did not contain any labelled neurons in the thalamus, despite

the presence of terminals in the lateral habenula (Figure 2L; see also Movie 20190422)

indicates that there is an additional or different source of the soma-targeted terminals.

Retrogradely labelled neurons in the entopeduncular nucleus were DsRed-negative (Fig 2C),

indicating that this is not the source.

It has been reported that the posterior hypothalamus provides input to the habenula (Turner *et*

al., 2016). Indeed, injection into the lateral habenula retrogradely labelled scattered cells in the

posterior hypothalamus of all fish where this area was imaged (Figure. 2H-K; n = 7 fish

imaged). Label was bilateral in all cases, and 20 out of 46 neurons were positive for DsRed,

indicating that this nucleus sends GABAergic projections to the lateral habenula. The labelled

neurons were located lateral to the posterior tuberal nucleus, and medial to the inferior lobe of

the hypothalamus (IHL), as judged by tracing experiments in the *Tg(76A:GAL4,*

UAS:GCaMP6s) line (Muto *et al.*, 2017) (Figure 2L-N). Neurons in this region expressed

GCaMP6s under the control of the *s1020t* GAL4 driver as well as DsRed under the control of

the *gad1b* promoter (Figure 2O-Q). Together, these observations suggest that GABAergic

terminals seen between cells in the lateral habenula of the s1020t fish derive from the posterior

hypothalamus, and imply that GABAergic terminals in the neuropils derive from the thalamus.

Blue light triggers activity in thalamic GABAergic neurons and afferents targeting the lateral habenula neuropil

It has been reported previously that cells in the anterior thalamus of zebrafish are reproducibly activated by pulses of blue light (Cheng *et al.*, 2017; Zhang *et al.*, 2017). To test whether the responding thalamic cells include GABAergic neurons, two photon calcium imaging was performed on larvae expressing GCaMP6s under the control of the s1020t GAL4 driver. An increase in GCaMP6s fluorescence was seen in an average of 46 ± 23 thalamic neurons (mean $\pm$ standard deviation; n = 6 fish). Of these, 20 ± 13 cells (mean $\pm$ standard deviation; approximately $43 \pm 18\%$) were GABAergic (Figure 3A-C). To determine whether terminals in the lateral habenula are also affected, imaging was also carried out in this region (n = 3 fish). In response to blue light, a rise in the fluorescence of this reporter was seen in a subset of terminals within the neuropil of the lateral habenula (Figure 3D-F). No response to blue light was seen in terminals between soma, although activity there was correlated to one another (Figure 3G-I'). These observations are consistent with the notion that blue light can influence the lateral habenula via GABAergic afferents from the thalamus that target the neuropils, but not the cell bodies.

Blue light transiently reduces intracellular calcium in a subset of lateral habenula neurons

If thalamic neurons provide GABAergic input to the neuropil of lateral habenula, then stimulation of these neurons may inhibit the lateral habenula. Previous two photon calcium imaging has shown that blue light causes both increase and decrease in ongoing activity throughout the habenula (Cheng *et al.*, 2017; Zhang *et al.*, 2017). In these recordings, however, it was not clear if inhibition also occurs in the lateral habenula. To determine this, we imaged fish expressing GCaMP6s under the control of the *dao* promoter (n = 6 fish). A minority of

cells (30/154) showed a loss of activity during exposure to light (Figure 4A-D). We also performed electrophysiology on intact animals. Both increase and decrease in membrane potential (Figure 4E, F) was detected in lateral habenula neurons by whole-cell recording. When loose-patch recording was employed, the onset of blue light was seen to either trigger or dampen neural spikes (Figure 4G, H). Hence, in addition to excitation, blue light-dependent reduction in the activity of lateral habenula cells is supported by both calcium imaging and electrophysiological recordings.

Discussion

We have examined the source of neurons that can influence neural activity in the lateral habenula of larval zebrafish through the release of GABA. GAD65/67 labelling and imaging of a transgenic line which expresses DsRed under the control of the *gad1b* promoter indicate that GABAergic terminals can be found in the habenula neuropils and also in between cell soma. This suggests that GABAergic neurons target both dendrites and the soma of lateral habenula neurons. Targeting onto the soma was not seen in the medial habenula of larval zebrafish, but only in the lateral habenula. Thus, while GABAergic neurons may influence the effect of excitatory input throughout the habenula by inhibition of dendrites, the lateral habenula could experience stronger inhibition due to signalling onto the soma. Tracing data indicates that GABAergic innervation targeting the neuropil and soma derive from different populations.

A recent study in mice demonstrated that GABAergic neurons from the lateral geniculate nucleus innervate and inhibit the lateral habenula in mice (Huang *et al.*, 2019). Consistent with these findings, we see that retrograde labelling of lateral habenula neurons in the zebrafish reveals *gad1b*-positive afferents from the thalamus. Further, we show that GABAergic

thalamic neurons as well as terminals in the lateral habenula are excited by light, while a

number of lateral habenula cells are inhibited. Electrical recordings of the lateral habenula

indicate that the onset of blue light led to a reduction in membrane potential and in the number

of spikes. The link between illumination levels and mood has been well-studied in animals and

humans (Viola *et al.*, 2008; Vandewalle *et al.*, 2010; Bedrosian *et al.*, 2011; LeGates *et al.*,

2014). Given that increased activity in the lateral habenula has negative effects (Lecca *et al.*,

2014; Proulx *et al.*, 2014; Lawson *et al.*, 2017; Howe & Kenny, 2018), inhibition of these

neurons by thalamic GABAergic neurons may contribute to the positive effects of light (Huang

et al., 2019). However, it should be noted that only a subset of neurons were inhibited by light,

indicating that the effect of blue light on inhibition of the lateral habenula, at least in the

conditions used here, are relatively modest.

Retrograde tracing experiments also revealed *gad1b*-positive inputs from a cluster of cells

lateral to the posterior tuberal nucleus and medial to the inferior lobe of the hypothalamus.

Projections from the posterior tuberculum (PT) to the habenula have been previously reported

(Hendricks & Jesuthasan, 2007; Turner *et al.*, 2016). However, neurons in the posterior

tuberculum have been characterized as largely non-GABAergic (Filippi *et al.*, 2014; Heap *et*

al., 2017). Similar scattered afferents have been identified by retrograde label in the habenula

of adult zebrafish (Turner *et al.*, 2016), where these were suggested to be located in the

posterior hypothalamus, and in the trout (Yañez & Anadón, 1996), where the cells were named

the "tuberculohabenular nucleus". We propose that these cells are located in the

tuberomammillary nucleus, which is located in the posterior hypothalamus (Chen *et al.*, 2017).

This would be consistent with the existence of histaminergic fibers in the lateral habenula

(Kaslin & Panula, 2001), the fact that neurons in the tuberomammillary nucleus in the posterior

hypothalamus are the sole source histamine in vertebrates, and that histaminergic neurons co-

release GABA (Sundvik & Panula, 2012; Yu *et al.*, 2015). Future experiments to define the

function of these cells should yield fresh insight into the mechanisms by which activity in the

lateral habenula is inhibited.

258 Acknowledgements

259   This research was supported by a Lee Kong Chian School of Medicine, Nanyang Technological

260   University Singapore Start-Up Grant, and the Singapore Ministry of Education under its

261   Academic Research Fund Tier 2 (MOE2017-T2-058) and Tier 1 (MOE2016-T1-001-152)

262   awards. We thank Vatsala Thirumalai and members of her lab for instruction on techniques in

263   zebrafish electrophysiology.

264 Conflict of Interest Statement

265   The authors declare no conflict of interest.

266 Author contributions

267   MR performed antibody label, retrograde tracing and microscopy. RKC performed

268   electrophysiology and functional imaging. SJ imaged transgenic lines, analyzed data and wrote

269   the manuscript.

270 Data Accessibility

271   All confocal stacks used here are available on Figshare.

     Figure captions

Figure 1. Detection of GABAergic terminals in the habenula. (A) A 6 day old fish, labelled

with an antibody to GAD65/67. Terminals are visible in the lateral habenula (yellow

arrowheads). The red arrows indicate GABAergic neurons in the pallium. (B) A 2 week old

fish, showing GAD65/67-labelled puncta in the neuropil of the medial habenula (white arrow),

and in between cells in the lateral habenula (yellow arrowhead). No habenula cell bodies are

labelled. This sample is slightly tilted. (C) A 7 day old *Tg(vGlut2:GFP), Tg(gad1b:DsRed)*

double transgenic fish. DsRed (magenta) is visible in terminals in the lateral habenula

(arrowheads), while habenula cell bodies express GFP (green). (D) An 8 day old fish *SqKR11Et*

fish, with afferents from the entopeduncular nucleus labelled in red. Terminals are restricted to

neuropil of the lateral habenula (white arrow), and do not overlap with GAD65/67 label

between cells (yellow arrowhead). (E) Afferents from thalamic neurons, labelled with Kaede

under the control of the *s1020tEt* driver. Terminals are visible in the neuropil (arrow) and also

between cells (arrowhead). (F-H) The left habenula of a 6 day old fish, with terminals labelled

by GCaMP6s, which is expressed under the control of the *s1020tEt* driver (G), and DsRed

under the control of the *gad1b* promoter (H). The white arrowhead in F indicates

autofluorescing pterinosomes in xanthophores (Le Guyader & Jesuthasan, 2002). Scale bar =

25 µm for A-E, 10 µm for F-H . All images are dorsal view, with anterior to the left.

Figure 2. Retrograde label of lateral habenula afferents. (A-C) An example of DiD

injection (cyan) into the left lateral habenula (A, red arrowhead) of a 7 day old

Tg(elavl3:GCaMP6f), Tg(gad1b:DsRed) fish. (B) Retrogradely labelled cell bodies

(arrowheads) in the thalamic region. (C) A deeper focal plane. Entopeduncular nucleus (EN)

cell bodies are also labelled by DiD but are *gad1b*-negative. (D-G) The boxed area in panel B

shown at higher magnification. The arrowheads indicate neurons in the thalamus expressing

GCaMP6f (E) and DsRed (F), which have been retrogradely labelled by DiD (H). (H-K) DiD

injection (cyan) into the right lateral habenula of another double transgenic 7 day old fish (H,

red arrowhead). Lateral habenula neurons express GCaMP6s under the *dao* promoter. (I)

Bilateral label of cell bodies in the posterior hypothalamus. (J, K) The boxed region in panel I

at higher magnification, showing retrogradely labelled neurons that are *gad1b* positive (yellow

arrowhead) or negative (green arrowhead). (L-N) DiD injection into the left lateral habenula

of a 7 day old *Tg(76A:GAL4, UAS:GCaMP6s, gad1b:DsRed)* fish. (L) The injection site (red

arrowhead). (M) Retrogradely labelled cells (yellow arrowheads), at the level of the inferior

lobe of the hypothalamus. (N) GCaMP6s expression in the inferior lobe of the hypothalamus.

(O-Q) Expression of GCaMP6s and DsRed in a neuron (yellow arrowhead) in the posterior

hypothalamus of 10 day old *s1020t:GAL4, UAS:GCaMP6s, gad1b:DsRed* fish. All panels are

dorsal views, with anterior to the left. Pa: pallium, OT: optic tectum, fr: fasciculus retroflexus,

rHb: right habenula, PT: posterior tuberculum, IPN: interpeduncular nucleus.

Figure 3. Effect of blue light on thalamic GABAergic neurons. (A-C) The thalamic response

of a 6 day old fish to 3 pulses of blue light. (A) Confocal image showing thalamic neurons,

some of which express DsRed under the control of the gad1b promoter (magenta). (B) Change

in GCaMP6f fluorescence of thalamic neurons, correlated to change in the AF4 neuropil (red

arrow), which receives retinal input. Cells indicated by arrowheads have a high correlation with

activity in this neuropil and express DsRed, as shown in panel A. (C) Rastermap of cells and

neuropils extracted from the time-series. Two major responses are phasic and tonic response

to light. (D - F) Calcium imaging of terminals in the neuropil of the ventral habenula of a 5 day

old fish exposed to 1 pulse of blue light. (D) Average projection of the time series, showing

neuropils (yellow arrowheads). (E) Correlation map of response, with red pixels showing

terminals with a highly correlated calcium increase in response to a pulse of blue light. (F)

Rastermap showing fluorescence change with time. (G - I) Response in terminals between cell

bodies. (G) Average of the time-series, showing the location of terminals (yellow arrowheads).

(H) Map of response, correlated to terminals in the thalamic neuropil AF4 (white arrowhead).

(H') Rastermap of 5 ROIs that are highly correlated with blue light pulses. (I) Map of response

correlated to terminals in the lateral habenula (white arrowhead). (I') Rastermap of 6 ROIs in

the lateral habenula that are highly correlated to one another. These have no relationship to the

light pulses. All panels are dorsal views, with anterior to the top. Th, thalamus. The wedge in

panel B indicates the lookup table used for correlation maps.

Figure 4. Calcium imaging and electrophysiological recordings in the lateral habenula.

(A-D Response of neurons in the lateral habenula of a 5 day old fish exposed to four pulses of

blue light. (A) Average of a time-series, showing distribution of cells labelled in a *dao:GAL4,*

UAS:GCaMP6s fish. (B) Correlation map of response to 4 pulses of light, with red showing a

highly correlated cell, and blue showing an anti-correlated cell. (C) z-scores of the two cells

indicated by the blue and red arrowheads in panel B. (D) Rastermap of the response of all cells.

Blue bars the presence of blue light. (E, F) Whole-cell recordings showing membrane

depolarization (E) or hyperpolarization (F) by blue light. The blue light was 13 seconds in

duration in both cases and the range of voltage change (peak-to-trough) was on average 2-3

mV in both cases. Solid horizontal lines represent the means of the membrane potentials before,

during and after the blue light in accordance with the trace colors. Each trace in panel E is a

separate cell. (G, H) Loose-patch recordings. (G) There are spikes of two different heights,

implying signals from two cells. Both show an increase in spikes in the presence of blue light.

(H) A recording in another fish, showing a reduction in spiking during blue light delivery. The

ticks in the upper trace in (G) and (H) were taken from spike detection from the main trace, as

indicated by the blue squares. Scale bar = 25 μm.

Supplementary Material

Supplementary Movie 1. Z-stack of a 5 day old SqKR11Et, elavl3:GCaMP6f fish. The

yellow arrowheads indicate the lateral forebrain bundle in the ventral telencephalon. The

white arrowhead indicates cells in the entopeduncular nucleus, which project to neuropils in

the habenula.

Supplementary Movie 2. Z-stack of a 6 day old Tg(s1020tGAL4, UAS:GCaMP6s,

gad1b:DsRed) fish. The yellow arrowheads indicate the lateral forebrain bundle in the ventral

telencephalon. The white arrowhead indicates the entopeduncular nucleus, which contains no

DsRed expressing cells. Cells in the thalamus are strongly labelled by the s1020t driver.

Supplementary Movie 3. Retrograde label of cells in the posterior hypothalamus. Z-stack

of a Tg(dao:GAL4, UAS:GCaMP6s, gad1b:DsRed) fish, after DiD label of the left lateral

habenula. Labelled cells are visible in the posterior hypothalamus. Habenula projections to the

raphe are also visible. There are no projections to the IPN. Note that GCaMP6s is expressed in

the posterior hypothalamus.

References

Amo, R., Aizawa, H., Takahoko, M., Kobayashi, M., Takahashi, R., Aoki, T., & Okamoto, H. (2010)

Identification of the zebrafish ventral habenula as a homolog of the mammalian lateral habenula.

J. Neurosci., **30**, 1566–1574.

Amo, R., Fredes, F., Kinoshita, M., Aoki, R., Aizawa, H., Agetsuma, M., Aoki, T., Shiraki, T.,

Kakinuma, H., Matsuda, M., Yamazaki, M., Takahoko, M., Tsuboi, T., Higashijima, S.-I.,

Miyasaka, N., Koide, T., Yabuki, Y., Yoshihara, Y., Fukai, T., & Okamoto, H. (2014) The

habenulo-raphe serotonergic circuit encodes an aversive expectation value essential for adaptive

active avoidance of danger. *Neuron*, **84**, 1034–1048.

Andalman, A.S., Burns, V.M., Lovett-Barron, M., Broxton, M., Poole, B., Yang, S.J., Grosenick, L., Lerner, T.N., Chen, R., Benster, T., Mourrain, P., Levoy, M., Rajan, K., & Deisseroth, K. (2019) Neuronal Dynamics Regulating Brain and Behavioral State Transitions. *Cell*, **177**, 970–985.e20.

Baño-Otálora, B. & Piggins, H.D. (2017) Contributions of the lateral habenula to circadian timekeeping. *Pharmacol. Biochem. Behav.*, **162**, 46–54.

Barker, D.J., Miranda-Barrientos, J., Zhang, S., Root, D.H., Wang, H.-L., Liu, B., Calipari, E.S., & Morales, M. (2017) Lateral Preoptic Control of the Lateral Habenula through Convergent Glutamate and GABA Transmission. *Cell Rep.*, **21**, 1757–1769.

Bedrosian, T.A., Fonken, L.K., Walton, J.C., Haim, A., & Nelson, R.J. (2011) Dim light at night provokes depression-like behaviors and reduces CA1 dendritic spine density in female hamsters. *Psychoneuroendocrinology*, **36**, 1062–1069.

Chang, S.-Y. & Kim, U. (2004) Ionic mechanism of long-lasting discharges of action potentials triggered by membrane hyperpolarization in the medial lateral habenula. *J. Neurosci.*, **24**, 2172–2181.

Cheng, R.-K., Krishnan, S., Lin, Q., Kibat, C., & Jesuthasan, S. (2017) Characterization of a thalamic nucleus mediating habenula responses to changes in ambient illumination. *BMC Biol.*, **15**, 104.

Chen, S., Reichert, S., Singh, C., Oikonomou, G., Rihel, J., & Prober, D.A. (2017) Light-Dependent Regulation of Sleep and Wake States by Prokineticin 2 in Zebrafish. *Neuron*, **95**, 153–168.e6.

Dreosti, E., Vendrell Llopis, N., Carl, M., Yaksi, E., & Wilson, S.W. (2014) Left-right asymmetry is required for the habenulae to respond to both visual and olfactory stimuli. *Curr. Biol.*, **24**, 440–445.

Fernandez, D.C., Fogerson, P.M., Lazzerini Ospri, L., Thomsen, M.B., Layne, R.M., Severin, D., Zhan, J., Singer, J.H., Kirkwood, A., Zhao, H., Berson, D.M., & Hattar, S. (2018) Light Affects Mood and Learning through Distinct Retina-Brain Pathways. *Cell*, **175**, 71–84.e18.

Filippi, A., Mueller, T., & Driever, W. (2014) vglut2 and gad expression reveal distinct patterns of dual GABAergic versus glutamatergic cotransmitter phenotypes of dopaminergic and noradrenergic neurons in the zebrafish brain. *J. Comp. Neurol.*, **522**, 2019–2037.

Folgueira, M., Bayley, P., Navratilova, P., Becker, T.S., Wilson, S.W., & Clarke, J.D.W. (2012)

Morphogenesis underlying the development of the everted teleost telencephalon. *Neural Dev.*, **7**,

32.

Freeman, J., Vladimirov, N., Kawashima, T., Mu, Y., Sofroniew, N.J., Bennett, D.V., Rosen, J.,

Yang, C.-T., Looger, L.L., & Ahrens, M.B. (2014) Mapping brain activity at scale with cluster

computing. *Nat. Methods*, **11**, 941–950.

Gao, D.M., Hoffman, D., & Benabid, A.L. (1996) Simultaneous Recording of Spontaneous Activities

and Nociceptive Responses from Neurons in the Pars Compacta of Substantia Nigra and in the

Lateral Habenula. *Eur. J. Neurosci.*, **8**, 1474–1478.

Gottesfeld, Z., Brandon, C., Jacobowitz, D.M., & Wu, J.-Y. (1979) The gaba system in the

mammalian habenula. *Brain Res. Bull.*, **4**, 693.

Heap, L.A., Vanwalleghem, G.C., Thompson, A.W., Favre-Bulle, I., Rubinsztein-Dunlop, H., &

Scott, E.K. (2017) Hypothalamic Projections to the Optic Tectum in Larval Zebrafish. *Front.*

Neuroanat., **11**, 135.

Hendricks, M. & Jesuthasan, S. (2007) Asymmetric innervation of the habenula in zebrafish. *J. Comp.*

Neurol., **502**, 611–619.

Herkenham, M. & Nauta, W.J. (1977) Afferent connections of the habenular nuclei in the rat. A

horseradish peroxidase study, with a note on the fiber-of-passage problem. *J. Comp. Neurol.*,

173, 123–146.

Hikosaka, O. (2010) The habenula: from stress evasion to value-based decision-making. *Nat. Rev.*

Neurosci., **11**, 503–513.

Howe, W.M. & Kenny, P.J. (2018) Burst firing sets the stage for depression. *Nature*,.

Huang, L., Xi, Y., Peng, Y., Yang, Y., Huang, X., Fu, Y., Tao, Q., Xiao, J., Yuan, T., An, K., Zhao,

H., Pu, M., Xu, F., Xue, T., Luo, M., So, K.-F., & Ren, C. (2019) A Visual Circuit Related to

Habenula Underlies the Antidepressive Effects of Light Therapy. *Neuron*, **102**, 128–142.e8.

Jetti, S.K., Vendrell-Llopis, N., & Yaksi, E. (2014) Spontaneous activity governs olfactory

representations in spatially organized habenular microcircuits. *Curr. Biol.*, **24**, 434–439.

Kaslin, J. & Panula, P. (2001) Comparative anatomy of the histaminergic and other aminergic systems

in zebrafish (Danio rerio). *J. Comp. Neurol.*, **440**, 342–377.

Kim, U. & Chang, S.-Y. (2005) Dendritic morphology, local circuitry, and intrinsic electrophysiology

of neurons in the rat medial and lateral habenular nuclei of the epithalamus. *J. Comp. Neurol.*,

483, 236–250.

Krishnan, S., Mathuru, A.S., Kibat, C., Rahman, M., Lupton, C.E., Stewart, J., Claridge-Chang, A.,

Yen, S.-C., & Jesuthasan, S. (2014) The right dorsal habenula limits attraction to an odor in

zebrafish. *Curr. Biol.*, **24**, 1167–1175.

Lawson, R.P., Nord, C.L., Seymour, B., Thomas, D.L., Dayan, P., Pilling, S., & Roiser, J.P. (2017)

Disrupted habenula function in major depression. *Mol. Psychiatry*, **22**, 202–208.

Lecca, S., Meye, F.J., & Mameli, M. (2014) The lateral habenula in addiction and depression: an

anatomical, synaptic and behavioral overview. *Eur. J. Neurosci.*, **39**, 1170–1178.

Lecourtier, L. & Kelly, P.H. (2007) A conductor hidden in the orchestra? Role of the habenular

complex in monoamine transmission and cognition. *Neurosci. Biobehav. Rev.*, **31**, 658–672.

Lee, A., Mathuru, A.S., Teh, C., Kibat, C., Korzh, V., Penney, T.B., & Jesuthasan, S. (2010) The

habenula prevents helpless behavior in larval zebrafish. *Curr. Biol.*, **20**, 2211–2216.

LeGates, T.A., Fernandez, D.C., & Hattar, S. (2014) Light as a central modulator of circadian

rhythms, sleep and affect. *Nat. Rev. Neurosci.*, **15**, 443–454.

Le Guyader, S. & Jesuthasan, S. (2002) Analysis of xanthophore and pterinosome biogenesis in

zebrafish using methylene blue and pteridine autofluorescence. *Pigment Cell Res.*, **15**, 27–31.

Li, B., Piriz, J., Mirrione, M., Chung, C., Proulx, C.D., Schulz, D., Henn, F., & Malinow, R. (2011)

Synaptic potentiation onto habenula neurons in the learned helplessness model of depression.

Nature, **470**, 535–539.

Matsumoto, M. & Hikosaka, O. (2009) Representation of negative motivational value in the primate

lateral habenula. *Nat. Neurosci.*, **12**, 77–84.

Mok, A.C. & Mogenson, G.J. (1974) Effects of electrical stimulation of the lateral hypothalamus,

hippocampus, amygdala and olfactory bulb on unit activity of the lateral habenular nucleus in the

rat. *Brain Res.*, **77**, 417–429.

Mueller, T. (2012) What is the Thalamus in Zebrafish? *Front. Neurosci.*, **6**, 64.

Mueller, T., Dong, Z., Berberoglu, M.A., & Guo, S. (2011) The dorsal pallium in zebrafish, Danio

rerio (Cyprinidae, Teleostei). *Brain Res.*, **1381**, 95–105.

Mueller, T., Vernier, P., & Wullimann, M.F. (2006) A phylotypic stage in vertebrate brain

development: GABA cell patterns in zebrafish compared with mouse. *J. Comp. Neurol.*, **494**,

620–634.

Muto, A., Lal, P., Ailani, D., Abe, G., Itoh, M., & Kawakami, K. (2017) Activation of the

hypothalamic feeding centre upon visual prey detection. *Nat. Commun.*, **8**, 15029.

Namboodiri, V.M.K., Rodriguez-Romaguera, J., & Stuber, G.D. (2016) The habenula. *Curr. Biol.*, **26**,

R873–R877.

Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Federico Rossi, L., Dalgleish, H., Carandini,

M., & Harris, K.D. (2017) Suite2p: beyond 10,000 neurons with standard two-photon

microscopy. *BioRxiv*,.

Pandey, S., Shekhar, K., Regev, A., & Schier, A.F. (2018) Comprehensive Identification and Spatial

Mapping of Habenular Neuronal Types Using Single-Cell RNA-Seq. *Curr. Biol.*, **28**, 1052–

1065.e7.

Proulx, C.D., Hikosaka, O., & Malinow, R. (2014) Reward processing by the lateral habenula in

normal and depressive behaviors. *Nat. Neurosci.*, **17**, 1146–1152.

Root, D.H., Mejias-Aponte, C.A., Zhang, S., Wang, H.-L., Hoffman, A.F., Lupica, C.R., & Morales,

M. (2014) Single rodent mesohabenular axons release glutamate and GABA. *Nat. Neurosci.*, **17**,

1543–1551.

Sakhi, K., Wegner, S., Belle, M.D.C., Howarth, M., Delagrange, P., Brown, T.M., & Piggins, H.D.

(2014) Intrinsic and extrinsic cues regulate the daily profile of mouse lateral habenula neuronal

activity. *J. Physiol.*, **592**, 5025–5045.

Satou, C., Kimura, Y., & Higashijima, S.-I. (2012) Generation of multiple classes of V0 neurons in

zebrafish spinal cord: progenitor heterogeneity and temporal control of neuronal diversity. *J.

Neurosci.*, **32**, 1771–1783.

Satou, C., Kimura, Y., Hirata, H., Suster, M.L., Kawakami, K., & -i. Higashijima, S. (2013)

Transgenic tools to characterize neuronal properties of discrete populations of zebrafish neurons.

Development,.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,

Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,

Tomancak, P., & Cardona, A. (2012) Fiji: an open-source platform for biological-image analysis.

Nat. Methods, **9**, 676–682.

Scott, E.K. & Baier, H. (2009) The cellular architecture of the larval zebrafish tectum, as revealed by

gal4 enhancer trap lines. *Front. Neural Circuits*, **3**, 13.

Scott, E.K., Mason, L., Arrenberg, A.B., Ziv, L., Gosse, N.J., Xiao, T., Chi, N.C., Asakawa, K.,

Kawakami, K., & Baier, H. (2007) Targeting neural circuitry in zebrafish using GAL4 enhancer

trapping. *Nature Methods*,.

Shabel, S.J., Proulx, C.D., Piriz, J., & Malinow, R. (2014) Mood regulation. GABA/glutamate co-

release controls habenula output and is modified by antidepressant treatment. *Science*, **345**,

1494–1498.

Smith, Y., Séguéla, P., & Parent, A. (1987) Distribution of GABA-immunoreactive neurons in the

thalamus of the squirrel monkey (Saimiri sciureus). *Neuroscience*, **22**, 579–591.

Sundvik, M. & Panula, P. (2012) Organization of the histaminergic system in adult zebrafish (Danio

rerio) brain: neuron number, location, and cotransmitters. *J. Comp. Neurol.*, **520**, 3827–3845.

Teh, C., Chudakov, D.M., Poon, K.-L., Mamedov, I.Z., Sek, J.-Y., Shidlovsky, K., Lukyanov, S., &

Korzh, V. (2010) Optogenetic in vivo cell manipulation in KillerRed-expressing zebrafish

transgenics. *BMC Dev. Biol.*, **10**, 110.

Thévenaz, P., Ruttimann, U.E., & Unser, M. (1998) A pyramid approach to subpixel registration

based on intensity. *IEEE Trans. Image Process.*, **7**, 27–41.

Turner, K.J., Hawkins, T.A., Yáñez, J., Anadón, R., Wilson, S.W., & Folgueira, M. (2016) Afferent

Connectivity of the Zebrafish Habenulae. *Front. Neural Circuits*, **10**, 30.

Vandewalle, G., Schwartz, S., Grandjean, D., Wuillaume, C., Balteau, E., Degueldre, C., Schabus, M.,

Phillips, C., Luxen, A., Dijk, D.J., & Maquet, P. (2010) Spectral quality of light modulates

emotional brain responses in humans. *Proc. Natl. Acad. Sci. U. S. A.*, **107**, 19549–19554.

Viola, A.U., James, L.M., Schlangen, L.J.M., & Dijk, D.-J. (2008) Blue-enriched white light in the

workplace improves self-reported alertness, performance and sleep quality. *Scand. J. Work*

Environ. Health, **34**, 297–306.

Wang, R.Y. & Aghajanian, G.K. (1977) Physiological evidence for habenula as major link between

forebrain and midbrain raphe. *Science*, **197**, 89–91.

Wolf, S., Dubreuil, A.M., Bertoni, T., Böhm, U.L., Bormuth, V., Candelier, R., Karpenko, S.,

Hildebrand, D.G.C., Bianco, I.H., Monasson, R., & Debrégeas, G. (2017) Sensorimotor

computation underlying phototaxis in zebrafish. *Nature Communications*,.

Yañez, J. & Anadón, R. (1996) Afferent and efferent connections of the habenula in the rainbow trout

(Oncorhynchus mykiss): an indocarbocyanine dye (DiI) study. *J. Comp. Neurol.*, **372**, 529–543.

Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S., & Hu, H. (2018) Ketamine blocks bursting in

the lateral habenula to rapidly relieve depression. *Nature*, **554**, 317–322.

Yu, X., Ye, Z., Houston, C.M., Zecharia, A.Y., Ma, Y., Zhang, Z., Uygun, D.S., Parker, S., Vyssotski,

A.L., Yustos, R., Franks, N.P., Brickley, S.G., & Wisden, W. (2015) Wakefulness Is Governed

by GABA and Histamine Cotransmission. *Neuron*, **87**, 164–178.

Zhang, B.-B., Yao, Y.-Y., Zhang, H.-F., Kawakami, K., & Du, J.-L. (2017) Left Habenula Mediates

Light-Preference Behavior in Zebrafish via an Asymmetrical Visual Pathway. *Neuron*, **93**, 914–

928.e4.

Zhang, L., Hernández, V.S., Swinny, J.D., Verma, A.K., Giesecke, T., Emery, A.C., Mutig, K.,

Garcia-Segura, L.M., & Eiden, L.E. (2018) A GABAergic cell type in the lateral habenula links

hypothalamic homeostatic and midbrain motivation circuits with sex steroid signaling. *Transl.*

Psychiatry, **8**, 50.

Zhao, H. & Rusak, B. (2005) Circadian firing-rate rhythms and light responses of rat habenular

nucleus neurons in vivo and in vitro. *Neuroscience*, **132**, 519–528.

Figure 1

Figure 2

Figure 3

Figure 4