The compact genome of *Giardia muris* reveals important steps in the evolution of intestinal protozoan parasites

Feifei Xu[1], Alejandro Jiménez-González[1], Elin Einarsson[1,#a], Ásgeir Ástvaldsson[1]

Dimitra Peirasmaki[1], Lars Eckmann[2], Jan O. Andersson[1], Staffan G. Svärd[1], Jon

Jerlström-Hultqvist[1*]

[1]Department of Cell and Molecular Biology, BMC, Box 596, Uppsala University,

SE-751 24 Uppsala, Sweden

[2]Department of Medicine, University of California, San Diego, La Jolla, California,

USA

[#a]Current address: Department of Biochemistry, University of Cambridge, Cambridge,

England

17 * Corresponding author: Dr. Jon Jerlström-Hultqvist

18 E-mail: jon.jerlstrom.hultqvist@icm.uu.se

20 **Running title:** Streamlined *Giardia* genome

21 **Key words:** parasite, diplomonad, *Giardia*, streamlined, intestinal colonization,

22 evolutionary biology, horizontal gene transfer, antigenic variation

Abstract

Diplomonad parasites of the genus *Giardia* have adapted to colonizing different hosts,
most notably the intestinal tract of mammals. The human-pathogenic *Giardia* species,
Giardia intestinalis, has been extensively studied at the genome and gene expression
level, but no such information is available for other *Giardia* species. Comparative data
would be particularly valuable for *Giardia muris*, which colonizes mice and is commonly
used as a prototypic *in vivo* model for investigating host responses to intestinal parasitic
infection. Here we report the draft-genome of *G. muris*. We discovered a highly
streamlined genome, amongst the most densely encoded ever described for a nuclear
eukaryotic genome. *G. muris* and *G. intestinalis* share many known or predicted virulence
factors, including cysteine proteases and a large repertoire of cysteine-rich surface
proteins involved in antigenic variation. Different to *G. intestinalis*, *G. muris* maintains
tandem arrays of pseudogenized surface antigens at the telomeres, whereas intact surface
antigens are present centrally in the chromosomes. The two classes of surface antigens
engage in genetic exchange. Reconstruction of metabolic pathways from the *G. muris*
genome suggest significant metabolic differences to *G. intestinalis.* Additionally, *G.
muris* encodes proteins that might be used to modulate the prokaryotic microbiota. The
responsible genes have been introduced in the *Giardia* genus via lateral gene transfer
from prokaryotic sources. Our findings point to important evolutionary steps in the
Giardia genus as it adapted to different hosts and it provides a powerful foundation for
mechanistic exploration of host-pathogen interaction in the *G. muris* – mouse
pathosystem.

Author summary

The *Giardia* genus comprises eukaryotic single-celled parasites that infect many
animals. The *Giardia intestinalis* species complex, which can colonize and cause

diarrheal disease in humans and different animal hosts has been extensively explored

at the genomic and cell biologic levels. Other *Giardia* species, such as the mouse

parasite *Giardia muris*, have remained uncharacterized at the genomic level,

hampering our understanding of *in vivo* host-pathogen interactions and the impact of

host dependence on the evolution of the *Giardia* genus. We discovered that the *G.*

muris genome encodes many of the same virulence factors as *G. intestinalis*. The *G.*

muris genome has undergone genome contraction, potentially in response to a more

defined infective niche in the murine host. We describe differences in metabolic and

microbiome modulatory gene repertoire, mediated mainly by lateral gene transfer, that

could be important for understanding infective success across the *Giardia* genus. Our

findings provide new insights for the use of *G. muris* as a powerful model for

exploring host-pathogen interactions in giardiasis.

Background

Many eukaryotes have evolved from free-living to parasitic lifestyles over evolutionary time, yet parasitism has developed independently in different taxonomic groups and is therefore characterized by many unique features[1,2]. Comparative genomics provides an opportunity to investigate the factors of parasitism such as loss of morphological, metabolic and genomic complexity, and consequently reduced evolutionary potential for a free-living lifestyle[2]. It can also identify the drivers and consequences of a parasitic lifestyle and generate new testable ecological and evolutionary hypotheses[3].

Giardia is a protozoan parasite that non-invasively colonizes the intestinal tract of many vertebrates. The human pathogen, *Giardia intestinalis,* is estimated to cause 300 million cases of giardiasis in the world each year, being a major cause of diarrheal disease[4]. Giardiasis is also a problem in domestic animals, and the zoonotic potential of *Giardia* has been highlighted in recent years[5]. *In vitro* models of the interaction of *G. intestinalis* with human cells have helped to unravel clues to how *Giardia* causes disease[6–8], such as the importance of, the adhesive disc for attachment[9], flagella for motility[10,11], secreted cysteine proteases for interference with host defenses[12–16], interactions with the intestinal microbiota[6,17], differentiation into cysts for transmission[4,18] and interference with nitric oxide (NO) production[19,20]. Despite this progress it remains uncertain whether these *in vitro* models are representative of the natural infection, particularly because animal models of *G. intestinalis* infection have significant limitations. For example, infection of mice, the most commonly used laboratory animals, with human *G. intestinalis* isolates is unreliable and requires manipulations such as antibiotic conditioning[6]. *Giardia muris*, one of six recognized species of *Giardia*[21], has been used as a

mouse model since the 1960s for exploring the pathogenesis and immunological

responses of the mammalian host to infection(22). The availability of knock-out mice

and other host-related resources makes *G. muris* a powerful model to investigate host-

pathogen interactions(6). The life cycle and infective process of *G. muris* is closely

related to infection by *G. intestinalis*(23). Major findings in *Giardia* biology such as

flagellar and disc function, cellular differentiation(22,24,25), and immunity(23,26–29)

have been pioneered with *G. muris*, and later been shown to be transferable to human

G. intestinalis infections(29,30). Unfortunately, research on *G. muris* has been

hampered by the lack of genome information and gene expression data(5).

Here we describe the draft genome of *G. muris*, representing the first genome

of any *Giardia* outside of the *G. intestinalis* species complex. We performed

comparative genomics with free-living (*Kipferlia bialata*(31)) and parasitic (*G.*

intestinalis(32–34) and *S. salmonicida*(35)) relatives to *G. muris* to determine how *G.*

muris may have evolved into an intestinal pathogen of rodents.

Results

Genome assembly

We extracted DNA from freshly excysted *G. muris* cysts purified from the feces of

infected mice (Fig. S1A) and assembled a high-quality draft genome using sequences

obtained by PacBio and Illumina technologies. In addition, we generated RNA-Seq

data for gene prediction and gene expression analyses with total RNA extracted from

cysts, recently excysted cells (excyzoites), and trophozoites isolated from the small

intestine of infected mice (Fig. S1A).

The *G. muris* draft genome consists of 59 contigs spanning 9.8 Mbp, which is

notably smaller than the *G. intestinalis* WB genome (12.6 Mbp, Table 1). Most of the

genome (9.0 Mbp, 92%) is found on five contigs (>1 Mbp). Of the remaining short

contigs (<30 kbp), ten are terminated in telomeric repeats (TAGGG), suggesting they

are the terminal points of five chromosomes. The karyotype of *G. muris* was

previously shown to consist of four separable chromosomes(36). We hypothesize that

our five major contigs represent a total of five chromosomes in *G. muris,* two of

which are so close in size (1.290 and 1.297 Mbp) that they were not readily resolved

using pulsed-field gels, and are named accordingly from 1 to 5 from largest to

smallest in size (Fig. 1). 42 of the 44 small contigs contain ribosomal DNA (rDNA)

clusters that encode 28S, 18S, and 5.8S rRNAs. In fact, rDNA clusters make up 2.0%

of the total genome, and account for 91.6% of the identified repeats (Methods S1).

Half of the contigs terminated by telomeric repeats have adjacent rDNA clusters (Fig.

S1B), suggesting that multiple of the *G. muris* chromosomes(36), like those in *G.*

intestinalis(37), have long repeats of rDNAs close to the telomeres. In contrast to *G.*

intestinalis chromosomes(37), no retrotransposon sequences were found in the

telomeric regions and overall very few retrotransposon sequences were detected in the

G. muris genome.

Allelic sequence heterozygosity (ASH) in the assembly was estimated to be

0.016% (Table 1), equivalent to the low level found in the *G. intestinalis* WB genome

(0.026%, Table 1). Distribution of ASH along chromosomes showed only weak

clustering in certain areas, particularly at the ends of chromosomes (Fig. 1).

Table 1 - Comparison of genome content between *G. muris, G. intestinalis, S.*

salmonicida and K. bialata

Species	*G. muris*	*G. intestinalis*	*S. salmonicida*	*K. bialata*

Genome size (Mbp)	9.8	12.6	12.9	51.0
Chromosomes (scaffolds)	5 (59)	5 (35)	9 (233)	ND (11,564)
G+C %	54.7	46.3	33.4	49.4
Number of protein encoding genes	4653	4963	8067	17,389
Mean / Median protein size (aa)	578 / 428	635 / 457	373	333
*Mean / Median intergenic size (bp)	264 / 37	470 / 81	421	597
§Coding density %	84.5 / 88.6	81.5 / 84.7	72.1	ND
Number of introns	3 *cis*,5 *trans*	8 *cis*,5 *trans*	3 *cis*	124,912
tRNA genes	68	65	145	ND
ASH %	0.016	0.028	0.15	ND
References	This study	Ref(32)	Ref(38)	Ref(31)

* Mean/Median intergenic distance is based on all RNAs (mRNAs, tRNAs, rRNAs),

but not pseudogenized genes.

§ Coding density: First value is based on all RNAs (mRNAs, tRNAs, rRNAs), but not

pseudogenized genes; Second value is based on all RNAs including ψVSP.

Table 2 - Lateral gene transfers in *G. muris (Gm)*, *G. intestinalis (Gi* WB, *Gi* GS, *Gi*

P15*), S. salmonicida (Ss), Trepomonas spp. (Trep), K. bialata (Kb)* and *T. vaginalis (Tv).*

Species	*Gm*	*Gi WB*	*Gi GS*	*Gi P15*	*Ss*	*Trep*	*Kb*	*Tv*
2.5-diketo-D-gluconic acid reductase	X		X				X	X

Arginase	X		X					X
Carboxymuconolactone decarboxylase	X		X					
Ferritin-like	X							X
Fructokinase	X							
Ketosteroid isomerase	X							
L-ascorbate-6-phosphate lactonase	X		X					
Maltose-O-acetyltransferase	X				X	X		
Mannose-6-phosphate isomerase	X				X			
Phosphopantothenate-cysteine ligase	X				X			
Quorum-quenching N-acyl-homoserine lactonase	X		X					
Ribonuclease 3	X						X	X
Tae4	X							
Tryptophanase	X				X	X		
β-phosphoglucomutase		X	X	X	X	X		
Extracellular nuclease		X	X	X	X	X	X	X
Flavohemoprotein		X	X	X				
Glycerol kinase		X	X	X	X	X		
Inositol-3-phosphate synthase		X	X	X				
Methyltransferase		X	X	X				

NADPH-ferrihemoprotein	X	X	X	X	X		
Purine nucleoside phosphorylase	X	X	X	X	X	X	X
Pyruvate phosphate dikinase	X	X	X		X		
Sugar/H$^+$ symporter	X	X	X				
Threonine dehydratase	X	X	X	X	X		

Genome streamlining and synteny

Gene prediction and manually curated annotation identified 4653 protein coding

genes in *G. muris* (Table 1). This makes 84.5% of the genome coding, counting also the

tRNAs and rRNAs (Table 1). Thus, the *G. muris* genome is an example of a very compact

eukaryotic genome. Consistent with that, the average intergenic size is 264 bp (Table 1),

with a prominent skew towards shorter intergenic regions for a high proportion of genes

(median size at 37 bp, Fig. 2C). The compactness of the genome is also illustrated by

multiple instances of overlapping genes, with 441 genes (9.5% of all genes) showing an

average overlapping size of 21 bp (spanning 1-327 bp) with neighboring genes.

The *G. muris* and the new improved *G. intestinalis* WB genome

(AACB03000000, F. Xu and S.G. Svärd) do not maintain clear chromosomal synteny

even though both are assembled as five near-complete chromosomes (Fig. S2).

However, local synteny (Fig. 2AB) was obtained among 3,043 one-to-one orthologs

(Table S1A) (an average amino acid similarity of 44.7%) shared by the two genomes.

Comparing local synteny, it becomes obvious that *G. muris* keeps shorter orthologous

gene and intergenic region sizes (Fig. 2BCD).

Gene regulation

We could not identify any universal, conserved promoter motifs shared by all

G. muris genes except for an enrichment of A residues around the start codon (Fig.

S3A), which resembles observations in *G. intestinalis*(39)*.* The streamlining of the *G.*

muris genome was also apparent at the 3' end of genes where the putative

polyadenylation signal, which is similar to the one described in *G. intestinalis*(39), is

overlapping with the stop codon for most genes (Fig. S3B). Genes up-regulated early

during encystation in *G. intestinalis* have specific promoter elements(40,41). Most of

these genes were also identified in *G. muris*, with one notable exception of cyst-wall

protein 3 (Fig. 3C). Encystation-related genes in *G. muris* share promoter motifs (Fig.

3A), similar to the Myb binding sites found in *G. intestinalis*(41), suggesting a similar

type of regulation. We also noted that the encystation-related genes are among the

most highly expressed genes in *G. muris* trophozoites *in vivo* (Fig. S3C, Table S1B),

similar to *G. intestinalis* infection in mice(42,43).

Very few genes in *G. intestinalis* contain introns, with only eight known *cis*-

spliced and four *trans*-spliced genes (five *trans*-introns)(44–46). Similarly, only three *cis*-

and no *trans*-introns were identified in the parasitic diplomonad *S. salmonicida*(35)*,*

whereas the free-living fornicate *K. bialata* has on average seven *cis*-introns per protein

encoding gene(31). *G. muris* maintains homologs to the eight *cis*-spliced *G. intestinalis*

genes, but has only three retained introns (Fig. S4A). All four *trans*-spliced genes in *G.*

intestinalis have homologs in *G. muris* with conserved splicing motifs (Fig. S4). Mining

genes with similar motifs did not reveal additional intron-containing genes in *G. muris*.

Similar to *G. intestinalis*(44) all the *trans*-spliced genes in *G. muris* preserve a similar

cleavage motif TCCTTTACTCAA (Fig. S4C) as the RNA processing sequence

motif(44). Thus, we observe a reduction of introns in *G. muris*, and *cis*-introns seem to be

easier to lose than *trans*-introns.

VSPs and antigenic variation in *G. muris*

Variant specific-surface proteins (VSPs) in *G. intestinalis* are characterized as

cysteine-rich proteins with frequent CXXC motifs and a conserved C-terminal

transmembrane (TM) domain followed by a cytoplasmic pentapeptide (CRGKA, Fig.

S5A). We identified 265 VSP homologs in *G. muris.* Their C-terminal pentapeptide

(GCRGK, Fig. S5A, Table S1C) differed slightly from that in *G. intestinalis*. However,

the cysteine and arginine residues in the pentapeptide, which are known to be post

translationally modified in *G. intestinalis*, are conserved(47,48). In addition, the

preceding 24 aa of the *G. muris* VSP TM domain show conservation to the TM domain of

G. intestinalis VSPs (Fig. S5A). Most *G. muris* VSPs contain the conserved GGCY motif

present in most *G. intestinalis* VSPs (Table S1C). Since *bona fide* VSPs need signal

peptides (SPs) at the N-terminus to guide VSPs to the parasite surface, we divided this

group into two subgroups; proteins with predicted SPs are called VSPs, whereas VSP

proteins without SPs are referred to as pseudogenized VSPs (ψVSPs). The 26 complete

VSP genes (16 unique at 98% identity to each other) are mostly located chromosome-

centrally (Fig. 1, Table 3). Seven pairs of VSP genes were identified with identical

sequences arranged either as head-to-head (2 pairs) or tail-to-tail (5 pairs) (Table 4).

Sequences in between the different VSP pairs resemble NimA (never in mitosis gene a)-

related kinase (NEKs), ankyrin repeat proteins (ARPs) and zinc-finger domains. There

are also 4 copies of identical VSPs clustering close to the 3' end of chromosome 2 (Fig.

2) with tandem repeats and sequences resembling NEKs and zinc-finger domains

interspersed (Table 4).

In contrast to complete VSPs, most ψVSPs (183 / 239) are found in linear

arrays (n=17) in *G. muris*, herein defined as having >3 ψVSPs genes (Fig. S5BC).

Strikingly, nine out of the ten ends of the main contigs have a ψVSP array at or close

to the ends of the chromosomes (telomere-adjacent) containing a total of 131 genes

(Table S1C). The only main contig that ends without an array has a cluster of two

ψVSPs close to the chromosome terminus. We found a single ψVSP array consisting

of 12 genes in a chromosome position that was non-telomere adjacent (on

chromosome 5) (Fig. 1). The ψVSP arrays vary in copy numbers (5-23 genes) and the

terminal part of the ψVSP array is always arranged with the tail-end towards the

chromosome terminus. The tandem arrangement of the gene arrays suggested that

they were generated by gene duplication. Two of the terminal arrays are scrambled

and have a shift in the ψVSP array directionality at the site of an intact VSP (Fig.

S5B).

We constructed a phylogeny of all the VSPs and ψVSPs in *G. muris* to

investigate their evolutionary dynamics. The phylogeny revealed relaxed clustering of

ψVSP genes originating in each linear array (Fig. S5C). However, the internal linear

array on chromosome 5 represents a noteworthy exception showcasing a very recent

gene duplication event. Interestingly, the great majority of full-length VSPs (23/28)

are clustered in the phylogeny, including the VSPs in the scrambled ψVSP arrays,

despite these genes being distributed in physically separate chromosomal locations

across all five primary scaffolds (Fig. S5C). The few non-clustered VSP genes in the

phylogeny that are not part of pairs or clusters are found directly adjacent to ψVSPs

genes. The relaxed clustering of VSPs and ψVSPs suggested that these genes might

be undergoing periodical recombination or gene conversion.

The VSPs and ψVSPs showed distinct expression patterns. Essentially all the

ψVSP genes were non-transcribed in the three surveyed life-stages (Fig. S5D). VSPs, on

the other hand, showed on average higher expression with one or a few loci displaying

dominant expression in the different life-stages (Fig. S5D).

There is also another, less characterized VSP-related cysteine-rich protein family

in *G. intestinalis*, High Cysteine Membrane Proteins (HCMPs)(49), with 62

members(32). Many are highly up-regulated during interaction with intestinal epithelial

cells(50). The HCMPs have several CXXC and CXC motifs, one VSP-like

transmembrane domain but with longer C-terminals than in the VSPs(50). The 34 genes

matching these criteria in the *G. muris* genome were named HCMPs after the

corresponding gene family in *G. intestinalis*. They are found spread-out on the five

chromosomes (Fig. 1).

Table 3 – Summaries of gene families within *Giardia*

	G. intestinalis *	*G. muris* *
NEK	184 (26)	216 (23)
ARP-1	269 (5)	298 (6)
ARP-2	33	86 (16)
ARP-3	8	33
VSP	133	26
ψVSP	208	239

247   *** Values in () indicate the number of pseudogenized copies.**

249   **Table 4** – Arrangement of VSP genes in the *G. muris* genome.

Chr	Geneid 1	Geneid 2	Arrangement	Gene size (aa)	Distance (bp)	Genes in between
1	20512	13275	-> <-	596	2025	ψARP-2
1	21145	21149	-> <-	594	1658	ψARP-2
1	13124	21374	-> <-	620	8400	ψARP-2, alpha-tubulin

2	16008	21957	<- ->	624	8794	NEK, ARP-2
2	22301	22304	-> <-	523	1656	ψARP-2
2	12920	22758 (22764, 22769)	<- ->	515	10612 (6824, 8428)	NEK, TRP, Zinc
4	24220	24228	-> <-	623	8303	ψARP-2, alpha-tubulin
5	24787	24792	<- ->	619	7158	NEK, Zinc, ARP-2

Multigene families in *G. muris*

The largest multigene families in *G. intestinalis* outside the VSPs and HCMPs are

the NEKs(51) and ankyrin repeat containing proteins (Protein 21.1)(32). There are 230

NEKs in *G. muris* (Fig. 1, Table 4), making up 71% of its kinome, slightly more than

what was found in *G. intestinalis*(51). We classified ankyrin repeat containing proteins

further into three groups. The ankyrin repeat protein-1 (ARP-1) with only ankyrin repeats,

ARP-2 with ankyrin repeats plus zinc finger domains, and ARP-3 with both ankyrin

repeats plus domains other than zinc finger domains. The NEKs and the different classes

of ARPs are scattered throughout the chromosomes without obvious clustering (Fig. 1). A

phylogenetic analysis revealed that 79 of the NEKs are conserved as 1:1 orthologs

between *G. muris* and *G. intestinalis* (Fig. S6A). Each species has one massively

expanded cluster of NEKs with *G. muris* having the largest with 104 members and the

one in *G. intestinalis* having 79 members. ARPs show a similar evolutionary stability

with 132 conserved 1:1 orthologs between species (Fig. S6B). *G. muris* shows a major

species-specific expansion of 91 genes whereas the largest expanded clusters in *G.*

intestinalis amounts to two groups of 15 genes each. The partly shared domain-structure

of NEKs and ARPs prompted us to investigate their relationship by a network analysis

employing reciprocal blastp (1e-05 cutoff). The two groups are not recovered as clearly

separated clusters but form partially overlapping networks, indicating that there might be

recombination or gene conversion in between (Fig. S6C). The larger numbers of genes in

these multigene families in *G. muris* compared to *G. intestinalis* and their evolutionary

dynamics are intriguing given the otherwise streamlined features of the *G. muris* genome,

perhaps suggesting that they have unique roles in adaptation to their murine hosts.

Virulence factors in *Giardia*

G. intestinalis is not known to possess classical virulence factors, such as

enterotoxins, but several genes are important for colonization of the host and thus for

pathogenesis. These include genes for motility(10), the adhesive disc for attachment(9),

secreted cysteine proteases that can degrade host defensive factors(12,13), and cysteine-

rich surface protein like the VSPs(52) and the HCMPs(49) that undergo antigenic

variation. The cytoskeletal protein repertoire in *G. muris* is very similar to *G. intestinalis*

apart from several fragmented alpha-tubulins (3 complete genes with homologs in *G.*

intestinalis and 9 incomplete gene fragments). The adhesive disc is a unique cytoskeletal

structure of *Giardia* parasites essential for attachment of the trophozoite in the small

intestine, but is missing in other fornicates(9). The first detailed studies of the adhesive

disc were performed on *G. muris* trophozoites(24), but more recent work has mostly

focused on *G. intestinalis*. The vast majority (82 of 85) of *G. intestinalis* disc proteins(9)

were also identified in *G. muris* (Table S1D)*;* 12 were NEK kinases and 27 were ARP-1

proteins. Two of the three *G. intestinalis* disc proteins not found in *G. muris* (Table S1D)

localize to a structure in the *G. intestinalis* disc on top of the ventral groove, but this

structure is missing in the *G. muris* disc(22), suggesting functional disc differences. Many

of the disc proteins that are immunodominant during *G. intestinalis* infections (e.g. alpha-

giardins, beta-giardin, SALP-1, alpha- and beta-tubulin(53) are highly expressed (here

defined as >500 FPKM) in *G. muris* trophozoites in the small intestine (Table S1B).

Proteases are important virulence factors in many pathogens and cysteine-protease

activities have been suggested to play a role in *Giardia* virulence(4,13). We identified 81

proteins classified as proteases in the *G. muris* genome, compared to 96 proteins

identified in an identical search in *G. intestinalis* WB (Table S1E). The largest family of

proteases in *G. muris*, with 15 members*,* are papain-like cysteine proteases (C1A family).

This protein family is also the largest protease group in *G. intestinalis* with 21

members(54,55). Several conserved groups of proteases were found to have been present

in the ancestor to *Giardia* and *Spironucleus*, although we also found evidence for lineage-

specific gene loss and expansion in *G. muris* (Fig. S7). The most highly expressed

cysteine protease of *G. muris in vivo* is the closest homolog to the highest expressed

protease in *G. intestinalis* WB(55).

Metabolic pathways in *G. muris*

Our metabolic reconstruction identified 95 metabolic pathways in *G. muris*

compared to 98 pathways detailed in *G. intestinalis* WB; four pathways were unique in *G.*

muris and eight in *G. intestinalis* WB. Even though the overall metabolism is highly

similar between *G. muris* and *G. intestinalis* WB, the genes for several specific enzymes

and their putative reactions show distinct differences. Thus, 18 unique reactions (14

enzymes) were predicted in *G. muris* and 25 in *G. intestinalis* WB (10 enzymes) (Table

2). Several of these unique proteins showed moderate-high identity to prokaryotic

proteins (Table S1F). Five of these prokaryote-like genes are present in the genome of

assemblage B strain *G. intestinalis* GS.

The potential utilization of carbohydrate sources for glycolysis is different in *G.*

muris compared to *G. intestinalis*. Fructokinase and mannose 6-phosphate isomerase

enable *G. muris* to use fructose and mannose 6-phosphate unlike *G. intestinalis*. In both

cases, the enzymes were acquired from bacteria via lateral gene transfer (Table S1F,

Figure S8AB). Curiously, the *G. muris* gene for mannose 6-phosphate isomerase is found

next to a bacterial transcriptional regulator/sugar kinase gene. Phylogenetic analyses

show that *G. muris* mannose 6-phosphate isomerase and the transcriptional

regulator/sugar kinase gene clusters deep in the Bacteroidetes group. This gene

arrangement is observed in bacteria of the genus *Alistipes*, whose genomes harbor the

most similar homologs, supporting the notion that a single event of lateral gene transfer

best explains the origin of these genes in *G. muris* (Figure S8BC). Both genes have the A-

rich initiator that precedes the start codon in most *G. muris* genes and are expressed in *G.*

muris trophozoites (Table S1B).

The utilization of glycerol for ATP synthesis via glycerol kinase has been

suggested in *G. intestinalis* upon depletion of primary carbon sources(56). This enzyme is

found in both *Spironucleus* and *Trichomonas* but has been lost in *G. muris*. Another

notable metabolic difference to *G. intestinalis* WB is the lack of pyrophosphate-

dependent pyruvate phosphate dikinase (PPDK) that leaves a single, less energy efficient,

route from phosphoenol pyruvate to pyruvate via pyruvate kinase in *G. muris*.

G. muris is predicted to synthesize coenzyme A from pantothenate, employing a

bifunctional phosphopantothenoyl decarboxylase-phosphopantothenate synthase. The

same pathway is described in *S. salmonicida*(35), but the complete pathway is missing in

G. intestinalis (Table 2).

As in all other studied metamonads, *G. muris* encodes the arginine dihydrolase

(ADH) pathway that enables the use of arginine as an energy source and at the same time,

reduces the available free arginine in the environment, preventing nitric oxide (NO)

production in host cells(57). NO efficiently kills *G. intestinalis* trophozoites and the main

scavenger enzyme for NO in *G. intestinalis* is Flavohemoprotein(58), which is lacking in

G. muris (Table 2). Arginine is an important modulator of virulence in many infectious

organisms since it interferes with NO production(59). *G. muris* encodes arginases which

converts arginine directly to ornithine and urea. Arginases are present in *G. intestinalis*

GS and *T. vaginalis,* representing an ancestral acquisition in Metamonada followed by

subsequent losses in *Spironucleus* and *G. intestinalis* assemblage A and E (Table 2 and

Fig. S8D).

We noticed that several of the bacterial derived genes that are shared with *G.*

intestinalis GS are clustered together on the chromosomes in *G. muris* in highly dynamic

genomic regions. For example, arginase genes are found in a four-gene genomic region

that is present in two adjacent copies in chromosome 2, two on chromosome 1 and one on

chromosome 5 (dark grey filled arrows, Fig. 4A). Intact arginase genes are only found on

chromosome 2 adjacent to the genes encoding 2,5-diketo-D-gluconic acid reductase. All

the duplication events have occurred between ARPs (red arrows) and NEKs (dark red

arrows). The homologous regions on the three chromosomes likely originated via two

duplication events.

Nucleotide substitutions have accumulated since the duplication events and in-

frame stop codons (marked by red asterisk in Fig. 4A) have rendered three of the arginase

genes pseudogenized. The small ORFs sitting at the other side of arginase are hypothetical

proteins and have similar homologs in other parts of the genome, but the sequences of their

duplicated homologs in those regions are pseudogenized (dotted light grey block in Fig.

4A). Both genes have their closest relatives in bacteria, even though the genes can be

found in other eukaryotes (Fig. S13E). Phylogenetic analyses indicate the genes might

have been transferred from different bacterial donors multiple times into different

eukaryotic lineages (Fig. S13E).

A second, distantly related 2,5-diketo-D-gluconic acid reductase gene copy (Fig.

4B, Fig. S8F) in the genome is present in three different genomic locations together with

two other enzymes, carboxymuconolactone decarboxylase (CMD) and ketosteroid

isomerase-like protein. All three genes, constitute putative lateral gene transfers (Fig.

S8GH). This three-gene region is close to ψVSPs (orange arrows) and ARPs.

372
Interaction with the intestinal microbiota

Giardia trophozoites colonize the intestinal lumen where they can potentially

interact with other intestinal microbes. Although little is functionally known about such

interactions and their consequences for parasite survival, four proteins encoded in the *G.*

muris genome could play a role in these interactions.

Bactericidal/permeability-increasing (BPI) proteins are innate immune defense

proteins that bind to lipopolysaccharide and display potent killing activity against gram-

negative bacteria by increasing membrane permeability. Beyond this basic function BPI

proteins might also act as effectors in controlling mutualistic symbioses(60). Homologs of

BPI proteins are found in *G. muris* and *G. intestinalis*(61), but it remains to be determined

if they have anti-microbial activity.

G. muris encodes tryptophanase, an enzyme that metabolizes tryptophan to

pyruvate with concomitant release of indole and ammonia. While pyruvate can be utilized

in energy metabolism, indole and its metabolites have been shown to affect gut

microbiota composition, possibly by interfering with quorum-sensing systems, and might

be able to influence host health(62). Phylogenetic analysis of this protein showed that this

enzyme represents an ancestral acquisition in diplomonads with subsequent loss in *G.*

intestinalis (Fig. S8I).

Two more proteins with potential importance for microbiota interactions are

encoded in *G. muris*: Tae4 and quorum-quenching N-acyl-homoserine lactonase. The

Tae4 proteins are wide-spread amidases that were first described in association with the

T6SS system effector Tae4 in *Salmonella* Typhimurium(63). The Tae4 proteins degrade

bacterial peptidoglycan by hydrolyzing the amide bond, γ-D-glutamyl-mDAP (DL-bond)

of Gram-negative bacteria(63), and is required for interbacterial antagonism and

successful gut colonization by *S.* Typhimurium(64). Quorum-quenching N-acyl-

homoserine lactonase degrades N-acyl-homoserine lactone, a molecule used by both

Gram-positive and Gram-negative bacteria, for quorum sensing(65). Our phylogenetic

analysis supports the lateral acquisition of both genes (Fig. S8JK). While Tae4 has been a

recent acquisition in *G. muris*, quorum-quenching N-acyl-homoserine lactonase was

present in the common ancestor of *G. muris* and *G. intestinalis* and lost in *G. intestinalis*

WB and P15 (Fig. S8K).

Discussion

Our data shows that *G. muris* has an even more compact genome than *G. intestinalis*,

whose genome is already known to be highly streamlined(32). Genome compaction

via reduction of mobile or repetitive elements have been seen in other eukaryotic

parasites(1,66). *G. muris* appears to fall into this category as it encodes no known

classes of mobile elements and repetitive elements are mostly confined to telomeric

contexts. The shortness of intergenic regions in *G. muris* ranks among the most

extreme recorded for any eukaryote, even shorter than Microsporidia which are

known as the most compact and reduced eukaryotic genomes(67). The global synteny

map of *G. muris* to *G. intestinalis* indicates many frequent small-scale genome

rearrangements that often favors a more efficient gene packing in *G. muris* thus

allowing shorter intergenic regions. This evidence of gene shuffling and the fact that

there is very little evidence of genome degradation would argue for optimization of

growth as the driving force of *G. muris* genome streamlining.

G. muris trophozoites have not been grown axenically *in vitro*, which has

hampered exploration of its genome, gene regulation and metabolism(5,21), and has

limited the use of *G. muris* as an *in vitro* model system for the human parasite *G.*

intestinalis and other intestinal protozoan parasites(5). We identified several

metabolic differences between *G. muris* and *G. intestinalis* that might indicate

avenues to successful strain axenization. Most of these differences are represented by

instances of lateral gene transfer of metabolic genes or losses thereof in either *G.*

intestinalis or *G. muris*. *G. intestinalis* isolates are typically poor at infecting mice.

Despite this, the assemblage B isolate GS, which shares more metabolic enzymes with

G. muris than the assemblage A isolate WB, is better able to establish infection in

mice than WB. This suggests that the shared metabolic capacity of *G. muris* and *G.*

intestinalis GS enables survival in the murine intestinal tract. Additionally, *G. muris*

might be able to interact or interfere with intestinal Gram-positive and Gram-negative

bacteria and this could be a key to establish successful infections. *G. intestinalis*,

which lacks some of the putative microbiome modulators, such as Tryptophanase and

Tae4, is dependent on reduction of the small intestinal microbiota in order to

efficiently infect mice(68). *G. muris* is cleared from the murine host by secretory

IgA(26), whereas the role of IgA in anti-giardial defense is less clear for *G.*

intestinalis GS(26,69). We speculate that *G. muris* is more resistant to elimination by

innate factors such as competition with the normal microbiota, or host production of

reactive oxygen species and/or NO, whereas GS is more sensitive to innate factors

and eliminated much faster within 1-3 weeks (while *G. muris* clearance requires 4-8

weeks). Future insights into the importance of innate factors in *G. muris* infection

should be facilitated by the availability of the complete genome sequence.

Sub-telomeric regions in parasitic protozoa often contain arrays of expanded

gene families that are under positive selection by the immune system(66). The relaxed

evolutionary pressure offered by keeping pseudogenized copies of surface antigens

might be an advantage for *G. muris* that allows genetic drift and recombination to

drive rapid and stealthy diversification, thus avoiding elimination by adaptive immune

defenses. It was previously reported that *G. muris* is capable of antigenic variation

and encodes VSP genes with high similarity and conserved structural features

(CRGKA pentapeptide) to those *G. intestinalis*(70). We failed to identify close

homologs to the previously sequenced *G. muris* VSPs in our *G. muris* genome. The

linear ψVSP arrays in *G. muri*s have previously been described in *G. intestinalis*(71).

Our phylogenetic analyses of *G. muris* VSPs and ψVSPs revealed evidence of

recombination or segmental gene conversion, as previously demonstrated in *G.*

intestinalis(71). However, we recognized two clear differences in the VSP repertoire

in *G. muris* and *G. intestinalis*. First, *G. muris* encodes a low number of intact VSP

loci that are located internally on the chromosomes. Second, the ψVSP arrays are

almost exclusively telomere adjacent, as opposed to *G. intestinalis* where this

tendency is not apparent(71). These aspects of the *G. muris* VSP repertoire resemble

the antigenic variation systems of *Pneumocystis* spp. and *Trypanosoma brucei*

(72,73). Despite clear mechanistic differences, all these systems have converged on

having large reservoirs of mostly telomeric positioned, arrayed genes that are

transcriptionally silent and are sources for recombination and gene conversion into

expression sites.

The function of ankyrin repeat proteins and NEK kinases remains mostly

unknown. They represent, together with VSPs, the most dynamic protein families in

the *Giardia* genomes(51). The *Giardia* NEK kinases lack transmembrane domains

and have been suggested to target and localize to different intracellular structures with

their ankyrin repeats(51) and many of the *G. intestinalis* NEK kinases localize to

cytoskeletal structures, including the flagella and adhesive disc(9). Rearrangements

and duplications in the *G. muris* genome are frequently associated with these large

gene families (Fig. 2A and Fig. 4), indicating they might serve as anchoring-points for

recombination.

Lateral gene transfer is an important shaping factor in the evolution of

metabolism in protists(74). The origins of laterally transferred genes in *G. muris* are

here inferred to be by prokaryotic sources that are members of the gastrointestinal

flora, in agreement to previous observations(75). Most of the putative differences in

metabolic potential in the *Giardia* genomes are attributable to lateral gene transfers,

either by lineage specific gene gain or loss. For example, the ability to utilize

mannose has been introduced from bacteria of the genus *Alistipes* via lateral gene

transfer. This event is supported by phylogenetic reconstruction, shows a high degree

of sequence conservation (>70% at the amino acid level) and displays maintained

gene order to the one seen in the closest related bacterial lineages (Figure S8B).

Interestingly, several lateral gene transfers were found clustered in amplified areas of

the *G. muris* chromosome. Curiously, *G. intestinalis* assemblage B also maintains

clustered copies of arginase and 2,5-diketo-D-gluconic acid reductase, while these

genes have both been lost in the *G. intestinalis* assemblage A and E lineages.

Anti-microbial peptides of several classes, such as defensins and trefoil-factor 3,

are up-regulated in the small intestine of *G. muris* infected mice(29). Secreted cysteine

proteases from *G. intestinalis* have been shown to be able to degrade defensins(12). We

detected prominent expression of several cysteine proteases in *G. muris*. The protease

with the highest expression is suggested to have a role in encystation and excystation(55).

Its *G. intestinalis* homolog is up-regulated and secreted during interactions with human

intestinal epithelial cells(8) and it cleaves chemokines, tight junction proteins and

defensins(12,76). Thus, this is most likely also an important virulence factor in *G. muris*.

Our results from this study are summarized in a model of the evolution of

Giardia's virulence traits in Figure 5. A number of characters important for *Giardia*'s

ability to infect the intestine of mammals are pre-parasitic inventions (such as

modified mitochondria and differentiation into transmissive cysts, Fig. 5) and some

are found in all diplomonad parasites (e.g. loss of metabolic functions, streamlined

microtubular cytoskeleton, expansion of gene families like ankyrins and cysteine

proteases and loss of introns, Fig. 5). *Giardia* specific innovations include the

adhesive disc for attachment, VSPs and High Cysteine Membrane Proteins for

antigenic variation and mitosomes involved in Fe-S complex synthesis (Fig. 5).

Whereas some are only found in *G. muris* (e.g. metabolic genes involved in

microbiota interactions, Fig. 5), suggesting adaptation to the intestinal environment of

mice. Our data shows that the environment in the host's intestine, most of all the

immune system and the microbiota, apply selective pressure for changes in the

genome, metabolic potential and the parasite surface proteome.

Materials and methods

Cell preparation and nucleic acid extraction.

4.5×10^7 muris trophozoites (day 7 post infection) were collected from small intestines

of three C57 mice, washed once in PBS and pellet frozen at -80°C (Biosample

SAMN11231832). Viable cysts of *G. muris* isolate Roberts-Thomson passaged

through mice were obtained from Waterborne Inc. These cysts had been purified from

fecal material using Percoll and sucrose gradients (Biosample SAMN11231833).

DNA and RNA were extracted from 1×10^7 cysts using standard methods.

1×10^8 cysts were excysted according to the procedure in Feely et al.(77) (Biosample

SAMN11231834). RNA was purified from cell material equivalent to 1×10^7 cysts.

DNA for long-read sequencing was prepared from the remaining cysts as described in

Methods S1.

Sequencing, assembly and annotation

Total genomic DNA was sequenced using both Illumina MiSeq and PacBio RS II

sequencers. The stranded transcriptome mRNA and the miRNA libraries were

sequenced with Illumina HiSeq 2000 system. The RNA samples extracted from

excysted cells and cysts prior excystation were prepared using the TruSeq stranded

mRNA sample preparation kit and sequenced by HiSeq 2500.

PacBio long reads were assembled de *novo* using the SMRT Analysis (v2.3.0)

pipeline. A detailed description of genome sequencing, assembly, annotation, synteny

analyses and RNA-Seq is available in Methods S1.

Pathway analysis

The metabolic pathways of *G. muris* and *G. intestinalis* WB were predicted with a

combination of BlastKOALA(78) implemented in KEGG(79), Pathway Tools

v21.5(80) and GiardiaDB(81). The different predictions were combined and manually

curated under Pathway Tools(80). Pathway Tools function pathway hole filler(82)

was used to further complete the pathway, and transport inference parser(83) was

used to infer transport reaction(s) for transporters which were then verified with

Conserved Domain databases(84).

Phylogenetic analysis

G. muris sequences were used as queries to retrieve at least 5000 hits with e-value

<0.001, using BLASTP against the nr database and the organism-specific proteomes.

The datasets were aligned in the forward and reverse orientation using MAFFT

v6.603b(85) and PROBCONS v1.12(86). The four resulting alignments were

combined with T-COFFEE(87) and trimmed by BMGE v1.12(88). Maximum

Likelihood (ML) trees were computed using IQtree v1.6.5(89) under LG4X

substitution model(90). Branch supports were assessed using ultrafast bootstrap

approximation (UFboot)(91) with 1,000 bootstrap replicates and 1,000 replicates for

SH-like approximate likelihood ratio test (SH-aLRT)(92). A detailed description of

the phylogenetic analyses is available in Methods S1.

Figures

Figure 1. **Circular representation of the *G. muris* chromosomes.** From outside

inward: five chromosomes, GC percent, unique genes (grey) including unique

metabolic genes in Table 2 (red), ARPs (greenblue) / NEKs (pink), VSPs (orange) /

ψVSPs (blue) / HCMPs (purple), Coding percent / 5 kbp (green if <= 0.5), # SNPs /

kbp >= 5 (red), BLASTN matches with >95% identity and > 1000 bp in size. Circular

plot was drawn with circlize(93).

Figure 2. **Examples of synteny between *G. muris* and *G. intestinalis*. A**) A 50 kbp

region on chromosome 3 which share synteny to a 58 kbp region on chromosome 5 in

WB. Synteny plot was plotted using genoplotR(94). Shades of red and blue represent

forward and inverted matches between orthologs. Genes are drawn as arrows in blue.

ARPs in red, NEKs in dark red, and VSPs in orange. Dark grey filled genes are

unique genes to that genome in comparison to the other. **B**) A 14 kbp region on

chromosome 1 which shares synteny to a 16 kbp region on chromosome 5 in *G.*

intestinalis. It uses the same color scheme as in A. **C)** Violin plots of intergenic sizes

of neighboring positional orthologs of *G. muris* and *G. intestinalis*, and the grey

vertical line represents the median intergenic size of *G. muris*. **D)** Violin plots of

positional orthologs sizes of *G. muris* and *G. intestinalis*, and the grey vertical line

represents the median ortholog size of *G. muris*. **E)** Histogram of the positional

ortholog size difference between *G. muris* and *G. intestinalis*.

Figure 3. **Gene regulation and organization of VSPs in *G. muris*. A**) Promoter

motifs shared by encystation-related genes. Motif 1 (gold in B) represents the general

promotor motif positioned directly adjacent to the start codon. Motif 2 (teal in B)

resembles the encystation-regulated promoter previously identified in *G. intestinalis*

(52). **B)** The distribution and position of motif 1 (gold) and motif 2 (teal) in chosen

genes regulated during encystation. **C)** *Giardia* cyst wall proteins. Cyst wall protein 3

is missing in *G. muris*. Signal peptide (pink). Acidic LRR-domain (grey). C-terminal

basic extension (green).

Figure 4. **Synteny plot of two duplicated regions in the genome. A, B)** Shades of

red and blue represent forward and inverted matches between neighboring sequences.

ARPs are drawn in red, NEKs in dark red and VSPs in orange. Pseudogenized genes

are drawn in dashed lines. Dark grey filled genes are unique genes to *G. muris* in

comparison with *G. intestinalis*. Point mutation in arginase is marked in red asterisk.

Homologous sequences that are not annotated in the genome are drawn in dashed box

on sides of the backbone grey line. Genes discussed in the paper: 21985 and 21992

encode the intact arginase genes, 21986 and 21992 encode the enzyme 2,5-diketo-D-

gluconic acid reductase, whereas 21984 and 21994 are hypothetical proteins in A;

14480 encodes 2,5-diketo-D-gluconic acid reductase, 24746 encodes CMD and 24745

encodes ketosteroid isomerase-like protein in B.

Figure 5. Model of the evolution of virulence traits in *Giardia* parasites. A set of

important diplomonad evolutionary innovations and their chronology is depicted at

relevant phylogenetic nodes. A) Free-living fornicate ancestor. B) Diplomonad ancestor.

C) *Giardia* ancestor. D) *Giardia muris*.

Supporting information

Figure S1. **A)** *G. muris* cells propagate as trophozoites in the mice small intestine.

The cell trophozoites undergo encystation into cysts and are shed in the feces. The

cysts are prompted to excyst as they encounter the conditions of the digestive tract.

The excyzoites transition into the vegetative trophozoites. Cell material for genomic

DNA was harvested from *G. muris* cysts from infected mice. **B)** Sub-telomeric

regions in *G. muris*. Sequence motifs in the 10 unitigs terminating in $(TAGGG)^n$

telomeric sequences (purple). Ribosomal DNA sequences (black). Satellite-like

repeated sequences (green).

Figure S2. Chromosome-scale synteny between *G. muris* and *G. intestinalis* WB.

G. muris chromosomes are color coded. The links between the two genomes

representing sequence similarity identified by MUMmer promer(95), are colored after

the *G. muris* chromosomal color. Circular plot was drawn with circlize(93).

Figure S3. Sequence logo around start codon (A) and stop codon (B) in *G. muris*.

C) *G. muris* gene expression from *in vivo* trophozoites. Genes showing top-ranked

FPKM ratios (log2 (Trophs/Cysts)) between *in vivo* derived trophozoites and cysts

(cut-off value ≥4). The value is indicated in the teal circle. An orange dot indicates

that the ortholog of this gene is upregulated (>2 fold) in *G. intestinalis* at 7 hrs, 12 hrs

or 22 hrs into encystation (40).

Figure S4. *Trans*- and *cis*-introns in *G. muris*. A) Aligned *trans*- and *cis*-splice

junctions in *G. muris*. **B)** Splice-site associated motifs in *trans*-spliced exons. Base-

pairing stems are underlined. Bases in red denote the predicted cleavage motif. **C)**

Consensus alignment of the cleavage motifs in *G. muris*. **D-E)** Predicted secondary

structures of **D)** *cis*-spliced and **E)** *trans*-spliced genes. The secondary structure was

predicted using the mfold webserver using default settings and manually curated. GU-rich

stretches and cleavage motifs are boxed.

Figure S5. A) Conservation of the VSP C-terminal in *G. muris* and *G. intestinalis*.

Sequence logos of conserved motifs in the C-terminal of VSPs in *G. muris* (n=25)

(upper panel) and *G. intestinalis* WB (n=183) (lower panel) were generated using

MEME v5.0.5. Transmembrane and cytosolic pentapeptides were predicted by the

Phobius webserver. **B) VSP and ψVSP evolutionary dynamics.** The ψVSP arrays

(defined as >3 ψVSP genes arrayed) in the *G. muris* genome arranged by

chromosome, termini arranged to the right. Arrays of ψVSPs without chromosomal

context are shown below as orphan arrays. **C)** VSP cluster mostly together in a

phylogeny whereas most ψVSP gene arrays show relaxed clustering. Intact VSP

genes are colored red, while ψVSP genes are colored according to their resident

ψVSP array in panel A. Genes with black label are not found in arrays. **D)**

Expression measurements of VSP and ψVSP in *G. muris*. VSP and ψVSP RNA-

seq read counts (FPKM) from trophozoites, cysts and excyzoites. Size of the circle

corresponds to expression level.

Figure S6. A) Phylogenetic tree of *G. muris* and *G. intestinalis* NEKs. B) Phylogenetic tree of *G. muris* and *G. intestinalis* ARPs. C) Network analysis of *G. muris* and *G. intestinalis* NEKs and ARPs. *G. muris* genes are represented in circle whereas *G. intestinalis* genes are shown in triangle. Blue indicates NEK and pink as ARP. Edges are weighted and scaled by reciprocal BLAST scores with evalue 1e-05 as the cutoff.

Figure S7. Phylogenetic tree of C1 family cysteine proteases in diplomonads. Black: *G. muris*, Red: *S. salmonicida*, Cyan: *G. intestinalis* GS/M, Green: *G. intestinalis* P15, Purple: *G. intestinalis* WB.

Figure S8. Phylogenetic analyses of HGT genes. A) Fructokinase. **B)** Mannose-6-phosphate isomerase. **C)** Transcriptional regulator/sugar kinase. **D)** Arginase. **E)** 2,5-diketo-D-gluconic acid reductase 1. **F)** 2,5-diketo-D-gluconic acid reductase 2. **G)** Carboxymuconolactone decarboxylase. **H)** Ketosteroid isomerase. **I)** Tryptophanase. **J)** Tae4. **K)** Quorum-quenching N-acyl-homoserine lactonase.

Table S1. A) One-to-one ortholog list between *G. muris* and *G. intestinalis* WB. Orthologs inferred by OrthoMCL. **B)** RNA-Seq expression values in excel file. Sorted according to FPKM values. **C)** VSP and ψVSP gene metadata. '-' in the pentapeptide motif represents lack of the 3' end sequence. **D)** List of ventral disc proteins in *G. muris*. **E)** Proteases detected in *G. muris* and *G. intestinalis*. **F)** Detailed information about hits of unique genes in *G. muris*.

655 **Methods S1. Additional methods section**

Acknowledgements

658 Elaine Hanson is acknowledged for technical support performing trophozoite

659 purifications from infected mice.

660 Phylogenetic analyses were performed on resources provided by the Swedish National

661 Infrastructure for Computing (SNIC) at Uppsala Multidisciplinary Center for Advanced

662 Computational Science (UPPMAX).

Declarations

Ethics approval and consent to participate

All animal studies were approved by the Institutional Animal Care and Use

Committees of the University of California, San Diego, USA.

Consent for publication

No applicable.

Availability of data and material

Raw DNA and RNA sequence reads are archived at NCBI Sequence Read Archive

(SRA) under accession number SRR8858297 - SRR8858305.

This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank

under the accession PRJNA524057. The version described in this paper is version

VDLU00000000.1. The datasets generated and analyzed in this project are also

available via GiardiaDB.

Competing interests

The authors declare that they have no competing interests.

Funding

The project was supported by funding from Vetenskapsrådet-M (www.vr.se) grant

(2017-02918) to SGS. LE was supported by National Institutes of Health

(www.nih.gov) grant DK120515. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

SGS, LE, JA and JJH conceived the study. JJH and EE performed RNA and DNA

extractions. FX assembled the genome. FX, AJG, JJH, AA, DP, JA, SGS and EE

annotated the genome. FX, JJH, AJG and SGS analyzed the data. All authors wrote

the paper.

References

1.   Poulin R, Randhawa HS. Evolution of parasitism along convergent lines: from ecology to genomics. Parasitology. 2015 Feb;142(S1):S6–15.

2.   Hupalo DN, Bradic M, Carlton JM. The impact of genomics on population genetics of parasitic diseases. Current Opinion in Microbiology. 2015 Feb;23:49–54.

3.   Jackson AP, Otto TD, Aslett M, Armstrong SD, Bringaud F, Schlacht A, et al. Kinetoplastid Phylogenomics Reveals the Evolutionary Innovations Associated with the Origins of Parasitism. Current Biology. 2016 Jan;26(2):161–72.

4.   Einarsson E, Ma'ayeh S, Svärd SG. An up-date on Giardia and giardiasis. Current Opinion in Microbiology. 2016 Dec;34:47–52.

5.   Cacciò SM, Lalle M, Svärd SG. Host specificity in the Giardia duodenalis species complex. Infect Genet Evol. 2018;66:335–45.

6.   Fink MY, Singer SM. The Intersection of Immune Responses, Microbiota, and Pathogenesis in Giardiasis. Trends in Parasitology. 2017 Nov;33(11):901–13.

7.   Ma'ayeh SY, Knörr L, Sköld K, Granham A, Ansell BRE, Jex AR, et al. Responses of the Differentiated Intestinal Epithelial Cell Line Caco-2 to Infection With the Giardia intestinalis GS Isolate. Front Cell Infect Microbiol. 2018;8:244.

8.   Ma'ayeh SY, Liu J, Peirasmaki D, Hörnaeus K, Bergström Lind S, Grabherr M, et al. Characterization of the Giardia intestinalis secretome during interaction with human intestinal epithelial cells: The impact on host cells. PLoS Negl Trop Dis. 2017;11(12):e0006120.

9.   Nosala C, Hagen KD, Dawson SC. 'Disc-o-Fever': Getting Down with Giardia's Groovy Microtubule Organelle. Trends in Cell Biology. 2018 Feb;28(2):99–112.

10. Nosala C, Dawson SC. The Critical Role of the Cytoskeleton in the Pathogenesis of Giardia. Current Clinical Microbiology Reports. 2015 Dec;2(4):155–62.

11. McInally SG, Dawson SC. Eight unique basal bodies in the multi-flagellated diplomonad Giardia lamblia. Cilia [Internet]. 2016 Dec [cited 2019 May 27];5(1). Available from: http://ciliajournal.biomedcentral.com/articles/10.1186/s13630-016-0042-4

12. Liu J, Ma'ayeh S, Peirasmaki D, Lundström-Stadelmann B, Hellman L, Svärd SG. Secreted Giardia intestinalis cysteine proteases disrupt intestinal epithelial cell junctional complexes and degrade chemokines. Virulence. 2018 Dec 31;9(1):879–94.

13. Ortega-Pierres G, Argüello-García R, Laredo-Cisneros MS, Fonseca-Linán R, Gómez-Mondragón M, Inzunza-Arroyo R, et al. Giardipain-1, a protease secreted

by Giardia duodenalis trophozoites, causes junctional, barrier and apoptotic damage in epithelial cell monolayers. International Journal for Parasitology. 2018 Jul;48(8):621–39.

14. Amat CB, Motta J-P, Fekete E, Moreau F, Chadee K, Buret AG. Cysteine Protease–Dependent Mucous Disruptions and Differential Mucin Gene Expression in Giardia duodenalis Infection. The American Journal of Pathology. 2017 Nov;187(11):2486–98.

15. Cotton JA, Bhargava A, Ferraz JG, Yates RM, Beck PL, Buret AG. Giardia duodenalis Cathepsin B Proteases Degrade Intestinal Epithelial Interleukin-8 and Attenuate Interleukin-8-Induced Neutrophil Chemotaxis. Appleton JA, editor. Infection and Immunity. 2014 Jul;82(7):2772–87.

16. Cotton JA, Motta J-P, Schenck LP, Hirota SA, Beck PL, Buret AG. Giardia duodenalis Infection Reduces Granulocyte Infiltration in an In Vivo Model of Bacterial Toxin-Induced Colitis and Attenuates Inflammation in Human Intestinal Tissue. Wallace J, editor. PLoS ONE. 2014 Oct 7;9(10):e109087.

17. Allain T, Amat CB, Motta J-P, Manko A, Buret AG. Interactions of Giardia sp. with the intestinal barrier: Epithelium, mucus, and microbiota. Tissue Barriers. 2017 Jan 2;5(1):e1274354.

18. Mendez TL, De Chatterjee A, Duarte T, De Leon J, Robles-Martinez L, Das S. Sphingolipids, Lipid Rafts, and Giardial Encystation: The Show Must Go On. Current Tropical Medicine Reports. 2015 Sep;2(3):136–43.

19. Stadelmann B, Merino MC, Persson L, Svärd SG. Arginine Consumption by the Intestinal Parasite Giardia intestinalis Reduces Proliferation of Intestinal Epithelial Cells. PLoS ONE. 2012;7(9).

20. Eckmann L, Laurent F, Langford TD, Hetsko ML, Smith JR, Kagnoff MF, et al. Nitric Oxide Production by Human Intestinal Epithelial Cells and Competition for Arginine as Potential Determinants of Host Defense Against the Lumen-Dwelling Pathogen Giardia lamblia. The Journal of Immunology. 2000 Feb 1;164(3):1478–87.

21. Helmy YA, Spierling NG, Schmidt S, Rosenfeld UM, Reil D, Imholt C, et al. Occurrence and distribution of Giardia species in wild rodents in Germany. Parasites & Vectors [Internet]. 2018 Dec [cited 2019 May 27];11(1). Available from: https://parasitesandvectors.biomedcentral.com/articles/10.1186/s13071-018-2802-z

22. Friend DS. THE FINE STRUCTURE OF GIARDIA MURIS. The Journal of Cell Biology. 1966 May 1;29(2):317–32.

23. Dann SM, Le CHY, Hanson EM, Ross MC, Eckmann L. Giardia Infection of the Small Intestine Induces Chronic Colitis in Genetically Susceptible Hosts. The Journal of Immunology. 2018 Jul 15;201(2):548–59.

24. Holberton DV. Fine structure of the ventral disk apparatus and the mechanism of attachment in the flagellate Giardia muris. J Cell Sci. 1973 Jul;13(1):11–41.

25. Schaefer FW, Rice EW, Hoff JC. Factors promoting in vitro excystation of Giardia muris cysts. Trans R Soc Trop Med Hyg. 1984;78(6):795–800.

26. Langford TD, Housley MP, Boes M, Chen J, Kagnoff MF, Gillin FD, et al. Central importance of immunoglobulin A in host defense against Giardia spp. Infect Immun. 2002 Jan;70(1):11–8.

27. Davids BJ, Palm JED, Housley MP, Smith JR, Andersen YS, Martin MG, et al. Polymeric immunoglobulin receptor in intestinal immune defense against the lumen-dwelling protozoan parasite Giardia. J Immunol. 2006 Nov 1;177(9):6281–90.

28. Dreesen L, De Bosscher K, Grit G, Staels B, Lubberts E, Bauge E, et al. Giardia muris infection in mice is associated with a protective interleukin 17A response and induction of peroxisome proliferator-activated receptor alpha. Infect Immun. 2014 Aug;82(8):3333–40.

29. Manko A, Motta J-P, Cotton JA, Feener T, Oyeyemi A, Vallance BA, et al. Giardia co-infection promotes the secretion of antimicrobial peptides beta-defensin 2 and trefoil factor 3 and attenuates attaching and effacing bacteria-induced intestinal disease. Adam RD, editor. PLOS ONE. 2017 Jun 16;12(6):e0178647.

30. Saghaug CS, Sørnes S, Peirasmaki D, Svärd S, Langeland N, Hanevik K. Human Memory CD4+ T Cell Immune Responses against Giardia lamblia. Wilkins PP, editor. Clinical and Vaccine Immunology. 2016 Jan;23(1):11–8.

31. Tanifuji G, Takabayashi S, Kume K, Takagi M, Nakayama T, Kamikawa R, et al. The draft genome of Kipferlia bialata reveals reductive genome evolution in fornicate parasites. Schierwater B, editor. PLOS ONE. 2018 Mar 28;13(3):e0194487.

32. Morrison HG, McArthur AG, Gillin FD, Aley SB, Adam RD, Olsen GJ, et al. Genomic minimalism in the early diverging intestinal parasite Giardia lamblia. Science. 2007 Sep 28;317(5846):1921–6.

33. Franzén O, Jerlström-Hultqvist J, Castro E, Sherwood E, Ankarklev J, Reiner DS, et al. Draft genome sequencing of giardia intestinalis assemblage B isolate GS: is human giardiasis caused by two different species? PLoS Pathog. 2009 Aug;5(8):e1000560.

34. Jerlström-Hultqvist J, Franzén O, Ankarklev J, Xu F, Nohýnková E, Andersson JO, et al. Genome analysis and comparative genomics of a Giardia intestinalis assemblage E isolate. BMC Genomics. 2010;11(1):543.

35. Xu F, Jerlström-Hultqvist J, Einarsson E, Ástvaldsson Á, Svärd SG, Andersson JO. The Genome of Spironucleus salmonicida Highlights a Fish Pathogen Adapted to Fluctuating Environments. PLoS Genetics. 2014;10(2).

36. Campbell SR, van Keulen H, Erlandsen SL, Senturia JB, Jarroll EL. Giardia sp.: Comparison of electrophoretic karyotypes. Experimental Parasitology. 1990 Nov;71(4):470–82.

37. Prabhu A, Morrison HG, Martinez CR, Adam RD. Characterisation of the subtelomeric regions of Giardia lamblia genome isolate WBC6. Int J Parasitol. 2007 Apr;37(5):503–13.

38. Xu F, Jerlström-Hultqvist J, Einarsson E, Ástvaldsson Á, Svärd SG, Andersson JO. The Genome of Spironucleus salmonicida Highlights a Fish Pathogen Adapted to Fluctuating Environments. PLOS Genetics. 2014 Feb;10(2):e1004053.

39. Franzén O, Jerlström-Hultqvist J, Einarsson E, Ankarklev J, Ferella M, Andersson B, et al. Transcriptome Profiling of Giardia intestinalis Using Strand-specific RNA-Seq. Tucker-Kellogg G, editor. PLoS Computational Biology. 2013 Mar 28;9(3):e1003000.

40. Einarsson E, Troell K, Hoeppner MP, Grabherr M, Ribacke U, Svärd SG. Coordinated Changes in Gene Expression Throughout Encystation of Giardia intestinalis. Hehl AB, editor. PLOS Neglected Tropical Diseases. 2016 Mar 25;10(3):e0004571.

41. Morf L, Spycher C, Rehrauer H, Fournier CA, Morrison HG, Hehl AB. The transcriptional response to encystation stimuli in Giardia lamblia is restricted to a small set of genes. Eukaryotic Cell. 2010 Oct;9(10):1566–76.

42. Barash NR, Nosala C, Pham JK, McInally SG, Gourguechon S, McCarthy-Sinclair B, et al. Giardia Colonizes and Encysts in High-Density Foci in the Murine Small Intestine. mSphere. 2017 Jun;2(3).

43. Pham JK, Nosala C, Scott EY, Nguyen KF, Hagen KD, Starcevich HN, et al. Transcriptomic Profiling of High-Density Giardia Foci Encysting in the Murine Proximal Intestine. Frontiers in Cellular and Infection Microbiology [Internet]. 2017 May 31 [cited 2019 May 27];7. Available from: http://journal.frontiersin.org/article/10.3389/fcimb.2017.00227/full

44. Hudson AJ, Moore AN, Elniski D, Joseph J, Yee J, Russell AG. Evolutionarily divergent spliceosomal snRNAs and a conserved non-coding RNA processing motif in Giardia lamblia. Nucleic Acids Research. 2012 Nov;40(21):10995–1008.

45. Kamikawa R, Inagaki Y, Tokoro M, Roger AJ, Hashimoto T. Split Introns in the Genome of Giardia intestinalis Are Excised by Spliceosome-Mediated trans-Splicing. Current Biology. 2011 Feb;21(4):311–5.

46. William Roy S. Transcriptomic analysis of diplomonad parasites reveals a trans-spliced intron in a helicase gene in Giardia. PeerJ. 2017 Jan 5;5:e2861.

47. Touz MC, Conrad JT, Nash TE. A novel palmitoyl acyl transferase controls surface protein palmitoylation and cytotoxicity in Giardia lamblia VSP palmitoylation. Molecular Microbiology. 2005 Nov;58(4):999–1011.

48. Touz MC, Ropolo AS, Rivero MR, Vranych CV, Conrad JT, Svard SG, et al. Arginine deiminase has multiple regulatory roles in the biology of Giardia lamblia. Journal of Cell Science. 2008 Sep 1;121(17):2930–8.

49. Davids BJ, Reiner DS, Birkeland SR, Preheim SP, Cipriano MJ, McArthur AG, et al. A new family of giardial cysteine-rich non-VSP protein genes and a novel cyst protein. PLoS ONE. 2006 Dec 20;1:e44.

50. Ringqvist E, Avesson L, Söderbom F, Svärd SG. Transcriptional changes in Giardia during host-parasite interactions. Int J Parasitol. 2011 Mar;41(3–4):277–85.

51. Manning G, Reiner DS, Lauwaet T, Dacre M, Smith A, Zhai Y, et al. The minimal kinome of Giardia lamblia illuminates early kinase evolution and unique parasite biology. Genome Biology. 2011;12(7):R66.

52. Gargantini PR, Serradell M del C, Ríos DN, Tenaglia AH, Luján HD. Antigenic variation in the intestinal parasite Giardia lamblia. Current Opinion in Microbiology. 2016 Aug;32:52–8.

53. Palm JED, Weiland ME-L, Griffiths WJ, Ljungström I, Svärd SG. Identification of immunoreactive proteins during acute human giardiasis. J Infect Dis. 2003 Jun 15;187(12):1849–59.

54. Liu J, Svärd SG, Klotz C. Giardia intestinalis cystatin is a potent inhibitor of papain, parasite cysteine proteases and, to a lesser extent, human cathepsin B. FEBS Letters. 2019 Jun;593(12):1313–25.

55. DuBois KN, Abodeely M, Sakanari J, Craik CS, Lee M, McKerrow JH, et al. Identification of the major cysteine protease of Giardia and its role in encystation. J Biol Chem. 2008 Jun 27;283(26):18024–31.

56. Ansell BRE, McConville MJ, Baker L, Korhonen PK, Young ND, Hall RS, et al. Time-Dependent Transcriptional Changes in Axenic Giardia duodenalis Trophozoites. PLoS Neglected Tropical Diseases. 2015;9(12):1–24.

57. Stadelmann B, Merino MC, Persson L, Svärd SG. Arginine Consumption by the Intestinal Parasite Giardia intestinalis Reduces Proliferation of Intestinal Epithelial Cells. Zilberstein D, editor. PLoS ONE. 2012 Sep 19;7(9):e45325.

58. Mastronicola D, Testa F, Forte E, Bordi E, Pucillo LP, Sarti P, et al. Flavohemoglobin and nitric oxide detoxification in the human protozoan parasite Giardia intestinalis. Biochem Biophys Res Commun. 2010 Sep 3;399(4):654–8.

59. Das P, Lahiri A, Lahiri A, Chakravortty D. Modulation of the arginase pathway in the context of microbial pathogenesis: A metabolic enzyme moonlighting as an immune modulator. PLoS Pathogens. 2010;6(6).

60. Krasity BC, Troll JV, Weiss JP, McFall-Ngai MJ. LBP/BPI proteins and their relatives: conservation over evolution and roles in mutualism. Biochemical Society Transactions. 2011 Aug 1;39(4):1039–44.

61. Ankarklev J, Franzén O, Peirasmaki D, Jerlström-Hultqvist J, Lebbad M, Andersson J, et al. Comparative genomic analyses of freshly isolated Giardia intestinalis assemblage A isolates. BMC Genomics [Internet]. 2015 Dec [cited 2018 Apr 29];16(1). Available from: http://www.biomedcentral.com/1471-2164/16/697

62. Lee J-H, Wood TK, Lee J. Roles of Indole as an Interspecies and Interkingdom Signaling Molecule. Trends in Microbiology. 2015 Nov;23(11):707–18.

63. Russell AB, Singh P, Brittnacher M, Bui NK, Hood RD, Carl MA, et al. A Widespread Bacterial Type VI Secretion Effector Superfamily Identified Using a Heuristic Approach. Cell Host & Microbe. 2012 May;11(5):538–49.

64. Sana TG, Flaugnatti N, Lugo KA, Lam LH, Jacobson A, Baylot V, et al. Salmonella Typhimurium utilizes a T6SS-mediated antibacterial weapon to establish in the host gut. Proceedings of the National Academy of Sciences. 2016 Aug 23;113(34):E5044–51.

65. Kim MH, Choi W-C, Kang HO, Lee JS, Kang BS, Kim K-J, et al. The molecular structure and catalytic mechanism of a quorum-quenching N-acyl-L-homoserine lactone hydrolase. Proceedings of the National Academy of Sciences. 2005;102(49):17606–11.

66. Leckenby A, Hall N. Genomic changes during evolution of animal parasitism in eukaryotes. Current Opinion in Genetics & Development. 2015 Dec;35:86–92.

67. Peyretaillade E, Boucher D, Parisot N, Gasc C, Butler R, Pombert J-F, et al. Exploiting the architecture and the features of the microsporidian genomes to investigate diversity and impact of these parasites on ecosystems. Heredity (Edinb). 2015 May;114(5):441–9.

68. Singer SM, Nash TE. The Role of Normal Flora in *Giardia lamblia* Infections in Mice. The Journal of Infectious Diseases. 2000;181(4):1510–2.

69. Singer SM, Nash TE. T-Cell-Dependent Control of Acute Giardia lamblia Infections in Mice. Infection and Immunity. 2000 Jan 1;68(1):170–5.

70. Ropolo AS, Saura A, Carranza PG, Lujan HD. Identification of Variant-Specific Surface Proteins in Giardia muris Trophozoites. Infection and Immunity. 2005 Aug 1;73(8):5208–11.

71. Adam RD, Nigam A, Seshadri V, Martens CA, Farneth GA, Morrison HG, et al. The Giardia lamblia vsp gene repertoire: characteristics, genomic organization, and evolution. BMC Genomics. 2010;11(1):424.

72. Schmid-Siegert E, Richard S, Luraschi A, Mühlethaler K, Pagni M, Hauser PM. Mechanisms of Surface Antigenic Variation in the Human Pathogenic Fungus Pneumocystis jirovecii. Weiss LM, editor. mBio [Internet]. 2017 Nov 7 [cited 2019 May 28];8(6). Available from: http://mbio.asm.org/lookup/doi/10.1128/mBio.01470-17

73. Mugnier MR, Stebbins CE, Papavasiliou FN. Masters of Disguise: Antigenic Variation and the VSG Coat in Trypanosoma brucei. PLoS Pathog. 2016;12(9):e1005784.

74. Husnik F, McCutcheon JP. Functional horizontal gene transfer from bacteria to eukaryotes. Nature Reviews Microbiology. 2018 Feb;16(2):67–79.

75. Alsmark C, Foster PG, Sicheritz-Ponten T, Nakjang S, Martin Embley T, Hirt RP. Patterns of prokaryotic lateral gene transfers affecting parasitic microbial eukaryotes. Genome Biology. 2013;14(2):R19.

76. Liu J, Fu Z, Hellman L, Svärd SG. Cleavage specificity of recombinant Giardia intestinalis cysteine proteases: Degradation of immunoglobulins and defensins. Molecular and Biochemical Parasitology. 2019 Jan;227:29–38.

77. Feely DE, Gardner MD, Hardin EL. Excystation of Giardia muris induced by a phosphate-bicarbonate medium: localization of acid phosphatase. J Parasitol. 1991 Jun;77(3):441–8.

78. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. Journal of Molecular Biology. 2016;428(4):726–31.

79. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Research. 2017 Jan 4;45(D1):D353–61.

80. Karp PD, Latendresse M, Paley SM, Krummenacker M, Ong QD, Billington R, et al. Pathway Tools version 19.0 update: software for pathway/genome informatics and systems biology. Briefings in Bioinformatics. 2016 Sep 1;17(5):877–90.

81. Aurrecoechea C, Brestelli J, Brunk BP, Carlton JM, Dommer J, Fischer S, et al. GiardiaDB and TrichDB: integrated genomic resources for the eukaryotic protist pathogens Giardia lamblia and Trichomonas vaginalis. Nucleic Acids Research. 2009 Jan 1;37(suppl_1):D526–30.

82. Green ML, Karp PD. A Bayesian method for identifying missing enzymes in predicted metabolic pathway databases. BMC Bioinformatics. 2004 Jun 9;5:76.

83. Lee TJ, Paulsen I, Karp P. Annotation-based inference of transporter function. Bioinformatics. 2008 Jul 1;24(13):i259-267.

84. Marchler-Bauer A, Bryant SH. CD-Search: protein domain annotations on the fly. Nucleic Acids Res. 2004 Jul 1;32(Web Server issue):W327-331.

85. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic acids research. 2002;30(14):3059–66.

86. Do CB, Mahabhashyam MSP, Brudno M, Batzoglou S. ProbCons: Probabilistic consistency-based multiple sequence alignment. Genome research. 2005;15(2):330–40.

87. Notredame C, Higgins DG, Heringa J. T-Coffee: A novel method for fast and accurate multiple sequence alignment. J Mol Biol. 2000;302(1):205–17.

88. Criscuolo A, Gribaldo S. BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. BMC Evol Biol. 2010 Jul 13;10:210.

89. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Molecular Biology and Evolution. 2015 Jan 1;32(1):268–74.

90. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods. 2017 May 8;14:587.

91. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution. 2018 Feb 1;35(2):518–22.

92. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Systematic Biology. 2010 May 1;59(3):307–21.

93. Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize implements and enhances circular visualization in R. Bioinformatics. 2014 Oct;30(19):2811–2.

94. Guy L, Kultima JR, Andersson SGE. genoPlotR: comparative gene and genome visualization in R. Bioinformatics. 2010 Sep 15;26(18):2334–5.

95. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile and open software for comparing large genomes. Genome Biol. 2004;5(2):R12.

Chromosomes
GC%
Unique genes
ARPs/NEKs
(ψ)VSPs/HCMPs
Coding%/
5kbp
#SNPs/
kbp

100
0

1
0
37
0

ARPs
NEKs
VSPs
ψVSPs
HCMPs

Coding%<0.5
#SNPs>=5

Diplomonad adaptations
Metabolic adaptation : gene loss / gain via HGT
Cysteine-rich gene families
Heavy intron loss

S. salmonicida

G. intestinalis

G. muris

B.

C.

D.

Giardia **adaptations**
Adhesive disc
VSPs and HCMPs
Mitosomes
Metabolic adaptation : gene loss / gain via HGT

Giardia muris **adaptation**
Further intron loss
Loss of retrotransposons
Genome stream-lining
Microbiome interaction: HGT
Metabolic adaptation : gene loss / gain via HGT

A.

Ancestral Fornicate characters
Cyst formation
Hydrogenosome

K. bialata