Research Article

Full Title:

Laboratory testing of very low-copper-treated water to prolong pupation and

emerging time of mosquito larvae: an alternative method to delay mosquito

breeding capability

Short Title:

Effects of exposure to very low concentrations of copper on mosquito larvae

Authors:

Mohamad Reza [a*], Cimi Ilmiawati [b]

[a] *Department of Biology, Faculty of Medicine, Andalas Unversity, Padang, West*

Sumatra, Indonesia

[b] *Department of Pharmacology, Faculty of Medicine, Andalas University, Padang,*

West Sumatra, Indonesia

*Corresponding Author:

Mohamad Reza, MD, PhD

Department of Biology

Faculty of Medicine, Andalas University

Main Campus at Limau Manis, Gedung C Lantai 2

Pauh, Padang, 25166

West Sumatra, Indonesia

e-mail: reza@med.unand.ac.id

ABSTRACT

Introduction: Larvicide application in ovitrap is one of the currently available methods used in mosquito eradication campaign because they eliminate the larval stage. We previously reported that copper in liquid form is a promising candidate due to its potent larvicide properties in a laboratory setting and even in the field. In the field study, several larvae survived in the ovitrap due to the dilution of copper concentration by the rain. The surviving larvae were smaller and less motile. This led our interest to study the effect of a sub-lethal dose of copper in ovitrap on larvae development, pupation time and lifespan in the adult stage.

Methods: First instar larvae of three species of mosquito (*Aedes albopictus*, *Culex pipiens* and *Anopheles stephensi*) were bred in copper-treated water at a concentration of 0.60 ppm, 0.30 ppm, and 0.15 ppm and compared with the control group. The surviving larvae were recorded every day in terms of pupa emerging time and adult emerging time. The number of adult mortality was recorded and compared with the control.

Results: Copper showed potent larvicide effect in the 0.60 ppm concentration and prolonged pupation time and caused a significantly lower number of emerging mosquitoes down to the lowest concentration of 0.15 ppm. The adult lifespan was not different compared to the control.

Conclusion: This study demonstrates the capability of copper below 1 ppm to prolong the pupation time and the emerging time of mosquito larvae. Our findings open the possibility of copper application to cut mosquito breeding capacity that eventually will reduce disease transmission.

Keywords: *Aedes albopictus*, *Anopheles stephensi*, *Culex pipiens*, copper, larval control, ovitrap, pupation time

INTRODUCTION

Mosquitoes are causing millions of deaths every year worldwide. More than half of the world's populations live in areas where mosquitoes are present as vectors and transmit malaria, dengue, Zika, Chikungunya, yellow fever and other diseases [1]. Mosquito-borne diseases are causing an enormous burden on the public health system in many countries with limited financial and human resources. Despite being considered as the most conceivable method to eradicate mosquito-borne diseases [1], global funding for malaria vector control is far below the budget endowed for exploring cures and vaccines for these diseases (US$ 56 *vs* 408 million in 2018) [2]. Lack of funding is considered as one of the challenges in achieving and sustaining malaria control [3].

Core methods for vector control, i.e. insecticide-treated nets (ITNs) and indoor residual spraying (IRS), rely on insecticides. Unfortunately, there is an increase in mosquito resistance to insecticides [4]. Therefore, there is a need to develop novel methods for vector control, particularly practical and economical ones, to be combined with the available methods in integrated vector management. Larval source management (LSM) is part of integrated vector control using strategies to modify or manipulate water bodies as the potential larval habitats of mosquitoes to prevent maturation of mosquito developmental stages. LSM strategies also include the introduction of larvicide (chemical and biological agents) and larvivorous fish (natural predator) into larval habitats [5-7]. Previous studies have proposed the utility of using metallic [8, 9] and liquid [5, 10, 11]

copper as a potential and affordable larvicide [7]. Copper is effective as a

mosquito larvicide at a concentration below 2 ppm [9], the threshold value deemed

safe for human drinking water [12], thus making copper a potential candidate for

use in the public health setting.

Our previous findings on the properties of copper to kill mosquito larvae

[10] and laboratory testing on the effect of liquid copper at a concentration of 10

ppm on three species of mosquito larvae [11] prompted us to perform a field test

in a malaria and dengue-endemic area of West Sumatra, Indonesia. Despite

confirming the larvicide effect of copper [7], we observed that dilution by

rainwater should be considered as confounding in the field since we noted that

some larvae survived in outdoor ovitraps. However, the survived larvae appeared

to be less motile and were smaller in size compared to unexposed larvae.

Consequently, further study is required to determine whether the survived larvae

capable to continue their life cycle until adulthood. Therefore, we performed a

laboratory test of sub-lethal concentrations of copper exposure to mosquito larvae

to verify its effect on mosquito development and breeding capacity.

MATERIALS AND METHODS

Ethical Approval

This research was approved by the ethics committee of the Faculty of

Medicine, Universitas Andalas (Approval No.036/KEP/FK/2019).

Mosquito Eggs and Mice

Anopheles stephensi (strain SDA 500), were reared in our laboratory under

room temperature (26°C), 50-70% relative humidity, and a 13:11 h light: dark

cycle. From the day of emergence, mosquitoes were provided with a 5% fructose

solution soaked in a filter paper. Females five days after emergence were allowed

to feed on anesthetized mice. Three days later, an oviposition dish was placed in a

cage containing gravid females. The eggs were laid on a filter paper soaking in

the dish. The filter paper with eggs was placed on a 12 x 20 cm hatching tray

containing 500 ml of water. After hatching, 3-5 mg of carp food per tray was

sprinkled on the surface of the water twice daily. Twelve days later, pupae were

collected daily and transferred to cages for adult emergence.

Eggs of *Aedes albopictus* and *Culex pipiens* were purchased and the first

instar newly hatched larvae were utilized in the experiment. Female BALB/c mice

were purchased from SLC (Shizuoka, Japan). The mice were fed *ad libitum* and

exposed to a 13:11 h of light: dark cycle. This mosquito-mouse cycle was used to

maintain mosquitoes in the laboratory.

Preparation of Copper Sulfate Solutions

Four concentrations of $CuSO_4$ solution were used for the experiment. The

copper concentrations were 0.60, 0.30, 0.15, and 0 ppm. Each solution was

allocated to a pre-washed square plastic plate and covered with a transparent

plastic plate. First, we prepared a standard solution of 100 mM of $CuSO_4$ by

mixing copper sulfate powder with filtered water (Fine Ceramic Filter NGK

insulators, Nagoya, Japan). Then, we prepared the four concentrations for the

experiment by diluting them from the standard solution and confirming the copper

level using a copper measuring device (Hanna Instruments, Tokyo, Japan) and Z-

5010 Polarized Zeeman flame atomic absorption spectrophotometer (Hitachi Ltd,

Tokyo, Japan).

Preparation of Mosquito Larvae

First instar newly hatched larvae were used in this experiment. Fifty larvae were separated and put inside each container. Three containers for each concentration of copper were prepared. Larvae were allowed to hatch and were observed every 24-hours under the microscope for mortality. A fine ceramic filter (NGK Insulators, Nagoya, Japan) was used to remove chlorine ions from tap water for the experiment.

RESULTS

To determine the effect of very low concentrations of copper on mosquito larval mortality, larvae of *A. albopictus*, *A. stephensi* and *C. pipiens* were exposed to 0.60, 0.30 and 0.15 ppm of copper solutions ($CuSO_4$) and were observed daily. Larvae of *A. stephensi* and *C. pipiens* exposed to 0.60 ppm of copper showed 100% mortality within a week, while a small number of *A. albopictus* larvae survived. Larvae from three species roughly showed 50% mortality when exposed to 0.30 ppm of copper for seven days. $CuSO_4$ at 0.15 ppm only had a statistically significant effect on *C. pipiens* larval mortality compared to control, started on the third day of exposure (**Figure 1**).

Figure 1. The observed mortality of *A. albopictus*, *A. stephensi*, and *C. pipiens* larvae on exposure to 0.15, 0.30, and 0.60 ppm of $CuSO_4$. *significantly different from control, $p < 0.05$ (t-test or Mann-Whitney U test)

To figure out the effect of very low concentrations of copper exposure to

larval pupation time, the number of emerging pupae from the three species were

counted. Pupae started to emerge at day eight to 10, depending on the mosquito

species. Surviving larvae exposed to 0.15 and 0.30 ppm of copper showed

prolonged pupation time in all species observed compared to the control group.

We also observed that the surviving larvae of *Ae. albopictus* on 0.60 ppm were not

turned into pupae (**Figure 2**).

Figure 2. The number of emerging pupae of *A. albopictus*, *A. stephensi*, and *C.*

***pipiens* on exposure to 0.15, 0.30, and 0.60 ppm of CuSO$_4$. *significantly**

different from control, p<0.05 (Mann-Whitney U test)

To determine the effect of very low concentrations of copper exposure on

mosquito ability to reach adulthood, we observed and counted the number of

emerging adults in three exposed species. Surviving pupae of all observed species

exposed to 0.15 ppm of CuSO$_4$ become adult mosquitoes, albeit at a statistically

significantly lower number compared to the control group. Adult mosquito of all

observed species emerged at later days compared to control and survived up to 22

days of observation. Almost none of the pupae of all observed species exposed to

0.30 ppm of CuSO$_4$ emerged as an adult mosquito (**Figure 3**).

Figure 3. The number of emerging *A. albopictus*, *A. stephensi*, and *C. pipiens*

mosquito on exposure to 0.15, 0.30, and 0.60 ppm of CuSO$_4$. *significantly

different from control, p<0.05 (Mann-Whitney U test)

It is noteworthy that all adults emerged in copper-treated groups and

control appeared to be viable and showed no difference in term of their lifespan. It

seemed that the copper effect on the survived larvae was limited to prolonging the

emergence of pupae and was not carried on to the adult stage.

DISCUSSION

The results of the present study demonstrate that copper adequately lead to

a high rate of larval mortality in at a concentration below 1 ppm (0.60 ppm). At a

further lower concentration (0.30 ppm), copper still kill half of the larvae, and

together with the lowest concentration of 0.15 ppm, prolongs pupation time and

the emerging time of adult mosquitoes compared to the control group. The number

of mosquitoes that emerged afterward is also significantly lower than those in the

control group.

However, the copper effect is limited in the larvae and pupae stage. Copper

shows no effect on the adult stage. The effect on the larvae probably caused by the

copper effect to commensal bacteria in the larval midgut, which disrupts their

intestinal function [13]. This prevents them to eat properly and to collect enough

energy for pupation and emerge as an adult mosquito. Once the larvae become

adult, this copper effect no longer plays a role in their life cycle.

This is an interesting finding, since at a very low concentration copper

prolongs breeding time of mosquito. This might lead to the possibility to utilize

copper in a tap water to jeopardize mosquitoes breeding capability.   The

mosquitoes will need a longer time to breed, which then reduce their capability to

transmit mosquito-borne diseases such as dengue hemorrhagic fever.

It is also known that *Aedes sp.* are resistant to insecticide in indoor or

outdoor spraying. A study conducted in Sri Lanka showed *Aedes aegypti* and

Aedes albopictus were highly resistant to DDT and they can oviposit indoors and

outdoors [14]. Another study in Bangladesh reported abundant potential larval

habitats for *Aedes sp.* in containers or jars spread around the city [15]. These

situations need another approach of vector control, such as laying ovitraps or

another effort to cut or delay mosquito breeding capability, or even provide

insecticide in the tap water if possible.

The US EPA is suggesting 1.3 ppm as a safe limit of copper concentration

in the drinking water [16], thus making the application of very low 0.15 ppm of

copper might be logically acceptable in the future.

We believed that, together with the application of copper in the ovitraps,

the concentration of 0.15 ppm of copper in the tap water in houses and large scale

water reservoir might be a potential strategy in cutting mosquito breeding

capability and their ability to transmit mosquito-borne diseases, especially dengue

hemorrhagic fever, which mainly breed in the city water reservoir and container.

This result might be a chance to re-evaluate the allowance of copper in drinking

water as one of the possible strategies in eradicating mosquito-borne diseases in

the future.

Authors` contributions: MR conceived and designed the study protocol. MR and

CI formulated the study proposal and obtained funding. MR conducted research in

the laboratory. MR carried out data analysis and interpretation and wrote the

manuscript, while CI conducted the statistical analysis. All authors read and

approved the final manuscript.

Acknowledgment: The authors thank Prof. Hirotomo Kato, Ph.D for his kind

permit to use his laboratory in Japan for this research.

Conflict of interest: The authors report no conflicts of interest.

References

1. WHO.   Mosquito-borne   diseases.   2019.   Available   from:

https://www.who.int/neglected_diseases/vector_ecology/mosquito-

borne-diseases/en/

2. WHO. Global malaria report 2019. Geneva: World Health Organization;

2019.

3. Bastiaens GJH, Bousema T, Leslie T. Scale-up of malaria rapid

diagnostic tests and artemisinin-based combination therapy: challenges

and perspectives in Sub-Saharan Africa. PLoS Med. 2014;11(1):

e1001590. doi:10.1371/journal.pmed.1001590.

4. Liu N. Insecticide resistance in mosquitoes: impact, mechanisms, and

research directions. Annu Rev Entomol. 2015; 60:537-559.

5. Reza M, Yamamoto DS, Matsuoka H. Laboratory testing of larvivorous

fish Japanese medaka (Oryzias latipes) predation ability to copper-

treated *Anopheles stephensi* larvae: an alternative method for vector

control under low concentration of copper. Med Entomol Zool.

2013;64(2):67-71.

6. WHO. Larval source management: a supplementary measure for malaria

vector control: an operational manual. Geneva: World Health

Organization; 2013.

7. Reza M, Ilmiawati C, Matsuoka H. Application of copper-based ovitraps

in local houses in West Sumatera, Indonesia: a field test of a simple and

affordable larvicide for mosquito control. Trop Med Health. 2016;44:11.

doi:10.1186%2Fs41182-016-0007-8.

8. Romi R, Di Luca M, Raineri W, Pesce M, Rey A, Giovannangeli S,

Zanasi F, Bella A. Laboratory and field evaluation of metallic copper on

Aedes albopictus (Diptera: Culicidae) larval development. J Med

Entomol. 2000; 37(2):281-285.

9. Becker N, Oo TT, Schork N. Metallic copper spray – a new control

technique to combat invasive container-inhabiting mosquitoes. Parasites

Vectors. 2015;8:575. doi:10.1186/s13071-015-1180-z.

10. Reza M, Yamamoto DS, Matsuoka H. Low concentration copper

solution jeopardizes larval movement and ability to survive predation:

new insight into malaria eradication via vector control. Med Entomol

Zool. 2012;63:217-222.

11. Reza M, Yamamoto DS, Matsuoka H. Larvicidal and ovipositional

preference test of copper solution for mosquitos. Med Entomol Zool.

2014; 65(3):147-150.

12. WHO. Trace elements in human nutrition and health. Geneva: World

Health Organization; 1996.

13. Beaty BJ, Mackie RS, Mattingly KS, Carlson JO, Rayms-Keller A. The

midgut epithelium of aquatic arthropods: a critical target organ in

environmental toxicology. Environ Health Perspect.

2002;110(suppl6):911-914.

14. Dharsini S, Vinobaba M, Jude PJ et al. Prevalence and insecticide

susceptibility of dengue vectors in the district of Batticaloa in Eastern Sri

Lanka. Trop Med Health. 2011;39:47-52.

15. Ferdousi F, Yoshimatsu S, Ma E et al. Identification of essential

containers for *Aedes* larval breeding to control dengue in Dhaka,

Bangladesh. Trop Med Health. 2015;43(4):253-264.

16. US EPA. Drinking Water Requirements for States and Public Water

Systems: Lead and Copper Rule. Available from:

https://www.epa.gov/dwreginfo/lead-and-copper-rule

Figure 1. The observed mortality of *A. albopictus*, *A. stephensi*, and *C. pipiens* larvae on exposure to 0.15, 0.30, and 0.60 ppm of CuSO$_4$. *significantly different from control, p<0.05 (t-test or Mann-Whitney U test)

Figure 2. The number of emerging pupae of *A. albopictus*, *A. stephensi*, and *C. pipiens* on exposure to 0.15, 0.30, and 0.60 ppm of $CuSO_4$. *significantly different from control, $p < 0.05$ (Mann-Whitney U test)

Figure 3. The number of emerging *A. albopictus*, *A. stephensi*, and *C. pipiens* mosquito on exposure to 0.15, 0.30, and 0.60 ppm of CuSO₄. *significantly different from control, p<0.05 (Mann-Whitney U test)