TITLE

Comparative proximity biotinylation implicates RAB18 in cholesterol mobilization and

biosynthesis

RUNNING TITLE

GEF-dependent RAB18 interactions

AUTHORS

Robert S. Kiss*[1], Jarred Chicoine[1], Youssef Khalil[2], Robert Sladek[1], He Chen[1],

Alessandro Pisaturo[1], Cyril Martin[1], Jessica D. Dale[3], Tegan A. Brudenell[3], Archith

Kamath[4,5], Emanuele Paci[6], Peter Clayton[2], Jimi C. Wills[4], Alex von Kriegsheim[4],

Tommy Nilsson[1], Eamonn Sheridan[3], Mark T. Handley*[3,7]

AFFILIATIONS

[1]Research Institute of the McGill University Health Centre

1001 boul Decarie

Glen Site Block E

Montreal, QC

H4A 3J1

Canada

[2] Genetics and Genomic Medicine

Great Ormond Street Institute of Child Health

University College London

30 Guilford Street

London

WC1N 1EH

United Kingdom

[3]Leeds Institute of Medical Research

St James's University Hospital

Leeds

LS9 7TF

United Kingdom

[4]Cancer Research UK Edinburgh Centre

MRC Institute of Genetics & Molecular Medicine

The University of Edinburgh

Western General Hospital

Edinburgh

EH4 2XR

United Kingdom

[5]Medical Sciences Division

University of Oxford

Oxford

OX3 9DU

United Kingdom

[6]Astbury Centre for Structural Molecular Biology

University of Leeds

Leeds

LS2 9JT

United Kingdom

[7]Faculty of Biological Sciences

University of Leeds

Leeds

LS2 9JT

United Kingdom.

*authors contributed equally

CORRESPONDING AUTHORS

Robert S. Kiss: robert.kiss@mcgill.ca

Mark T. Handley: m.handley@leeds.ac.uk

KEYWORDS

SUMMARY STATEMENT

We used proximity biotinylation together with guanine nucleotide exchange factor

(GEF)-null cell lines to discriminate functional RAB18-interactions. This approach

revealed that RAB18 mediates lathosterol mobilization and cholesterol biosynthesis.

ABSTRACT

Loss of functional RAB18 causes the autosomal recessive condition Warburg Micro

syndrome. To better understand this disease, we used proximity biotinylation in

HEK293 and HeLa cells to generate an inventory of potential RAB18 effectors. A

restricted set of 25 RAB18-interactions were regulated by the binary RAB3GAP1-

RAB3GAP2 RAB18-guanine nucleotide exchange factor (GEF) complex. These

included three groups of functionally interrelated proteins: a group of microtubule-

interacting/membrane shaping proteins; a group of proteins involved in membrane

tethering and docking; and a group of lipid-modifying/lipid transport proteins. We

provide confirmatory evidence for several of the interactors (SPG20/SPART,

SEC22A, TMCO4). Further we provide functional evidence that RAB18 links the Δ8-

Δ7 sterol isomerase emopamil binding protein (EBP) to a molecular machinery

mobilizing the products of EBP-catalysis. The cholesterol precursor lathosterol

accumulates in RAB18-null cells, and *de novo* cholesterol biosynthesis is reduced.

Our data demonstrate that GEF-dependent Rab-interactions are highly amenable to

interrogation by proximity biotinylation and suggest that Micro syndrome is a

cholesterol biosynthesis disorder.

INTRODUCTION

Rab Proteins are a large subfamily of small GTPases with discrete roles in
coordinating membrane trafficking (Zhen & Stenmark, 2015). They associate with
cellular membranes as a result of their C-terminal prenylation and like other small
GTPases, adopt different conformations and enter into different protein-protein
interactions according to whether they are GDP-, or GTP-bound. For Rab proteins,
cycles of GTP binding and hydrolysis are accompanied by cycles of membrane
association and dissociation that serve to promote the targeting of particular Rab
protein isoforms to particular membrane compartments. Although they possess
some intrinsic GTP-hydrolysis activity, their *in vivo* nucleotide-bound state is tightly
governed in cells by two classes of regulatory proteins. Guanine-nucleotide
exchange factors (GEFs) catalyse the exchange of bound GDP for GTP while
GTPase-activating proteins (GAPs) promote the hydrolysis of bound GTP to GDP
(Barr & Lambright, 2010, Lamber et al., 2019). The dissociation of Rab proteins from
membranes is mediated by GDP-dissociation inhibitor (GDI) proteins. GDIs
sequester GDP-bound Rabs in the cytosol, and are also involved in their
reassociation with membranes (Zhen & Stenmark, 2015).

Rab proteins have a variety of roles in the regulation of processes required to confer
compositional identity to membranous organelles and to subdomains within them.
These include membrane remodelling and the establishment of membrane contact
sites (Bui et al., 2010, Raiborg et al., 2015, Rocha et al., 2009, Sobajima et al.,
2018). Where exchange between organelles is mediated by carrier vesicles, these
processes are important in vesicle budding and transport, tethering at the
appropriate target membrane, and fusion (Cai et al., 2007). Under other
circumstances, membrane contact sites may be established to mediate direct fusion
between organelles or to facilitate the transfer of lipids and ions (Langemeyer et al.,
2018, Wickner, 2010, Wu et al., 2018). Rab proteins fulfil their roles by way of
protein-protein interactions with interacting partners termed 'effectors'. These
comprise an array of phylogenetically unrelated protein classes and can serve a
range of molecular functions. As such, they are most usually identified biochemically.

Biochemical identification of Rab effectors is challenging; Rab-effector interactions
are usually GTP-dependent and are often highly transient. Immunoprecipitation,
affinity purification and yeast-2-hybrid approaches have each been used but may be

more or less effective depending on the Rab isoform studied (Christoforidis et al.,

1999, Fukuda et al., 2008). One newer approach that has yielded identification of a

number of novel interactions is 'BioID' proximity biotinylation utilizing Rab proteins

fused to mutant forms of the promiscuous biotin ligase BirA*; the Rab fusion protein

biotinylates proximal proteins which are then purified on streptavidin and identified

through mass spectrometry (Gillingham et al., 2019, Liu et al., 2018, Roux et al.,

2012). Biotin labelling occurs in a relatively physiological context, and prospective

effectors can be purified under high stringency conditions. However, a drawback of

the technique is that it does not discriminate between close associations resulting

from functional protein-protein interactions and those resulting from overlapping

localizations.

RAB18 is a ubiquitously expressed ancestral Rab protein that localizes to the *cis*-

Golgi, endoplasmic reticulum (ER) and lipid droplets (LDs)(Gerondopoulos et al.,

2014, Handley et al., 2015, Martin et al., 2005, Ozeki et al., 2005). Previous work

has suggested that it functions in the regulation of lipolysis, and lipogenesis (Martin

et al., 2005, Ozeki et al., 2005, Pulido et al., 2011), trafficking between the Golgi and

endoplasmic reticulum (ER) (Dejgaard et al., 2008, Handley et al., 2015), ER

structure (Gerondopoulos et al., 2014), exocytosis (Vazquez-Martinez et al., 2007)

and autophagy (Bekbulat et al., 2018, Feldmann et al., 2017). In order to address

how RAB18 coordinates and performs its roles, we used BioID to generate an

inventory of its potential effectors. To discriminate functional interactions, we used

complementary comparative analyses in HEK293 and HeLa cells. Known RAB18

effectors were more strongly labelled by a fusion of BirA* and wild-type RAB18 than

by one incorporating an inactive form, RAB18(Ser22Asn), which is deficient in

nucleotide-binding. Similarly, BirA*-RAB18 labelled known effectors more strongly in

wild-type cells than in cells in which RAB18-GEF activity was disrupted with

CRISPR. Interestingly, disruption of different GEF complexes largely affected

different sets of RAB18 interactions.

Biallelic loss-of-function variants in *RAB18* or in the genes encoding the 'RAB3GAP'

RAB18-GEF complex cause the autosomal recessive condition Warburg Micro

syndrome (Aligianis et al., 2005, Bem et al., 2011, Borck et al., 2011, Handley &

Sheridan, 2018, Liegel et al., 2013)(MIMs 600118, 614222, 614225, 615663,

212720). We found that a restricted set of RAB18-interactions were RAB3GAP-

dependent, including known and novel effectors in three functional groups. We

present direct validation for several examples including the microtubule-binding

protein SPG20/SPART, the SNARE protein homologue SEC22A and an orphan

lipase TMCO4. Our data strongly support a previous suggestion (Xu et al., 2018) that

RAB18 effectors act collectively in lipid transfer between closely apposed

membranes. Further, we provide evidence that RAB18 serves to coordinate the

generation of the cholesterol precursor lathosterol by the EBP enzyme and its

subsequent mobilization by the lipid transfer protein (LTP) ORP2/OSBPL2. Because

of the clinical overlap between Micro syndrome and conditions including

lathosterolosis (MIM 607330) and Smith-Lemli-Opitz syndrome (SLOS; MIM

270400), our data suggest that impaired cholesterol biosynthesis may partly underlie

Micro syndrome pathology.

RESULTS

<u>An inventory of nucleotide-binding-dependent RAB18-associated proteins in HEK293</u>
<u>cells</u>

We first generated HEK293 cells stably expressing BirA*-tagged fusion proteins of
wild-type RAB18, nucleotide-binding deficient RAB18(Ser22Asn) or GTP-hydrolysis
deficient RAB18(Gln67Leu) using the Flp-In system (Figure S1). We then carried out
proximity-labelling, affinity purification and mass spectrometry of biotinylated proteins
as previously described (Roux et al., 2018, Roux et al., 2012). Following removal
from the dataset of previously identified non-specific binders from an in-house
database, a total of 98 proteins were identified as associating with RAB18 across all
samples (see Table S1).

The most comprehensive annotation of candidate RAB18 effectors thus far was
made in the 2014 paper by Gillingham et al., which utilized an affinity purification-
mass spectrometry (AP-MS) approach and the *Drosophila* RAB18 orthologue
(Gillingham et al., 2014). In that study, a total of 456 proteins were identified as
interacting with RAB18. However, only 14 of these were well represented in terms of
spectral counts, exhibited low non-specific binding to GST/Sepharose and showed
low binding to other Rab protein isoforms. We took these 14 proteins as the most
plausible physiological RAB18 interactors. Orthologues of 7 of these 14 proteins
were also represented in our BioID data. The number of spectral counts recorded for
these 7 proteins did not clearly distinguish them from the remainder of this BioID
dataset. We therefore proceeded by comparing their differential labelling by the
different RAB18 fusion proteins.

We began analyses by exploring the possibility that BirA*-RAB18(Gln67Leu) fusion
protein would produce enhanced biotinylation of RAB18-effectors (as compared to
BirA*-RAB18(WT)). We normalized total spectral counts between the BirA*-
RAB18(Gln67Leu) and BirA*-RAB18(WT) datasets and then calculated mutant:wild-
type ratios for each RAB18-associated protein (Table S1). Association ratios for
known RAB18-interactors ranged from 0.1-1.49 indicating that RAB18 associations
were altered by the Gln67Leu variant, but not predictably so.

We next compared the BirA*-RAB18(Ser22Asn) and BirA*-RAB18(WT) datasets

(Figure 1, Table S1). The RAB18-GEF subunits RAB3GAP1 and RAB3GAP2

showed association ratios >1 consistent with the high affinity of Rab-GEFs for

cognate Rabs in their nucleotide-free state. Since most effector-interactions are

GTP-dependent, we ranked other prospective RAB18-interactors according to those

that showed the lowest association ratios. Figure 1B shows the 28 of 98 proteins

with BirA*-RAB18:BirA*-RAB18(Ser22Asn) association ratios <0.5. These include

the remaining 5 common interactors from the Gillingham et al. study, several of

which have association ratios of zero, indicating that labelling by BirA*-

RAB18(Ser22Asn) was absent. These data suggest that comparison of labelling by

wild-type and nucleotide-binding-deficient BirA*-Rab fusion proteins can be an

effective means to distinguish putative effectors.

<u>An inventory of RAB18-GEF-dependent RAB18-associated proteins in HeLa cells</u>

We had previously used CRISPR to generate a panel of clonal, otherwise isogenic,

HeLa cell lines, null for RAB18 and a number of its regulators (see Figure S2).

Having shown that the BirA*-RAB18(WT):BirA*-RAB18(Ser22Asn) comparison in

HEK293 cells was informative, we carried out similar comparisons between BirA*-

labelling in wild-type and RAB18-GEF deficient HeLa cells (Figure 2A). Since GEF

activity promotes Rab GTP binding, and this is usually necessary for effector

interactions, these interactions will be attenuated in GEF-null cells. RAB3GAP1 and

RAB3GAP2 are each essential subunits of a binary RAB18-GEF complex whereas

TRAPPC9 is reported to be essential for the RAB18-GEF activity of a different GEF,

the multisubunit TRAPPII complex (Gerondopoulos et al., 2014, Li et al., 2017). We

therefore carried out proximity labelling using transient expression of the same

exogenous BirA*-RAB18 construct in wild-type cells and in RAB3GAP1-,

RAB3GAP2- and TRAPPC9-null cell lines.

Prior to mass-spec analysis, samples from each of the streptavidin pull-downs were

subjected to Western blotting to ensure comparable BirA*-RAB18 expression (Figure

S3A). Label-free quantitative proteomics analyses were used to calculate 'LFQ

intensities' (Cox et al., 2014) for each RAB18-associated protein, which were then

normalized in each experiment according to the quantity of RAB18 found in each

sample. Samples from three independent experiments were analysed, and pull-down
samples from untransfected biotin-treated cells were used as controls in each case.
After filtering the data to remove known mass-spec contaminants, and any protein
identified at a high level in control samples, a total of 584, 483 and 506 RAB18-
associated proteins were identified in each experiment. A total of 457 proteins were
present in two or more of the replicate experiments (see Table S2). Orthologues of
10 of the 14 putative RAB18-interacting proteins identified by Gillingham et al. were
identified in the HeLa cell BioID dataset including all 7 of those also identified in the
HEK293 cell dataset. However, as in that dataset, these were not distinguished by
their comparative abundance.

Different Rab-GEF complexes may operate in distinct subcellular localizations and
coordinate associations with different effectors (Carney et al., 2006). Therefore, we
assessed whether non-zero intensities for each RAB18-associated protein correlated
between samples (Figure 2B, Figure S3B). Very strong correlations between protein
intensities from RAB3GAP1- and RAB3GAP2-null cells indicated that loss of either
protein had a functionally equivalent effect (R^2=0.99, see Figure 2B). In contrast,
intensities from RAB3GAP1- and TRAPPC9-null cells were much more poorly
correlated (R^2=0.73, see Figure S3B). We therefore considered RAB3GAP- and
TRAPPC9-dependent RAB18-associations separately.

Of the 457 proteins identified in two or more independent experiments, only 25
showed an association ratio <0.5 in the absence of functional RAB3GAP (Figure 2C-
D, Table S2). These included orthologues of 9 of the 10 proteins identified in the
Gillingham et al. study. Thus, our approach was extremely powerful in discriminating
putative effector proteins. 133 proteins showed an association ratio <0.5 in the
absence of functional TRAPPC9 including the remaining common RAB18-interactor
from the Gillingham et al. study (see Table S2). There was only limited overlap
between RAB3GAP- and TRAPPC9-dependent associations (Figure 2C). Indeed,
among the 28 nucleotide-binding-dependent RAB18 associations identified in
HEK293 cells, 6 were also RAB3GAP-dependent, and 7 were TRAPPC9-dependent
in HeLa cells respectively, but none were both (Figure 1B). Among the 25
RAB3GAP-dependent associations in HeLa cells, only 5 were also TRAPPC9
dependent (Figure 2C, Table S2).

One of the TRAPPC9-dependent RAB18 associations was that with TBC1D5, a Tre-
2/Bub/Cdc16 (TBC) domain-containing RAB-GAP with a well characterised role in
regulation of RAB7 (Jia et al., 2016, Jimenez-Orgaz et al., 2018, Seaman et al.,
2009). This was a strong candidate as a RAB18 regulator or effector since it had
also been identified in the HEK293 dataset as well as in several previous studies
(Gillingham et al., 2019, Gillingham et al., 2014). We generated TBC1D5-null HeLa
cells, and first tested whether RAB7 and RAB18 dynamics were altered in these cells
using fluorescence recovery after photobleaching (FRAP). RAB7 dynamics were
substantially different in the TBC1D5-null cells as compared to those in wild-type
cells, consistent with reduced RAB7 GTP-hydrolysis resulting in its reduced GDI-
mediated exchange between membrane and cytosolic compartments (Figure S4A).
In contrast, RAB18 dynamics were unchanged in TBC1D5-null cells compared to
controls (Figure S4B). Further, RAB7 dynamics were unchanged in RAB18-null cells
compared to controls, indicating that RAB18 is not required for TBC1D5 activity
(Figure S4A). These data do not exclude TBC1D5 as a potential RAB18 effector, but
argue against a role for RAB18 in RAB7-regulation under resting conditions.

<u>Validation screening of RAB3GAP-dependent RAB18 associations</u>

Our continued study focused on the 25 RAB3GAP-dependent RAB18 associations
identified in HeLa cells on the basis that these included the majority of known RAB18
effectors together with a number of promising candidate effectors not previously
identified. Encouragingly, many of these also appeared to share interconnected
functions and fell into three main groups (Figure 2D). A group of proteins involved in
membrane shaping, cytoskeletal remodelling and in membrane-microtubule contacts
included SPG20, BICD2, REEP4, CAMSAP1 and FAM134B. Of these, the RAB18-
SPG20 interaction was previously validated (Gillingham et al., 2014), BICD2
interacts with RAB18 among a number of Rab isoforms (Gillingham et al., 2019,
Gillingham et al., 2014) and REEP4 was previously shown to interact with RAB3GAP
(Tinti et al., 2012). Although not previously linked to RAB18, CAMSAP1 and
FAM134B were each identified in both HEK293 and HeLa datasets. Next, a group of
proteins involved in establishing membrane contacts included components of the
NRZ/Dsl1 membrane-tethering complex, ZW10, RINT1 and NBAS, the Sec1/Munc18
(SM) protein SCFD2 and SNAP-REceptor (SNARE) proteins STX18 and BNIP1.

These proteins have been previously studied in the context of RAB18 (Gillingham et

al., 2019, Gillingham et al., 2014, Li et al., 2019, Xu et al., 2018, Zhao & Imperiale,

2017). Also in this group were the SNARE protein homologue SEC22A and the ER-

resident multispanning transmembrane protein WFS1, which regulates membrane

contacts between the ER and mitochondria (Angebault et al., 2018). These latter

proteins had not been linked to RAB18 previously. The third group of proteins

associated with RAB18 were a number of lipid transport, exchange and modifying

proteins, C2CD2L, C2CD2, ORP2/OSBPL2, INPP5B, EBP and TMCO4.

Interestingly, all but TMCO4 have related lipid species as their known substrates.

C2CD2 and C2CD2L are thought to mediate phosphatidylinositol (PI) transfer

between apposed membranes (Lees et al., 2017), ORP2 has recently been shown to

exchange PI(4,5)P_2 and cholesterol and INPP5B hydrolyses PI(4,5)P_2 to PI(4)P

(Wang et al., 2019). EBP is a Δ8-Δ7 Sterol Isomerase involved in cholesterol

biosynthesis (Silve et al., 1996). Various members of the OSBP family interact with

Rab proteins (Gillingham et al., 2019, Johansson et al., 2005), and a number of Rab

isoforms interact with INPP5B (Fukuda et al., 2008, Williams et al., 2007), though

associations with RAB18 have not been previously reported.

For initial validation of our HeLa dataset, we first carried out an additional

independent BioID experiment with wild-type and RAB3GAP1-null cells and

subjected the resulting samples to Western blotting for selected RAB18-associated

proteins (Figure 3A). As with the mass spectrometry, these proteins showed either

complete (RAB3GAP1, RAB3GAP2, ZW10) or partial (SPG20, STX18) dependence

on RAB3GAP for their RAB18 association.

Since RAB18 and its regulators are linked to Warburg Micro syndrome, we asked

whether the putative RAB18 effectors were linked to any diseases with overlapping

clinical features (Figure 3B). Micro syndrome is a clinically distinctive disorder

characterised by intellectual disability (ID), postnatal microcephaly, brain

malformations, ascending spastic paraplegia, neuropathy, hypogonadism and eye

abnormalities that include congenital bilateral cataracts, microphthalmia,

microcornea and optic atrophy (Handley & Sheridan, 2018). Among the RAB3GAP-

dependent RAB18 associations identified in the BioID screen, we noted that several

are encoded by disease-associated genes or their homologues. In common with

Micro syndrome, diseases linked to *SPG20*, *BICD2*, and the *REEP4*-homologues

REEP1 and *REEP2* are associated with ascending paraplegia (hereditary spastic
paraplegia/HSP). *FAM134B* is associated with sensory and autonomic neuropathy,
while the eye features of Micro syndrome overlap a different set of genes. *EBP* is
linked to microphthalmia, microcornea and cataracts as well as to brain
malformations like those in Micro syndrome. Both *WFS1* and *NBAS* have been
linked to conditions associated with optic atrophy, and both *WFS1* and the INPP5B
homologues OCRL1 and INPP5K are linked to conditions associated with congenital
cataracts.

Given the suggestive convergences in protein function and gene-disease-
associations, we proceeded to examine the subcellular localizations of 12 putative
effectors for which antibodies were available (Figure 3C-D). To determine whether
the localization of these proteins was altered in cells lacking RAB18, we analysed
wild type and RAB18-null lines in each case. In order to directly compare cells of
different genotypes under otherwise identical conditions, we labelled them with
CellTrace-Violet and CellTrace-Far Red reagents before seeding, immunostaining
and imaging them together. Since RAB18 can localize to LDs, we analysed both
untreated cells (Figure 3C) and cells loaded with oleic acid and labelled with
BODIPY-558/568-C12 (Figure 3D).

We observed a variety of staining patterns for the different putative effector proteins.
These ranged from staining that was enriched at the perinuclear region of cells, to
staining that appeared reticular, to staining that appeared more diffuse (Figure 3C).
Each staining pattern was compatible with the known localization of RAB18, which is
distributed between *cis*-Golgi, ER and cytosolic compartments (Handley et al., 2015).
Staining patterns for individual proteins were similar in the HeLa cells and also in
wild-type and RAB18-null RPE1 cells generated to provide biological replicates
(Figure S5). In lipid-loaded cells, we observed that the localizations of proteins with
reticular staining patterns overlapped with LDs, but they did not obviously shift to
adopt an LD localization. However, the two proteins that showed the most diffuse
staining patterns in untreated cells - ZW10 and SPG20 - appeared enriched in the
vicinity of LDs in lipid-loaded cells (Figure 3D).

We saw no evidence for dramatic changes in protein localizations in RAB18-null
cells as compared to their wild-type counterparts. Fluorescence intensities in RAB18-
null and wild-type cells were also generally similar, except in the case of staining for

SPG20, in which it appeared lower in RAB18-null HeLa cells than in wild-type cells

(Figure 3C).

<u>Levels of SPG20 are significantly reduced in RAB18-null and TBC1D20-null cells</u>

In a RAB18-null mouse model of Warburg Micro syndrome, peripheral nerves have a

disordered cytoskeleton, and there is a striking accumulation of microtubules at

motor nerve terminals (Carpanini et al., 2014). Therefore, interactions between

RAB18 and microtubule-binding proteins is of particular interest. To confirm the

reduction in levels of SPG20 we observed in RAB18-null HeLa cells, we used

quantitative fluorescence microscopy (Figure 4A-C). The SPG20 antibody used in

this study has previously been used for this purpose (Nicholson et al., 2015).

However, to confirm its specificity and also to determine the levels of non-specific

background produced in our experiments, we first analysed SPG20-null cells (Figure

4A-B). Measured fluorescence intensity of SPG20-null cells provided a baseline

level, above which fluorescence levels are proportional to levels of SPG20. In

RAB18-null cells, SPG20 fluorescence was reduced to 67.16±3.77% (s.e.m.,

$p<0.001$) of that in wild-type cells (Figure 4C).

To determine whether levels of SPG20 were altered by disrupted RAB18 regulation,

we next compared SPG20 fluorescence of wild type, RAB3GAP1-, RAB3GAP2- and

TBC1D20-null cells. Loss of the RAB18-GEF subunits RAB3GAP1 or RAB3GAP2

did not significantly affect levels of SPG20, whereas loss of the RAB18-GAP

TBC1D20 led to a reduction comparable to that in RAB18-null cells (57.48%±2.57

(s.e.m., $p<0.00005$). To rule-out the possibility that reduced SPG20 levels in RAB18-

and TBC1D20-null HeLa cells were the result of clonal variation, we analysed the

corresponding panel of RPE1 cell lines. However, because the RPE1 cells were less

amenable to comparative immunofluorescence experiments than HeLa cells, we

used LFQ analysis of whole cell lysates. As in the HeLa cells, levels of SPG20 were

significantly reduced in RAB18- and TBC1D20-null RPE1 cells compared to wild-

type controls ($p<0.05$ following FDR correction), but not in the other genotypes

tested (Figure 4D, Table S3).

RAB18, TBC1D20 and the RAB3GAP complex have all been linked to roles in

proteostasis and autophagy (Bekbulat et al., 2018, Feldmann et al., 2017, Sidjanin et

al., 2016, Spang et al., 2014). It was therefore possible that reduced SPG20 levels in

RAB18- and TBC1D20-null cells were the result of widespread dysregulation of

proteostasis. To assess this possibility, we compared LFQ data from wild-type and

TBC1D20-null RPE1 and HeLa cells (Tables S3 and S4). Following FDR correction,

only a small number of proteins showed significantly altered levels in each cell type

and there was limited overlap between cell types. This is in-line with a recent study

showing a compensatory mechanism maintains levels of basal autophagy when

RAB18 is absent or dysregulated (Bekbulat et al., 2018). Thus, dysregulation of

SPG20 levels most likely arises from a discrete mechanism.

Discrete changes in SPG20 levels in RAB18- and TBC1D20-null cells together with

the previous report of a RAB18-SPG20 interaction (Gillingham et al., 2014) provided

strong evidence for a functional relationship between these proteins. We therefore

carried out co-expression experiments to determine whether they colocalize in cells.

Coexpression of mCherry-RAB18 and mEmerald-SPG20 in HeLa cells showed that,

as previously reported, mCherry-RAB18 adopts a largely reticular localization

(Gerondopoulos et al., 2014) whereas mEmerald-SPG20 appears largely diffuse

(Eastman et al., 2009)(Figure 4E). We did not observe any clear colocalization

between the proteins or any relocalization of mEmerald-SPG20 in cells expressing

mCherry-RAB18. Since both RAB18 and SPG20 localize to LDs, we next explored

whether the localization of either protein to LDs was dependent on the other. In oleic

acid/BODIPY-558/568-C12-loaded HeLa cells, we found that mEmerald-SPG20

became concentrated around LDs, but that its localization was similar in both wild-

type and RAB18-null cells (Figure 4F, left panels). Similarly, EGFP-RAB18 was

enriched around LDs in both wild-type and SPG20-null cells (Figure 4F, right

panels).

Collectively, our findings suggest that RAB18 and SPG20 become localized to LDs

independently, and that any RAB18-SPG20 interaction is likely to be transient. The

reduced levels of SPG20 in RAB18- and TBC1D20-null cells seem likely to result

from its reduced stability, and it is intriguing that a transient interaction could

influence this. Nevertheless, this is consistent with the involvement of TBC1D20-

catalysed RAB18 GTP-hydrolysis and accompanying conformational change.

<u>SEC22A associates with RAB18 and its knockdown causes altered LD morphology</u>

The most studied group of RAB18 effector proteins to date are the tethering factors

ZW10, NBAS and RINT1, which together comprise the NRZ/Dsl1 complex

(Gillingham et al., 2014, Li et al., 2019, Xu et al., 2018, Zhao & Imperiale, 2017). The

NRZ complex regulates assembly of an ER SNARE complex containing STX18. The

canonical, fusogenic, form of this complex contains STX18, BNIP1, USE1 and

SEC22B (Spang, 2012, Tagaya et al., 2014). However, it has been proposed that

SEC22B is dispensable for the functions of RAB18, the NRZ complex, STX18,

BNIP1 and USE1 in regulating LDs (Xu et al., 2018). Rather than promoting

membrane fusion it is suggested that, in the absence of SEC22B, these proteins can

mediate the close apposition of membranes to facilitate lipid transfer.

In the HEK293 and HeLa BioID datasets, RAB18-associations with all three NRZ

components were nucleotide-binding-dependent and RAB3GAP-dependent

respectively (Figures 1B, 2D, Tables S1, S2). In the HeLa dataset, the ER SNARE

proteins STX18 and BNIP1 were also identified as associating with RAB18 in a

RAB3GAP-dependent manner (Figure 2D, Table S2). Interestingly, also among the

RAB3GAP-dependent RAB18 associations was SEC22A, a poorly studied

homologue of SEC22B.

SEC22A is one of two SEC22B homologues in humans, the other being SEC22C.

Like SEC22B, SEC22A and SEC22C possess N-terminal Longin domains and C-

terminal transmembrane (TM) domains. However, they lack the central coiled-coil

SNARE domain through which SEC22B mediates membrane fusion as part of the

STX18 complex. SEC22B localizes to the ER–Golgi intermediate compartment,

whereas different isoforms of SEC22C localize to the ER or *cis*-Golgi (Ge et al.,

2013, Yamamoto et al., 2017, Zhang et al., 1999). The localization of SEC22A had

not been addressed. In the absence of commercially available antibodies for

SEC22A, we examined its localization through expression of an mEmerald-SEC22A

fusion protein (Figure 5A). mEmerald-SEC22A produced a characteristic reticular

staining pattern and colocalized with an exogenous ER marker suggesting that

SEC22A localizes to the ER.

We next sought to compare the localization of SEC22A and RAB18 and to determine

whether they interact. However, coexpression of mEmerald-SEC22A and mCherry-

RAB18 appeared to disrupt normal ER morphology and to produce vesicular

structures and inclusions positive for both proteins (Figure S6). Although this was not

inconsistent with a protein-protein interaction, it precluded the use of coexpressed

exogenous proteins to test such an interaction. We therefore used a BirA*-SEC22A

fusion protein to verify the RAB18-SEC22A association, to identify other SEC22A-

associations, and to determine whether these associations were influenced by the

absence of RAB18 or its regulators. To minimize potential toxicity while increasing

biotin-ligase activity, we used BioID2 (Kim et al., 2016) with a p.Gly40Ser active site

modification (Branon et al., 2018) and reduced biotin incubation time to 6 hours. LFQ

analysis following streptavidin pull-down and mass spectrometry indicated that

BioID2(Gly40S)-SEC22A was present at much lower levels than the BirA*-RAB18 in

the prior experiments (3.79±0.96%). However, after adjusting for non-specific binding

and normalizing the data according to the quantity of BioID2(Gly40S)-SEC22A in

each sample, the construct appeared to label RAB18 in a RAB3GAP-dependent

manner (Figure 5B). RAB3GAP1 and RAB3GAP2, as well as the putative RAB18-

associated proteins REEP4 and BICD2, were among 55 SEC22A-associated

proteins present in samples from wild-type cells in >2 replicate experiments and

represented by >3 spectral counts (Table S5). Furthermore, also among these

proteins, a subset of 9 SEC22A-associations were attenuated (association ratios

<0.5) in samples from both RAB18-null and RAB3GAP-null cells. Broadly, these data

were consistent with a functional SEC22A-RAB18 interaction.

Given the involvement of RAB18, the NRZ complex, and a STX18 complex lacking

SEC22B in the regulation of LDs, we asked whether SEC22A might also be involved.

Multiple studies have shown that LD morphology is altered in lipid-loaded cells in

which RAB18 expression - or that of its regulators - is disrupted or silenced, with

fewer and/or larger LDs are observed in these cells compared to controls (Bekbulat

et al., 2018, Carpanini et al., 2014, Gerondopoulos et al., 2014, Li et al., 2017, Liegel

et al., 2013, Xu et al., 2018). Similar observations have been made in ZW10-, NBAS-

, STX18-, BNIP1- and USE1-null cells, whereas LD size distribution is unaltered

when SEC22B expression is silenced (Xu et al., 2018). We examined the effects of

silencing ZW10, NBAS and SEC22A in oleic acid-loaded immortalized human

hepatocyte (IHH) cells (Figure 5C). ZW10 and NBAS silencing provided positive

controls.

ZW10 silencing led to a significant reduction in LD number (p<0.001) compared to
controls, whereas NBAS silencing led to both a significant reduction in LD number
and a significant increase in LD size (p<0.001 in each case). The effects of SEC22A
silencing mirrored those of NBAS silencing, producing a significant reduction in LD
number (p<0.001) and a significant increase in LD size (p<0.001). Together, these
data implicate SEC22A in RAB18-mediated LD regulation.

<u>RAB18 recruits the orphan lipase TMCO4 to the ER membrane in a RAB3GAP-</u>
<u>dependent manner</u>

Among the lipid modifying/mobilizing proteins identified as potential RAB18 effectors
in HeLa cells, TMCO4 was identified in all three replicate experiments and its
association with RAB18 was highly RAB3GAP-dependent (association ratio 0.06).
TMCO4 (<u>t</u>rans<u>m</u>embrane and <u>co</u>iled-coil domains <u>4</u>) is annotated as containing
transmembrane and coiled-coil domains, but is orthologous to the Yeast protein Mil1/
Yfl034w. Mil1 was found to be peripherally membrane-associated/soluble, and is
thought to be an α/β hydrolase and a probable lipase based on structural modelling
(Whitfield et al., 2016). Consistent with this, a catalytic triad within its predicted active
site was shown to be required to confer tolerance to the membrane-intercalating
cationic amphipathic drug sertraline (Whitfield et al., 2016). The best template match
produced for TMCO4 by the structural modelling tool Phyre2 (Kelley et al., 2015) is
an acylglycerol/diacyclglycerol lipase (Figure S7). To explore the localization of
TMCO4, we expressed TMCO4-EGFP in HeLa cells (Figure 6A). This construct
showed a diffuse appearance consistent with a largely cytosolic localization. In
contrast, EGFP-RAB18 partly localizes to the ER, as shown by its colocalization with
an ER marker.

To assess the potential interaction between RAB18 and TMCO4, we coexpressed
mCherry-RAB18 and TMCO4-EGFP (Figure 6B). As in our previous experiments, we
used Celltrace reagents to distinguish cells of wild-type and mutant genotypes and
imaged these on the same dishes. In wild-type HeLa cells, we found that the
coexpression of mCherry-RAB18 led to a dramatic redistribution of TMCO4-EGFP to
the ER membrane suggesting that RAB18 is involved in the recruitment of TMCO4 to
this compartment (Figure 6B, upper panels). This redistribution was completely

absent in RAB3GAP1- and RAB3GAP2-null cells but unaffected in TRAPPC9-null

cells consistent with the BioID data. As a means of verifying the interaction, we

carried out immunoprecipitation experiments using exogenous HA-RAB18 and

TMCO4-EGFP (Figure 6C). As expected, TMCO4-EGFP copurified with HA-RAB18

when expressed in wild-type or TRAPPC9-null cells, but not when expressed in

RAB3GAP1-null cells. These data indicate that RAB18 and TMCO4 interact directly

or indirectly as part of a protein complex in a RAB3GAP-dependent manner. Further,

both the microscopy and the immunoprecipitation data support the suggestion that

different GEFs can promote different RAB18-interactions.

<u>RAB18 is involved in cholesterol mobilization and biosynthesis</u>

The lipid-related RAB3GAP-dependent RAB18-associated proteins in the HeLa

BioID dataset included ORP2/OSBPL2 and INPP5B, which are robustly linked to a

role in cholesterol mobilization. ORP2 is thought to function as a lipid transfer protein

that delivers cholesterol to the plasma membrane (PM) in exchange for $PI(4,5)P_2$,

whereas INPP5B is implicated in the hydrolysis of ORP2-bound $PI(4,5)P_2$,

presumably driving the exchange process (Wang et al., 2019). Two other proteins in

this group, C2CD2L/TMEM24 and C2CD2, have not been linked to cholesterol, but

C2CD2L is characterised as a PI transporter and found to promote PM $PI(4,5)P_2$

production (Lees et al., 2017). On the basis of these findings, we investigated the

potential role of RAB18 in cholesterol uptake and efflux.

We performed loading and efflux experiments to demonstrate the flux of

cholesterol/cholesteryl ester (CE) based on the activity of RAB18. Chinese hamster

ovary (CHO) cells were generated to stably express RAB18(WT), RAB18(Gln67Leu),

or RAB18(Ser22Asn)(Figure S8). These cells were then preincubated with

lipoprotein deficient serum (LPDS) before addition of [^{14}C]-oleate for 24 hours. [^{14}C]-

oleate was added in the presence of LPDS (Figure 7A, left panel) or 10% FBS

(Figure 7A, right panel). [^{14}C]-CE levels were measured at t=0, and efflux was

assessed by measuring CE levels 4 and 8 hours following the addition of high

density lipoprotein (HDL) to the cells. In each case, [^{14}C]-CE was isolated by thin

layer chromatography (TLC) and quantified by scintillation counting. Efflux was also

assessed directly by loading the cells with [^{3}H]-cholesterol, then incubating them with

apolipoprotein (apo) A-I. [^{3}H]-cholesterol associated with apoA-I in the medium, as a

percentage of total cellular radioactivity, is shown in Figure 7B.

In cells loaded with [^{14}C]-oleate/LPDS, levels of CE were comparable in

RAB18(Ser22Asn) and RAB18(WT) cells, whereas RAB18(Gln67Leu) cells stored

significantly more (Figure 7A, left panel). In cells loaded with [^{14}C]-oleate/FBS, levels

of CE in RAB18(Ser22Asn) remained unchanged, whereas its storage was elevated

in RAB18(WT) cells and RAB18(Gln67Leu) cells (Figure 7A, right panel).

Interestingly, in both [^{14}C]-oleate/LPDS-loaded and [^{14}C]-oleate/FBS-loaded cells,

the addition of HDL led to rapid depletion of CE in RAB18(Gln67Leu) cells, but not in

RAB18(Ser22Asn) or RAB18(WT) cells (Figure 7A). Consistently, [^{3}H]-cholesterol

also underwent significantly more rapid efflux from these cells (Figure 7B).

The above data indicated that 'activated' GTP-bound RAB18 strongly promotes the

turnover and mobilization of CE and were consistent with a role for RAB18 in

mediating cholesterol mobilization via ORP2. Given that RAB18 also associated with

EBP in our HeLa dataset, we next explored whether RAB18 might be necessary for

the mobilization of substrates or products of EBP-catalysis. EBP is involved in *de*

novo cholesterol biosynthesis (Silve et al., 1996). In the Bloch pathway, it catalyses

the conversion of 5α-cholesta-8, 24-dien-3β-ol (zymosterol) to 5α-cholesta-7, 24-

dien-3β-ol (24-dehydrolathosterol). In the Kandutsch-Russel pathway, it catalyses

the conversion of 5α-cholest-8(9)-en-3β-ol (zymostenol) to 5α-cholest-7-en-3β-ol

(lathosterol)(Platt et al., 2014). To identify any alteration in sterol levels in RAB18-

null HeLa cells as compared to wild-type cells, we first incubated them for 48 hours

in media supplemented with LPDS, then subjected samples to analysis by GC-MS

(Figure 7C).

In RAB18-null cells, we found that levels of the EBP substrate zymostenol were not

significantly different from those in wild-type cells. In contrast, levels of its product

lathosterol were significantly increased (p<0.01). These data are consistent with

intracellular accumulation of lathosterol resulting from a failure in its mobilization

away from the site of its generation by EBP (Figure 7D). Levels of the downstream

intermediate desmosterol were significantly reduced in RAB18-null (p<0.01),

consistent with impaired delivery of substrates to post-EBP biosynthetic enzymes.

We reasoned that impaired delivery of cholesterol precursors to post-EBP enzymes

might reduce cholesterol biosynthesis in cells in which RAB18 is absent or

dysregulated. We therefore assessed cholesterol biosynthesis in the HeLa cell lines

by incubating them for 24 hours in media supplemented with LPDS, then treating

them for 24 hours with $[^3H]$-acetate or $[^3H]$-mevalonate (Figure 7E-F). Free

cholesterol was separated by TLC and labelled cholesterol was quantified by

scintillation counting. Under both conditions, labelling was similar in two clonal wild-

type cell lines, but was reduced in RAB18-, RAB3GAP1-, RAB3GAP2-, TBC1D20-

and TRAPPC9-null cells. Labelling was lowest in RAB18-null cells (6.8±0.5% of wild-

type controls for acetate, 39.5±2.5% for mevalonate) and slightly higher in the

RAB3GAP1-, RAB3GAP2-, TBC1D20- and TRAPPC9-null lines (23±2%-43±3% for

acetate, 46±2.5%-73±5% for mevalonate). These data strongly suggest that RAB18

and its regulators are required for normal cholesterol biosynthesis.

Our model for the involvement of RAB18 in cholesterol biosynthesis is that it is

facilitates mobilization of EBP-products including lathosterol via ORP2 (Figure 7D).

Structural data are available for the ORP2 OSBP-related domain (ORD)(Wang et al.,

2019). Therefore, to test our model, we generated molecular dynamics simulations of

the ORP2-ORD incorporating either cholesterol or lathosterol (Figure 7G-H, Figure

S9, Videos S1-S2). Over 300ns, both simulations remained comparably stable in

terms of positional root-mean-square deviation (RSMD)(Figure S9A). The pattern of

positional root mean square fluctuations (RSMFs) for ORP2-ORD amino acid

residues in each simulation was also largely similar although some differences were

evident (Figure S9B). Among residues in the vicinity of the sterol, only a single Tyr

residue (Tyr110 in NP_653081) showed a clear difference. This residue appeared to

fluctuate less in the presence of lathosterol than in the presence of cholesterol.

Examination of the simulations suggested that Tyr110 adopts different positions in

the presence of the different sterols (Figure 7G-H, Figure S9C, Videos S1-S2). The

3-hydroxyl of cholesterol maintains a relatively stable distance from the Tyr. In

contrast, the 3-hydroxyl of lathosterol is frequently closer to the Tyr-hydroxyl, though

this varies over the timecourse (Figure S9C). In crystal structures of the OSBP-

related protein Osh4 and sterols, it was noted that water-mediated interactions were

prominent (Im et al., 2005). Therefore, we explored the positioning of water

molecules with respect to lathosterol and cholesterol in our simulations. Interestingly,

water molecules were much more frequently found in close proximity to the

lathosterol 3-hydroxyl than to the cholesterol 3-hydroxyl (Figure S9D). Together,

these data suggest that the ligand-binding tunnel of ORP2 adopts a relatively stable

but distinct conformations when bound to cholesterol or lathosterol. Despite their

minor structural differences, stabilization of the lathosterol 3-hydroxyl appears to

involve water-mediated interactions to a greater degree than that of the cholesterol

3-hydroxyl. Plots of the RMSFs of the atoms in each sterol (Figure S9E) show

generally higher fluctuations with lathosterol than with cholesterol. Speculatively, this

may suggest that the interaction with lathosterol is entropically more favourable.

DISCUSSION

The data presented in this study provide preliminary evidence that Warburg Micro

syndrome/RAB18 deficiency is associated with disrupted *de novo* cholesterol

biosynthesis. We have shown that lathosterol accumulates in RAB18-null HeLa cells

suggesting that its delivery to lathosterol oxidase (SC5DL) is impaired. Further, we

have shown that levels of cholesterol biosynthesis are reduced when RAB18 or its

regulators are absent. Pathogenic variants in *SC5DL* cause the rare recessive

disease lathosterolosis, which shares overlapping features with Micro syndrome

including microcephaly, intellectual disability, micrognathia, high arched palate and

cataract (Anderson et al., 2019, Brunetti-Pierri et al., 2002, Ho et al., 2014,

Krakowiak et al., 2003, Rossi et al., 2007). Similarly, the next enzyme in the pathway

is encoded by *DHCR7*, in which pathogenic variants cause Smith-Lemli-Opitz

syndrome (SLOS)(Nowaczyk & Wassif, 1998). While the symptoms of this disorder

can be highly variable, it is among the top differential diagnoses for Micro syndrome

(Handley & Sheridan, 2018).

Confirmation that features of Micro syndrome pathology arise from impaired

cholesterol biosynthesis could spur development of therapeutic strategies to treat the

disease. In SLOS, cholesterol supplementation has been widely used and more

recently is being combined with antioxidants with the aim of reducing toxicity from

aberrant cholesterol metabolites (Fliesler et al., 2018, Korade et al., 2014, Svoboda

et al., 2012). Supplementation with fat soluble vitamins may balance deficiencies.

Finally, paradoxical treatment with certain statins has been used in both SLOS and

lathosterolosis and may increase the expression of DHCR7 (Correa-Cerro et al.,

2006, Ho et al., 2014, Wassif et al., 2017). Careful clinical research will be required

to determine whether such interventions are safe, and whether they have any

efficacy in Micro syndrome.

In this study, we have complemented previous work showing that proximity

biotinylation is a powerful means of identifying candidate Rab effectors (Gillingham

et al., 2019). Further – at least in the case of RAB18 - we have found that comparing

biotin-labelling produced by a BirA*-Rab in wild-type and GEF-deficient cells can be

particularly informative. We found that marked reductions in RAB18-association in

RAB3GAP-null cells were restricted to a relatively small number of proteins and that

these comprised known and/or plausible interactors. By identifying these, we were

able to exclude ~95% of RAB18-associations from consideration as more likely to

represent 'noise' from bystander proteins. Interestingly, we found that the disruption

of different GEF complexes affected largely distinct subsets of RAB18 associations.

This may indicate that different RAB18-GEFs control different aspects of RAB18

function.

It has been proposed that RAB18 functions to coordinate lipid transfer between

apposed membranes, and our RAB3GAP-dependent interaction data are consistent

with this model (Xu et al., 2018). The 25 RAB3GAP-dependent RAB18 interactors

we identified include microtubule-interacting/membrane-remodelling proteins,

proteins involved in bringing membranes into close apposition, and proteins involved

in lipid modification and mobilization. It seems likely that membrane remodelling

precedes the engagement of tethers and SNAREs and the establishment of

membrane-contacts. Lipid transporter/exchange proteins are required for lipid

transfer to occur at these contact points, while lipid-modifying proteins may serve to

generate substrates for transport and/or concentration gradients necessary to

promote this transport.

Three key RAB18 interactors identified in our study were ORP2, INPP5B and EBP.

ORP2 is thought to act as sterol-PI(4,5)P_2 exchanger, whereas INPP5B is thought to

drive the exchange process by catalysing PI(4,5)P_2 hydrolysis (Wang et al., 2019).

The finding that these proteins each associate with RAB18 may suggest that they

function in an analogous manner to OSBP, the phosphatase SACM1L (SAC1) and

the ARF1 GTPase in mediating sterol exchange (Antonny et al., 2018). Because of

the role EBP in cholesterol biosynthesis, we were prompted to explore whether

substrates or products of EBP-catalysis accumulated in the absence of RAB18. The

finding that lathosterol but not zymostenol accumulates suggests that RAB18 is

required to coordinate the mobilization of EBP-products via ORP2 (Figure 7E).

The potential role of the other lipid-related proteins identified in our screen, C2CD2L,

C2CD2 and TMCO4, is less clear. However, since C2CD2L is found to transport PI

and to promote PI(4,5)P_2 generation (Lees et al., 2017), it is possible it functions in

concert with ORP2 and INPP5B. TMCO4 is poorly characterized. However, it was

identified in one study as one of a number of genes with upregulated expression in

rabbit cerebral arteries under conditions of hypercholesterolemia (Ong et al., 2013).

Another study suggests it is present on lipid rafts (Jin et al., 2012).

The regulated transport of specific sterol intermediates by RAB18 raises the
possibility that other Rab proteins might regulate transport of other intermediates or
regulate the biosynthetic pathways of other lipid species. In addition, it is possible
that particular intermediates might be delivered to different subcellular locations via
different Rabs. Interestingly, INPP5B has been shown in several studies to have a
broad Rab-binding specificity (Fukuda et al., 2008, Williams et al., 2007). Further,
many other phosphoinositide phosphatase enzymes, including members from
distinct protein families, bind to Rab proteins (Gillingham et al., 2019). Similarly,
multiple OSBP-homologues also bind to Rabs (Gillingham et al., 2019, Rocha et al.,
2009, Sobajima et al., 2018, Zhao & Ridgway, 2017). Thus, it is possible that
different Rab isoforms may specify intermembrane transfer of different sterols by
association with different combinations of phosphatase and OSBP proteins.

Differential mobilization of intermediates by discrete OSBP-INPP-Rab combinations
is one possible explanation for prominent features of lathosterolosis and SLOS that
do not overlap with Micro syndrome. Several such features resemble those caused
by impaired ciliogenesis, ciliary function or Sonic Hedgehog (SHH) deficiency, and
are ascribed to a deficit in cholesterol at the cilium (Blassberg et al., 2016, Nowaczyk
& Wassif, 1998). It may be that cholesterol biosynthesis at, or its delivery to cilia, is
less affected in Micro syndrome because alternative lipid transfer mechanisms are
involved.

One of the clinical features present in Micro syndrome but not in lathosterolosis or
SLOS is ascending spastic paraplegia. In this context, the reduced levels of SPG20
in RAB18- and TBC1D20-null cells may be relevant. Biallelic loss-of-function variants
in *SPG20* cause Troyer syndrome, a complex HSP also characterized by ascending
paraplegia (Baple & Crosby, 2004). SPG20 is a protein of enigmatic function, but
one possible avenue for future research could be the potential relationship between
SPG20 and the other RAB18-associated microtubule (MT)-binding proteins identified
in this study. Like SPG20, CAMSAP1 is found to associate with poles of the mitotic
spindle (Hueschen et al., 2017, Lind et al., 2011). CAMSAP1 is a MT minus-end
binding protein whereas BICD2 is a component of a minus-end-directed dynein-
dynactin motor complex (Hendershott & Vale, 2014, Hueschen et al., 2017,
Urnavicius et al., 2015). REEP4 is localized to the ER, and like RAB18 is linked to
the regulation of ER structure (Gerondopoulos et al., 2014, Kumar et al., 2019).

However, it also contributes to the clearance of ER from metaphase chromatin

(Schlaitz et al., 2013). This role is dependent on its MT-binding, and therefore

spindle-binding during metaphase. Thus, these four MT-binding proteins all appear

to function in the same spatiotemporally defined compartment during mitosis (Figure

S10A).

Among the membrane tethering and docking proteins associated with RAB18, we

investigated SEC22A on the basis of its homology to SEC22B. SEC22B is a

component of the canonical, fusogenic, syntaxin 18 SNARE complex. Other

components of this complex interact with RAB18 to regulate membrane contacts,

though SEC22B itself is not necessary for this activity (Xu et al., 2018). Reciprocal

proximity biotinylation with a BioID2-SEC22A fusion was consistent with a

RAB3GAP-dependent RAB18-SEC22A interaction, and suggested that other

SEC22A interactions are altered when RAB18 is absent or dysregulated. We also

found that the silencing of SEC22A led to altered LD morphology, similar to that

previously observed when expression of other RAB18-associated tethering/docking

proteins was disrupted. Though our data are tentative, it is tempting to speculate that

SEC22A contributes to a non-fusogenic SNARE complex, since it lacks the SNARE

domain usually required to provide the mechanical force for fusion (Figure S10B).

SNARE proteins can be relatively promiscuous in their interactions, and there are

numerous examples of individual SNAREs contributing to distinct SNARE complexes

(Petkovic et al., 2014, Wang et al., 2017). There are also examples in which SNARE

complexes mediate the stable association of membranes rather than their fusion

(Petkovic et al., 2014).

To summarise, we have used complementary proximity ligation approaches together

with CRISPR gene-editing to inventory RAB18-associated proteins. The RAB3GAP-

dependent RAB18-associations we have highlighted are supported by additional

validation and functional evidence as well as by correlative information from the

literature. Broadly, the latter suggests a role for Rab proteins distinct from that in

mediating vesicular membrane traffic. Another role of Rabs may be to assemble a

network of diverse membrane-membrane contact sites and then orchestrate highly

regulated metabolism and flow of lipid species within a discontinuous lipid phase.

Disruption of specific contact sites can produce metabolic deficiency and may cause

human disease.

 MATERIALS AND METHODS

 <u>Plasmids</u>

 Generation of recombinant pcDNA5 FRT/TO FLAG-BirA(Arg118Gly) vectors for
 preparation of stable T-Rex-293 cell lines is described below. Generation of
 recombinant pX461 and pX462 plasmids for CRISPR gene-editing is described
 below. Generation of recombinant pCMV vectors for preparation of stable CHO cell
 lines is described below. The EGFP-RAB18 construct has been described previously
 (Gerondopoulos et al., 2014). The RAB18 sequence was excised from this construct
 using BamHI and HindIII restriction enzymes (New England Biolabs, Hitchin, UK),
 and used to generate constructs encoding mEmerald-RAB18 and mCherry-RAB18
 by ligation into mEmerald-C1 and mCherry-C1 vectors (Addgene, Watertown, MA)
 using HC T4 Ligase and rapid ligation buffer (Promega, Southampton, UK).
 Constructs encoding BirA*-RAB18, BioID2(Gly40Ser)-SEC22A, mEmerald-SEC22A,
 mEmerald-SPG20 and mEmerald-RAB7A were generated following PCR
 amplification from template and subcloning into an intermediate pCR-Blunt II-TOPO
 vector using a Zero Blunt TOPO PCR Cloning Kit (ThermoFisher Scientific,
 Waltham, MA) according to manufacturer's instructions. Fragments were excised
 from intermediate vectors and then subcloned into target vectors using restriction-
 ligation, as above. A construct encoding mCherry-ER was obtained from Addgene,
 and a construct encoding TMCO4-EGFP was synthesised and cloned by GeneWiz
 (Leipzig, Germany). Details of PCR templates, primers and target vectors are listed
 in Table S6.

 <u>Antibodies and reagents</u>

 A custom polyclonal antibody to RAB18 generated by Eurogentec (Southampton,
 UK) has been described previously (Handley et al., 2015). An antibody to
 RAB3GAP1 was obtained from Bethyl Labs (Montgomery, TX), an antibody to GFP
 was obtained from Takara Bio (Saint-Germain-en-Laye, France), an antibody to β-
 Tubulin was obtained from Abcam (Cambridge, UK) and an antibody to β-Actin was
 obtained from ThermoFisher. Antibodies to hemagglutinin (HA), RAB3GAP2 and
 TBC1D20 were obtained from Merck (Gillingham, UK). Antibodies to ZW10, STX18,

SPG20, RINT1, REEP4, BNIP1, C2CD2, TRIM13, WFS1, INPP5B, OSBPL2 and

NBAS were obtained from Proteintech (Manchester, UK). Antibody catalogue

numbers and the dilutions used in this study are listed in Table S6.

<u>Cell culture</u>

T-REx-293,HeLa and IHH cells were maintained in DMEM media, RPE1 cells in

DMEM/F12 media and CHO cells in alpha-MEM media (ThermoFisher). In each

case, media was supplemented with 10% foetal calf serum (FCS) and 1% penicillin-

streptomycin (PS). Cells were maintained at 37°C and 5% CO_2.

<u>Generation of stable T-Rex-293 and CHO cell lines</u>

PCR products encoding mouse RAB18, RAB18(Gln67Leu) and RAB18(Ser22Asn)

were subcloned into NotI-linearized pcDNA5 FRT/TO FLAG-BirA(Arg118Gly) vector

using the In-Fusion HD EcoDry Cloning Plus kit (Takara Bio) according to

manufacturer's instructions. Details of PCR templates, primers and target vectors are

listed in Table S6. 1.5ug of each recombinant vector together with 13.5ug of pOG44

plasmid (ThermoFisher) were used in cotransfections of T-REx-293 cells, in 10cm

dishes, with TransIT-LT1 Transfection Reagent (Mirus Bio, Madison, WI).16 hours

following transfection, media was replaced and cells were allowed to recover for 24

hours. Each dish was then split to 4x 10cm dishes in selection media containing 10

ug/ml Blasticidin and 50 ug/ml Hygromycin B. Resistant clones were pooled and

passaged once prior to use.

A PCR product encoding mouse RAB18 was subcloned into an intermediate TOPO

vector using a TOPO PCR Cloning Kit (ThermoFisher) according to manufacturer's

instructions. The RAB18 fragment was then excised and subcloned into the pCMV

vector. PCR-based site-directed mutagenesis using a GeneArt kit (ThermoFisher)

was then used to generate pCMV-RAB18(Gln67Leu) and pCMV-RAB18(Ser22Asn)

constructs. CHO cells were transfected using Lipofectamine 2000 reagent

(ThermoFisher) and cells stably-expressing each construct were selected-for with

blasticidin. Under continued selection, clonal cell-lines were grown from single cells

and then RAB18 protein expression was assessed. Cell lines comparably expressing

RAB18 constructs at levels 2.5-5x higher than those wild-type cells were used in

subsequent experiments.

<u>Generation of clonal 'knockout' HeLa and RPE1 cell lines</u>

CRISPR/Cas9 gene-editing was carried out essentially as described in Ran et al.,

2013 (Ran et al., 2013). Guide RNA (gRNA) sequences are shown in (Table S6). A

list of the clonal cell lines generated for this study, together with the loss-of-function

variants they carry is shown in (Figure S2A). Western blot validation is shown in

(Figure S2B-E). Briefly, for each targeted exon, pairs of gRNA sequences were

selected using the online CRISPR design tool (http://crispr.mit.edu/). Oligonucleotide

pairs incorporating these sequences (Sigma) were annealed (at 50mM ea.) in 10mM

Tris pH8, 50mM NaCl and 1mM EDTA by incubation at 95ºC for 10 minutes followed

by cooling to room temperature. Annealed oligonucleotides were diluted and ligated

into BbsI-digested pX461 and pX462 plasmids (Addgene) using HC T4 Ligase and

rapid ligation buffer (Promega). Sequences of all recombinant plasmids were verified

by direct sequencing. Pairs of plasmids were contransfected into cells using

Lipofectamine 2000 reagent according to manufacturer's instructions. Cells were

selected for puromycin resistance (conferred by pX462) using 24 hours puromycin-

treatment. Following 12 hours recovery, they were selected for GFP fluorescence

(conferred by pX461) and cloned using FACSAria2 SORP, Influx or FACSMelody

instruments (BD, Wokingham, UK). After sufficient growth, clones were analysed by

PCR of the targeted exons (Primers are listed in Table S6). In order to sequence

individual gene edited-alleles, PCR products from each clone were first cloned into

ZeroBlunt TOPO vector (ThermoFisher) and then subjected to colony PCR. These

PCR products were then analysed by direct sequencing. Sequencing data was

assessed using BioEdit software (http://www.mbio.ncsu.edu/BioEdit/bioedit.html).

<u>BirA/BioID proximity labelling (T-REx-293 cells)</u>

The T-REx-293 Cell Lines (described above) were seeded onto 3x 15cm plates each

and allowed to adhere. Expression of BirA*-RAB18 fusion proteins was induced by

treatment with 20ng/ml Tetracycline for 16 hours. Media was then replaced with

media containing 20% FBS, 20 ng/ml Tetracycline and 50 uM Biotin and the cells

were incubated for a further 8 hours, washed with warmed PBS and pelleted in ice-

cold PBS. Cell pellets were snap-frozen and stored at -80ºC prior to lysis. Lysis was

carried out in 3ml of ice-cold RIPA buffer (150 mM NaCl, 1% NP40, 0.5% Sodium

Deoxycholate, 0.1% SDS, 1mM EDTA, 50mM Tris, pH 7.4) supplemented with

complete-mini protease inhibitor cocktail (Roche, Basel, Switzerland), 1mM PMSF,

and 62.5 U/ml Benzonase (Merck). Lysates were incubated for 1 hour at 4ºC then

sonicated in an ice bath (four 10 second bursts on low power). They were then

clarified by centrifugation, and the supernatants transferred to tubes containing pre-

washed streptavidin-sepharose (30µl bed-volume)(Merck). The beads were

incubated for 3 hours at 4ºC, then washed x5 in RIPA buffer and x4 in buffer

containing 100mM NaCl, 0.025% SDS and 25 mM Tris, pH7.4.

<u>BirA*/BioID proximity labelling (HeLa cells)</u>

Proximity-labelling in HeLa cells was carried out largely as described by Roux et al.

(Roux et al., 2012), but with minor modifications. HeLa cells were grown to 80%

confluence in T75 flasks and then each flask was transfected with 1-1.5µg of the

BirA*-RAB18 construct or 1µg of the BioID2(Gly40Ser)-SEC22A construct using

Lipofectamine 2000 reagent in Optimem serum-free medium (ThermoFisher) for 4

hours, according to manufacturer's instructions. 24 hours post-transfection, media

was replaced with fresh media containing 50µM Biotin (Merck) and the cells were

incubated for a further 24 or 6 hours (for BirA*-RAB18 and BioID2(Gly40Ser)-

SEC22A experiments respectively). Cells were then trypsinised and washed twice in

PBS before pellets were transferred to 2ml microcentrifuge tubes and snap-frozen.

For each pellet, lysis was carried out in 420µl of a buffer containing 0.2% SDS, 6%

Triton-X-100, 500mM NaCl, 1mM DTT, EDTA-free protease-inhibitor solution

(Expedeon, Cambridge, UK), 50mM Tris pH7.4. Lysates were sonicated for 10

minutes using a Bioruptor device together with protein extraction beads (Diagenode,

Denville, NJ). Each lysate was diluted with 1080µl 50mM Tris pH7.4, and they were

then clarified by centrifugation at 20 000xg for 30 minutes at 4ºC. Affinity purification

of biotinylated proteins was carried out by incubation of clarified lysates with

streptavidin-coated magnetic Dynabeads (ThermoFisher) for 24 hours at 4ºC. Note

that a mixture of Dynabeads - MyOne C1, MyOne T1, M270 and M280 – was used

to overcome a problem with bead-clumping observed when MyOne C1 beads were

used alone. Successive washes were carried out at room temperature with 2% SDS,

a buffer containing 1% Triton-X-100, 1mM EDTA, 500mM NaCl, 50mM HEPES

pH7.5, a buffer containing 0.5% NP40, 1mM EDTA, 250mM LiCl, 10mM Tris pH7.4,

50mM Tris pH7.4 and 50mM ammonium bicarbonate.

<u>Preparation of cell lysates for label-free quantitative proteomics</u>

RPE1 and HeLa cells were grown to confluence in T75 flasks. They were then

trypsinised, and cell pellets were washed with PBS and snap-frozen prior to use.

RPE1 pellets were resuspended in 300µl 6M GnHCl, 75mM Tris, pH=8.5. HeLa

pellets were resuspended in 300µl 8M urea, 75mM NaCl, 50mM Tris, pH=8.4. In

each case, samples were sonicated for 10 minutes using a Bioruptor device together

with protein extraction beads (Diagenode). RPE1 samples were heated for 5 minutes

at 95ºC. Samples were clarified by centrifugation.

<u>Mass spectrometry</u>

Washed beads from BioID experiments with T-Rex-293 cell lines were resuspended

in 50µl 6M urea, 2M thiourea, 10mM Tris, pH=8.5 and DTT was added to 1mM. After

30 minutes incubation at 37ºC, samples were alkylated with 5mM iodoacetamide

(IAA) in the dark for 20minutes. DTT was increased to 5mM and 1µg lysC was

added, then samples were incubated at 37ºC for 6 hours. Samples were diluted to

1.4M urea, then digested with trypsin (Promega), overnight at 37°C, according to

manufacturer's instructions. Samples were acidified by the addition of 0.9% formic

acid and 5% acetonitrile.

Washed beads from BioID experiments with HeLa cells were subjected to limited

proteolysis by trypsin (0.3 ug) at 27°C for 6.5hours in 2mM urea, 1mM DTT, 75mM

Tris, pH=8.5, then supernatants were incubated overnight at 37°C. Samples were

alkylated with 50mM IAA in the dark for 20minutes, then acidified by addition of 8µl

10% trifluoroacetic acid (TFA). RPE1 lysates were reduced and alkylated through

addition of tris(2-carboxyethyl)phosphine (TCEP) and 2-chloroacetamide (CAA) to

5mM and 10mM respectively and then incubated at 95°C for 5 minutes. After

cooling, samples were diluted to 3M guanidine and 0.5µg lysC added with incubation

overnight at 37°C. A further dilution to 1M guanidine was followed by digest with

0.3µg trypsin at 37°C for 4 hours. Samples were acidified with TFA. HeLa lysates

were reduced and alkylated by addition of DTT to 10mM, then by addition of IAA to

25mM, then further addition of DTT to 25mM, with incubation at room temperature

for 30-60 minutes following each step. Samples were digested with lysC, overnight at

37°C. They were then diluted to 2M urea, and further digested, overnight at 37°C.

Samples were acidified with TFA.

For BioID samples from T-REx-293 cells, LC-MS was carried out as previously

described (Brunet et al., 2016). Briefly, peptides in an aqueous solution containing

5% acetonitrile and 0.1% formic acid were loaded onto a 3 µm PepMap100, 2 cm, 75

µm diameter sample column using an Easy nLC 1000 ultrahigh pressure liquid

chromatography system (ThermoFisher). They were eluted with acetonitrile/formic

acid into an in-line 50 cm separating column (2 µm PepMap C18, 75 µm diameter) at

40°C. Separated peptides were ionized using an Easy Spray nano source and

subjected to MS/MS analysis using a Velos Orbitrap instrument (ThermoFisher).

Following acquisition, data were analysed using SEAQUEST software.

For other samples, peptides were loaded on to activated (methanol), equilibrated

(0.1% TFA) C18 stage tips before being washed with 0.1% TFA and eluted with

0.1% TFA/80 acetonitrile. The organic was dried off, 0.1% TFA added to 15 µl and 5

µl injected onto LC-MS. Peptides were separated on an Ultimate nano HPLC

instrument (ThermoFisher), and analysed on either an Orbitrap Lumos or a Q

Exactive Plus instrument (ThermoFisher). After data-dependent acquisition of HCD

fragmentation spectra, data were analysed using MaxQuant and the uniprot human

reference proteome. Versions, releases, parameters and gradients used for

separation are provided in table S6.

<u>Cell labelling</u>

In order to distinguish cells of different genotypes within the same well/on the same

coverslip, CellTrace Violet and CellTrace Far Red reagents (ThermoFisher) were

used to label cells before they were seeded. Cells of different genotypes were first

trypsinised and washed with PBS separately. They were then stained in suspension

by incubation with either 1µM CellTrace Violet or 200nM CellTrace Far Red for 20

minutes at 37°C. Remaining dye was removed by addition of a ten-fold excess of full

media, incubation for a further 5 minutes, and then by centrifugation and

resuspension of the resulting pellets in fresh media. Differently-labelled cells were

combined prior to seeding.

<u>Immunofluorescence microscopy</u>

HeLa or RPE1 cells were seeded in 96-well glass-bottom plates (PerkinElmer,

Waltham, MA) coated with Matrigel (Corning, Amsterdam, Netherlands) according to

manufacturer's instructions, and allowed to adhere for 48 hours prior to fixation. In

lipid-loading experiments, cells were treated with 200µM oleic acid complexed to

albumin (Merck) and 1µg/ml BODIPY-558/568-C12 (ThermoFisher) for 15 hours

prior to fixation. Cells were fixed using a solution of 3% deionised Glyoxal, 20%

EtOH, 0.75% acetic acid, pH=5 (Richter et al., 2018), for 20 minutes at room

temperature. They were then washed with PBS containing 0.9mM $CaCl_2$ and 0.5mM

$MgCl_2$ and blocked with a sterile-filtered buffer containing 1% Milk, 2% donkey serum

(Merck), 0.05% Triton-X-100 (Merck), 0.9mM $CaCl_2$ and 0.5mM $MgCl_2$ in PBS

pH=7.4 for at least 1 hour prior to incubation with primary antibody. Primary

antibodies were added in blocking buffer without Triton-X-100, and plates were

incubated overnight at 4ºC. Antibody dilutions are listed in Table S6. Following

washing in PBS, cells were incubated with 1:2000 Alexa 488-conjugated secondary

antibody (ThermoFisher) in blocking buffer at room temperature for 1-2 hours.

Following further washing in PBS, cells were imaged using an Operetta High Content

Imaging System (PerkinElmer) equipped with Harmony software. In comparative

fluorescence quantitation experiments, at least 18 frames – each containing >5 wild-

type and >5 mutant cells – were analysed per genotype. ImageJ software was used

to produce regions of interest (ROIs) corresponding to each cell using thresholding

tools and images from the 405nm and 645nm channels. Median 490nm fluorescence

intensity was measured for each cell and mutant fluorescence intensity (as %wild-

type) was calculated for each frame and combined for each genotype.

<u>Confocal microscopy – Live cell imaging</u>

HeLa or RPE1 cells were seeded on glass-bottom dishes (World Precision

Instruments, Hitchin, UK) coated with Matrigel (Corning) and allowed to adhere for

24 hours prior to transfection. Transfections and cotransfections were carried out

with 0.5µg of each of the indicated constructs using Lipofectamine 2000 reagent in

Optimem serum-free medium for 4 hours, according to manufacturer's instructions.

Media were replaced and cells were allowed to recover for at least 18 hours prior to

imaging. Imaging was carried out on a Nikon A1R confocal microscope equipped

with the Nikon Perfect Focus System using a 60× oil immersion objective with a 1.4

numerical aperture. In immunofluorescence experiments, the pinhole was set to

airy1. CellTrace Violet was excited using a 403.5nm laser, and emitted light was

collected at 425–475nm. EGFP and mEmerald were excited using a 488 nm laser,

and emitted light was collected at 500–550 nm. BODIPY-558/568-C12 and mCherry

were excited using a 561.3 nm laser, and emitted light was collected at 570–620 nm.

CellTrace Far Red was excited using a 638nm laser, and emitted light was collected

at 663-738nm. In fluorescence recovery after photobleaching (FRAP) experiments,

the pinhole was set to airy2 and digital zoom parameters were kept constant.

Bleaching was carried out using 90% laser power.

1005

<u>Immunoprecipitation</u>

HeLa cells were seeded onto 10cm dishes and allowed to adhere for 24 hours prior

to transfection. Transfections and cotransfections were carried out with 0.5µg of

each of the indicated constructs using Lipofectamine 2000 reagent in Optimem

serum-free medium for 4 hours, according to manufacturer's instructions. 24 hours

post-transfection cells were trypsinised, washed with PBS, then lysed in a buffer

containing 150mM NaCl, 0.5% Triton-X-100 and EDTA-free protease-inhibitor

solution (Expedeon), 10mM Tris, pH=7.4. Lysates were clarified by centrifugation,

input samples taken, and the remaining supernatants then added to 4µg rabbit anti-

HA antibody (Merck). After 30 minutes incubation at 4ºC on a rotator, 100µl washed

protein G-coupled Dynabeads (ThermoFisher) were added and samples were

incubated for a further 1 hour. The Dynabeads were washed x3 with buffer

containing 150mM NaCl, 0.1% Triton-X-100, 10mM Tris, pH=7.4, then combined with

a reducing loading buffer and subjected to SDS–PAGE.

 Lipid loading experiments

 For LD number and diameter measurements, IHH cells were seeded onto glass

 coverslips. siRNA transfections were carried out using FuGene reagent (Promega)

 according to manufacturer's instructions. siRNAs targeting ZW10 and NBAS were

 obtained from IDT, Coralville, IA; siRNA targeting SEC22A was obtained from

 Horizon Discovery, Cambridge, UK. 48 hours following transfection, cells were

 treated with 200nM BSA conjugated oleate for 24 hours. Coverslips were washed,

 fixed with 3% paraformaldehyde and stained with 1ug/mL BODIPY and 300nM DAPI.

 Fluorescence images were captured on a Zeiss LSM 780 confocal microscope

 equipped with a 100x objective. Images were analysed using ImageJ software. Data

 are derived from measurements from >100 cells/condition and are representative of

 three independent experiments.

 For cholesterol storage and efflux experiments with [^{14}C]-oleate, CHO cell lines

 (described above) were seeded onto 12-well plates and then grown to 60-75%

 confluence in Alpha media supplemented with 10% LPDS. Cells were grown in the

 presence of 10% LPDS for at least 24 hours prior to the addition of oleate. 1 µCi/ml

 [^{14}C]-oleate (Perkin Elmer) was added in the presence of 10% LPDS or 10% FBS for

 24 hours. Cells were then washed and incubated with 50µg/ml HDL for 0, 4 or 8

 hours. Cellular lipids were extracted with hexane. Lipids were then dried-down and

 separated by thin layer chromatography (TLC) in a hexane:diethyl ether:acetic acid

 (80:20:2) solvent system. TLC plates were obtained from Analtech, Newark, NJ.

 Bands corresponding to cholesteryl ester (CE) were scraped from the TLC plate, and

 radioactivity was determined by scintillation counting in a Beckman Coulter LS6500

 Scintillation Counter using BetaMax ES Liquid Scintillation Cocktail (ThermoFisher).

 Three independent experiments were carried out, each with four replicates of each

 condition. Data from a representative experiment are shown.

 For cholesterol efflux experiments with [^{3}H]-cholesterol, CHO cells were seeded onto

 12-well plates and then grown to 60% confluence in Alpha media supplemented with

 10% FBS. 5 µCi/ml [^{3}H]-cholesterol (PerkinElmer) was added in the presence of 10%

 FBS. After 3x PBS washes, cells were incubated with serum-free media containing

 25µg/ml of human apolipoprotein A-I (ApoA-I) for 5 hours. ApoA-I was a kind gift of

 Dr. Paul Weers (California State University, Long Beach). Radioactivity in aliquots of

1052   media were determined by scintillation counting in a Beckman Coulter LS6500

1053   Scintillation Counter using LSC Cocktail (PerkinElmer). Cell lysates were produced

1054   by addition of 0.1N NaOH for 1 hour, and their radioactivity was determined as

1055   above. Cholesterol efflux was calculated as an average (+/- SD) of the % cholesterol

1056   efflux (as a ratio of the media cpm/(media + cellular cpm) x 100%).

1057

<u>Sterol analysis</u>

HeLa cells were grown to 80% confluence in T75 flasks, washed twice in PBS and

then grown for a further 48 hours in DMEM supplemented with 10% LPDS. They

were then trypsinised and washed twice in PBS before pellets were transferred to

microcentrifuge tubes and snap-frozen. Pellets were resuspended in 200µl deionised

water, sonicated for 20 seconds using an ultrasonic processor (Sonics & Materials

Inc., CT, USA), then placed on ice. 750 µl of isopropanol containing 4 µmol/L 5α-

cholestane as an internal standard was added to each sample, and then each was

sonicated for a further 10 seconds. Lysates were transferred to 7ml glass vials and

mixed with 250 µL tetramethylammonium hydroxide for alkalyine saponification at

80°C for 15 min, then cooled down for 10 minutes at room temperature. Sterols were

extracted by addition of 500 µL tetrachloroethylene/methyl butyrate (1:3) and 2 ml

deionised water, then thorough mixing. Samples were centrifuged for 10 minutes at

3000 rpm, and the organic phase containing the sterols was transferred to 300 µl GC

vials. Extracts were dried under a stream of nitrogen, then sterols were silylated with

50 µL Tri-Sil HTP (HDMS:TMCS:Pyridine) Reagent (ThermoFisher) at 60°C for 1

hour.

Chromatography separation was performed on an Agilent gas chromatography-mass

spectrometry (GC-MS) system (6890A GC and 5973 MS) (Agilent Technologies,

Inc., CA, USA) with an HP-1MS capillary column (30 m length. x 250 µm diameter x

0.25 µm film thickness). The GC temperature gradient was as follows: Initial

temperature of 120°C increased to 200°C at a rate of 20°C/min, then increased to

300°C at a rate of 2°C/min with a 15 minute solvent delay. Injection was at 250°C in

splitless mode with ultrapurified helium as the carrier gas and the transfer line was

280°C. The mass spectra were acquired by electron impact at 70 eV using selected

ion monitoring as follows: Lathosterol-TMS, cholesterol-TMS, and cholest8(9)-enol-

TMS: m/z 458; 5α-cholestane and desmosterol-TMS: m/z 372; Lanosterol-TMS: m/z

393; and 7-dehydrocholesterol-TMS: m/z 325. The data were analysed using

MassHunter Workstation Quantitative Analysis Software (Agilent Technologies, Inc.)

and OriginPro 2017 (OriginLab Corp, MA, USA).

1088

<u>Cholesterol biosynthesis assays</u>

HeLa cell lines (described above) were seeded onto 10 cm plates and then grown in

DMEM supplemented with 10% LPDS for 24 hours. Cells were washed and then

incubated with 10µCi/ml [^{3}H]-acetate or 5µCi/ml [^{3}H]-mevalonate (Perkin Elmer) in

the presence of 10% LPDS for another 24 hours. Cellular lipids were extracted with

hexane and then evaporated to dryness. Cell lysates were produced by addition of

0.1N NaOH for 1 hour. Free cholesterol was separated by TLC in a hexane:diethyl

ether:methanol:acetic acid (80:20:3:1.5) solvent system. TLC plates were obtained

from Analtech, Newark, NJ. Bands corresponding to free cholesterol were scraped

from the TLC plate, and radioactivity was determined by scintillation counting in a

Beckman Coulter LS6500 Scintillation Counter using BetaMax ES Liquid Scintillation

Cocktail (ThermoFisher).

1101

<u>Molecular dynamics</u>

Simulations were constructed as previously described, starting from a modified

version of the OSBPL2 lipid-binding domain structure (PDB 5ZM8)(Wang et al.,

2019). Parameters for lathosterol were obtained by modifying those of cholesterol to

substituting the double bond between C5-C6 with one between C7-C8. Simulations

were performed with the CHARMM36 force field.

1108

<u>Western blotting</u>

Cell lysates were made with a buffer containing 150mM NaCl, 0.5% Triton-X-100

and EDTA-free protease-inhibitor solution (Expedeon), 50mM Tris, pH=7.4. Cell

lysates and input samples from BioID and immunoprecipitation experiments were

combined 1:1 with a 2x reducing loading buffer; a reducing loading buffer containing

10mM EDTA was added directly to Dynabead samples. SDS–PAGE and Western

blotting were carried out according to standard methods.

1116

ACKNOWLEDGEMENTS

1118 We thank the Warburg Micro syndrome children and their families. We thank

1119 Professor C. A. Johnson and Dr J. A. Poulter for a critical reading of the manuscript

1120 and Professor Jia-Wei Wu for sharing structural data.

1121

COMPETING INTERESTS

1123 No competing interests declared.

1124

FUNDING

1126 MH is supported by the University of Leeds and by a Programme Grant from the

1127 Newlife Foundation for Disabled Children (Grant Reference Number: 17-18/23).

1128

DATA AVAILABILITY

The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2019) partner

repository with the dataset identifiers PXD016631, PXD016336, PXD016326,

PXD016233 and PXD016404.

REFERENCES

Aligianis IA, Johnson CA, Gissen P, Chen D, Hampshire D, Hoffmann K, Maina EN,
Morgan NV, Tee L, Morton J, Ainsworth JR, Horn D, Rosser E, Cole TR, Stolte-
Dijkstra I, Fieggen K, Clayton-Smith J, Megarbane A, Shield JP, Newbury-Ecob R et
al. (2005) Mutations of the catalytic subunit of RAB3GAP cause Warburg Micro
syndrome. *Nat Genet* 37: 221-3

Anderson R, Rust S, Ashworth J, Clayton-Smith J, Taylor RL, Clayton PT, Morris
AAM (2019) Lathosterolosis: A Relatively Mild Case with Cataracts and Learning
Difficulties. *JIMD Rep* 44: 79-84

Angebault C, Fauconnier J, Patergnani S, Rieusset J, Danese A, Affortit CA,
Jagodzinska J, Megy C, Quiles M, Cazevieille C, Korchagina J, Bonnet-Wersinger D,
Milea D, Hamel C, Pinton P, Thiry M, Lacampagne A, Delprat B, Delettre C (2018)
ER-mitochondria cross-talk is regulated by the Ca(2+) sensor NCS1 and is impaired
in Wolfram syndrome. *Sci Signal* 11

Antonny B, Bigay J, Mesmin B (2018) The Oxysterol-Binding Protein Cycle: Burning
Off PI(4)P to Transport Cholesterol. *Annu Rev Biochem* 87: 809-837

Baple E, Crosby A (2004) Troyer Syndrome. In *GeneReviews((R))*, Adam MP,
Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A (eds)
Seattle (WA):

Barr F, Lambright DG (2010) Rab GEFs and GAPs. *Curr Opin Cell Biol* 22: 461-70

Bekbulat F, Schmitt D, Feldmann A, Huesmann H, Eimer S, Juretschke T, Beli P,
Behl C, Kern A (2018) RAB18 impacts autophagy via lipid droplet-derived lipid
transfer and is rescued by ATG9A. *bioRxiv*: 421677

Bem D, Yoshimura S, Nunes-Bastos R, Bond FC, Kurian MA, Rahman F, Handley
MT, Hadzhiev Y, Masood I, Straatman-Iwanowska AA, Cullinane AR, McNeill A,
Pasha SS, Kirby GA, Foster K, Ahmed Z, Morton JE, Williams D, Graham JM,
Dobyns WB et al. (2011) Loss-of-function mutations in RAB18 cause Warburg micro
syndrome. *Am J Hum Genet* 88: 499-507

Blassberg R, Macrae JI, Briscoe J, Jacob J (2016) Reduced cholesterol levels impair
Smoothened activation in Smith-Lemli-Opitz syndrome. *Hum Mol Genet* 25: 693-705

Borck G, Wunram H, Steiert A, Volk AE, Korber F, Roters S, Herkenrath P, Wollnik
B, Morris-Rosendahl DJ, Kubisch C (2011) A homozygous RAB3GAP2 mutation
causes Warburg Micro syndrome. *Human genetics* 129: 45-50

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL,
Perrimon N, Ting AY (2018) Efficient proximity labeling in living cells and organisms
with TurboID. *Nat Biotechnol* 36: 880-887

Brunet S, Saint-Dic D, Milev MP, Nilsson T, Sacher M (2016) The TRAPP Subunit
Trs130p Interacts with the GAP Gyp6p to Mediate Ypt6p Dynamics at the Late Golgi.
Front Cell Dev Biol 4: 48

Brunetti-Pierri N, Corso G, Rossi M, Ferrari P, Balli F, Rivasi F, Annunziata I,
Ballabio A, Russo AD, Andria G, Parenti G (2002) Lathosterolosis, a novel multiple-
malformation/mental retardation syndrome due to deficiency of 3beta-
hydroxysteroid-delta5-desaturase. *Am J Hum Genet* 71: 952-8

Bui M, Gilady SY, Fitzsimmons RE, Benson MD, Lynes EM, Gesson K, Alto NM,
Strack S, Scott JD, Simmen T (2010) Rab32 modulates apoptosis onset and
mitochondria-associated membrane (MAM) properties. *The Journal of biological*
chemistry 285: 31590-602

Cai H, Reinisch K, Ferro-Novick S (2007) Coats, tethers, Rabs, and SNAREs work
together to mediate the intracellular destination of a transport vesicle. *Developmental*
cell 12: 671-82

Carney DS, Davies BA, Horazdovsky BF (2006) Vps9 domain-containing proteins:
activators of Rab5 GTPases from yeast to neurons. *Trends Cell Biol* 16: 27-35

Carpanini SM, McKie L, Thomson D, Wright AK, Gordon SL, Roche SL, Handley MT,
Morrison H, Brownstein D, Wishart TM, Cousin MA, Gillingwater TH, Aligianis IA,
Jackson IJ (2014) A novel mouse model of Warburg Micro syndrome reveals roles
for RAB18 in eye development and organisation of the neuronal cytoskeleton.
Disease models & mechanisms 7: 711-22

Christoforidis S, McBride HM, Burgoyne RD, Zerial M (1999) The Rab5 effector
EEA1 is a core component of endosome docking. *Nature* 397: 621-5

Correa-Cerro LS, Wassif CA, Kratz L, Miller GF, Munasinghe JP, Grinberg A, Fliesler
SJ, Porter FD (2006) Development and characterization of a hypomorphic Smith-
Lemli-Opitz syndrome mouse model and efficacy of simvastatin therapy. *Hum Mol*
Genet 15: 839-51

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M (2014) Accurate proteome-
wide label-free quantification by delayed normalization and maximal peptide ratio
extraction, termed MaxLFQ. *Mol Cell Proteomics* 13: 2513-26

Dejgaard SY, Murshid A, Erman A, Kizilay O, Verbich D, Lodge R, Dejgaard K, Ly-
Hartig TB, Pepperkok R, Simpson JC, Presley JF (2008) Rab18 and Rab43 have
key roles in ER-Golgi trafficking. *J Cell Sci* 121: 2768-81

Eastman SW, Yassaee M, Bieniasz PD (2009) A role for ubiquitin ligases and
Spartin/SPG20 in lipid droplet turnover. *The Journal of cell biology* 184: 881-94

Feldmann A, Bekbulat F, Huesmann H, Ulbrich S, Tatzelt J, Behl C, Kern A (2017)
The RAB GTPase RAB18 modulates macroautophagy and proteostasis. *Biochem*
Biophys Res Commun 486: 738-743

Fliesler SJ, Peachey NS, Herron J, Hines KM, Weinstock NI, Ramachandra Rao S,
Xu L (2018) Prevention of Retinal Degeneration in a Rat Model of Smith-Lemli-Opitz
Syndrome. *Sci Rep* 8: 1286

Fukuda M, Kanno E, Ishibashi K, Itoh T (2008) Large scale screening for novel rab
effectors reveals unexpected broad Rab binding specificity. *Mol Cell Proteomics* 7:
1031-42

Ge L, Melville D, Zhang M, Schekman R (2013) The ER-Golgi intermediate
compartment is a key membrane source for the LC3 lipidation step of
autophagosome biogenesis. *Elife* 2: e00947

Gerondopoulos A, Bastos RN, Yoshimura S, Anderson R, Carpanini S, Aligianis I,
Handley MT, Barr FA (2014) Rab18 and a Rab18 GEF complex are required for
normal ER structure. *The Journal of cell biology* 205: 707-20

Gillingham AK, Bertram J, Begum F, Munro S (2019) In vivo identification of GTPase
interactors by mitochondrial relocalization and proximity biotinylation. *Elife* 8

Gillingham AK, Sinka R, Torres IL, Lilley KS, Munro S (2014) Toward a
comprehensive map of the effectors of rab GTPases. *Developmental cell* 31: 358-73

Handley M, Sheridan E (2018) RAB18 Deficiency. In *GeneReviews((R))*, Adam MP,
Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A (eds)
Seattle (WA):

Handley MT, Carpanini SM, Mali GR, Sidjanin DJ, Aligianis IA, Jackson IJ,
FitzPatrick DR (2015) Warburg Micro syndrome is caused by RAB18 deficiency or
dysregulation. *Open biology* 5: 150047

Hendershott MC, Vale RD (2014) Regulation of microtubule minus-end dynamics by
CAMSAPs and Patronin. *Proc Natl Acad Sci U S A* 111: 5860-5

Ho AC, Fung CW, Siu TS, Ma OC, Lam CW, Tam S, Wong VC (2014)
Lathosterolosis: a disorder of cholesterol biosynthesis resembling smith-lemli-opitz
syndrome. *JIMD Rep* 12: 129-34

Hueschen CL, Kenny SJ, Xu K, Dumont S (2017) NuMA recruits dynein activity to
microtubule minus-ends at mitosis. *Elife* 6

Im YJ, Raychaudhuri S, Prinz WA, Hurley JH (2005) Structural mechanism for sterol
sensing and transport by OSBP-related proteins. *Nature* 437: 154-8

Jia D, Zhang JS, Li F, Wang J, Deng Z, White MA, Osborne DG, Phillips-Krawczak
C, Gomez TS, Li H, Singla A, Burstein E, Billadeau DD, Rosen MK (2016) Structural
and mechanistic insights into regulation of the retromer coat by TBC1d5. *Nat
Commun* 7: 13305

Jimenez-Orgaz A, Kvainickas A, Nagele H, Denner J, Eimer S, Dengjel J, Steinberg
F (2018) Control of RAB7 activity and localization through the retromer-TBC1D5
complex enables RAB7-dependent mitophagy. *EMBO J* 37: 235-254

Jin J, Sison K, Li C, Tian R, Wnuk M, Sung HK, Jeansson M, Zhang C, Tucholska M,
Jones N, Kerjaschki D, Shibuya M, Fantus IG, Nagy A, Gerber HP, Ferrara N,
Pawson T, Quaggin SE (2012) Soluble FLT1 binds lipid microdomains in podocytes
to control cell morphology and glomerular barrier function. *Cell* 151: 384-99

Johansson M, Lehto M, Tanhuanpaa K, Cover TL, Olkkonen VM (2005) The
oxysterol-binding protein homologue ORP1L interacts with Rab7 and alters
functional properties of late endocytic compartments. *Mol Biol Cell* 16: 5480-92

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ (2015) The Phyre2 web
portal for protein modeling, prediction and analysis. *Nat Protoc* 10: 845-58

Kim DI, Jensen SC, Noble KA, Kc B, Roux KH, Motamedchaboki K, Roux KJ (2016)
An improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell* 27: 1188-
96

Korade Z, Xu L, Harrison FE, Ahsen R, Hart SE, Folkes OM, Mirnics K, Porter NA
(2014) Antioxidant supplementation ameliorates molecular deficits in Smith-Lemli-
Opitz syndrome. *Biol Psychiatry* 75: 215-22

Krakowiak PA, Wassif CA, Kratz L, Cozma D, Kovarova M, Harris G, Grinberg A,
Yang Y, Hunter AG, Tsokos M, Kelley RI, Porter FD (2003) Lathosterolosis: an

inborn error of human and murine cholesterol synthesis due to lathosterol 5-
desaturase deficiency. *Hum Mol Genet* 12: 1631-41

Kumar D, Golchoubian B, Belevich I, Jokitalo E, Schlaitz AL (2019) REEP3 and
REEP4 determine the tubular morphology of the endoplasmic reticulum during
mitosis. *Mol Biol Cell* 30: 1377-1389

Lamber EP, Siedenburg AC, Barr FA (2019) Rab regulation by GEFs and GAPs
during membrane traffic. *Curr Opin Cell Biol* 59: 34-39

Langemeyer L, Perz A, Kummel D, Ungermann C (2018) A guanine nucleotide
exchange factor (GEF) limits Rab GTPase-driven membrane fusion. *The Journal of*
biological chemistry 293: 731-739

Lees JA, Messa M, Sun EW, Wheeler H, Torta F, Wenk MR, De Camilli P, Reinisch
KM (2017) Lipid transport by TMEM24 at ER-plasma membrane contacts regulates
pulsatile insulin secretion. *Science* 355

Li C, Luo X, Zhao S, Siu GK, Liang Y, Chan HC, Satoh A, Yu SS (2017) COPI-
TRAPPII activates Rab18 and regulates its lipid droplet association. *EMBO J* 36:
441-457

Li D, Zhao YG, Li D, Zhao H, Huang J, Miao G, Feng D, Liu P, Li D, Zhang H (2019)
The ER-Localized Protein DFCP1 Modulates ER-Lipid Droplet Contact Formation.
Cell Rep 27: 343-358 e5

Liegel RP, Handley MT, Ronchetti A, Brown S, Langemeyer L, Linford A, Chang B,
Morris-Rosendahl DJ, Carpanini S, Posmyk R, Harthill V, Sheridan E, Abdel-Salam
GM, Terhal PA, Faravelli F, Accorsi P, Giordano L, Pinelli L, Hartmann B, Ebert AD
et al. (2013) Loss-of-function mutations in TBC1D20 cause cataracts and male
infertility in blind sterile mice and Warburg micro syndrome in humans. *Am J Hum*
Genet 93: 1001-14

Lind GE, Raiborg C, Danielsen SA, Rognum TO, Thiis-Evensen E, Hoff G,
Nesbakken A, Stenmark H, Lothe RA (2011) SPG20, a novel biomarker for early
detection of colorectal cancer, encodes a regulator of cytokinesis. *Oncogene* 30:
3967-78

Liu X, Salokas K, Tamene F, Jiu Y, Weldatsadik RG, Ohman T, Varjosalo M (2018)
An AP-MS- and BioID-compatible MAC-tag enables comprehensive mapping of
protein interactions and subcellular localizations. *Nat Commun* 9: 1188

Martin S, Driessen K, Nixon SJ, Zerial M, Parton RG (2005) Regulated localization of
Rab18 to lipid droplets: effects of lipolytic stimulation and inhibition of lipid droplet
catabolism. *The Journal of biological chemistry* 280: 42325-35

Nakamura H, Yamashita N, Kimura A, Kimura Y, Hirano H, Makihara H, Kawamoto
Y, Jitsuki-Takahashi A, Yonezaki K, Takase K, Miyazaki T, Nakamura F, Tanaka F,
Goshima Y (2016) Comprehensive behavioral study and proteomic analyses of
CRMP2-deficient mice. *Genes Cells* 21: 1059-1079

Nicholson JM, Macedo JC, Mattingly AJ, Wangsa D, Camps J, Lima V, Gomes AM,
Doria S, Ried T, Logarinho E, Cimini D (2015) Chromosome mis-segregation and
cytokinesis failure in trisomic human cells. *Elife* 4

Nowaczyk MJM, Wassif CA (1998) Smith-Lemli-Opitz Syndrome. In
GeneReviews((R)), Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH,
Stephens K, Amemiya A (eds) Seattle (WA):

Ong WY, Ng MP, Loke SY, Jin S, Wu YJ, Tanaka K, Wong PT (2013)
Comprehensive gene expression profiling reveals synergistic functional networks in
cerebral vessels after hypertension or hypercholesterolemia. *PloS one* 8: e68335

Ozeki S, Cheng J, Tauchi-Sato K, Hatano N, Taniguchi H, Fujimoto T (2005) Rab18
localizes to lipid droplets and induces their close apposition to the endoplasmic
reticulum-derived membrane. *J Cell Sci* 118: 2601-11

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ,
Inuganti A, Griss J, Mayer G, Eisenacher M, Perez E, Uszkoreit J, Pfeuffer J,
Sachsenberg T, Yilmaz S, Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF et al.
(2019) The PRIDE database and related tools and resources in 2019: improving
support for quantification data. *Nucleic Acids Res* 47: D442-D450

Petkovic M, Jemaiel A, Daste F, Specht CG, Izeddin I, Vorkel D, Verbavatz JM,
Darzacq X, Triller A, Pfenninger KH, Tareste D, Jackson CL, Galli T (2014) The
SNARE Sec22b has a non-fusogenic function in plasma membrane expansion. *Nat*
Cell Biol 16: 434-44

Platt FM, Wassif C, Colaco A, Dardis A, Lloyd-Evans E, Bembi B, Porter FD (2014)
Disorders of cholesterol metabolism and their unanticipated convergent mechanisms
of disease. *Annu Rev Genomics Hum Genet* 15: 173-94

Pulido MR, Diaz-Ruiz A, Jimenez-Gomez Y, Garcia-Navarro S, Gracia-Navarro F,
Tinahones F, Lopez-Miranda J, Fruhbeck G, Vazquez-Martinez R, Malagon MM
(2011) Rab18 dynamics in adipocytes in relation to lipogenesis, lipolysis and obesity.
PloS one 6: e22931

Raiborg C, Wenzel EM, Pedersen NM, Olsvik H, Schink KO, Schultz SW, Vietri M,
Nisi V, Bucci C, Brech A, Johansen T, Stenmark H (2015) Repeated ER-endosome
contacts promote endosome translocation and neurite outgrowth. *Nature* 520: 234-8

Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F (2013) Genome
engineering using the CRISPR-Cas9 system. *Nat Protoc* 8: 2281-2308

Richter KN, Revelo NH, Seitz KJ, Helm MS, Sarkar D, Saleeb RS, D'Este E, Eberle
J, Wagner E, Vogl C, Lazaro DF, Richter F, Coy-Vergara J, Coceano G, Boyden ES,
Duncan RR, Hell SW, Lauterbach MA, Lehnart SE, Moser T et al. (2018) Glyoxal as
an alternative fixative to formaldehyde in immunostaining and super-resolution
microscopy. *EMBO J* 37: 139-159

Rocha N, Kuijl C, van der Kant R, Janssen L, Houben D, Janssen H, Zwart W,
Neefjes J (2009) Cholesterol sensor ORP1L contacts the ER protein VAP to control
Rab7-RILP-p150 Glued and late endosome positioning. *The Journal of cell biology*
185: 1209-25

Rossi M, D'Armiento M, Parisi I, Ferrari P, Hall CM, Cervasio M, Rivasi F, Balli F,
Vecchione R, Corso G, Andria G, Parenti G (2007) Clinical phenotype of
lathosterolosis. *Am J Med Genet A* 143A: 2371-81

Roux KJ, Kim DI, Burke B, May DG (2018) BioID: A Screen for Protein-Protein
Interactions. *Curr Protoc Protein Sci* 91: 19 23 1-19 23 15

Roux KJ, Kim DI, Raida M, Burke B (2012) A promiscuous biotin ligase fusion
protein identifies proximal and interacting proteins in mammalian cells. *The Journal*
of cell biology 196: 801-10

Schlaitz AL, Thompson J, Wong CC, Yates JR, 3rd, Heald R (2013) REEP3/4
ensure endoplasmic reticulum clearance from metaphase chromatin and proper
nuclear envelope architecture. *Developmental cell* 26: 315-23

Seaman MN, Harbour ME, Tattersall D, Read E, Bright N (2009) Membrane
recruitment of the cargo-selective retromer subcomplex is catalysed by the small
GTPase Rab7 and inhibited by the Rab-GAP TBC1D5. *J Cell Sci* 122: 2371-82

Sidjanin DJ, Park AK, Ronchetti A, Martins J, Jackson WT (2016) TBC1D20
mediates autophagy as a key regulator of autophagosome maturation. *Autophagy*: 1-
17

Silve S, Dupuy PH, Labit-Lebouteiller C, Kaghad M, Chalon P, Rahier A, Taton M,
Lupker J, Shire D, Loison G (1996) Emopamil-binding protein, a mammalian protein
that binds a series of structurally diverse neuroprotective agents, exhibits delta8-
delta7 sterol isomerase activity in yeast. *The Journal of biological chemistry* 271:
22434-40

Sobajima T, Yoshimura SI, Maeda T, Miyata H, Miyoshi E, Harada A (2018) The
Rab11-binding protein RELCH/KIAA1468 controls intracellular cholesterol
distribution. *The Journal of cell biology* 217: 1777-1796

Spang A (2012) The DSL1 complex: the smallest but not the least CATCHR. *Traffic*
13: 908-13

Spang N, Feldmann A, Huesmann H, Bekbulat F, Schmitt V, Hiebel C, Koziollek-
Drechsler I, Clement AM, Moosmann B, Jung J, Behrends C, Dikic I, Kern A, Behl C
(2014) RAB3GAP1 and RAB3GAP2 modulate basal and rapamycin-induced
autophagy. *Autophagy* 10: 2297-309

Svoboda MD, Christie JM, Eroglu Y, Freeman KA, Steiner RD (2012) Treatment of
Smith-Lemli-Opitz syndrome and other sterol disorders. *Am J Med Genet C Semin*
Med Genet 160C: 285-94

Tagaya M, Arasaki K, Inoue H, Kimura H (2014) Moonlighting functions of the NRZ
(mammalian Dsl1) complex. *Front Cell Dev Biol* 2: 25

Tinti M, Johnson C, Toth R, Ferrier DE, Mackintosh C (2012) Evolution of signal
multiplexing by 14-3-3-binding 2R-ohnologue protein families in the vertebrates.
Open biology 2: 120103

Urnavicius L, Zhang K, Diamant AG, Motz C, Schlager MA, Yu M, Patel NA,
Robinson CV, Carter AP (2015) The structure of the dynactin complex and its
interaction with dynein. *Science* 347: 1441-1446

Vazquez-Martinez R, Cruz-Garcia D, Duran-Prado M, Peinado JR, Castano JP,
Malagon MM (2007) Rab18 inhibits secretory activity in neuroendocrine cells by
interacting with secretory granules. *Traffic* 8: 867-82

Wang H, Ma Q, Qi Y, Dong J, Du X, Rae J, Wang J, Wu WF, Brown AJ, Parton RG,
Wu JW, Yang H (2019) ORP2 Delivers Cholesterol to the Plasma Membrane in
Exchange for Phosphatidylinositol 4, 5-Bisphosphate (PI(4,5)P2). *Mol Cell* 73: 458-
473 e7

Wang T, Li L, Hong W (2017) SNARE proteins in membrane trafficking. *Traffic* 18:
767-775

Wassif CA, Kratz L, Sparks SE, Wheeler C, Bianconi S, Gropman A, Calis KA,
Kelley RI, Tierney E, Porter FD (2017) A placebo-controlled trial of simvastatin
therapy in Smith-Lemli-Opitz syndrome. *Genet Med* 19: 297-305

Whitfield ST, Burston HE, Bean BD, Raghuram N, Maldonado-Baez L, Davey M,
Wendland B, Conibear E (2016) The alternate AP-1 adaptor subunit Apm2 interacts
with the Mil1 regulatory protein and confers differential cargo sorting. *Mol Biol Cell*
27: 588-98

Wickner W (2010) Membrane fusion: five lipids, four SNAREs, three chaperones,
two nucleotides, and a Rab, all dancing in a ring on yeast vacuoles. *Annu Rev Cell*
Dev Biol 26: 115-36

Williams C, Choudhury R, McKenzie E, Lowe M (2007) Targeting of the type II
inositol polyphosphate 5-phosphatase INPP5B to the early secretory pathway. *J Cell*
Sci 120: 3941-51

Wu H, Carvalho P, Voeltz GK (2018) Here, there, and everywhere: The importance
of ER membrane contact sites. *Science* 361

Xu D, Li Y, Wu L, Li Y, Zhao D, Yu J, Huang T, Ferguson C, Parton RG, Yang H, Li
P (2018) Rab18 promotes lipid droplet (LD) growth by tethering the ER to LDs
through SNARE and NRZ interactions. *The Journal of cell biology* 217: 975-995

Yamamoto Y, Yurugi C, Sakisaka T (2017) The number of the C-terminal
transmembrane domains has the potency to specify subcellular localization of
Sec22c. *Biochem Biophys Res Commun* 487: 388-395

Zhang T, Wong SH, Tang BL, Xu Y, Hong W (1999) Morphological and functional
association of Sec22b/ERS-24 with the pre-Golgi intermediate compartment. *Mol*
Biol Cell 10: 435-53

Zhao K, Ridgway ND (2017) Oxysterol-Binding Protein-Related Protein 1L Regulates
Cholesterol Egress from the Endo-Lysosomal System. *Cell Rep* 19: 1807-1818

Zhao L, Imperiale MJ (2017) Identification of Rab18 as an Essential Host Factor for
BK Polyomavirus Infection Using a Whole-Genome RNA Interference Screen.
mSphere 2

Zhen Y, Stenmark H (2015) Cellular functions of Rab GTPases at a glance. *J Cell*
Sci 128: 3171-6

1426

1427

A

BirA*-RAB18(Ser22Asn)

BirA*-RAB18(wild-type)

wild-type HEK293 cells

wild-type HEK293 cells

B

Protein	Ratio	Orthologue PPI (Gillingham et al., 2014)	Additional evidence	
SCFD2	0.00	Slh	Gillingham et al., 2019	RAB3GAP-dependent association in HeLa cells
ZW10	0.00	mit(1)15	Xu et al., 2018; Gillingham et al., 2019	
NBAS	0.00	rod	Xu et al., 2018; Gillingham et al., 2019	
CAMSAP1	0.04			
FAM134B	0.36			
RINT1	0.41	CG8605	Xu et al., 2018	
RAB3GAP2	1.58	rab3-GAP	Gerondopoulos et al., 2014	
RAB3GAP1	1.59	CG31935	Gerondopoulos et al., 2014	
TBC1D5	0.00	CG8449	Gillingham et al., 2019	TRAPPII-dependent association in HeLa cells
c15orf38	0.00			
SEC23IP	0.00			
ATP6AP2	0.29			
GORASP2	0.36			
GIGYF2	0.41			
TPR	0.41			
USP15	0.00			Present in HeLa dataset (n≥2)
TMPO	0.00			
ARFGAP3	0.20			
ARFGAP2	0.31			
SLK	0.34			
NUP153	0.39			
ATG2B	0.43			
SCFD1	0.45	Slh	Gillingham et al., 2019 Xu et al., 2018	
USE1	0.45			
TMX1	0.45			
CORO1B	0.47			
PREB	0.47			
SLC25A4	0.00			Absent in HeLa dataset (n≥2)
LUZP1	0.00			
DPYSL2	0.34		Nakamura et al., 2016	

1428

Figure 1. Nucleotide-binding-dependent RAB18-associations in HEK293 cells.

(A) Schematic to show BirA*-RAB18(Ser22Asn):BirA*-RAB18(WT) comparison.

RAB18 crystal structure from RCSB PDB code 1X3S. (B) Table to show putative

nucleotide-binding-dependent RAB18-associations with BirA*-

RAB18(Ser22Asn):BirA*-RAB18(WT) association ratios <0.5. Proteins orthologous

to interactors identified by Gillingham et al. (2014) are indicated. Previous studies

providing supporting evidence for interactions are indicated. Proteins are grouped

according to their attributes in the HeLa cell dataset (Figure 2 and Table S2).

Association ratios were derived individually following normalization by total spectral

counts per condition. The full dataset is provided in table S1.

1439

A

BirA*-RAB18(wild-type)   BirA*-RAB18(wild-type)

GTP   GTP
GAP   GAP
GEF
GDP   GDP

BirA*   BirA*

GEF-null HeLa cells   wild-type HeLa cells

B

RAB3GAP2-null / RAB3GAP1-null $R^2=0.99$

C

All (457)
TRAPPII-dependent (133)
RAB3GAP-dependent (25)

D

Protein	n	Ratio	Orthologue PPI (Gillingham et al., 2014)	Additional evidence	
CAMSAP1	3	0.26			
REEP4	3	0.35		Tinti et al., 2012*	Microtubule-associated/Membrane remodelling
BICD2	2	0.25	BicD	Gillingham et al., 2019	
SPG20	**2**	**0.42**	**CG12001**	**This study (Figure 4)**	
FAM134B	2	0.46			
ZW10	3	0.02	mit(1)15	Xu et al., 2018; Gillingham et al., 2019	
RINT1	3	0.16	CG8605	Xu et al., 2018	
NBAS	3	0.17	rod	Xu et al., 2018; Gillingham et al., 2019	
SCFD2	3	0.41	Slh	Gillingham et al., 2019	Membrane tethering/docking
SEC22A	**3**	**0.46**		**This study (Figure 5)**	
WFS1	2	0.00			
BNIP1	2	0.36		Xu et al., 2018	
STX18	2	0.40	Syx18	Xu et al., 2018	
TMCO4	**3**	**0.06**		**This study (Figure 6)**	
OSBPL2	**3**	**0.35**		**This study (Figure 7)**	
EBP	**3**	**0.38**		**This study (Figure 7)**	Lipid modifying/mobilising
INPP5B	**2**	**0.00**		**This study (Figure 7)**	
C2CD2L	2	0.22			
C2CD2	2	0.36			
RAB3GAP2	3	0.00	rab3-GAP	Gerondopoulos et al., 2014	
RAB3GAP1	3	0.01	CG31935	Gerondopoulos et al., 2014	
SSR3	3	0.49			Other
MFHAS1	2	0.00	lrrk		
TRIM13	2	0.29			
SCARA3	2	0.42			

1440

Figure 2. RAB3GAP-dependent RAB18-associations in HeLa cells. (A)

Schematic to show comparison of BirA*-RAB18(WT) in wild-type and guanine

nucleotide exchange factor (GEF)-null cells. RAB18 crystal structure from RCSB

PDB code 1X3S. (B) Plot to show correlation between non-zero LFQ intensities of

individual proteins identified in samples purified from RAB3GAP1- and RAB3GAP2-

null cells. (C) Venn diagram illustrating all RAB18-associations, TRAPPII-dependent

interactions (TRAPPC9-null:wild-type association ratios <0.5) and RAB3GAP-

dependent associations (RAB3GAP1/2-null:wild-type association ratios <0.5). (D)

Table to show putative RAB18-associations with RAB3GAP1/2-null:wild-type

association ratios <0.5. Proteins orthologous to interactors identified by Gillingham et

al. (2014) are indicated. Previous studies providing supporting evidence for

interactions are indicated. Proteins are grouped according to their reported functions.

Association ratios were derived individually following normalization by RAB18 LFQ

intensity in each replicate experiment. The full dataset is provided in table S2.

1455

1456

1457

Figure 3. Initial validation of RAB3GAP-dependent RAB18-associations in

HeLa cells. (A) Western blotting of samples purified from wild-type and RAB3GAP1-

null cells in an independent BioID experiment. Levels of selected proteins are

consistent with association ratios in Figure 1C. (B) Table to show diseases with

features overlapping those of Warburg Micro syndrome linked to genes encoding

putative effector proteins or their homologues. (C) Comparative fluorescence

microscopy of selected RAB18-associated proteins in wild-type and RAB18-null

HeLa cells. Cells of different genotypes were labelled with CellTrace-Violet and

CellTrace-Far Red reagents, corresponding to blue and magenta channels

respectively. Cells were stained with antibodies against indicated proteins in green

channel panels. (D) Comparative fluorescence microscopy of selected RAB18-

associated proteins in lipid-loaded wild-type and RAB18-null HeLa cells. Cells were

stained as above but were treated for 15 hours with 200μM oleic acid, 1μg/ml

BODIPY-558/568-C12 (Red channel) prior to fixation.

1472

1473

Figure 4. Levels of SPG20 are significantly reduced in RAB18-null and

TBC1D20-null HeLa and RPE1 cells. (A) Comparative fluorescence microscopy of

SPG20 in wild-type and mutant HeLa cell lines. Wild-type and mutant cells of the

indicated genotypes were labelled with CellTrace-Far Red and CellTrace-Violet and

reagents respectively (magenta and blue channels). Cells were stained with an

antibody against SPG20 (green channel) (B) Quantification of SPG20 fluorescence

in wild-type cells by direct comparison with SPG20-null cells. (C) Quantification of

SPG20 fluorescence (%wt) in cells of different genotypes. Data were derived from

analysis of at least 18 frames – each containing >5 wild-type and >5 mutant cells –

per genotype. $*p<0.001$. (D) LFQ intensities for SPG20 (Q8N0X7) and β-Actin

(P60709) in whole-cell lysates of RPE1 cells of the indicated genotypes. n=3;

$*p<0.05$ following FDR correction. Full dataset provided in table S3. Error bars

represent s.e.m. (E) Confocal micrograph to show localization of exogenous

mEmerald-SPG20 (Green) and mCherry-RAB18 (Red) in HeLa cells. (F) Confocal

micrographs to show localization of exogenous mEmerald-SPG20 (Green; left

panels) and EGFP-RAB18 (Green; right panels) in HeLa cells loaded with 200μM

oleic acid, 1μg/ml BODIPY-558/568-C12 (Red channel). Wild-type and mutant cells

of the indicated genotypes were labelled with CellTrace-Violet and CellTrace-Far

Red reagents respectively (magenta and blue channels).

1493

1494

Figure 5. SEC22A associates with RAB18 and influences LD morphology. (A)
Confocal micrograph to show overlapping localization of exogenous mEmerald-
SEC22A (Green) and mCherry-ER (Red) in HeLa cells. (B) RAB18 LFQ intensities
from a reciprocal BioID experiment showing a reduced association between
BioID2(Gly40Ser)-SEC22A and endogenous RAB18 in RAB3GAP-null compared to
wild-type HeLa cells. Data were adjusted to account for non-specific binding of
RAB18 to beads and normalized by SEC22A LFQ intensities in each replicate
experiment. Error bars represent s.e.m. Data for other BioID2(Gly40Ser)-SEC22A-
associated proteins are provided in table S5. (C) Bar graphs to show effects of
ZW10, NBAS and SEC22A knockdowns on lipid droplet number and diameter.
siRNA-treated IHH cells were loaded with 200nM BSA-conjugated oleate, fixed and
stained with BODIPY and DAPI, and imaged. Images were analysed using ImageJ.
Data are derived from measurements from >100 cells/condition and are
representative of three independent experiments. Error bars represent SD. *p<0.001

A

B

C

1509

Figure 6. mCherry-RAB18 recruits TMCO4-EGFP to the ER membrane in a

RAB3GAP-dependent manner. (A) Confocal micrographs to show diffuse

localization of exogenous TMCO4-EGFP (Green) compared to mCherry-ER (Red)

and overlapping localization of exogenous EGFP-RAB18 (Green) and mCherry-ER

in HeLa cells. (B) Confocal micrographs to show localization of exogenous mCherry-

RAB18 and TMCO4-EGFP in wild-type cells and in mutant cells of different

genotypes. Wild-type and mutant cells of the indicated genotypes were labelled with

CellTrace-Violet and CellTrace-Far Red reagents respectively (magenta and blue

channels). (C) Immunoprecipitation of exogenous HA-RAB18 from HeLa cells of

different genotypes. Cells were transfected with the indicated constructs and lysed

24 hours post-transfection. Anti-HA immunoprecipitates and input samples were

subjected to SDS-PAGE and immunostaining for HA and GFP.

1522

Figure 7. RAB18 is involved in the mobilization and biosynthesis of

cholesterol. (A) Plots to show cholesteryl ester (CE) loading and efflux. CHO cells,

stably expressing RAB18(WT), RAB18(Gln67Leu) and RAB18(Ser22Asn), were

incubated with [^{14}C]-oleate, for 24 hours, in the presence of lipoprotein depleted

serum (LPDS)(Left panel) or FBS (Right panel). Following lipid extraction, thin layer

chromatography (TLC) was used to separate CE, and radioactivity was measured by

scintillation counting. Measurements were made at t=0 and at 4 and 8 hours

following the addition of 50µg/ml high density lipoprotein (HDL) to the cells. (B) Bar

graph to show cholesterol efflux. The CHO cells were incubated with [^{3}H]-

cholesterol, for 24 hours, in the presence of FBS. After washing, they were then

incubated with 25µg/ml apolipoprotein A-I for 5 hours. The quantity of [^{3}H]-

cholesterol in the media is shown as a percentage of the total cellular radioactivity

(mean+/-SD). (C) Bar graph to show sterols profile in wild-type and RAB18-null HeLa

cells. Cells were grown in media supplemented with LPDS for 48 hours. Extracted

sterols were analysed by gas chromatography-mass spectrometry (GC-MS). %

Sterol was calculated as a proportion of total quantified sterols, excluding

cholesterol, following normalization to a 5α-cholestane internal standard. n=3 (D)

Model for lathosterol mobilization mediated by RAB18. ORP2 binds PI(4,5)P$_2$ on an

apposed membrane. RAB18 interacts with ORP2 and INPP5B promoting the

hydrolysis of PI(4,5)P$_2$ to PI(4)P and maintaining a PI(4,5)P$_2$ concentration gradient.

RAB18 coordinates the production of lathosterol by EBP and its subsequent

mobilization by ORP2. (E) Bar graph to show incorporation of [^{3}H]-acetate into

cholesterol in a panel of HeLa cell lines. Cells were grown in LPDS-media for 24

hours then incubated with 10µCi/well [^{3}H]-acetate for 24 hours. TLC was used to

separate free cholesterol and radioactivity was quantified by scintillation counting

(n=4; mean±SD). (F) Bar graph to show incorporation of [^{3}H]-mevalonate into

cholesterol in the cell lines. Cells were grown LPDS-media for 24 hours, then

incubated with 5µCi/well [^{3}H]-mevalonate for 24 hours. TLC was used to separate

free cholesterol and radioactivity was quantified by scintillation counting (n=4;

mean±SD). (G) Still image from a molecular dynamics model of the ORP2 OSBP-

related domain (ORD) complexed with cholesterol. Position of ORP2 Tyr110

(NP_653081) is shown. (H) Still image from a molecular dynamics model of the

ORP2-ORD complexed with lathosterol. Positions of ORP2 Tyr110 and proximal

water molecules are shown. *p<0.01, **p<0.001, ***p<0.0005, ****p<0.005.