TITLE

Comparative proximity biotinylation implicates RAB18 in cholesterol mobilization and

biosynthesis

RUNNING TITLE

GEF-dependent RAB18 interactions

AUTHORS

Robert S. Kiss*[1], Jarred Chicoine[1], Youssef Khalil[2], Robert Sladek[1], He Chen[1],

Alessandro Pisaturo[1], Cyril Martin[1], Jessica D. Dale[3], Tegan A. Brudenell[3], Archith

Kamath[4,5], Emanuele Paci[6], Peter Clayton[2], Jimi C. Wills[4], Alex von Kriegsheim[4],

Tommy Nilsson[1], Eamonn Sheridan[3], Mark T. Handley*[3,7]

AFFILIATIONS

[1]Research Institute of the McGill University Health Centre

1001 boul Decarie

Glen Site Block E

Montreal, QC

H4A 3J1

Canada

[2] Genetics and Genomic Medicine

Great Ormond Street Institute of Child Health

University College London

30 Guilford Street

London

WC1N 1EH

United Kingdom

[3]Leeds Institute of Medical Research

St James's University Hospital

Leeds

LS9 7TF

United Kingdom

[4]Cancer Research UK Edinburgh Centre

MRC Institute of Genetics & Molecular Medicine

The University of Edinburgh

Western General Hospital

Edinburgh

EH4 2XR

United Kingdom

[5]Medical Sciences Division

University of Oxford

Oxford

OX3 9DU

United Kingdom

[6]Astbury Centre for Structural Molecular Biology

University of Leeds

Leeds

LS2 9JT

United Kingdom

[7]Faculty of Biological Sciences

University of Leeds

Leeds

LS2 9JT

United Kingdom.

*authors contributed equally

CORRESPONDING AUTHORS

Robert S. Kiss: robert.kiss@mcgill.ca

Mark T. Handley: m.handley@leeds.ac.uk

KEYWORDS

RAB18, BioID, Cholesterol, Lathosterol, EBP, ORP2

SUMMARY STATEMENT

We used proximity biotinylation together with guanine nucleotide exchange factor

(GEF)-null cell lines to discriminate functional RAB18-interactions. Our data suggest

that RAB18 mediates lathosterol mobilization and cholesterol biosynthesis.

ABSTRACT

Loss of functional RAB18 causes the autosomal recessive condition Warburg Micro

syndrome. To better understand this disease, we used proximity biotinylation to

generate an inventory of potential RAB18 effectors. A restricted set of 25 RAB18-

interactions were dependent on the binary RAB3GAP1-RAB3GAP2 RAB18-guanine

nucleotide exchange factor (GEF) complex. Consistent with a role for RAB18 in

regulating membrane contact sites (MCSs), interactors included groups of

microtubule/membrane-remodelling proteins, membrane-tethering and docking

proteins, and lipid-modifying/transporting proteins. We provide evidence validating

novel interactions with SEC22A and TMCO4. We also provide functional evidence

that RAB18 links the Δ8-Δ7 sterol isomerase emopamil binding protein (EBP) to a

molecular machinery mobilizing the products of EBP-catalysis. The cholesterol

precursor lathosterol accumulates in RAB18-null cells, and *de novo* cholesterol

biosynthesis is reduced. Our data demonstrate that GEF-dependent Rab-interactions

are highly amenable to interrogation by proximity biotinylation and suggest that Micro

syndrome is a cholesterol biosynthesis disorder.

INTRODUCTION

Rab Proteins are a large subfamily of small GTPases with discrete roles in coordinating membrane trafficking (Zhen & Stenmark, 2015). Like other small GTPases, they adopt different conformations and enter into different protein-protein interactions according to whether they are GDP-, or GTP-bound. Although they possess some intrinsic GTP-hydrolysis activity, their *in vivo* nucleotide-bound state is tightly governed in cells by two classes of regulatory proteins. Guanine-nucleotide exchange factors (GEFs) catalyse the exchange of bound GDP for GTP while GTPase-activating proteins (GAPs) promote the hydrolysis of bound GTP to GDP (Barr & Lambright, 2010, Lamber et al., 2019).

Biallelic loss-of-function variants in *RAB18*, *RAB3GAP1*, *RAB3GAP2*, or *TBC1D20*, cause the autosomal recessive condition Warburg Micro syndrome (Aligianis et al., 2005, Bem et al., 2011, Borck et al., 2011, Handley & Sheridan, 2018, Liegel et al., 2013)(MIMs 600118, 614222, 614225, 615663, 212720). *RAB3GAP1* and *RAB3GAP2* encode subunits of the binary RAB18-GEF complex, 'RAB3GAP', whereas *TBC1D20* encodes a RAB18-GAP (Gerondopoulos et al., 2014, Handley et al., 2015). Thus, the same pathology is produced when functional RAB18 is absent or when its normal regulation is disrupted. However, it is unclear how RAB18 dysfunction contributes to disease pathology at a molecular level.

Rab proteins fulfil their roles by way of protein-protein interactions with interacting partners termed 'effectors'. The identification of these proteins can therefore provide insight into these roles. However, biochemical identification of Rab effectors is challenging; Rab-effector interactions are usually GTP-dependent and are often highly transient. Immunoprecipitation, affinity purification and yeast-2-hybrid approaches have each been used, but may be more or less effective depending on the Rab isoform studied (Christoforidis et al., 1999, Fukuda et al., 2008).

One newer approach is 'BioID' proximity biotinylation utilizing Rab proteins fused to mutant forms of the biotin ligase BirA. The Rab fusion protein biotinylates proximal proteins which are then purified on streptavidin and identified through mass spectrometry (Gillingham et al., 2019, Liu et al., 2018, Roux et al., 2012). Biotin labelling occurs in a relatively physiological context and prospective effectors can be purified under high stringency conditions. However, a drawback of the technique is

that it does not distinguish between close associations resulting from functional

protein-protein interactions and those resulting from overlapping localizations.

To discriminate functional RAB18 interactions, we compared BirA*-RAB18 labelling

of protein in wild-type HeLa cells to that in cells in which RAB18-GEF activity was

disrupted with CRISPR. Known and novel effectors were more strongly labelled in

the wild-type cells. In particular, 25 RAB18-interactions were RAB3GAP-dependent.

These proteins comprised several groups. Proteins within each group were clearly

interrelated through involvement in connected biological processes. Moreover, gene-

disease associations within the set included multiple overlapping phenotypes. We

present additional evidence for several examples of RAB3GAP-dependent RAB18

interactions: SPG20/SPART; SEC22A; TMCO4; EBP; ORP2/OSBPL2 and INPP5B.

Our data elaborate an existing model suggesting that RAB18 effectors act

collectively in lipid transfer at membrane contact sites (Xu et al., 2018). We identify

multiple proteins already implicated in the establishment and maintenance of

membrane-contacts. Importantly, we also identify novel RAB18-interactors involved

in lipid biosynthesis, mobilization and metabolism. We find that RAB18 appears to

coordinate generation of the cholesterol precursor lathosterol by the $\Delta 8$-$\Delta 7$ sterol

isomerase enzyme EBP, and its subsequent mobilization by the lipid transfer protein

(LTP) ORP2/OSBPL2. Consistently, cholesterol biosynthesis is impaired in model

cell lines lacking RAB18 or its regulators. Taken together, these findings suggest that

Micro syndrome is a cholesterol biosynthesis disorder. Further, that therapeutic

interventions might be feasible in this disease.

RESULTS

<u>An inventory of RAB18-GEF-dependent RAB18-associated proteins in HeLa cells</u>

We first used CRISPR to generate a panel of clonal, otherwise isogenic, HeLa cell

lines null for RAB18 and a number of its regulators (see Figure S1). We then carried

out proximity labelling using transient expression of the same exogenous BirA*-

RAB18 construct in RAB3GAP1-, RAB3GAP2- and TRAPPC9-null cell lines and in

wild-type cells (Figure 1A). RAB3GAP1 and RAB3GAP2 are each essential subunits

of a binary RAB18-GEF complex (Gerondopoulos et al., 2014). TRAPPC9 is

reported to be essential for the RAB18-GEF activity of a different GEF, the

multisubunit TRAPPII complex (Li et al., 2017).

Proximity-labelling, affinity purification and mass spectrometry of biotinylated

proteins were carried out essentially as previously described (Roux et al., 2018,

Roux et al., 2012). Prior to mass-spec analysis, samples from each of the

streptavidin pull-downs were subjected to Western blotting to ensure comparable

BirA*-RAB18 expression (Figure S2A). Label-free quantitative proteomics analyses

were used to calculate 'LFQ intensities' for each RAB18-associated protein (Cox et

al., 2014). These were then normalized in each experiment according to the quantity

of RAB18 found in each sample. Samples from three independent experiments were

analysed. Pull-downs from untransfected biotin-treated cells were used as controls.

After filtering the data to remove known mass-spec contaminants and any protein

identified at a high level in control samples, a total of 584, 483 and 506 RAB18-

associated proteins were identified in each experiment. A total of 457 proteins were

present in two or more of the replicate experiments (see Table S1).

Different Rab-GEF complexes may operate in distinct subcellular localizations and

coordinate associations with different effectors (Carney et al., 2006). Therefore, we

assessed whether non-zero intensities for each RAB18-associated protein correlated

between samples (Figure 1B, Figure S2B). Very strong correlations between protein

intensities from RAB3GAP1- and RAB3GAP2-null cells indicated that loss of either

protein had a functionally equivalent effect (R^2=0.99, Figure 1B). In contrast,

intensities from RAB3GAP1/2- and TRAPPC9-null cells were much more poorly

correlated (R^2=0.73, R^2=0.74, Figure S2B). We therefore considered RAB3GAP- and

TRAPPC9-dependent RAB18-interactions separately. Intensities from wild-type and
RAB3GAP-null samples correlated with an R^2=0.87, but a number of proteins
showed reduced intensities in the RAB3GAP-null samples (Figure 1C).

GEF activity promotes Rab GTP binding and this is usually necessary for effector
interactions. We therefore reasoned that levels of true effector proteins would be
reduced in samples from GEF-null cells as compared to those from wild-type cells.
We calculated GEF-null:wild-type intensity ratios for each RAB18-associated protein
(Table S1). Only 25 proteins showed a RAB3GAP-null:wild-type ratio <0.5 (Figure
1D, Table 1, Table S1). 133 proteins showed a TRAPPII-null:wild-type intensity ratio
<0.5 (Figure 1D, Table S1). There was only limited overlap between RAB3GAP- and
TRAPPC9-dependent associations (Figure 1D).

The most comprehensive annotation of candidate RAB18 effectors thus far was
made in the 2014 paper by Gillingham et al., which utilized an affinity purification-
mass spectrometry (AP-MS) approach and the *Drosophila* RAB18 orthologue
(Gillingham et al., 2014). In that study, a total of 456 proteins were identified as
interacting with RAB18. However, only 14 of these were well represented in terms of
spectral counts, exhibited low non-specific binding to GST/sepharose and showed
low binding to other Rab protein isoforms. We took these 14 proteins as the most
plausible physiological RAB18 interactors and searched for these in our datasets.

Orthologues of 11 of the 14 putative RAB18-interacting proteins identified by
Gillingham et al. were identified in our combined dataset. 10/14 were among the 25
RAB3GAP-dependent associations (Table 1). 2/14 were among the TRAPPII-
dependent associations (Table S1).

For initial validation of our dataset, we carried out an additional independent BioID
experiment with wild-type and RAB3GAP1-null cells and subjected the resulting
samples to Western blotting for selected RAB18-associated proteins (Figure 1E). As
with the mass spectrometry, these proteins showed either complete (RAB3GAP1,
RAB3GAP2, ZW10) or partial (SPG20, STX18) dependence on RAB3GAP for their
RAB18 association.

We further validated our approach with additional proximity biotinylation experiments
in HEK293 cells. We used cells stably expressing BirA*-tagged RAB18 fusions
incorporating wild-type RAB18, GTP-hydrolysis deficient RAB18(Gln67Leu), or

nucleotide-binding deficient RAB18(Ser22Asn) mutants (Figure S3A-B). A total of 98

proteins were identified as associating with RAB18 across all samples (Table S2).

Gln67Leu:wild-type intensity ratios for known RAB18-interactors ranged from 0.1-

1.49 indicating that RAB18 associations were altered by the Gln67Leu variant, but

not predictably so. In contrast, Ser22Asn:wild-type intensity ratios were <0.5 for the

majority of these proteins. 28 nucleotide-binding-dependent RAB18 associations

included 6 of the RAB3GAP-dependent associations and 7 of the TRAPPII-

dependent associations seen in the HeLa cells (Figure S3C). These data confirm

that the loss of GEFs has similar effects on RAB18-interactions to direct loss of

nucleotide binding. In addition, they support the differing regulation of specific

RAB18-interactions by different GEFs.

<u>Validation screening of RAB3GAP-dependent RAB18 associations reveals reduced</u>

<u>levels of SPG20 in RAB18-null and TBC1D20-null cells</u>

Our continued study focused on the 25 RAB3GAP-dependent RAB18 associations

identified in HeLa cells. Encouragingly, these appeared to share interconnected

functions and fell into discrete groups (Table 1). Furthermore, genes encoding 11 of

the 25 proteins or their homologues are associated with inherited diseases that

share features with Micro syndrome (Table 2).

Given the suggestive convergences in protein function and gene-disease-

associations, we examined the subcellular localizations of 12 putative effectors for

which antibodies were available (Figure 2A-B). To determine whether the localization

of these proteins was altered in cells lacking RAB18, we analysed wild type and

RAB18-null lines in each case. In order to directly compare cells of different

genotypes under otherwise identical conditions, we labelled them with CellTrace-

Violet and CellTrace-Far Red reagents before seeding, immunostaining and imaging

them together. Since RAB18 can localize to lipid droplets (LDs), we analysed both

untreated cells (Figure 2A) and cells loaded with oleic acid and labelled with

BODIPY-558/568-C12 (Figure 2B).

The putative effector proteins showed various staining patterns. These ranged from

staining that was enriched at the perinuclear region of cells, to staining that appeared

reticular, to staining that appeared more diffuse. Staining patterns were similar in the

HeLa cells and also in RPE1 cells generated to provide biological replicates (Figure

S4A). Each pattern was compatible with the known localization of RAB18, which is

distributed between *cis*-Golgi, ER and cytosolic compartments (Handley et al., 2015).

In lipid-loaded cells, localizations of proteins with reticular staining patterns

overlapped with LDs but they did not obviously shift to adopt a LD localization. Two

proteins that showed diffuse staining patterns in untreated cells - ZW10 and SPG20 -

appeared enriched in the vicinity of LDs (Figure 2B, bottom right panels).

We saw no evidence for dramatic changes in protein localizations in RAB18-null

cells as compared to their wild-type counterparts. Fluorescence intensities in RAB18-

null and wild-type cells were also generally similar, except in the case of staining for

SPG20, which appeared lower in RAB18-null HeLa cells than in wild-type cells

(Figure 2A, bottom right panels).

To confirm the reduction in levels of SPG20 we observed in RAB18-null HeLa cells

and to determine the effects of other genotypes, we used quantitative fluorescence

microscopy (Figure 2C). To establish SPG20 antibody specificity we first analysed

SPG20-null cells (Figure 2D, left panels). Measured fluorescence intensity of these

cells also provided a baseline level, above which fluorescence levels are proportional

to levels of SPG20 (Figure 2E). In RAB18-null cells, SPG20 fluorescence was

reduced to 67.16±3.77% (p<0.001) of that in wild-type cells (Figure 2F). Loss of the

RAB18-GEF subunits RAB3GAP1 or RAB3GAP2 did not significantly affect levels of

SPG20, whereas loss of the RAB18-GAP TBC1D20 led to a reduction comparable to

that in RAB18-null cells (57.48%±2.57, p<0.00005)(Figure 2F).

We analysed levels of SPG20 in the corresponding panel of RPE1 cell lines using

LFQP analysis of whole cell lysates (Figure S4B, Table S3). As in the HeLa cells,

levels of SPG20 were significantly reduced in RAB18- and TBC1D20-null RPE1 cells

compared to wild-type controls (p<0.05 following FDR correction), but not in the

other genotypes tested. These data suggest that reduced SPG20 levels result from

specific genotypes and are not the result of clonal variation. A comparison between

LFQP data from wild-type and TBC1D20-null RPE1 and HeLa cells (Tables S3 and

S4) showed limited overlap between differentially expressed proteins. This indicates

that reduced SPG20 levels are unlikely to have resulted from widespread

dysregulation of proteostasis. The RAB18-SPG20 interaction has been previously

reported (Gillingham et al., 2014), and our findings (above) provide further support
for a physiological relationship between these proteins.

<u>SEC22A associates with RAB18 and its knockdown causes altered LD morphology</u>

The most studied group of RAB18 effector proteins to date are the tethering factors
that comprise the NRZ/Dsl complex (ZW10, NBAS and RINT1), and the ER SNARE
proteins that comprise the Syntaxin18 complex (STX18, BNIP1, USE1 and SEC22B)
(Gillingham et al., 2014, Li et al., 2019, Xu et al., 2018, Zhao & Imperiale, 2017).
Although SNARE complexes typically mediate membrane fusion, it has been
proposed that RAB18 interacts with these proteins to mediate the close apposition of
membranes to facilitate lipid transfer (Xu et al., 2018). It has also been suggested
that SEC22B is dispensable for this function (Xu et al., 2018).

Our screen for RAB3GAP-dependent RAB18-interactors identified all of the NRZ
complex components, as well as the SNARE proteins STX18 and BNIP1 (Table 1).
Interestingly, we did not identify SEC22B but did identify SEC22A among these
proteins. SEC22A is one of the two SEC22B homologues in humans that lack the
central coiled-coil SNARE domain through which SEC22B mediates membrane
fusion (Rossi et al., 2004). Since it had not been previously described as a RAB18-
interacting protein, we investigated this further.

In the absence of commercially available antibodies for SEC22A, we examined its
localization through expression of a mEmerald-SEC22A fusion protein (Figure 3A).
mEmerald-SEC22A produced a characteristic reticular staining pattern and
colocalized with an exogenous ER marker suggesting that SEC22A localizes to the
ER. We next sought to compare the localization of SEC22A and RAB18 and to
determine whether they interact. However, coexpression of mEmerald-SEC22A and
mCherry-RAB18 disrupted normal ER morphology produced vesicular structures and
inclusions positive for both proteins (Figure S5). Although this was not inconsistent
with a protein-protein interaction, it precluded the use of coexpressed exogenous
proteins in continued testing.

As an alternative means of assessing SEC22A-interactions, we used proximity
biotinylation with a BirA*-SEC22A fusion protein in the HeLa cell panel. To minimize
potential toxicity while increasing biotin-ligase activity, we used BioID2 (Kim et al.,

2016) with a p.Gly40Ser active site modification (Branon et al., 2018) and reduced

biotin incubation time. Despite a low level of BioID2(Gly40S)-SEC22A expression,

the construct appeared to label RAB18 in a RAB3GAP-dependent manner (Figure

3B). 55 SEC22A-associated proteins were present in samples from wild-type cells in

>2 replicate experiments and represented by >3 spectral counts (Table S5). Further,

a subset of 9 SEC22A-associations were attenuated (intensity ratios <0.5) in

samples from both RAB18-null and RAB3GAP-null cells. These data are consistent

with a physiological SEC22A-RAB18 interaction.

A phenotype of altered LD morphology in lipid-loaded cells has been widely reported

in cells deficient in RAB18 (Bekbulat et al., 2019, Carpanini et al., 2014,

Gerondopoulos et al., 2014, Li et al., 2017, Liegel et al., 2013, Xu et al., 2018).

Similar observations have been made in cells deficient in some components of the

NRZ or Syntaxin18 complexes, but not in cells deficient in SEC22B (Xu et al., 2018).

To test whether SEC22A expression influences LD morphology, we examined the

effects of its silencing in oleic acid-loaded induced human hepatocyte (IHH) cells

(Figure 3C). ZW10 and NBAS silencing provided positive controls in our

experiments. ZW10 silencing led to a significant reduction in LD number (p<0.001)

compared to controls, whereas NBAS silencing led to both a significant reduction in

LD number and a significant increase in LD size (p<0.001 in each case). The effects

of SEC22A silencing mirrored those of NBAS silencing, producing a significant

reduction in LD number (p<0.001) and a significant increase in LD size (p<0.001).

Together, these data implicate SEC22A as involved in the same RAB18-mediated

process(es) as the NRZ and SNARE proteins.

<u>RAB18 recruits the orphan lipase TMCO4 to the ER membrane in a RAB3GAP-</u>

<u>dependent manner</u>

The most novel group of putative RAB18 effectors identified in our study were the

lipid modifying/mobilizing proteins, none of which had been reported to associate

with RAB18 previously. Among these, TMCO4 was identified in all three replicate

experiments and its association with RAB18 was highly RAB3GAP-dependent

(intensity ratio 0.06). Although annotated as containing transmembrane and coiled-

coil domains, it is orthologous to the Yeast protein Mil1/ Yfl034w, and likely to be a

partly soluble lipase (Whitfield et al., 2016). Consistently, TMCO4-EGFP expressed

in HeLa cells showed a diffuse localization. In contrast, EGFP-RAB18 partly localizes

to the ER, as shown by its colocalization with an ER marker (Figure 4A).

To assess the potential RAB18-TMCO4 interaction, we coexpressed mCherry-

RAB18 and TMCO4-EGFP (Figure 4B). As in our previous experiments, Celltrace

reagents were used to distinguish cells of wild-type and mutant genotypes. In wild-

type HeLa cells, coexpression of mCherry-RAB18 led to a dramatic redistribution of

TMCO4-EGFP to the ER membrane suggesting that RAB18 mediates recruitment of

TMCO4 to this compartment. Redistribution was completely absent in RAB3GAP1-

and RAB3GAP2-null cells but unaffected in TRAPPC9-null cells, consistent with the

BioID data.

As a means of verifying the RAB18-TMCO4 interaction, we carried out

immunoprecipitation experiments using exogenous HA-RAB18 and TMCO4-EGFP

(Figure 4C). As expected, TMCO4-EGFP copurified with HA-RAB18 when

expressed in wild-type or TRAPPC9-null cells, but not when expressed in

RAB3GAP1-null cells. These data indicate that RAB18 and TMCO4 interact directly

or indirectly as part of a protein complex in a RAB3GAP-dependent manner. Further,

both the microscopy and the immunoprecipitation data support the suggestion that

different GEFs can promote different RAB18-interactions.

<u>RAB18 is involved in cholesterol mobilization and biosynthesis</u>

Other putative RAB18 effectors with lipid-related functions included ORP2/OSBPL2,

INPP5B and EBP. ORP2 and INPP5B are robustly linked to a role in cholesterol

mobilization. ORP2 is thought to function as a lipid transfer protein that exchanges

cholesterol and PI(4,5)P_2 (Wang et al., 2019). INPP5B is implicated in the hydrolysis

of ORP2-bound PI(4,5)P_2, presumably driving the exchange process (Wang et al.,

2019). EBP is involved in *de novo* cholesterol biosynthesis (Silve et al., 1996). In the

Bloch pathway, it catalyses the conversion of 5α-cholesta-8, 24-dien-3β-ol

(zymosterol) to 5α-cholesta-7, 24-dien-3β-ol (24-dehydrolathosterol). In the

Kandutsch-Russel pathway, it catalyses the conversion of 5α-cholest-8(9)-en-3β-ol

to 5α-cholest-7-en-3β-ol (lathosterol)(Platt et al., 2014). On the basis of these

findings, we investigated the potential role of RAB18 in cholesterol uptake, efflux and

biosynthesis.

We first performed loading and efflux experiments to measure the flux of

cholesterol/cholesteryl ester (CE) while modifying the activity of RAB18. Chinese

hamster ovary (CHO) cells were generated to stably express RAB18(WT),

RAB18(Gln67Leu), or RAB18(Ser22Asn) (Figure S6A). In cells labelled with [^{14}C]-

oleate, but cholesterol depleted with lipoprotein-depleted serum (LPDS), levels of CE

were comparable in RAB18(Ser22Asn) and RAB18(WT) cells, whereas

RAB18(Gln67Leu) cells stored significantly more (Figure 5A, left panel). In cells

labelled with [^{14}C]-oleate and cholesterol-loaded FBS, levels of CE in

RAB18(Ser22Asn) remained unchanged, whereas its storage was elevated in

RAB18(WT) cells and RAB18(Gln67Leu) cells (Figure 5A, right panel). Interestingly,

in both [^{14}C]-oleate/LPDS and [^{14}C]-oleate/FBS cells, the addition of high density

lipoprotein (HDL, a vehicle mediating removal of cellular cholesterol) led to rapid

depletion of CE in RAB18(Gln67Leu) cells, but not in RAB18(Ser22Asn) or

RAB18(WT) cells (Figure 5A). Consistently, RAB18(Gln67Leu) cells also effluxed

significantly more [^{3}H]-cholesterol upon their incubation with apolipoprotein (apo) A-I

than the other cell types (Figure 5B). These observations were not explained by

altered expression of ABCA1, the transporter responsible for the rate-limiting step of

cholesterol efflux (Figure S6B). These data suggest that 'activated' GTP-bound

RAB18 strongly promotes the storage, turnover and mobilization of CE stored in

LDs. A plausible explanation for this is that active RAB18 promotes cholesterol

mobilization via ORP2 and INPP5B.

Given that ORP2 and INPP5B function in sterol mobilization whereas EBP functions

in sterol biosynthesis, we reasoned that RAB18 might coordinate their activities; that

ORP2 might act as an exchanger for the substrates or products of EBP-catalysis as

well as for cholesterol. In this case, defective mobilization of cholesterol precursors

might lead to their accumulation in RAB18-null cells. To test this hypothesis, we

incubated wild-type and RAB18-null cells for 48 hours in media supplemented with

LPDS, then subjected samples to analysis by GC-MS (Figure 5C-D). In RAB18-null

cells, we found that levels of the EBP-substrate cholest-8(9)-en-3β-ol were not

significantly different from those in wild-type cells. In contrast, levels of EBP-product

lathosterol were significantly increased (p<0.01). This suggests that RAB18

facilitates ORP2-mediated mobilization of lathosterol away from the site of its

generation by EBP. Consistently, levels of desmosterol - downstream of 24-

dehydrolathosterol in the Bloch pathway - were significantly reduced in RAB18-null

cells (p<0.01). These data suggest that delivery of substrates to post-EBP

biosynthetic enzymes is impaired.

To further test our hypothesis, we used molecular dynamics (MD) simulations to

model lathosterol binding to ORP2. We compared simulations of the ORP2 OSBP-

related domain (ORD) incorporating cholesterol and lathosterol over 1µs (Figure 5E-

G, Figure S7, Videos S1-2). Over this time, both simulations remained comparatively

stable in terms of positional root-mean-square deviation (RMSD) (Figure S7A), with

both termini highly dynamic. The pattern of positional root-mean-square fluctuations

(RMSFs) for atoms in each sterol showed higher conformational heterogeneity for

bound lathosterol than bound cholesterol, suggesting a less constrained interaction

(Figure S7B). RMSFs for the ORP2-ORD amino acid residues show similar patterns

for the two systems, although Tyr52 (Tyr110 in NP_653081), proximal to the sterol 3-

hydroxyl, fluctuated less in the presence of lathosterol (Figure 5E). This residue

appeared differently oriented with respect to the sterol in each simulation (Figure 5F,

Videos S1-2). Indeed, for a large proportion of the span of the simulation with

lathosterol, the Tyr52 hydroxyl and the sterol 3-hydroxyl were located close together

and this distance remained relatively stable (Figure 5G).

In crystal structures of sterols with the OSBP-related protein Osh4, water-mediated

interactions were prominent (Im et al., 2005). Therefore, we explored whether

positioning of water molecules might differ between the two simulations.

Interestingly, we found that water molecules were more frequently found in close

proximity to lathosterol than the cholesterol (Figure S7C). Together, the MD

simulations suggest that the ligand-binding tunnel of the ORP2-ORD adopts distinct

conformations when bound to lathosterol or cholesterol despite their minor structural

differences. The interactions appear relatively stable in each case, but speculatively,

the interaction with lathosterol may be more entropically favourable.

We next reasoned that impaired delivery of substrates to post-EBP enzymes might

reduce cholesterol biosynthesis in cells in which RAB18 is absent or dysregulated.

To explore this possibility, we cultured the panel of HeLa cell lines for 24 hours in

media supplemented with LPDS, treated them for 24 hours with [3H]-mevalonate or

[^{3}H]-acetate, then quantified labelled cholesterol (Figure 5H). Under both conditions,

labelling in two clonal wild-type controls was comparable, but was reduced in

RAB18-, RAB3GAP1-, RAB3GAP2-, TBC1D20- and TRAPPC9-null cells. Labelling

was lowest in the RAB18-null cells ($39.5\pm2.5\%$ and $6.8\pm0.5\%$ of controls for [^{3}H]-

mevalonate and [^{3}H]-acetate respectively). Levels in the cells of other genotypes

were between $46\pm2.5\%$-$73\pm5\%$ for [^{3}H]-mevalonate and $23\pm2\%$-$43\pm3\%\pm.0.5\%$ for

[^{3}H]-acetate. These data strongly suggest that RAB18 and its regulators are required

for normal cholesterol biosynthesis.

DISCUSSION

In this study, we have complemented previous work showing that proximity

biotinylation is a powerful means of identifying candidate Rab effectors (Gillingham

et al., 2019). Further – at least in the case of RAB18 - we have found that comparing

biotin-labelling produced by a BirA*-Rab in wild-type and GEF-deficient cells can be

particularly informative. We found that marked reductions in RAB18-association in

RAB3GAP-null cells were restricted to only 25 proteins and that these comprised

known and/or plausible effectors. We were able to exclude ~95% of RAB18-

associations from consideration as more likely to represent 'noise' from bystander

proteins.

Prior evidence supported 12 of the 25 interactions we detected. Independent

experiments with a mutant RAB18 fusion protein confirmed nucleotide-binding-

dependence of several interactors, and immunofuorescence confirmed compatible

localizations of several more. The known functions of the proteins were consistent

with previous work implicating RAB18 in coordination of lipid exchange between

apposed membranes (Xu et al., 2018). Further, gene-disease associations showed

substantial overlap with RAB18-deficiency/Warburg Micro syndrome. We have

presented additional validation for six interactions, five of which are novel.

Our protein-interaction data implicate RAB18 in regulation of a stepwise process in

which membrane/cytoskeletal remodelling precedes the engagement of tethering

proteins and then SNAREs to establish membrane contact sites. At these sites, it

then appears to couple the generation of lathosterol by EBP and its subsequent

mobilization and delivery to downstream biosynthetic enzymes. The effectors most

directly responsible for sterol mobilization are likely to be ORP2 and INPP5B. We

propose that RAB18, ORP2 and INPP5B function in an analogous manner to the

well studied ARF1-OSBP-SACM1L axis (Antonny et al., 2018). The GTPase

(RAB18/ARF1) facilitates ORP (ORP2/OSBP)-mediated exchange of sterol

(lathosterol/cholesterol) and phosphatidylinositde ($PI(4,5)P_2$/PI(4)P) between

apposed membranes. The phosphatase (INPP5B/SACM1L) acts to maintain the

necessary phosphoinositide concentration gradient (Figure 6).

The role of Rab proteins in mediating vesicular membrane trafficking is well known.

Emerging evidence also suggests that regulation of nonvesicular lipid exchange at

membrane contact sites may likewise be a broadly conserved function. Several Rab

isoforms are shown to interact with ORPs (Gillingham et al., 2019, Levin-Konigsberg

et al., 2019, Rocha et al., 2009, Sobajima et al., 2018). Many are also found to

interact with phospholipid phosphatases, including phosphatases from distinct

protein families (Fukuda et al., 2008, Gillingham et al., 2019, Williams et al., 2007).

Thus, the large number of possible GTPase-OSBP-phosphatase combinations

provides a potential means to orchestrate mobilization of a multitude of lipid species.

In this context, our finding that RAB18 also interacts with a biosynthetic enzyme and

mediates mobilization of a specific sterol intermediate is suggestive. The connection

between the enzyme and the mobilization machinery provides a mechanism

conferring increased selectivity for the lipid cargo (Nishimura & Stefan, 2020).

Resolving the functional and compositional diversity of membrane contact sites is a

challenge for future research. Similarly, the scope and precise mechanics of RAB18-

regulated lipid exchange are not yet understood. Although our experiments have

helped to validate our proximity-labelling approach, the roles of most of the RAB18-

effectors identified require additional analyses. Among the lipid-related proteins,

TMCO4 may potentially be directly or indirectly associated with sterol metabolism.

Although its substrate(s) are unknown, its expression is found to be upregulated in

hypercholesterolemia (Ong et al., 2013) and it is present on lipid rafts (Jin et al.,

2012). The remaining lipid-related proteins, C2CD2L/TMEM24 and C2CD2, might

potentially function in concert with ORP2 and/or INPP5B, since C2CD2L is found to

mediate PI transport and to facilitate generation of PI(4,5)P_2 (Lees et al., 2017).

The possible substitution of SEC22B for SEC22A in a RAB18-regulated Syntaxin18

SNARE complex (Figure S8A), and a possible role for this complex in promoting

membrane contacts rather than membrane fusion, is consistent with previous data

(Xu et al., 2018). Further, it would be in keeping with roles for the NRZ/Dsl1 complex

and SCFD2/Sly1 in dynamically orchestrating SNARE complex assembly (Ren et al.,

2009, Spang, 2012, Yamaguchi et al., 2002). More ambiguously, the RAB18-

interacting microtubule binding proteins have not previously been reported to work

together but do function in compatible locations (Figure S8B). SPG20 and

CAMSAP1 each associate with mitotic spindle poles, REEP4 participates in spindle-

dependent ER clearance from metaphase chromatin and BICD2 is a component of

the minus-end-directed dynein-dynactin motor complex (Hendershott & Vale, 2014,

Hueschen et al., 2017, Kumar et al., 2019, Lind et al., 2011, Nicholson et al., 2015,

Schlaitz et al., 2013, Urnavicius et al., 2015). Our TRAPPII-dependent RAB18

interaction data indicate that different GEF complexes affect largely distinct subsets

of interactions. However, more work will be required to determine whether these

regulators mediate independent or interdependent functions.

Our objective in studying RAB18 was to better understand the molecular pathology

of Warburg Micro syndrome. Though our protein-interaction data are relatively

preliminary, our functional findings represent good progress towards this goal. Our

key finding is that disrupted *de novo* cholesterol biosynthesis may contribute to

disease pathogenesis. Strongly supporting this suggestion, genes encoding multiple

cholesterol biosynthesis enzymes are linked to similar disorders (Platt et al., 2014).

For example, pathogenic variants in the lathosterol oxidase gene, *SC5D*, cause

lathosterolosis, which is associated with microcephaly, intellectual disability,

micrognathia, high arched palate and cataract (Anderson et al., 2019, Brunetti-Pierri

et al., 2002, Ho et al., 2014, Krakowiak et al., 2003, Rossi et al., 2007). Pathogenic

variants in the 7-dehydrocholesterol reductase gene, *DHCR7*, cause Smith-Lemli-

Opitz syndrome (SLOS), which has a similar spectrum of features and is among the

top differential diagnoses for Micro syndrome (Handley & Sheridan, 2018, Nowaczyk

& Wassif, 1998). Indeed, the similarities with SLOS were noted in the report first

identifying RAB18 as a disease-associated gene nearly a decade ago (Bem et al.,

2011).

Therapeutics have been trialled in lathosterolosis and SLOS. Cholesterol

supplementation has been widely used and more recently is being combined with

antioxidants with the aim of reducing toxicity from aberrant cholesterol metabolites

(Fliesler et al., 2018, Korade et al., 2014, Svoboda et al., 2012). Additionally,

paradoxical treatment with statins may increase the expression of sterol synthesizing

enzymes (Correa-Cerro et al., 2006, Ho et al., 2014, Jira et al., 1997, Wassif et al.,

2017). Careful clinical research will be required to determine the safety and efficacy

of such interventions in Warburg Micro syndrome.

MATERIALS AND METHODS

<u>Plasmids</u>

The EGFP-RAB18 construct has been described previously (Gerondopoulos et al.,

2014). The RAB18 sequence was excised from this construct using BamHI and

HindIII restriction enzymes (New England Biolabs, Hitchin, UK), and used to

generate constructs encoding mEmerald-RAB18 and mCherry-RAB18 by ligation

into mEmerald-C1 and mCherry-C1 vectors (Addgene, Watertown, MA) using HC T4

Ligase and rapid ligation buffer (Promega, Southampton, UK). Constructs encoding

BirA*-RAB18, BioID2(Gly40Ser)-SEC22A and mEmerald-SEC22A were generated

following PCR amplification from template and subcloning into an intermediate pCR-

Blunt II-TOPO vector using a Zero Blunt TOPO PCR Cloning Kit (ThermoFisher

Scientific, Waltham, MA) according to manufacturer's instructions. Fragments were

excised from intermediate vectors and then subcloned into target vectors using

restriction-ligation, as above. A construct encoding mCherry-ER was obtained from

Addgene, and a construct encoding TMCO4-EGFP was synthesised and cloned by

GeneWiz (Leipzig, Germany). Generation of recombinant pX461 and pX462

plasmids for CRISPR gene-editing, and recombinant pCMV vectors for preparation

of stable CHO cell lines are described below. Generation of recombinant pcDNA5

FRT/TO FLAG-BirA(Arg118Gly) vectors for preparation of stable T-Rex-293 cell lines

is described in supplementary methods. Details of PCR templates, primers and

target vectors are listed in Table S6.

<u>Antibodies and reagents</u>

A custom polyclonal antibody to RAB18 generated by Eurogentec (Southampton,

UK) has been described previously (Handley et al., 2015). An antibody to

RAB3GAP1 was obtained from Bethyl Labs (Montgomery, TX), an antibody to GFP

was obtained from Takara Bio (Saint-Germain-en-Laye, France), an antibody to β-

Tubulin was obtained from Abcam (Cambridge, UK) and an antibody to β-Actin was

obtained from ThermoFisher. Antibodies to hemagglutinin (HA), RAB3GAP2 and

TBC1D20 were obtained from Merck (Gillingham, UK). Antibodies to ZW10, STX18,

SPG20, RINT1, REEP4, BNIP1, C2CD2, TRIM13, WFS1, INPP5B, OSBPL2 and

NBAS were obtained from Proteintech (Manchester, UK). Antibody catalogue

numbers and the dilutions used in this study are listed in Table S6.

<u>Cell culture</u>

HeLa, T-REx-293 and IHH cells were maintained in DMEM media, RPE1 cells in

DMEM/F12 media and CHO cells in alpha-MEM media (ThermoFisher). In each

case, media was supplemented with 10% foetal calf serum (FCS) and 1% penicillin-

streptomycin (PS). Cells were maintained at 37°C and 5% CO_2.

<u>Generation of clonal 'knockout' HeLa and RPE1 cell lines</u>

CRISPR/Cas9 gene-editing was carried out essentially as described in Ran et al.,

2013 (Ran et al., 2013). Guide RNA (gRNA) sequences are shown in Table S6. A list

of the clonal cell lines generated for this study, together with the loss-of-function

variants they carry is shown in Figure S1A. Western blot validation is shown in

Figure S1B-E. Briefly, for each targeted exon, pairs of gRNA sequences were

selected using the online CRISPR design tool (http://crispr.mit.edu/). Oligonucleotide

pairs incorporating these sequences (Sigma) were annealed (at 50mM ea.) in 10mM

Tris pH8, 50mM NaCl and 1mM EDTA by incubation at 95ºC for 10 minutes followed

by cooling to room temperature. Annealed oligonucleotides were diluted and ligated

into BbsI-digested pX461 and pX462 plasmids (Addgene) using HC T4 Ligase and

rapid ligation buffer (Promega). Sequences of all recombinant plasmids were verified

by direct sequencing. Pairs of plasmids were contransfected into cells using

Lipofectamine 2000 reagent according to manufacturer's instructions. Cells were

selected for puromycin resistance (conferred by pX462) using 24 hours puromycin-

treatment. Following 12 hours recovery, they were selected for GFP fluorescence

(conferred by pX461) and cloned using FACSAria2 SORP, Influx or FACSMelody

instruments (BD, Wokingham, UK). After sufficient growth, clones were analysed by

PCR of the targeted exons (Primers are listed in Table S6). In order to sequence

individual gene-edited alleles, PCR products from each clone were first cloned into

ZeroBlunt TOPO vector (ThermoFisher) and then subjected to colony PCR. These

PCR products were then analysed by direct sequencing. Sequencing data was

assessed using BioEdit software (http://www.mbio.ncsu.edu/BioEdit/bioedit.html).

<u>BirA*/BioID proximity labelling (HeLa cells)</u>

Proximity-labelling in HeLa cells was carried out largely as described by Roux et al.

(Roux et al., 2012), but with minor modifications. HeLa cells were grown to 80%

confluence in T75 flasks and then each flask was transfected with 1-1.5µg of the

BirA*-RAB18 construct or 1µg of the BioID2(Gly40Ser)-SEC22A construct using

Lipofectamine 2000 reagent in Optimem serum-free medium (ThermoFisher) for 4

hours, according to manufacturer's instructions. 24 hours post-transfection, media

was replaced with fresh media containing 50µM Biotin (Merck) and the cells were

incubated for a further 24 or 6 hours (for BirA*-RAB18 and BioID2(Gly40Ser)-

SEC22A experiments respectively). Cells were then trypsinised and washed twice in

PBS before pellets were transferred to 2ml microcentrifuge tubes and snap-frozen.

For each pellet, lysis was carried out in 420µl of a buffer containing 0.2% SDS, 6%

Triton-X-100, 500mM NaCl, 1mM DTT, EDTA-free protease-inhibitor solution

(Expedeon, Cambridge, UK), 50mM Tris pH7.4. Lysates were sonicated for 10

minutes using a Bioruptor device together with protein extraction beads (Diagenode,

Denville, NJ). Each lysate was diluted with 1080µl 50mM Tris pH7.4, and they were

then clarified by centrifugation at 20 000xg for 30 minutes at 4ºC. Affinity purification

of biotinylated proteins was carried out by incubation of clarified lysates with

streptavidin-coated magnetic Dynabeads (ThermoFisher) for 24 hours at 4ºC. Note

that a mixture of Dynabeads - MyOne C1, MyOne T1, M270 and M280 – was used

to overcome a problem with bead-clumping observed when MyOne C1 beads were

used alone. Successive washes were carried out at room temperature with 2% SDS,

a buffer containing 1% Triton-X-100, 1mM EDTA, 500mM NaCl, 50mM HEPES

pH7.5, a buffer containing 0.5% NP40, 1mM EDTA, 250mM LiCl, 10mM Tris pH7.4,

50mM Tris pH7.4, and 50mM ammonium bicarbonate.

<u>Mass spectrometry</u>

Washed beads from BioID experiments with HeLa cells were subjected to limited

proteolysis by trypsin (0.3 µg) at 27°C for 6.5hours in 2mM urea, 1mM DTT, 75mM

Tris, pH=8.5, then supernatants were incubated overnight at 37°C. Samples were

alkylated with 50mM iodoacetamide (IAA) in the dark for 20minutes, then acidified by

addition of 8µl 10% trifluoroacetic acid (TFA).

Peptides were loaded on to activated (methanol), equilibrated (0.1% TFA) C18 stage

tips before being washed with 0.1% TFA and eluted with 0.1% TFA/80 acetonitrile.

The organic was dried off, 0.1% TFA added to 15 µl and 5 µl injected onto LC-MS.

Peptides were separated on an Ultimate nano HPLC instrument (ThermoFisher), and

analysed on either an Orbitrap Lumos or a Q Exactive Plus instrument

(ThermoFisher). After data-dependent acquisition of HCD fragmentation spectra,

data were analysed using MaxQuant and the uniprot human reference

proteome. Versions, releases, parameters and gradients used for separation are

provided in Table S6.

Cell labelling

In order to distinguish cells of different genotypes within the same well/on the same

coverslip, CellTrace Violet and CellTrace Far Red reagents (ThermoFisher) were

used to label cells before they were seeded. Cells of different genotypes were first

trypsinised and washed with PBS separately. They were then stained in suspension

by incubation with either 1µM CellTrace Violet or 200nM CellTrace Far Red for 20

minutes at 37°C. Remaining dye was removed by addition of a ten-fold excess of full

media, incubation for a further 5 minutes, and then by centrifugation and

resuspension of the resulting pellets in fresh media. Differently-labelled cells were

combined prior to seeding.

Immunofluorescence microscopy

Cells were seeded in 96-well glass-bottom plates (PerkinElmer, Waltham, MA)

coated with Matrigel (Corning, Amsterdam, Netherlands) according to manufacturer's

instructions, and allowed to adhere for 48 hours prior to fixation. In lipid-loading

experiments, cells were treated with 200µM oleic acid complexed to albumin (Merck)

and 1µg/ml BODIPY-558/568-C12 (ThermoFisher) for 15 hours prior to fixation. Cells

were fixed using a solution of 3% deionised Glyoxal, 20% EtOH, 0.75% acetic acid,

pH=5 (Richter et al., 2018), for 20 minutes at room temperature. They were then

washed with PBS containing 0.9mM $CaCl_2$ and 0.5mM $MgCl_2$ and blocked with a

sterile-filtered buffer containing 1% Milk, 2% donkey serum (Merck), 0.05% Triton-X-

100 (Merck), 0.9mM $CaCl_2$ and 0.5mM $MgCl_2$ in PBS pH=7.4 for at least 1 hour prior

to incubation with primary antibody. Primary antibodies were added in blocking buffer

without Triton-X-100, and plates were incubated overnight at 4ºC. Antibody dilutions

are listed in Table S6. Following washing in PBS, cells were incubated with 1:2000

Alexa 488-conjugated secondary antibody (ThermoFisher) in blocking buffer at room

temperature for 1-2 hours. Following further washing in PBS, cells were imaged

using an Operetta High Content Imaging System (PerkinElmer) equipped with

Harmony software. In comparative fluorescence quantitation experiments, at least 18

frames – each containing >5 wild-type and >5 mutant cells – were analysed per

genotype. ImageJ software was used to produce regions of interest (ROIs)

corresponding to each cell using thresholding tools and images from the 405nm and

645nm channels. Median 490nm fluorescence intensity was measured for each cell

and mutant fluorescence intensity (as %wild-type) was calculated for each frame and

combined for each genotype.

<u>Confocal microscopy – Live cell imaging</u>

HeLa or RPE1 cells were seeded on glass-bottom dishes (World Precision

Instruments, Hitchin, UK) coated with Matrigel (Corning) and allowed to adhere for

24 hours prior to transfection. Transfections and cotransfections were carried out

with 0.5µg of each of the indicated constructs using Lipofectamine 2000 reagent in

Optimem serum-free medium for 4 hours, according to manufacturer's instructions.

Media were replaced and cells were allowed to recover for at least 18 hours prior to

imaging. Imaging was carried out on a Nikon A1R confocal microscope equipped

with the Nikon Perfect Focus System using a 60✕ oil immersion objective with a 1.4

numerical aperture. The pinhole was set to airy1. CellTrace Violet was excited using

a 403.5nm laser, and emitted light was collected at 425–475nm. EGFP and

mEmerald were excited using a 488 nm laser, and emitted light was collected at

500–550 nm. mCherry was excited using a 561.3 nm laser, and emitted light was

collected at 570–620 nm. CellTrace Far Red was excited using a 638nm laser, and

emitted light was collected at 663-738nm.

<u>Immunoprecipitation</u>

HeLa cells were seeded onto 10cm dishes and allowed to adhere for 24 hours prior

to transfection. Transfections and cotransfections were carried out with 0.5μg of

each of the indicated constructs using Lipofectamine 2000 reagent in Optimem

serum-free medium for 4 hours, according to manufacturer's instructions. 24 hours

post-transfection cells were trypsinised, washed with PBS, then lysed in a buffer

containing 150mM NaCl, 0.5% Triton-X-100 and EDTA-free protease-inhibitor

solution (Expedeon), 10mM Tris, pH=7.4. Lysates were clarified by centrifugation,

input samples taken, and the remaining supernatants then added to 4μg rabbit anti-

HA antibody (Merck). After 30 minutes incubation at 4ºC on a rotator, 100μl washed

protein G-coupled Dynabeads (ThermoFisher) were added and samples were

incubated for a further 1 hour. The Dynabeads were washed x3 with buffer

containing 150mM NaCl, 0.1% Triton-X-100, 10mM Tris, pH=7.4, then combined with

a reducing loading buffer and subjected to SDS–PAGE.

<u>Generation of stable CHO cell lines</u>

A PCR product encoding mouse RAB18 was subcloned into an intermediate TOPO

vector using a TOPO PCR Cloning Kit (ThermoFisher) according to manufacturer's

instructions. The RAB18 fragment was then excised and subcloned into the pCMV

vector. PCR-based site-directed mutagenesis using a GeneArt kit (ThermoFisher)

was then used to generate pCMV-RAB18(Gln67Leu) and pCMV-RAB18(Ser22Asn)

constructs. CHO cells were transfected using Lipofectamine 2000 reagent

(ThermoFisher) and cells stably-expressing each construct were selected-for with

blasticidin. Under continued selection, clonal cell-lines were grown from single cells

and then RAB18 protein expression was assessed. Cell lines comparably expressing

RAB18 constructs at levels 2.5-5x higher than those wild-type cells were used in

subsequent experiments.

<u>Lipid loading experiments</u>

For LD number and diameter measurements, IHH cells were seeded onto glass

coverslips. siRNA transfections were carried out using FuGene reagent (Promega)

according to manufacturer's instructions. siRNAs targeting ZW10 and NBAS were

obtained from IDT, Coralville, IA; siRNA targeting SEC22A was obtained from

Horizon Discovery, Cambridge, UK. 48 hours following transfection, cells were

treated with 200nM BSA conjugated oleate for 24 hours. Coverslips were washed,

fixed with 3% paraformaldehyde and stained with 1μg/mL BODIPY and 300nM DAPI.

Fluorescence images were captured on a Zeiss LSM 780 confocal microscope

equipped with a 100x objective. Images were analysed using ImageJ software. Data

are derived from measurements from >100 cells/condition and are representative of

three independent experiments.

For cholesterol storage and efflux experiments with [^{14}C]-oleate, CHO cell lines

(described above) were seeded onto 12-well plates and then grown to 60-75%

confluence in Alpha media supplemented with 10% LPDS. Cells were grown in the

presence of 10% LPDS for at least 24 hours prior to the addition of oleate. 1 μCi/ml

[^{14}C]-oleate (Perkin Elmer) was added in the presence of 10% LPDS or 10% FBS for

24 hours. Cells were then washed and incubated with 50μg/ml HDL for 0, 4 or 8

hours. Cellular lipids were extracted with hexane. Lipids were then dried-down and

separated by thin layer chromatography (TLC) in a hexane:diethyl ether:acetic acid

(80:20:2) solvent system. TLC plates were obtained from Analtech, Newark, NJ.

Bands corresponding to cholesteryl ester (CE) were scraped from the TLC plate, and

radioactivity was determined by scintillation counting in a Beckman Coulter LS6500

Scintillation Counter using BetaMax ES Liquid Scintillation Cocktail (ThermoFisher).

Three independent experiments were carried out, each with four replicates of each

condition. Data from a representative experiment are shown.

For cholesterol efflux experiments with [^{3}H]-cholesterol, CHO cells were seeded onto

12-well plates and then grown to 60% confluence in Alpha media supplemented with

10% FBS. 5 μCi/ml [^{3}H]-cholesterol (PerkinElmer) was added in the presence of 10%

FBS. After 3x PBS washes, cells were incubated with serum-free media containing

25μg/ml of human apolipoprotein A-I (ApoA-I) for 5 hours. ApoA-I was a kind gift of

Dr. Paul Weers (California State University, Long Beach). Radioactivity in aliquots of

media were determined by scintillation counting in a Beckman Coulter LS6500

Scintillation Counter using LSC Cocktail (PerkinElmer). Cell lysates were produced

by addition of 0.1N NaOH for 1 hour, and their radioactivity was determined as

above. Cholesterol efflux was calculated as an average (+/- SD) of the % cholesterol

efflux (as a ratio of the media cpm/(media + cellular cpm) x 100%).

<u>Sterol analysis</u>

HeLa cells were grown to 80% confluence in T75 flasks, washed twice in PBS and then grown for a further 48 hours in DMEM supplemented with 10% LPDS. They were then trypsinised and washed twice in PBS before pellets were transferred to microcentrifuge tubes and snap-frozen. Pellets were resuspended in 200µl deionised water, sonicated for 20 seconds using an ultrasonic processor (Sonics & Materials Inc., CT, USA), then placed on ice. 750 µl of isopropanol containing 4 µmol/L 5α-cholestane as an internal standard was added to each sample, and then each was sonicated for a further 10 seconds. Lysates were transferred to 7ml glass vials and mixed with 250 µl tetramethylammonium hydroxide for alkalyine saponification at 80°C for 15 min, then cooled down for 10 minutes at room temperature. Sterols were extracted by addition of 500 µl tetrachloroethylene/methyl butyrate (1:3) and 2 ml deionised water, then thorough mixing. Samples were centrifuged for 10 minutes at 3000 rpm, and the organic phase containing the sterols was transferred to 300 µl GC vials. Extracts were dried under a stream of nitrogen, then sterols were silylated with 50 µL Tri-Sil HTP (HDMS:TMCS:Pyridine) Reagent (ThermoFisher) at 60°C for 1 hour.

Chromatography separation was performed on an Agilent gas chromatography-mass spectrometry (GC-MS) system (6890A GC and 5973 MS) (Agilent Technologies, Inc., CA, USA) with an HP-1MS capillary column (30 m length. x 250 µm diameter x 0.25 µm film thickness). The GC temperature gradient was as follows: Initial temperature of 120°C increased to 200°C at a rate of 20°C/min, then increased to 300°C at a rate of 2°C/min with a 15 minute solvent delay. Injection was at 250°C in splitless mode with ultrapurified helium as the carrier gas and the transfer line was 280°C. The mass spectra were acquired by electron impact at 70 eV using selected ion monitoring as follows: Lathosterol-TMS, cholesterol-TMS, and cholest8(9)-enol-TMS: m/z 458; 5α-cholestane and desmosterol-TMS: m/z 372; Lanosterol-TMS: m/z 393; and 7-dehydrocholesterol-TMS: m/z 325. The data were analysed using MassHunter Workstation Quantitative Analysis Software (Agilent Technologies, Inc.) and OriginPro 2017 (OriginLab Corp, MA, USA).

Molecular dynamics

Simulations were constructed as previously described, starting from a modified

version of the OSBPL2 lipid-binding domain structure (PDB 5ZM8)(Wang et al.,

2019). Parameters for lathosterol were obtained by modifying those of cholesterol to

substituting the double bond between C5-C6 with one between C7-C8. Simulations

were performed with the CHARMM36 force field.

Western blotting

Cell lysates were made with a buffer containing 150mM NaCl, 0.5% Triton-X-100

and EDTA-free protease-inhibitor solution (Expedeon), 50mM Tris, pH=7.4. Cell

lysates and input samples from BioID and immunoprecipitation experiments were

combined 1:1 with a 2x reducing loading buffer; a reducing loading buffer containing

10mM EDTA was added directly to Dynabead samples. SDS–PAGE and Western

blotting were carried out according to standard methods.

ACKNOWLEDGEMENTS

We thank the Warburg Micro syndrome children and their families. We thank

Professor C. A. Johnson and Dr J. A. Poulter for a critical reading of the manuscript.

COMPETING INTERESTS

No competing interests declared.

FUNDING

MH is supported by the University of Leeds and by a Programme Grant from the

Newlife Foundation for Disabled Children (Grant Reference Number: 17-18/23).

DATA AVAILABILITY

The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2019) partner

repository with the dataset identifiers PXD016631, PXD016336, PXD016326,

PXD016233 and PXD016404.

REFERENCES

Aligianis IA, Johnson CA, Gissen P, Chen D, Hampshire D, Hoffmann K, Maina EN,
Morgan NV, Tee L, Morton J, Ainsworth JR, Horn D, Rosser E, Cole TR, Stolte-
Dijkstra I, Fieggen K, Clayton-Smith J, Megarbane A, Shield JP, Newbury-Ecob R et
al. (2005) Mutations of the catalytic subunit of RAB3GAP cause Warburg Micro
syndrome. *Nat Genet* 37: 221-3

Anderson R, Rust S, Ashworth J, Clayton-Smith J, Taylor RL, Clayton PT, Morris
AAM (2019) Lathosterolosis: A Relatively Mild Case with Cataracts and Learning
Difficulties. *JIMD Rep* 44: 79-84

Antonny B, Bigay J, Mesmin B (2018) The Oxysterol-Binding Protein Cycle: Burning
Off PI(4)P to Transport Cholesterol. *Annu Rev Biochem* 87: 809-837

Barr F, Lambright DG (2010) Rab GEFs and GAPs. *Curr Opin Cell Biol* 22: 461-70

Bekbulat F, Schmitt D, Feldmann A, Huesmann H, Eimer S, Juretschke T, Beli P,
Behl C, Kern A (2019) RAB18 Loss Interferes With Lipid Droplet Catabolism and
Provokes Autophagy Network Adaptations. *J Mol Biol* 432: 1216-34

Bem D, Yoshimura S, Nunes-Bastos R, Bond FC, Kurian MA, Rahman F, Handley
MT, Hadzhiev Y, Masood I, Straatman-Iwanowska AA, Cullinane AR, McNeill A,
Pasha SS, Kirby GA, Foster K, Ahmed Z, Morton JE, Williams D, Graham JM,
Dobyns WB et al. (2011) Loss-of-function mutations in RAB18 cause Warburg micro
syndrome. *Am J Hum Genet* 88: 499-507

Borck G, Wunram H, Steiert A, Volk AE, Korber F, Roters S, Herkenrath P, Wollnik
B, Morris-Rosendahl DJ, Kubisch C (2011) A homozygous RAB3GAP2 mutation
causes Warburg Micro syndrome. *Human genetics* 129: 45-50

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL,
Perrimon N, Ting AY (2018) Efficient proximity labeling in living cells and organisms
with TurboID. *Nat Biotechnol* 36: 880-887

Brunet S, Saint-Dic D, Milev MP, Nilsson T, Sacher M (2016) The TRAPP Subunit
Trs130p Interacts with the GAP Gyp6p to Mediate Ypt6p Dynamics at the Late Golgi.
Front Cell Dev Biol 4: 48

Brunetti-Pierri N, Corso G, Rossi M, Ferrari P, Balli F, Rivasi F, Annunziata I,
Ballabio A, Russo AD, Andria G, Parenti G (2002) Lathosterolosis, a novel multiple-
malformation/mental retardation syndrome due to deficiency of 3beta-
hydroxysteroid-delta5-desaturase. *Am J Hum Genet* 71: 952-8

Carney DS, Davies BA, Horazdovsky BF (2006) Vps9 domain-containing proteins:
activators of Rab5 GTPases from yeast to neurons. *Trends Cell Biol* 16: 27-35

Carpanini SM, McKie L, Thomson D, Wright AK, Gordon SL, Roche SL, Handley MT,
Morrison H, Brownstein D, Wishart TM, Cousin MA, Gillingwater TH, Aligianis IA,
Jackson IJ (2014) A novel mouse model of Warburg Micro syndrome reveals roles
for RAB18 in eye development and organisation of the neuronal cytoskeleton.
Disease models & mechanisms 7: 711-22

Christoforidis S, McBride HM, Burgoyne RD, Zerial M (1999) The Rab5 effector
EEA1 is a core component of endosome docking. *Nature* 397: 621-5

Correa-Cerro LS, Wassif CA, Kratz L, Miller GF, Munasinghe JP, Grinberg A, Fliesler
SJ, Porter FD (2006) Development and characterization of a hypomorphic Smith-
Lemli-Opitz syndrome mouse model and efficacy of simvastatin therapy. *Hum Mol*
Genet 15: 839-51

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M (2014) Accurate proteome-
wide label-free quantification by delayed normalization and maximal peptide ratio
extraction, termed MaxLFQ. *Mol Cell Proteomics* 13: 2513-26

Fliesler SJ, Peachey NS, Herron J, Hines KM, Weinstock NI, Ramachandra Rao S,
Xu L (2018) Prevention of Retinal Degeneration in a Rat Model of Smith-Lemli-Opitz
Syndrome. *Sci Rep* 8: 1286

Fukuda M, Kanno E, Ishibashi K, Itoh T (2008) Large scale screening for novel rab
effectors reveals unexpected broad Rab binding specificity. *Mol Cell Proteomics* 7:
1031-42

Gerondopoulos A, Bastos RN, Yoshimura S, Anderson R, Carpanini S, Aligianis I,
Handley MT, Barr FA (2014) Rab18 and a Rab18 GEF complex are required for
normal ER structure. *The Journal of cell biology* 205: 707-20

Gillingham AK, Bertram J, Begum F, Munro S (2019) In vivo identification of GTPase
interactors by mitochondrial relocalization and proximity biotinylation. *Elife* 8: e45916

Gillingham AK, Sinka R, Torres IL, Lilley KS, Munro S (2014) Toward a
comprehensive map of the effectors of rab GTPases. *Developmental cell* 31: 358-73

Handley M, Sheridan E (2018) RAB18 Deficiency. In *GeneReviews((R))*, Adam MP,
Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A (eds)
Seattle (WA): NBK475670

Handley MT, Carpanini SM, Mali GR, Sidjanin DJ, Aligianis IA, Jackson IJ,
FitzPatrick DR (2015) Warburg Micro syndrome is caused by RAB18 deficiency or
dysregulation. *Open biology* 5: 150047

Hendershott MC, Vale RD (2014) Regulation of microtubule minus-end dynamics by
CAMSAPs and Patronin. *Proc Natl Acad Sci U S A* 111: 5860-5

Ho AC, Fung CW, Siu TS, Ma OC, Lam CW, Tam S, Wong VC (2014)
Lathosterolosis: a disorder of cholesterol biosynthesis resembling smith-lemli-opitz
syndrome. *JIMD Rep* 12: 129-34

Hueschen CL, Kenny SJ, Xu K, Dumont S (2017) NuMA recruits dynein activity to
microtubule minus-ends at mitosis. *Elife* 6: e29328

Im YJ, Raychaudhuri S, Prinz WA, Hurley JH (2005) Structural mechanism for sterol
sensing and transport by OSBP-related proteins. *Nature* 437: 154-8

Jin J, Sison K, Li C, Tian R, Wnuk M, Sung HK, Jeansson M, Zhang C, Tucholska M,
Jones N, Kerjaschki D, Shibuya M, Fantus IG, Nagy A, Gerber HP, Ferrara N,
Pawson T, Quaggin SE (2012) Soluble FLT1 binds lipid microdomains in podocytes
to control cell morphology and glomerular barrier function. *Cell* 151: 384-99

Jira P, Wevers R, de Jong J, Rubio-Gozalbo E, Smeitink J (1997) New treatment
strategy for Smith-Lemli-Opitz syndrome. *The Lancet* 349: 1222

Kim DI, Jensen SC, Noble KA, Kc B, Roux KH, Motamedchaboki K, Roux KJ (2016)
An improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell* 27: 1188-
96

Korade Z, Xu L, Harrison FE, Ahsen R, Hart SE, Folkes OM, Mirnics K, Porter NA
(2014) Antioxidant supplementation ameliorates molecular deficits in Smith-Lemli-
Opitz syndrome. *Biol Psychiatry* 75: 215-22

Krakowiak PA, Wassif CA, Kratz L, Cozma D, Kovarova M, Harris G, Grinberg A,
Yang Y, Hunter AG, Tsokos M, Kelley RI, Porter FD (2003) Lathosterolosis: an
inborn error of human and murine cholesterol synthesis due to lathosterol 5-
desaturase deficiency. *Hum Mol Genet* 12: 1631-41

Kumar D, Golchoubian B, Belevich I, Jokitalo E, Schlaitz AL (2019) REEP3 and
REEP4 determine the tubular morphology of the endoplasmic reticulum during
mitosis. *Mol Biol Cell* 30: 1377-1389

Lamber EP, Siedenburg AC, Barr FA (2019) Rab regulation by GEFs and GAPs
during membrane traffic. *Curr Opin Cell Biol* 59: 34-39

Lees JA, Messa M, Sun EW, Wheeler H, Torta F, Wenk MR, De Camilli P, Reinisch
KM (2017) Lipid transport by TMEM24 at ER-plasma membrane contacts regulates
pulsatile insulin secretion. *Science* 355: eaah6171

Levin-Konigsberg R, Montaño-Rendón F, Keren-Kaplan T, Li R, Ego B, Mylvaganam
S, DiCiccio JE, Trimble WS, Bassik MC, Bonifacino JS, Fairn GD, Grinstein S (2019)
Phagolysosome resolution requires contacts with the endoplasmic reticulum and
phosphatidylinositol-4-phosphate signalling. *Nat Cell Biol* 20: 1234-47

Li C, Luo X, Zhao S, Siu GK, Liang Y, Chan HC, Satoh A, Yu SS (2017) COPI-
TRAPPII activates Rab18 and regulates its lipid droplet association. *EMBO J* 36:
441-457

Li D, Zhao YG, Li D, Zhao H, Huang J, Miao G, Feng D, Liu P, Li D, Zhang H (2019)
The ER-Localized Protein DFCP1 Modulates ER-Lipid Droplet Contact Formation.
Cell Rep 27: 343-358 e5

Liegel RP, Handley MT, Ronchetti A, Brown S, Langemeyer L, Linford A, Chang B,
Morris-Rosendahl DJ, Carpanini S, Posmyk R, Harthill V, Sheridan E, Abdel-Salam
GM, Terhal PA, Faravelli F, Accorsi P, Giordano L, Pinelli L, Hartmann B, Ebert AD
et al. (2013) Loss-of-function mutations in TBC1D20 cause cataracts and male
infertility in blind sterile mice and Warburg micro syndrome in humans. *Am J Hum
Genet* 93: 1001-14

Lind GE, Raiborg C, Danielsen SA, Rognum TO, Thiis-Evensen E, Hoff G,
Nesbakken A, Stenmark H, Lothe RA (2011) SPG20, a novel biomarker for early
detection of colorectal cancer, encodes a regulator of cytokinesis. *Oncogene* 30:
3967-78

Liu X, Salokas K, Tamene F, Jiu Y, Weldatsadik RG, Ohman T, Varjosalo M (2018)
An AP-MS- and BioID-compatible MAC-tag enables comprehensive mapping of
protein interactions and subcellular localizations. *Nat Commun* 9: 1188

Nakamura H, Yamashita N, Kimura A, Kimura Y, Hirano H, Makihara H, Kawamoto
Y, Jitsuki-Takahashi A, Yonezaki K, Takase K, Miyazaki T, Nakamura F, Tanaka F,
Goshima Y (2016) Comprehensive behavioral study and proteomic analyses of
CRMP2-deficient mice. *Genes Cells* 21: 1059-1079

Ng BG, Lourenço CM, Losfeld ME, Buckingham KJ, Kircher M, Nickerson DA,
Shendure J, Bamshad MJ; University of Washington Center for Mendelian
Genomics, Freeze HH (2019) Mutations in the translocon-associated protein

complex subunit SSR3 cause a novel congenital disorder of glycosylation. *J Inherit*
Metab Dis 42: 993-7

Nicholson JM, Macedo JC, Mattingly AJ, Wangsa D, Camps J, Lima V, Gomes AM,
Doria S, Ried T, Logarinho E, Cimini D (2015) Chromosome mis-segregation and
cytokinesis failure in trisomic human cells. *Elife* 4: e05068

Nowaczyk MJM, Wassif CA (1998) Smith-Lemli-Opitz Syndrome. In
GeneReviews((R)), Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH,
Stephens K, Amemiya A (eds) Seattle (WA): NBK1143

Ong WY, Ng MP, Loke SY, Jin S, Wu YJ, Tanaka K, Wong PT (2013)
Comprehensive gene expression profiling reveals synergistic functional networks in
cerebral vessels after hypertension or hypercholesterolemia. *PloS one* 8: e68335

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ,
Inuganti A, Griss J, Mayer G, Eisenacher M, Perez E, Uszkoreit J, Pfeuffer J,
Sachsenberg T, Yilmaz S, Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF et al.
(2019) The PRIDE database and related tools and resources in 2019: improving
support for quantification data. *Nucleic Acids Res* 47: D442-D450

Platt FM, Wassif C, Colaco A, Dardis A, Lloyd-Evans E, Bembi B, Porter FD (2014)
Disorders of cholesterol metabolism and their unanticipated convergent mechanisms
of disease. *Annu Rev Genomics Hum Genet* 15: 173-94

Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F (2013) Genome
engineering using the CRISPR-Cas9 system. *Nat Protoc* 8: 2281-2308

Ren Y, Yip CK, Tripathi A, Huie D, Jeffrey PD, Walz T, Hughson FM (2009) A
structure-based mechanism for vesicle capture by the multisubunit tethering complex
Dsl1. *Cell* 139: 1119-29

Richter KN, Revelo NH, Seitz KJ, Helm MS, Sarkar D, Saleeb RS, D'Este E, Eberle
J, Wagner E, Vogl C, Lazaro DF, Richter F, Coy-Vergara J, Coceano G, Boyden ES,
Duncan RR, Hell SW, Lauterbach MA, Lehnart SE, Moser T et al. (2018) Glyoxal as
an alternative fixative to formaldehyde in immunostaining and super-resolution
microscopy. *EMBO J* 37: 139-159

Rocha N, Kuijl C, van der Kant R, Janssen L, Houben D, Janssen H, Zwart W,
Neefjes J (2009) Cholesterol sensor ORP1L contacts the ER protein VAP to control
Rab7-RILP-p150 Glued and late endosome positioning. *The Journal of cell biology*
185: 1209-25

Rossi M, D'Armiento M, Parisi I, Ferrari P, Hall CM, Cervasio M, Rivasi F, Balli F,
Vecchione R, Corso G, Andria G, Parenti G (2007) Clinical phenotype of
lathosterolosis. *Am J Med Genet A* 143A: 2371-81

Rossi V, Banfield DK, Vacca M, Dietrich LE, Ungermann C, D'Esposito M, Galli T,
Filippini F (2004) Longins and their longin domains: regulated SNAREs and
multifunctional SNARE regulators. *Trends Biochem Sci* 29: 682-8

Roux KJ, Kim DI, Burke B, May DG (2018) BioID: A Screen for Protein-Protein
Interactions. *Curr Protoc Protein Sci* 91: 19.23.1-19.23.15

Roux KJ, Kim DI, Raida M, Burke B (2012) A promiscuous biotin ligase fusion
protein identifies proximal and interacting proteins in mammalian cells. *The Journal*
of cell biology 196: 801-10

Schlaitz AL, Thompson J, Wong CC, Yates JR, 3rd, Heald R (2013) REEP3/4
ensure endoplasmic reticulum clearance from metaphase chromatin and proper
nuclear envelope architecture. *Developmental cell* 26: 315-23

Silve S, Dupuy PH, Labit-Lebouteiller C, Kaghad M, Chalon P, Rahier A, Taton M,
Lupker J, Shire D, Loison G (1996) Emopamil-binding protein, a mammalian protein
that binds a series of structurally diverse neuroprotective agents, exhibits delta8-
delta7 sterol isomerase activity in yeast. *The Journal of biological chemistry* 271:
22434-40

Sobajima T, Yoshimura SI, Maeda T, Miyata H, Miyoshi E, Harada A (2018) The
Rab11-binding protein RELCH/KIAA1468 controls intracellular cholesterol
distribution. *The Journal of cell biology* 217: 1777-1796

Spang A (2012) The DSL1 complex: the smallest but not the least CATCHR. *Traffic*
13: 908-13

Svoboda MD, Christie JM, Eroglu Y, Freeman KA, Steiner RD (2012) Treatment of
Smith-Lemli-Opitz syndrome and other sterol disorders. *Am J Med Genet C Semin*
Med Genet 160C: 285-94

Tinti M, Johnson C, Toth R, Ferrier DE, Mackintosh C (2012) Evolution of signal
multiplexing by 14-3-3-binding 2R-ohnologue protein families in the vertebrates.
Open biology 2: 120103

Urnavicius L, Zhang K, Diamant AG, Motz C, Schlager MA, Yu M, Patel NA,
Robinson CV, Carter AP (2015) The structure of the dynactin complex and its
interaction with dynein. *Science* 347: 1441-1446

Wang H, Ma Q, Qi Y, Dong J, Du X, Rae J, Wang J, Wu WF, Brown AJ, Parton RG,
Wu JW, Yang H (2019) ORP2 Delivers Cholesterol to the Plasma Membrane in
Exchange for Phosphatidylinositol 4, 5-Bisphosphate (PI(4,5)P2). *Mol Cell* 73: 458-
473 e7

Wassif CA, Kratz L, Sparks SE, Wheeler C, Bianconi S, Gropman A, Calis KA,
Kelley RI, Tierney E, Porter FD (2017) A placebo-controlled trial of simvastatin
therapy in Smith-Lemli-Opitz syndrome. *Genet Med* 19: 297-305

Whitfield ST, Burston HE, Bean BD, Raghuram N, Maldonado-Baez L, Davey M,
Wendland B, Conibear E (2016) The alternate AP-1 adaptor subunit Apm2 interacts
with the Mil1 regulatory protein and confers differential cargo sorting. *Mol Biol Cell*
27: 588-98

Williams C, Choudhury R, McKenzie E, Lowe M (2007) Targeting of the type II
inositol polyphosphate 5-phosphatase INPP5B to the early secretory pathway. *J Cell*
Sci 120: 3941-51

Xu D, Li Y, Wu L, Li Y, Zhao D, Yu J, Huang T, Ferguson C, Parton RG, Yang H, Li
P (2018) Rab18 promotes lipid droplet (LD) growth by tethering the ER to LDs
through SNARE and NRZ interactions. *The Journal of cell biology* 217: 975-995

Yamaguchi T, Dulubova I, Min SW, Chen X, Rizo J, Südhof TC (2002) Sly1 binds to
Golgi and ER syntaxins via a conserved N-terminal peptide motif. *Dev Cell* 2: 295-
305

Zhao L, Imperiale MJ (2017) Identification of Rab18 as an Essential Host Factor for
BK Polyomavirus Infection Using a Whole-Genome RNA Interference Screen.
mSphere 2: e00291-17

Zhen Y, Stenmark H (2015) Cellular functions of Rab GTPases at a glance. *J Cell*
Sci 128: 3171-6
1047

Figure 1. RAB3GAP-dependent RAB18-interactions in HeLa cells. (A) Schematic
to show experimental approach. Proximity biotinylation of guanine nucleotide
exchange factor (GEF)-dependent interactors by BirA*-RAB18 (ID-RAB18) is
disrupted in GEF-null cells. GEF-independent interactors are biotinylated in both
GEF-null and wild-type cells. Following affinity purification, GEF-dependent

interactions are determined by LFQ intensity ratios. (B) Plot to show correlation

between Log_2 LFQ intensities of individual proteins identified in samples purified

from RAB3GAP1- and RAB3GAP2-null cells. (C) Plot to show correlation between

Log_2 LFQ intensities of individual proteins identified in samples purified from wild-

type and RAB3GAP-null cells. Highlighted datapoints correspond to proteins later

found to have RAB3GAP-null:wild-type intensity ratios <0.5. (D) Venn diagram to

show overlap between all RAB18-associations, TRAPPII-dependent interactions

(TRAPPC9-null:wild-type intensity ratios <0.5) and RAB3GAP-dependent

associations (RAB3GAP-null:wild-type intensity ratios <0.5). (E) Western blotting of

samples purified from wild-type and RAB3GAP1-null cells in an independent BioID

experiment. Levels of selected proteins are consistent with RAB3GAP-null:wild-type

intensity ratios {braces}.

1067

1068

1069

Protein	n	Ratio	Orthologue PPI (Gillingham et al., 2014)	Additional evidence	
CAMSAP1	3	0.26			Microtubule/membrane remodelling
REEP4	3	0.35		Tinti et al., 2012*	
BICD2	2	0.25	BicD	Gillingham et al., 2019	
SPG20	**2**	**0.42**	**CG12001**	**This study (Figure 2)**	
ZW10	3	0.02	mit(1)15	Xu et al., 2018; Gillingham et al., 2019	Membrane tethering/docking
RINT1	3	0.16	CG8605	Xu et al., 2018	
NBAS	3	0.17	rod	Xu et al., 2018; Gillingham et al., 2019	
SCFD2	3	0.41	Slh	Gillingham et al., 2019	
SEC22A	**3**	**0.46**		**This study (Figure 3)**	
BNIP1	2	0.36		Xu et al., 2018	
STX18	2	0.40	Syx18	Xu et al., 2018	
TMCO4	**3**	**0.06**		**This study (Figure 4)**	Lipid modifying/ mobilising
OSBPL2	**3**	**0.35**		**This study (Figure 5)**	
EBP	**3**	**0.38**		**This study (Figure 5)**	
INPP5B	**2**	**0.00**		**This study (Figure 5)**	
C2CD2L	2	0.22			
C2CD2	2	0.36			
TRIM13	2	0.29			Autophagy receptors
FAM134B	2	0.46			
RAB3GAP2	3	0.00	rab3-GAP	Gerondopoulos et al., 2014	Other
RAB3GAP1	3	0.01	CG31935	Gerondopoulos et al., 2014	
SSR3	3	0.49			
MFHAS1	2	0.00	lrrk		
WFS1	2	0.00			
SCARA3	2	0.42			

1070
Table 1. RAB3GAP-dependent RAB18-interactions in HeLa cells. 25 proteins

with mean RAB3GAP-null:wild-type intensity ratios <0.5, identified in two or more

independent proximity biotinylation experiments. Orthologous proteins identified by

Gillingham et al., 2014, and other studies providing supporting evidence for

interactions are shown. Proteins are grouped according to their reported functions.

The full dataset is provided in Table S1.

Gene(s)	Homologue(s)	Syndrome(s)	Inheritance	OMIM	Overlapping features
RAB3GAP1, RAB3GAP2, RAB18, TBC1D20	-	Warburg Micro syndrome; Martsolf syndrome	AR	600118, 614222, 614225, 615663; 212720	intellectual disability (ID), microcephaly (M), ascending spastic paraplegia (ASP), cataract (C), microphthalmia (Mo), microcornea (Mc), optic atrophy (OA), seizures (S), corpus callosum hypogenesis (CCH), cerebellar vermis hypoplasia (CVH), genital abnormalities (GA), Neuropathy (N)
EBP	-	CDPX2; MEND syndrome	XLD; XLR	302960; 300960	ID, M, C, Mo, Mc, S, CCH, CVH, GA
INPP5B	*OCRL*; *INPP5K*	Lowe syndrome; MDCCAID	XLR; AR	309000; 607875	ID, M, C, S
SSR3	-	Congenital disorder of glycosylation	AR	Ng et al., 2019	ID, M, CCH, GA
WFS1	-	Wolfram syndrome; CTRCT41	AR; AD	222300; 116400	C, OA, GA
SPG20	-	Troyer syndrome (SPG20)	AR	275900	ID, M, ASP
BICD2	-	Spinal muscular atrophy	AD	615290; 615291	ASP
REEP4	*REEP1*; *REEP2*	SPG31; SPG72	AD	610250; 615625	ASP
NBAS	-	SOPH syndrome	AR	614800	OA
FAM134B	-	HSAN IIB	AR	613115	N

1077

1078

1079 **Table 2. Genes encoding putative RAB18 effectors or their homologues are**

1080 **associated with diseases that share overlapping features with Warburg Micro**

1081 **syndrome.** AR, autosomal recessive; XLD, X-linked dominant; XLR, X-linked

1082 recessive; AD, autosomal dominant.

1083

Figure 2. Initial screening of putative RAB18 effectors reveals that levels of SPG20 are significantly reduced in RAB18-null and TBC1D20-null cells. (A) Comparative fluorescence microscopy of selected RAB18-associated proteins in wild-type and RAB18-null HeLa cells. Cells of different genotypes were labelled with

CellTrace-Violet and CellTrace-Far Red reagents, corresponding to blue and

magenta channels respectively. Cells were stained with antibodies against indicated

proteins in green channel panels. (B) Comparative fluorescence microscopy of

selected RAB18-associated proteins in lipid-loaded wild-type and RAB18-null HeLa

cells. Cells were stained as above but were treated for 15 hours with 200µM oleic

acid, 1µg/ml BODIPY-558/568-C12 (Red channel) prior to fixation. (C) Schematic to

show method for quantification of protein levels by fluorescence intensity. In each

frame, cell areas for each genotype are generated by thresholding CellTrace

channels, intensity of antibody-staining is measured for each cell in multiple frames.

(D) Example frames showing wild-type and mutant cells of the indicated genotypes,

labelled with CellTrace-Far Red and CellTrace-Violet and reagents respectively, then

stained for SPG20. (E) Quantification of SPG20-specific fluorescence in wild-type

cells by direct comparison with SPG20-null cells. (F) Quantification of SPG20

fluorescence (%wt) in cells of different genotypes. Data were derived from analysis

of at least 18 frames – each containing >5 wild-type and >5 mutant cells – per

genotype. Error bars represent s.e.m. *$p<0.001$.

1106

1107

Figure 3. SEC22A associates with RAB18 and influences LD morphology. (A)
Confocal micrograph to show overlapping localization of exogenous mEmerald-
SEC22A (Green) and mCherry-ER (Red) in HeLa cells. (B) RAB18 LFQ intensities
from a reciprocal BioID experiment showing a reduced association between
BioID2(Gly40Ser)-SEC22A and endogenous RAB18 in RAB3GAP-null compared to
wild-type HeLa cells. Data were adjusted to account for non-specific binding of
RAB18 to beads and normalized by SEC22A LFQ intensities in each replicate
experiment. Error bars represent s.e.m. Data for other BioID2(Gly40Ser)-SEC22A-
associated proteins are provided in table S5. (C) Bar graphs to show effects of
ZW10, NBAS and SEC22A knockdowns on lipid droplet number and diameter.
siRNA-treated IHH cells were loaded with 200nM BSA-conjugated oleate, fixed and
stained with BODIPY and DAPI, and imaged. Images were analysed using ImageJ.

1120   Data are derived from measurements from >100 cells/condition and are

1121   representative of three independent experiments. Error bars represent SD. *p<0.001

1122

1123

1124

Figure 4. mCherry-RAB18 recruits TMCO4-EGFP to the ER membrane in a

RAB3GAP-dependent manner. (A) Confocal micrographs to show diffuse

localization of exogenous TMCO4-EGFP (Green) compared to mCherry-ER (Red)

and overlapping localization of exogenous EGFP-RAB18 (Green) and mCherry-ER

in HeLa cells. (B) Confocal micrographs to show localization of exogenous mCherry-

RAB18 and TMCO4-EGFP in wild-type cells and in mutant cells of different

genotypes. Wild-type and mutant cells of the indicated genotypes were labelled with

CellTrace-Violet and CellTrace-Far Red reagents respectively (magenta and blue

channels). (C) Immunoprecipitation of exogenous HA-RAB18 from HeLa cells of

different genotypes. Cells were transfected with the indicated constructs and lysed

24 hours post-transfection. Anti-HA immunoprecipitates and input samples were

subjected to SDS-PAGE and immunostaining for HA and GFP.

1137

1138

Figure 5. RAB18 is involved in the mobilization and biosynthesis of

cholesterol. (A) Plots to show cholesteryl ester (CE) loading and efflux. CHO cells,

stably expressing RAB18(WT), RAB18(Gln67Leu) and RAB18(Ser22Asn), were

incubated with $[^{14}C]$-oleate, for 24 hours, in the presence of lipoprotein depleted

serum (LPDS)(Left panel) or FBS (Right panel). Following lipid extraction, thin layer

chromatography (TLC) was used to separate CE, and radioactivity was measured by

scintillation counting. Measurements were made at t=0 and at 4 and 8 hours

following the addition of 50µg/ml high density lipoprotein (HDL) to the cells. (B) Bar

graph to show cholesterol efflux. CHO cells were incubated with [^{3}H]-cholesterol, for

24 hours, in the presence of FBS. After washing, they were incubated with 25µg/ml

apolipoprotein A-I for 5 hours. The quantity of [^{3}H]-cholesterol in the media is shown

as a percentage of the total cellular radioactivity (mean±SD). (C) Schematic of post-

squaline cholesterol biosynthesis pathway with the sterols quantified by GC-MS

named. Solid arrows indicate biosynthetic steps catalysed by EBP, SC5D and

DHCR7. (D) Bar graph of sterols profile in wild-type and RAB18-null HeLa cells.

Cells were grown in media supplemented with LPDS for 48 hours. Extracted sterols

were analysed by gas chromatography-mass spectrometry (GC-MS). % Sterol was

calculated as a proportion of total quantified sterols, excluding cholesterol, following

normalization to a 5α-cholestane internal standard. n=3; ±s.e.m. (E) Positional root

mean square fluctuations (RSMFs) for amino acid residues in simulations of ORP2-

ORD complexed with cholesterol and lathosterol. Box and arrow show position of

Tyr52 (Tyr110 in NP_653081). Simulations used a structure of the ORP2-ORD

based on PDB 5ZM8. Parameters for lathosterol substituted the cholesterol double

bond between C5-C6 with one between C7-C8. Simulations were performed with the

CHARMM36 force field. (F) Still images from molecular dynamics simulations of the

ORP2-ORD complexed with cholesterol (left) and lathosterol (right). Position of

ORP2 Tyr110 and sterol-proximal water molecules are shown. (G) Plot to show the

distance between Tyr52 and sterol 3' hydroxyl groups over time in each simulation.

(H) Bar graphs to show incorporation of [^{3}H]-mevalonate and [^{3}H]-acetate into

cholesterol in a panel of HeLa cell lines. Cells were grown in media supplemented

with LPDS for 24 hours, then incubated with 5µCi/well [^{3}H]-mevalonate or 10µCi/well

[^{3}H]-acetate for 24 hours. TLC was used to separate free cholesterol and

radioactivity was quantified by scintillation counting (n=4; mean±SD). *p<0.01,

**p<0.001, †p<0.0005.

1173

1174 **Figure 6. Model for lathosterol mobilization mediated by RAB18.** ORP2 binds

1175 PI(4,5)P2 on an apposed membrane. RAB18 interacts with ORP2 and INPP5B

1176 promoting the hydrolysis of PI(4,5)P2 to PI(4)P and maintaining a PI(4,5)P2

1177 concentration gradient. RAB18 then coordinates the biosynthesis of lathosterol by

1178 EBP and subsequent lathosterol mobilization by ORP2.