Bacterial Major Vault Protein homologs shed new light on origins of the enigmatic

organelle

Tymofii H. Sokolskyi[1]

[1] Trinity College of Arts and Sciences, Duke university, Durham, NC, USA

Abstract

Vaults are large cone-shaped and highly conservative ribonucleoprotein complexes present in the

cells of most major eukaryote clades. However, despite their wide distribution, their functions

and evolutionary dynamics still remain enigmatic. Several minor functions in modulating

signaling cascades and multidrug resistance phenotypes were previously discovered for

eukaryotic vaults, yet nothing is known about bacterial homologs of the major vault protein

(MVP), a protein that comprises the entirety of vault external surface. Using gene and protein

BLAST searches in NCBI and UniProt databases we identified a number of bacterial species in

prokaryotic orders Myxococcales, Cytophagales and Oscillatoriales with >50% identity to

eukaryotic MVP sequences. Interestingly, all of these species are characterized by one common

feature – gliding type of motility. Secondary structures of the identified proteins were predicted

using RAPTORX web service and aligned via jFATCAT-flexible algorithm in the RCSB PDB

Java Structure Alignment tool to elucidate structural identity. Coiled coil domain at the MVP C-

terminus of all studied bacterial species resembled TolA protein of *Escherichia coli* by both

structure and sequence. We also showed that MVP sequences from chemotrophic bacteria

Myxococcales and Cytophagales contain a domain homologous to eukaryotic band-7 domain,

unlike cyanobacterial and eukaryotic major vault proteins. As expected, maximum-likelihood

phylogenetic trees for MVP sequences separate studied taxa into two clades – first clade contains

Oscillatoriales (Cyanobacteria) and Eukaryotes and the second one contains chemotrophic

bacteria. In addition, binding prediction via RAPTORX showed great multiplicity GMP and

CMP nucleoside monophosphate binding pockets in Myxococcales and Cytophagales MVP,

unlike eukaryotic and cyanobacterial proteins which had much lower affinity to these substrates.

Due to high similarity of eukaryotic and cyanobacterial MVP sequences and a pattern of its

phylogenetic distribution, we can speculate that the most likely scenario for vault appearance in

eukaryotes is horizontal gene transfer from cyanobacteria. Presence of GMP and CMP binding

pockets in MVP could also point to a function in depleting cytosolic nucleotide concentration

which would be beneficial, for instance, during a viral infection. Further research is necessary to

uncover potential functions of this enigmatic protein in bacteria and to determine its evolutionary

patterns. In addition, a correlation between MVP presence and gliding motility in bacteria could also lead to elucidating selective pressures on the early evolution of this protein.

Introduction

Vaults are large 13 MDa ribonucleoprotein complexes present in cells of many Eukaryota species (Kedersha et al., 1991). They consist of 3 types of proteins (major vault protein (MVP), vault poly-ADP ribose polymerase (vPARP), telomerase-associated protein (TEP1)) and vault RNA. TEP1 is also shared with the telomerase complex. Vaults seem to be very conservative structures present among various Metazoa, Fungi and Protozoa taxa (Kong et al., 1999), meaning they could perform or could have performed a globally significant function. Multiple specific functions of this structure known in Metazoa are implicated in signaling pathway regulation, multidrug resistance, immunity, etc. (Berger et al., 2009). They include:

1. Nuclear import of phosphoinositide phosphatase PTEN.
2. Scaffold in EGFR/MAPK cascade.
3. Apoptotic suppression via interactions with COP1 protein.
4. Nuclear import of activated estrogen receptors.
5. Unidentified role in axonal transport, possibly RNA transport (Li et al., 1999).
6. Possible regulation of poly-ADP ribosylation to facilitate DNA reparation.
7. Unidentified role in sea urchin ontogenesis (Stewart et al., 2005).
8. Multidrug resistance in mammal cells (Izquierdo et al., 1996).
9. Possible role in Epstein-Barr virus immunity.

A hypothesis was proposed for vaults to be an efficient way of amino acid and nucleotide storage (Shaik, 2013). This could explain their weird phylogenetic distribution – mostly in taxa that lost essential amino acid biosynthesis pathways and their upregulation during pathogen invasion – they could act as nutrient sequesters from the cytoplasm (Shaik, 2013). However, if this explanation is viable, then many questions regarding vault composition arise – what is the purpose of TEP1 and vPARP presence in the complex, why vaults contain relatively few RNAs, why are they shaped like they are and specifically why are they hollow inside.

Vault's outer surface completely consists of major vault protein monomers. Recently its homologs were identified in several bacterial taxa, namely in some representatives of Cyanobacteria, Deltaproteobacteria and Bacteriodetes (Shaik, 2013). However, until now they have not been investigated neither *in silico* nor *in vivo*. Analysis of structure and phylogeny of bacterial MVP-like proteins may shed new light on vault function and evolution.

Materials and methods

BLAST searches for the Homo sapiens MVP in Uniprot KB Bacteria database uncovered various

proteins, several of which, representing different bacterial clades were selected for further

investigation: *Moorea producens* uncharacterized protein F4Y3B4_9CYAN, *Saprospira grandis*

uncharacterized protein H6L4P8_SAPGL, *Microscilla marina* uncharacterized protein

A1ZGE7_9BACT, and *Enhygromyxa salina* uncharacterized protein A0A0C2D5V5_9DELT.

Subsequent searches for these proteins found similarities with TolA proteins for various

Proteobacteria and Bacteroidetes, and band 7 (SPFH) domain-containing proteins.

Then chosen MVP-like sequences from the first BLAST iteration were aligned with various

TolA protein sequences found in the UniProt database for the respective taxa using local

pairwise alignment tool EMBOSS Water (default settings; Smith-Waterman algorithm). Most

common region of similarity was identified – sequence of approximately 200-250 amino acids.

Maximum-likelihood phylogenetic tree with 1000 bootstrap replications were constructed in

MEGA 7.0, including some sequences of eukaryotic MVP.

Secondary structure and binding prediction for isolated regions of some of the studied proteins

was performed with the help of RAPTORX web service (Källberg et al., 2012). Acquired

structure data in the PDB format was then subjected for pairwise structure alignment via

jFATCAT-flexible algorithm in the RCSB PDB Java Structure Alignment tool to confirm

structural identity (Prlić et al., 2010; Ye & Godzik, 2003).

Results

Comparison to TolA

BLAST searches for these sequences uncovered various TolA proteins; highest identity of 36%

had *Gilliamella* TolA A0A1B9K2Y3_9GAMM. Series of alignments between the chosen

sequences and TolA and TolA-like proteins of different bacterial species (*Enhygromyxa,*

Myxococcus, Sandaracinus, Escherichia, Haemophilius, Methylophaga, Mangrovibacter,

Gilliamella, Frischella, Galibacterium) showed that the highest similarity is located in the coiled

coil region of the selected proteins and domain II of TolA. Identity variation between 22% and

37% and alignment of TolA sequences not only with the coiled coil region, but also with

adjacent fragments of MVP shoulder and N-terminal domain suggests that it is most likely a

result of homologous relationship between TolA domain II and MVP.

Results of the MOTIF search service on Genome Net (Bioinformatics Center at Kyoto

University, http://www.genome.jp/tools/motif/), Pfam database (Finn et al., 2013) support this

hypothesis showing TolA domains with E-values up to 1e-07 when analyzing bacterial MVP-

like proteins. Schematic structure comparisons of *Saprospira* MVP-like protein and *Escherichia*

TolA are shown in **Fig.1, A**.

Fig.1 Structural features of MVP-like proteins. **A.** Schematic representation of *Saprospira*

grandis uncharacterized protein (upper block) and *Escherichia coli* TolA protein (lower block)

sequences. Aligned regions are marked with red line (portion of domain II from *E. coli* TolA and

coiled coil with a small portion of MVP shoulder from *S. grandis* protein). Identity 27,6%,

similarity 42,0%. **B.** Location of band 7 domain in *Saprospira* protein H6L4P8_SAPGL (result

of MOTIF search). **C.** Predicted GMP-binding site of *Saprospira* MVP-like protein.

Structure prediction

Some of the most similar sequence fragments, corresponding to coiled coils of MVP-like

proteins and domain II of TolA for some of the studied proteins were subjected to structure

prediction via RAPTORX web service. Pairwise structural alignment was then conducted to

support the idea of homology between major vault protein and TolA. Chosen sequences include:

Haemophilius TolA TOLA_HAEIN (70-275); *Homo* Major Vault Protein MVP_HUMAN (729-

805); *Moorea* MVP-like protein F4Y3B4_9CYAN (750-820); *Saprospira* uncharacterized

protein H6L4P8_SAPGL (701-780). Acquired results, summarized in **table 1**, confirm the

conclusion of evolutionary relationship between TolA and MVP-like proteins because of

relatively low P-values and RMSD.

	Moorea	*Homo*	*Haemophilius*	*Saprospira*
Moorea				
Homo	1,79e-0,4; 3,17			
Haemophilius	2,03e-05; 2,96	9,93e-04; 1,56		

Saprospira	1,94e-08; 3,05	2,19e-06; 1,17	5,34e-06; 1,76	

Table 1. Results of jFATCAT-flexible structure alignment of structures, explained in text**.** First number represents P-value, second – RMSD value.

Comparison with band 7 domain

Other than TolA, some BLAST searches for MVP shoulder-like domains in KB Bacteria database showed more than 30% identity with band 7 proteins (*Halothermothrix* protein B8D0R4_HALOH showed the highest result of 33%). In addition, results of the MOTIF search service also identified SPFH / Band 7 domain in 509-598 region of *Saprospira* sequence H6L4P8_SAPGL with E-value=0.00039 (**Fig. 1, B**). Interestingly, same results were observed with other studied bacterial proteins, except cyanobacterial MVP-like proteins and eukaryotic MVP.

Secondary structures for corresponding fragments of these two proteins along with *Pongo* MVP MVP_PONAB shoulder domain (519-647) and *Homo* stomatin STOM_HUMAN band 7 domain (56-227) were predicted and aligned using jFATCAT-flexible algorithm**.** Resulting RMSD and P-value are shown in **table 2**. P-values and RMSD are low enough to conclude likely homology. *Saprospira* MVP shoulder-like domain exhibits significantly higher structural similarity to band 7 domain of stomatin and *Halothermothrix* protein than to MVP shoulder of *Pongo* MVP.

	Halothermothrix	Homo	Pongo	Saprospira
Halothermothrix				
Homo	1,18e-12; 1,55			
Pongo	1,57e-01; 3,01	1,55e-01; 3,03		
Saprospira	1,18e-14; 1,36	6,38e-10; 1,82	2,68e-01; 3,26	

Table 2. Results of jFATCAT-flexible alignment of structures, explained in text. First number represents P-value, second – RMSD value.

Binding prediction

Binding prediction via RAPTOR-X server was conducted for *Saprospira*, *Enhygromyxa*, *Moorea* and *Homo* MVP-like proteins. *Saprospira* protein showed high-multiplicity binding with GMP and CMP. Multiplicity represents the frequency with which the selected pocket was found in a set of ligand-binding protein structures – generally, if the value is above 40 pocket is highly

likely true. Multiplicity for the first GMP-binding site is 102, for the second – 35 (shown on **Fig.**

1, C), for CMP-binding site – 68. For *Ehyngromyxa* it is 102 and 28 for GMP and 72 for CMP.

For *Moorea* protein it is only 46 and 12 for GMP and 28 for CMP; *Homo* MVP shows 37 and 14

for 2 possible GMP-binding sites, 38 for UMP and none for CMP. All of the possible nucleotide

binding sites are located in 500-700 sequence, that corresponds to the part that aligns with MVP

shoulder/band 7 domain. TolA proteins were also subjected for similar analysis. Neither

Haemophilius nor *Gilliamella* TolA show any nucleotide binding *in silico*.

Discussion

Origin of vaults. We found 2 likely homologies for bacterial MVP-like proteins: TolA and band

7 domain proteins. A maximum-likelihood phylogenetic tree was constructed for complete

protein sequences of various bacterial MVP-like proteins (**fig. 2**). Two macroclades can be

defined: first one contains Bacteroidetes and Deltaproteobacteria, the second one – eukaryotic

MVP and cyanobacterial MVP-like proteins. The main structural difference between them is

presence of band 7 domain in the sequences of all first clade taxa, and its absence in all second

clade taxa. *Hyalangium* protein A0A085WQF1_9DELT is the only found non-cyanobacterial

MVP-like protein lacking band 7 domain, no similar deltaproteobacterial sequences were found.

Band 7 domains are generally known for association with lipid membranes (Tavernarakis et al.,

1999).

Fig. 2 Maximum-likelihood phylogenetic tree for various bacterial MVP-like proteins. *Haemophilius* TolA sequence was used as an outgroup.

One of these bacteria, specifically *Saprospira grandis*, displays an unusual characteristic – rod-shaped protein structures in the cytoplasm, called rhaphidosomes (Saw et al., 2012). Their functions are currently unknown. One of the rhaphidosome proteins, SGRA_0791, was shown to contain Band 7-like domain (Saw et al., 2012). However, alignment of its 25-232 sequence that corresponds to Band 7 domain, according to Uniprot, with 500-600 sequence of *Saprospira* Band 7-like domain displayed only 20% identity, that is very low for such a short sequence, especially compared to >50% identities of MVP-like proteins between each other. At least for now there is no evidence of connection between rhaphidosomes and MVP-like proteins.

Major vault protein homologs were not found in archaea, that are closely related to eukaryotes. Therefore, MVP-like proteins were likely to appear in the bacterial lineage after Bacteria/Archaea divergence and then experience secondary loss in some lineages. Alternatively, they MVP-like proteins were present in the Last Universal Common Ancestor and then lost in Archaea and retained in Bacteria. However, if that is the case, then the loss should have occurred after the divergence of Eukarya and Archaea and then simultaneously happen in all known archaeal clades which seems improbable. Therefore, we suggest that the most likely scenario of MVP appearance in Eukarya is lateral transfer from a bacterial lineage, most likely Cyanobacteria (Oscillatoriales), because they possess MVP-like protein with more than 50% identity to MVP. This event probably has happened before the divergence of major protist groups, animals, plants and fungi (Daly et al., 2013).

Collected information suggests a probable evolutionary scenario for major vault protein, schematically depicted in **fig. 3**. Pre-MVP appeared in the common ancestor of most bacterial phyla and contained band 7-like (SPFH) and TolAII-like domains (possibility of independent appearance of both domain types in many distantly related phyla is very low). Consequently, a duplication and a series of deletions occurred resulting in TolA protein, that lost band 7 domain in favor of a short transmembrane helix and acquired a variable N-terminal TolAIII domain. Some lineages, including Bacteroidetes, Cyanobacteria (at least Oscillatoriales) and Deltaproteobacteria retained the original pre-MVP. Ability to bind NMP appeared prior to Cyanobacteria/Bacteroidetes/Proteobacteria divergence, because it can be found in both *Moorea* and *Saprospira* proteins.

Cyanobacterial pre-MVP evolution made it move away from cell membrane transforming its band 7 domain into MVP shoulder. Cyanobacterial proteins could represent the first evolutionary

steps towards the vault complex formation. Then, TolAII-like domain evolved to vault coiled

coils that facilitate MVP aggregation into vaults, as demonstrated in modern eukaryotes (van

Zon et al., 2002). Similar structure of *Hyalangium* protein could have been acquired

convergently or due to a separate lateral gene transfer event. Then sequence of the resulting

protein was laterally transmitted from cyanobacteria to early eukaryotes (most likely due to

phagotrophic lifestyle of the latter ones (Shaik, 2013)) or due to an endosymbiotic event and was

subsequently lost in various lineages possibly including plants and fungi (Daly et al., 2013).

Fig. 3 Most likely scenario of major vault protein evolution. Probably, NMP-binding appeared in

Cyanobacteria and Deltaproteobacteria/Bacteroidetes MVP common ancestor, and then started

waning in Cyanobacteria.

Future directions. All bacteria with MVP-like proteins identified here despite belonging to

highly different taxa share a common characteristic: gliding motility (Engene et al., 2012; Iizuka

et al., 2003 [1, 2]; Reichenbach, 2015). Specifically, these taxa include Deltaproteobacteria (only

order Myxococcales), Bacteroidetes (specifically Cytophagales) and Cyanobacteria (only

Oscillatoriales). This type of locomotion usually is prevalent among bacteria living in low-water

content environments, including biofilms, mats, soils or hypersaline habitats, that is also true for

the species studied here (Spormann, 1999). Interestingly, TolA protein that is partially

homologous to bacterial MVP-like proteins is involved in flagellar locomotion (Cascales et al.,

2001). Therefore MVP-like proteins could have played a role in the transition to gliding motility

in selected taxa, mechanism of which is poorly studied (McBride, 2001).

Observation of MVP-like proteins NMP binding also supports the idea of vaults functioning as sequesters and regulators, as was proposed by Shaik, 2013. Bacterial MVP-like proteins could perform the same function vaults do, but without necessary aggregation into vaults. One of the interesting possible early functions of MVP-like proteins can be pathogen defense. By sequestering nucleotides and amino acids from the cytoplasm they can at least decrease the rate of bacteriophage reproduction. Nucleotides are required for bacteriophage replication, while they are not immediately necessary for bacterium during the infection process because no DNA replication is occurring. Membrane-association via band 7 domains also could help them maintain this function. It is possible that MVP demonstrates a lot weaker nucleotide binding than MVP-like proteins because of eukaryotic compartmentalization – deoxynucleotide concentration is a lot smaller in eukaryotic cytosol than in bacterial, meaning there is less sense in sequestering it during bacterial/viral infection. However, eukaryotes face wider systematic range of pathogens, meaning that a new way of spatial nutrient isolation should have been developed – that is aggregation into large hollow structures (vaults). However, this hypothesis is purely speculative and requires experimental verification.

References

Asfa Alli Shaik. The unique phylogenetic distribution of vault particles reveals its functional roles. A thesis submitted for the degree of doctor of Philosophy, Department of biological sciences, National university of Singapore 2013

Berger, W., Steiner, E., Grusch, M., Elbling, L., & Micksche, M. (2009). Vaults and the major vault protein: novel roles in signal pathway regulation and immunity. *Cellular and molecular life sciences*, *66*(1), 43.

Cascales, E., Lloubès, R., & Sturgis, J. N. (2001). The TolQ–TolR proteins energize TolA and share homologies with the flagellar motor proteins MotA–MotB. *Molecular microbiology*, *42*(3), 795-807.

Corda, D., & Di Girolamo, M. (2003). Functional aspects of protein mono☐ADP☐ribosylation. *The EMBO journal*, *22*(9), 1953-1958.

Daly, T. K., Sutherland-Smith, A. J., & Penny, D. (2012). Beyond BLASTing: tertiary and quaternary structure analysis helps identify major vault proteins. *Genome biology and evolution*, *5*(1), 217-232.

246   Daly, T. K., Sutherland-Smith, A. J., & Penny, D. (2013). In silico resurrection of the Major
247   Vault Protein suggests it is ancestral in modern eukaryotes. *Genome biology and*
248   *evolution*, *5*(8), 1567-1583.

249   Engene, N., Rottacker, E. C., Kaštovský, J., Byrum, T., Choi, H., Ellisman, M. H., ... &
250   Gerwick, W. H. (2012). Moorea producens gen. nov., sp. nov. and Moorea bouillonii
251   comb. nov., tropical marine cyanobacteria rich in bioactive secondary
252   metabolites. *International journal of systematic and evolutionary microbiology*, *62*(Pt
253   5), 1171.

254   Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., ... &
255   Sonnhammer, E. L. (2013). Pfam: the protein families database. *Nucleic acids*
256   *research*, *42*(D1), D222-D230.

257   Iizuka, T., Jojima, Y., Fudou, R., Tokura, M., Hiraishi, A., & Yamanaka, S. (2003).
258   Enhygromyxa salina gen. nov., sp. nov., a slightly halophilic myxobacterium isolated
259   from the coastal areas of Japan. *Systematic and applied microbiology*, *26*(2), 189-196.

260   Iizuka, T., Jojima, Y., Fudou, R., Hiraishi, A., Ahn, J. W., & Yamanaka, S. (2003).
261   Plesiocystis pacifica gen. nov., sp. nov., a marine myxobacterium that contains
262   dihydrogenated menaquinone, isolated from the Pacific coasts of Japan. *International*
263   *journal of systematic and evolutionary microbiology*, *53*(1), 189-195.

264   Izquierdo, M. A., Scheffer, G. L., Flens, M. J., Giaccone, G., Broxterman, H. J., Meijer, C. J.,
265   ... & Scheper, R. J. (1996). Broad distribution of the multidrug resistance-related vault
266   lung resistance protein in normal human tissues and tumors. *The American journal of*
267   *pathology*, *148*(3), 877.

268   Källberg, M., Wang, H., Wang, S., Peng, J., Wang, Z., Lu, H., & Xu, J. (2012). Template-
269   based protein structure modeling using the RaptorX web server. *Nature*
270   *protocols*, *7*(8), 1511.

271   Kedersha, N. L., Heuser, J. E., Chugani, D. C., & Rome, L. H. (1991). Vaults. III. Vault
272   ribonucleoprotein particles open into flower-like structures with octagonal
273   symmetry. *The Journal of cell biology*, *112*(2), 225-235.

274   Kong, L. B., Siva, A. C., Rome, L. H., & Stewart, P. L. (1999). Structure of the vault, a
275   ubiquitous celular component. *Structure*, *7*(4), 371-379.

276   Lewin, R. A. (1997). Saprospira grandis: A Flexibacterium That Can Catch Bacterial Prey
277   by``Ixotrophy''. *Microbial ecology*, *34*(3), 232-236.

278   Li, J. Y., Volknandt, W., Dahlstrom, A., Herrmann, C., Blasi, J., Das, B., & Zimmermann, H.
279   (1999). Axonal transport of ribonucleoprotein particles (vaults). *Neuroscience*, *91*(3),
280   1055-1065.

281   McBride, M. J. (2001). Bacterial gliding motility: multiple mechanisms for cell movement
282   over surfaces. *Annual Reviews in Microbiology*, *55*(1), 49-75.

283   Prlić, A., Bliven, S., Rose, P. W., Bluhm, W. F., Bizon, C., Godzik, A., & Bourne, P. E.
284   (2010). Pre-calculated protein structure alignments at the RCSB PDB
285   website. *Bioinformatics*, *26*(23), 2983-2985.

286   Reichenbach, H. 2015. Hyalangium gen. nov. Bergey's Manual of Systematics of Archaea
287   and Bacteria. 1–4.

288   Saw, J. H., Yuryev, A., Kanbe, M., Hou, S., Young, A. G., Aizawa, S. I., & Alam, M.
289   (2012). Complete genome sequencing and analysis of Saprospira grandis str. Lewin, a
290   predatory marine bacterium. *Standards in Genomic Sciences*, *6*(1), 84.

291   Spormann, A. M. (1999). Gliding Motility in Bacteria: Insights from Studies ofMyxococcus
292   xanthus. *Microbiol. Mol. Biol. Rev.*, *63*(3), 621-641.

293   Stewart, P. L., Makabi, M., Lang, J., Dickey-Sims, C., Robertson, A. J., Coffman, J. A., &
294   Suprenant, K. A. (2005). Sea urchin vault structure, composition, and differential
295   localization during development. *BMC developmental biology*, *5*(1), 3.

296   Tavernarakis, N., Driscoll, M., & Kyrpides, N. C. (1999). The SPFH domain: implicated in
297   regulating targeted protein turnover in stomatins and other membrane-associated
298   proteins. *Trends in biochemical sciences*, *24*(11), 425-427.

299   van Zon, A., Mossink, M. H., Schoester, M., Scheffer, G. L., Scheper, R. J., Sonneveld, P., &
300   Wiemer, E. A. (2002). Structural domains of vault proteins: a role for the coiled coil
301   domain in vault assembly. *Biochemical and biophysical research
302   communications*, *291*(3), 535-541.

303   Ye, Y., & Godzik, A. (2003). Flexible structure alignment by chaining aligned fragment pairs
304   allowing twists. *Bioinformatics*, *19*(suppl_2), ii246-ii255.