Peri-Head Distance Coding in the Mouse Brainstem

Wenxi Xiao[1]*, Kyle S. Severson[1]*, Hao Zheng[1], Ke Chen[1], P. M. Thompson[1], Manuel S. Levy[1],

Seonmi Choi[1], Shengli Zhao[2], Jun Takatoh[1], Vincent Prevosto[1&], and Fan Wang[1&]

Affiliations

1. Department of Brain and Cognitive Sciences, McGovern Institute for Brain Research,

Massachusetts Institute of Technology, Cambridge, MA, USA.

2. Department of Neurobiology, Duke University, Durham, NC, USA.

* These two authors contributed equally

& Address correspondence to prevosto@mit.edu and fan_wang@mit.edu

SUMMARY

Perceiving object distance in peri-personal space is essential for guiding movement and avoiding

danger. During active sensation, distance information is often anchored to the body via touch; yet

how early somatosensory circuits extract distance information from tactile inputs remains unclear.

Here, we investigate how second-order neurons in the mouse whisker brainstem encode peri-

head distance. Using *in vivo* extracellular recordings in awake mice in a naturalistic wall-passing

paradigm, we find brainstem neurons employ two distance-coding schemes: a "proximity" code,

where firing increases monotonically as objects approach the face; and a "map" code, where

neurons exhibit peak tuning at specific distances to collectively tile peri-head space. The map

code outperforms proximity code in population decoding of distance. Perturbation experiments

reveal multi-whisker integration and internuclear inhibition contribute to the generation of map-

like tuning. These findings highlight a previously underappreciated computational role for

brainstem circuits, where inhibition acts as a neural comparator to transform proximity-based

sensory inputs into a map-like representation of peri-personal space.

MAIN TEXT

The ability to perceive and represent space is fundamental for animals to interact with the external world. Estimating the distance of a predator, planning reach-to-grasp movements for food, or navigating complex terrains with obstacles and gaps all require accurate spatial processing. Of particular ethological importance is the space immediately surrounding the body, peripersonal space (PPS), which serves as a critical interface for sensorimotor integration, enabling object interaction, social engagement, and defensive behaviors (Serino, 2019). Classic primate studies identified multisensory neural substrates for PPS, where bimodal neurons respond both to tactile stimulation on specific body parts and to visual or auditory stimuli presented *near* those same body surfaces (Graziano & Cooke, 2006).

PPS has been proposed as a contact-related action field (Bufacchi & Iannetti, 2018), in which an accurate representation of relative distance to nearby entities directly constrains motor planning. For example, reaching for a cup beside the hand versus one placed farther away requires distinct patterns of arm and trunk muscle activation; likewise, crossing a narrow versus wide gap imposes different mechanical and postural demands. Some PPS-encoding neurons increase their firing as visual stimuli move into near space and loom toward the body surface, consistent with a proximity-based rate code (Graziano & Cooke, 2006). However, neuropsychological evidence indicates that distance from the body—in effect, what lies within PPS—is a fundamental component of spatial representation (Cowey et al., 1994; Làdavas et al., 1998), suggesting that a simple monotonic proximity code may be insufficient as the sole format for encoding distances in PPS. Clarifying how distance is represented in PPS is therefore crucial for uncovering the sensory computations that support spatial awareness and action execution.

To estimate distances in the surrounding environment, humans often rely on physical references—most commonly, the body itself—for example, describing something as being "an arm's length away". This body-referenced approach to spatial perception dates back to ancient

civilizations, which formalized units of measurement based on body parts, including the "palm",

"foot", "step", and "fathom" (the span of outstretched arms). Compared to continuous proximity

cues, body-part references provide discrete spatial scales that can be more readily interpreted

and used for motor planning. This intuitive measurement strategy raises the question of whether

the brain implements an analogous "internal ruler" that systematically encodes distances relative

to specific body parts. If so, what is the representational format of this distance code? Addressing

these questions may provide important insights into how spatial representations within PPS are

constructed and used by the brain to guide behavior.

The rodent whisker sensory system provides a natural framework to study distance

encoding, with grid-like whiskers serving as a built-in coordinate system for peri-head space

(Brecht et al., 1997). Rodents are nocturnal animals that rely extensively on their whiskers to

explore their surroundings (Diamond et al., 2008). The whisker array is organized into arcs whose

lengths form a gradient along the anterior-posterior axis (Belli et al., 2017), mapping whisker

identity onto specific regions of peri-head space (Figure 1A). Each whisker thus serves as a

potential distance reference, allowing the brain to integrate inputs across the array into a head-

centric representation of peri-head space (Pluta et al., 2017). Supporting this idea, a subset of

neurons in the whisker barrel cortex (wS1) exhibit non-monotonic tuning to specific distances

between the mouse's head and surrounding walls during virtual corridor navigation (Sofroniew &

Vlasov et al., 2015). Such tuning can be interpreted as a "map code", where the peaks of the

tuning curves map the distance to the head like tick marks on a ruler. At the population level, these

preferred distances spanned the full range of whisker reach, effectively forming a head-centered

representation of peri-head distance.

However, the map code for peri-head distance has not been observed in the periphery.

Whiskers are innervated by mechanosensory neurons with soma residing in the trigeminal

ganglion (TG), each of which has a single-whisker receptive field (Petersen, 2007). These

neurons respond to mechanical stresses on the whisker follicle generated when the whisker

bends upon contact with an object (Mitchinson et al., 2004; Bush et al., 2016; Campagner et al.,

2016; Severson & Xu et al., 2017). Consequently, extracellular recordings from TG neurons, using

a vertical pole positioned at various locations along the length of a single whisker, revealed that

firing rates increase as the point of contact moves closer to the face (Szwed et al., 2006). Thus,

in the periphery, TG neurons encode radial distance via a proximity-based rate code, referenced

to the base of a single whisker (Ahissar & Knutsen, 2008). This implies that a sensory code

transformation occurs along the whisker sensory pathway to convert whisker-centered proximity

code in the periphery into the map-like peri-head distance code observed in wS1.

We hypothesize that a transformation from proximity to map-like representation of peri-

head distance occurs at the level of second-order neurons in the principal trigeminal nucleus (PrV)

of the brainstem. PrV contains neurons with either single- or multi-whisker receptive fields

(Minnery & Simons, 2003) and receives convergent inputs from multiple subtypes of TG

mechanosensory neurons, including both rapidly and slowly adapting afferents (Sakurai et al.,

2013; Jacquin et al., 2015). In addition, PrV circuits incorporate local glycinergic inhibitory

interneurons (Avendaño et al., 2005; Bellavance et al., 2010) and receive topographically

organized long-range GABAergic input from the spinal trigeminal nucleus interpolaris (SpVi;

Jacquin et al., 2015, Furuta et al., 2008). These circuit motifs, supporting both multi-whisker

integration (Minnery & Simons, 2003; Furuta et al., 2008) and feedforward inhibition, make PrV a

candidate site for implementing the transformation of whisker-centered proximity signals into a

head-centered map code. Inspired by Porter and Groh (2006), we propose that a simple circuit

(Figure 1B) could transform monotonic proximity-based inputs from TG neurons into a non-

monotonic map-like readout of peri-head distance. Importantly, this circuit requires inhibition:

when proximity-based inputs from two TG neurons with different mechanical sensitivity are

subtracted via inhibition, a peak response at a specific peri-head distance emerges. To test these

hypotheses, we performed *in vivo* electrophysiological recordings from PrV neurons in a wall-

passing paradigm.

Results

A wall-passing paradigm to study peri-head distance encoding in the whisker brainstem

To evoke tactile sensation similar to that experienced during naturalistic wall-following behavior,

we devised a wall-passing paradigm for awake, behaving mice head-fixed on a linear treadmill

(Severson & Xu et al., 2017; Figure 1C, left; STAR Methods). In this setup, each stimulus

presentation consists of a wall moving past the mouse's head at a fixed radial distance from

anterior to posterior, delivering controlled tactile stimulation to the whisker array. A linear actuator

moved the wall along the anteroposterior axis at a constant velocity of 10 mm/s, operating

independently of the animal's locomotion (open loop). Before each wall pass, the radial distance

between the wall and the mouse's face (wall distance) was randomly set. Each pass lasted

approximately 14 seconds, with wall movement onset at t = 0 and whisker-wall contacts typically

occurring between 3-11 seconds. In total, each distance was presented for 15-20 wall stimulus

presentations. The wall position was aligned to the midline of the head, with wall distances

measured from the wall to the C2 whisker follicle (Figure 1C, right).

After habituation to the wall-passing stimulus, mice exhibited predominantly passive

whisker-wall interactions. To assess the degree of active retraction, we compared wall-induced

whisker bending angles of a single row of whiskers in awake versus anesthetized conditions (see

Supplemental Figure S1). Under anesthesia, the moving wall pushed whiskers in the retraction

direction, producing larger absolute bending angles at closer wall distances (Supplemental Figure

S1D, right). Awake, behaving mice in the same wall-passing paradigm showed similar whisker

bending angles (mean difference: 2.28°), with no statistically significant difference from the

anesthetized condition across whiskers (Supplemental Figure S1F). This observation suggests

that mice engaged in little active whisker retraction to avoid contact, instead allowing the whiskers

to passively glide along the passing wall. Given this minimal wall-induced active retraction,

whisker bending mostly increased monotonically with wall proximity, supporting proximity-based

distance encoding in the periphery.

PrV contains both proximity and map codes for peri-head distance

To examine how PrV neurons respond to the passing wall stimulus at varying distances, we first

performed *in vivo* extracellular recordings in mice with only the C-row whiskers intact, using single

tungsten electrodes (see Supplemental Figure S2A-C; STAR Methods). Based on our single-unit

isolation metrics (see STAR Methods; Supplemental Figure S3J-M), we isolated 46 units from 14

mice (see Table S1 for mouse metadata), including 18 units with single-whisker and 28 units with

multi-whisker receptive fields. We analyzed single-unit responses using peri-event time

histograms (PETHs) by computing the averaged firing rates for wall contact (3-11 s) and baseline

periods (-4.5-0 s) across trials. To determine wall responsiveness, we applied one-tailed Wilcoxon

signed-rank tests (see STAR Methods). Of the 46 units, 41/46 (89.1%) were wall-activated and

3/46 (6.5%) were wall-suppressed; one unit showed both activation and suppression at different

distances, yielding a total of 43/46 (93.5%) wall-responsive (presumably touch-sensitive) units.

To assess distance tuning, we performed two-tailed Wilcoxon signed-rank tests across all

distance pairs ($p < 0.05$, corrected using the Benjamini-Hochberg false discovery rate). Units with

significant differences in mean firing rates for at least one distance pair were classified as

"distance-tuned" (84.8%, 39/46). For each distance-tuned unit, we computed a z-scored tuning

curve (see STAR Methods). The majority of units (52.2%, 24/46) showed a broadly monotonic,

inversely proportional relationship with wall distance (e.g., Supplemental Figure S2D), consistent

with PrV neurons inheriting proximity-based TG inputs. Interestingly, a distinct subset of units

(13.0%, 6/46) clearly preferred intermediate or far wall distances (e.g., Supplemental Figure S2E),

revealing a heterogeneous representation of peri-head distance in PrV (see Supplemental Figure

S2F).

Because high-impedance tungsten electrodes preferentially capture high-amplitude

spikes, unit isolation was biased toward larger neurons (Buzsáki et al., 2012), raising the

possibility that under-sampling biased our estimate of the proportion of cells tuned to intermediate

wall distances. Furthermore, the low-yield of single electrode recordings resulted in a small

sample size (n=46 units well-isolated units over 14 mice). To overcome these limitations, we

shifted to 32-channel single-shank silicon probes for acute recordings of PrV single-unit activity

during the wall-passing paradigm (Figure 1C-E; See STAR Methods). We isolated 1,141 single

units across 36 mice (Supplemental Figure S3A-E). Of the 1,141 units, 67.3% (768/1141) were

significantly modulated by the passing-wall stimulus ("wall-responsive"). Among these wall-

responsive units, 532/768 (69.3%) showed significant activation at one or more distances,

333/768 (43.4%) showed significant suppression, and 97 units met both criteria (activated at some

distances and suppressed at others). Given that our probe spanned the entire dorsoventral axis

of PrV (Figure 1D) and that whisker-responsive neurons are specifically localized to ventral PrV

(barrelettes, Bechara et al., 2015), the prominence of unresponsive units, likely corresponding to

unstimulated microvibrissa-, jaw-, or lip-related neurons in dorsal PrV, is not unexpected.

Representative spike rasters, PETHs, and z-scored wall-distance tuning curves from six

wall-responsive units are shown in Figure 2A-C. These examples illustrate the diversity of tuning

responses: Unit 1 showed monotonic increases in firing with decreasing wall distance ("proximity"

tuning); Units 2-4 exhibited peak firing at specific intermediate distances ("map" tuning); Unit 5

displayed a complex response profile with both suppression and activation depending on wall

distance; and Unit 6 was suppressed during wall contact. In addition to sustained responses, wall-

touch onset and offset responses were often observed in the same PrV neurons, suggesting

convergence of multiple afferent types (i.e., slowly and rapidly adapting) onto PrV neurons,

consistent with anatomical evidence that different types of TG afferent terminals are intermingled

within PrV barrelettes (Sakurai et al., 2013). However, our analyses focused on mean firing rates

across the entire wall-contact period (8 s) and did not further dissect these transient dynamics.

We identified 656/1,141 (57.5%) units that were tuned to wall distance. Among these

distance-tuned units, 472/656 (72.0%)) were significantly activated by at least one distance

relative to baseline, 213/656 (32.5%) were significantly suppressed by at least one distance

relative to baseline, and 29/656 (4.4%) met both criteria (activated at some distances and

suppressed at others). Interestingly, "wall-activated", "distance-tuned" units exhibited lower

baseline firing rates than untuned units (5.46 ± 0.670 vs. 6.90 ± 0.682 sp/s; two-tailed

Kolmogorov-Smirnov (K-S) test, p = 1.91 × 10^{-10}), while "wall-suppressed", "distance-tuned" units

exhibited baseline firing rates comparable to untuned units (7.31 ± 1.124 vs. 6.90 ± 0.682 sp/s;

two-tailed K-S test, p = 0.27; Supplemental Figure S4A). During wall contact, "wall-activated",

"distance-tuned" units showed significant increase in firing relative to baseline (16.44 ± 1.640 vs.

6.36 ± 0.704 sp/s; one-tailed K-S test, p = 9.26 × 10^{-28}; Supplemental Figure S4B, left), "wall-

suppressed", "distance-tuned" units showed significant decrease in firing relative to baseline

(4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s; one-tailed K-S test, p = 3.06 × 10^{-4}; Supplemental Figure

S4B, right), whereas untuned units showed no difference (6.65 ± 0.687 vs. 6.90 ± 0.682 sp/s; two-

tailed K-S test, p = 0.74; Supplemental Figure S4C), validating our classification criteria.

To visualize tuning at the population-level, we concatenated normalized tuning curves for

wall-activated, distance-tuned units (Figure 2D) and wall-suppressed, distance-tuned units

(Supplemental Figure S4D) into heatmaps, sorted by each unit's preferred wall distance. While

many units exhibited tuning to the closest wall position (7 mm), others showed tuning to

intermediate or far distances (Figure 2E, Supplemental Figure S4F). To further characterize wall-

activated distance-tuned units, we compared mean firing rates at each non-closest distance (≥ 9

mm) to the closest distance (7 mm; one-tailed Wilcoxon signed-rank tests, see STAR Methods).

Units with peak firing at the closest distance were classified as "proximity" units (Supplemental

Figure S4G, left), while units with significantly higher firing at any non-closest distance were

categorized as "map" units (Supplemental Figure S4G, right). This analysis identified 266

proximity units (23.3%) and 118 map units (10.3%) out of the 1,141 total recorded units (Figure

2F). An additional 88 units (7.7%) were tuned but did not meet criteria for either map or proximity

and were labeled "ambiguous" (Supplemental Figure S4G, middle) and excluded from further

analysis. Suppressed-only units comprised a small portion of the population (119/1141; 10.4%,

Figure 2F).

We note that classification as "proximity" versus "map" is based on relative peak firing

rates within the tested range (7-23 mm). It remains possible that some proximity units would

exhibit decreased firing at even shorter distances, thus revealing map-like tuning beyond the

tested range. Therefore, our categorization serves as an operational definition to capture distinct

response properties within the presented distance range, rather than strict functional classes.

To assess the reliability of proximity and map codes across repetitions, we performed 50%

cross-validation by sorting units using half the wall passes and applying the sorting order to the

remaining half. The resulting distance-tuning heatmaps showed little difference in preferred

distance between cross-validated partitions (Supplemental Figure S4E). Together, these results

reveal that PrV robustly encodes peri-head distance using both a proximity code—characterized

by increasing firing at close distances—and a distributed map code. Though the map coding is

more prominent in wS1 (Sofroniew & Vlasov et al., 2015), its presence in PrV points to an

underappreciated role for the whisker brainstem in active sensory processing, beyond simple

relay.

The map code in PrV allows more accurate readout of peri-head distance

We next asked which encoding scheme, proximity or map, allows more accurate readout of peri-

head distance from a given neural population. To address this, we trained maximum correlation

coefficient decoders (see STAR Methods) on either the proximity or map subpopulation activity,

with untuned and shuffled subpopulations used as controls.

As expected, decoding accuracy during the wall-contact period was higher in both map

and proximity populations compared to controls. Notably, the map population (n = 118 units)

outperformed the proximity population (subsampled n = 118 of 266 units) in overall distance

decoding accuracy over the entire wall passing period averaged across all wall distances (Figure

3A). Untuned units still performed slightly above chance due to residual distance tuning that fell

below our stringent statistical threshold for wall-tuning classification.

Confusion matrices revealed that map units were particularly more accurate at decoding

intermediate and far distances (Figure 3B, right), whereas proximity units showed stronger

decoding performance only at the closest distance (Figure 3B, left). To quantify this, we compared

classification accuracy across nine wall distances for each population (Figure 3C; Table S2). At 7

and 9 mm, proximity units were significantly more accurate than map units. In contrast, map units

were significantly more accurate at decoding more distal wall positions (11-23 mm). To assess

decoding precision in millimeters, we computed a distance-classification error (in mm) by

multiplying the confusion matrix with a distance error matrix (Figure 3D; Table S3). While

classification error increased with distances, both populations showed submillimeter error in

decoding the closest distances (7 and 9 mm). However, only the map population showed

submillimeter error in decoding intermediate distances (from 11 to 15 mm, Figure 3E). These

results are consistent with the performance of single-neuron decoders trained on map and

proximity units and averaged across those populations (see STAR Methods, Supplemental Figure

S5; Table S4; Table S5), although their overall decoding accuracy is lower than population

decoders.

Together, these results suggest that proximity and map codes provide complementary

encoding schemes to read out peri-head distance: the proximity code supports highly accurate

decoding at close distances, while the map code enables finer discrimination across intermediate

and far distances. Thus, optimal distance readout by downstream circuits would integrate

information across both proximity and map subpopulations.

Multi-whisker integration contributes to the generation of the PrV map code

In the wall-passing paradigm, closer wall distances induced greater whisker bending, driving

stronger activation of whisker TG afferents. TG therefore transmits to PrV a monotonic, proximity-

based input, with firing rates increasing as stimulus distance to the head decreases. The presence

of PrV neurons tuned to specific wall distances (i.e., map cells) suggests a transformation of this

peripheral proximity code into a representation of peri-head distance. Because the passing wall

stimulates multiple whiskers of different lengths, we hypothesized that PrV could generate the

map code by integrating multi-whisker inputs. To test this hypothesis, we recorded PrV single-unit

activity during the wall-passing paradigm under two conditions within the same recording session:

first, with the mouse's full whisker array intact, and second, after acutely cutting all but a single

whisker, thus eliminating multi-whisker integration (see STAR Methods; Figure 4A). Continuous

recording of neural activity allowed us to track each neuron's distance tuning profile before and

after the acute whisker cutting manipulation.

Across the recorded population (n = 195 units from 18 mice), acute whisker cutting did not

significantly alter baseline firing rates (measured before wall-movement onset for each wall pass;

Wilcoxon signed-rank test, full array: 7.09 ± 1.034 vs. single whisker: 7.10 ± 1.038 sp/s, $p = 0.70$).

In contrast, firing rates for wall-activated, distance-tuned units (n = 86) during wall-passing periods

were significantly reduced following cutting down to a single whisker (full array: 11.58 ± 1.338 vs.

single whisker: 8.18 ± 2.602 sp/s, $p = 4.43 \times 10^{-8}$, Figure 4B). Of the 86 wall-activated and

distance-tuned units, 66 showed significant changes in firing rates during wall-passing periods

between cutting conditions (two-sided Wilcoxon signed-rank test, $p < 0.05$). Among these, 52 units

showed a significant decrease in firing rates, consistent with loss of excitatory input from whiskers

within their receptive fields, while 14 units displayed increased firing rates, likely reflecting

disinhibition due to removal of surround-whisker inputs (Furuta et al., 2008). Note that since we

could not identify the receptive field of each recorded PrV unit on the high channel count probe,

and that distance tuning was computed after the recording session, in many cases the principal

whisker driving a given neuron of interest was inevitably cut, leading to a near-complete loss of

response.

Of 21 units classified as map cells before whisker cutting, 10 of these remained tuned to

wall distance after cutting down to a single whisker. After cutting, we observed shifts in preferred

wall distance: two map cells shifted their preferred distance from an intermediate to the closest

wall distance (Figure 4C,D; population summary in Figure 4K). The whisker cutting manipulation

effectively converted these map cells to proximity cells, indicating that in these cases multi-

whisker integration was essential for generating the non-monotonic map code for peri-head

distance. On the other hand, four map cells shifted their preferred distance closer but did not fully

convert to proximity cells (e.g., Figure 4E,F). Two map cells retained their preferred distance after

whisker cutting but showed changes in response amplitude (Figure 4G,H). These results showed

that map cells could be generated with inputs from just a single whisker although multi-whisker

inputs still modulate the gain or the preferred distance tuning in map cells. Together, these findings

demonstrated that multi-whisker integration contributes to, but is not required for, transforming

proximity-based inputs into a non-monotonic map code in PrV.

We also examined how proximity cells were affected by the loss of multi-whisker input. 45

units were classified as proximity cells before whisker cutting, and only 15 of these remained

tuned to wall distance after cutting down to a single whisker. In remaining proximity cells, reducing

input to a single whisker changed the gain of their responses: nine units showed decreased

activity (e.g., Figure 4I,J, Top), while four exhibited increased activity (e.g., Figure 4I,J, Bottom).

Inhibition shapes the PrV map code

In addition to integrating inputs from different whiskers, the sensory code transformation occurring in PrV may also involve the subtraction of TG inputs derived from the same whisker but with differing distance sensitivity profiles. Prior research indicates that TG touch neurons exhibit heterogeneity in mechanosensory thresholds as well as in their adaptation properties (Abraira & Ginty, 2013; Furuta et al., 2020). This suggests that TG touch neurons innervating the same whisker may exhibit different proximity-tuning profiles. Whether involving single- or multi-whisker integration, a common feature of the hypothesized transformation model (Figure 1B) is the engagement of inhibitory mechanisms. Next, we aimed to directly test this by combining *in vivo* electrophysiology with optogenetic or chemogenetic silencing approaches.

The intrinsic circuitry of PrV is shaped by two major sources of inhibition: local glycinergic and GABAergic interneurons within PrV itself, as well as profuse long-range GABAergic projections from the interpolar part of the spinal trigeminal nucleus (SpVi; Furuta et al., 2008; Bellavance et al., 2010). To assess the role of local inhibitory neurons in encoding of peri-head distance within PrV, we unilaterally injected AAV-DIO-GtACR (Govorunova et al., 2015; Takatoh & Prevosto et al., 2021) into PrV of vGat-Cre mice (N=2), enabling Cre-dependent expression of the inhibitory opsin GtACR in PrV inhibitory neurons (Supplemental Figure S6A). We then performed acute optogenetic silencing of these local inhibitory neurons while recording single-unit activity in PrV during the wall-passing paradigm (Supplemental Figure S6B,D). For each wall distance, the "Laser OFF" and "Laser ON" conditions were randomly interleaved. Each distance was repeated 15-20 times in each condition to allow the comparison of distance tuning profiles with and without local inhibition.

Among the 112 well-isolated units, 18 were putative inhibitory neurons expressing GtACR, as they were strongly inhibited during the "Laser ON" period (modulation index ≤ -0.5; see STAR Methods; Supplemental Figure S6C,E). Of distance-tuned units (71/112), 17 were opto-inhibited,

exhibiting a decrease in firing rates during laser stimulation (modulation index < 0; see STAR Methods), while preserving their distance tuning profile (i.e., proximity or map, data not shown). Sixteen units showed disinhibition under the "Laser ON" condition (modulation index > 0; see STAR Methods). Nonetheless, these opto-disinhibited neurons, regardless of whether they were proximity or map cells, retained their preferred distance (e.g., Supplemental Figure S6F,G). These observations suggest that PrV local inhibitory neurons primarily modulate the gain of individual PrV neurons' firing rates, rather than altering their distance tuning.

We next investigated the role of long-range internuclear inhibitory inputs from SpVi to PrV. Due to the extensive size of SpVi, optogenetic silencing of its inhibitory neurons using a single optical fiber is impractical. Instead, we employed an AAV strategy to achieve sustained silencing by expressing tetanus toxin light chain (TeLC; Takatoh & Prevosto et al., 2021) to permanently block synaptic transmission. AAV-DIO-TeLC-EYFP was unilaterally injected into SpVi of vGat-Cre mice (N=11) to target inhibitory neurons known to send long-range projections to PrV (Figure 5A; Furuta et al., 2008). Post-hoc histology confirmed TeLC-EYFP expression in SpVi (Supplemental Figure S6H), as well as labeled axon terminals and boutons in PrV originating from SpVi inhibitory neurons (Figure 5B). Note that the large size of SpVi limited viral access to all PrV-projecting neurons, thereby constraining the extent of silencing.

We recorded single-unit activity in PrV 10-15 days after TeLC injection during the wall-passing paradigm and isolated 587 units (Supplemental Figure S3F-I). To assess the impact of suppressing long-range inhibitory inputs from SpVi, we compared wall-distance tuning in SpVi-TeLC mice to that in wild-type mice. Side-by-side comparison of normalized tuning curves of all wall-activated, distance-tuned units revealed a clear reduction in the proportion of map units in SpVi-TeLC mice (Figure 5C, Left: data previously presented in Figure 2, Right: TeLC experiment), shifting the population to prefer closer wall distances (Figure 5D). Tuning profile classification confirmed a significant shift in the distribution of tuning types (Figure 5E). Most importantly, the proportion of map units was reduced by more than half in SpVi-TeLC mice compared to wild-type

mice (3.9% vs 10.34%; Z-statistic: 4.64, p = 1.8×10^{-6}), while the proportion of proximity cells

increased (32.09% vs 23.31%; Z-statistic: -3.93, p = 4.27×10^{-5}). Our silencing manipulation did

not fully abolish map cells, possibly due to incomplete silencing of all long-range inhibitory

intersubnuclear projections. The proportion of suppressed units decreased modestly (7.3% vs

10.43%; Z-statistic: 2.12, p = 0.02), whereas the proportions of untuned (49.9% vs 48.2%; Z-

statistic: -0.68, p = 0.50) and ambiguous units (6.79% vs 7.71%; Z-statistic: 0.69, p = 0.49)

remained unchanged.

During both baseline and wall-passing periods, untuned units in SpVi-TeLC mice exhibited

significantly higher firing rates than those in the wild-type population, consistent with disinhibition

following loss of inhibitory inputs from SpVi (baseline: 9.34 ± 1.330 vs 6.90 ± 0.682 sp/s, p = 7.63

× 10^{-5}; wall passes: 9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s, p =1.06 × 10^{-4}; one-tailed K-S tests;

Supplemental Figure S6I). In contrast, distance-tuned units in SpVi-TeLC mice showed

comparable baseline activity (in the absence of wall contact) to those in wild-type mice (5.60 ±

0.845 vs. 5.46 ± 0.670 sp/s; one-tailed K-S test, p = 0.14), but significantly elevated firing rates

during wall contact (20.25 ± 2.263 vs. 16.44 ± 1.640 sp/s; one-tailed K-S test, p = 1.64 × 10^{-3}

Supplemental Figure S6J). These findings suggest that TeLC-mediated silencing of SpVi inputs

altered the tuning preferences of PrV neurons (i.e., shifting them toward closer wall distances)

without affecting their spontaneous activity level. Thus, long-range inhibition from SpVi plays a

key role in transforming feedforward proximity-based inputs from TG into a map-like peri-head

distance representation in PrV. More broadly, our findings extend the role of SpVi-PrV inhibitory

projections beyond simple sensory gating (Furuta et al., 2008; Mohar et al., 2013), implicating

them in a sensory code transformation.

Finally, to demonstrate how this SpVi-PrV inhibitory circuit enables the proximity-to-map

transformation, we implemented a simple model constrained by known anatomical motifs (Figure

6A; see STAR Methods). In our model, TG afferents send proximity-based inputs to neurons in

PrV via two distinct pathways: a direct excitatory (lemniscal) pathway and an indirect inhibitory

pathway mediated by SpVi inhibitory neurons. TG proximity-based inputs were modeled as

sigmoidal functions with thresholds skewed toward proximal distances (Figure 6B), and PrV

neurons received sparse, random combinations of excitatory inputs from TG neurons and

inhibitory inputs from SpVi (see STAR Methods). The model transformed monotonic inputs into a

heterogeneous population of PrV tuning curves, including both proximity- and map-like responses

to distance (Figure 6C). Across the modeled population (n = 1,225), PrV neurons exhibited a wide

range of preferred distances, reproducing qualitative features of the empirical tuning curves

(Figure 6D-F, left). Reducing inhibitory input per PrV neuron by 50% led to noticeable loss of map

units and expansion of proximity units (Figure 6D-F, right). These simulations show that simple

feedforward motifs can transform monotonic sensory input into a non-monotonic map-like code,

enabling extraction of an abstract representation of peri-head distance, and that inhibition is

necessary for generating intermediate and far distance tuning in PrV. Together, these results

highlight complementary roles for local and long-range inhibition in the brainstem trigeminal

complex: local inhibitory circuits primarily regulate sensory gain, whereas long-range SpVi inputs

reshape tuning to generate the map code for peri-head distance.

DISCUSSION

Here, we revealed a previously unrecognized sensory code transformation in the whisker

brainstem: from proximity-based whisker sensory inputs to map-like peri-head distance

representation. Using *in vivo* electrophysiological recordings from awake, behaving mice in a

naturalistic wall-passing paradigm, we discovered two encoding schemes for distance in PrV: a

proximity code in which neuronal firing rates increase monotonically as the distance becomes

closer, and a map code in which neurons exhibit peak firing at specific distances. Both encoding

schemes are robust and stable, invariant to the multiplexed and highly variable whisker inputs

across trials. Mechanistically, we provide evidence that both multi-whisker integration and long-

range inhibitory inputs from SpVi play critical roles in generating the non-monotonic map code.

While most prior research on whisker-based object localization has examined neural activities of

TG afferents (Ahissar & Knutsen, 2008) and cortical areas (Barnéoud et al., 1991; O'Connor et

al., 2010; Sofroniew & Vlasov et al., 2015) in awake, behaving rodents, the brainstem trigeminal

complex has remained largely unexplored. To our knowledge, only two prior studies have reported

in vivo extracellular single-unit activity in PrV (Moore et al., 2015; Chakrabarti & Schwarz, 2018).

Our findings thus fill a critical gap in understanding early stages of tactile processing, revealing

an anatomically-defined brainstem circuit that integrates and transforms peripheral signals to

construct representations of peri-head space. More broadly, this work highlights the role of PrV

as a processing center that converts afferent inputs into meaningful representations for

downstream computation.

Functional roles of proximity and map codes in PrV

Our discovery of PrV map cells challenges a long-held assumption of PrV as a simple relay of

trigeminal input. The map code, in which neurons fire maximally at a preferred distance and

decrease their response as the stimulus deviates from that value, mirrors a common encoding

motif observed across sensory systems (Porter & Groh, 2006). Such "peaked" tuning could be

computationally advantageous, as it allows distance information to be extracted from a relatively

small subset of neurons (Pouget et al., 2000; Jazayeri & Movshon, 2006), resulting in a sparse,

metabolically efficient representation capable of attaining high precision (Laughlin, 2001;

Olshausen & Field, 2004). The presence of a map code in the brainstem allows feature extraction

from multiplexed tactile signals during active sensation (e.g., contact forces, stick-slips,

vibrations), reducing computational burden for downstream structures and enabling rapid

extraction of behaviorally relevant features.

Functionally, the map and proximity codes could be specialized for different operating

regimes of PPS. Our decoder analysis showed that map cells were significantly better at decoding

intermediate and far distances, suggesting that they are optimized for fine-grained spatial

localization over a broader distance range within PPS. This fine-grained distance representation

is likely critical for tasks in which rodents use their whiskers to judge whether gaps are crossable

(Azarfar et al., 2018), detect apertures just large enough to pass through (Harris et al., 1999;

Grant et al., 2018), follow walls at a consistent distance (Sofroniew et al., 2014; Sofroniew &

Vlasov et al., 2015), or avoid obstacles during locomotion (Sofroniew et al., 2014; Warren et al.,

2021). The presence of map cells in the brainstem may significantly enhance the speed and

precision of these sensorimotor computations. Consistent with this idea, PrV projects directly to

the superior colliculus (Bosman et al., 2011), providing a subcortical pathway for distance signals

from the brainstem to premotor circuits that bypasses the thalamus and cortex. This direct routing

could support particularly rapid sensorimotor coordination, enabling swift orienting, approach, or

avoidance responses based on peri-head distance signals. Additionally, map cells could be better

suited for 3D shape perception, allowing the brain to detect small variations in contact distance

across the whisker array to reconstruct an object's surface geometry. In contrast, proximity cells

excel at decoding the closest distance, making them well suited for threshold-like detection of

imminent contact in peri-head space and for triggering rapid, reflexive responses, while

simultaneously encoding distance-invariant features such as texture, softness, or compliance.

While proximity and map cells may lie on a continuum of distance tuning rather than

forming strictly discrete classes, the distinction provides a useful framework for comparing

representational strategies along the lemniscal pathway. In TG, radial distance encoding is

mechanically driven and predominantly proximity-based (Szwed et al., 2006; Ahissar & Knutsen,

2008; Pammer & O'Connor et al., 2013; Birdwell et al., 2007; Bush et al., 2016). In PrV, we

identified the presence of map cells, but proximity cells are overrepresented. In wS1, a more

balanced distribution of distance preferences was reported (Sofroniew & Vlasov et al., 2015). This

gradient suggests a progressive transformation from low-level mechanical inputs to higher-order,

feature-selective representations of peri-head space that are increasingly aligned with the

demands of sensorimotor behavior.

Feedforward inhibition transforms proximity inputs into a map code

The results from SpVi-TeLC experiments and computational modeling support a canonical E-I-E

motif—monosynaptic excitation from TG combined with long-range disynaptic inhibition from

SpVi—that converts monotonic, proximity-based TG input into non-monotonic, distance-selective

responses in PrV map cells. Specifically, inhibitory inputs from SpVi act as "neural comparators":

by subtracting proximity signals with different distance sensitivity profiles, they allow PrV map

cells to exhibit peak activity at specific distances. Our findings extend beyond prior work showing

that SpVi inhibition sharpens angular tuning in PrV (Bellavance et al., 2010)—a form of gain

modulation—by demonstrating that the same projection can fundamentally change how features

are encoded, revealing a broader function for disynaptic inhibition in sensory code transformation.

More broadly, these findings situate the SpVi-PrV pathway within a growing class of

microcircuits involving inhibitory neurons that implement core sensory computations. Across

modalities, feedforward inhibition does not simply limit excitation but actively extracts features by

implementing subtractive comparison operations to transform stimulus representations.

Examples include binaural circuits in the lateral superior olive of the auditory brainstem, where

excitatory and inhibitory inputs carrying sound level information from the two ears are compared

to encode interaural level differences (Benichoux & Tollin, 2018), inhibitory subnetworks that

sculpt orientation and higher-order tuning in visual cortex (Niell & Scanziani, 2021), and lateral

inhibition in olfactory circuits that decorrelates odor representations (Geramita et al., 2016).

Recent connectomic analyses have revealed that inhibitory neurons form stereotyped, feature-

targeted connectivity patterns (Schneider-Mizell et al., 2025), supporting the idea that inhibition is

a commonly repeated anatomical substrate poised to perform computations underlying neural

code transformations. By showing that this convergent E-I-E motif (Figure 1B; Figure 6A) in the

first central processing center of the whisker system can transform monotonic inputs from the

periphery into a discrete distance map, our work supports a unifying view of this motif as a

computational primitive for constructing higher-order sensory representations.

Map code generated with and without multi-whisker integration

We initially hypothesized that the PrV map code emerges through integration of inputs from

multiple whiskers. Our whisker cutting experiments showed that a subset of PrV map cells

retained map-like distance tuning even when only a single whisker remained. These observations

indicate that multi-whisker integration supports, but is not required for, the generation of map-like

tuning in a subset of PrV neurons. Together with the diversity of mechanosensory thresholds

among TG touch neurons (Hao & Delmas, 2010; Furuta et al., 2020), these observations refined

our circuit model whereby PrV constructs the map code by subtracting inputs from TG afferents

(from same or different whiskers) with different proximity-tuning profiles.

Our study does not rule out peripheral mechanisms that may contribute to the map code.

Mechanosensory receptors in the whisker pad skin (Rice et al., 1986; Severson et al., 2019),

including low-threshold mechanoreceptors (LTMRs) that encode skin stretch (Abraira & Ginty,

2013; Grigg, 1996) and C-LTMRs in the surrounding hairy skin (Li et al., 2011), could convey

additional distance information to PrV. These signals may encode whisker pad tension and its

dynamic changes during wall contact, thereby shaping distance representations in ways not

captured by our current understanding of TG input. Notably, conventional extracellular recordings

in TG predominantly sample large-diameter neurons, making it technically difficult to assess the

contribution of C-LTMRs and other small-diameter afferents. Future work will be necessary to

disentangle the relative contributions of central and peripheral mechanisms in constructing peri-

head distance representations.

AUTHOR CONTRIBUTIONS

F.W. conceived the study. F.W., K.S.S, V.P., and W.X. designed the experiments. W.X. and K.S.S. performed all experiments. P.M.T. and V.P. provided training and guidance to W.X. on in vivo extracellular recording in the brainstem. K.S.S. and V.P. developed analysis code with input from W.X. H.Z. implemented the brainstem circuitry simulation. K.C. and M.L. advised on data analysis and statistical methods. J.T advised on viral constructs and viral injection surgeries. K.S.S., W.X., and V.P. analyzed the data. S.Z. generated GtACR and TeLC viruses. S.C. managed the mouse colony, performed genotyping, and bred all animals. W.X., K.S.S, and F.W. wrote the manuscript with input from all authors.

ACKNOWLEDGMENTS

We thank Drs. Jennifer Groh, David Kleinfeld, Lindsey Glickfeld, and John Pearson, as well as members of the Wang Lab, for insightful discussions of this study. We are grateful to Julia Dziubek, Ina Bando, and He Jiang for technical assistance, and to Priyadarshini Dutta and Gabrielle Sevrain for assistance with mouse colony maintenance. This work was supported by grants NS077986, DP1 NS137188, and a subcontract of U19NS137920 (to F.W.).

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1. Abraira, V.E., and Ginty, D.D. (2013). The sensory neurons of touch. *Neuron* **79**, 618–639.
2. Ahissar, E., and Knutsen, P.M. (2008). Object localization with whiskers. *Biol. Cybern.* **98**, 449–458.
3. Avendaño, C., Machín, R., Bermejo, P.E., and Lagares, A. (2005). Neuron numbers in the sensory trigeminal nuclei of the rat: a GABA- and glycine-immunocytochemical and stereological analysis. *J. Comp. Neurol.* **493**, 538–553.
4. Azarfar, A., Zhang, Y., Alishbayli, A., Miceli, S., Kepser, L., van der Wielen, D., van de Moosdijk, M., Homberg, J., Schubert, D., et al. (2018). An open-source high-speed infrared videography database to study the principles of active sensing in freely navigating rodents. *Gigascience* **7**, giy134.
5. Bechara, A., Laumonnerie, C., Vilain, N., Kratochwil, C.F., Cankovic, V., Maiorano, N.A., Kirschmann, M.A., Ducret, S., and Rijli, F.M. (2015). Hoxa2 selects barrelette neuron identity and connectivity in the mouse somatosensory brainstem. *Cell Rep.* **13**, 783–797.
6. Barnéoud, P., Gyger, M., Andrés, F., and van der Loos, H. (1991). Vibrissa-related behavior in mice: transient effect of ablation of the barrel cortex. *Behav. Brain Res.* **44**, 87–99.
7. Bellavance, M.A., Demers, M., and Deschênes, M. (2010). Feedforward inhibition determines the angular tuning of vibrissal responses in the principal trigeminal nucleus. *J. Neurosci.* **30**, 1057–1063.

8. Belli, H.M., Yang, A.E., Bresee, C.S., and Hartmann, M.J. (2017). Variations in vibrissal geometry across the rat mystacial pad: base diameter, medulla, and taper. *J. Neurophysiol.* **117**, 1807–1820.

9. Benichoux, V., and Tollin, D.J. (2018). Sound localization: These are not the neurons you are looking for. *eLife* **7**, e39244.

10. Birdwell, J.A., Solomon, J.H., Thajchayapong, M., Taylor, M.A., Cheely, M., Towal, R.B., Conradt, J., and Hartmann, M.J. (2007). Biomechanical models for radial distance determination by the rat vibrissal system. *J. Neurophysiol.* **98**, 2439–2455.

11. Bosman, L.W., Houweling, A.R., Owens, C.B., Tanke, N., Shevchouk, O.T., Rahmati, N., Teunissen, W.H., Ju, C., Gong, W., Koekkoek, S.K**.,** et al. (2011). Anatomical pathways involved in generating and sensing rhythmic whisker movements. *Front. Integr. Neurosci.* **5**, 53.

12. Brecht, M., Preilowski, B., and Merzenich, M.M. (1997). Functional architecture of the mystacial vibrissae. *Behav. Brain Res.* **84**, 81–97.

13. Bufacchi, R.J., and Iannetti, G.D. (2018). An action field theory of peripersonal space. *Trends Cogn. Sci.* **22**, 1076–1090.

14. Bush, N.E., Schroeder, C.L., Hobbs, J.A., Yang, A.E., Huet, L.A., Solla, S.A., and Hartmann, M.J. (2016). Decoupling kinematics and mechanics reveals coding properties of trigeminal ganglion neurons in the rat vibrissal system. *eLife* **5**, e13969.

15. Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and currents—EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420.

16. Campagner, D., Evans, M.H., Bale, M.R., Erskine, A., and Petersen, R.S. (2016). Prediction of primary somatosensory neuron activity during active tactile exploration. *eLife* **5**, e10696.

17. Chakrabarti, S., and Schwarz, C. (2018). Cortical modulation of sensory flow during active touch in the rat whisker system. *Nat. Commun.* **9**, 3907.

18. Cowey, A., Small, M., and Ellis, S. (1994). Left visuo-spatial neglect can be worse in far than in near space. *Neuropsychologia* **32**, 1059–1066.

19. Diamond, M.E., von Heimendahl, M., Knutsen, P.M., Kleinfeld, D., and Ahissar, E. (2008). "Where" and "what" in the whisker sensorimotor system. *Nat. Rev. Neurosci.* **9**, 601–612.

20. Furuta, T., Bush, N.E., Yang, A.E., Ebara, S., Miyazaki, N., Murata, K., Hirai, D., Shibata, K.I., and Hartmann, M.J.Z. (2020). The cellular and mechanical basis for response characteristics of identified primary afferents in the rat vibrissal system. *Curr. Biol.* **30**, 815–826.e5.

21. Furuta, T., Timofeeva, E., Nakamura, K., Okamoto-Furuta, K., Togo, M., Kaneko, T., and Deschênes, M. (2008). Inhibitory gating of vibrissal inputs in the brainstem. *J. Neurosci.* **28**, 1789–1797.

22. Geramita, M.A., Burton, S.D., and Urban, N.N. (2016). Distinct lateral inhibitory circuits drive parallel processing of sensory information in the mammalian olfactory bulb. *eLife* **5**, e16039.

23. Govorunova, E.G., Sineshchekov, O.A., Janz, R., Liu, X., and Spudich, J.L. (2015). Natural light-gated anion channels: a family of microbial rhodopsins for advanced optogenetics. *Science* **349**, 647–650.

24. Grant, R.A., Breakell, V., and Prescott, T.J. (2018). Whisker touch sensing guides locomotion in small, quadrupedal mammals. *Proc. Biol. Sci.* **285**, 20180592.

25. Graziano, M.S., and Cooke, D.F. (2006). Parieto-frontal interactions, personal space, and defensive behavior. *Neuropsychologia* **44**, 845–859.

26. Grigg, P. (1996). Stretch sensitivity of mechanoreceptor neurons in rat hairy skin. *J. Neurophysiol.* **76**, 2886–2895.

27. Harris, J.A., Petersen, R.S., and Diamond, M.E. (1999). Distribution of tactile learning and its neural basis. *Proc. Natl. Acad. Sci. USA* **96**, 7587–7591.

28. Jacquin, M.F., Arends, J.J., Renehan, W.E., Waite, P.M., and Shortland, P.J. (2015). Whisker-related circuitry in the trigeminal nucleus principalis: topographic precision. *Somatosens. Mot. Res.* **32**, 8–20.

29. Jazayeri, M., and Movshon, J.A. (2006). Optimal representation of sensory information by neural populations. *Nat. Neurosci.* **9**, 690–696.

30. Làdavas, E., Zeloni, G., and Farnè, A. (1998). Visual peripersonal space centred on the face in humans. *Brain* **121**, 2317–2326.

31. Laughlin, S.B. (2001). Energy as a constraint on the coding and processing of sensory information. *Curr. Opin. Neurobiol.* **11**, 475–480.

32. Li, L., Rutlin, M., Abraira, V.E., Cassidy, C., Kus, L., Gong, S., Jankowski, M.P., Luo, W., Heintz, N**., et al.** (2011). The functional organization of cutaneous low-threshold mechanosensory neurons. *Cell* **147**, 1615–1627.

33. Minnery, B.S., and Simons, D.J. (2003). Response properties of whisker-associated trigeminothalamic neurons in rat nucleus principalis. *J. Neurophysiol.* **89**, 40–56.

34. Mitchinson, B., Gurney, K.N., Redgrave, P., Melhuish, C., Pipe, A.G., Pearson, M., Gilhespy, I., and Prescott, T.J. (2004). Empirically inspired simulated electro-mechanical model of the rat mystacial follicle-sinus complex. *Proc. Biol. Sci.* **271**, 2509–2516.

35. Mohar, B., Katz, Y., and Lampl, I. (2013). Opposite adaptive processing of stimulus intensity in two major nuclei of the somatosensory brainstem. *J. Neurosci.* **33**, 15394–15400.

36. Moore, J.D., Mercer Lindsay, N., Deschênes, M., and Kleinfeld, D. (2015). Vibrissa self-motion and touch are reliably encoded along the same somatosensory pathway from brainstem through thalamus. *PLoS Biol.* **13**, e1002253.

37. Niell, C.M., and Scanziani, M. (2021). How cortical circuits implement cortical computations: mouse visual cortex as a model. *Annu. Rev. Neurosci.* **44**, 517–546.

38. O'Connor, D.H., Peron, S.P., Huber, D., and Svoboda, K. (2010). Neural activity in barrel cortex underlying vibrissa-based object localization in mice. *Neuron* **67**, 1048–1061.

39. Olshausen, B.A., and Field, D.J. (2004). Sparse coding of sensory inputs. *Curr. Opin. Neurobiol.* **14**, 481–487.

40. Pammer, L., O'Connor, D.H., Hires, S.A., Clack, N.G., Huber, D., Myers, E.W., and Svoboda, K. (2013). The mechanical variables underlying object localization along the axis of the whisker. *J. Neurosci.* **33**, 6726–6741.

41. Petersen, C.C. (2007). The functional organization of the barrel cortex. *Neuron* **56**, 339–355.

42. Pluta, S.R., Lyall, E.H., Telian, G.I., Ryapolova-Webb, E., and Adesnik, H. (2017). Surround integration organizes a spatial map during active sensation. *Neuron* **94**, 1220–1233.e5.

43. Porter, K.K., and Groh, J.M. (2006). The "other" transformation required for visual–auditory integration: representational format. *Prog. Brain Res.* **155**, 313–323.

44. Pouget, A., Dayan, P., and Zemel, R. (2000). Information processing with population codes. *Nat. Rev. Neurosci.* **1**, 125–132.

45. Rice, F.L., Mance, A., and Munger, B.L. (1986). A comparative light microscopic analysis of the sensory innervation of the mystacial pad. I. Innervation of vibrissal follicle-sinus complexes. *J. Comp. Neurol.* **252**, 154–174.

46. Schneider-Mizell, C.M., Bodor, A.L., Brittain, D., Buchanan, J., Bumbarger, D.J., Elabbady, L., Gamlin, C., Kapner, D., Kinn, S., Mahalingam, G., et al. (2025). Inhibitory specificity from a connectomic census of mouse visual cortex. *Nature* **640**, 448–458.

47. Sakurai, K., Akiyama, M., Cai, B., Scott, A., Han, B.X., Takatoh, J., Sigrist, M., Arber, S., and Wang, F. (2013). The organization of submodality-specific touch afferent inputs in the vibrissa column. *Cell Rep.* **5**, 87–98.

48. Serino, A. (2019). Peripersonal space (PPS) as a multisensory interface between the individual and the environment, defining the space of the self. *Neurosci. Biobehav. Rev.* **99**, 138–159.

49. Severson, K.S., Xu, D., Van de Loo, M., Bai, L., Ginty, D.D., and O'Connor, D.H. (2017). Active touch and self-motion encoding by Merkel cell-associated afferents. *Neuron* **94**, 666–676.

50. Severson, K.S., Xu, D., Yang, H., and O'Connor, D.H. (2019). Coding of whisker motion across the mouse face. *eLife* **8**, e41535.

51. Sofroniew, N.J., Cohen, J.D., Lee, A.K., and Svoboda, K. (2014). Natural whisker-guided behavior by head-fixed mice in tactile virtual reality. *J. Neurosci.* **34**, 9537–9550.

52. Sofroniew, N.J., Vlasov, Y.A., Hires, S.A., Freeman, J., and Svoboda, K. (2015). Neural coding in barrel cortex during whisker-guided locomotion. *eLife* **4**, e12559.

53. Szwed, M., Bagdasarian, K., Blumenfeld, B., Barak, O., Derdikman, D., and Ahissar, E. (2006). Responses of trigeminal ganglion neurons to the radial distance of contact during active vibrissal touch. *J. Neurophysiol.* **95**, 791–802.

54. Takatoh, J., Prevosto, V., Thompson, P.M., Lu, J., Chung, L., Harrahill, A., Li, S., Zhao, S., He, Z., Golomb, D., et al. (2022). The whisking oscillator circuit. *Nature* **609**, 560–568.

55. Warren, R.A., Zhang, Q., Hoffman, J.R., Li, E.Y., Hong, Y.K., Bruno, R.M., and Sawtell, N.B. (2021). A rapid whisker-based decision underlying skilled locomotion in mice. *eLife* **10**, e63596.

Figure 1. Acute PrV recordings in a wall passing paradigm

(**A**) Conceptual illustration showing how whiskers map peri-head space in head-centered coordinates.

(**B**) Left, schematic of a hypothetical brainstem circuit to compute non-monotonic distance code. Two TG neurons (A and B, red) with different distance sensitivities project to the brainstem: neuron A provides excitatory input to a PrV output neuron (C, black), whereas neuron B synapses onto an inhibitory neuron (blue) in the trigeminal brainstem complex. Middle, tuning curves for neurons A and B; the shaded region indicates the difference between their distance-tuning profiles. Right, linear summation yields a non-monotonic tuning curve in neuron C, with a peak at an intermediate peri-head distance.

(**C**) Left, Schematic of the experimental setup. A head-fixed mouse ran on a treadmill while a wall moved from anterior to posterior past the mouse's face. Wall distance from the face was randomly varied between 7-23 mm in 2-mm increments on each pass. Each recording contains 15-20 repetitions per distance. Right, Example video frame showing multiple whiskers contacting the wall (moving left to right). Red arrow indicates measured distance between face and wall.

(**D**) Example coronal brain section (4.8 mm posterior to Bregma) showing two silicon probe tracks through PrV labeled with DiI (red); cell bodies stained with Nissl (blue).

(**E**) Example extracellular voltage trace from one recording session. Colored bars indicate wall-passing periods.

Figure 2. PrV contains multiple neural codes for peri-head distance

(**A**) Example spike rasters of six single units (units 1-6) in PrV during repeated wall pass trials. Each row corresponds to one wall pass trial, and each dot corresponds to a time bin (50 ms) containing at least one spike, colored according to wall distances. Wall passing begins at time 0. Wall contact (grey bar) begins between 2-3 seconds and ends between 9-11 seconds, depending on the wall distance and whisker length. Scale bar: 20 passes.

(**B**) Peri-event time histograms (PETHs) computed as spike density functions aligned to wall onset time (t=0 s), for the same example units shown in A. Error shading: SEM.

(**C**) Z-scored tuning curves (mean ± 95% CI) to wall distance for same example units shown in A, B.

Figure 2 (continued)

(**D**) Heatmap of normalized tuning curves for wall-activated, distance-tuned units (n=472 from 36 mice), sorted by preferred wall distance from close (top) to far (bottom). Tuning curves were linearly interpolated between presented distances; unpresented distances are shown as white space.

(**E**) Histogram of preferred wall distances, measured as the location of the tuning curve peak, from units in D.

(**F**) Fraction of recorded single units (n=1141) in PrV defined as either untuned (gray), suppressed only (purple), map (yellow, prefer ≥ 8 mm), proximity (pink, prefer < 8 mm), or ambiguous (orange), which could not be statistically defined as either proximity or map (See Supplemental Figure S3).

Figure 3. The map code enables more accurate decoding of peri-head distance
(**A**) Decoding accuracy over time averaged across all distances for population decoder models trained on subsamples of 118 units from either proximity (pink, n=266), map (yellow, n=118), untuned (black, n=550), or trial-shuffled control (gray, n=384) subpopulations. Shaded area: 95% confidence interval. Wall contact occurs around 3-11s (gray bar) after the start of wall movement at t=0 s.
(**B**) Average confusion matrices showing proportion of wall distance decoded for each actual distance from proximity (left) and map (right) populations.
(**C**) Bar plot showing mean ± SEM population decoding accuracy for proximity (pink), map (yellow), untuned (gray), and shuffled control (white); *, $p<0.05$, Wilcoxon rank-sum test with BHFDR correction.
(**D**) Matrix quantifying error, in mm, for each coordinate in the confusion matrix.
(**E**) Bar plot showing mean decoding error ± SEM in mm for decoders trained on each population (same colors as D); *, $p<0.05$, Wilcoxon rank-sum test with BHFDR correction. Error was computed by multiplying the confusion matrix with the absolute value of the error matrix and summing across rows.

Figure 4. Multi-whisker input supports but is not required for the distance map code

(**A**) Top, Example video frame showing multiple whiskers during wall-passing stimulation (wall moving left to right). Middle, same session during whisker cutting. Bottom, same session after cutting to a single B1 whisker.

(**B**) Pairwise comparisons of firing rates (log scale) during wall-passing periods for wall-activated and distance-tuned units (n=86 units from 18 mice) in the full-whisker versus single-whisker condition. Lines connect the same unit across conditions. **, Population mean firing rates: 11.58 ± 2.986 sp/s (full array) vs. 8.18 ± 2.602 sp/s (single whisker), $p = 4.43 \times 10^{-8}$); *, individual neuron, $p < 0.05$; Wilcoxon signed-rank tests.

Figure 4 (continued)

(**C,D**) Two example map units that converted to proximity units after cutting down to a single whisker (B1, top; γ, bottom): (C) Spike density functions (SDFs); (D) corresponding tuning curves.

(**E,F**) Example map unit that retained map-like tuning but shifted its preferred wall distance closer after cutting down to a single C2 whisker.

(**G,H**) Two example map units showing gain modulation after cutting down to a single whisker (C1, top; β, bottom): Top, decreased activity; Bottom, increased activity (disinhibition).

(**I,J**) Two proximity units showing similar gain modulation effects: Top, decreased activity; Bottom, increased activity (disinhibition).

(**K**) Pairwise comparison of preferred wall distances for wall-activated units (n=33 units from 13 mice) that remained distance-tuned after cutting down to a single whisker. Lines connect the same unit across conditions. Open circles: map units; closed circles: proximity units. Color indicates tuning category: proximity (red), ambiguous (orange), map (yellow).

Figure 5. Silencing intersubnuclear inhibitory projections from SpVi diminishes the PrV map code

(**A**) Viral-genetic strategy to express TeLC in inhibitory cells in caudal SpVi, which send long-range projections to PrV.

(**B**) Example coronal section at 4.6 mm posterior to Bregma, showing axons of TeLC-labeled inhibitory cells (TeLC-EYFP, green) in PrV and a silicon probe recording track labeled with DiI (red). Cell bodies were stained with Nissl (blue).

(**C**) Heatmaps of normalized tuning curves for wall-activated, distance-tuned units in PrV. Left, wild-type mice (same data as Fig. 2D, reproduced here for side-by-side comparison). Right, a separate cohort of vGat-Cre mice with TeLC-mediated silencing of PrV-projecting inhibitory neurons (n=252 units from 11 mice). Units are ordered by preferred wall distance, from near (top) to far (bottom). The color bar to the right of each heatmap indicates unit tuning category. Tuning curves were linearly interpolated between presented wall distances; unpresented distances appear as white space.

(**D**) Cumulative distributions of preferred distances for the same units in C, comparing wild-type (black) and SpVi-TeLC (green) populations (p = 8.02 × 10^{-5}, one-tailed K-S test).

(**E**) Comparison of the proportions of distance-tuned units in wild-type (n=1141 from 36 mice) population and in SpVi-TeLC (n=587 from 11 mice) population. TeLC-mediated silencing of PrV-projecting inhibitory neurons in SpVi significantly reduced the prevalence of map units (3.90 vs 10.34%; Z-statistic: 4.64, p = 1.80 × 10^{-6}, one-tailed Z-test) and suppressed units (7.30 vs 10.43%; Z-statistic: 2.12, p = 0.02, one-tailed Z-test), but increased the prevalence of proximity units (32.09 vs 23.31%; Z-statistic: -3.93, p = 4.27 × 10^{-5}, one-tailed Z-test). The proportions of untuned and ambiguous units were not significantly different between samples from wild-type and TeLC mice. *, significance.

Figure 6. Simulation of brainstem circuitry to generate distance-map code

(**A**) Schematic diagram of the modeled neural circuit. The input layer consists of a pool of TG neurons (gray-filled circles), whose tuning to wall distance was modeled as sigmoidal functions with varying thresholds (indicated by the outline color of each circle). These TG neurons project to PrV neurons (black circles, output layer) via two distinct pathways: a direct excitatory (lemniscal) pathway (red) and an indirect inhibitory pathway (blue). The inhibitory pathway is mediated by a population of inhibitory neurons located in SpVi (blue circles), which project to PrV neurons.

(**B**) Top, TG tuning curves (color-coded to match TG neurons outlined in A) are modeled as sigmoidal functions with thresholds spanning the full range of wall distances. Bottom, The sigmoid threshold distribution is exponentially skewed to emphasize the encoding of proximal wall distances.

(**C**) Example tuning curves of modeled PrV neurons. Y-axis: baseline-subtracted activity predicted by the model (A.U.). Wall distances are normalized between 0 and 1, and nine uniformly spaced distance bins (labeled 0-8) are used for analysis and visualization.

(**D**) Left, Heatmap of normalized tuning curves for distance-tuned units (n=590) in the modeled PrV population, sorted by preferred distance bin from close (top) to far (bottom). Right, Same analysis after reducing inhibitory inputs to each PrV neuron by 50%, n=636 distance-tuned units.

(**E**) Left, Histogram of preferred distance bins, defined as the peak (max) of the tuning curve, for modeled PrV neurons. Right, Same analysis after reducing inhibitory inputs to each PrV neuron by 50%.

(**F**) Left, Proportion of modeled PrV neurons classified as proximity (pink, prefer distance bin 0) or map (yellow, prefer > distance bin 0). Right, Same classification following 50% reduction of inhibitory inputs per modeled PrV neuron.

Figure S1. Wall-passing stimulus induced minimal active whisker retraction

(**A**) In a subset of mice (N=4), high-speed video was acquired during the wall-passing paradigm under both awake and anesthetized conditions. Example video frame with tracked whiskers color-coded left-to-right of an awake, behaving mouse: C3 (green), C2 (yellow), C1 (orange), and γ (red). Whisker angle (θ) was measured relative to the mediolateral axis (dotted line); example shown for C1 (orange).

(**B**) Example trace of whisker angle (θ, black) and its midpoint (θ_{mid}, gray) from an awake, behaving mouse; θ_{mid} is the slowly varying centerline of θ.

(**C**) Baseline-subtracted midpoint angle for the C1 whisker of 20 repetitions of one wall distance, aligned to wall movement onset (t=0 s) in an awake, behaving mouse. Baseline was defined during the pre-wall period. Gray bar: wall contact period (3-11 s).

(**D**) Comparison of whisker midpoint angle between awake and anesthetized conditions in the wall-passing paradigm. Peri-event time histograms (PETHs) show changes in baseline-subtracted midpoint angle ($\Delta\theta_{mid}$) for C1, aligned to wall onset (t=0 s) in one mouse. Left: awake, behaving; right: anesthetized. Each trace is the mean of 20 repetitions of each wall distance (color) ±SEM shading.

(**E**) Mean change in $\Delta\theta_{mid}$ versus wall distance for the same C1 whisker shown in D. Green: awake, behaving; blue: anesthetized.

(**F**) Same as E, but averaged across mice (N=4). Subpanels show mean $\Delta\theta_{mid}$ versus wall distance for left: C1, middle: C2, right: Greek whiskers. Lines indicate across-mouse means ±SEM. Green: awake, behaving; blue: anesthetized. Significance reflects mouse-level paired comparisons (awake vs. anesthetized) with Bonferroni correction (α = 0.017): C1, p = 0.03 (n.s.); C2, p = 0.04 (n.s.), Greek: p = 0.0199 (n.s.).

Figure S2. Acute single-electrode recordings from PrV during the wall-passing paradigm

(**A**) Schematic of the experimental setup for acute single-unit recordings in PrV using tungsten electrodes. The wall stimulus moved past the mouse's face at randomized distances ranging from 4 to 20 mm in 1 mm increments (10–20 repetitions per distance). The mouse has C-row whiskers intact.

(**B**) Example coronal brain section at 4.8 mm posterior to Bregma showing one recording track through PrV labeled with DiI (red). Cell bodies were stained with Nissl (blue).

(**C**) Example extracellular voltage trace from one recording session, showing two wall pass events.

(**D**) Left, PETHs computed as spike density functions aligned to wall onset time (t=0 s), for an example PrV unit preferring the closest distance presented. Error shading: SEM. Right, Z-scored tuning curve (mean ± 95% CI) to wall distance for the same unit.

(**E**) Same as D, but for a PrV unit preferring an intermediate distance.

(**F**) Heatmap of normalized tuning curves for all distance-tuned units (n=39 from 14 mice), sorted by preferred distance from close (top) to far (bottom). Tuning curves were linearly interpolated between presented distances; unpresented distances are shown as white space.

Figure S3. Spike sorting quality metrics

(**A**) Inter-spike-interval (ISI) histograms for the example units in Figure 2A-C. The grey shaded region indicates ± 0.8 ms refractory period. Insets show the corresponding mean spike waveforms; shaded areas represent ± STD. Scale bars: 0.5 ms (horizontal), 0.25 mV (vertical).

(**B**) Scatter plot of waveform signal-to-noise ratio (SNR) versus ISI false alarm (FA) rate for wild-type dataset collected with silicon probes (WT-Si Probe; n=1358 total units). The area within the grey square contains acceptable units with SNR > 5 and false alarm rate < 2% (n=1301 units from 36 mice).

(**C**) Z-scored waveforms of a subset of acceptable units from the WT-Si Probe dataset (n=100 wave-forms shown).

(**D**) Histogram of spike widths of all acceptable units from the WT-Si Probe dataset.

(**E**) Scatter plot of spike widths, in ms, versus mean firing rate, in spikes per second, for baseline peri-ods (prior to wall pass) of acceptable units.

(**F-I**) Same as (B-E), but for tetanus toxin (TeLC) dataset (n=665 total units from 11 mice; n=631 accept-able units in green shaded area).

(**J-M**) Same as (B-E), but for tungsten dataset (n=46 total units from 14 mice; n=46 acceptable units in grey shaded area; n=46 waveforms shown in K).

Figure S4. Co-existence of proximity and map codes for encoding peri-head distance
(**A**) Cumulative distribution functions (CDFs; log scale) of baseline firing rates for wall-activated, distance-tuned units (red), wall-suppressed, distance-tuned units (blue), and untuned units (black). Wall-activated units had significantly lower baseline firing rates than untuned units (5.46 ± 0.670 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 1.91 × 10^{-10}, two-tailed K-S test), while wall-suppressed units had firing rates comparable to untuned units (7.31 ± 1.124 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 0.27, two-tailed K-S test).

Figure S4 (continued)

(**B**) Left, CDFs of firing rates for wall-activated, distance-tuned neurons during wall passes (red) versus baseline periods (black). Firing rates were significantly higher during wall contact at the preferred distance (16.44 ± 1.640 vs. 6.36 ± 0.704 sp/s, mean ± SEM; p = 9.26 × 10^{-28}, one-tailed K-S test). Right, CDFs of firing rates for wall-suppressed, distance-tuned neurons during wall passes (blue) versus baseline periods (black). Firing rates were significantly lower during wall contact at the preferred distance (4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s, mean ± SEM; p = 3.06 × 10^{-4}, one-tailed K-S test).

(**C**) CDFs of firing rates for untuned units during wall passes (gray) and baseline periods (black). No significant difference was observed (6.65 ± 0.687 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 0.74, two-tailed KS test).

(**D**) Heatmap of normalized tuning curves for suppressed units (n=213), sorted by preferred wall distance from close (top) to far (bottom). Units that were both activated and suppressed appear both in here and in Figure 2D. Tuning curves are linearly interpolated between distances presented and omitted (white space) for distances not presented.

(**E**) Cross-validated heatmaps of normalized tuning curves for activated units designated as either "map" (n=118) or "proximity" (n=266), for which trials were randomly partitioned into two equal partitions. Units in left panel were sorted by their preferred wall distances (close-top, far-bottom). Units in the right panel were sorted using the same order as the left panel.

(**F**) Histogram of the location of tuning curve troughs from suppressed units in D.

(**G**) Z-scored tuning curve (mean ± 95% CI) to wall distance for an example proximity unit (left), ambiguous unit (middle) determined by (i) not significant on the map test (one-tailed Wilcoxon signed-rank test, p = 0.09; n.s., not significant), and (ii) not having the highest firing rate at the closest distance, and map unit (right) determined by one-tailed Wilcoxon signed-rank tests comparing firing at each non-closest distance to the closest distance (p = 1.27 × 10^{-4}; *, p < 0.05 with BHFDR correction).

Figure S5. Single-neuron decoding performance

(**A**) Confusion matrices showing proportion of predicted wall distances for each actual distance from two example single neurons. Top, Example proximity unit (Unit 1 in Fig. 2A-C). Bottom, Example map unit (Unit 3 in Fig. 2A-C).

(**B**) Mean decoding accuracy (± SEM) of single-neuron decoders grouped by tuning category: proximity (pink, n=266), map (yellow, n=118), untuned (gray, n=550), and trial-shuffled control (white, n=1141). *, $p < 0.05$; n.s., not significant (Wilcoxon rank-sum test with BHFDR correction).

(**C**) Mean decoding error (± SEM) in mm by tuning category (same color scheme as B). Error was computed as the elementwise product of the confusion matrix and absolute error matrix (Fig. 3D), summed across rows. *, $p < 0.05$; n.s., not significant (Wilcoxon rank-sum test with BHFDR correction).

Figure S6. Inhibitory circuit mechanisms contribute to the peri-head distance map code

(**A**) Schematic for expression of the inhibitory opsin, GtACR, in local inhibitory cells in PrV.

(**B**) Schematic of acute silicon probe recording paired with optogenetic inhibition of local inhibitory inter-neurons in PrV. Continuous light stimulation (470 nm, 2 mW) was applied during the entire wall pass with 5s inter-trial interval and randomized laser on/off trials.

(**C**) SDFs (in spikes/second) aligned to wall onset time of an example putative inhibitory neuron displaying laser silencing effect during wall passes for "Laser OFF" (left) and "Laser ON" (right) trials. Error shading: SEM.

(**D**) Example coronal section at 5.1 mm posterior to Bregma, showing GtACR-labeled inhibitory cells (green) and two silicon probe tracks through PrV, outlined with white dashed line. Cell bodies were stained with Nissl (blue).

Figure S6 (continued)

(**E**) Distribution of modulation index (ML) for all acceptable units (n=112) in the GtACR dataset. Units were classified as strongly inhibited (MI ≤ −0.5, dark blue), inhibited (−0.5 < MI < 0, light blue), disinhibited (MI > 0, red) when their bootstrap MI confidence interval excluded zero, or as no change when the confidence intervals included zero (gray).

(**F**) Top, SDFs of an example proximity unit displaying laser-induced disinhibition in Laser ON (right) condition. Bottom, Same but for an example map unit.

(**G**) Z-scored tuning curves for the same units in F. Error bars: 95% CI.

(**H**) Example coronal section at 7.4 mm posterior to Bregma, showing the TeLC-labeled inhibitory cells (TeLC-EYFP, green) in SpVi, outlined with white dashed line. Cell bodies were stained with Nissl (blue).

(**I**) Left, CDFs of baseline firing rates for untuned units in SpVi-TeLC mice (green) and wild-type mice (black). Firing rates were significantly higher in SpVi-TeLC mice (9.34 ± 1.330 vs. 6.90 ± 0.682 sp/s; p = 7.63 × 10^{-5}, one-tailed K-S test). Right, CDFs of mean firing rates for the same populations during wall passes. Untuned units in SpVi-TeLC mice showed significantly elevated firing compared to wild-type mice (9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s; one-tailed K-S test, p = 1.06 × 10^{-4}).

(**J**) Left, CDFs of baseline firing rates for wall-activated, distance-tuned units in SpVi-TeLC mice (green) and wild-type mice (black). No significant difference was observed (5.60 ± 0.845 vs. 5.46 ± 0.670 sp/s; one-tailed K-S test, p = 0.14). Right, During wall contact at the preferred distance, these same SpVi-TeLC units exhibited significantly elevated firing (20.25 ± 2.263 vs. 16.44 ± 1.640 sp/s; one-tailed K-S test, p = 1.64 × 10^{-3}). *, p < 0.0125 (Bonferroni-corrected, four comparisons); n.s., not significant.

STAR* METHODS

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Fan Wang (fan_wang@mit.edu)

Materials Availability

This study did not generate new unique reagents.

DATA AND SOFTWARE AVAILABILITY

MATLAB scripts and analysis code used in this study will be made available on Github, and processed datasets in NWB (Neurodata Without Borders) format are deposited on DANDI.

EXPERIMENTAL MODEL DETAILS

Adult C57BL/6J (JAX stock #000664) and vGat-IRES-Cre (JAX stock #016962; Slc32a1$^{tm2(cre)Lowl/J}$) mice were obtained from the Jackson Laboratory. During behavioral and electrophysiological experiments, mice were singly housed under a 12-hour light/dark cycle with ad libitum access to food and water. All experiments were conducted during the light phase. Details regarding the sex and age of all animals are provided in Table S1. All procedures were performed in accordance with protocols approved by the Institutional Animal Care and Use Committees at the Massachusetts Institute of Technology.

METHOD DETAILS

Headpost Implantation and Craniotomy

Mice were anesthetized with 3% isoflurane in 0.8 L/min oxygen and positioned in a stereotaxic frame (Model 963, David Kopf Instruments). Anesthesia was maintained at 1.5% isoflurane throughout the surgery. A custom stainless steel headbar (Pololu; 0.060″ ± 0.004″, type 304) and a stainless-steel ground screw were affixed to the skull using C&B Metabond (Parkell, #S380) and reinforced with dental cement (Stoelting). Mice were allowed at least seven days to recover following head-post implantation surgery. On the electrophysiological recording day, a 1 mm × 1 mm craniotomy was created 4.7 mm posterior and 2.0 mm lateral to Bregma using a dental drill (Aseptico). The dura was carefully removed to provide access to the principal trigeminal nucleus (PrV). Craniotomies were acutely sealed with silicone elastomer (Kwik-Cast, WPI).

Viral Vectors

The AAVs used included AAV-EF1a-DIO-GtACR2-EYFP (Takatoh & Prevosto et al., 2021), AAV-hSyn-FLEX-TeLC-P2A-EYFP-WPRE (Addgene, #135391).

Viral Injections

For optogenetic silencing experiments, vGat-Cre mice received stereotaxic injections of 300 nl AAV-EF1a-DIO-GtACR2-eYFP into PrV at coordinates 4.7 mm posterior, 2.0 mm lateral to Bregma, and 4.5 mm below the brain surface. For chronic silencing of SpVi neurons projecting to PrV, 300 nl of either AAV-hSyn-FLEX-TeLC-P2A-EYFP-WPRE was stereotaxically injected into caudal SpVi (7.4 mm posterior, 2.0 mm lateral to Bregma, and 5.6 mm below the brain surface).

573 Injections were performed using beveled glass micropipettes connected to a microsyringe pump
574 (UMP3, WPI) controlled by a Micro4 controller (WPI) at a rate of 30 nl/min. Following injections,
575 craniotomies were sealed with bone wax (Surgical Specialties, #902), and the scalp was closed
576 using nonabsorbable silk sutures (LOOK, #774B).
577
Wall-Passing Paradigm
After recovering from headpost implantation, mice were head-fixed using plate clamps (Thorlabs)
positioned over a custom linear treadmill (Severson & Xu et al., 2017) to promote running and
whisking. Mice were habituated to the treadmill for at least two days, followed by at least three
days of habituation on a wall-pass apparatus prior to recording. The passing wall was driven in
the mediolateral and anteroposterior dimensions by two linear actuators (Zaber, Models
LSM200B-T4 rev 2 and LSM050B-T4 rev 2). During each pass, the wall moved from anterior to
posterior across the mouse's face at a constant velocity of 10 mm/s. In the silicon probe
experiments, wall distance from the whisker pad (C2 follicle) was randomly varied across trials
between 7 and 23 mm in 2-mm increments (4-mm increments for optogenetic silencing
experiments), using a custom MATLAB script. In the tungsten electrode experiments, wall
distance was randomly varied between 4 mm and 20 mm in 1 mm increments. All experiments
were performed in the dark. After the mediolateral Zaber motor set the wall distance, a
randomized delay of 0.1 to 1 s preceded wall movement.
592
Each recording session in the silicon probe experiments included 15-20 repetitions per wall
distance, during which mice freely whisked and interacted with the wall. Tungsten electrode
experiments included 10-20 repetitions per distance. Whiskers were illuminated using an infrared
LED backlight (Roithner Lasertechnik, #LED660N-66-60) and imaged with a high-speed camera
(Basler, Model acA1300-200um) equipped with a telecentric lens (Edmund Optics, #55-349). Prior
to each recording, the wall apparatus was aligned to the midline of the mouse's head, and a
calibration image was acquired and converted to physical units (31 pixels/mm). The closest wall
distance was defined as the minimum Euclidean distance between the C2 whisker follicle and the
wall. Absolute wall distance (in mm) was calculated by adding the calibrated scaling factor to the
horizontal position (in 2-mm steps, relative to the closest position) of the stepper motor controlling
the wall's mediolateral position.
604
High-Speed Videography
High-speed whisker videos (688 × 688 pixels at 500 Hz) were acquired with a machine-vision
camera (Basler acA1300-200um) controlled by custom Python software (CamPy). Illumination
was provided by a 940 nm LED (Roithner Lasertechnik) collimated with a condenser lens and
diffused to form an even backlight. The whiskers appeared in silhouette against this background
and were imaged via a front-surface mirror into a 0.25× telecentric lens (Edmund Optics).
Frame timing was triggered by a Teensy microcontroller, and TTL pulses were recorded
alongside electrophysiology to synchronize spikes and video.
613
Awake and Anesthetized Wall-Passing Recordings
In a subset of mice (N = 4), high-speed video was continuously acquired during the wall-passing
paradigm under both awake and anesthetized conditions. Mice were head-fixed throughout,

keeping their position constant relative to the moving wall stimulus. To enable whisker tracking,
all whiskers except a single row (C1-C4 and γ or β) were trimmed. While awake, each mouse
experienced wall passes with randomized wall distances (7-23 mm in 2-mm increments),
repeated for 20 passes per distance. Following the awake session, the same animals were
anesthetized via intraperitoneal injection of a fentanyl-based cocktail (0.05 mg/kg fentanyl,
5 mg/kg midazolam, 0.5 mg/kg dexmedetomidine). After a 10-minute induction period, the same
wall-passing protocol was repeated under anesthesia. Comparisons between awake and
anesthetized conditions were performed within-subject across matched wall distances.

Electrophysiology Recording: Silicon Probe

Following habituation to the wall-passing paradigm, mice were head-fixed and allowed to run on
the treadmill. The craniotomy was exposed and covered with artificial cerebrospinal fluid (aCSF).
For wild-type and TeLC experiments, the tip of a 32-channel silicon probe (NeuroNexus, A1x32-
Poly2-10mm-50s-177-A32 or A1x32-Poly3-10mm-50s-177) was coated with the fluorescent dye
DiI (Invitrogen™, #V22885). For optogenetic silencing experiments, a 473 nm laser (Cobolt, 06-
MLD 473) was connected to a ferrule attached to a 32-channel silicon optrode (NeuroNexus,
A1x32-Poly2-10mm-50s-413, 105 µm/125 µm, 0.22 NA). The probe or optrode was advanced
approximately 5 mm into the brain using a micromanipulator (Sutter Instruments, Model MPC-
325) until it reached ventral PrV and well-isolated units responsive to manual whisker stimulation
were encountered. The brain was allowed to relax for at least 40 minutes to ensure stable
recordings. Neural signals were hardware-filtered on the headstage (Blackrock Microsystems,
Cereplex M, #10532) between 0.3 Hz and 7.5 kHz for spikes and digitized at 30 kHz using
Cereplex Direct software (Blackrock Microsystems, Version 7.0.5). Analog voltage signals from
the anterior-posterior stepper motor were simultaneously acquired at 30 kHz. Wall pass start and
end times were determined from the rising and falling edges of these voltage signals to
synchronize electrophysiological recordings with wall movement. Recording sites were
subsequently confirmed by post hoc histological analysis.

Electrophysiology Recording: Single Tungsten Electrode

One day prior to recording, whiskers and facial fur on the left side of the face were trimmed using
fine forceps and microdissection scissors (FST, #91500-09) under isoflurane anesthesia (1.5%),
sparing the C-row whiskers (γ or β, C1–C5). On the recording day, mice were head-fixed and
allowed to run on the treadmill. The craniotomy was exposed and kept covered with aCSF. The
tip of a tungsten microelectrode (0.005″ diameter, 5 MΩ, 8° tip angle; A-M Systems, #575300)
was coated with the fluorescent dye DiI and lowered approximately 5 mm into the brain using a
micromanipulator until it reached ventral PrV. Electrical activity was monitored via audio output
from the amplifier (WPI, DAM80). Manual whisker stimulation was performed to identify a well-
isolated, whisker-responsive unit. The brain was allowed to relax for at least 20 minutes to achieve
stable recordings. Differential signals between the recording and reference electrodes were
amplified 10,000× and bandpass filtered between 300 and 3,000 Hz (WPI, DAM80). Following
recording, receptive fields and response types (rapidly adapting [RA] or slowly adapting [SA])
were manually classified. Analog voltage signals from the amplifier and the anterior-posterior
stepper motor were acquired using Cereplex Direct at a sampling rate of 30 kHz. Wall pass start
and end times were extracted from the rising and falling edges of the stepper motor TTL signals

to synchronize electrophysiological recordings with wall movement. All recording sites were
confirmed by post hoc histological analysis.

Isolation of Single Unit Activity
For silicon probe data, initial single-unit isolation (spike sorting) was performed using Kilosort2.5
(Pachitariu et al., 2016), followed by manual curation in Phy (github.com/cortex-lab/phy; Rossant,
2025). Spike times and waveforms for isolated clusters were exported from Phy. Most units
exhibited narrow spike widths, defined by a trough-to-peak latency of less than 0.4 ms. Duplicate
units, identified as those with high Pearson's correlation coefficients (> 0.1) in spike counts binned
at 1 ms resolution, were excluded from further analysis. For tungsten electrode data, spike
waveforms were extracted by thresholding high-pass filtered traces (500 Hz) and clustered using
MClust-4.4 (A.D. Redish, http://redishlab.neuroscience.umn.edu/MClust/MClust.html).

Isolation quality was assessed using the following metrics: signal-to-noise ratio (SNR), defined as
the mean waveform amplitude divided by the standard deviation (SD) of waveform noise; and
false alarm (FA) rate, defined as the percentage of spikes with inter-spike intervals (ISI) shorter
than 0.8 ms. Units with an SNR less than 5 SDs or an FA rate greater than 2% did not meet single-
unit isolation quality criteria and were excluded from subsequent analyses. For silicon probe data,
in wild-type mice, 1141 of 1358 units (36 mice), in TeLC experiments, 587 of 665 units (11 mice),
and for Tungsten recordings (46 of 46 units (14 mice), passed isolation quality and minimum firing
rate (>1 Hz) criteria and were subsequently entered into population analyses. For acute whisker
cutting experiments, only units demonstrating stable waveform properties both before and after
cutting were included in analyses.

Acute Whisker Cutting Manipulation
During recording sessions involving acute whisker cutting, neural responses were first recorded
while the full whisker array was intact. Each session included 15 to 20 repetitions of wall passes
over approximately one hour. To best guess the primary whisker to spare during acute trimming,
the receptive fields of PrV neurons were mapped by manually stimulating individual whiskers with
a cotton-tipped applicator while monitoring spike responses. All whiskers except the designated
spared whisker were then cut down to the base of the shaft using microdissection scissors (FST,
Cat# 91500-09). After trimming, the spared whisker and surrounding whisker pad were brushed
with a cotton-tipped applicator to stimulate all touch-sensitive recorded units. The wall-passing
procedure was then repeated for an additional 15 to 20 trials. After completing the post-cutting
wall passes, the whisker and pad were brushed once more to confirm the continued presence of
responsive units. Electrophysiological signals were continuously recorded throughout the pre-cut,
cutting, brushing, and post-cut stimulation periods.

Optogenetic Silencing of PrV Local Interneurons
Light pulses were generated using a pulse generator (Doric, #OTPG-4), with laser timing
synchronized to wall movement via Doric software. Continuous 473 nm light stimulation (2 mW)
was delivered by a laser (Cobolt, #06-MLD 473) throughout each wall pass, with a 5-second inter-
trial interval. Each recording session consisted of 30 to 40 repetitions at each wall distance,

ranging from 7 to 23 mm in 4 mm increments, conducted over approximately two hours with
randomized "Laser Off" and "Laser On" trials.
706

Histology
Following recordings, mice were deeply anesthetized and transcardially perfused with phosphate-
buffered saline (PBS), followed by 4% paraformaldehyde in PBS. Brains were post-fixed overnight
and then cryoprotected in 30% sucrose in PBS at 4°C. Tissue was embedded in OCT compound
(Sakura Finetek) and coronally sectioned at 100 µm using a cryostat (Leica Biosystems). For wild-
type, optogenetic inhibition, and tungsten recording experiments, sections containing PrV were
stained with Neurotrace Blue (1:500; Thermo Fisher Scientific, #N21479) in 0.3% Triton X-
100/PBS. For TeLC experiments, sections containing PrV and SpVi underwent immunostaining
with goat anti-GFP (1:500; Rockland, #600-101-215) in blocking buffer, followed by either Alexa
Fluor 488 (1:500; Jackson ImmunoResearch, #705-545-147) or Alexa Fluor 647 (1:500; Jackson
ImmunoResearch, #705-606-147) donkey anti-goat secondary antibodies and Neurotrace Blue
(1:500; Thermo Fisher Scientific, #N21479) in blocking buffer. The sections were then briefly
rinsed with 0.3% Triton X-100/PBS and mounted on slides with a custom-made mounting medium.
All histological images were acquired with either a Keyence microscope (BZ-X800) or a Zeiss
LSM700 confocal microscope.
722

QUANTIFICATION AND STATISTICAL ANALYSIS
All analyses were performed in MATLAB (MathWorks, versions 2021b and 2023b). Data are
reported as mean ± standard errors of the mean (SEM) unless otherwise noted. Measures of
central tendency, variability, test statistics, p-values, degree of freedom, and figure panel
assignments for all analyses are detailed in Table S6. We analyzed single-unit
electrophysiological recordings from n = 1,187 units in PrV across 63 mice, as well as high-speed
video recordings from four sessions in four mice. The video data comprised 15,679,458 frames,
totaling 522.6 minutes. Sample sizes were not predetermined.
731

Video Analysis
The backbone of each whisker was tracked at subpixel-resolution using the Janelia Whisker
Tracker (Clack et al., 2012), yielding a set of "trajectories" (tracked objects in image X-Y
coordinates) for each frame. All subsequent processing of whisker angle (θ) was conducted in
MATLAB according to published methods (Pammer & O'Connor et al., 2013; Severson & Xu et
al., 2017), with several modifications described below in "Video analysis" subsections.
738

Video Analysis: Identifying Tracked Whiskers
To identify the same whiskers across frames (linking), whiskers were ordered by follicle x-position.
Trajectories with follicle x- or y-position outside the region of interest (bounding box), with arc
length shorter than 150 pixels (4.65 mm), or whisker "score" (Janelia whisker tracker
measurements) less than 600 were ignored. Post-hoc median filtering (3-frame window) was
applied to smooth over single-frame linking errors.
745

Video Analysis: Face Masking
Computing time series of whisker angle relies on being able to reliably track the base of the
whisker near the follicle location across all video frames. As previously described (Pammer &
O'Connor et al., 2013; Severson & Xu et al., 2017), we used a dynamic "mask" to truncate the
tracked whisker trajectories near the face to remove "noise" due to occlusion by facial fur. The
follicle location was estimated by extrapolating past the intersection of the whisker and the mask
using the angle at the whisker base (Pammer & O'Connor et al., 2013). We fit a separate mask
for each frame, obtained using a custom algorithm that fitted a smoothing spline to the contour of
the face. We blurred the image using a Gaussian filter, binarized the image, and detected the
contour by taking a spatial derivative along the y-dimension. The pixel locations of the facial
contour were thus detected as the intersection between the white background and black shadow
cast by the face. A smoothing spline was then fitted to the facial contour positions and shifted
upward by ~10 pixels to yield a 20-point facemask that intersected with the whisker trajectories.
Frames in which a tracked whisker did not intersect the facemask were considered missing data.
760
Data Analysis: Wall-Induced Whisker Retraction, Awake vs Anesthetized
For each of the tracked whiskers (C1, C2, and Greek), whisker angle (θ) was measured relative
to the mediolateral axis of the face. The midpoint of the whisker angle trace (θ_{mid}) was defined as
the slowly varying centerline of θ. To obtain θ_{mid}, θ was first smoothed using a Savitsky-Golay filter
(21 frame, third order), followed by interpolation of the peaks and valleys to obtain the upper and
lower envelope of the θ trace. θ_{mid} was then calculated as the mean of this envelope. For each
wall pass, θ_{mid} was computed during both the baseline period (-4.5 to 0 s, with t=0 marking wall
onset) and the wall-passing period (3 to 11 s). Baseline-subtracted θ_{mid} ($\Delta\theta_{mid}$) was then calculated
for each pass. For each mouse (N = 4 mice), $\Delta\theta_{mid}$ was averaged across wall distances and
repetitions to obtain a single value per whisker and condition (awake or anesthetized). Data were
aggregated across mice into a summary table (mouse × whisker × condition), and paired
comparisons were made between awake and anesthetized conditions for each whisker.
773
To compare the effect of wall-induced whisker retraction of awake versus anesthetized conditions,
we performed two-tailed paired t-tests across mice (within-subjects) for each whisker. The null
hypothesis was that mean $\Delta\theta_{mid}$ was equal in awake and anesthetized conditions. Effect size was
computed as the mean difference between conditions (awake minus anesthetized), and
Bonferroni correction was applied to control for multiple comparisons across whiskers
(α =0.0167).
780
Data Analysis: Spike Density Function Calculation
Spike density functions (SDFs) were computed by aligning spike times to the onset of wall-pass
trials, with time zero defined as the moment when the wall began moving toward the mouse. For
all trial conditions, two epochs were defined consistently: a baseline period from −4.5 to 0 s, and
a wall-pass period from 3 to 11 s. Spike rasters were generated from the aligned spike times using
1 ms bins. SDF curves were then produced by convolving these spike rasters with a Gaussian
filter (5 ms SD, full kernel). SEMs were calculated as the SD of the SDF across trials divided by
the square root of the number of trials in each condition. The resulting SDFs and SEMs were
averaged across repetitions of the same wall distance and stored for each cell.

Data Analysis: Wall Distance Tuning Calculation

Wall distance tuning curves were generated by uniformly binning distance measurements into nine 2-mm-wide bins with centers at 7, 9, 11, 13, 15, 17, 19, 21, and 23 mm. Distances falling outside the bin edges were excluded. Peri-event time histograms (PETHs) were computed by binning instantaneous firing rates in 200-ms intervals, aligned to the onset of anterior-posterior wall movement as defined by the stepper motor TTL pulse (t=0). Each wall pass trial lasted approximately 14 seconds. Average firing rates during the wall pass period (3–11 s) were computed from the PETH and compared to baseline firing rates calculated from −4.5 to 0 s. For each distance, mean firing rates were averaged across repeated passes to construct the tuning curve. Standard errors were calculated by dividing the SD across repetitions by the square root of the number of repetitions for each distance. CIs were estimated using bootstrap resampling (10,000 iterations) and defined as the 95th percentile range. Z-scored tuning curves were computed by subtracting the mean from firing rate across all baseline periods in the recording session and dividing by the corresponding SD. Uniform binning enabled concatenation of data across sessions for population-level analyses, including decoding analyses described later. Sessions with fewer than nine presented distances were excluded from population analyses. For population heatmaps, each neuron's tuning curve was normalized to its own minimum and maximum firing rates, and the color scale represents the resulting normalized mean spike rate.

Data Analyses: Characterizing Single Unit Responses

To assess responsiveness to wall stimulation, one-tailed Wilcoxon signed-rank tests ($p < 0.05$) were performed on each of 1,187 units. A unit was classified as "wall-activated" if its mean firing rate during the wall pass period (3-11 s) at any distance was significantly higher than its baseline firing rate (−4.5-0 s). Conversely, a unit was classified as "wall-suppressed" if its mean firing rate during the wall pass period at any distance was significantly lower than baseline. Units exhibiting significant increases at some distances and decreases at others were classified as both "wall-activated" and "wall-suppressed."

Wall distance tuning was assessed using two-tailed Wilcoxon signed-rank tests across all pairs of distances. A unit was classified as "distance-tuned" if the mean firing rate at one or more distances significantly differed from that at other distances ($p < 0.05$, corrected for multiple comparisons using the Benjamini-Hochberg false discovery rate method).

For comparisons of baseline firing rate distributions between distance-tuned and untuned units, the mean and 95% confidence intervals (CI) were calculated using bootstrap resampling (10,000 iterations). The CI was determined by calculating half the range between the upper and lower bounds of the bootstrap interval. Results were reported as mean ± CI. Statistical differences between distributions were assessed using two-sample Kolmogorov-Smirnov (K-S) tests ('kstest2' function in MATLAB), performed with one- or two-tailed hypotheses as dictated by the experimental design (see Table S6). Significance was determined at $p < 0.05$, with Bonferroni correction for multiple comparisons. The same procedures were applied for comparing average firing rates during wall passes and baseline periods between distance-tuned and untuned populations.

To further characterize tuning profile within "wall-activated" and "distance-tuned" units, we
performed one-tailed Wilcoxon signed-rank tests comparing mean firing rates at each non-closest
distance (≥ 9 mm) to the closest distance (7 mm). Units with significantly higher firing rates at any
non-closest distance were classified as "map units," while those exhibiting maximal firing at the
closest distance (7 mm) were classified as "proximity units." Units that met criteria for "wall-
activated" and "distance-tuned" but did not fulfill the criteria for either map or proximity were
labeled "ambiguous" and excluded from decoding and population cross-validation analyses.

For analyses of acute whisker cutting experiments, mean firing rates during baseline and wall-
pass periods were compared between full whisker array (control) and single-whisker (cut)
conditions. Pairwise comparisons of firing rates were conducted using two-tailed Wilcoxon signed-
rank tests. To assess individual neuron responses to whisker cutting, the proportions of units
exhibiting significant increases, decreases, or no changes in firing rates were computed.
Significance thresholds were corrected for multiple comparisons using Bonferroni adjustment. To
evaluate time-dependent plasticity, firing rates for each unit were partitioned into three temporal
segments: the first half of wall-pass trials (full whisker array), the second half (full whisker array),
and the post-manipulation period. For each unit, mean firing rates were computed separately for
these periods. Adaptation was assessed by comparing firing rates between the first and second
halves using two-tailed Wilcoxon signed-rank tests. A non-significant result ($p > 0.05$) indicated
stable firing prior to whisker cutting. Firing rates from the second half of the full whisker array
period were then compared to post-cutting periods using two-sided Wilcoxon signed-rank tests,
where significant differences indicated effects attributable to whisker cutting.

To compare the proportions of map, proximity, ambiguous, suppressed, or untuned cells between
SpVi-TeLC mice and wild-type mice, a one-tailed Z-test for proportions was performed. The null
hypothesis (H_0) stated that the proportion of each cell type in the SpVi-TeLC population (p_2) was
less than or equal to that in the wild-type population (p_1), while the alternative hypothesis (H_a)
posited that $p_2 > p_1$. Proportions (p_1 & p_2) were calculated as the number of cells of a given type
(x_1 & x_2) divided by the total number of recorded units (n_1 and n_2) in each group. A pooled
proportion (p_{pooled}) was used to estimate the standard error (SE) of the difference between groups:

865
$$p_{pooled} = \frac{x_1 + x_2}{n_1 + n_2}, \quad SE = \sqrt{p_{pooled}(1 - p_{pooled})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$$

866 The Z-statistic was computed as:

867
$$Z = \frac{p_1 - p_2}{SE}$$

P-values were calculated using MATLAB's cumulative normal distribution function ('normcdf' in
MATLAB) and evaluated at $\alpha = 0.05$. A p-value below this threshold indicated a significantly higher
proportion of the specified cell type in SpVi-TeLC mice compared to wild-type mice.

To compare the firing rate distributions between SpVi-TeLC and wild-type populations (Group 1
and Group 2), we computed the mean and 95% CI for each group using bootstrap resampling
(10,000 iterations). For each group, the results were reported as mean ± CI. To assess the
statistical significance of differences between the distributions of the two groups, we performed a

two-sample KS test. The test was conducted with a specified directional hypothesis (one- or two-
tailed, depending on the experimental design), and the resulting p-value was reported to
determine the significance ($p < 0.05$, with Bonferroni correction for multiple comparisons) of the
observed differences.
880
Data Analysis: Decoding of Wall Distance from Population or Single Neurons Activity
For population decoding of wall distance, we applied a population decoder (Neural Decoding
Toolbox, www.readout.info; Meyers, 2013; Vincis & Chen et al., 2020). Units from sessions with
all nine distances presented that passed spike sorting quality metrics (1,141 total units) were
concatenated into a pseudo-population. Populations were subsampled into sub-populations of
equal size (118 units), containing units designated as either "map", "proximity", or "untuned" cells.
Units in the "shuffled" subpopulation were randomly sampled from map and proximity
subpopulations and had their trial distance labels shuffled with replacement. Spike timestamps
for each unit were aligned to the onset of anterior-posterior wall movement (time 0) and were
binned using a 50 ms bin size with a 150 ms sliding window to create a firing rate matrix, where
each row corresponds to a trial and each column corresponds to a time bin. The firing rate matrix
spans from time 0 to 14 seconds. To ensure equal trial numbers across conditions (i.e., wall
distances), trials were subsampled five times to match 15 trials per condition. The binned data
and their corresponding condition labels were randomly split into 10 subsets—9 used for training
the classifier and 1 reserved for validation. The process was repeated 10 times for each unique
partition for 10-fold cross-validation to assess decoding accuracy. The classifier operates by
learning a mean population vector (template) for each class from the training set. During testing,
it calculates the Pearson correlation coefficient between a test point and the learned templates.
During validation and testing, the class with the highest correlation value is assigned as the
predicted label. Decoding accuracy (true positive rate) was averaged over the 3-11 second wall-
pass period to obtain the overall decoding accuracy for the sampling epoch. In addition to the
decoding accuracy, confusion matrices were computed at each time step. To assess variability,
the entire subsampling procedure was repeated 10 times, and standard error was calculated by
computing the SD of decoding accuracy across runs and dividing by the square root of 10.
905
For single-neuron decoders, we quantified how well individual neurons discriminated wall
distance. For each unit, spike trains were grouped according to nine discrete wall distances (7, 9,
11, 13, 15, 17, 19, 21, and 23 mm). For each distance, spike times were aligned to trial onset and
converted into spike count vectors using 200 ms non-overlapping bins spanning the 3–11 s wall
contact period. These binned firing rate profiles were used to construct condition-specific
templates for distance classification. Decoding was performed using a Euclidean distance
classifier, which assigned each test trial to the distance condition whose mean firing-rate vector
(template) was closest in Euclidean space. Cross-validation was implemented as a leave-two-out
procedure (xval = 2), in which two trials per condition were held out as test data, and the remaining
trials were used for training. This was repeated exhaustively across all combinations of held-out
pairs, such that every trial pair served as a test set at least once. Classification accuracy was
computed as the fraction of correctly identified test trials, averaged across all cross-validation
iterations. For shuffled controls, we performed a trial-shuffled permutation in which class labels
were randomly reassigned across conditions (n = 1000 iterations).

For both population and single-neuron decoders, differences in mean decoding accuracy and
error across wall distances were assessed using Wilcoxon rank-sum tests. Separate tests were
performed for each of the nine wall distances, and p-values were corrected for multiple
comparisons using BH-FDR to compare proximity and map populations.
924

Data Analysis: Acute Whisker Cutting Manipulation
To test for the effect for acute whisker cutting on firing rates, we computed each unit's mean firing
rate during baseline and wall-passing periods on each repetition for the full whisker array (control)
or single-whisker condition (manipulation). Pairwise comparison of mean firing rate for neural
population and individual units between conditions was performed via two-sided Wilcoxon signed-
rank test. To evaluate how individual neurons responded to whisker cutting, we computed the
proportion of units that exhibited significant increases, decreases, or no change in firing rate
between conditions. P-values were corrected for multiple comparisons across units using the BH-
FDR procedure.
934

935 ***Data Analysis: Modulation Index in Acute Optogenetic Experiments***
936
937 We quantified the acute optogenetic effects on firing rates using a modulation index (MI)
938 computed for each unit as:

939
$$MI = \frac{\text{FR}_{LASER\ ON} - \text{FR}_{LASER\ OFF}}{\text{FR}_{LASER\ ON} + \text{FR}_{LASER\ OFF}}$$

940
where $\text{FR}_{LASER\ ON}$ and $\text{FR}_{LASER\ OFF}$ denote the mean firing rates during matched laser-on and laser-
off wall-passing periods, respectively. For each unit, we obtained a bootstrap CI on the MI and
classified the unit as significantly modulated if the CI did not include zero. Significantly modulated
units were further assigned to one of three categories based on their MI: strongly inhibited (MI ≤
-0.5), inhibited (-0.5 < MI < 0), or disinhibited (MI > 0). Units whose MI confidence intervals
overlapped zero were classified as no change. All analyses were restricted to units that passed
our unit inclusion criteria (see Isolation of Single Unit Activity).
948

Computational Simulation of Brainstem Circuitry
To model the tuning properties of PrV neurons, we simulated a pool of TG afferents whose
activities decreased monotonically with wall distance, consistent with their mechanical sensitivity.
These proximity-based TG inputs projected to PrV neurons via a direct excitatory (lemniscal)
pathway and an indirect inhibitory pathway (SpVi inhibitory projections to PrV). TG tuning curves
were modeled as sigmoidal functions with thresholds spanning the distance range. The activity of
each PrV neuron j (v_j^{PrV}) in response to wall distance $x \in [0,1]$ was defined as:

956
$$v_j^{\text{PrV}} = f\left(\sum_{i \in C_j^E} w_{ij}\sigma_{\text{TG}}(x; \beta_i, \theta_i) - \sum_{l \in C_j^I} w_{lj}\sigma_{\text{TG}}(x; \beta_l, \theta_l) + I_j^{bg}\right)$$

957 where

958
$$\sigma_{\text{TG}}(x; \beta, \theta) = \frac{1}{1 + \exp(\beta(x - \theta))}$$

is the TG neuron's sigmoidal tuning function. Here, β and θ control sensitivity and selectivity to

wall distance, respectively. $f(x) = \max\{x, 0\}$ is the non-linear activation function for PrV neurons

and $I_j^{bg} \sim U(0,1)$ is the background excitatory input to PrV neuron j. C_j^E denotes the set of TG

neurons providing direct excitatory input to PrV neuron j, with synaptic weight $w_{ij} \sim U(0, |C_j^E|^{-\frac{1}{2}})$;

C_j^I denotes TG neurons projecting indirectly to PrV neuron j via inhibitory neurons in SpVi, with

weight $w_{lj} \sim U(0, |C_j^I|^{-\frac{1}{2}})$. Subscript i and l denote each TG neuron in set C_j^E and set C_j^I. For

simplicity, inhibitory neurons were modeled as providing a sign-inverted copy of TG activity, such

that v_j^{PrV} reflect a linear combination of excitatory and inhibitory streams.

Intuitively, the subtraction of two sigmoidal functions can yield a non-monotonic, Gaussian-like

function. Here we generalize this by considering a pool of sigmoidal functions (TG neurons).

Specifically, we implemented several assumptions:

(1) Sparse connectivity: Both C_j^E and C_j^I represented small subsets of TG neurons.

972   Parameter $\beta_i, \theta_i, \beta_l, \theta_l, w_{ij}, w_{lj}$ were independently sampled for each PrV neuron. The

973   number of inhibitory inputs $|C_j^I|$ per PrV neuron j was treated as a random variable

974   sampling from an exponential distribution parametrized by λ_c (truncated to the interval

975   [1,30] to preserve E/I balance):

976      $$P(|C_j^I| = k) = \frac{e^{-\lambda_c k}}{\sum_{m=1}^{30} e^{-\lambda_c m}}, k = 1,2,3, \ldots 30, \qquad \lambda_c = 0.5$$

977   The number of excitatory inputs was fixed at $|C_j^E| = 20$.

978   (2) Skewed distribution of TG tuning. The TG threshold parameter θ was drawn from a

979   truncated exponential distribution within [0,1]:

980      $$\theta \sim f(x; \lambda_\theta) = \frac{\lambda_\theta e^{-\lambda_\theta x}}{1 - e^{\lambda_\theta}}, \qquad x \in [0,1]$$

981   where λ_θ describes the skewness of the distribution. This skewed distribution increases

982   the density of high-threshold TG neurons that only fire at proximal distances, thereby

983   enhancing encoding resolution and reducing overall metabolic cost (efficient coding). We

984   used $\lambda_\theta = 2$ for the excitatory inputs and $\lambda_\theta = 4$ for the more skewed inhibitory pathway.

985   Sensitivity parameters were sampled as $\beta_i, \beta_l \sim N(20, 5)$. Untuned neurons were modeled

986   by explicitly weakening all weights for a fraction of neurons to 1% (i.e., p(untuned) = 0.4).

The input (distance) range x was normalized to [0,1] for simplicity. We sampled 99 uniformly

spaced points corresponding to wall-distance presentations. To evaluate tuning curves, we

averaged simulated activity into nine discrete distance bins [0, …, 8]. All neurons with flat

tuning curves (max – min < 0.1) were considered untuned. Neurons with max activity below

zero were considered suppressed. Among the remainder, proximity cells were defined as

neurons with maximal responses at x = 0, while map cells were defined as neurons with peak

responses at intermediate or far distances (x > 0). In total, we modeled 1,225 PrV neurons.

To simulate partial silencing of inhibitory projections, the distribution of inhibitory inputs was
truncated to approximately 50% of its original range [1,15], such that $\left|C_j^I\right|$ for each PrV neuron
j is decreased by:

998
$$P\left(\left|C_j^I\right| = k\right) = \frac{e^{-\lambda_c k}}{\sum_{m=1}^{16} e^{-\lambda_c m}}, \qquad k = 1,2,3, \dots 15.$$

999
1000
1001　**STAR* METHODS REFERENCES**

56. Clack, N.G., O'Connor, D.H., Huber, D., Petreanu, L., Hires, A., Peron, S., Svoboda, K., and Myers, E.W. (2012). Automated tracking of whiskers in videos of head-fixed rodents. *PLoS Comput. Biol.* **8**, e1002591.
57. Meyers, E.M. (2013). The neural decoding toolbox. *Front. Neuroinform.* **7**, 8.
58. Pachitariu, M., Steinmetz, N.A., Kadir, S.N., Carandini, M., and Harris, K.D. (2016). Fast and accurate spike sorting of high-channel count probes with KiloSort. *Adv. Neural Inf. Process. Syst.* **29**.
59. Rossant, C. (n.d.). Phy: spike sorting software for extracellular electrophysiology. GitHub repository. Available at: https://github.com/cortex-lab/phy.
60. Vincis, R., Chen, K., Czarnecki, L., Chen, J., and Fontanini, A. (2020). Dynamic representation of taste-related decisions in the gustatory insular cortex of mice. *Curr. Biol.* **30**, 1834–1844.e5.

Table S1. Mouse Metadata and Figure Appearances.

Mouse ID	Sex	Mouse Line	Birth Date	Recording Dates	Experiments	Appearances - Main Figures					
						1	2	3	4	5	6
Ephys1	M	C57BL/6J	11/08/22	02/15/23-02/17/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys2	M	C57BL/6J	11/08/22	02/08/23-02/10/23	PrV Si probe, WT	E	A-F	B-D,F		C-E	
Ephys3	M	C57BL/6J	09/20/22	02/01/23-02/03/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys5	M	C57BL/6J	06/22/21	08/19/21-08/24/21	PrV Tungsten						
Ephys6	M	C57BL/6J	06/22/21	09/17/21-09/20/21	PrV Tungsten						
Ephys7	M	C57BL/6J	06/22/21	09/13/21-09/14/21	PrV Tungsten						
Ephys8	F	C57BL/6J	01/25/22	06/15/22	PrV Tungsten						
Ephys9	M	C57BL/6J	08/24/21	10/15/21-10/19/21	PrV Tungsten						
Ephys12	M	C57BL/6J	05/14/22	08/08/22	PrV Tungsten						
Ephys13	F	C57BL/6J	04/12/22	06/29/22-07/06/22	PrV Tungsten						
Ephys14	F	C57BL/6J	04/12/22	07/27/22-08/04/22	PrV Tungsten						
Ephys15	M	C57BL/6J	05/14/22	08/11/22-08/25/22	PrV Tungsten						
Ephys16	M	C57BL/6J	05/14/22	09/01/22-09/06/22	PrV Tungsten						
Ephys17	M	C57BL/6J	05/14/22	09/09/22	PrV Tungsten						
Ephys18	M	C57BL/6J	07/12/22	09/29/22-10/04/22	PrV Tungsten						
Ephys19	M	C57BL/6J	07/12/22	10/17/22	PrV Tungsten						
Ephys20	M	C57BL/6J	07/12/22	10/27/22-11/01/22	PrV Tungsten						
Ephys24	F	C57BL/6J	10/25/22	01/31/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys25	M	C57BL/6J	11/08/22	02/22/23-02/25/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys26	M	C57BL/6J	11/08/22	02/26/23-02/28/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys27	F	C57BL/6J	11/15/22	03/21/23-03/23/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys28	F	C57BL/6J	11/15/22	03/24/23-03/27/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys29	F	C57BL/6J	11/29/22	03/28/23-03/30/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys30	M	C57BL/6J	02/27/23	05/24/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys32	F	C57BL/6J	03/14/23	06/15/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys33	F	C57BL/6J	03/14/23	06/02/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys34	M	C57BL/6J	02/27/23	07/11/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys36	F	C57BL/6J	03/14/23	06/08/23	PrV Si probe, WT	D	D-F	B-D,F		C-E	
Ephys40	M	C57BL/6J	03/28/23	06/26/23	PrV Si probe, WT		A-F	B-D,F		C-E	
Ephys41	M	C57BL/6J	03/28/23	06/27/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys42	F	C57BL/6J	05/23/23	07/25/23	PrV Si probe, WT		D-F	B-D,F		C-E	
beh104	M	C57BL/6J	06/06/23	12/04/23	whisker behavior						
beh101	F	C57BL/6J	09/26/23	12/12/23	whisker behavior						
beh102	M	C57BL/6J	09/26/23	12/13/23	whisker behavior						
beh103	F	C57BL/6J	08/15/23	12/12/23	whisker behavior						
Ephys72	M	vGAT-ires-Cre	11/27/23	03/07/23-03/08/23	PrV optrode local GtACR						
Ephys73	F	vGAT-ires-Cre	11/27/23	03/11/23-03/12/23	PrV optrode local GtACR						
Ephys75	M	C57BL/6J	04/02/24	05/22/24	PrV Si probe, WT + 1WC	C	D-F	B-D,F	B,K	C-E	
Ephys76	F	C57BL/6J	04/02/24	06/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B-F,K	C-E	
Ephys77	F	C57BL/6J	04/22/24	08/14/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys78	F	C57BL/6J	04/02/24	07/02/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys79	M	C57BL/6J	04/02/24	06/24/24	PrV Si probe, WT + 1WC		D-F	B-D,F	A-D,K	C-E	
Ephys81	F	C57BL/6J	04/02/24	07/03/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,G,H,K	C-E	
Ephys83	M	vGAT-ires-Cre	03/09/24	07/09/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys85	M	vGAT-ires-Cre	03/09/24	07/11/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys87	F	C57BL/6J	03/12/24	07/24/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys88	M	C57BL/6J	05/14/24	09/05/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,I,J,K	C-E	
Ephys89	F	C57BL/6J	04/22/24	07/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys92	F	C57BL/6J	04/22/24	08/15/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys94	M	vGAT-ires-Cre	05/22/24	08/17/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys96	F	vGAT-ires-Cre	05/18/24	08/20/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys97	F	vGAT-ires-Cre	03/01/24	11/14/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys100	M	vGAT-ires-Cre	06/20/24	09/08/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys101	M	C57BL/6J	05/14/24	09/09/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys102	F	vGAT-ires-Cre	06/20/24	09/12/24	PrV Si probe, TeLC in SpVi					C-E	

Ephys105	M	C57BL/6J	07/02/24	09/23/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys106	M	C57BL/6J	07/02/24	09/24/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,I,J,K	C-E	
Ephys107	M	C57BL/6J	07/02/24	09/25/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys108	M	C57BL/6J	07/02/24	09/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,E,F,K	C-E	
Ephys114	F	vGAT-ires-Cre	05/22/24	11/18/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys119	M	vGAT-ires-Cre	08/20/24	12/08/24-12/09/24	PrV Si probe, TeLC in SpVi					B-E	
Ephys121	F	vGAT-ires-Cre	08/20/24	12/14/24-12/15/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys122	M	vGAT-ires-Cre	09/09/24	12/16/24-12/17/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys126	M	C57BL/6J	12/03/24	03/20/25-03/21/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys127	F	C57BL/6J	11/03/24	03/24/25-03/25/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys128	F	C57BL/6J	12/31/24	03/26/25-03/27/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	

Mouse ID	Appearances - Supplemental Figures						Other Notes
	S1	S2	S3	S4	S5	S6	
Ephys1			B-E	A-D	B,C	H,I	
Ephys2			A-E	A-E	A-C	H,I	Unit 5 in 20230208_s1 is 1st example in Fig. 2A-C, S3A, S4E, S5A
Ephys3			B-E	A-D	B,C	H,I	
Ephys5		F	J-M				
Ephys6		F	J-M				
Ephys7		F	J-M				
Ephys8		B,F	J-M				
Ephys9		C-F	J-M				
Ephys12		F	J-M				
Ephys13		D-F	J-M				
Ephys14		F	J-M				
Ephys15		F	J-M				
Ephys16		F	J-M				
Ephys17		F	J-M				
Ephys18		F	J-M				
Ephys19		F	J-M				
Ephys20		F	J-M				
Ephys24			B-E	A,C	B,C	H,I	
Ephys25			B-E	A,C	B,C	H,I	
Ephys26			B-E	A-D	B,C	H,I	
Ephys27			B-E	A-D	B,C	H,I	
Ephys28			B-E	A,C	B,C	H,I	
Ephys29			B-E	A,C	B,C	H,I	
Ephys30			B-E	A,C	B,C	H,I	
Ephys32			B-E	A-D	B,C	H,I	
Ephys33			B-E	A-D	B,C	H,I	
Ephys34			B-E	A,C	B,C	H,I	
Ephys36			B-E	A,C	B,C	H,I	
Ephys40			A-E	A,C	B,C	H,I	Unit 8 in 20230626_s1 is 2nd example in Fig. 2A-C, S3A
Ephys41			B-E	A-D	B,C	H,I	
Ephys42			B-E	A,C	B,C	H,I	
beh104	C,F						
beh101	F						
beh102	D-F						
beh103	A,B,F						
Ephys72						D,E	Unit 2 in 20240307_s1 is 1st example in Fig. S6E
Ephys73						C,E	Unit 14 in 20240311_s1 is 2nd example in Fig. S6E
Ephys75			B-E	A-D	B,C	H,I	
Ephys76			B-E	A,C	B,C	H,I	
Ephys77			B-E	A,C	B,C	H,I	
Ephys78			B-E	A,C	B,C	H,I	
Ephys79			B-E	A-D	B,C	H,I	
Ephys81			A-E	A,C	B,C	H,I	Unit 10 in 20240703_s1 is 4th example in Fig. 2A-C
Ephys83			F-I			H,I	
Ephys85			F-I			H,I	
Ephys87			B-E	A,C	B,C	H,I	
Ephys88			A-E	A,C	A-C	H,I	Unit 5 in 20240905_s1 is 3rd example in Fig. 2A-C, S3A; 2nd in S5A
Ephys89			B-E	A,C	B,C	H,I	
Ephys92			B-E	A-D	B,C	H,I	
Ephys94			F-I			H,I	
Ephys96			F-I			H,I	
Ephys97			F-I			H,I	
Ephys100			F-I			H,I	
Ephys101			B-E	A-D	B,C	H,I	
Ephys102			F-I		B,C	H,I	

Ephys105			B-E	A-E	B,C	H,I	
Ephys106			A-E	A-D	B,C	H,I	Unit 7 in 20240924_s1 is 6th example in Fig. 1A-C, S3A
Ephys107			B-E	A,C,E	B,C	H,I	Unit 1 in 20240925_1 is 2nd example in Fig. S4E
Ephys108			B-E	A,C	B,C	H,I	
Ephys114			F-I			H,I	
Ephys119			F-I			G-I	
Ephys121			F-I			H,I	
Ephys122			F-I			H,I	
Ephys126			B-E	A-D	B,C	H,I	
Ephys127			B-E	A-D	B,C	H,I	
Ephys128			B-E	A-D	B,C	H,I	

Table S2. Population Decoder Accuracy

| Distance (mm) | Decoder Accuracy Mean ± SEM (%) | | | | P-Value |
	Proximity	Map	Untuned	Shuffled	Proximity vs Map
7	87.84 ± 1.210	81.14 ± 0.239	32.30 ± 1.767	11.17 ± 0.979	2.06×10^{-4}, *
9	78.87 ± 1.484	74.68 ± 0.330	15.89 ± 0.946	10.85 ± 0.738	3.30×10^{-4}, *
11	64.14 ± 2.238	82.57 ± 0.237	13.41 ± 1.045	11.42 ± 1.016	2.06×10^{-4}, *
13	50.60 ± 3.327	83.90 ± 0.213	13.04 ± 0.567	10.62 ± 1.008	2.06×10^{-4}, *
15	36.29 ± 1.824	82.50 ± 0.148	12.27 ± 0.789	10.78 ± 0.669	2.06×10^{-4}, *
17	30.73 ± 1.372	74.29 ± 0.377	12.01 ± 0.554	10.95 ± 0.690	2.06×10^{-4}, *
19	28.37 ± 1.284	63.16 ± 0.408	11.81 ± 0.384	10.69 ± 0.954	2.06×10^{-4}, *
21	28.91 ± 1.108	49.38 ± 0.439	11.63 ± 0.314	10.54 ± 0.828	2.06×10^{-4}, *
23	34.01 ± 0.670	52.12 ± 0.344	12.18 ± 1.011	11.09 ± 1.111	2.06×10^{-4}, *

*: significant ($p < 0.05$); n.s.: not significant

Table S3. Population Decoder Error

| Distance (mm) | Decoder Accuracy Mean ± SEM (mm) | | | | P-Value |
	Proximity	Map	Untuned	Shuffled	Proximity vs Map
7	0.33 ± 0.034	0.50 ± 0.012	4.90 ± 0.196	8.03 ± 0.137	2.00×10^{-4}, *
9	0.68 ± 0.076	0.51 ± 0.011	5.33 ± 0.122	6.45 ± 0.157	1.30×10^{-3}, *
11	1.34 ± 0.138	0.46 ± 0.011	4.96 ± 0.090	5.30 ± 0.087	2.00×10^{-4}, *
13	2.04 ± 0.107	0.70 ± 0.012	4.50 ± 0.063	4.74 ± 0.077	2.00×10^{-4}, *
15	2.50 ± 0.084	0.78 ± 0.011	4.24 ± 0.052	4.46 ± 0.066	2.00×10^{-4}, *
17	2.50 ± 0.040	1.02 ± 0.012	4.28 ± 0.056	4.70 ± 0.097	2.00×10^{-4}, *
19	2.25 ± 0.039	1.25 ± 0.017	4.81 ± 0.069	5.36 ± 0.122	2.00×10^{-4}, *
21	2.06 ± 0.108	1.20 ± 0.008	5.71 ± 0.081	6.54 ± 0.085	2.00×10^{-4}, *
23	2.32 ± 0.111	1.33 ± 0.012	7.21 ± 0.165	8.03 ± 0.156	2.00×10^{-4}, *

*: significant ($p < 0.05$); n.s.: not significant

Table S4. Single Neuron Decoder Accuracy

Distance (mm)	Decoder Accuracy Mean ± SEM (%)		P-Value
	Proximity	Map	Proximity vs Map
7	65.74 ± 1.319	59.09 ± 1.889	0.02, *
9	40.83 ± 1.662	39.26 ± 2.253	0.73, n.s.
11	31.44 ± 1.447	37.95 ± 2.203	0.02, *
13	24.50 ± 1.312	37.34 ± 2.387	0, *
15	18.75 ± 1.099	34.72 ± 2.277	0, *
17	20.27 ± 1.195	27.63 ± 1.886	0, *
19	20.68 ± 1.143	23.54 ± 1.824	0.15, n.s.
21	20.69 ± 1.216	23.68 ± 1.722	0.05, n.s.
23	26.44 ± 1.505	31.55 ± 2.313	0.03, *

*: significant ($p < 0.05$); n.s.: not significant

Table S5. Single Neuron Decoder Error

| Distance (mm) | Decoder Accuracy Mean ± SEM (mm) | | P-Value |
	Proximity	Map	Proximity vs Map
7	2.11 ± 0.115	2.94 ± 0.198	0, *
9	3.34 ± 0.144	2.78 ± 0.166	0.21, n.s.
11	3.89 ± 0.127	2.89 ± 0.157	0, *
13	4.02 ± 0.101	2.85 ± 0.145	0, *
15	3.76 ± 0.076	3.02 ± 0.146	0, *
17	3.42 ± 0.072	3.32 ± 0.141	0.21, n.s.
19	3.31 ± 0.077	3.45 ± 0.160	0.84, n.s.
21	3.71 ± 0.103	3.76 ± 0.225	0.32, n.s.
23	4.63 ± 0.143	4.58 ± 0.274	0.46, n.s.

*: significant ($p < 0.05$); n.s.: not significant

Table S6. Statistics

Fig.	Comparison	Groups Tested	Measure (Mean ± SEM)	Statistical Test	Test Statistic	P-Value	Significance	DoF
3D	Population decoding accuracy	Proximity vs Map	see Table S2	Wilcoxon rank-sum test	N/A	see Table S2	see Table S2	N/A
3F	Population decoding error	Proximity vs Map	see Table S3	Wilcoxon rank-sum test	N/A	see Table S3	see Table S3	N/A
4B	Firing rates during wall passes	Full array vs. One whisker	12.37 ± 2.986 vs. 8.61 ± 2.487 sp/s	Wilcoxon signed-rank	N/A	1.27×10^{-9}	*	N/A
5D	Preferred distance CDFs	Wild-type vs. TeLC populations	N/A	One-tailed K-S	N/A	8.02×10^{-5}	*	N/A
5E	Proportion of map units	Wild-type vs. TeLC populations	10.34% vs. 3.90%	One-tailed Z-test	Z = 4.64	1.80×10^{-6}	*	N/A
5E	Proportion of proximity units	Wild-type vs. TeLC populations	23.31% vs. 32.09%	One-tailed Z-test	Z = -3.93	4.27×10^{-5}	*	N/A
5E	Proportion of untuned units	Wild-type vs. TeLC populations	48.20% vs. 49.92%	Two-tailed Z-test	Z = -0.68	0.5	n.s.	N/A
5E	Proportion of supressed units	Wild-type vs. TeLC populations	10.43% vs. 7.30%	One-tailed Z-test	Z = 2.12	0.02	*	N/A
5E	Proportion of ambiguous units	Wild-type vs. TeLC populations	7.71% vs. 6.79%	Two-tailed Z-test	Z = 0.69	0.49	n.s.	N/A
S1F, Left	C1 whisker midpoint	Awake vs. Anesthetized	-8.52 ± 0.633° vs. -5.88 ± 0.910°	Two-tailed paired t-test	t = -3.86	0.03	n.s. (α = 0.0167)	3
S1F, Middle	C2 whisker midpoint	Awake vs. Anesthetized	-6.81 ± 0.867° vs. -4.72 ± 0.927°	Two-tailed paired t-test	t = -3.57	0.04	n.s. (α = 0.0167)	3
S1F, Right	Greek whisker midpoint	Awake vs. Anesthetized	-4.76 ± 1.041° vs. -2.63 ± 0.714°	Two-tailed paired t-test	t = -4.55	0.0199	n.s. (α = 0.0167)	3
S4B, Left	Firing rates of wall-activated, distance-tuned units	Wall passes vs. Baseline	16.44 ± 1.640 vs. 6.36 ± 0.704 sp/s	One-tailed K-S	N/A	9.26×10^{-28}	*	N/A
S4B, Right	Firing rates of wall-suppressed, distance-tuned units	Wall passes vs. Baseline	4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s	One-tailed K-S	N/A	3.06×10^{-4}	*	N/A
S4C	Untuned unit firing rates	Wall passes vs. Baseline	6.65 ± 0.687 vs. 6.90 ± 0.682 sp/s	Two-tailed K-S	N/A	0.74	n.s.	N/A
S4G, Middle	Wall-tuning category (ambiguous unit)	Non-closest vs. Closest distance	32.59 ± 4.253 vs. 26.53 ± 4.107 sp/s	One-tailed Wilcoxon signed-rank test	N/A	0.09	n.s.	N/A
S4G, Right	Wall-tuning category (map unit)	Non-closest vs. Closest distance	51.04 ± 2.49 vs. 26.09 ± 1.033 sp/s	One-tailed Wilcoxon signed-rank test	N/A	1.27×10^{-4}	*	N/A
S5B	Single-neuron decoder accuracy	Proximity vs Map	see Table S4	Wilcoxon rank-sum test	N/A	see Table S4	see Table S4	N/A
S5C	Single-neuron decoder error	Proximity vs Map	see Table S5	Wilcoxon rank-sum test	N/A	see Table S5	see Table S5	N/A
S6H, Left	Baseline firing rates (untuned units)	TeLC vs. Wild-type populations	9.34 ± 1.330 vs. 6.90 ± 0.682 sp/s	One-tailed K-S	N/A	7.63×10^{-5}	* (α = 0.0125)	N/A
S6H, Right	Wall-pass firing rates (untuned units)	TeLC vs. Wild-type populations	9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s	One-tailed K-S	N/A	1.06×10^{-4}	* (α = 0.0125)	N/A
S6I, Left	Baseline firing rates (tuned units)	TeLC vs. Wild-type populations	5.60 ± 0.845 vs. 5.46 ± 0.670 sp/s	One-tailed K-S	N/A	0.14	n.s. (α = 0.0125)	N/A
S6I, Right	Wall-pass firing rates (tuned units)	TeLC vs. Wild-type populations	20.25 ± 2.263 vs. 16.44 ± 1.640 sp/s	One-tailed K-S	N/A	1.64×10^{-3}	* (α = 0.0125)	N/A
N/A	Firing rates during baseline	Fullarray vs. One whisker	7.08 ± 1.424 vs. 7.08 ± 1.418 sp/s	Wilcoxon signed-rank	N/A	0.59	n.s.	N/A