Peri-Head Distance Coding in the Mouse Brainstem

Wenxi Xiao[1]*, Kyle S. Severson[1]*, Hao Zheng[1], Ke Chen[1], Paul M. Thompson[1], Manuel S.

Levy[1], Seonmi Choi[1], Shengli Zhao[2], Jun Takatoh[1], Vincent Prevosto[1&], and Fan Wang[1&]

Affiliations

1. Department of Brain and Cognitive Sciences, McGovern Institute for Brain Research,

Massachusetts Institute of Technology, Cambridge, MA, USA.

2. Department of Neurobiology, Duke University, Durham, NC, USA.

* These two authors contributed equally

& Address correspondence to prevosto@mit.edu and fan_wang@mit.edu

SUMMARY

Perceiving object distance in peri-personal space (PPS) is essential for guiding movement and

avoiding danger, yet how early somatosensory circuits extract distance information from tactile

inputs remains unclear. Here, we investigate how second-order neurons in the mouse whisker

brainstem encode peri-head distance. Using *in vivo* extracellular recordings in awake mice in a

naturalistic wall-passing paradigm, we find brainstem neurons employ two distance-coding

schemes: a "proximity" code, where firing increases monotonically as objects approach the face;

and a "map" code, where neurons exhibit peak tuning at specific distances to collectively tile peri-

head space. The map code outperforms proximity code in population decoding of distance.

Perturbation experiments revealed multi-whisker integration and intersubnuclear inhibition

contribute to the generation of map-like tuning. These findings highlight the previously

underappreciated brainstem circuit for sensory code transformation, suggesting inhibition acts as

a neural comparator, subtracting peripheral inputs with differing sensitivity to transform proximity-

based sensory inputs into a map-like representation of PPS.

MAIN TEXT

The ability to perceive and represent space is fundamental for animals to interact with the external

world. Estimating the distance of a predator, planning reach-to-grasp movements for food, or

navigating complex terrains with obstacles and gaps all require accurate spatial processing. Of

particular ethological importance is the space immediately surrounding the body, peripersonal

space (PPS), which serves as a critical interface for sensorimotor integration, enabling object

interaction, social engagement, and defensive behaviors (Serino, 2019). Classic primate studies

identified a multisensory neural substrate for PPS in the ventral premotor cortex (PMv), where

bimodal neurons respond both to tactile stimulation on specific body parts and to visual or auditory

stimuli presented *near* those same body surfaces (Rizzolatti et al., 1981; Gentilucci et al., 1983;

Rizzolatti & Gentilucci, 1988).

PPS has been proposed as a contact-related action field (Serino et al., 2009; Bufacchi &

Iannetti, 2018), where accurate representation of relative distance to nearby entities directly

informs motor planning. For example, reaching for a cup beside the hand versus one placed

farther away requires distinct patterns of arm and trunk muscle activation; likewise, crossing a

narrow versus wide gap requires different physical demands. Some bimodal neurons in PMv

increase their activity as visual stimuli approach the body surface (Rizzolatti et al., 1981; Rizzolatti

& Gentilucci, 1988), consistent with a proximity-based rate code. However, whether this proximity

code is the sole encoding scheme for representing distances within PPS remains unclear.

Clarifying how distance is represented in PPS is therefore crucial for uncovering the sensory

computations that support spatial awareness and action execution.

To estimate distances in the surrounding environment, humans often rely on physical

references—most commonly, the body itself—for example, describing something as being "an

arm's length away". This body-referenced approach to spatial perception dates back to ancient

civilizations, which formalized units of measurement based on body parts, including the "palm",

"foot", "step", and "fathom" (the span of outstretched arms). Compared to continuous proximity

cues, body-part references provide discrete spatial scales that can be more readily interpreted

and used for motor planning. This intuitive measurement strategy raises the question of whether

the brain implements an analogous "internal ruler" that systematically encodes distances relative

to specific body parts. If so, what is the representational format of this distance code? Addressing

these questions may provide important insights into how spatial representations within PPS are

constructed and used by the brain to guide behavior.

The rodent whisker sensory system provides a natural framework to study distance

encoding, with grid-like whiskers serving as a built-in coordinate system for peri-head space.

Rodents are nocturnal animals that rely primarily on their whiskers to explore their surroundings

in the dark (Diamond et al., 2008; Kleinfeld & Deschênes, 2011; Mitchinson & Prescott, 2013).

The whisker array is organized into arcs of systematically varying lengths, with longer whiskers

located posterior and shorter whiskers anterior (Belli et al., 2017; Severson & Xu et al., 2017;

Bresee & Belli et al., 2023). This arrangement forms an anatomical gradient in which whisker

length and position correspond systematically to specific regions of peri-head space (Figure 1A).

As a result, each whisker serves as a spatial reference point, allowing the brain to integrate inputs

across the array into a head-centric representation of peri-head space. Supporting this idea, a

recent study found that during virtual corridor navigation a subset of neurons in the whisker barrel

cortex (wS1) exhibited non-monotonic tuning to the distance between the mouse's head and

surrounding walls (Sofroniew & Vlasov et al., 2015). Non-monotonic tuning can be interpreted as

a "map code", where the peak of each tuning curve maps a specific location in peri-head space

onto a neural axis. At the population level, these preferred distances spanned the full range of

whisker reach, effectively forming a head-centered representation of peri-head distance.

However, the map code for peri-head distance has not been observed in the periphery.

Whiskers are innervated by mechanosensory neurons with soma residing in the trigeminal

ganglion (TG), each of which has a single-whisker receptive field (Szwed et al., 2006; Khatri et

al., 2009; Bale & Maravall, 2018). These neurons respond to mechanical stresses on the whisker

follicle generated when the whisker bends upon contact with an object (Mitchinson et al., 2004;

Bush et al., 2016; Campagner et al., 2016; Severson & Xu et al., 2017). Consequently,

extracellular recordings from TG neurons, using a vertical pole positioned at various locations

along the length of a single whisker, revealed that firing rates increase as the point of contact

moves closer to the face (Szwed et al., 2006). Thus, in the periphery, TG neurons encode radial

distance via a proximity-based rate code, referenced to the base of a single whisker (Ahissar &

Knutsen, 2008). This implies that a sensory code transformation occurs along the whisker sensory

pathway to convert whisker-centered proximity code in the periphery into the map-like peri-head

distance code observed in wS1.

We hypothesize that a transformation from proximity to map-like representation of peri-

head distance occurs at the level of second-order neurons in the principal trigeminal nucleus (PrV)

of the brainstem. PrV contains neurons with either single- or multi-whisker receptive fields

(Shipley, 1974; Jacquin et al., 1989; Veinante & Deschênes, 1999; Minnery et al., 2003; Minnery

& Simons, 2003; Timofeeva et al., 2004, 2005) and receives convergent inputs from multiple

subtypes of TG mechanosensory neurons, including both rapidly and slowly adapting afferents

(Sakurai et al., 2013; Jacquin et al., 2015). In addition, PrV circuits incorporate local glycinergic

inhibitory interneurons (Avendaño et al., 2005; Bellavance et al., 2010) and receive

topographically organized long-range GABAergic input from the spinal trigeminal nucleus

interpolaris (SpVi; Jacquin et al., 2015, Furuta et al., 2008). These circuit motifs, supporting both

multi-whisker integration (Minnery & Simons, 2003; Furuta et al., 2008) and feedforward inhibition,

make PrV a candidate site for implementing the transformation of whisker-centered proximity

signals into a head-centered map code. Inspired by Porter and Groh (2006), we propose that a

simple circuit (Figure 1B) could transform monotonic proximity-based inputs from TG neurons into

a non-monotonic map-like readout of peri-head distance. Importantly, this circuit requires inhibition: when proximity-based inputs from two TG neurons with different mechanical sensitivity are subtracted via inhibition, a peak response at a specific peri-head distance emerges. To test these hypotheses, we performed *in vivo* electrophysiological recordings from PrV neurons in a wall-passing paradigm.

Results

A wall-passing paradigm to study peri-head distance encoding in the whisker brainstem

To evoke tactile sensation similar to that experienced during naturalistic wall-following behavior, we devised a wall-passing paradigm for awake, behaving mice head-fixed on a linear treadmill (Severson & Xu, 2017; Figure 1C, right; STAR Methods). In this setup, each stimulus presentation consists of a wall moving past the mouse's head at a fixed radial distance from anterior to posterior, delivering controlled tactile stimulation to the whisker array. A linear actuator moved the wall along the anteroposterior axis at a constant velocity of 10 mm/s, operating independently of the animal's locomotion (open loop). Before each wall pass, the radial distance between the wall and the mouse's face (wall distance) was randomly set. Each pass lasted approximately 14 seconds, with wall movement onset at t = 0 and whisker-wall contacts typically occurring between 3-11 seconds. In total, each distance was presented for 15-20 wall stimulus presentations. The wall position was aligned to the midline of the head, with wall distances measured from the wall to the C2 whisker follicle (Figure 1C, left).

After habituation to the wall-passing stimulus, mice exhibited predominantly passive whisker-wall interactions. To assess the degree of active retraction, we compared wall-induced whisker bending angles of a single row of whiskers in awake versus anesthetized conditions (see Supplemental Figure S1). Under anesthesia, the moving wall pushed whiskers in the retraction

direction, producing larger absolute bending angles at closer wall distances (Supplemental Figure

S1D, right). Awake, behaving mice in the same wall-passing paradigm showed similar whisker

bending angles (mean difference: 2.28°), with no statistically significant difference from the

anesthetized condition across whiskers (Supplemental Figure S1F). This observation suggests

that mice engaged in little active whisker retraction to avoid contact, instead allowing the whiskers

to passively glide along the passing wall. Given this minimal wall-induced active retraction,

whisker bending mostly increased monotonically with wall proximity, supporting proximity-based

distance encoding in the periphery.

PrV contains both a proximity and a map code of peri-head distance

To examine how PrV neurons respond to the passing wall stimulus at varying distances, we first

performed *in vivo* extracellular recordings in mice with only the C-row whiskers intact, using single

tungsten electrodes (see Supplemental Figure S2A-C; STAR Methods). Based on our single-unit

isolation metrics (see STAR Methods; Supplemental Figure S3K-M), we isolated 46 units from 14

mice (see Table S1 for mouse metadata), including 18 units with single-whisker and 28 units with

multi-whisker receptive fields. We analyzed single-unit responses using peri-event time

histograms (PETHs) by computing the averaged firing rates for wall contact (3-11 s) and baseline

periods (-4.5-0 s) across trials. To determine wall responsiveness, we applied one-tailed Wilcoxon

signed-rank tests (see STAR Methods). Units with significantly higher firing rates at any tested

distance relative to baseline were classified as "wall-activated," while those with significantly lower

firing were classified as "wall-suppressed." Using these criteria, 93.5% (43/46) of units recorded

with single tungsten electrodes were identified as wall-responsive (presumably touch-sensitive).

To assess distance tuning, we performed two-tailed Wilcoxon signed-rank tests across all

distance pairs (p < 0.05, corrected using the Benjamini-Hochberg false discovery rate; Storey,

2002). Units with significant differences in mean firing rates for at least one distance pair were

classified as "distance-tuned" (84.8%, 39/46). For each distance-tuned unit, we computed a z-scored tuning curve (see STAR Methods). The majority of units (52.2%, 24/46) showed a broadly monotonic, inversely proportional relationship with wall distance (e.g., Supplemental Figure S2D), consistent with PrV neurons inheriting proximity-based TG inputs. Interestingly, a distinct subset of units (13.0%, 6/46) clearly preferred intermediate or far wall distances (e.g., Supplemental Figure S2E), revealing a heterogeneous representation of peri-head distance in PrV (see Supplemental Figure S2F).

Because high-impedance tungsten electrodes preferentially capture high-amplitude spikes, unit isolation was biased toward larger neurons (Henze et al., 2000; Buzsáki et al., 2012), raising the possibility that smaller PrV neurons, that were potentially tuned to intermediate wall distances, were under-sampled. Furthermore, single electrode recordings were low-yield, resulting in small sample size (n=46 units). To overcome these limitations, we adopted 32-channel single-shank silicon probes for acute recordings of PrV single-unit activity during the wall-passing paradigm (Figure 1C; See STAR Methods). We isolated 1,141 single units across 36 mice (e.g., Figure 1D; Supplemental Figure S3A-E). Of the 1,141 units, 67.3% (768/1141) responded to the passing-wall stimulus, either being activated (532/768, 69.3%) or suppressed (333/768, 43.3%). Given that our probe spanned the entire dorsoventral axis of PrV (Figure 1D) and that whisker-innervating neurons are specifically localized to ventral PrV (Bechara et al., 2015), the prominence of unresponsive units, likely corresponding to microvibrissa-, jaw-, or lip-related neurons in dorsal PrV, is not unexpected.

Representative spike rasters, PETHs, and z-scored wall-distance tuning curves from six wall-responsive units are shown in Figure 2A-C. These examples illustrate the diversity of tuning responses: Unit 1 showed monotonic increases in firing with decreasing wall distance ("proximity" tuning); Units 2-4 exhibited peak firing at specific intermediate distances ("map" tuning); Unit 6 was suppressed during wall contact; and Unit 5 displayed a complex response profile with both suppression and activation depending on wall distance. In addition to sustained responses, wall-

touch onset and offset responses were often observed in the same PrV neurons, suggesting

convergence of multiple afferent types (i.e., slowly and rapidly adapting) onto PrV neurons,

consistent with anatomical evidence that different types of TG afferent terminals are intermingled

within PrV barrelettes (Sakurai et al., 2013). However, our analyses focused on mean firing rates

across the entire wall-contact period (8s) and did not further dissect these transient dynamics.

We systematically identified a total of 656 units out of 1,141 (57.5%) that were tuned to

wall distance. Among these distance-tuned neurons, 72.0% (472/656) were significantly activated

by at least one wall distance relative to baseline, while 32.5% (213/656) were significantly

suppressed by at least one wall distance relative to baseline. Interestingly, wall-activated

distance-tuned units exhibited lower baseline firing rates than untuned units (5.46 ± 0.670 vs.

6.90 ± 0.682 sp/s; two-tailed K-S test, p = 1.91 × 10^{-10}), while wall-suppressed and distance-

tuned units exhibited baseline firing rates comparable to untuned units (7.31 ± 1.124 vs. 6.90 ±

0.682 sp/s; two-tailed K-S test, p = 0.27; Supplemental Figure S4A). During wall contact, "wall-

activated" and "distance-tuned" units showed significant increase in firing relative to baseline

(16.44 ± 1.640 vs. 6.36 ± 0.704 sp/s; one-tailed K-S test, p = 9.26 × 10^{-28}; Supplemental Figure

S4B, left), "wall-suppressed" and "distance-tuned" units showed significant decrease in firing

relative to baseline (4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s; one-tailed K-S test, p = 3.06 × 10^{-4};

Supplemental Figure S4A, right), whereas untuned units showed no difference (6.65 ± 0.687 vs.

6.90 ± 0.682 sp/s; two-tailed K-S test, p = 0.74; Supplemental Figure S4C), validating our

classification criteria.

To visualize tuning at the population-level, we concatenated normalized tuning curves for

"wall-activated" and "wall-suppressed" units into heatmaps, sorted by each unit's preferred wall

distance (Figure 2D, S3D). While many units exhibited tuning to the closest wall position (7 mm),

others showed tuning to intermediate or far distances (Figure 2E, S3F). To further characterize

wall-activated distance-tuned units, we compared mean firing rates at each non-closest

distance (≥ 9 mm) to the closest distance (7 mm; one-tailed Wilcoxon signed-rank tests, see

STAR Methods). Units with significantly higher firing at any non-closest distance were

categorized as "map" units (Supplemental Figure S4G, right), while units with peak firing at the

closest distance were classified as "proximity" units (Supplemental Figure S4G, left). This

analysis identified 266 proximity units (23.3%) and 118 map units (10.3%) out of the 1,141 total

recorded units (Figure 2F). An additional 88 units (7.7%) were tuned but did not meet criteria for

either map or proximity and were labeled "ambiguous" (Supplemental Figure S4G, middle) and

excluded from further analysis. Suppressed-only units comprised a small portion of the population

(119/1141; 10.4%, Figure 2F).

We note that classification as "proximity" versus "map" is based on relative peak firing

rates within the tested range (7-23 mm). It remains possible that some proximity units would

exhibit decreased firing at even shorter distances, thus revealing map-like tuning beyond the

tested range. Therefore, our categorization serves as an operational definition to capture distinct

response properties within the presented distance range, rather than strict functional classes.

To assess the reliability of proximity and map codes across repetitions, we performed 50%

cross-validation by sorting units using half the wall passes and applying the order to the remaining

half. The resulting distance-tuning heatmaps showed little difference in preferred distance

between cross-validated partitions (Supplemental Figure S4E). Together, these results reveal that

PrV robustly encodes peri-head distance using both a proximity code—characterized by

increasing firing at close distances—and a distributed map code. Though the map coding is more

prominent in wS1, its presence in PrV points to an underappreciated role for the whisker brainstem

in active sensory processing, beyond simple relay.

The map code in PrV is more accurate at decoding peri-head distance

We next asked which encoding scheme, proximity or map, allows more accurate readout of peri-

head distance from a given neural population. To address this, we trained maximum correlation

coefficient decoders (Meyers, 2013; Vincis & Chen et al., 2020) on either the proximity or map

subpopulation activity (see STAR Methods), with untuned and shuffled subpopulations used as

controls.

As expected, decoding accuracy during the wall-contact period was higher in both map

and proximity populations compared to controls. Notably, the map population (n = 118 units)

outperformed the proximity population (subsampled n = 118 of 266 units) in overall decoding

accuracy over the entire wall passing period averaged across all wall distances (Figure 3A).

Untuned units still performed slightly above chance due to residual distance tuning that fell below

our stringent statistical threshold for wall-tuning classification.

Confusion matrices revealed that map units were particularly more accurate at decoding

intermediate and far distances (Figure 3B, bottom), whereas proximity units showed stronger

decoding performance only at the closest distance (Figure 3B, top). To quantify this, we compared

classification accuracy across nine wall distances for each population (Figure 3C; Table S2). At 7

and 9 mm, proximity units were significantly more accurate than map units. In contrast, map units

were significantly more accurate at decoding more distal wall positions (11-23 mm). To assess

decoding precision in millimeters, we computed a distance-classification error (in mm) by

multiplying the confusion matrix with a distance error matrix (Figure 3D; Table S3). While

classification error increased with distances, both populations showed submillimeter error in

decoding the closest distances (7 and 9 mm). However, only the map population showed

submillimeter error in decoding intermediate distances (from 11 to 15 mm, Figure 3E). These

results are consistent with the performance of single-neuron decoders trained on map and

proximity units and averaged across those populations (see STAR Methods, Supplemental Figure

S5; Table S4; Table S5), although their overall decoding accuracy is lower than population

decoders.

Together, these results suggest that proximity and map codes provide complementary

encoding schemes to read out peri-head distance: the proximity code supports highly accurate

decoding at close distances, while the map code enables finer discrimination across intermediate

and far distances. Thus, optimal distance readout by downstream circuits would integrate

information across both proximity and map subpopulations.

Multi-whisker integration contributes to the generation of PrV map code

In the wall-passing paradigm, closer wall distances induced greater whisker bending, driving

stronger activation of whisker TG afferents. TG therefore transmits a monotonic, proximity-based

input to PrV. The presence of PrV neurons tuned to specific wall distances (i.e., map cells)

suggests a transformation of the peripheral proximity code into a proximity-map representation of

peri-head distance. Because the passing wall stimulates multiple whiskers of different lengths, we

hypothesized that PrV could generate the map code by integrating multi-whisker inputs. To test

this hypothesis, we recorded PrV single-unit activity during the wall-passing paradigm under two

conditions within the same recording session: first, with the mouse's full whisker array intact, and

second, after acutely cutting all but a single whisker, thus eliminating multi-whisker integration

(see STAR Methods; Figure 4A). Continuous recording of neural activity allowed us to track each

neuron's distance tuning profile before and after the acute whisker cutting manipulation.

Across the recorded population (n = 195 units from 18 mice), acute whisker cutting did not

significantly alter baseline firing rates (measured before wall-movement onset for each wall pass;

Wilcoxon signed-rank test, full array: 7.09 ± 1.034 vs. single whisker: 7.10 ± 1.038 sp/s, p = 0.70).

In contrast, firing rates for wall-activated and distance-tuned units (n = 86) during wall-passing

periods were significantly reduced following cutting down to a single whisker (full array: 11.58 ±

1.338 vs. single whisker: 8.18 ± 2.602 sp/s, p = 4.43 × 10^{-8}, Figure 4B). Of the 86 wall-activated

and distance-tuned units, 66 showed significant changes in firing rates during wall-passing

periods between cutting conditions (two-sided Wilcoxon signed-rank test, $p < 0.05$). Among these,

52 units showed a significant decrease in firing rates, consistent with loss of excitatory input from

whiskers within their receptive fields, while 14 units displayed increased firing rates, likely

reflecting disinhibition due to removal of surround-whisker inputs (Furuta et al., 2008; Furuta et

al., 2010). Note that due to the use of multi-channel recordings in mice with intact whisker arrays,

it was difficult to precisely identify the receptive field of each recorded PrV unit. As a result, in

many cases the principal whisker driving a given neuron was inadvertently cut, leading to a near-

complete loss of response.

Before whisker cutting, 21 units were classified as map cells, and only 10 of these

remained tuned to wall distance after cutting down to a single whisker. After cutting, we observed

shifts in preferred wall distance: two map cells shifted their preferred distance from an

intermediate to the closest wall distance (Figure 4C,D, summarized in the population in Figure

4K). The whisker cutting manipulation effectively converted these map cells to proximity cells,

indicating that in these cases multi-whisker integration was essential for generating the non-

monotonic map code for peri-head distance. On the other hand, four map cells shifted their

preferred distance closer but did not fully convert to proximity cells (e.g., Figure 4E,F). Two map

cells retained their preferred distance after whisker cutting but showed changes in response

amplitude (Figure 4G,H). These results showed that map cells could be generated with inputs

from just a single whisker although multi-whisker inputs still modulate the gain or the preferred

distance tuning in map cells. Together, these findings demonstrated that multi-whisker integration

contributes to, but is not required for, transforming proximity-based inputs into a non-monotonic

map code in PrV.

We also examined how proximity cells were affected by the loss of multi-whisker input. 45

units were classified as proximity cells before whisker cutting, and only 15 of these remained

tuned to wall distance after cutting down to a single whisker. In remaining proximity cells, reducing

input to a single whisker changed the gain of their responses: nine units showed decreased

activity (e.g., Figure 4I,J, Top), while four exhibited increased activity (e.g., Figure 4I,J, Bottom).

Inhibition shapes the PrV map code

In addition to integrating inputs from different whiskers, the sensory code transformation occurring

in PrV may also involve the subtraction of TG inputs from the same whisker but differ in their

distance sensitivity profiles. Prior research indicates that TG touch neurons exhibit heterogeneity

in mechanosensory thresholds as well as in their adaptation properties. (Hao & Delmas, 2010;

Abraira & Ginty, 2013; Furuta et al., 2020). This suggests that TG touch neurons innervating the

same whisker may exhibit different proximity-tuning profiles. Whether involving single- or multi-

whisker integration, a common feature of the hypothesized transformation model (Figure 1B) is

the engagement of inhibitory mechanisms. Next, we aim to directly test this by combining *in vivo*

electrophysiology with optogenetic or chemogenetic silencing approaches.

The intrinsic circuitry of PrV is shaped by two major sources of inhibition: local glycinergic

and GABAergic interneurons within PrV itself, as well as profuse long-range GABAergic

projections from spinal trigeminal nucleus interpolaris (SpVi; Furuta et al., 2008; Bellavance et al.,

2010). To assess the role of local inhibitory neurons in encoding of peri-head distance within PrV,

we unilaterally injected AAV-DIO-GtACR (Govorunova et al., 2015; Takatoh & Prevosto et al.,

2021) into PrV of vGat-Cre mice (N=2), enabling Cre-dependent expression of the inhibitory opsin

GtACR in PrV inhibitory neurons (Supplemental Figure S6A). We then performed acute

optogenetic silencing of these local inhibitory neurons while recording single-unit activity in PrV

during the wall-passing paradigm (Supplemental Figure S6B,D). For each wall distance, the

"Laser OFF" and "Laser ON" conditions were randomly interleaved. Each distance was repeated

15-20 times in each condition to allow the comparison of distance tuning profiles with and without

local inhibition.

Among the 112 well-isolated units, 18 were putative inhibitory neurons expressing GtACR,

as they were strongly inhibited during the "Laser ON" period (modulation index ≤ -0.5; see STAR

Methods; Supplemental Figure S6C,E). Of distance-tuned units (71/112), 17 were opto-inhibited,

exhibiting a decrease in firing rates during laser stimulation (modulation index < 0; see STAR

Methods), while preserving their distance tuning profile (i.e., proximity or map, data not shown).

Sixteen units showed disinhibition under the "Laser ON" condition (modulation index > 0; see

STAR Methods). Nonetheless, these opto-disinhibited neurons, regardless of whether they were

proximity or map cells, retained their preferred distance (e.g., Supplemental Figure S6F,G). These

observations suggest that PrV local inhibitory neurons primarily modulate the gain of individual

PrV neurons' firing rates, rather than altering their preferred distance.

We next investigated the role of long-range intersubnuclear inhibitory inputs from SpVi to

PrV. Due to the extensive size of SpVi, optogenetic silencing of its inhibitory neurons using a

single optical fiber is impractical. Instead, we employed a AAV strategy to achieve sustained

silencing by expressing tetanus toxin light chain (TeLC; Link et al., 1992; Schiavo et al., 1992;

Zhang et al., 2015; Takatoh & Prevosto et al., 2021) to permanently block synaptic transmission.

AAV-DIO-TeLC-EYFP was unilaterally injected into SpVi of vGat-Cre mice (N=11) to target

inhibitory neurons known to send long-range projections to PrV (Figure 5A; Furuta et al., 2008).

Post-hoc histology confirmed TeLC-EYFP expression in SpVi (Supplemental Figure S6H), as well

as labeled axon terminals and boutons in PrV originating from SpVi inhibitory neurons (Figure

5B). Note that the large size of SpVi limited viral access to all PrV-projecting neurons, thereby

constraining the extent of silencing.

We recorded single-unit activity in PrV 10-15 days after TeLC injection during the wall-

passing paradigm and isolated 587 units. To assess the impact of suppressing long-range

inhibitory inputs from SpVi, we compared wall-distance tuning in SpVi-TeLC mice to that in wild-

type mice (data previously presented in Figure 2). Side-by-side comparison of normalized tuning

curves of all "wall-activated," "distance-tuned" units revealed a clear reduction in the proportion of

map units in SpVi-TeLC mice (Figure 5C), shifting the population to prefer closer wall distances

(Figure 5D). Tuning profile classification confirmed a significant shift in the distribution of tuning

types (Figure 5E). Most importantly, the proportion of map units was reduced by more than half

in SpVi-TeLC mice compared to wild-type mice (3.9% vs 10.34%; Z-statistic: 4.64, p = 1.8×10^{-6}),

while the proportion of proximity cells increased (32.09% vs 23.31%; Z-statistic: -3.93, p =

4.27×10^{-5}). Our silencing manipulation did not fully abolish map cells, possibly due to incomplete

silencing of all long-range inhibitory intersubnuclear projections. The proportion of suppressed

units decreased modestly (7.3% vs 10.43%; Z-statistic: 2.12, p = 0.02), whereas the proportions

of untuned (49.9% vs 48.2%; Z-statistic: -0.68, p = 0.50) and ambiguous units (6.79% vs 7.71%;

Z-statistic: 0.69, p = 0.49) remained unchanged.

During both baseline and wall-passing periods, untuned units in SpVi-TeLC mice exhibited

significantly higher firing rates than those in the wild-type population, consistent with disinhibition

following loss of inhibitory inputs from SpVi (baseline: 9.34 ± 1.330 vs 6.90 ± 0.682 sp/s, p = 7.63

× 10^{-5}; wall passes: 9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s, p =1.06 × 10^{-4}; one-tailed K-S tests;

Supplemental Figure S6I). In contrast, distance-tuned units in SpVi-TeLC mice showed

comparable baseline activity to those in wild-type mice (5.60 ± 0.845 vs. 5.46 ± 0.670 sp/s; one-

tailed K-S test, p = 0.14), but significantly elevated firing rates during wall contact (20.25 ± 2.263

vs. 16.44 ± 1.640 sp/s; one-tailed K-S test, p = 1.64 × 10^{-3} Supplemental Figure S6J). These

findings suggest that TeLC-mediated silencing of SpVi inputs altered the tuning preferences of

PrV neurons (i.e., shifting them toward closer wall distances) without affecting their spontaneous

activity level. Thus, long-range inhibition from SpVi plays a key role in transforming feedforward

proximity-based inputs from TG into a map-like peri-head distance representation in PrV. More

broadly, our findings extend the role of SpVi-PrV inhibitory projections beyond simple sensory

gating (Furuta et al., 2008; Mohar et al., 2013), implicating them in a sensory code transformation.

Finally, to demonstrate how this inhibitory circuit enables the proximity-to-map transformation, we

implemented a simple model constrained by known anatomical motifs (Figure 6A; see STAR

Methods). In our model, TG afferents send proximity-based inputs to neurons in PrV via two

distinct pathways: a direct excitatory (lemniscal) pathway and an indirect inhibitory pathway

mediated by SpVi inhibitory neurons. TG proximity-based inputs were modeled as sigmoidal

functions with thresholds skewed toward proximal distances (Figure 6B), and PrV neurons

received sparse, random combinations of excitatory inputs from TG neurons and inhibitory inputs

from SpVi (see STAR Methods). The model transformed monotonic inputs into a heterogeneous

population of PrV tuning curves, including both proximity- and map-like responses to distance

(Figure 6C). Across the modeled population (n = 1,225), PrV neurons exhibited a wide range of

preferred distances, reproducing qualitative features of the empirical tuning curves (Figure 6D,E,

left). Reducing inhibitory input per PrV neuron by 50% led to noticeable loss of map units and

expansion of proximity units (Figure 6D-F). These simulations show that simple feedforward

motifs can transform monotonic sensory input into a non-monotonic map-like code, enabling

extraction of an abstract representation of peri-head distance, and that inhibition is necessary for

generating intermediate and far distance tuning in PrV. Together, these results highlight

complementary roles for local and long-range inhibition in the brainstem trigeminal complex: local

inhibitory circuits primarily regulate sensory gain, whereas long-range SpVi inputs reshape tuning

to generate the map code for peri-head distance.

DISCUSSION

Here, we revealed a previously unrecognized sensory code transformation in the whisker

brainstem: from proximity-based whisker sensory inputs to map-like peri-head distance

representation. Using *in vivo* electrophysiological recordings from awake, behaving mice in a

naturalistic wall-passing paradigm, we discovered two encoding schemes for distance in PrV: a

proximity code in which neuronal firing rates increase monotonically as the distance becomes

closer, and a map code in which neurons exhibit peak firing at specific distances. Both encoding

schemes are robust and stable, invariant to the multiplexed and highly variable whisker inputs

across trials. Mechanistically, we provide evidence that both multi-whisker integration and long-

range inhibitory inputs from SpVi play critical roles in generating the non-monotonic map code.

While most prior research on whisker-based object localization has examined TG afferents

(Ahissar & Knutsen, 2008) and cortical areas (Barnéoud et al., 1991; O'Connor et al., 2010;

Sofroniew & Vlasov et al., 2015; Heimburg, Mari Saluti & Timm et al., 2025) in awake, behaving

rodents, the brainstem trigeminal complex has remained largely unexplored. To our knowledge,

only two prior studies have reported *in vivo* extracellular single-unit activity in PrV (Moore et al.,

2015; Chakrabarti & Schwarz, 2018). Our findings thus fill a critical gap in understanding early

stages of tactile processing, revealing an anatomically-defined brainstem circuit that integrates

and transforms peripheral signals to construct representations of peri-head space. More broadly,

this work highlights the role of PrV as a processing center that converts afferent inputs into

meaningful representations for downstream computation.

Map vs. Proximity code of peri-head distance

Our discovery of PrV map cells challenges a long-held assumption of PrV as a simple relay of

trigeminal input. The map code, in which neurons fire maximally at a preferred distance and

decrease their response as the stimulus deviates from that value, mirrors a common encoding

motif observed across sensory and motor systems (Hubel & Wiesel, 1962; O'Keefe & Dostrovsky,

1971; Georgopoulos et al., 1986). Such "peaked" tuning could be computationally advantageous,

as it allows distance information to be extracted from a relatively small subset of neurons (Pouget

et al., 2000; Jazayeri & Movshon, 2006), resulting in a sparse, metabolically efficient

representation capable of attaining high precision (Laughlin, 2001; Olshausen & Field, 2004). The

presence of a map code in the brainstem allows feature extraction from multiplexed tactile signals

during active sensation (e.g., contact forces, stick-slips, vibrations), reducing computational

burden for downstream structures and enabling rapid extraction of behaviorally relevant features.

Functionally, the map and proximity codes could be specialized for different operating

regimes of PPS. Our decoder analysis showed that map cells were significantly better at decoding

intermediate and far distances, suggesting that they are optimized for fine-grained spatial

localization over a broader distance range within PPS. This fine-grained distance representation

is likely critical for tasks in which rodents use their whiskers to judge whether gaps are crossable

(Hutson & Masterton, 1986; Barnéoud et al., 1991; Azarfar et al., 2018), detect apertures just

large enough to pass through (Harris et al., 1999; Krupa et al., 2001; Grant et al., 2018; Heimburg,

Mari Saluti & Timm et al., 2025), follow walls at a consistent distance (Sofroniew et al., 2014;

Sofroniew & Vlasov et al., 2015), or avoid obstacles during locomotion (Warren et al., 2021). The

presence of map cells in the brainstem may significantly enhance the speed and precision of

these sensorimotor computations. Additionally, map cells could be better suited for 3D shape

perception, allowing the brain to detect small variations in contact distance across the whisker

array to reconstruct an object's surface geometry. In contrast, proximity cells excel at decoding

the closest distance, making them well suited for threshold-like detection of imminent contact in

peri-head space and for triggering rapid, reflexive responses, while simultaneously encoding

distance-invariant features such as texture, softness, or compliance.

While proximity and map cells may lie on a continuum of distance tuning rather than

forming strictly discrete classes, the distinction provides a useful framework for comparing

representational strategies along the lemniscal pathway. In TG, radial distance encoding is

mechanically driven and predominantly proximity-based (Szwed et al., 2006; Ahissar & Knutsen,

2008; Pammer & O'Connor et al., 2013; Birdwell et al., 2007; Solomon & Hartmann, 2011; Bush

et al., 2016). In PrV, we identified the presence of map cells, but proximity cells are

overrepresented. In wS1, a more balanced distribution of distance preferences was reported

(Sofroniew & Vlasov et al., 2015). This gradient suggests a progressive transformation from low-

level mechanical inputs to higher-order, feature-selective representations of peri-head space that

are increasingly aligned with the demands of sensorimotor behavior.

Transforming proximity inputs into a map code via feedforward inhibition

Our results identify a circuit mechanism by which the whisker brainstem transforms a simple

proximity code into a map code. SpVi-TeLC experiments and computational modeling support a

canonical E-I-E motif—monosynaptic excitation from TG combined with long-range disynaptic

inhibition from SpVi—that converts monotonic, proximity-based TG input into non-monotonic,

distance-selective responses in PrV map cells. Specifically, inhibitory inputs from SpVi act as

"neural comparators": by subtracting proximity signals with different distance sensitivity profiles,

they allow PrV map cells to exhibit peak activity at specific distances. Our findings extend beyond

prior work showing that SpVi inhibition sharpens angular tuning in PrV (Bellavance et al., 2010)—

a form of gain modulation—by demonstrating that the same projection can fundamentally change

how features are encoded, revealing a broader function for disynaptic inhibition in sensory code

transformation.

More broadly, these findings situate the SpVi-PrV pathway within a growing class of

microcircuits involving inhibitory neurons that implement core sensory computations. Across

modalities, feedforward inhibition does not simply limit excitation but actively extracts features by

implementing subtractive comparison operations to transform stimulus representations.

Examples include binaural circuits in the lateral superior olive of the auditory brainstem, where

excitatory and inhibitory inputs carrying sound level information from the two ears are compared

to encode interaural level differences (ILDs; Benichoux & Tollin, 2018), inhibitory subnetworks that

sculpt orientation and higher-order tuning in visual cortex (Lee et al., 2012; Znamenskiy et al.,

2024), and lateral inhibition in olfactory circuits that decorrelates odor representations (Giridhar

et al., 2011; Geramita et al., 2016). Recent connectomic analyses have revealed that inhibitory

neurons form stereotyped, feature-targeted connectivity patterns (Schneider-Mizell et al., 2025),

supporting the idea that inhibition is a commonly repeated anatomical substrate poised to perform

computations underlying neural code transformations. By showing that this convergent E-I-E motif

(Figure 1B; Figure 6A) in the first central processing center of the whisker system can transform

monotonic inputs from the periphery into a discrete distance map, our work supports a unifying

view of this motif as a computational primitive for constructing higher-order sensory

representations.

Map code formed with and without multi-whisker integration

We initially hypothesized that the PrV map code emerges through integration of inputs from

multiple whiskers. Our whisker cutting experiments, although technically challenging and low yield

due to inadvertent removal of inputs from the recorded neurons' whisker receptive field, showed

that a subset of PrV map cells retained map-like distance tuning even when only a single whisker

remained. These observations indicate that multi-whisker integration supports, but is not required

for, the generation of map-like tuning in a subset of PrV neurons. Together with the diversity of

mechanosensory thresholds among TG touch neurons (Hao & Delmas, 2010; Furuta et al., 2020),

these observations refined our circuit model whereby PrV constructs the map code by subtracting

inputs from TG afferents (from same or different whiskers) with different proximity-tuning profiles.

Our study does not rule out peripheral mechanisms that may contribute to the map code.

Mechanosensory receptors embedded in the whisker pad skin (Rice et al., 1986; Fundin et al.,

1994; Rice et al., 1997; Ebara et al., 2002; Severson et al., 2019), including Ruffini-like and other

low-threshold mechanoreceptors that known to encode skin stretch in dorsal root ganglion and

TG (Nordin & Hagbarth, 1989; Edin & Johansson, 1995; Roudaut et al., 2012; Abraira & Ginty,

2013) and C-low-threshold mechanoreceptors (C-LTMRs) innervating the surrounding hairy skin

(Grigg, 1996; Li et al., 2011; Usoskin et al., 2015; Reynders et al., 2015), could convey additional

distance information to PrV. These signals may encode whisker pad tension and its dynamic

changes during wall contact, thereby shaping distance representations in ways not captured by

our current understanding of TG input. Notably, conventional extracellular recordings in TG

predominantly sample large-diameter neurons because high-impedance electrodes are biased

toward detecting large-amplitude spikes from cells with large soma (Buzsáki et al., 2012), making

it technically difficult to assess the contribution of C-LTMRs and other small-diameter afferents.

Future work will be necessary to disentangle the relative contributions of central and peripheral

mechanisms in constructing peri-head distance representations.

AUTHOR CONTRIBUTIONS

F.W. conceived the study. F.W., K.S.S, V.P., and W.X. designed the experiments. W.X. and K.S.S. performed all experiments. P.M.T. and V.P. provided training and guidance to W.X. on in vivo extracellular recording in the brainstem. K.S.S. and V.P. developed analysis code with input from W.X. H.Z. implemented the brainstem circuitry simulation. K.C. and M.L. advised on data analysis and statistical methods. J.T advised on viral constructs and viral injection surgeries. K.S.S., W.X., and V.P. analyzed the data. S.Z. generated GtACR and TeLC viruses. S.C. managed the mouse colony, performed genotyping, and bred all animals. W.X., K.S.S, and F.W. wrote the manuscript with input from all authors.

ACKNOWLEDGMENTS

We thank Drs. Jennifer Groh, David Kleinfeld, Lindsey Glickfeld, and John Pearson, as well as members of the Wang Lab, for insightful discussions of this study. We are grateful to Julia Dziubek, Ina Bando, and He Jiang for technical assistance, and to Priyadarshini Dutta and Gabrielle Sevrain for assistance with mouse colony maintenance. This work was supported by grants NS077986, DP1 NS137188, and a subcontract of U19NS137920 (to F.W).

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURES and FIGURE LEGENDS

Figure 1. Acute PrV recordings in a wall passing paradigm
(**A**) Conceptual illustration showing how whiskers maps peri-head space in a head-centered coordinate frame, shown from a top view. (**B**) Left, schematic of a hypothetical brainstem circuit. Two TG neurons (A and B, red) project to the brainstem, where neuron A provides excitatory input to a PrV output neuron (C, black), and neuron B synapses onto an inhibitory neuron (blue) in the brainstem trigeminal complex. Middle, simulated tuning curves for neurons A and B with different mechanical sensitivity profiles; shaded region represents the difference between the tuning curves. Right, The PrV output neuron C integrates these inputs via linear summation, resulting in a non-monotonic tuning curve with a peak at an intermediate peri-head distance. (**C**) Left, Schematic of the experimental setup. A head-fixed mouse ran on a treadmill while a wall moved from anterior to posterior past the mouse's face. Wall distance from the face was randomly varied between 7-23 mm in 2-mm increments on each pass. Each recording contains 15-20 repetitions per distance. Right. Example video frame showing multiple whiskers contacting the wall (moving left to right). Red arrow indicates measured distance between face and wall. Scale bar: 2 mm. (**D**) Example coronal brain section (4.8 mm posterior to Bregma) showing two silicon probe tracks through PrV labeled with DiI (red); cell bodies stained with Nissl (blue). (**E**) Example extracellularvoltage trace from one recording session. Colored bars indicate wall-passing periods.

Figure 2. PrV contains multiple neural codes for peri-head distance

(**A**) Example spike rasters of six single units (units 1-6) in PrV during repeated wall pass trials. Each row corresponds to one wall pass trial, and each dot corresponds to a time bin (50 ms) containing at least one spike, colored according to wall distances. Wall passing begins at time 0. Wall contact (grey bar) begins between 2-3 seconds and ends between 9-11 seconds, depending on the wall distance and whisker length. Scale bar: 20 passes. (**B**) Peri-event time histograms (PETHs) computed as spike density functions aligned to wall onset time (t=0 s), for the same example units shown in A. Error shading: SEM. (**C**) Z-scored tuning curves (mean ± 95% CI) to wall distance for the same example units shown in A, B. (**D**) Heatmap of normalized tuning curves for wall-activated, distance-tuned units

Figure 2 (continued)
(n=472), sorted by preferred wall distance from close (top) to far (bottom). Tuning curves were linearly interpolated between presented distances; unpresented interpolated distances are shown as white space. (**E**) Histogram of preferred wall distances, measured as the location of the tuning curve peak, from units in D. (**F**) Fraction of recorded single units (N=1141) in PrV that were defined as either untuned (gray), suppressed only (purple), map (yellow, prefer ≥ 8 mm), proximity (red, prefer < 8 mm), or ambiguous (orange), which could not be statistically defined as either proximity or map (See Supplemental Figure S3).

Figure 3. The map code enables more accurate decoding of peri-head distance

(**A**) Decoding accuracy over time averaged across all distances for population decoder models trained on subsampled to 118 units from either proximity (pink, n=266), map (yellow, n=118), untuned (black, n=550), or trial-shuffled control (gray, n=384) subpopulations. Shaded area: 95% confidance interval. Wall contact occurs around 3-11s (gray bar) after the start of wall movement at t=0. (**B**) Average confusion matrices showing proportion of wall distance decoded for each actual distance from proximity (left) and map (right) populations. (**C**) Bar plot showing mean ± SEM population decoding accuracy for proximity (red), map (yellow), untuned (grey), and shuffled control (white); *, $p<0.05$, Wilcoxon rank-sum test with BHFDR correction. (**D**) Matrix quantifying error, in mm, for each coordinate in the confusion matrix. (**E**) Bar plot showing mean decoding error ± SEM in mm for decoders trained on each population (same colors as D); *, $p<0.05$, Wilcoxon rank-sum test with BHFDR correction. Error was computed by multiplying the confusion matrix multiplied with the absolute value of the error matrix and summing across rows.

Figure 4. Multi-whisker input supports, but is not required for, the distance map code

(**A**) Top, Example video frame showing multiple whiskers during wall-passing stimulation (wall moving left to right). Middle, same session during whisker cutting. Bottom, same session after cutting to a single B1 whisker. Scale bars: 2 mm. (**B**) Pairwise comparisons of firing rates (log scale) during wall-passing periods for wall-activated and distance-tuned units (n=86) in the full-whisker versus single-whisker condition. Gray lines connect the same unit across conditions. **, Population mean firing rates: 11.58 ± 2.986 sp/s (full array) vs. 8.18 ± 2.602 sp/s (single whisker), $p = 4.43 × 10^{-8}$). *, individual neuron, $p < 0.05$, Wilcoxon signed-rank test. (**C**,**D**) Two example map units that converted to proximity units after cutting down to a single whisker (B1, top; γ, bottom): (C) Spike density functions (SDFs); (D) corresponding tuning curves. (**E**,**F**) Example map unit that retained map-like tuning but

Figure 4 (continued)

shifted its preferred wall distance closer after cutting down to a single C2 whisker. (**G**,**H**) Two example map units showing gain modulation after cutting down to a single whisker (C1, top; β, bottom): Top, decreased activity; Bottom, increased activity (disinhibition). (**I**,**J**) Two proximity units showing similar gain modulation effects: Top, decreased activity; Bottom, increased activity (disinhibition). (**K**) Pairwise comparison of preferred wall distances for wall-activated units (n=33) that remained distance-tuned after cutting down to a single whisker. Open circles: map units; closed circles: proximity units. Color indicates tuning category: proximity (red), ambiguous (orange), map (yellow).

Figure 5. Silencing intersubnuclear inhibitory projections from SpVi diminishes the PrV map code

(**A**) Viral-genetic strategy to express TeLC in inhibitory cells in caudal SpVi with long-range projections to PrV. (**B**) Example coronal section at 4.6 mm posterior to Bregma, showing axons of TeLC-labeled inhibitory cells (TeLC-EYFP, green) in PrV and a silicon probe recording track labeled with DiI (red). Scale bar: 200 μm. Cell bodies were stained with Nissl (blue). (**C**) Heatmaps of normalized tuning curves for wall-activated, distance-tuned units in PrV. Left, wild-type mice (same data as Fig. 2D, reproduced here for side-by-side comparison). Right, a separate cohort of Vgat-Cre mice with TeLC-mediated silencing of PrV-projecting inhibitory neurons (n=252 units). Units are ordered by preferred wall distance, from near (top) to far (bottom). The color bar to the left of each heatmap indicates unit tuning category. Tuning curves were linearly interpolated between tested wall distances; untested (interpolated) distances appear as white space. (**D**) Cumulative distributions of preferred distances for the same units shown in C, comparing wild-type (black) and SpVi-TeLC (green) populations ($p = 8.02 \times 10^{-5}$, one-tailed KS test). (**E**) Comparison of the proportions of distance-tuned units recorded in wild-type mice (n=1141) and in SpVi-TeLC (n=587) mice. TeLC-mediated silencing of PrV-projecting inhibitory neurons in SpVi significantly reduced the prevalence of map units (3.90 vs 10.34%; Z-statistic: 4.64, $p = 1.80 \times 10^{-6}$, one-tailed Z-test) and suppressed units (7.30 vs 10.43%; Z-statistic: 2.12, $p = 0.02$, one-tailed Z-test), but increased the prevalence of proximity units (32.09 vs 23.31%; Z-statistic: -3.93, $p = 4.27 \times 10^{-5}$, one-tailed Z-test). The proportions of untuned and ambiguous units were not significantly different between samples from wild-type and TeLC mice. Asterisks indicate significance.

Figure 6. Simulation of brainstem circuitry to generate distance-map code

(**A**) Schematic diagram of the modeled neural circuit. The input layer consists of a pool of TG neurons (gray-filled circles), whose tuning to wall distance was modeled as sigmoidal functions with varying thresholds (indicated by the outline color of each circle). These TG neurons project to PrV neurons (black circles, output layer) via two distinct pathways: a direct excitatory (lemniscal) pathway (red) and an indirect inhibitory pathway (blue). The inhibitory pathway is mediated by a population of inhibitory neurons located in SpVi (blue circles), which project to PrV neurons. (**B**) Top, TG tuning curves (color-coded to match TG neurons outlined in A) are modeled as sigmoidal functions with thresholds spanning the full range of wall distances. Bottom, The sigmoid threshold distribution is exponentially skewed to emphasize the encoding of proximal wall distances. (**C**) Example tuning curves of modeled PrV neurons. Y-axis: baseline-subtracted activity predicted by the model (A.U.). Wall distances are normalized between 0 and 1, and nine uniformly spaced distance bins (labeled 0-8) are used for analysis and visualization. (**D**) Left, Heatmap of normalized tuning curves for distance-tuned units in the moddled PrV population, sorted by preferred distance bin from close (top) to far (bottom). Right, Same analysis after reducing inhibitory inputs to each PrV neuron by 50%. (**E**) Left, Histogram of preferred distance bins, defined as the peak (max) of the tuning curve, for modeled PrV neuron. Right, Same analysis after reducing inhibitory inputs to each PrV neuron by 50%. (**F**) Left, Proportion of modeled PrV neurons classified as proximity (red, prefer distance bin 0) or map (yellow, prefer > distance bin 0). Right, Same classification following 50% reduction of inhibitory inputs per modeled PrV neuron.

Figure S1. Wall-passing stimulus induced minimal active whisker retraction

(**A**) In a subset of mice (N=4), high-speed video was acquired during the wall-passing paradigm under both awake and anesthetized conditions. Example video frame with tracked whiskers color-coded left-to-right of an awake, behaving mouse: C3 (green), C2 (yellow), C1 (orange), and γ (red). Whisker angle (θ) was measured relative to the mediolateral axis (dotted line); example shown for C1 (orange). Scale bar, 2 mm. (**B**) Example trace of whisker angle (θ, black) and its midpoint (θ_{mid}, gray) from an awake, behaving mouse; θ_{mid} is the slowly varying centerline of θ. (**C**) Baseline-subtracted midpoint angle for the C1 whisker of 20 repetitions of one wall distance, aligned to wall movement onset (t = 0 s) in an awake, behaving mouse. Baseline was defined during the pre-wall period. Gray bar: wall contact period (3-11 s). (**D**) Comparison of whisker position between awake and anesthetized conditions in the wall-passing paradigm. Peri-event time histograms (PETHs) show changes in baseline-subtracted midpoint angle ($\Delta\theta_{mid}$) for C1, aligned to wall onset (t = 0 s) in one mouse. Left: awake, behaving; right: anesthetized. Each trace is the mean of 20 repetitions of the same wall distance with ±SEM shading. (**E**) Mean change in baseline-subtracted midpoint angle versus wall distance for the same C1 whisker shown in D. Green: awake, behaving; blue: anesthetized. (**F**) Same as E, but averaged across mice. Subpanels show mean change in midpoint angle versus wall distance for left: C1, middle: C2, right: Greek whiskers. Lines indicate across-mouse means ±SEM. Green: awake, behaving; blue: anesthetized. Significance reflects mouse-level paired comparisons (awake vs. anesthetized) with Bonferroni correction (α = 0.017): C1, p = 0.03 (n.s.); C2, p = 0.04 (n.s.), Greek: p = 0.0199 (n.s.).

Figure S2. Acute single-electrode recordings from PrV during the wall-passing paradigm

(**A**) Schematic of the experimental setup for acute single-unit recordings in PrV using tungsten electrodes. The wall stimulus moved past the mouse's face at randomized distances ranging from 4 to 20 mm in 1 mm increments, with 10–20 repetitions per distance. The mouse has the C-row whiskers intact. (**B**) Example coronal brain section at 4.8 mm posterior to Bregma showing one recording track through PrV labeled with DiI (red). Cell bodies were stained with Nissl (blue). (**C**) Example extracellular voltage trace from one recording session, showing two wall pass events. (**D**) Right, Peri-event time histograms (PETHs) computed as spike density functions aligned to wall onset time (t=0 s), for an example PrV unit preferring the closest distance presented. Error shading: SEM. Left, Z-scored tuning curve (mean ± 95% CI) to wall distance for the same unit. (**E**) Same as D, but for a PrV unit preferring an intermediate distance. (**F**) Heatmap of normalized tuning curves for all distance-tuned units (n=39), sorted by preferred distance from close (top) to far (bottom). Tuning curves were linearly interpolated between presented distances; unpresented interpolated distances are shown as white space.

Figure S3. Spike sorting quality metrics
(**A**) Inter-spike-interval (ISI) histograms for the example units in Figure 2A-C. The grey shaded region indicates ± 0.8 ms refractory period. Insets show the corresponding mean spike waveforms; shaded areas represent ± STD. Scale bars: 0.5 ms (horizontal), 0.25 mV (vertical). (**B**) Scatter plot of waveform signal-to-noise ratio (SNR) versus ISI false alarm (FA) rate for wild-type (WT) dataset collected with silicon probes (WT-Si Probe; n=1358 total units). The area within the grey square contains acceptable units with SNR > 5 and false alarm rate < 2% (n=1301 units from 36 mice). (**C**) Z-scored waveforms of a subset of acceptable units from the WT-Si Probe dataset (n=100 waveforms shown). (**D**) Histogram of spike widths of all acceptable units from the WT-Si Probe dataset. (**E**) Scatter plot of spike widths, in ms, versus mean firing rate, in spikes per second, for baseline periods (prior to wall pass) of acceptable units. (**F-I**) Same as (B-E), but for tetanus toxin (TeLC) dataset (n=665 total units from 11 mice; n=631 acceptable units in green shaded area). (**J-M**) Same as (B-E), but for tungsten dataset (n=46 total units from 14 mice; n=46 acceptable units in grey shaded area; n=46 waveforms shown in K).

Figure S4. Co-existence of proximity and map codes for encoding peri-head distance

(**A**) Cumulative distribution functions (CDFs; log scale) of baseline firing rates for wall-activated, distance-tuned units (red), wall-suppressed, distance-tuned units (blue), and untuned units (black). Wall-activated units had significantly lower baseline firing rates than untuned units (5.46 ± 0.670 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 1.91 × 10^{-10}, two-tailed Kolmogorov–Smirnov test), while wall-suppressed units had firing rates comparable to untuned units (7.31 ± 1.124 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 0.27, two-tailed KS test). (**B**) Left, CDFs of firing rates for wall-activated, distance-tuned neurons during wall passes (red) versus baseline periods (black). Firing rates were significantly higher during wall contact at the preferred distance (16.44 ± 1.640 vs. 6.36 ± 0.704 sp/s, mean ± SEM; p = 9.26 × 10^{-28}, one-tailed KS test). Right, CDFs of firing rates for wall-suppressed, distance-tuned neurons during wall passes (blue) versus baseline periods (black). Firing rates were significantly lower

Figure S4 (continued)
during wall contact at the preferred distance (4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s, mean ± SEM; p = 3.06 × 10^{-4}, one-tailed KS test). (**C**) CDFs of firing rates for untuned neurons during wall passes (gray) and baseline periods (black). No significant difference was observed (6.65 ± 0.687 vs. 6.90 ± 0.682 sp/s, mean ± SEM; p = 0.74, two-tailed KS test). (**D**) Heatmap of normalized tuning curves for suppressed units (n=213), sorted by preferred wall distance from close (top) to far (bottom). Units that were both activated and suppressed appear both in here and in Figure 2D. Tuning curves are linearly interpolated for between distances presented and omitted (white space) for interpolated distances not presented. (**E**) Cross-validated heatmaps of normalized tuning curves for activated units designated as either "map" (n=118) or "proximity" (n=266), for which trials were randomly partitioned into two equal partitions. Units in left panel were sorted by their preferred wall distances (close-top, far-bottom). Units in the right panel were sorted using the same order as the left panel. (**F**) Histogram of the location of tuning curve troughs from suppressed units in D. (**G**) Z-scored tuning curve (mean ± 95% CI) to wall distance for an example proximity unit (left), ambiguous unit (middle) determined by (i) not significant on the map test (one-tailed Wilcoxon signed-rank test, p = 0.09; n.s., not significant), and (ii) not having the highest firing rate at the closest distance, and map unit (right) determined by one-tailed Wilcoxon signed-rank tests comparing firing at each non-closest distance to the closest distance (p = 1.27 × 10^{-4}; *, p < 0.05 with BHFDR correction).

Figure S5. Single-neuron decoding performance
(**A**) Confusion matrices showing proportion of predicted wall distances for each actual distance from two example single-neuron decoders. Left, Example proximity unit (Unit 1 in Fig. 2A-C). Right, Example map unit (Unit 3 in Fig. 2A-C). (**B**) Mean decoding accuracy (± SEM) of single-neuron decoders grouped by tuning category: proximity (pink, n=266), map (yellow, n=118), untuned (grey, n=550), and trial-shuffled control (white, n=1141). *, $p < 0.05$; n.s., not significant (Wilcoxon rank-sum test with BHFDR correction). (**C**) Mean decoding error (± SEM) in mm by tuning category (same colors scheme as in B). Error was computed as the elementwise product of the confusion matrix and the absolute error matrix (Fig. 3D), summed across rows. *, $p < 0.05$; n.s., not significant (Wilcoxon rank-sum test with BHFDR correction).

Figure S6. Inhibitory circuit mechanisms contribute to the peri-head distance map code
(**A**) Schematic for expression of the inhibitory opsin, GtACR, in local inhibitory cells in PrV. (**B**) Schematic of acute silicon probe recording paired with optogenetic inhibition of local inhibitory interneurons in PrV. Continuous light stimulation (470 nm, 2 mW) was applied during the entire wall pass with 5s inter-trial interval and randomized laser on/off trials. (**C**) SDFs (in spikes/second) aligned to wall onset time of an example putative inhibitory neurons displaying laser silencing effect during wall passes. "Laser OFF" (left) and "Laser ON" (right) trials were randomly interleaved and repeated for each wall distance. Error shading: SEM. (**D**) Example coronal section at 5.1 mm posterior to Bregma, showing the GtACR-labeled inhibitory cells (green) and two silicon probe tracks through PrV, outlined with white dashed line. Cell bodies were stained with Nissl (blue). Scale bar: 1 mm. (**E**) Distribution of modulation index (ML) for all acceptable units (n=112) in the GtACR dataset. Units were classified as strongly inhibited (MI ≤ −0.5, dark blue), inhibited (−0.5 < MI < 0, light blue), disinhibited (MI > 0, red) when their bootstrap MI confidence interval excluded zero, and as no

Figure S6 (continued)

change when the confidence interval included zero (gray). (**F**) Top, SDFs of an example proximity cell displaying laser-induced disinhibition across all wall distances presented. (**G**) Top, Z-scored tuning curves showing mean firing rate (± 95% CI) for each wall distance for the same example unit shown in F top for Laser OFF and ON conditions. Bottom, Z-scored tuning curves for the same example unit shown in F bottom. (**H**) Example coronal section at 7.4 mm posterior to Bregma, showing the TeLC-labeled inhibitory cells (TeLC-EYFP, green) in SpVi, outlined with white dashed line. Cell bodies were stained with Nissl (blue). Scale bar: 200 μm. (**I**) Left, CDFs of baseline firing rates for untuned units in SpVi-TeLC mice (green) and wild-type mice (black). Firing rates were significantly higher in SpVi-TeLC mice (9.34 ± 1.330 vs. 6.90 ± 0.682 sp/s; p = 7.63 × 10^{-5}, one-tailed K-S test). Right, CDFs of mean firing rates for the same populations during wall passes. Untuned units in SpVi-TeLC mice showed significantly elevated firing compared to wild-type mice (9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s; one-tailed K-S test, p =1.06 × 10^{-4}). (**J**) Left, CDFs of baseline firing rates for wall-activated, distance-tuned units in SpVi-TeLC mice (green) and wild-type mice (black). No significant difference was observed (5.60 ± 0.845 vs. 5.46 ± 0.670 sp/s; one-tailed K-S test, p = 0.14). Right, During wall contact at the preferred distance, these same SpVi-TeLC units exhibited significantly elevated firing (20.25 ± 2.263 vs. 16.44 ± 1.640 sp/s; one-tailed K-S test, p = 1.64 × 10^{-3}). Asterisks indicate p < 0.0125 (Bonferroni-corrected, four comparisons). n.s., not significant.

STAR* METHODS

Lead Contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Fan Wang (fan_wang@mit.edu)

Materials Availability
This study did not generate new unique reagents.

DATA AND SOFTWARE AVAILABILITY
MATLAB scripts and analysis code used in this study, as well as processed datasets in NWB (Neurodata Without Borders) format, will be made available on Github and DANDI.

EXPERIMENTAL MODEL DETAILS
Adult C57BL/6J (JAX stock #000664) and vGat-IRES-Cre (JAX stock #016962; Slc32a1^tm2(cre)Lowl/J) mice were obtained from the Jackson Laboratory. During behavioral and electrophysiological experiments, mice were singly housed under a 12-hour light/dark cycle with ad libitum access to food and water. All experiments were conducted during the light phase. Details regarding the sex and age of all animals are provided in Table S1. All procedures were performed in accordance with protocols approved by the Institutional Animal Care and Use Committees at the Massachusetts Institute of Technology.

METHOD DETAILS

Headpost Implantation and Craniotomy
Mice were anesthetized with 3% isoflurane in 0.8 L/min oxygen and positioned in a stereotaxic frame (Model 963, David Kopf Instruments). Anesthesia was maintained at 1.5% isoflurane throughout the surgery. A custom stainless steel headbar (Pololu; 0.060″ ± 0.004″, type 304) and a stainless-steel ground screw were affixed to the skull using C&B Metabond (Parkell, #S380) and reinforced with dental cement (Stoelting). Mice were allowed at least seven days to recover following head-post implantation surgery. On the electrophysiological recording day, a 1 mm × 1 mm craniotomy was created 4.7 mm posterior and 2.0 mm lateral to Bregma using a dental drill (Aseptico). The dura was carefully removed to provide access to the principal trigeminal nucleus (PrV). Craniotomies were acutely sealed with silicone elastomer (Kwik-Cast, WPI).

Viral Vectors
The AAVs used included AAV-EF1a-DIO-GtACR2-EYFP (Takatoh & Prevosto et al., 2021), AAV-hSyn-FLEX-TeLC-P2A-EYFP-WPRE (Addgene, #135391).

Viral Injections
For optogenetic silencing experiments, vGat-Cre mice received stereotaxic injections of 300 nl AAV-EF1a-DIO-GtACR2-eYFP into PrV at coordinates 4.7 mm posterior, 2.0 mm lateral to Bregma, and 4.5 mm below the brain surface. For chronic silencing of SpVi neurons projecting to PrV, 300 nl of either AAV-hSyn-FLEX-TeLC-P2A-EYFP-WPRE (diluted 1:3 in saline) was stereotaxically injected into caudal SpVi (7.4 mm posterior, 2.0 mm lateral to Bregma, and 5.6 mm

579 below the brain surface). Injections were performed using beveled glass micropipettes connected
580 to a microsyringe pump (UMP3, WPI) controlled by a Micro4 controller (WPI) at a rate of 30 nl/min.
581 Following injections, craniotomies were sealed with bone wax (Surgical Specialties, #902), and
582 the scalp was closed using nonabsorbable silk sutures (LOOK, #774B).
583
Wall-Passing Paradigm
After recovering from headpost implantation, mice were head-fixed using plate clamps (Thorlabs)
positioned over a custom linear treadmill (Severson & Xu et al., 2017) to promote running and
whisking. Mice were habituated to the treadmill for at least two days, followed by at least three
days of habituation on a wall-pass apparatus prior to recording. The passing wall was driven in
the mediolateral and anteroposterior dimensions by two linear actuators (Zaber, Models
LSM200B-T4 rev 2 and LSM050B-T4 rev 2). During each pass, the wall moved from anterior to
posterior across the mouse's face at a constant velocity of 10 mm/s. In the silicon probe
experiments, wall distance from the whisker pad (C2 follicle) was randomly varied across trials
between 7 and 23 mm in 2-mm increments (4-mm increments for optogenetic silencing
experiments), using a custom MATLAB script. In the tungsten electrode experiments, wall
distance was randomly varied between 4 mm and 20 mm in 1 mm increments. All experiments
were performed in the dark. After the mediolateral Zaber motor set the wall distance, a
randomized delay of 0.1 to 1 s preceded wall movement.
598
Each recording session in the silicon probe experiments included 15-20 repetitions per wall
distance, during which mice freely whisked and interacted with the wall. Tungsten electrode
experiments included 10-20 repetitions per distance. Whiskers were illuminated using an infrared
LED backlight (Roithner Lasertechnik, #LED660N-66-60) and imaged with a high-speed camera
(Basler, Model acA1300-200um) equipped with a telecentric lens (Edmund Optics, #55-349). Prior
to each recording, the wall apparatus was aligned to the midline of the mouse's head, and a
calibration image was acquired and converted to physical units (31 pixels/mm). The closest wall
distance was defined as the minimum Euclidean distance between the C2 whisker follicle and the
wall. Absolute wall distance (in mm) was calculated by adding the calibrated scaling factor to the
horizontal position (in 2-mm steps, relative to the closest position) of the stepper motor controlling
the wall's mediolateral position.
610
High-Speed Videography
Video frames (640 pixels x 480 pixels, 32 µm/pixel) were acquired at 500 Hz using a PhotonFocus
DR1-D1312-200-G2-8 camera (90 µs exposure time) and Streampix 5 software (Norpix). Light
from a 940 nm LED (Roithner Laser) was passed through a condenser lens (Thorlabs) and
directed into a 0.25X telecentric lens (Edmund Optics) after passing through the whisker field.
Ephus triggered individual camera frames (5 s, 2,500 frames per trial) synchronized with
electrophysiological recordings.
618
619 ***Awake and Anesthetized Wall-Passing Recordings***
620 In a subset of mice (N = 4), high-speed video was continuously acquired during the wall-passing
621 paradigm under both awake and anesthetized conditions. Mice were head-fixed throughout,
622 keeping their position constant relative to the moving wall stimulus. To enable whisker tracking,
623 all whiskers except a single row (C1-C4 and γ or β) were trimmed. While awake, each mouse

experienced wall passes with randomized wall distances (7-23 mm in 2-mm increments), repeated for 20 passes per distance. Following the awake session, the same animals were anesthetized via intraperitoneal injection of a fentanyl-based cocktail (0.05 mg/kg fentanyl, 5 mg/kg midazolam, 0.5 mg/kg dexmedetomidine). After a 10-minute induction period, the same wall-passing protocol was repeated under anesthesia. Comparisons between awake and anesthetized conditions were performed within-subject across matched wall distances.

Electrophysiology Recording: Silicon Probe

Following habituation to the wall-passing paradigm, mice were head-fixed and allowed to run on the treadmill. The craniotomy was exposed and covered with artificial cerebrospinal fluid (aCSF). For wild-type and TeLC experiments, the tip of a 32-channel silicon probe (NeuroNexus, A1x32-Poly2-10mm-50s-177-A32 or A1x32-Poly3-10mm-50s-177) was coated with the fluorescent dye DiI (Invitrogen™, #V22885). For optogenetic silencing experiments, a 473 nm laser (Cobolt, 06-MLD 473) was connected to a ferrule attached to a 32-channel silicon optrode (NeuroNexus, A1x32-Poly2-10mm-50s-413, 105 µm/125 µm, 0.22 NA). The probe or optrode was advanced approximately 5 mm into the brain using a micromanipulator (Sutter Instruments, Model MPC-325) until it reached ventral PrV and well-isolated units responsive to manual whisker stimulation were encountered. The brain was allowed to relax for at least 40 minutes to ensure stable recordings. Neural signals were hardware-filtered on the headstage (Blackrock Microsystems, Cereplex M, #10532) between 0.3 Hz and 7.5 kHz for spikes and digitized at 30 kHz using Cereplex Direct software (Blackrock Microsystems, Version 7.0.5). Analog voltage signals from the anterior-posterior stepper motor were simultaneously acquired at 30 kHz. Wall pass start and end times were determined from the rising and falling edges of these voltage signals to synchronize electrophysiological recordings with wall movement. Recording sites were subsequently confirmed by post hoc histological analysis.

Electrophysiology Recording: Single Tungsten Electrode

One day prior to recording, whiskers and facial fur on the left side of the face were trimmed using fine forceps and microdissection scissors (FST, #91500-09) under isoflurane anesthesia (1.5%), sparing the C-row whiskers (γ or β, C1–C5). On the recording day, mice were head-fixed and allowed to run on the treadmill. The craniotomy was exposed and kept covered with aCSF. The tip of a tungsten microelectrode (0.005″ diameter, 5 MΩ, 8° tip angle; A-M Systems, #575300) was coated with the fluorescent dye DiI and lowered approximately 5 mm into the brain using a micromanipulator until it reached ventral PrV. Electrical activity was monitored via audio output from the amplifier (WPI, DAM80). Manual whisker stimulation was performed to identify a well-isolated, whisker-responsive unit. The brain was allowed to relax for at least 20 minutes to achieve stable recordings. Differential signals between the recording and reference electrodes were amplified 10,000× and bandpass filtered between 300 and 3,000 Hz (WPI, DAM80). Following recording, receptive fields and response types (rapidly adapting [RA] or slowly adapting [SA]) were manually classified. Analog voltage signals from the amplifier and the anterior-posterior stepper motor were acquired using Cereplex Direct at a sampling rate of 30 kHz. Wall pass start and end times were extracted from the rising and falling edges of the stepper motor TTL signals to synchronize electrophysiological recordings with wall movement. All recording sites were confirmed by post hoc histological analysis.

Isolation of Single Unit Activity

For silicon probe data, initial single-unit isolation (spike sorting) was performed using Kilosort2.5 (Pachitariu et al., 2016), followed by manual curation in Phy (github.com/cortex-lab/phy; Rossant, 2025). Spike times and waveforms for isolated clusters were exported from Phy. Most units exhibited narrow spike widths, defined by a trough-to-peak latency of less than 0.4 ms. Duplicate units, identified as those with high Pearson's correlation coefficients (> 0.1) in spike counts binned at 1 ms resolution, were excluded from further analysis. For tungsten electrode data, spike waveforms were extracted by thresholding high-pass filtered traces (500 Hz) and clustered using MClust-4.4 (A.D. Redish, http://redishlab.neuroscience.umn.edu/MClust/MClust.html).

Isolation quality was assessed using the following metrics: signal-to-noise ratio (SNR), defined as the mean waveform amplitude divided by the standard deviation (SD) of waveform noise; and false alarm (FA) rate, defined as the percentage of spikes with inter-spike intervals (ISI) shorter than 0.8 ms. Units with an SNR less than 5 SDs or an FA rate greater than 2% did not meet single-unit isolation quality criteria and were excluded from subsequent analyses. For silicon probe data, in wild-type mice, 1141 of 1358 units (36 mice), in TeLC experiments, 587 of 665 units (11 mice), and for Tungsten recordings (46 of 46 units (14 mice), passed isolation quality and minimum firing rate (>1 Hz) criteria and were subsequently entered into population analyses. For acute whisker cutting experiments, only units demonstrating stable waveform properties both before and after cutting were included in analyses.

Acute Whisker Cutting Manipulation

During recording sessions involving acute whisker cutting, neural responses were first recorded while the full whisker array was intact. Each session included 15 to 20 repetitions of wall passes over approximately one hour. To best guess the primary whisker to spare during acute trimming, the receptive fields of PrV neurons were mapped by manually stimulating individual whiskers with a cotton-tipped applicator while monitoring spike responses. All whiskers except the designated spared whisker were then cut down to the base of the shaft using microdissection scissors (FST, Cat# 91500-09). After trimming, the spared whisker and surrounding whisker pad were brushed with a cotton-tipped applicator to stimulate all touch-sensitive recorded units. The wall-passing procedure was then repeated for an additional 15 to 20 trials. After completing the post-cutting wall passes, the whisker and pad were brushed once more to confirm the continued presence of responsive units. Electrophysiological signals were continuously recorded throughout the pre-cut, cutting, brushing, and post-cut stimulation periods.

Optogenetic Silencing of PrV Local Interneurons

Light pulses were generated using a pulse generator (Doric, #OTPG-4), with laser timing synchronized to wall movement via Doric software. Continuous 473 nm light stimulation (2 mW) was delivered by a laser (Cobolt, #06-MLD 473) throughout each wall pass, with a 5-second inter-trial interval. Each recording session consisted of 30 to 40 repetitions at each wall distance, ranging from 7 to 23 mm in 4 mm increments, conducted over approximately two hours with randomized "Laser Off" and "Laser On" trials.

Histology
Following recordings, mice were deeply anesthetized and transcardially perfused with phosphate-
buffered saline (PBS; Gibco), followed by 4% paraformaldehyde (PFA) in PBS. Brains were post-
fixed overnight and then cryoprotected in 30% sucrose in PBS at 4°C. Tissue was embedded in
OCT compound (Sakura Finetek) and coronally sectioned at 100 μm using a cryostat (Leica
Biosystems). For wild-type, optogenetic inhibition, and tungsten recording experiments, sections
containing PrV were stained with Neurotrace Blue (1:500; Thermo Fisher Scientific, #N21479) in
0.3% Triton X-100/PBS. For TeLC experiments, sections containing PrV and SpVi underwent
immunostaining with goat anti-GFP (1:500; Rockland, #600-101-215) in blocking buffer, followed
by either Alexa Fluor 488 (1:500; Jackson ImmunoResearch, #705-545-147) or Alexa Fluor 647
(1:500; Jackson ImmunoResearch, #705-606-147) donkey anti-goat secondary antibodies and
Neurotrace Blue (1:500; Thermo Fisher Scientific, #N21479) in blocking buffer. The sections were
then briefly rinsed with 0.3% Triton X-100/PBS and mounted on slides with a custom-made
mounting medium. All histological images were acquired with either a Keyence microscope (BZ-
X800) or a Zeiss LSM700 confocal microscope.
727

QUANTIFICATION AND STATISTICAL ANALYSIS
All analyses were performed in MATLAB (MathWorks, versions 2021b and 2023b). Data are
reported as mean ± standard errors of the mean (SEM) unless otherwise noted. Measures of
central tendency, variability, test statistics, p-values, degree of freedom, and figure panel
assignments for all analyses are detailed in Table S6. We analyzed single-unit
electrophysiological recordings from n = 1,187 units in PrV across 63 mice, as well as high-speed
video recordings from four sessions in four mice. The video data comprised 15,679,458 frames,
totaling 522.6 minutes. Sample sizes were not predetermined.
736

Video Analysis
The backbone of each whisker was tracked at subpixel-resolution using the Janelia Whisker
Tracker (Clack et al., 2012), yielding a set of "trajectories" (tracked objects in image X-Y
coordinates) for each frame. All subsequent processing of whisker angle (θ) was conducted in
MATLAB according to published methods (Pammer & O'Connor et al., 2013; Severson & Xu et
al., 2017), with several modifications described below in "Video analysis" subsections.
743

Video Analysis: Identifying Tracked Whiskers
To identify the same whiskers across frames (linking), whiskers were ordered by follicle x-position.
Trajectories with follicle x- or y-position outside the region of interest (bounding box), with arc
length shorter than 150 pixels (4.65 mm), or whisker "score" (Janelia whisker tracker
measurements) less than 600 were ignored. Post-hoc median filtering (3-frame window) was
applied to smooth over any remaining linking errors.
750

Video Analysis: Face Masking
Computing time series of whisker angle relies on being able to reliably track the base of the
whisker near the follicle location across all video frames. As previously described (Pammer &
O'Connor et al., 2013), we used a "mask" to truncate the tracked whisker trajectories near the
face to remove "noise" due to occlusion by facial fur. The follicle location was then estimated by
extrapolating the angle of the whisker base past the intersection of the whisker and the mask
(Pammer & O'Connor et al., 2013). We fit a separate mask for each frame, obtained using a
custom algorithm that fitted a smoothing spline to the contour of the face. We blurred the image
using a Gaussian filter, binarized the image, and detected the contour by taking a spatial derivative

along the y-dimension. The pixel locations of the facial contour were thus detected as the
intersection between the white background and black shadow cast by the face. A smoothing spline
was then fitted to the facial contour positions and shifted upward by ~10 pixels to yield a 20-point
facemask that intersected with the whisker trajectories. Frames in which a tracked whisker did not
intersect the facemask were considered missing data.
765
Data Analysis: Wall-Induced Whisker Retraction, Awake vs Anesthetized
For each of the tracked whiskers (C1, C2, and Greek), whisker angle (θ) was measured relative
to the mediolateral axis of the face. The midpoint of the whisker angle trace (θ_{mid}) was defined as
the slowly varying centerline of θ. To obtain θ_{mid}, θ was first smoothed using a Savitsky-Golay filter
(21 frame, third order), followed by interpolation of the peaks and valleys to obtain the upper and
lower envelope of the θ trace. θ_{mid} was then calculated as the mean of this envelope. For each
wall pass, θ_{mid} was computed during both the baseline period (-4.5 to 0 s, with t=0 marking wall
onset) and the wall-passing period (3 to 11 s). Baseline-subtracted θ_{mid} ($\Delta\theta_{mid}$) was then calculated
for each pass. For each mouse (N = 4 mice), $\Delta\theta_{mid}$ was averaged across wall distances and
repetitions to obtain a single value per whisker and condition (awake or anesthetized). Data were
aggregated across mice into a summary table (mouse × whisker × condition), and paired
comparisons were made between awake and anesthetized conditions for each whisker.
778
To compare the effect of wall-induced whisker retraction of awake versus anesthetized conditions,
we performed two-tailed paired t-tests across mice (within-subjects) for each whisker. The null
hypothesis was that mean $\Delta\theta_{mid}$ was equal in awake and anesthetized conditions. Effect size was
computed as the mean difference between conditions (awake minus anesthetized), and
Bonferroni correction was applied to control for multiple comparisons across whiskers
(α =0.0167).
785
Data Analysis: Spike Density Function Calculation
Spike density functions (SDFs) were computed by aligning spike times to the onset of wall-pass
trials, with time zero defined as the moment when the wall began moving toward the mouse. For
all trial conditions, two epochs were defined consistently: a baseline period from −4.5 to 0 s, and
a wall-pass period from 3 to 11 s. Spike rasters were generated from the aligned spike times using
1 ms bins. SDF curves were then produced by convolving these spike rasters with a Gaussian
filter (5 ms SD, full kernel). SEMs were calculated as the SD of the SDF across trials divided by
the square root of the number of trials in each condition. The resulting SDFs and SEMs were
averaged across repetitions of the same wall distance and stored for each cell.
795
Data Analysis: Wall Distance Tuning Calculation
Wall distance tuning curves were generated by uniformly binning distance measurements into
nine 2-mm-wide bins with centers at 7, 9, 11, 13, 15, 17, 19, 21, and 23 mm. Distances falling
outside the bin edges were excluded. Peri-event time histograms (PETHs) were computed by
binning instantaneous firing rates in 200-ms intervals, aligned to the onset of anterior-posterior
wall movement as defined by the stepper motor TTL pulse (t=0). Each wall pass trial lasted
approximately 14 seconds. Average firing rates during the wall pass period (3–11 s) were
computed from the PETH and compared to baseline firing rates calculated from −4.5 to 0 s. For
each distance, mean firing rates were averaged across repeated passes to construct the tuning

curve. Standard errors were calculated by dividing the SD across repetitions by the square root
of the number of repetitions for each distance. CIs were estimated using bootstrap resampling
(10,000 iterations) and defined as the 95th percentile range. Z-scored tuning curves were
computed by subtracting the mean from firing rate across all baseline periods in the recording
session and dividing by the corresponding SD. Uniform binning enabled concatenation of data
across sessions for population-level analyses, including decoding analyses described later.
Sessions with fewer than nine presented distances were excluded from population analyses. For
population heatmaps, each neuron's tuning curve was normalized to its own minimum and
maximum firing rates, and the color scale represents the resulting normalized mean spike rate.
814
Data Analyses: Characterizing Single Unit Responses
To assess responsiveness to wall stimulation, one-tailed Wilcoxon signed-rank tests ($p < 0.05$)
were performed on each of 1,187 units. A unit was classified as "wall-activated" if its mean firing
rate during the wall pass period (3-11 s) at any distance was significantly higher than its baseline
firing rate (−4.5-0 s). Conversely, a unit was classified as "wall-suppressed" if its mean firing rate
during the wall pass period at any distance was significantly lower than baseline. Units exhibiting
significant increases at some distances and decreases at others were classified as both "wall-
activated" and "wall-suppressed."
823
824   Wall distance tuning was assessed using two-tailed Wilcoxon signed-rank tests across all pairs
825   of distances. A unit was classified as "distance-tuned" if the mean firing rate at one or more
826   distances significantly differed from that at other distances ($p < 0.05$, corrected for multiple
827   comparisons using the Benjamini-Hochberg false discovery rate method).
828
For comparisons of baseline firing rate distributions between distance-tuned and untuned units,
the mean and 95% confidence intervals (CI) were calculated using bootstrap resampling (10,000
iterations). The CI was determined by calculating half the range between the upper and lower
bounds of the bootstrap interval. Results were reported as mean ± CI. Statistical differences
between distributions were assessed using Kolmogorov-Smirnov (KS) tests, performed with one-
or two-tailed hypotheses as dictated by the experimental design (see Table S6). Significance was
determined at $p < 0.05$, with Bonferroni correction for multiple comparisons. The same procedures
were applied for comparing average firing rates during wall passes and baseline periods between
distance-tuned and untuned populations.
838
To further characterize tuning profile within "wall-activated" and "distance-tuned" units, we
performed one-tailed Wilcoxon signed-rank tests comparing mean firing rates at each non-closest
distance (≥ 9 mm) to the closest distance (7 mm). Units with significantly higher firing rates at any
non-closest distance were classified as "map units," while those exhibiting maximal firing at the
closest distance (7 mm) were classified as "proximity units." Units that met criteria for "wall-
activated" and "distance-tuned" but did not fulfill the criteria for either map or proximity were
labeled "ambiguous" and excluded from decoding and population cross-validation analyses.
846
For analyses of acute whisker cutting experiments, mean firing rates during baseline and wall-
pass periods were compared between full whisker array (control) and single-whisker (cut)
conditions. Pairwise comparisons of firing rates were conducted using two-tailed Wilcoxon signed-
rank tests. To assess individual neuron responses to whisker cutting, the proportions of units
exhibiting significant increases, decreases, or no changes in firing rates were computed.
Significance thresholds were corrected for multiple comparisons using Bonferroni adjustment. To

evaluate time-dependent plasticity, firing rates for each unit were partitioned into three temporal
segments: the first half of wall-pass trials (full whisker array), the second half (full whisker array),
and the post-manipulation period. For each unit, mean firing rates were computed separately for
these periods. Adaptation was assessed by comparing firing rates between the first and second
halves using two-tailed Wilcoxon signed-rank tests. A non-significant result ($p > 0.05$) indicated
stable firing prior to whisker cutting. Firing rates from the second half of the full whisker array
period were then compared to post-cutting periods using two-sided Wilcoxon signed-rank tests,
where significant differences indicated effects attributable to whisker cutting.

To compare the proportions of map, proximity, ambiguous, suppressed, or untuned cells between
SpVi-TeLC mice and wild-type mice, a one-tailed Z-test for proportions was performed. The null
hypothesis (H_0) stated that the proportion of each cell type in the SpVi-TeLC population (p_2) was
less than or equal to that in the wild-type population (p_1), while the alternative hypothesis (H_a)
posited that $p_2 > p_1$. Proportions (p_1 & p_2) were calculated as the number of cells of a given type
(x_1 & x_2) divided by the total number of recorded units (n_1 and n_2) in each group. A pooled
proportion (p_{pooled}) was used to estimate the standard error (SE) of the difference between groups:

869
$$p_{pooled} = \frac{x_1 + x_2}{n_1 + n_2}, \quad SE = \sqrt{p_{pooled}(1 - p_{pooled})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$$

870 The Z-statistic was computed as:

871
$$Z = \frac{p_1 - p_2}{SE}$$

P-values were calculated using MATLAB's cumulative normal distribution function (normcdf) and
evaluated at $\alpha = 0.05$. A p-value below this threshold indicated a significantly higher proportion of
the specified cell type in SpVi-TeLC mice compared to wild-type mice.

To compare the firing rate distributions between SpVi-TeLC and wild-type populations (Group 1
and Group 2), we computed the mean and 95% CI for each group using bootstrap resampling
(10,000 iterations). For each group, the results were reported as mean ± CI. To assess the
statistical significance of differences between the distributions of the two groups, we performed a
two-sample KS test. The test was conducted with a specified directional hypothesis (one- or two-
tailed, depending on the experimental design), and the resulting p-value was reported to
determine the significance ($p < 0.05$, with Bonferroni correction for multiple comparisons) of the
observed differences.

Data Analysis: Decoding of Wall Distance from Population or Single Neurons Activity
For population decoding of wall distance, we applied a population decoder (Neural Decoding
Toolbox, www.readout.info; Meyers, 2013; Vincis & Chen et al., 2020). Units from sessions with
all nine distances presented that passed spike sorting quality metrics (1,141 total units) were
concatenated into a pseudo-population. Populations were subsampled into sub-populations of
equal size (118 units), containing units designated as either "map", "proximity", or "untuned" cells.
Units in the "shuffled" subpopulation were randomly sampled from map and proximity
subpopulations and had their trial distance labels shuffled with replacement. Spike timestamps
for each unit were aligned to the onset of anterior-posterior wall movement (time 0) and were
binned using a 50 ms bin size with a 150 ms sliding window to create a firing rate matrix, where
each row corresponds to a trial and each column corresponds to a time bin. The firing rate matrix

spans from time 0 to 14 seconds. To ensure equal trial numbers across conditions (i.e., wall distances), trials were subsampled five times to match 15 trials per condition. The binned data and their corresponding condition labels were randomly split into 10 subsets—9 used for training the classifier and 1 reserved for validation. The process was repeated 10 times for each unique partition for 10-fold cross-validation to assess decoding accuracy. The classifier operates by learning a mean population vector (template) for each class from the training set. During testing, it calculates the Pearson correlation coefficient between a test point and the learned templates. During validation and testing, the class with the highest correlation value is assigned as the predicted label. Decoding accuracy (true positive rate) was averaged over the 3-11 second wall-pass period to obtain the overall decoding accuracy for the sampling epoch. In addition to the decoding accuracy, confusion matrices were computed at each time step. To assess variability, the entire subsampling procedure was repeated 10 times, and standard error was calculated by computing the SD of decoding accuracy across runs and dividing by the square root of 10.

For single-neuron decoders, we quantified how well individual neurons discriminated wall distance based on methods used in Jezzini et al., 2013. For each unit, spike trains were grouped according to nine discrete wall distances (7, 9, 11, 13, 15, 17, 19, 21, and 23 mm). For each distance, spike times were aligned to trial onset and converted into spike count vectors using 200 ms non-overlapping bins spanning the 3–11 s wall contact period. These binned firing rate profiles were used to construct condition-specific templates for distance classification. Decoding was performed using a Euclidean distance classifier, which assigned each test trial to the distance condition whose mean firing-rate vector (template) was closest in Euclidean space. Cross-validation was implemented as a leave-two-out procedure (xval = 2), in which two trials per condition were held out as test data, and the remaining trials were used for training. This was repeated exhaustively across all combinations of held-out pairs, such that every trial pair served as a test set at least once. Classification accuracy was computed as the fraction of correctly identified test trials, averaged across all cross-validation iterations. For shuffled controls, we performed a trial-shuffled permutation in which class labels were randomly reassigned across conditions (n = 1000 iterations).

For both population and single-neuron decoders, differences in mean decoding accuracy and error across wall distances were assessed using Wilcoxon rank-sum tests. Separate tests were performed for each of the nine wall distances, and p-values were corrected for multiple comparisons using BH-FDR to compare proximity and map populations.

Data Analysis: Acute Whisker Cutting Manipulation
To test for the effect for acute whisker cutting on firing rates, we computed each unit's mean firing rate during baseline and wall-passing periods on each repetition for the full whisker array (control) or single-whisker condition (manipulation). Pairwise comparison of mean firing rate for neural population and individual units between conditions was performed via two-sided Wilcoxon signed-rank test. To evaluate how individual neurons responded to whisker cutting, we computed the proportion of units that exhibited significant increases, decreases, or no change in firing rate between conditions. P-values were corrected for multiple comparisons across units using the BH-FDR procedure.

Data Analysis: Modulation Index in Acute Optogenetic Experiments

We quantified the acute optogenetic effects on firing rates using a modulation index (MI) computed for each unit as:

$$MI = \frac{\text{FR}_{LASER\ ON} - \text{FR}_{LASER\ OFF}}{\text{FR}_{LASER\ ON} + \text{FR}_{LASER\ OFF}}$$

where $\text{FR}_{LASER\ ON}$ and $\text{FR}_{LASER\ OFF}$ denote the mean firing rates during matched laser-on and laser-off wall-passing periods, respectively. For each unit, we obtained a bootstrap CI on the MI and classified the unit as significantly modulated if the CI did not include zero. Significantly modulated units were further assigned to one of three categories based on their MI: strongly inhibited (MI ≤ -0.5), inhibited (-0.5 < MI < 0), or disinhibited (MI > 0). Units whose MI confidence intervals overlapped zero were classified as no change. All analyses were restricted to units that passed our unit inclusion criteria (see Isolation of Single Unit Activity).

Computational Simulation of Brainstem Circuitry

To model the tuning properties of PrV neurons, we simulated a pool of TG afferents whose activities decreased monotonically with wall distance, consistent with their mechanical sensitivity. These proximity-based TG inputs projected to PrV neurons via a direct excitatory (lemniscal) pathway and an indirect inhibitory pathway (SpVi inhibitory projections to PrV). TG tuning curves were modeled as sigmoidal functions with thresholds spanning the distance range. The activity of each PrV neuron j (v_j^{PrV}) in response to wall distance $x \in [0,1]$ was defined as:

$$v_j^{PrV} = \sum_{i \in C_j^E} w_{ij} \sigma_{TG}(x; \beta_i, \theta_i) - \sum_{k \in C_j^I} w_{kj} \sigma_{TG}(x; \beta_k, \theta_k)$$

where

$$\sigma_{TG}(x; \beta, \theta) = \frac{1}{1 + \exp(\beta(x - \theta))}$$

is the TG neuron's sigmoidal tuning function. Here, β and θ control sensitivity and selectivity to wall distance, respectively. C_j^E denotes the set of TG providing direct excitatory input to PrV neuron j, with synaptic weight w_{ij}; C_j^I denotes TG neurons projecting indirectly to PrV via inhibitory neurons in SpVi, with weight w_{lj}. Subscript i and l denote each TG neuron in set C_j^E and set C_j^I. For simplicity, inhibitory neurons were modeled as providing a sign-inverted copy of TG activity, such that v_j^{PrV} reflect a linear combination of excitatory and inhibitory streams.

Intuitively, the subtraction of two sigmoidal functions can yield a non-monotonic, Gaussian-like function. Here we generalize this by considering a pool of sigmoidal functions (TG neurons). Specifically, we implemented several assumptions:

(1) Sparse connectivity: Both C_j^E and C_j^I represented small subsets of TG neurons. Parameter $\beta_i, \theta_i, \beta_l, \theta_l, w_{ij}, w_{lj}$ were independently sampled for each PrV neuron. The number of inhibitory inputs $|C_j^I|$ per PrV neuron j was treated as a random variable sampling from an exponential distribution parametrized by λ_c (truncated to the interval [1,30] in the spirit of preserving E/I balance):

$$P\left(\left|C_j^I\right| = k\right) = \frac{e^{-\lambda_c k}}{\sum_{m=1}^{30} e^{-\lambda_c m}}, k = 1,2,3,\dots 30, \qquad \lambda_c = 0.03$$

The number of excitatory inputs was fixed at $\left|C_j^E\right| = 20$.

(2) Skewed distribution of TG tuning. The TG threshold parameter θ was drawn from a
truncated exponential distribution within $[0,1]$:

$$\theta \sim f(x; \lambda_\theta) = \frac{\lambda_\theta e^{-\lambda_\theta x}}{1 - e^{\lambda_\theta}}, \qquad x \in [0,1]$$

where λ_θ describes the skewness of the distribution. This skewed distribution increases
the density of high-threshold TG neurons that only fire at proximal distances, thereby
enhancing encoding resolution and reducing overall metabolic cost (efficient coding). We
used $\lambda_\theta = 3$ for the excitatory inputs and $\lambda_\theta = 6$ for the more skewed inhibitory pathway.
Sensitivity parameters were sampled as $\beta_i, \beta_l \sim N(20, 5)$. Synaptic weights were sampled
from $w_{ij} \sim N\left(0, \frac{0.1}{|C_j^E|}\right)$ for the excitatory pathway and $w_{lj} \sim N\left(0, \frac{0.1}{|C_j^I|}\right)$ for the inhibitory
pathway.
The input range x is $[0,1]$ for simplicity. We sampled nine uniformly spaced points between 0
and 1 (labeled as 7,9,11,13,15,17,19,21,23 to correspond to empirical wall-distance
measurements for easy comparison) and evaluated tuning curves at these points. Neurons
with flat tuning curves (max - min < 0.5, untuned) were excluded. Among the remainder,
proximity cells were defined as neurons with maximal responses at x = 0, while map cells
were defined as neurons with peak responses at x > 0. In total, we modeled 1,225 PrV
neurons.

To simulate partial silencing of inhibitory projections, the distribution of inhibitory inputs was
truncated to approximately half of its original range [1,16], such that $\left|C_j^I\right|$ for each PrV neuron
j is decreased by:

$$P\left(\left|C_j^I\right| = k\right) = \frac{e^{-\lambda_c k}}{\sum_{m=1}^{16} e^{-\lambda_c m}}, \qquad k = 1,2,3,\dots 16$$

This reduced the expected number of inhibitory inputs per PrV neuron by half.
1004
1005
1006
REFERENCES

Abraira, V.E., and Ginty, D.D. (2013). The sensory neurons of touch. *Neuron* **79**, 618–639.

Ahissar, E., and Knutsen, P.M. (2008). Object localization with whiskers. *Biol. Cybern.* **98**, 449–
458.

Avendaño, C., Machín, R., Bermejo, P.E., and Lagares, A. (2005). Neuron numbers in the sensory
trigeminal nuclei of the rat: a GABA- and glycine-immunocytochemical and stereological analysis.
J. Comp. Neurol. **493**, 538–553.

Azarfar, A., Zhang, Y., Alishbayli, A., Miceli, S., Kepser, L., van der Wielen, D., van de Moosdijk,
M., Homberg, J., Schubert, D., et al. (2018). An open-source high-speed infrared videography
database to study the principles of active sensing in freely navigating rodents. *Gigascience* **7**,
giy134.

Bale, M.R., and Maravall, M. (2018). Organization of sensory feature selectivity in the whisker
system. *Neuroscience* **368**, 70–80.

Bechara, A., Laumonnerie, C., Vilain, N., Kratochwil, C.F., Cankovic, V., Maiorano, N.A.,
Kirschmann, M.A., Ducret, S., and Rijli, F.M. (2015). Hoxa2 selects barrelette neuron identity and
connectivity in the mouse somatosensory brainstem. *Cell Rep.* **13**, 783–797.

Barnéoud, P., Gyger, M., Andrés, F., and van der Loos, H. (1991). Vibrissa-related behavior in
mice: transient effect of ablation of the barrel cortex. *Behav. Brain Res.* **44**, 87–99.

Bellavance, M.A., Demers, M., and Deschênes, M. (2010). Feedforward inhibition determines the
angular tuning of vibrissal responses in the principal trigeminal nucleus. *J. Neurosci.* **30**, 1057–
1063.

Belli, H.M., Yang, A.E., Bresee, C.S., and Hartmann, M.J. (2017). Variations in vibrissal geometry
across the rat mystacial pad: base diameter, medulla, and taper. *J. Neurophysiol.* **117**, 1807–
1820.

Benichoux, V., and Tollin, D.J. (2018). Sound localization: These are not the neurons you are
looking for. *eLife* **7**, e39244.

Birdwell, J.A., Solomon, J.H., Thajchayapong, M., Taylor, M.A., Cheely, M., Towal, R.B., Conradt,
J., and Hartmann, M.J. (2007). Biomechanical models for radial distance determination by the rat
vibrissal system. *J. Neurophysiol.* **98**, 2439–2455.

Bresee, C.S., Belli, H.M., Luo, Y., and Hartmann, M.J.Z. (2023). Comparative morphology of the
whiskers and faces of mice (*Mus musculus*) and rats (*Rattus norvegicus*). *J. Exp. Biol.* **226**,
jeb245597.

Bufacchi, R.J., and Iannetti, G.D. (2018). An action field theory of peripersonal space. *Trends
Cogn. Sci.* **22**, 1076–1090.

Bush, N.E., Schroeder, C.L., Hobbs, J.A., Yang, A.E., Huet, L.A., Solla, S.A., and Hartmann, M.J.
(2016). Decoupling kinematics and mechanics reveals coding properties of trigeminal ganglion
neurons in the rat vibrissal system. *eLife* **5**, e13969.

Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and
currents—EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* **13**, 407–420.

Campagner, D., Evans, M.H., Bale, M.R., Erskine, A., and Petersen, R.S. (2016). Prediction of
primary somatosensory neuron activity during active tactile exploration. *eLife* **5**, e10696.

Chakrabarti, S., and Schwarz, C. (2018). Cortical modulation of sensory flow during active touch
in the rat whisker system. *Nat. Commun.* **9**, 3907.

Clack, N.G., O'Connor, D.H., Huber, D., Petreanu, L., Hires, A., Peron, S., Svoboda, K., and
Myers, E.W. (2012). Automated tracking of whiskers in videos of head-fixed rodents. *PLoS
Comput. Biol.* **8**, e1002591.

Diamond, M.E., von Heimendahl, M., Knutsen, P.M., Kleinfeld, D., and Ahissar, E. (2008). "Where"
and "what" in the whisker sensorimotor system. *Nat. Rev. Neurosci.* **9**, 601–612.

Ebara, S., Kumamoto, K., Matsuura, T., Mazurkiewicz, J.E., and Rice, F.L. (2002). Similarities and
differences in the innervation of mystacial vibrissal follicle-sinus complexes in the rat and cat: a
confocal microscopic study. *J. Comp. Neurol.* **449**, 103–119.

Edin, B.B., and Johansson, N. (1995). Skin strain patterns provide kinaesthetic information to the
human central nervous system. *J. Physiol.* **487**, 243–251.

Fundin, B.T., Rice, F.L., Pfaller, K., and Arvidsson, J. (1994). The innervation of the mystacial pad
in the adult rat studied by anterograde transport of HRP conjugates. *Exp. Brain Res.* **99**, 233–
246.

Furuta, T., Bush, N.E., Yang, A.E., Ebara, S., Miyazaki, N., Murata, K., Hirai, D., Shibata, K.I., and
Hartmann, M.J.Z. (2020). The cellular and mechanical basis for response characteristics of
identified primary afferents in the rat vibrissal system. *Curr. Biol.* **30**, 815–826.e5.

Furuta, T., Timofeeva, E., Nakamura, K., Okamoto-Furuta, K., Togo, M., Kaneko, T., and
Deschênes, M. (2008). Inhibitory gating of vibrissal inputs in the brainstem. *J. Neurosci.* **28**, 1789–
1797.

Gentilucci, M., Scandolara, C., Pigarev, I.N., and Rizzolatti, G. (1983). Visual responses in the
postarcuate cortex (area 6) of the monkey that are independent of eye position. *Exp. Brain Res.*
50, 464–468.

Georgopoulos, A.P., Schwartz, A.B., and Kettner, R.E. (1986). Neuronal population coding of
movement direction. *Science* **233**, 1416–1419.

Geramita, M.A., Burton, S.D., and Urban, N.N. (2016). Distinct lateral inhibitory circuits drive
parallel processing of sensory information in the mammalian olfactory bulb. *eLife* **5**, e16039.

Giridhar, S., Doiron, B., and Urban, N.N. (2011). Timescale-dependent shaping of correlation by
olfactory bulb lateral inhibition. *Proc. Natl. Acad. Sci. USA* **108**, 5843–5848.

Govorunova, E.G., Sineshchekov, O.A., Janz, R., Liu, X., and Spudich, J.L. (2015). Natural light-
gated anion channels: a family of microbial rhodopsins for advanced optogenetics. *Science* **349**,
647–650.

Grant, R.A., Breakell, V., and Prescott, T.J. (2018). Whisker touch sensing guides locomotion in
small, quadrupedal mammals. *Proc. Biol. Sci.* **285**, 20180592.

Grigg, P. (1996). Stretch sensitivity of mechanoreceptor neurons in rat hairy skin. *J. Neurophysiol.*
76, 2886–2895.

Hao, J., and Delmas, P. (2010). Multiple desensitization mechanisms of mechanotransducer channels shape firing of mechanosensory neurons. *J. Neurosci.* **30**, 13384–13395.

Harris, J.A., Petersen, R.S., and Diamond, M.E. (1999). Distribution of tactile learning and its neural basis. *Proc. Natl. Acad. Sci. USA* **96**, 7587–7591.

Heimburg, F., Saluti, N.M., Timm, J., Adlakha, A., Bortolozzo-Gleich, M.H., Martín-Cortecero, J., Castelanelli, M., Klumpp, M., Embray, L., et al. (2025). A tactile discrimination task to study neuronal dynamics in freely-moving mice. *Nat. Commun*. **16**, 6421.

Henze, D.A., Borhegyi, Z., Csicsvari, J., Mamiya, A., Harris, K.D., and Buzsáki, G. (2000). Intracellular features predicted by extracellular recordings in the hippocampus *in vivo*. *J. Neurophysiol.* **84**, 390–400.

Hubel, D.H., and Wiesel, T.N. (1962). Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol.* **160**, 106–154.

Hutson, K.A., and Masterton, R.B. (1986). The sensory contribution of a single vibrissa's cortical barrel. *J. Neurophysiol.* **56**, 1196–1223.

Jacquin, M.F., Arends, J.J., Renehan, W.E., Waite, P.M., and Shortland, P.J. (2015). Whisker-related circuitry in the trigeminal nucleus principalis: topographic precision. *Somatosens. Mot. Res.* **32**, 8–20.

Jacquin, M.F., Barcia, M., and Rhoades, R.W. (1989). Structure–function relationships in rat brainstem subnucleus interpolaris: IV. Projection neurons. *J. Comp. Neurol.* **282**, 45–62.

Jazayeri, M., and Movshon, J.A. (2006). Optimal representation of sensory information by neural populations. *Nat. Neurosci.* **9**, 690–696.

Jezzini, A., Mazzucato, L., La Camera, G., and Fontanini, A. (2013). Processing of hedonic and chemosensory features of taste in medial prefrontal and insular networks. *J. Neurosci.* **33**, 18966–18978.

Khatri, V., Bermejo, R., Brumberg, J.C., Keller, A., and Zeigler, H.P. (2009). Whisking in air: encoding of kinematics by trigeminal ganglion neurons in awake rats. *J. Neurophysiol.* **101**, 1836–1846.

Kleinfeld, D., and Deschênes, M. (2011). Neuronal basis for object location in the vibrissa scanning sensorimotor system. *Neuron* **72**, 455–468.

Krupa, D.J., Matell, M.S., Brisben, A.J., Oliveira, L.M., and Nicolelis, M.A. (2001). Behavioral properties of the trigeminal somatosensory system in rats performing whisker-dependent tactile discriminations. *J. Neurosci.* **21**, 5752–5763.

Laughlin, S.B. (2001). Energy as a constraint on the coding and processing of sensory information. *Curr. Opin. Neurobiol.* **11**, 475–480.

Lee, S.-H., Kwan, A.C., Zhang, S., Phoumthipphavong, V., Flannery, J.G., Masmanidis, S.C.,
Taniguchi, H., Huang, Z.J., Zhang, F., Boyden, E.S., **et al.** (2012). Activation of specific
interneurons improves V1 feature selectivity and visual perception. *Nature* **488**, 379–383.

Li, L., Rutlin, M., Abraira, V.E., Cassidy, C., Kus, L., Gong, S., Jankowski, M.P., Luo, W., Heintz,
N.**,** et al. **(**2011). The functional organization of cutaneous low-threshold mechanosensory
neurons. *Cell* **147**, 1615–1627.

Link, E., Edelmann, L., Chou, J.H., Binz, T., Yamasaki, S., Eisel, U., Baumert, M., Südhof, T.C.,
Niemann, H., and Jahn, R. (1992). Tetanus toxin action: inhibition of neurotransmitter release
linked to synaptobrevin proteolysis. *Biochem. Biophys. Res. Commun.* **189**, 1017–1023.

Meyers, E.M. (2013). The neural decoding toolbox. *Front. Neuroinform.* **7**, 8.

Minnery, B.S., Bruno, R.M., and Simons, D.J. (2003). Response transformation and receptive-
field synthesis in the lemniscal trigeminothalamic circuit. *J. Neurophysiol.* **90**, 1556–1570.

Minnery, B.S., and Simons, D.J. (2003). Response properties of whisker-associated
trigeminothalamic neurons in rat nucleus principalis. *J. Neurophysiol.* **89**, 40–56.

Mitchinson, B., Gurney, K.N., Redgrave, P., Melhuish, C., Pipe, A.G., Pearson, M., Gilhespy, I.,
and Prescott, T.J. (2004). Empirically inspired simulated electro-mechanical model of the rat
mystacial follicle-sinus complex. *Proc. Biol. Sci.* **271**, 2509–2516.

Mitchinson, B., and Prescott, T.J. (2013). Whisker movements reveal spatial attention: a unified
computational model of active sensing control in the rat. *PLoS Comput. Biol.* **9**, e1003236.

Mohar, B., Katz, Y., and Lampl, I. (2013). Opposite adaptive processing of stimulus intensity in
two major nuclei of the somatosensory brainstem. *J. Neurosci.* **33**, 15394–15400.

Moore, J.D., Mercer Lindsay, N., Deschênes, M., and Kleinfeld, D. (2015). Vibrissa self-motion
and touch are reliably encoded along the same somatosensory pathway from brainstem through
thalamus. *PLoS Biol.* **13**, e1002253.

Nordin, M., and Hagbarth, K.E. (1989). Mechanoreceptive units in the human infra-orbital nerve.
Acta Physiol. Scand. **135**, 149–161.

O'Connor, D.H., Peron, S.P., Huber, D., and Svoboda, K. (2010). Neural activity in barrel cortex
underlying vibrissa-based object localization in mice. *Neuron* **67**, 1048–1061.

O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence
from unit activity in the freely moving rat. *Brain Res.* **34**, 171–175.

Olshausen, B.A., and Field, D.J. (2004). Sparse coding of sensory inputs. *Curr. Opin. Neurobiol.*
14, 481–487.

Pachitariu, M., Steinmetz, N.A., Kadir, S.N., Carandini, M., and Harris, K.D. (2016). Fast and
accurate spike sorting of high-channel count probes with KiloSort. *Adv. Neural Inf. Process. Syst.*
29.

Pammer, L., O'Connor, D.H., Hires, S.A., Clack, N.G., Huber, D., Myers, E.W., and Svoboda, K. (2013). The mechanical variables underlying object localization along the axis of the whisker. *J. Neurosci.* **33**, 6726–6741.

Porter, K.K., and Groh, J.M. (2006). The "other" transformation required for visual–auditory integration: representational format. *Prog. Brain Res.* **155**, 313–323.

Pouget, A., Dayan, P., and Zemel, R. (2000). Information processing with population codes. *Nat. Rev. Neurosci.* **1**, 125–132.

Reynders, A., Mantilleri, A., Malapert, P., Rialle, S., Nidelet, S., Laffray, S., Beurrier, C., Bourinet, E., and Moqrich, A. (2015). Transcriptional profiling of cutaneous MRGPRD free nerve endings and C-LTMRs. *Cell Rep.* **10**, 1007–1019.

Rice, F.L., Fundin, B.T., Arvidsson, J., Aldskogius, H., and Johansson, O. (1997). Comprehensive immunofluorescence and lectin binding analysis of vibrissal follicle-sinus complex innervation in the mystacial pad of the rat. *J. Comp. Neurol.* **385**, 149–184.

Rice, F.L., Mance, A., and Munger, B.L. (1986). A comparative light microscopic analysis of the sensory innervation of the mystacial pad. I. Innervation of vibrissal follicle-sinus complexes. *J. Comp. Neurol.* **252**, 154–174.

Rizzolatti, G., and Gentilucci, M. (1988). Motor and visual-motor functions of the premotor cortex. In *Neurobiology of Neocortex*, P. Rakic and W. Singer, eds. (Chichester, UK: John Wiley & Sons), pp. 269–284.

Rizzolatti, G., Scandolara, C., Matelli, M., and Gentilucci, M. (1981). Afferent properties of periarcuate neurons in macaque monkeys. II. Visual responses. *Behav. Brain Res.* **2**, 147–163.

Rossant, C. (n.d.). Phy: spike sorting software for extracellular electrophysiology. GitHub repository. Available at: https://github.com/cortex-lab/phy.

Roudaut, Y., Lonigro, A., Coste, B., Hao, J., Delmas, P., and Crest, M. (2012). Touch sense: functional organization and molecular determinants of mechanosensitive receptors. *Channels (Austin)* **6**, 234–245.

Schiavo, G., Benfenati, F., Poulain, B., Rossetto, O., Polverino de Laureto, P., DasGupta, B.R., and Montecucco, C. (1992). Tetanus and botulinum-B neurotoxins block neurotransmitter release by proteolytic cleavage of synaptobrevin. *Nature* **359**, 832–835.

Schneider-Mizell, C.M., Bodor, A.L., Brittain, D., Buchanan, J., Bumbarger, D.J., Elabbady, L., Gamlin, C., Kapner, D., Kinn, S., Mahalingam, G., et al. (2025). Inhibitory specificity from a connectomic census of mouse visual cortex. *Nature* **640**, 448–458.

Sakurai, K., Akiyama, M., Cai, B., Scott, A., Han, B.X., Takatoh, J., Sigrist, M., Arber, S., and Wang, F. (2013). The organization of submodality-specific touch afferent inputs in the vibrissa column. *Cell Rep.* **5**, 87–98.

Serino, A. (2019). Peripersonal space (PPS) as a multisensory interface between the individual
and the environment, defining the space of the self. *Neurosci. Biobehav. Rev.* **99**, 138–159.

Serino, A., Annella, L., and Avenanti, A. (2009). Motor properties of peripersonal space in humans.
PLoS One **4**, e6582.

Severson, K.S., Xu, D., Van de Loo, M., Bai, L., Ginty, D.D., and O'Connor, D.H. (2017). Active
touch and self-motion encoding by Merkel cell-associated afferents. *Neuron* **94**, 666–676.

Severson, K.S., Xu, D., Yang, H., and O'Connor, D.H. (2019). Coding of whisker motion across
the mouse face. *eLife* **8**, e41535.

Shipley, M.T. (1974). Response characteristics of single units in the rat's trigeminal nuclei to
vibrissa displacements. *J. Neurophysiol.* **37**, 73–90.

Sofroniew, N.J., Cohen, J.D., Lee, A.K., and Svoboda, K. (2014). Natural whisker-guided behavior
by head-fixed mice in tactile virtual reality. *J. Neurosci.* **34**, 9537–9550.

Sofroniew, N.J., Vlasov, Y.A., Hires, S.A., Freeman, J., and Svoboda, K. (2015). Neural coding in
barrel cortex during whisker-guided locomotion. *eLife* **4**, e12559.

Solomon, J.H., and Hartmann, M.J. (2011). Radial distance determination in the rat vibrissal
system and the effects of Weber's law. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **366**, 3049–3057.

Storey, J.D. (2002). A direct approach to false discovery rates. *J. R. Stat. Soc. B* **64**, 479–498.

Szwed, M., Bagdasarian, K., Blumenfeld, B., Barak, O., Derdikman, D., and Ahissar, E. (2006).
Responses of trigeminal ganglion neurons to the radial distance of contact during active vibrissal
touch. *J. Neurophysiol.* **95**, 791–802.

Takatoh, J., Prevosto, V., Thompson, P.M., Lu, J., Chung, L., Harrahill, A., Li, S., Zhao, S., He, Z.,
Golomb, D., et al. (2022). The whisking oscillator circuit. *Nature* **609**, 560–568.

Timofeeva, E., Dufresne, C., Sík, A., Zhang, Z.W., and Deschênes, M. (2005). Cholinergic
modulation of vibrissal receptive fields in trigeminal nuclei. *J. Neurosci.* **25**, 9135–9143.

Timofeeva, E., Lavallée, P., Arsenault, D., and Deschênes, M. (2004). Synthesis of multiwhisker
receptive fields in subcortical stations of the vibrissa system. *J. Neurophysiol.* **91**, 1510–1515.

Veinante, P., and Deschênes, M. (1999). Single- and multi-whisker channels in the ascending
projections from the principal trigeminal nucleus in the rat. *J. Neurosci.* **19**, 5085–5095.

Vincis, R., Chen, K., Czarnecki, L., Chen, J., and Fontanini, A. (2020). Dynamic representation of
taste-related decisions in the gustatory insular cortex of mice. *Curr. Biol.* **30**, 1834–1844.e5.

Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., Hjerling-Leffler, J.,
Haeggström, J., Kharchenko, O., Kharchenko, P.V., et al. (2015). Unbiased classification of
sensory neuron types by large-scale single-cell RNA sequencing. *Nat. Neurosci.* **18**, 145–153.

Warren, R.A., Zhang, Q., Hoffman, J.R., Li, E.Y., Hong, Y.K., Bruno, R.M., and Sawtell, N.B.
(2021). A rapid whisker-based decision underlying skilled locomotion in mice. *eLife* **10**, e63596.

Zhang, Y., Zhao, S., Rodriguez, E., Takatoh, J., Han, B.X., Zhou, X., and Wang, F. (2015).
Identifying local and descending inputs for primary sensory neurons. *J. Clin. Invest.* **125**, 3782–
3794.

Znamenskiy, P., Kim, M.H., Muir, D.R., Iacaruso, M.F., Hofer, S.B., and Mrsic-Flogel, T.D. (2024).
Functional specificity of recurrent inhibition in visual cortex. *Neuron* **112**, 991–1000.e8.

Table S1. Mouse Metadata and Figure Appearances.

Mouse ID	Sex	Mouse Line	Birth Date	Recording Dates	Experiments	Appearances - Main Figures					
						1	2	3	4	5	6
Ephys1	M	C57BL/6J	11/08/22	02/15/23-02/17/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys2	M	C57BL/6J	11/08/22	02/08/23-02/10/23	PrV Si probe, WT	E	A-F	B-D,F		C-E	
Ephys3	M	C57BL/6J	09/20/22	02/01/23-02/03/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys5	M	C57BL/6J	06/22/21	08/19/21-08/24/21	PrV Tungsten						
Ephys6	M	C57BL/6J	06/22/21	09/17/21-09/20/21	PrV Tungsten						
Ephys7	M	C57BL/6J	06/22/21	09/13/21-09/14/21	PrV Tungsten						
Ephys8	F	C57BL/6J	01/25/22	06/15/22	PrV Tungsten						
Ephys9	M	C57BL/6J	08/24/21	10/15/21-10/19/21	PrV Tungsten						
Ephys12	M	C57BL/6J	05/14/22	08/08/22	PrV Tungsten						
Ephys13	F	C57BL/6J	04/12/22	06/29/22-07/06/22	PrV Tungsten						
Ephys14	F	C57BL/6J	04/12/22	07/27/22-08/04/22	PrV Tungsten						
Ephys15	M	C57BL/6J	05/14/22	08/11/22-08/25/22	PrV Tungsten						
Ephys16	M	C57BL/6J	05/14/22	09/01/22-09/06/22	PrV Tungsten						
Ephys17	M	C57BL/6J	05/14/22	09/09/22	PrV Tungsten						
Ephys18	M	C57BL/6J	07/12/22	09/29/22-10/04/22	PrV Tungsten						
Ephys19	M	C57BL/6J	07/12/22	10/17/22	PrV Tungsten						
Ephys20	M	C57BL/6J	07/12/22	10/27/22-11/01/22	PrV Tungsten						
Ephys24	F	C57BL/6J	10/25/22	01/31/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys25	M	C57BL/6J	11/08/22	02/22/23-02/25/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys26	M	C57BL/6J	11/08/22	02/26/23-02/28/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys27	F	C57BL/6J	11/15/22	03/21/23-03/23/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys28	F	C57BL/6J	11/15/22	03/24/23-03/27/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys29	F	C57BL/6J	11/29/22	03/28/23-03/30/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys30	M	C57BL/6J	02/27/23	05/24/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys32	F	C57BL/6J	03/14/23	06/15/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys33	F	C57BL/6J	03/14/23	06/02/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys34	M	C57BL/6J	02/27/23	07/11/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys36	F	C57BL/6J	03/14/23	06/08/23	PrV Si probe, WT	D	D-F	B-D,F		C-E	
Ephys40	M	C57BL/6J	03/28/23	06/26/23	PrV Si probe, WT		A-F	B-D,F		C-E	
Ephys41	M	C57BL/6J	03/28/23	06/27/23	PrV Si probe, WT		D-F	B-D,F		C-E	
Ephys42	F	C57BL/6J	05/23/23	07/25/23	PrV Si probe, WT		D-F	B-D,F		C-E	
beh104	M	C57BL/6J	06/06/23	12/04/23	whisker behavior						
beh101	F	C57BL/6J	09/26/23	12/12/23	whisker behavior						
beh102	M	C57BL/6J	09/26/23	12/13/23	whisker behavior						
beh103	F	C57BL/6J	08/15/23	12/12/23	whisker behavior						
Ephys72	M	vGAT-ires-Cre	11/27/23	03/07/23-03/08/23	PrV optrode local GtACR						
Ephys73	F	vGAT-ires-Cre	11/27/23	03/11/23-03/12/23	PrV optrode local GtACR						
Ephys75	M	C57BL/6J	04/02/24	05/22/24	PrV Si probe, WT + 1WC	C	D-F	B-D,F	B,K	C-E	
Ephys76	F	C57BL/6J	04/02/24	06/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B-F,K	C-E	
Ephys77	F	C57BL/6J	04/22/24	08/14/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys78	F	C57BL/6J	04/02/24	07/02/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys79	M	C57BL/6J	04/02/24	06/24/24	PrV Si probe, WT + 1WC		D-F	B-D,F	A-D,K	C-E	
Ephys81	F	C57BL/6J	04/02/24	07/03/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,G,H,K	C-E	
Ephys83	M	vGAT-ires-Cre	03/09/24	07/09/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys85	M	vGAT-ires-Cre	03/09/24	07/11/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys87	F	C57BL/6J	03/12/24	07/24/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E	
Ephys88	M	C57BL/6J	05/14/24	09/05/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,I,J,K	C-E	
Ephys89	F	C57BL/6J	04/22/24	07/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys92	F	C57BL/6J	04/22/24	08/15/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys94	M	vGAT-ires-Cre	05/22/24	08/17/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys96	F	vGAT-ires-Cre	05/18/24	08/20/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys97	F	vGAT-ires-Cre	03/01/24	11/14/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys100	M	vGAT-ires-Cre	06/20/24	09/08/24	PrV Si probe, TeLC in SpVi					C-E	
Ephys101	M	C57BL/6J	05/14/24	09/09/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E	
Ephys102	F	vGAT-ires-Cre	06/20/24	09/12/24	PrV Si probe, TeLC in SpVi					C-E	

Ephys105	M	*C57BL/6J*	07/02/24	09/23/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E
Ephys106	M	*C57BL/6J*	07/02/24	09/24/24	PrV Si probe, WT + 1WC		A-F	B-D,F	B,I,J,K	C-E
Ephys107	M	*C57BL/6J*	07/02/24	09/25/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E
Ephys108	M	*C57BL/6J*	07/02/24	09/26/24	PrV Si probe, WT + 1WC		D-F	B-D,F	B,E,F,K	C-E
Ephys114	F	*vGAT-ires-Cre*	05/22/24	11/18/24	PrV Si probe, TeLC in SpVi					C-E
Ephys119	M	*vGAT-ires-Cre*	08/20/24	12/08/24-12/09/24	PrV Si probe, TeLC in SpVi					B-E
Ephys121	F	*vGAT-ires-Cre*	08/20/24	12/14/24-12/15/24	PrV Si probe, TeLC in SpVi					C-E
Ephys122	M	*vGAT-ires-Cre*	09/09/24	12/16/24-12/17/24	PrV Si probe, TeLC in SpVi					C-E
Ephys126	M	*C57BL/6J*	12/03/24	03/20/25-03/21/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B	C-E
Ephys127	F	*C57BL/6J*	11/03/24	03/24/25-03/25/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E
Ephys128	F	*C57BL/6J*	12/31/24	03/26/25-03/27/25	PrV Si probe, WT + 1WC		D-F	B-D,F	B,K	C-E

Mouse ID	Appearances - Supplemental Figures						Other Notes
	S1	S2	S3	S4	S5	S6	
Ephys1			B-E	A-D	B,C	H,I	
Ephys2			A-E	A-E	A-C	H,I	Unit 5 in 20230208_s1 is 1st example in Fig. 2A-C, S3A, S4E, S5A
Ephys3			B-E	A-D	B,C	H,I	
Ephys5		F	J-M				
Ephys6		F	J-M				
Ephys7		F	J-M				
Ephys8		B,F	J-M				
Ephys9		C-F	J-M				
Ephys12		F	J-M				
Ephys13		D-F	J-M				
Ephys14		F	J-M				
Ephys15		F	J-M				
Ephys16		F	J-M				
Ephys17		F	J-M				
Ephys18		F	J-M				
Ephys19		F	J-M				
Ephys20		F	J-M				
Ephys24			B-E	A,C	B,C	H,I	
Ephys25			B-E	A,C	B,C	H,I	
Ephys26			B-E	A-D	B,C	H,I	
Ephys27			B-E	A-D	B,C	H,I	
Ephys28			B-E	A,C	B,C	H,I	
Ephys29			B-E	A,C	B,C	H,I	
Ephys30			B-E	A,C	B,C	H,I	
Ephys32			B-E	A-D	B,C	H,I	
Ephys33			B-E	A-D	B,C	H,I	
Ephys34			B-E	A,C	B,C	H,I	
Ephys36			B-E	A,C	B,C	H,I	
Ephys40			A-E	A,C	B,C	H,I	Unit 8 in 20230626_s1 is 2nd example in Fig. 2A-C, S3A
Ephys41			B-E	A-D	B,C	H,I	
Ephys42			B-E	A,C	B,C	H,I	
beh104	C,F						
beh101	F						
beh102	D-F						
beh103	A,B,F						
Ephys72						D,E	Unit 2 in 20240307_s1 is 1st example in Fig. S6E
Ephys73						C,E	Unit 14 in 20240311_s1 is 2nd example in Fig. S6E
Ephys75			B-E	A-D	B,C	H,I	
Ephys76			B-E	A,C	B,C	H,I	
Ephys77			B-E	A,C	B,C	H,I	
Ephys78			B-E	A,C	B,C	H,I	
Ephys79			B-E	A-D	B,C	H,I	
Ephys81			A-E	A,C	B,C	H,I	Unit 10 in 20240703_s1 is 4th example in Fig. 2A-C
Ephys83			F-I			H,I	
Ephys85			F-I			H,I	
Ephys87			B-E	A,C	B,C	H,I	
Ephys88			A-E	A,C	A-C	H,I	Unit 5 in 20240905_s1 is 3rd example in Fig. 2A-C, S3A; 2nd in S5A
Ephys89			B-E	A,C	B,C	H,I	
Ephys92			B-E	A-D	B,C	H,I	
Ephys94			F-I			H,I	
Ephys96			F-I			H,I	
Ephys97			F-I			H,I	
Ephys100			F-I			H,I	
Ephys101			B-E	A-D	B,C	H,I	
Ephys102			F-I		B,C	H,I	

Ephys105			B-E	A-E	B,C	H,I	
Ephys106			A-E	A-D	B,C	H,I	Unit 7 in 20240924_s1 is 6th example in Fig. 1A-C, S3A
Ephys107			B-E	A,C,E	B,C	H,I	Unit 1 in 20240925_1 is 2nd example in Fig. S4E
Ephys108			B-E	A,C	B,C	H,I	
Ephys114			F-I			H,I	
Ephys119			F-I			G-I	
Ephys121			F-I			H,I	
Ephys122			F-I			H,I	
Ephys126			B-E	A-D	B,C	H,I	
Ephys127			B-E	A-D	B,C	H,I	
Ephys128			B-E	A-D	B,C	H,I	

Table S2. Population Decoder Accuracy

Distance (mm)	Decoder Accuracy Mean ± SEM (%)				P-Value
	Proximity	Map	Untuned	Shuffled	Proximity vs Map
7	87.84 ± 1.210	81.14 ± 0.239	32.30 ± 1.767	11.17 ± 0.979	2.06×10^{-4}, *
9	78.87 ± 1.484	74.68 ± 0.330	15.89 ± 0.946	10.85 ± 0.738	3.30×10^{-4}, *
11	64.14 ± 2.238	82.57 ± 0.237	13.41 ± 1.045	11.42 ± 1.016	2.06×10^{-4}, *
13	50.60 ± 3.327	83.90 ± 0.213	13.04 ± 0.567	10.62 ± 1.008	2.06×10^{-4}, *
15	36.29 ± 1.824	82.50 ± 0.148	12.27 ± 0.789	10.78 ± 0.669	2.06×10^{-4}, *
17	30.73 ± 1.372	74.29 ± 0.377	12.01 ± 0.554	10.95 ± 0.690	2.06×10^{-4}, *
19	28.37 ± 1.284	63.16 ± 0.408	11.81 ± 0.384	10.69 ± 0.954	2.06×10^{-4}, *
21	28.91 ± 1.108	49.38 ± 0.439	11.63 ± 0.314	10.54 ± 0.828	2.06×10^{-4}, *
23	34.01 ± 0.670	52.12 ± 0.344	12.18 ± 1.011	11.09 ± 1.111	2.06×10^{-4}, *

*: significant ($p < 0.05$); n.s.: not significant

Table S3. Population Decoder Error

| Distance (mm) | Decoder Accuracy Mean ± SEM (mm) | | | | P-Value |
	Proximity	Map	Untuned	Shuffled	Proximity vs Map
7	0.33 ± 0.034	0.50 ± 0.012	4.90 ± 0.196	8.03 ± 0.137	2.00×10^{-4}, *
9	0.68 ± 0.076	0.51 ± 0.011	5.33 ± 0.122	6.45 ± 0.157	1.30×10^{-3}, *
11	1.34 ± 0.138	0.46 ± 0.011	4.96 ± 0.090	5.30 ± 0.087	2.00×10^{-4}, *
13	2.04 ± 0.107	0.70 ± 0.012	4.50 ± 0.063	4.74 ± 0.077	2.00×10^{-4}, *
15	2.50 ± 0.084	0.78 ± 0.011	4.24 ± 0.052	4.46 ± 0.066	2.00×10^{-4}, *
17	2.50 ± 0.040	1.02 ± 0.012	4.28 ± 0.056	4.70 ± 0.097	2.00×10^{-4}, *
19	2.25 ± 0.039	1.25 ± 0.017	4.81 ± 0.069	5.36 ± 0.122	2.00×10^{-4}, *
21	2.06 ± 0.108	1.20 ± 0.008	5.71 ± 0.081	6.54 ± 0.085	2.00×10^{-4}, *
23	2.32 ± 0.111	1.33 ± 0.012	7.21 ± 0.165	8.03 ± 0.156	2.00×10^{-4}, *

*: significant ($p < 0.05$); n.s.: not significant

Table S4. Single Neuron Decoder Accuracy

| Distance (mm) | Decoder Accuracy Mean ± SEM (%) | | P-Value |
	Proximity	Map	Proximity vs Map
7	65.74 ± 1.319	59.09 ± 1.889	0.02, *
9	40.83 ± 1.662	39.26 ± 2.253	0.73, n.s.
11	31.44 ± 1.447	37.95 ± 2.203	0.02, *
13	24.50 ± 1.312	37.34 ± 2.387	0, *
15	18.75 ± 1.099	34.72 ± 2.277	0, *
17	20.27 ± 1.195	27.63 ± 1.886	0, *
19	20.68 ± 1.143	23.54 ± 1.824	0.15, n.s.
21	20.69 ± 1.216	23.68 ± 1.722	0.05, n.s.
23	26.44 ± 1.505	31.55 ± 2.313	0.03, *

*: significant ($p < 0.05$); n.s.: not significant

Table S5. Single Neuron Decoder Error

Distance (mm)	Decoder Accuracy Mean ± SEM (mm)		P-Value
	Proximity	Map	Proximity vs Map
7	2.11 ± 0.115	2.94 ± 0.198	0, *
9	3.34 ± 0.144	2.78 ± 0.166	0.21, n.s.
11	3.89 ± 0.127	2.89 ± 0.157	0, *
13	4.02 ± 0.101	2.85 ± 0.145	0, *
15	3.76 ± 0.076	3.02 ± 0.146	0, *
17	3.42 ± 0.072	3.32 ± 0.141	0.21, n.s.
19	3.31 ± 0.077	3.45 ± 0.160	0.84, n.s.
21	3.71 ± 0.103	3.76 ± 0.225	0.32, n.s.
23	4.63 ± 0.143	4.58 ± 0.274	0.46, n.s.

*: significant ($p < 0.05$); n.s.: not significant

Table S6. Statistics

Fig.	Comparison	Groups Tested	Measure (Mean ± SEM)	Statistical Test	Test Statistic	P-Value	Significance	DoF
3D	Population decoding accuracy	Proximity vs Map	see Table S2	Wilcoxon rank-sum test	N/A	see Table S2	see Table S2	N/A
3F	Population decoding error	Proximity vs Map	see Table S3	Wilcoxon rank-sum test	N/A	see Table S3	see Table S3	N/A
4B	Firing rates during wall passes	Full array vs. One whisker	12.37 ± 2.986 vs. 8.61 ± 2.487 sp/s	Wilcoxon signed-rank	N/A	1.27×10^{-9}	*	N/A
5D	Preferred distance CDFs	Wild-type vs. TeLC populations	N/A	One-tailed K-S	N/A	8.02×10^{-5}	*	N/A
5E	Proportion of map units	Wild-type vs. TeLC populations	10.34% vs. 3.90%	One-tailed Z-test	Z = 4.64	1.80×10^{-6}	*	N/A
5E	Proportion of proximity units	Wild-type vs. TeLC populations	23.31% vs. 32.09%	One-tailed Z-test	Z = -3.93	4.27×10^{-5}	*	N/A
5E	Proportion of untuned units	Wild-type vs. TeLC populations	48.20% vs. 49.92%	Two-tailed Z-test	Z = -0.68	0.5	n.s.	N/A
5E	Proportion of supressed units	Wild-type vs. TeLC populations	10.43% vs. 7.30%	One-tailed Z-test	Z = 2.12	0.02	*	N/A
5E	Proportion of ambiguous units	Wild-type vs. TeLC populations	7.71% vs. 6.79%	Two-tailed Z-test	Z = 0.69	0.49	n.s.	N/A
S1F, Left	C1 whisker midpoint	Awake vs. Anesthetized	-8.52 ± 0.633° vs. -5.88 ± 0.910°	Two-tailed paired t-test	t = -3.86	0.03	n.s. (α = 0.0167)	3
S1F, Middle	C2 whisker midpoint	Awake vs. Anesthetized	-6.81 ± 0.867° vs. -4.72 ± 0.927°	Two-tailed paired t-test	t = -3.57	0.04	n.s. (α = 0.0167)	3
S1F, Right	Greek whisker midpoint	Awake vs. Anesthetized	-4.76 ± 1.041° vs. -2.63 ± 0.714°	Two-tailed paired t-test	t = -4.55	0.0199	n.s. (α = 0.0167)	3
S4B, Left	Firing rates of wall-activated, distance-tuned units	Wall passes vs. Baseline	16.44 ± 1.640 vs. 6.36 ± 0.704 sp/s	One-tailed K-S	N/A	9.26×10^{-28}	*	N/A
S4B, Right	Firing rates of wall-suppressed, distance-tuned units	Wall passes vs. Baseline	4.47 ± 1.000 vs. 6.19 ± 1.076 sp/s	One-tailed K-S	N/A	3.06×10^{-4}	*	N/A
S4C	Untuned unit firing rates	Wall passes vs. Baseline	6.65 ± 0.687 vs. 6.90 ± 0.682 sp/s	Two-tailed K-S	N/A	0.74	n.s.	N/A
S4G, Middle	Wall-tuning category (ambiguous unit)	Non-closest vs. Closest distance	32.59 ± 4.253 vs. 26.53 ± 4.107 sp/s	One-tailed Wilcoxon signed-rank test	N/A	0.09	n.s.	N/A
S4G, Right	Wall-tuning category (map unit)	Non-closest vs. Closest distance	51.04 ± 2.49 vs. 26.09 ± 1.033 sp/s	One-tailed Wilcoxon signed-rank test	N/A	1.27×10^{-4}	*	N/A
S5B	Single-neuron decoder accuracy	Proximity vs Map	see Table S4	Wilcoxon rank-sum test	N/A	see Table S4	see Table S4	N/A
S5C	Single-neuron decoder error	Proximity vs Map	see Table S5	Wilcoxon rank-sum test	N/A	see Table S5	see Table S5	N/A
S6H, Left	Baseline firing rates (untuned units)	TeLC vs. Wild-type populations	9.34 ± 1.330 vs. 6.90 ± 0.682 sp/s	One-tailed K-S	N/A	7.63×10^{-5}	* (α = 0.0125)	N/A
S6H, Right	Wall-pass firing rates (untuned units)	TeLC vs. Wild-type populations	9.04 ± 1.390 vs. 6.65 ± 0.687 sp/s	One-tailed K-S	N/A	1.06×10^{-4}	* (α = 0.0125)	N/A
S6I, Left	Baseline firing rates (tuned units)	TeLC vs. Wild-type populations	5.60 ± 0.845 vs. 5.46 ± 0.670 sp/s	One-tailed K-S	N/A	0.14	n.s. (α = 0.0125)	N/A
S6I, Right	Wall-pass firing rates (tuned units)	TeLC vs. Wild-type populations	20.25 ± 2.263 vs. 16.44 ± 1.640 sp/s	One-tailed K-S	N/A	1.64×10^{-3}	* (α = 0.0125)	N/A
N/A	Firing rates during baseline	Fullarray vs. One whisker	7.08 ± 1.424 vs. 7.08 ± 1.418 sp/s	Wilcoxon signed-rank	N/A	0.59	n.s.	N/A