Disentangling Cephalopod Chromatophores Motor Units with Computer Vision

Mathieu D. M. Renard#[1], Johann Ukrow#[1], Margot Elmaleh[1], Dominic A. Evans[1],
Yifan Wu[2], Xitong Liang[2-5], Gilles Laurent*[1]

[1]Max Planck Institute for Brain Research, Frankfurt, Germany

[2]State Key Laboratory of Membrane Biology, School of Life Sciences, Peking University, Beijing 100871, China

[3]IDG/McGovern Institute for Brain Research, Peking University, Beijing 100871, China.

7
[4]Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China.

[5]Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China.

#joint first authors

*corresponding author

Abstract:

Cephalopod chromatophores are skin pigment organs that enable unmatched
camouflage through rapid, flexible and neurally controlled deformation. Although
their morphology is well known, the organization of their motor control is not entirely
understood. Here, we combine high-resolution videography with a dedicated
computer-vision pipeline (CHROMAS) to investigate chromatophore control and their
likely innervation in *Euprymna berryi* and *Sepia officinalis.* By segmenting
chromatophores into radial slices and analyzing anisotropic deformations, we applied
dimensionality reduction (PCA) and source separation (ICA) to estimate the number
and spatial influence of motor neurons responsible for the control of individual and
groups of chromatophores. On average, four independent components were
detected (suggesting innervation by four motor neurons), each forming contiguous
petal-shaped domains rather than causing uniform expansion. Clustering thousands
of components revealed motor units spanning multiple chromatophores, most
involving fewer than 14 but occasionally spanning more widely. These motor units
displayed a wide variety of geometries, ranging from compact local groups to
elongated or fragmented structures; they often overlapped, with repeated co-
innervation of chromatophore pairs occurring more often than expected by chance.
Expansion was consistently faster and more stereotyped than relaxation, consistent
with active contraction (corresponding to chromatophore expansion) and passive
recoil (chromatophore contraction). Together, these results show that individual
chromatophores are not singular or uniform pixels, but rather contrast elements that
can be fractionated into smaller territories, themselves coordinated with those of
other chromatophores. This geometry of neural control enables, among others, the
generation of "virtual" chromatophores (by convergence of adjacent territories of
neighbouring chromatophores) as well as that of noise in the distribution of pixel
shapes.

Introduction

Coleoids, a subgroup of the cephalopod taxon which includes octopuses, squids, and cuttlefish, possess one of the most sophisticated camouflage systems found in nature. This extraordinary ability is mediated by arrays of pigment cells, known as chromatophores, which expand and contract to alter skin coloration and statistical patterning in response to motor command from the brain. The collective behavior of thousands or millions of chromatophores thus creates adaptive patterns that enable the animals to blend with their surroundings, evade detection by predators or prey and communicate with conspecifics (Hanlon & Messenger, 1988).

Cephalopod chromatophores differ from pigment cells in other animals: each chromatophore is a neuromuscular organ composed of a central pigment cell containing a cytoelastic sacculus, rimmed by 15–25 slender radial muscle fibers, themselves attached to an extracellular semi-stiff fiber mesh at their distal end (Messenger, 2001). The pigment granules residing in the elastic sacculus thus spread into a thin disc when the radial muscles contract due to motorneuron excitation, expanding the pigmented area. When the motorneuron input ceases, the chromatophore muscles relax, the elasticity of the sacculus causes it to recoil (and the cytoplasmic membrane to invaginate and fold onto itself), reducing visible coloration (or the size of this biological, expandable pixel). These muscle-driven, active expansion and elastic, passive retraction enable very rapid patterning and color changes, seen in no other biological coloration system. In addition to their unique neuromuscular structure, chromatophores occur in distinct color classes (typically yellow, red, brown, and black), each defined by the pigment (e.g, xanthommatin for yellow, Reiter et al., 2018) and density of pigment granules they contain. These classes are linked to the age of the chromatophore (starting yellow and darkening over several weeks, refs), and usually migrate in the thickness of the skin, thus forming superimposed chromatic layers. Together they broaden the chromatic range available for patterning.

The basic neural circuit for camouflage patterning is well established: motor commands flow from visual processing centers (optic lobes) through basal lobes to the chromatophore lobes, where the somata of the motor neurons that directly

innervate the muscles of the skin chromatophores reside (Boycott, 1961; Messenger,
2001). What remains unclear is how the axonal terminals of these motor neurons
map onto individuals and groups of chromatophores. Anatomical and physiological
studies from Florey and colleagues (Cloney & Florey, 1968; Florey, 1969), as well as
Dubas and Boyle (1985), provided evidence for polyneuronal innervation, with
individual muscles or chromatophores under the control of several motor neurons.
However, the number of inputs, their spatial organization, and the logic of motor unit
formation remain unresolved.

This uncertainty limits our understanding of how the distributed neural activity that
controls   camouflage   generates   the   details   of   body   patterns.   Treating
chromatophores as uniform "pixels" ignores the possibility that subregions within
each chromatophore could act as semi-independent effectors. Likewise, the degree
to which chromatophores are coordinated in groups—forming motor units that span
several chromatophores—has only been estimated qualitatively (Florey, 1969;
Dubas & Boyle, 1985). These early anatomical and physiological studies suggested
that individual motor neurons may innervate dozens to hundreds of chromatophores,
but such estimates were indirect and limited in scope. More recently, computational
approaches have begun to address this issue (Reiter et al., 2018), but these
analyses considered chromatophores as indivisible units and provided only indirect
evidence of motor-unit structure. No systematic, quantitative framework yet exists to
describe the number, geometry, and overlap of chromatophore motor units on a
large scale. We addressed this gap using high-resolution tracking of chromatophore
dynamics combined with computational tools capable of disentangling overlapping
motor inputs.

Results

Methodological development with Euprymna berryi

The hummingbird bobtail squid *E. berryi* was used for initial experimental configuration and setup. This species was chosen for its compact size, large chromatophores, and rapid growth. Video recordings (10-60 min durations) were taken from 14 animals of 1-60 days of age (see Methods). In Fig 1, we analyzed a 30-s video dataset (1080 × 1080 px, 20 frames per second or fps) from a sedated hatchling that exhibited spontaneous chromatophore activity (Fig. 1a). Mild and "noisy" spontaneous activity was critical to enable measurements of motion correlation between chromatophores. Chromatophores were segmented, each divided in polar (radial) slices and the areal variations of each slice across all successive time frames were measured, as developed and described in Ukrow & Renard (2025). Chromatophore slices were then clustered using HDBSCAN (Campello et al., 2013), based on the temporal correlation of their size variations; in Fig. 1b, slices belonging to the same motion cluster (whether it belonged to one or more chromatophores) are represented in the same color. This analysis revealed that clusters were composed of slices that lay both within and across individual chromatophores (Fig. 1c), with some chromatophores contributing to multiple clusters (Fig. 1d-f). Within a cluster, the orientation of each slice tended to align centripetally towards a virtual point located between the contributing chromatophores (Fig. 1g), effectively forming a closed "virtual chromatophore" rather than ring-like patterns. An example is shown in Fig 1e, in which a single chromatophore (red ring) was divided into two adjacent slice clusters: one half was grouped with neighboring chromatophores located above (cyan); the other half with chromatophores located below (purple). Notably, smaller (yellow, less mature) chromatophores in the same region did not deform and therefore did not cluster, despite lying in the same apparent innervation field. These inactive chromatophores proved to be very useful, serving as stable anchors for motion tracking. Whereas *E. berryi* proved extremely useful to define the important parameters of our analyses and troubleshoot our techniques, this nocturnal species' camouflaging behavior was very limited and only very moderately interesting. We thus ran our higher-resolution analyses in

subsequent experiments, conducted with *Sepia officinalis*, a diurnal master of camouflage with large numbers and density of chromatophores.

Figure 1: Identification and characterization of putative chromatophore motor units in *E. berryi*. a) Example HD-video frame extracted from a *E. berryi* recording. b) Visual representation of presumed motor units. Colors represent motion clusters of chromatophore slices grouped together by covariation using the HDBSCAN algorithm. Clusters represent presumed motor units. c) Number of chromatophores per cluster size, and frequency of chromatophores belonging to multiple clusters. d) Zoom on the frame shown in a, with detected slices overlaid in magenta. e) Zoom on two overlapping motor units from b. The central chromatophore (red circle) contributes slices to both the blue and the purple clusters. Cluster centers of mass, calculated from the epicenters of the chromatophores belonging to each cluster, are shown as colored crosses. f) Surface areas of slices belonging to each cluster over time. In bold, the surface areas of the slices highlighted in white in e. g) Distribution of slice orientations relative to the motor unit's center of mass.

Analysis of chromatophore motion in Sepia officinalis

Building on our analyses with *E. berryi*, we applied CHROMAS to the skin of *S. officinalis*, where chromatophore density is roughly ten times higher and chromatophore dynamics are correspondingly harder to resolve. Animals were recorded under head fixation with and without sedation (HF+S and HF), conditions chosen to elicit spontaneous chromatophore activity required for correlation analysis. Comparisons across these two conditions revealed no difference, suggesting that

the dynamics observed were inherent to the chromatophore control system. Using
8K video recordings and visible implant elastomer (VIE) markers for consistent
region identification across days or weeks (see methods), we carried out
experiments with 12 animals between 5 days and 10 months of age, and ranging
from 5 to 100 mm in mantle length. We here analyze sections (0.5 by 0.5 mm) of a
108-s HF video segment (2160 images) and a 20-s HF+S segment (400 images),
that together yielded 3,285 segmented individual chromatophores. These datasets
provide the basis for the following quantification of chromatophore anisotropic
expansion and motor unit identification.

Interpretation of single-chromatophore fine motion

We performed PCA on the 36- slice time series of each chromatophore (see Fig. 2a
for a single chromatophore) and from it estimated the number of PCs necessary to
explain most of the variance in the motion data. The number of relevant components
was determined by identifying the point of maximum curvature in the cumulative
explained variance curve, as implemented in the CHROMAS pipeline (ref, see
Methods). In Fig 2b, this point corresponds to PC5, such that PCs 1-5 account for
about 95% of the variance in the data . Over 1,829 chromatophores in the HF + S
dataset, this approach yielded 3.715 ± 1.211 PCs per chromatophore on average
(Fig. 2c). To test for an effect of sedation, we repeated the same analysis on 1,456
chromatophores from the head-fixed, not sedated condition (HF), obtaining 3.683 ±
1.224 PCs. A Welch's t-test reveals no significant difference between these two
conditions (t = 0.21, p = 0.83; Cohen's d = –0.03) (Fig. 6d).

The number of dominant explanatory components identified by PCA was then used
to initialise an Independent Component Analysis (ICA). This operation extracted the
dynamics of the putative underlying sources (Fig. 2d) and revealed their spatial
influence on individual chromatophore slices (Fig. 2e–f). Each independent
component—interpreted as the signature of firing activity in a single motor neuron—
was overlaid onto the 36 equal polar slices to generate a detailed influence map for
each chromatophore (Fig 2e-f). Across our dataset, the putative motor neurons so-
identified innervated between three and sixteen slices per chromatophore.
Importantly, these targeted slices consistently formed contiguous domains around

the chromatophore's circumference. The influence profiles frequently adopted a
petal-like shape, characterised by a peak in one polar position and a gradual
tapering of influence across the adjacent slices. This parcelated representation
suggests the topography of motor innervation of single chromatophores, and
provides the most likely explanation for the observed anisotropic expansion patterns
of single chromatophores.

Figure 2: Multiple innervation of individual chromatophores. a) Activity traces (surface area as
functions of time) of 36 polar slices from a single chromatophore, offset vertically pairwise by 5 pixels
for clarity. Each trace represents the positive, detrended change in surface area over time,
highlighting expansion dynamics while excluding negative (contraction) phases. b) Cumulative
explained variance plot for Principal Component Analysis. Relationship between number of principal
components and explained variance, with PCs ranked by decreasing contribution . Red stippled line at
PC corresponding to greatest slope change (see Methods) (here PC5). c) Histogram showing the

number of principal components identified per chromatophore across the dataset. d) Independent sources acting on the 36 slices of a single chromatophore, extracted with Independent Component Analysis. In this case, the chromatophore is controlled by 5 independent sources that likely correspond to 5 distinct motor neurons. e) Polar plot representing the influence of each source (motor neuron activity) on each slice. Black line indicates the first radial slice; influence of a source on a slice expressed in % (radial scale). f) Examples of PCA-ICA analyses run on six other chromatophores.

Structure and size of chromatophore motor units

Affinity-propagation clustering of the 6,795 independent components (ICs) extracted from 1,829 chromatophores (HF+S dataset) yielded 754 clusters, and thus putative motor units (MUs). Each putative MU comprised a mean of 9.003 ± 4.755 ICs, and 95.877 ± 7.566 % of these MUs spanned multiple chromatophores, indicating extensive multi-chromatophore innervation (Fig. 3a-b). For this spatially bounded dataset (Fig 3b), the cluster- size distribution was skewed, with 90 % of motor units innervating **fewer than 14 chromatophores** (Fig. 3c). A Welch's t-test revealed no significant difference in the number of components per MU between the two groups (HF = 8.551 ± 5.753, n = 626, vs. HF+S = 9.003 ± 4.755, n = 754; $t \approx -1.57$, $p = 0.12$).

Figure 3: Identification of putative motor units. a) Schematic representation of derived chromatophore innervation by motor neurons. Each chromatophore is depicted as a circle, with arcs around them representing independent components identified by ICA, each interpreted as corresponding to the zone of influence of a single motor neuron on that particular chromatophore. One goal of this clustering was to reveal shared chromatophore innervation and thus, motor units. Colors and lines connecting IC arcs indicate that they belong to the same motor unit, highlighting the shared innervation pattern across multiple chromatophores. b) A small number of clusters of size 3 to 5 are mapped back on the original image for visualisation. c) Distribution of cluster size (size in number of IC). d) Distribution of cluster convex hull areas, showing a right-skewed profile with most clusters occupying smaller areas. All values are reported in pixel² (px²); note that the main text reports corresponding measurements in µm² after conversion (10 pixels = 1 µm, thus 1 px² = 0.01 µm²). e) Relationship between chromatophore count and convex hull area for each cluster. f) Chromatophore density (chromatophores per pixel²) as a function of convex hull area, plotted on a log-log scale. The inverse relationship reflects decreasing density in larger clusters. g) Histogram showing the distribution of second-order co-occurrence counts—defined as the number of chromatophore pairs co-occurring in at least two clusters—across 1000 randomised datasets (blue). Randomisations preserved both the number of clusters and the number of memberships per chromatophore. The red dashed line indicates the observed experimental value, significantly higher than expected by chance.

The spatial relationship between chromatophores belonging to the same putative MU was assessed by calculating pairwise distances and convex hull areas of putative MU clusters. The global nearest-neighbor distance (NND)—defined as the average shortest distance from each chromatophore to its closest neighbor across the entire dataset—was 6.130 ± 1.589 µm. In contrast, the mean within-cluster NND was

40.548 ± 46.946 µm, indicating that chromatophores grouped within the same motor

unit are not limited to immediate neighboring chromatophores. The corresponding

furthest distances—defined as the greatest distance between any two

chromatophores within a cluster—was 228.96 ± 94.36 µm. The corresponding global

value (361.50 ± 49.80 µm) depends directly on the field of view of the camera and is

thus less informative.

To quantify spatial coverage, the convex hull area was measured for each cluster.

(The convex hull area refers to the area of the smallest convex polygon that fully

encloses a given set of points.) The distribution of these areas is shown in Fig. 3d,

with most clusters occupying relatively small regions. Across 754 clusters, convex

hull areas ranged from 25.18 µm² to 915,087.66 µm², with a median of

136,199.10 µm² and a mean of 209,793.70 µm². Ninety percent of clusters had a

convex hull area smaller than 51,513.25 µm² (or 0.0515 mm², the equivalent of a fine

grain of sand). Figure 3e-f illustrates the relationships between spatial and

compositional features of motor units. A positive correlation ($r = 0.66$) was observed

between cluster area and chromatophore count (Fig 3e), while a weak negative

correlation was found between area and chromatophore density ($r = -0.14$) (Fig 3f),

and between chromatophore count and density ($r = -0.14$). These relationships

suggest that clusters occupying larger areas tend to have more chromatophores

(Fig. 3e), but at lower density (Fig. 3f).

We next examined the second- order co- occurrence, defined as the number of

unique chromatophore pairs that were parts of at least two distinct motor units. This

number was 1,073 in our dataset and compared to that expected with a null

hypothesis of random paired assignments. A set of 1,000 random permutations

yielded a distribution shown in Fig 3g, and an expected mean value of 263.72 ±

16.94 pairings . The highly significant difference between these two values ($Z =$

49.36, $p < 0.0001$) indicates that co-innervations are not random, and suggests the

existence of cooperative or joint-causality mechanisms for the innervation and

control of nearby chromatophores.

The spatial structure of putative motor units displayed a wide range of configurations

(Fig. 4). Some clusters were compact, with chromatophores arranged closely

together and spanning small distances (e.g., panels a–d). In many cases, clusters

(i.e., MUs) shared chromatophores, resulting in overlapping network structures (e.g.,

panels e–f). Other clusters exhibited more elongated or linear profiles (e.g., panel g),

with chromatophores extending over a longer axis. In addition, mixed profiles were

observed, where a core group of closely spaced chromatophores was connected to

one or several distant chromatophore(s) (e.g., panels g and h). Because our field of

view did not encompass the mantle in its entirety, our analysis will have missed large

motor units spread over very large areas, if they exist.

Figure 4: Spatial organization of putative motor units. Chromatophores belonging to the same
correlated-motion cluster, or putative MU, are shown as colored dots connected by lines. a-d)

Compact clusters. e-f) Overlapping clusters with shared chromatophores. g) Elongated cluster with a
linear structure. h) Mixed profile featuring a core group of adjacent chromatophores connected to a
single, distant chromatophore.

Dynamics of chromatophore expansion and contraction

Because chromatophore deformations are caused by active expansion (radial
muscle contraction) and passive (elastic) recovery (radial muscle relaxation), the two
phases of chromatophore deformation have different dynamics (fast expansion,
slower recovery). We tested whether these kinetic differences could be detected in
our videographic measurements at 20 fps, i.e., at sampling rates low compared with
those of electrophysiology (Florey, 1969). We analysed expansion and recovery
processes on individual slices, so as to isolate the activity of different presumed
radial-muscle groups within each chromatophore. We focused on chromatophores
with high activity levels to ensure clear detection of expansion and contraction
phases and accurate estimation of their respective speeds and durations.

Although very variable, our data showed that chromatophore expansion is
significantly faster and more stereotyped than contraction across chromatophores
(Fig. 5 and S1). These results indicate that these kinetic features can be detected
with the temporal resolution of our video-imaging. Specifically, across a subset of 10
active chromatophores (from the HF+S dataset), expansion phases occurred at an
average speed of 0.667 ± 0.590 pixels/frame, compared with 0.422 ± 0.557
pixels/frame for the contraction (n=2157 events). This difference was significant
(paired t-test, $t = 19.50$, $p < 0.001$). In our datasets, 10 aligned pixels span 1 µm.
Expansion events were spread on average over 4.41 ± 1.91 frames vs. contractions,
5.90 ± 3.50 frames. These results are thus consistent with the basic mechanisms of
chromatophore motor activation.

Figure 5: Composite summary of chromatophore slices expansion–contraction dynamics.
Events are first filtered to remove very small peaks (expansion amplitude ≤ 15% of the observed

amplitude range). Remaining events are split by expansion amplitude percentiles: SMALL = lower tail,

MEDIUM = central band, LARGE = upper tail. Within each tail band, outliers in either amplitude or

speed are removed using Tukey's IQR rule, and statistics are computed on the remaining events (see

Methods). a) Peak-aligned event segments (10 randomly selected events per column) for each group.

Traces are shifted to align at the peak (t=0). Same y calibration in the four panels. b) Mean event
profile for each size group, where each segment is color-coded by instantaneous size change
($|\Delta radius|$ for pairs of consecutive frames); LUT at right. Same y calibration in the four panels c)
Means ± SDs of chromatophore size change (px/frame) for expansion (green) and contraction (red)
phases. Statistical significance evaluated using paired t-tests between expansion and contraction
phases (n = 2157 event pairs). Asterisks indicate significant paired differences $p < 0.05$ (*), $p < 0.01$
(**), $p < 0.001$ (***), and $p < 0.0001$ (****). d) Same as in c but for event duration (in frames).

Direct activation of single chromatophore motor neurons (*Euprymna berryi*)

The above observations in *Sepia* were based on high-resolution video image data of

spontaneous resting activity, enabling the simultaneous monitoring of large regions

of skin and thus, the simultaneous tracking of large numbers of chromatophores and
measurements of their motion correlations. To test for the validity of the approach,
we also carried out combined video imaging and whole-cell patch-clamp recordings
of single motor neurons, thus enabling the direct control of single-motor neuron
discharge. Our goal was to test, on a small scale, the validity of the large-scale
motion-correlations approach based on measurements of spontaneous activity. In
particular, we wished to check directly whether individual chromatophores are
anisotropically controlled by multiple motor neurons with polarised action (as
suggested by the deformation "petals" in figure 2), and whether the action of
individual motor neurons on a field of nearby chromatophores is usually directional,
with expansion vectors oriented towards a virtual center of mass.

To this end, we returned to *Euprymna* (to increase the probability of finding both
motor neurons and corresponding chromatophores, because the number of
chromatophores in this species is small), patched a motor neuron in the posterior
chromatophore lobe and looked for the patch of skin containing the innervated
chromatophores. Once those had been identified, we injected a 1s-long direct
current in the patched motor neuron to cause sustained firing (Fig. 6a), and observed
the effect of this activation on the chromatophores (Fig. 6b and Fig. SV1). Here the
motor neuron innervates five nearby chromatophores, causing a partial, directed,
petal-like expansion of each one towards a virtual common center located near their
common center of mass. Detecting this innervation pattern in this example proved
difficult using spontaneous activity alone (*i.e.*, excluding the electrophysiological
recording), due to the small amplitude of the chromatophore deformations caused by
single motor neuron action potentials. These observations were repeated in 16 motor
neurons from 12 different animals (Fig. 6c). Note however that the expansion of
chromatophores by single- motor neuron electrical activation was not always
anisotropic (Fig. 6c - bottom right), suggesting that some chromatophores in
Euprymna are controlled by single motor neurons (or that motor neurons with a
common chromatophore target can be electrically coupled). We conclude that our
quantitative videographic approach based on the analysis of spontaneous
chromatophore deformations in *Sepia* reveals features that are consistent, but on a
large scale, with those triggered by focal motor stimulation experiments.

Figure 6: Patch clamp stimulation of chromatophore-lobe motor neuron. a) "Patch-clamp recordings from a chromatophore-lobe motor neuron. Ten consecutive sweeps are shown, vertically offset for clarity and downsampled x4 (from 100 to 25 kHz). b) Sequence of video stills showing chromatophore expansion elicited by stimulation of the same motor neuron as in a. In orange, chromatophore epicenters, defined as the center of mass at the frame where the chromatophore is in its most contracted state. Each chromatophore expands anisotropically toward a common center of innervation (approximated as a red cross). c) Composite images showing chromatophore outlines before (blue) and during stimulation (green). The first three examples display anisotropic expansion across several chromatophores within a single motor unit—the first example is the same motor unit shown in a–b. The last example shows isotropic expansion of a motor unit composed of a single chromatophore.

figureSV1.mp4

Figure SV1: Video example of patch clamp stimulation. On the left, a video showing the expansion of five chromatophores composing a motor unit—the same motor unit shown in 6b. On the right, membrane-potential trace from the corresponding motor neuron, synchronized with the video on the left.

Discussion

Our results extend early focal electrophysiological observations to large physical scales, thanks to the quantitative and easy-to-use videographic and quantitative methods that we developed recently (Ukrow & Renard, 2025) . Messenger (2001), drawing on Florey's electrophysiological studies (Florey, 1969), suggested that chromatophores are controlled by at least four motor neurons. Our decomposition consistently revealed three to four independent causes for individual chromatophore deformation. Our results match and support these early inferences, but are here

based on direct analysis of thousands of chromatophores in parallel. Likewise,
Dubas (1985, 1986) estimated that a single motor neuron may innervate tens to
hundreds of chromatophores, typically of the same color. Our clustering revealed
units of variable size, typically around ten chromatophores in *S. officinalis* but
sometimes extending much further (up to thirty), matching her conclusion that motor
fields are both distributed and overlapping. We never detected, however, motor units
containing many tens to hundreds of chromatophores with our technique. This may
be due to the generally small expansions caused by spontaneous motor neuron
discharge, presumably usually limited to single action potentials; it is possible also
that the gain of motor neuron-chromatophore transfer varies over the spatial extent
of an innervation field. If so, only some of the targets of a motor neuron would be
identifiable via our methods.

Contrary to mature skeletal muscle innervation in vertebrates (Henneman, 1957;
Kernell, 2006), individual invertebrate muscle fibers often receive convergent
excitatory inputs from more than one motor neuron (Hoyle, 1955; Bullock & Horridge,
1965; Sasaki & Burrows, 1998, Florey, 1969; Dubas & Boyle, 1985). The nature of
our videographic methods was not sufficient to resolve chromatophore fiber poly-
innervation, if it exists. If anything, however, the adjacent deformation "petals" we
identified rather suggest adjacent, but non overlapping territories of muscle-fiber
innervation by individual excitatory motor neurons. The patterns of correlated
deformation of multiple chromatophores were consistent with earlier findings (Dubas
& Boyle, 1985; Reiter et al., 2018) that individual motor neurons innervate
chromatophores of the same colour. Given the systematic intercalation of yellow and
brown chromatophores in the skin (refs) (in which chromatophore color is correlated
with chromatophore age, Reiter), the monochromatic innervation patterns, combined
with their great spatial precision, indicate a remarkable degree of local control of
chromatophore innervation by motor neurons in *Sepia*.

Most strikingly, our results reveal a phase offset between the anatomical matrix of
chromatophores and their functional innervation. Each chromatophore can be
subdivided into several (typically three to four) independently controlled zones, each
containing an adjacent subset of the 15–25 radial muscles that control each
chromatophore; these zones typically cut across neighboring chromatophores. This

results in the possible generation of "virtual chromatophores", or chromatic zones
formed by the convergence of the quadrants of nearby chromatophores. This
suggests that the smallest addressable units of cephalopod skin are not necessarily
single chromatophores but rather topographically ordered subsets of radial muscles,
spread over multiple chromatophores and grouped by their shared motor neuron.
This view modifies and extends Packard's (1982) billboard analogy: chromatophores
are not isolated "billboard lightbulbs" but overlapping zones wired through multiple
control lines. Such an arrangement implies strong developmental constraints on
wiring while simultaneously enhancing the flexibility of body-pattern generation.

Our analyses suggest that chromatophore control operates through overlapping
innervation fields that often do not coincide with the anatomical boundaries of
individual chromatophores. In *E. berryi*, motor units tended to be compact and
converged toward points between neighboring chromatophores, effectively creating
"virtual chromatophores" composed of partial contributions from adjacent cells.
Several mechanisms could underlie this organization. Developmentally, axons may
innervate radial muscles opportunistically within restricted skin regions rather than
selectively targeting individual chromatophores, leading to fields that span across
cells. The inactivity of younger orange chromatophores embedded within active
fields and the independent control of yellow chromatophores suggest that innervation
may also be stratified by depth, consistent with the layered organization of
chromatophore colors (Messenger, 2001) and the developmental processes shaping
this system (Packard, 1982). As chromatophores age and change color, vertical
migration within the skin might bring them within other motor innervation territories
and lead to their reinnervation. Our techniques will now enable the study of the
development of chromatophore innervation and possibly, of reinnervation as
chromatophores age and change pigment.

Functional implications

Although motor units are often composed of multiple chromatophore sectors rather
than whole chromatophores, the resulting pigment fields can be visually
indistinguishable from those produced by full-chromatophore control. This
fragmented organization could simply reflect wiring constraints, but it may also

confer functional advantages. By recruiting sectors from different chromatophores,
motor units introduce irregular contours, asymmetries, and non-uniform shapes,
expanding the expressive range of the skin and reducing geometric repetitiveness in
pattern textures. Sector-based units may therefore permit finer spatial modulation or
realistic pattern "noise", particularly during partial activation states.

Most motor units we observed fell below 51,513 µm² in convex-hull area, roughly
equivalent to a ~227 µm square. This scale matches the grain size of very fine sand
(125–250 µm; Wentworth, 1922; Blott & Pye, 2012), which dominates Mediterranean
coastal habitats where *S. officinalis* is common (Sandulli et al., 2010; Bettoso et al.,
2016; Dauvin et al., 2017; Asensio-Montesinos et al., 2022). Such correspondence
suggests that motor-unit resolution may be ecologically tuned to substrate textures,
providing effective camouflage without incurring the energetic costs of finer-grained
control.

Shared innervation of chromatophore sectors by multiple motor neurons may enable
gradual, fluid transitions between patterns. As one unit ramps down and another
ramps up, overlapping territories smoothen otherwise abrupt boundaries. These
overlaps could also underlie dynamic displays such as the "passing cloud" (How et
al., 2017; Laan et al., 2014; Mather & Mather, 2003), in which waves of activity
propagate seamlessly across the skin.

Motor units exhibited diverse geometries, from compact clusters to elongated or
fragmented ensembles. Such variability may be functionally specialized: tight units
supporting sharp edges or high-contrast motifs, while dispersed units contribute to
gradients or subtle textures. This architecture resembles receptive-field mosaics in
sensory systems. Just as retinal ganglion cells (Masland, 2012) or cutaneous
mechanoreceptors (Mountcastle, 2005) vary in size and density to capture distinct
features, chromatophore motor units form an outward-facing mosaic of overlapping
"projective fields" that compose complex visual patterns.

Technical issues

Our non-invasive recordings and large-scale analyses rely on indirect inference
based on spontaneous correlated motion rather than direct electrophysiological

measurements. The great advantage of this approach is that it enables large-scale
measurements in freely behaving animals. This would be impossible with
electrophysiological recordings. Chromatophore expansion served as a proxy for
radial muscle contraction, itself a result of motor neuron activity. One limitation is that
the inferences we make about motor neuron activation are indirect. Using
simultaneous motor neuron recordings and chromatophore imaging in reduced
preparations of the brain and mantle skin, we could nevertheless confirm several of
our interpretations using direct motor neuron activation. The non-isotropic action of
individual motor neurons on individual chromatophores and the converging
expansion of nearby chromatophores under the action of a shared motor neuron, for
example, could each be confirmed directly. We observed, however, that the direct-
current injections needed to generate a detectable chromatophore expansion were
often prolonged, causing the production of trains of action potentials in the patched
neuron. Spontaneous chromatophore expansion, however, is probably caused by
spontaneous and isolated (i.e., single) motor neuron action potentials. It is possible,
therefore, that what can be detected using our image analysis approach on
spontaneous activity underestimates the true patterns of innervation. This will be
especially true if the gains of the many neuromuscular junctions made by a single
motor neuron vary across its output field. Only the stronger outputs will lead to
detectable motion. (Note that the petal shape of the chromatophore expansion
quadrants suggests a gradient of innervation strength—central peak flanked by
decreasing motion amplitude) over each corresponding quadrant.) We also observe
that, to be interpretable, the patterns of chromatophore expansion during
spontaneous "noisy" activity should be as decorrelated as possible. If for example,
two motor neurons share a presynaptic excitatory drive, they may often fire together
when their common presynaptic neuron is active, leading to the erroneous
delineation of motor units. Our approach must therefore be applied in conditions of
minimal activity, ideally during the apparent spontaneous flicker that is usually
observed in resting animals.

Chromatophore slice dynamics were estimated from membrane deformation rather
than direct detection of muscle insertion points; similarly, putative motor units were
defined by clustering of correlated activity profiles. While such proxy-based
approaches involve multiple inferential steps, they are routinely used in neuroscience

and physiology to extract functional information from imaging data—for example,
inferring spiking from calcium fluorescence or estimating muscle activation from
high-speed video recordings. Our strategy applies the same principle to
chromatophore dynamics, providing scalable, non-invasive access to living animals.

The two principal clustering strategies which we settled on and used— HDBSCAN,
with *E. berryi*, and Affinity Propagation, with *S. officinalis*— make different
assumptions about cluster structure. Although both approaches yielded consistent
biological insights, it should be noted that inferred motor units are not insensitive to
algorithmic choices and should be interpreted with this in mind. More generally,
clustering constitutes a form of model selection, and alternative algorithms may
reveal complementary aspects of motor-unit organization.

Just as in any imaging approach, spatial and temporal sampling considerations
represent important constraints. In *S. officinalis*, recordings were restricted to small
regions of the dorsal mantle, precluding whole-body comparisons and leaving open
whether motor-unit organization is conserved across skin regions. Developmental
factors further narrowed the available time windows for observation: chromatophore
overlap in growing *E. berryi* hatchlings and low spontaneous activity in young *S.*
officinalis juveniles restricted the stages suitable for analysis. In this context, the
large, spatially dispersed motor units observed in *S. officinalis* may not represent a
universal architecture; they could reflect the developmental stage of the individuals
we studied or species-specific ecological requirements. Previous reports based on
more limited, but electrophysiological stimulation experiments, suggest considerable
diversity across cephalopods, with Dubas (1985) describing widely distributed motor
units in octopus and Florey (1969) reporting more compact units in squid. The extent
to which motor unit size and shape vary as functions of species, position on the
body, and developmental stage, will need to be examined.

The complex mechanics and innervation patterns of chromatophores introduce
ambiguities. Elastic recoil of the pigment sac and mechanical coupling between
adjacent radial muscles likely propagate forces beyond the site of neural activation,
producing secondary deformations (of unknown amplitude) in neighboring slices. In
addition, overlapping motor neuron axons could converge onto the same set of radial
muscles, in which case a single slice would at times reflect the combined drive of

multiple neurons. Together, these factors make it difficult to determine where the
influence of one motor neuron ends and that of another begins. As a result, our
estimates should be viewed as functional partitions of activity rather than definitive
anatomical boundaries.

In conclusion, we developed an image analysis pipeline (Ukrow & Renard, 2025) and
used it to analyse the structure of chromatophore expansion and shape control in
two species of coleoid cephalopods, from high resolution movies of freely moving
animals. This approach revealed many fine features at the distal end of this highly
specialized motor system. By identifying first the motor units, as done here, it may
become possible to reveal important aspects of their presynaptic drive and thus
make precise and testable predictions about the organization of this remarkable
neural system.

Methods

Chromatophore video acquisition and analysis

Animals

All research and animal care procedures related to video acquisition of chromatophores and analysis were carried out in Frankfurt, in accordance with the institutional guidelines that are in compliance with national and international laws and policies (DIRECTIVE 2010/63/EU; German animal welfare act; FELASA guidelines). The study was approved by the appropriate animal welfare authority (Regierungspräsidium Darmstadt) under approval number V54-19c20/15-F126/2012.

Hummingbird bobtail squids *Euprymna berryi* and European cuttlefish *Sepia officinalis* were hatched from eggs laid in house and reared in a seawater system. The closed system contained artificial seawater (ASW; Instant Ocean) with a salinity of 33-36% and pH of 8.1-8.4. Water quality was monitored continuously and was tested regularly. The housing temperature was different in function of the species. Aqua Medic *Tri Complex* supplied macro- and trace elements continuously. A constant water through-flow resulted in 5 complete water exchanges per hour. Room illumination provided a 12 h–12 h light–dark light cycle with gradual on- and off-sets at 07:00 and 19:00. Enrichment consisted of natural fine-grained sand substrate, artificial plastic plants and translucent red plastic houses. Both species were fed live food (either *Neomysis* spp. or small *Palaemonetes* spp.) twice per day.

E. berryi were reared in a seawater system at 20-26 °C. The filtration systems included filter bags, protein skimmer, biofilter, UVC-sterilisator and heater. The adult animal groups were housed in 300-L plastic tanks, juveniles in plastic boxes of 18-36 L. Experimental animals of unknown sex, just a few days after hatching, ranging from 3 to 6 mm in mantle length, were selected for healthy appearance and calm behaviour.

S. officinalis were reared at 18-20 °C. The filtration systems included drum filter, protein skimmer, biofilter, UVC-sterilisator and chiller. The animals were housed individually in 30-90 L PVC tanks with a constant water through-flow resulting in 5

complete water exchanges per hour. Animals above 50 mm in ML were fed once per day defrosted fish or shrimp. Experimental animals of unknown sex, 5 days to 10 months after hatching, ranging from 5 to 100 mm in mantle length, were selected for healthy appearance and calm behaviour.

Arena and filming equipment

E. berryi: the arena design aimed to confine the hatchlings within an area fully covered by the camera's field of view. The resulting arena measured 10 cm × 10 cm × 3 cm in height and included a conical sub-arena with a direct water inflow (Fig. S2a). This cone featured small holes allowing water to outflow into a larger surrounding chamber. The design incorporated a glass lid to reduce visual distortion by enabling direct water-glass contact and improving image clarity. Additionally, the arena floor included a glass surface to permit imaging from below. A peripheral "trench" area surrounding the chamber directed excess water through outflow tubing, ensuring continuous circulation. For recording, we used a Basler Ace 2 (a2A2590-60ucPRO) equipped with a Kowa lens (LM25JC10M). The imaging configuration enabled 1920 × 1920 px recordings corresponding to a 1 × 1 cm field of view, large enough to encompass the whole body of *E. berryi*. Illumination was provided by a custom-built LED ring mounted above the imaging arena and controlled through a custom software to modulate light intensity cyclically during recordings. Recordings were performed inside a custom sound- and light-proof enclosure lined with acoustic foam to dampen external vibrations and reflections. The box contained the imaging arena, camera, and LED ring, ensuring stable sensory conditions during recordings.

S. officinalis: the arena measured 20 cm x 30 cm x 10 cm, was made of acrylic glass material and featured a height-adjustable cover along with designated water entry and drainage points. Four LED lamps were mounted on adjustable side brackets around the arena, allowing control over their height, position, and angle (Fig. S2b). For recording we used the Basler CoaXPress 2.0 boA9344-70cc with the GMAX3265 CMOS sensor, capable of delivering 70 frames per second with a 65 MP resolution (9344 × 7000 pixels). It is equipped with a macro lens, the Qioptiq Apo-Rodagon-D 2x 4/75. The camera was installed on motorised linear rails above the arena for X-Y translation, and its position was controlled using a joystick to keep the

area of interest within the frame. The camera could be moved on the z-axis through a fine translation stage (Fig. S3). Acquiring videos at the camera's native full-sensor resolution (9344 × 7000 px; 65.4 MP) exceeded our data-throughput capacity: software compression was too slow and hardware compression introduced visible artifacts. We therefore recorded a square ROI covering ~25% of the sensor (3968 × 3968 px at 20 fps), corresponding to a 400 × 400 µm field of view (≈ 9.9 px/µm; 0.10 µm/px). This adjustment reduced the total image size but did not alter spatial sampling density, as the physical magnification and pixel size remained identical. This adjustment was not considered a significant loss of information, as the majority of the full frame was out of focus, due to the non-flat curvature of the animal's mantle.

Sedation

We used ethanol as a sedative during imaging. To achieve a state of sedation, the animal was gradually exposed to ethanol dissolved in artificial seawater, with the concentration incrementally increased up to 1.5%. The procedure was carried out in a large, dark bucket to reduce visual stimulation. This controlled approach minimised stress on the animal while ensuring effective sedation.

Ethanol is widely used in cephalopod research for its accessibility and reversibility (Andrews et al., 2013; Gleadall, 2013), but could in principle suppress neural activity or alter muscle responsiveness. In our dataset, however, PCA revealed near-identical component counts in sedated and awake head-fixed animals, and motor-unit organization remained unchanged. Thus, ethanol neither artificially inflated nor suppressed the apparent number of motor subunits, validating its use as a non-invasive means to stabilize animals during high-speed recordings.

Marking procedure

Because the entire mantle of *Sepia* could not fit under the camera view, it was essential to consistently image the same skin region across recording sessions. To

solve this issue we used Visible Implant Elastomer Tags (Northwest Marine
Technology, Inc. n.d.), a small and practical silicon-based tagging method. To apply
the tag, animals were anesthetised with 1.2 % ethanol and a drop of silicon-base tag
was injected through a syringe in the skin (Fig. S4). The silicon was coupled with a
curing agent and the tag solidified within 24 hours. This type of tag has been shown
to be very effective in cephalopods (Barry et al., 2011; Brewer et al., 2012; Zeeh et
al., 2009). In our case, the skin of both *Euprymna* and *Sepia* retained the tag for the
entire lifespan and veterinarian analyses showed that the tagging did not cause any
form of infection. A blue light lamp attached to the camera helped us find the tag by
making it fluoresce. We then always filmed the area of skin around the same tag.

Head fixation

To enable long filming sessions, we utilised head fixation, allowing us to film a single
area continuously with only minimal camera adjustments during recording (Fig. S5).
Animals were transferred to a bucket containing 1.7% ethanol in ASW, and
transferred to a shallow tank (1.5% ethanol in ASW, continuously bubbled with
oxygen) at the onset of surgical plane anesthesia. The dorsal head was raised 1cm
above the water line by a silicone-coated head rest, and the gills were superfused
with tank water at a rate of 60 ml/min via soft silicone tubes inserted bilaterally into
the mantle cavity. Breathing rate was maintained by observation of the collar
musculature and changes in ethanol concentration. Lidocaine (2%, 0.2ml) was
injected subcutaneously before exposing the dorsal head cartilage, reapplying
lidocaine and removing overlying connective tissue. A thin layer of Vetbond (3M) was
then applied to the dorsal head cartilage before attaching a custom 3D-printed head
plate (biocompatible resin) with dental cement (Venus, Kulzer) before sealing the
wound with surgical glue and transferring the animal to a recovery tank.

Filming sessions

E. berryi: 14 animals were filmed. To maximize spontaneous chromatophore activity,
individuals were lightly sedated with 1–1.5% ethanol, which also induced slight
chromatophore contraction. Expansion was triggered by cyclically varying the
ambient light intensity with a custom, remotely controlled LED ring positioned above
the arena, taking advantage of the light-dependent dynamics of chromatophores

(Zylinski et al., 2011). Filming sessions varied in duration depending on the
experimental condition. Sedated animals were recorded for up to 20 minutes before
being returned to normal seawater for recovery, whereas non-sedated individuals
could be filmed for up to 1 hour with continuous seawater circulation. Whenever
possible, the same individuals were recorded every three days over the course of
development, spanning a total period of approximately two months. The results
presented here are based on recordings from a lightly sedated individual exposed to
cyclic changes in ambient light intensity. This condition produced the most stable
and interpretable chromatophore dynamics, minimizing motion artifacts while
maintaining spontaneous activity.

S. officinalis: Over the course of the study, 12 individuals ranging in age from 5 days
to 10 months post-hatching were filmed under various experimental conditions. A
range of experimental configurations was explored to optimize chromatophore
visibility and recording stability, including evoked camouflage using printed or e-ink
patterns presented beneath a transparent tank, light ethanol sedation (up to 1.5 %),
and head fixation. In selected individuals, the same skin region was imaged at three-
day intervals for a maximum duration of six months. The dataset presented here was
obtained from a head-fixed individual filmed on days 1, 2, and 6 post-surgery, both
with and without light ethanol sedation. Head fixation provided stable, high-quality
recordings with precise spatial correspondence between frames. Non-sedated
sessions lasted up to 2 hours, while sedated sessions were limited to 20-minute
bouts before recovery in normal seawater. Comparable results could be obtained
from lightly sedated animals when movement was compensated by a tracking
camera or other gentle immobilization.

Computational analyses

Chromatophore activity was quantified using the CHROMAS software package
(Ukrow & Renard, 2025), which processes video data of cephalopod skin to identify
individual chromatophores and extract their spatiotemporal dynamics. CHROMAS
provides per-chromatophore metrics such as surface area, epicentre coordinates,
and anisotropic deformation. The resulting dataset provided spatiotemporal maps of

chromatophore activity, forming the basis for subsequent neural and biomechanical
analyses.

CHROMAS was employed as described in Ukrow & Renard, 2025, with the following
parameter and model choices.

For chunking, which filters out frames likely to impair analysis, we chose the
difference-of-Gaussians (DoG) focus metric with kernel sizes k1=11, k2=5, and
standard deviations s1=2 and s2=1 pixels.

To generate training data, manually annotated images (3968x3968 pixels) were
divided into 512×512 pixel patches. Of these, 90% were randomly selected for
training and the remaining 10% were held out for testing. To improve model
generalisation, training data augmentation followed Ukrow & Renard, 2025 with the
following parameters: spatial translations (±20%), scaling (±20%), rotation (±30°),
RGB channel shifts (±25), brightness variation (±10%), contrast alteration (±30%),
perspective warping (scale 0.05–0.5), and vertical/horizontal flipping. Each
transformation was applied independently with a probability of 25%, except flipping,
which was applied with a 50% probability.

Segmentation employed the neural-network option of CHROMAS using a Fully
Convolutional Network (FCN) architecture with a ResNet-50 backbone (Long et al.,
2015). Training was performed for 500 epochs with validation every 5 epochs, using
a composite loss function combining Dice loss and cross-entropy loss in equal
proportion (50:50). Optimisation used the Adam optimiser with an initial learning rate
of 0.001. A *ReduceLROnPlateau* scheduler (mode: 'min', factor: 0.1, patience: 50
epochs) adaptively reduced the learning rate when validation performance
plateaued. The model with the lowest recorded validation error was retained for
analysis.

For registration, 900 randomly selected tracking points were initialised on a 100×100
pixel grid. Displacement vectors exceeding 10 pixels were excluded. The
interpolation parameter of the moving-least-squares algorithm was set to $\alpha = 3.0$.

For stitching, we applied the ellipse-fit option for *Euprymnia berryi* and the manual

option for *Sepia officinalis* as detailed in Ukrow & Renard, 2025. The interpolation

parameter of the moving-least-squares algorithm was set to $\alpha = 3.0$.

All other parameters were set to their default values.

Chromatophore anisotropic deformation tracking

Chromatophore anisotropic deformation tracking provides a detailed characterisation

of the directional and irregular expansion patterns of chromatophores. This analysis

begins with a stabilisation step that compensates for both global and local

deformations of the skin. To achieve this, the CHROMAS pipeline identifies stable

points on the skin, referred to as motion markers, which correspond to

chromatophores that remain small and constant in size across frames. These motion

markers are used to derive skin deformation over time, creating a stabilised

coordinate system that eliminates artifacts caused by skin movement. Each

chromatophore is divided into 36 radial slices to enable precise monitoring of its

anisotropic deformations. This number of slices, determined based on the Nyquist–

Shannon theorem (Shannon, 1948) and histological analysis (Fig. S6), was set to 36,

corresponding to twice the upper bound of histologically confirmed muscle fibers.

Using a higher resolution would mainly amplify small segmentation imperfections,

the primary source of error. The epicenter of each chromatophore is used as the

origin for radial slicing and is calculated from the chromatophore's center of mass in

its most contracted state. This epicenter remains stable across frames due to its

relative positioning within the local coordinate system of motion markers. Slice areas

are calculated by averaging distances from the epicenter to the chromatophore's

edges, followed by squaring and scaling. This method ensures accuracy even in the

presence of pixel discretization errors. Orientation of slices is maintained consistently

across frames using the local coordinates derived from motion markers, allowing for

the reliable tracking of individual slices over time.

Expansion and contraction speeds

We analyzed the expansion and contraction dynamics of chromatophores by

tracking 36 radial slices for each of ten particularly active chromatophores. For each

recording, area values were reshaped into a matrix of dimensions "number of frames

× 36 slices" and converted to radii, assuming equal-angle sectors, using the relation r
= sqrt(36 × area / π). Peak-centered events were defined for each slice after
applying a light Savitzky–Golay smoothing (window size 5, polynomial order 2).
Local maxima were detected on the smoothed radius traces using the
scipy.signal.find_peaks function with a prominence threshold of 1.5. For each
detected peak (at frame p), the onset of expansion was defined as the frame
immediately before the first positive slope encountered when scanning left from the
peak, and the end of contraction as the last frame with a negative slope when
scanning right. Slopes were computed from the first difference of the radius signal,
with a small threshold (epsilon = 1e-5) used to avoid spurious sign flips due to
numerical jitter.

For each event, we calculated expansion amplitude ($A_{exp} = r_p − r_{start}$), duration ($D_{exp} =$

$p − start$), and mean speed ($S_{exp} = A_{exp} / D_{exp}$), as well as the corresponding

contraction amplitude, duration, and speed ($A_{con} = r_p − r_{end}$, $D_{con} = end − p$, $S_{con} =$

A_{con} / D_{con}). To suppress small fluctuations that likely represented noise, we excluded

events with an expansion amplitude smaller than 15% of the full amplitude range.

Events were then grouped by expansion amplitude into three categories based on

percentile tails (tail width = 33%): "small" (lowest 33%), "center" (middle 33–67%),

and "large" (highest 33%). Within each group, we applied Tukey's interquartile range

(IQR) method to remove outliers based on both amplitude and speed, keeping only

events that fell within [Q1 − k × IQR, Q3 + k × IQR], with k = 1.

Statistical comparisons of expansion versus contraction speeds and durations at the
event level were performed using paired t-tests. Statistical significance was reported
as follows: $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****). Durations
were expressed in frames, equivalent to milliseconds when converted at 20 frames
per second (e.g., 50 ms per frame).

Number and influence of motor neuron per chromatophore

To estimate the number of distinct motor neurons controlling each chromatophore,
time series representing the expansion dynamics of radial slices were processed
through a dimensionality reduction and source separation pipeline. Each
chromatophore's area trace matrix (frames × slices) was first detrended using
scipy.signal.detrend. Principal Component Analysis (PCA) was performed using
sklearn.decomposition. PCA, and the number of meaningful components was
estimated via the elbow method, with the elbow point identified using the
kneed.KneeLocator package (Satopaa, 2011). This number was then used as the
n_components parameter for Independent Component Analysis (ICA), performed
using *sklearn.decomposition.FastICA*.

The resulting ICA mixing matrix provides the contribution of each independent
component to each radial slice. To quantify this influence, the absolute values of the
mixing matrix were normalised per slice, yielding a percentage influence profile
across components. These influence profiles were visualised using polar plots to
assess spatial distribution. The number of components (ICs) retained per
chromatophore serves as an estimate of the number of distinct motor neuron inputs
driving that chromatophore's expansion.

Number of chromatophores per motor unit

Motor units were inferred by clustering independent components (ICs) extracted from
chromatophore activity traces using the FastICA algorithm
(*sklearn.decomposition.FastICA*). To group ICs with similar temporal profiles, we
applied Affinity Propagation (*sklearn.cluster.AffinityPropagation*) using pairwise
Pearson correlation as the similarity metric. Each resulting cluster was interpreted as
a putative motor unit, consisting of ICs presumed to originate from the same motor
neuron. Chromatophores were then assigned to motor units based on their strongest
IC loading. Motor unit size was defined as the number of unique chromatophores
associated with each cluster. All steps were implemented in Python using *NumPy*
(Harris et al., 2020), *pandas* (McKinney, 2010), and *scikit-learn* (Pedregosa et al.,
2011).

Motor unit shape and structure

To characterize chromatophore clusters, we developed a Python script using *pandas*
and *NumPy* for data handling, *SciPy* for distance calculations and minimum
spanning tree construction, and *matplotlib* for visualization (Hunter, 2007). To
assess the spatial distribution of chromatophores quantitatively, we computed
nearest-neighbor distances (NND) using Python. Epicenter coordinates [epicenter_x,
epicenter_y] were extracted from our dataset. For the global analysis, we computed
pairwise Euclidean distances between all chromatophore epicenters and recorded
the minimum distance to identify each chromatophore's nearest neighbor. For the
within-cluster analysis, we grouped chromatophores by cluster label and repeated
the NND computation within each group. The output metrics included the mean
NND, standard deviation, and coefficient of variation (CV), which were used to
describe both overall spatial density and intra-cluster compactness. Results were
visualised by plotting chromatophore positions and connecting each to its nearest
neighbor with a line segment. In addition to nearest-neighbor distances (NND), we
calculated furthest-neighbor distances (FND) and convex hull areas (via
scipy.spatial.ConvexHull) to estimate cluster extent. From these, density metrics
were derived by relating chromatophore counts to convex hull area, and Pearson
correlation coefficients (*scipy.stats*) were used to assess relationships between area,
density, and number of chromatophores.

In order to visualize the putative motor units ramifications, chromatophores were
grouped by cluster label and plotted according to their epicenter coordinates, with
multi-labeled chromatophores displayed as multicolored wedges, and internal cluster
geometry was quantified by computing pairwise Euclidean distances and reducing
them to a minimum spanning tree. This framework enabled both visualization and
quantitative comparison of cluster compactness, extent, and density.

Second order innervation

To assess second-order innervation, we first extracted all [chrom_id, label] pairs
from the cluster dataset, yielding a unique set of cluster memberships. A pairwise co-
occurrence count was computed by iterating over all clusters and recording how
often each pair of chromatophores appeared together; we then counted how many

pairs occurred together in two or more clusters. For the null model, we implemented a constrained permutation in which chromatophore IDs were randomly reassigned across clusters while preserving the number of unique IDs per cluster. This procedure maintains the original cluster sizes but not the per-chromatophore memberships. Across 1,000 simulations, we repeated the pairwise counting procedure and recorded the number of pairings that reached the second-order threshold. This produced a distribution of null values, against which the real count was compared using a z-test to determine statistical significance.

Electrophysiology

Animals

Electrophysiological experiments were carried out in Beijing, China. At the time of the experiments, there were no specific national regulations governing the use of cephalopods in research in China. Nevertheless, all procedures performed at Peking University were designed to comply, as far as possible, with the principles of the European Directive 2010/63/EU and FELASA recommendations.

For electrophysiological experiments, bobtail squids *Euprymna berryi* were either hatched from eggs laid in house or purchased as eggs and naturally hatched in the lab. Both adult squids and eggs were sourced from Guangdong and Hainan provinces in China. *E. berryi* were rared in a artificial seawater (ASW) system around 24℃, with a salinity of 1.021 and a pH of 8.3. Water quality was monitored continuously and was tested regularly. The ASW filtration system consists of filter floss, filter bags, biofilter (for nitrifying bacteria), a protein skimmer, and UV-lamp. Room light was set as a 12h-12h light-dark cycle with gradual on- and off-sets at 12:00 and 24:00.

Brain and skin preparation

Juvenile *E. berryi* ranging from 5-10 mm in body length were used for electrophysiological experiments. Animals were first anesthetized by 2% ethanol in ASW, and then transferred into an acrylic chamber with a sloped silicone rubber base, filled with oxygenized calcium-free saline (460 mM NaCl, 10 mM KCl, 10 mM

glucose, 10 mM HEPES, 55 mM MgCl$_2$, and with a pH of 7.4.) at room temperature.
The arms, beak, and both eyes were removed. The mantle was opened from the
ventral side and all visceral organs were removed. The skin of the head was
removed to expose the posterior sub-esophageal mass. The posterior
chromatophore lobe was de-sheathed gently with fine forceps. The mantle was
pinned on a 70° sloped silicone rubber plane, while the brain was placed on a flat
silicone rubber base. The mantle and the brain were connected solely through pallial
nerves. The solution was then replaced by oxygenated (100% O2) calcium-
containing saline (460 mM NaCl, 10 mM KCl, 10 mM glucose, 10 mM HEPES, 55
mM MgCl$_2$, 11 mM CaCl$_2$, and with a pH of 7.4).

Electrophysiological recording and imaging setup

The patch clamp recording was performed under an upright fluorescence
microscope (BX51W1, Olympus) with a 20 water-immersion objective (UMPlanFI 20,
Olympus). The brain was visualized by a microscope camera (Moment) . The mantle
chromatophore was recorded by a camera (a2A4096-30ucBAS, Basler) with a
manual lens (MLM-3X-MP, Computar) at 30 Hz. The glass electrodes (~8 M) were
made from borosilicate glass (BF150-86-10, WPI) by a puller (P-1000, Sutter
Instrument). The pipettes were filled with internal solution (450 mM K-gluconate, 10
mM NaCl, 4 mM MgCl$_2$, 3 mM EGTA, 20 mM HEPES, 2 mM Mg-ATP, 0.2 mM Na-
GTP, 2% neurobiotin, and with pH of 7.4). Recordings were performed in current
clamp mode with a HEKA amplifier (ESC100-USB). For intracellular stimulation, step
or constant currents were injected repetitively at 0.5 Hz. The onset of the stimulation
train simultaneously triggered the camera to record chromatophore activity on the
mantle skin.

Acknowledgments

913   We thank F. Bayer for assistance in building the experimental setup; F. Kretschmer
914   for optimizing the camera control, recording software and Git workflows; F. Vollrath
915   and S. Junek for help with image acquisition; P. Musset for help with high

performance computing; L. Jürgens, S. Schwind, L. E. Reyes de Frey, D. Burgard, M. Landler, M. Minde, S. Kranz, P. Dominiczak, N. K. Vogt, G. Wexel, S. Dizdarevic and E. Northrup for animal care; and T. Woo and other members of the Laurent laboratory for constructive exchanges. We thank Alice Perenzin and Antje Berken for grant management and scientific coordination. This research was funded by the Max Planck Society (GL), by the LOEWE Schwerpunkt CMMS (State of Hesse) (GL), by the Louis Jeantet Foundation (GL), by the European Union (ERC grant CAMOUFLAGE, 10114150) (GL), by National Natural Science Foundation of China (NSFC grant 32371215) (XL), State Key Laboratory of Membrane Biology, Peking University (XL); and Peking-Tsinghua Center for Life Sciences (XL).

References

Andrews, P. L., Darmaillacq, A., Dennison, N., Gleadall, I. G., Hawkins, P., Messenger, J. B., Osorio, D., Smith, V. J., & Smith, J. A. (2013). The identification and management of pain, suffering and distress in cephalopods, including anaesthesia, analgesia and humane killing. Journal of Experimental Marine Biology and Ecology, 447, 46–64. https://doi.org/10.1016/j.jembe.2013.02.010

Asensio-Montesinos, F., Blaya-Valencia, G., Corbí, H., Beltrán-Sanahuja, A., & Sanz-Lázaro, C. (2022). Microplastic accumulation dynamics in two Mediterranean beaches with contrasting inputs. Journal of Sea Research, 188, 102269. https://doi.org/10.1016/j.seares.2022.102269

Atwood, H. L. (1977). Crustacean neuromuscular systems: past, present, and future. In Identified neurons and behavior of arthropods (pp. 9-29). Boston, MA: Springer US. https://doi.org/10.1007/978-1-4684-6967-7_2

Barry, P. D., Tamone, S. L., & Tallmon, D. A. (2011). A comparison of tagging methodology for North Pacific giant octopus Enteroctopus dofleini. Fisheries Research, 109(2-3), 370-372. https://doi.org/10.1016/j.fishres.2011.02.011

Bettoso, N., Borme, D., Faresi, L., Aleffi, I., Orlando-Bonaca, M., & Lipej, L. (2016). New insights on the biological parameters of the exploited cuttlefish Sepia officinalis

L.(Mollusca: Cephalopoda) in the northern Adriatic Sea in relation to the main fishing gears employed. Mediterranean Marine Science, 17(1), 152-162. https://doi.org/10.12681/mms.1311

Blott, S. J., & Pye, K. (2012). Particle size scales and classification of sediment types based on particle size distributions: Review and recommended procedures. Sedimentology, 59(7), 2071-2096. https://doi.org/10.1111/j.1365-3091.2012.01335.x

Boycott, B. B. (1961). The functional organization of the brain of the cuttlefish Sepia officinalis. Proceedings of the Royal Society of London. Series B, Biological Sciences, 153(953), 503–534. https://doi.org/10.1098/rspb.1961.0015

Brewer, R. S., & Norcross, B. L. (2012). Long-term retention of internal elastomer tags in a wild population of North Pacific giant octopus (Enteroctopus dofleini). Fisheries Research, 134, 17-20. https://doi.org/10.1016/j.fishres.2012.07.020

Bullock, T., & Horridge, G. A. (1965). Structure and function in the nervous systems of invertebrates.

Campello, R. J. G. B., Moulavi, D., & Sander, J. (2013). Density-Based Clustering Based on Hierarchical Density Estimates. In J. Pei, V. S. Tseng, L. Cao, H. Motoda, & G. Xu (Eds.), Advances in Knowledge Discovery and Data Mining (pp. 160–172). Springer. https://doi.org/10.1007/978-3-642-37456-2_14

Cloney, R. A., & Florey, E. (1968). Ultrastructure of cephalopod chromatophore bergorgans. Zeitschrift Für Zellforschung Und Mikroskopische Anatomie, 89(2), 250–280. https://doi.org/10.1007/BF00347297

Dauvin, J. C., Bakalem, A., Baffreau, A., Delecrin, C., Bellan, G., Lardicci, C., ... & Grimes, S. (2017). The well sorted fine sand community from the western Mediterranean Sea: A resistant and resilient marine habitat under diverse human pressures. Environmental pollution, 224, 336-351. https://doi.org/10.1016/j.envpol.2017.02.013

Dubas, F., & Boyle, P. R. (1985). Chromatophore motor units in Eledone cirrhosa (Cephalopoda: Octopoda). Journal of experimental biology, 117(1), 415-431. https://doi.org/10.1242/jeb.117.1.415

Dubas, F.., Hanlon, R. T., Ferguson, G. P., & Pinsker, H. M. (1986). Localization and
stimulation of chromatophore motor neurones in the brain of the squid, Lolliguncula
brevis.   Journal   of   Experimental   Biology,   121(1),   1-25.
https://doi.org/10.1242/jeb.121.1.1

Florey, E. (1969). Ultrastructure and function of cephalopod chromatophores.
American Zoologist, 9(2), 429–442. https://doi.org/10.1093/icb/9.2.429

Gleadall, I. G. (2013). The effects of prospective anaesthetic substances on
cephalopods: summary of original data and a brief review of studies over the last two
decades. Journal of Experimental Marine Biology and Ecology, 447, 23-30.
https://doi.org/10.1016/j.jembe.2013.02.008

Hanlon, R. T., & Messenger, J. B. (1988). Adaptive coloration in young cuttlefish
(Sepia officinalis L.): the morphology and development of body patterns and their
relation to behaviour. Philosophical Transactions of the Royal Society of London. B,
Biological Sciences, 320(1200), 437-487. https://doi.org/10.1098/rstb.1988.0087

Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P.,
Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M.,
Hoyer, S., Van Kerkwijk, M. H., Brett, M., Haldane, A., Del Río, J. F., Wiebe, M.,
Peterson, P., . . . Oliphant, T. E. (2020). Array programming with NumPy. Nature,
585(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Henneman, E. (1957). Relation between Size of Neurons and Their Susceptibility to
Discharge.   Science,   126(3287),   1345–1347.
https://doi.org/10.1126/science.126.3287.1345

Hoyle, G. (1955). The anatomy and innervation of locust skeletal muscle.
Proceedings of the Royal Society of London. Series B-Biological Sciences, 143(911),
281-292. https://doi.org/10.1098/rspb.1955.0011

How, M. J., Norman, M. D., Finn, J., Chung, W. S., & Marshall, N. J. (2017). Dynamic
skin   patterns   in   cephalopods.   Frontiers   in   physiology,   8,   393.
https://doi.org/10.3389/fphys.2017.00393

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. Computing in science & engineering, 9(03), 90-95. https://doi.org/10.1109/MCSE.2007.55

Kernell, D. (2006). The motor neurone and its Muscle Fibres. https://doi.org/10.1093/acprof:oso/9780198526551.001.0001

Laan, A., Gutnick, T., Kuba, M. J., & Laurent, G. (2014). Behavioral analysis of cuttlefish traveling waves and its implications for neural control. Current biology, 24(15), 1737-1742. https://doi.org/10.1016/j.cub.2014.06.027

Long, J., Shelhamer, E., & Darrell, T. (2015). Fully convolutional networks for semantic segmentation. In Proceedings of the IEEE conference on computer vision and pattern recognition (pp. 3431-3440). https://doi.org/10.1109/TPAMI.2016.2572683

Lucas, B. D., & Kanade, T. (1981). An Iterative Image Registration Technique with an Application to Stereo Vision. IJCAI'81: 7th International Joint Conference on Artificial Intelligence, 2, 674–679. https://hal.science/hal-03697340

McKinney, W. (2010). Data Structures for Statistical Computing in Python. Scipy. https://doi.org/10.25080/Majora-92bf1922-00a

Masland, R. H. (2012). The neuronal organization of the retina. Neuron, 76(2), 266-280. https://doi.org/10.1016/j.neuron.2012.10.002

Mather, J. A., & Mather, D. L. (2004). Apparent movement in a visual display: the 'passing cloud' of Octopus cyanea (Mollusca: Cephalopoda). Journal of Zoology, 263(1), 89-94. https://doi.org/10.1017/S0952836904004911

Messenger, J. B. (2001). Cephalopod chromatophores: neurobiology and natural history. Biological Reviews, 76(4), 473-528. https://doi.org/10.1017/S1464793101005772

Mountcastle, V. B. (2005). The sensory hand: neural mechanisms of somatic sensation. Harvard University Press.

Northwest Marine Technology, Inc. (n.d.). Visible implant elastomer. Retrieved November 21, 2024, from https://www.nmt.us/visible-implant-elastomer/

Packard, A. (1982). Morphogenesis of chromatophore patterns in cephalopods: are
morphological and physiological units the same? Malacologia, 23(1), 193-201.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., ... &
Duchesnay, É. (2011). Scikit-learn: Machine learning in Python. the Journal of
machine Learning research, 12, 2825-2830.

Reiter, S., Hülsdunk, P., Woo, T., Lauterbach, M. A., Eberle, J. S., Akay, L. A.,
Longo, A., Meier-Credo, J., Kretschmer, F., Langer, J. D., Kaschube, M., & Laurent,
G. (2018). Elucidating the control and development of skin patterning in cuttlefish.
Nature, 562(7727), 361–366. https://doi.org/10.1038/s41586-018-0591-3

Sandulli, R., De Leonardis, C., & Vanaverbeke, J. (2010). Meiobenthic communities
in the shallow subtidal of three Italian Marine Protected Areas. Italian Journal of
Zoology, 77(2), 186-196. https://doi.org/10.1080/11250000903476616

Sasaki, K., & Burrows, M. (1998). Innervation pattern of a pool of nine excitatory
motor neurons in the flexor tibiae muscle of a locust hind leg. Journal of
experimental biology, 201(12), 1885-1893. https://doi.org/10.1242/jeb.201.12.1885

Satopaa, V., Albrecht, J., Irwin, D., & Raghavan, B. (2011, June). Finding a" kneedle"
in a haystack: Detecting knee points in system behavior. In 2011 31st international
conference on distributed computing systems workshops (pp. 166-171). IEEE.
https://doi.org/10.1109/ICDCSW.2011.20

Shannon, C. E. (1948). A mathematical theory of communication. The Bell System
Technical Journal, 27(3), 379–423. The Bell System Technical Journal.
https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

Ukrow, J., Renard, M. D., Moghimi, M., & Laurent, G. (2025). A computational
pipeline to track chromatophores and analyze their dynamics eLife 14:RP106509.
https://doi.org/10.7554/eLife.106509.3

Wentworth, C. K. (1922). A scale of grade and class terms for clastic sediments. The
journal of geology, 30(5), 377-392.

Zylinski, S., & Johnsen, S. (2011). Mesopelagic cephalopods switch between
transparency and pigmentation to optimize camouflage in the deep. Current Biology,
21(22), 1937-1941. https://doi.org/10.1016/j.cub.2011.10.014

Zeeh, K. M., & Wood, J. B. (2009). Impact of visible implant elastomer tags on the
growth rate of captive Caribbean reef squid Sepioteuthis sepioidea. Fisheries
Research, 95(2-3), 362-364. https://doi.org/10.1016/j.fishres.2008.08.002

1063

1064

1065

1066

1067

Supplementary materials

1069

Grouped by amplitude, percentile tails (33.0%) — Time-in-state within events

1070

Figure S1: **Relative temporal composition of chromatophore events across amplitude groups.** For each amplitude category (SMALL, MEDIUM, LARGE), the proportion of frames spent in the expansion (green) versus contraction (red) phase was computed across all valid events (n = 2157). Percentages were calculated relative to the total duration of each event (expansion + contraction). This analysis corresponds to the same dataset shown in Fig. 5.

1076

Figure S2: Designs of experimental arenas. a) Arena for *Euprymna berryi* hatchlings. b) Rig for *Sepia officinalis*. Four adjustable LED lamps in yellow. Transparent tank in cyan. Structure in gray.

1079

1080 **Figure S3: Sepia filming setup**. a) Remotely controlled linear rails. b) Z-axis manual translation
1081 stage. c) Camera and lens. d) LED lights.

1082

1083

1084 **Figure S4: Visible implant elastomer tagging.** a) Sepia officinalis marked at the center of the
1085 mantle with tags of various colors in a row. b) Marked animal under fluorescent light. c) Fluorescent
1086 lamp mounted next to the camera on the filming rig. d) Footage captured by the high-resolution
1087 camera. Close-up on the blue fluorescent tag indicated by an arrow.

1088

1089

1090

Figure S5: Schematics of experimental setup with head fixation. The camera (green) is
positioned above the tank, which is filled with artificial seawater (blue). A constant flow of water
(indicated by arrows) ensures proper oxygenation. LEDs (yellow) surrounding the tank provide
uniform illumination. A bracket (black) supports a thick glass panel (gray) to minimize water distortion
during filming. The same bracket also holds a head-fixation mount (white) to stabilize the animal's
head.

1097

1098

1099

1100

Figure S6: Confocal image of *E. berryi*'s stained skin captured with ZEISS LSM 980. DAPI in
blue (nuclei), phalloidin in magenta (muscle fibers), and tubulin in green (nerve fibers). A single
chromatophore is highlighted (white box) to illustrate its cellular organization. Radial muscle fibers
(magenta) extend outward from the central pigment sacculus, with nuclei (blue) arranged in a rosette
at the base of each fiber. Tubulin-labeled nerve fibers (green) approach the muscles, suggesting
potential neuromuscular connections, though the precise contact points cannot be confirmed in the
absence of a neuromuscular junction marker.

1108

1109