Physical proximity, social familiarity, and acoustic environment modulate heart rate synchrony in real-world social interactions

Hanlu He[1*], Jeppe Høy Christensen[2], A. Josefine Munch Sørensen[2], Ivana Konvalinka[1]

[1*]Section for Cognitive Systems, Department of Applied Mathematics and Computer Science, Technical University of Denmark, Denmark.
[2]Eriksholm Research Centre, Oticon A/S,Denmark.

*Corresponding author(s). E-mail(s): hahea@dtu.dk;
Contributing authors: jych@eriksholm.com; jsnn@eriksholm.com; ivako@dtu.dk;

Abstract

Human social behaviour unfolds in complex real-world environments shaped by social and acoustic factors, yet markers of social engagement and connection remain elusive. Interpersonal physiological synchrony (IPS) has been proposed as one such marker, but its occurrence in everyday settings is not well established. We continuously measured heart rate, GPS, and acoustic environments from 72 participants across three multi-day trips to New York City, capturing naturalistic social behaviour. Heart rates synchronized when participants were in close physical proximity, indicating that shared environmental context was sufficient to elicit IPS. IPS was stronger among socially familiar peers, and context dependent, emerging during close-proximity interactions and joint attention to shared stimuli, but not dispersed interactions. IPS was also modulated by acoustic context, with low-to-medium sound pressure levels and medium-to-high signal-to-noise ratios enhancing synchrony. These findings demonstrate that IPS emerges in naturalistic social settings, and establish IPS as a marker of real-world social engagement.

Keywords: Interpersonal physiological synchrony, Social interaction, Acoustic environment, Real-world engagement

1 Introduction

Human social behaviour unfolds in complex, dynamic real-world environments, where individuals continuously coordinate their actions, perceptions, and internal physiological signals during shared experiences. One well-documented form of such coordination is interpersonal synchrony, which spontaneously emerges between people's movements and physiological signals when they interact with one another [1–4], when they passively perceive each other[5] or when they merely have the same sensory input[6–8]. For example, people unconsciously synchronize their strides when walking side-by-side [9], and audiences spontaneously synchronize their clapping following a theater or opera performance [10]. Similarly, interpersonal physiological synchrony (IPS) - simultaneous changes in heart rates or phase-coupling between people's periodic cardiac rhythms - accompanies behavioural coordination during face-to-face interactions between mothers and infants and among romantic or marital partners[9, 11, 12], as well as during shared coordinated experiences such as choir singing[13] or playful joint actions[14, 15]. IPS can emerge even without overt behavioural coordination, as shown by the synchronization of heart rhythms between performers and socially affiliated spectators during collective rituals such as fire-walking[16].

Despite widespread evidence for interpersonal synchrony across bodily signals and social contexts, the mechanisms underlying IPS and its potential functional role remain elusive, particularly within the complex, multisensory environments in which everyday social interactions occur. Prior research has consistently linked behavioural synchrony to social bonding and prosocial behaviour, showing that people like each other more, feel greater affiliation, and act more cooperatively after synchronizing their movements[17–21]. Similarly, IPS has been shown to be enhanced by social affiliation[22, 23]: during the aforementioned fire-walking ritual, only spectators and performers with pre-existing social ties exhibited heart-rate coordination, whereas unrelated spectator-performer dyads did not[16]; socially close pairs going through a high-intensity haunted house synchronized their heart rates more than less close or unaffiliated pairs[23]; and autonomic synchrony between individuals meeting on blind dates has been shown to predict subsequent mutual attraction[24].

IPS can also arise without direct social interaction or shared social context, such as when people independently watch the same films or listen to the same narratives[25, 26]. In such cases, neural and physiological alignment is driven primarily by shared sensory input, and is amplified by simultaneous attentional engagement. For instance, distracted participants exposed to narrative stimuli exhibit reduced inter-subject correlation of heart rates[26]. Co-presence, such as when individuals attend to the same movies at the same time, further amplifies IPS[5]. This is consistent with accounts proposing that IPS and neural synchrony may be markers of joint attention or collective engagement[27–31], which are enhanced by shared social context[32, 33].

However, whether and how IPS emerges in everyday real-world interactions, outside of controlled laboratory environments, remains largely unknown. Measuring engagement in naturalistic settings

is particularly important, yet challenging, as subtle disengagement is difficult to detect in social interactions and may indicate sensory difficulty. This issue is especially relevant for individuals with hearing loss or degraded sensory input, who often experience reduced listening engagement in complex sensory environments due to increased cognitive demands [34–36]. In the absence of overt behavioural synchrony, it remains unknown whether IPS emerges in low arousal, real-world settings, or is instead driven by simultaneous surges in heart rates during high arousal events. Investigating whether IPS emerges during spontaneous face-to-face social exchanges and natural episodes of joint attention is essential for understanding its role in real-world social behaviour. Finally, establishing whether IPS is modulated by social affiliation and by the quality of the sensory input - particularly the acoustic properties of the environment - is necessary for evaluating its utility as a marker of real-world social engagement.

To address these questions, we conducted a study in ecologically valid, real-world settings by continuously recording heart rates from groups of students engaged in various social activities across three separate 4-day trips to New York (Trip 1: N = 23, Trip 2: N = 24, Trip 3: N = 25), organized as part of Oticon's annual Audioexplorer events[37]. Across the three trips, continuous monitoring generated a large volume of multimodal data, with approximately 293, 367, and 385 total hours recorded per day for Trips 1–3, respectively — equivalent to 14–17 hours of physiological, spatial, and acoustic data per participant per day. We investigated effects of physical proximity, activity type, social affiliation, and environmental noise on IPS. We categorized interactions into three distinct activity types: events involving shared sensory stimuli (e.g., attending the same lecture), face-to-face interactions (e.g., group games), and dispersed group interactions where the participants were together but not necessarily engaged in close interactions with all members of the group (e.g., having dinner across numerous small tables). In addition, we tracked participants' real-time GPS coordinates to quantify physical proximity, in order to investigate whether IPS emerges more broadly when people are together, as a result of the shared environment.

We further examined whether social affiliation modulates IPS by comparing heart rate synchrony between dyads with pre-existing social ties versus those without, while controlling for physical proximity. Given the cognitive demands of processing auditory stimuli in noisy environments - which are often associated with heightened cognitive load and reduced listening engagement [38–40] - we examined how such acoustic challenges encountered during various activities in New York City impacted IPS. Listening engagement may diminish when the perceived value of communication does not outweigh the effort required to understand the other person. To capture the acoustic context, we employed Oticon's hearing aids attached to participants' clothing, enabling real-time recording of environmental sound features. We then investigated whether elevated sound pressure levels (SPL) and lower signal-to-noise ratios (SNR) in the sound environment, which may contribute to disengagement, attenuate IPS.

Specifically, we hypothesized that significant levels of IPS, operationalized using measures of inter-subject correlation of heart rates (ISC-HR)[26], emerge when people are in close physical proximity, and that IPS is modulated by the type of activity people engage in. We expected IPS to be higher during face-to-face interactions involving reciprocal exchange of social signals, as well as during joint attention episodes to shared sensory stimuli, compared to dispersed interactions or physical separation. In addition, we hypothesized that social affiliation amplifies ISC-HR, through heightened shared engagement, resulting in higher IPS between socially affiliated individuals. Finally, we hypothesized that ISC-HR would be higher in auditory environments characterized by higher SNR and lower SPL, facilitating listening and social engagement. Thus, taking into account the physical and social proximity, the type of social activity and hence amount of shared sensory input, and the acoustic environment, we aimed to establish IPS more broadly as a socially and acoustically modulated marker of engagement in real-world environments.

2 Results

Across three separate trips to New York, groups of 23, 24, and 25 student participants, respectively, took part in scheduled activities and had free time to explore the city. Throughout each trip, participants wore hearing aids with microphones and Garmin wristbands synchronized to their mobile phones, which continuously measured features of the sound environment and heart rates at various locations (Figure 1). Data were collected continuously across four days for each trip. First, we examined whether inter-subject correlation of heart rates (ISC-HR), a proxy for IPS, was modulated by physical proximity between participants. GPS data were used to objectively determine the physical distance between individuals, and to assess ISC-HR when people were together in groups and in pairs. We further explored whether IPS was also modulated by social proximity, evaluated as pairs who knew each other beforehand and came to the trip in groups, while controlling for physical proximity. Finally, we investigated IPS across different planned activities to see whether IPS depended on the amount of shared social or environmental signals, such as settings with shared sensory information, close proximity face-to-face interactions, and dispersed interactions.

Fig. 1: **Study design and multimodal data collection during real-world interactions across three trips to New York City. (a)** Participants wore wearable sensors throughout four-day trips to New York City: Garmin wristbands to measure heart rate (HR), hearing aid microphones clipped to the collar to estimate the sound environment, and mobile phones to record GPS coordinates and synchronize sensor streams. **(b)** Example of a "sound-stress" map generated from hearing aid microphones, with colored dots indicating sound pressure levels (SPL) across locations in the city. **(c)** Example illustrations of the three main event types: a lecture with shared sensory input (stimulus-locked), outdoor group work in Central Park (close proximity face-to-face interaction), and a restaurant setting (dispersed social interaction). **(d)** Example of continuous HR time series from two participants over a 5-minute window. **(e)** Inter-subject correlation of heart rate (ISC-HR) was computed by correlating HR time series across participants in 5-minute epochs, illustrated here as correlation matrices across an event. Across three separate trips (n = 23, 24, and 25 participants), data were collected continuously from 9:00–22:00 each day. This enabled us to examine how interpersonal physiological synchrony (IPS), indexed by ISC-HR, varied as a function of shared sensory environments, face-to-face proximity, and more dispersed interactions, while controlling for the acoustic environment and physical proximity of participants.

2.1 People's heart rates synchronize when they are together in close physical proximity

To examine whether ISC-HR arises from mere physical co-presence, independent of social contexts [5], we used GPS data to identify periods when participants were physically close (within 20 m of each other) during daily activities (9:00–22:00) and segmented them into 5-minute windows. Across all three trips, ISC-HR was significantly higher during close-proximity group periods compared to shuffled controls, reflecting IPS between time-misaligned signals, with large effect sizes (Trip 1: $t(22) = 14.36$, $p = 1.18 \times 10^{-12}$, $d = 2.94$; Trip 2: $t(23) = 9.29$, $p = 2.98 \times 10^{-9}$, $d = 1.9$; Trip 3: $t(24) = 6.76$, $p = 5.41 \times 10^{-7}$, $d = 1.35$) (Figure 2a)).

We then examined ISC-HR at the dyadic level to better control for the physical proximity. Each pair was compared across three conditions: close proximity (within 20 m), far apart (more than 1 km), and time-misaligned shuffled controls. Repeated-measures ANOVAs revealed a consistent main effect of proximity across all trips (Trip 1: $F(2, 44) = 7.21$, $p = 0.002$, partial $\eta^2 = 0.18$; Trip 2, $F(2, 184) = 20.48$, $p = 9.29 \times 10^{-9}$, partial $\eta^2 = 0.13$; Trip 3, $F(2, 88) = 8.92$, $p = 0.001$, partial $\eta^2 = 0.12$.). Post hoc tests confirmed that ISC-HR was reliably higher when pairs were in close proximity compared to time-misaligned shuffled controls (Trip 1: $t = 4.03$, $p_{\text{Bonf.}} = 0.002$, $d = 1.08$; Trip 2: $t = 6.26$, $p_{\text{Bonf.}} = 3.54 \times 10^{-8}$, $d = 1$; Trip 3: $t = 3.51$, $p_{\text{Bonf.}} = 0.003$, $d = 0.74$).

For comparisons between close and far conditions, Trips 2 and 3 showed significant differences (Trip 2: $t = 4.24$, $p_{\text{Bonf.}} = 1.61 \times 10^{-4}$, $d = 0.57$; Trip 3: $t = 3.4$, $p_{\text{Bonf.}} = 0.004$, $d = 0.79$), while for Trip 1, the difference was not significant ($t = 2.32$, $p_{\text{Bonf.}} = 0.09$, $d = 0.76$). Further, no significant differences emerged between the far and shuffled conditions (Trip 1: $t = -1.18$, $p_{\text{Bonf.}} = 0.75$, $d = -0.34$; Trip 2: $t = -2.3$, $p_{\text{Bonf.}} = 0.07$, $d = -0.35$; Trip 3: $t = 1.55$, $p_{\text{Bonf.}} = 0.38$, $d = 0.3$) (Figure 2b).

Together, these results demonstrate that heart rate synchrony reliably emerges when individuals are in close physical proximity, both at the group and dyadic level, suggesting that shared environmental input facilitates physiological synchrony in real-world environments.

(a)

(b)

Fig. 2: **Heart rate synchrony emerges when people are in close physical proximity. (a)** Group-level ISC-HR across all 5-minute segments when participants were physically together, defined using GPS proximity data during daily activities (9:00–22:00). Average ISC-HR values were significantly greater than those obtained from time-misaligned shuffled controls (see Section 4.4.2) across all three trips. **(b)** Pairwise ISC-HR comparisons under three physical proximity conditions: close (<20 m), far apart (>1 km), and time-misaligned shuffled control. Across all three trips, ISC-HR was significantly higher when individuals were in close proximity compared to shuffled controls, with moderate to large effect sizes. In Trips 2 and 3, ISC-HR was also significantly higher in close proximity compared to far proximity. No differences were observed between far and shuffled conditions across all trips. These findings show that IPS emerges between people in close physical proximity, both at the group and dyadic level. Significant differences are denoted with asterisks ($*: p < 0.05$, $**: p < 0.01$, $***: p < 0.001$).

2.2 ISC-HR is modulated by social familiarity

The effect of social familiarity — defined as whether participants knew each other prior to the trip — was examined to test whether ISC-HR was modulated by pre-existing relationships beyond physical co-presence. Pairs were categorized as familiar (participants who knew and worked with each other before the trip) or unfamiliar (participants who were strangers prior to the trip). To control for physical proximity, pairwise ISC-HR values were computed only for segments when both participants were physically together identified by GPS coordinates in the same group and then averaged across those group segments.

The results revealed significantly higher ISC-HR for familiar pairs compared to unfamiliar pairs across all three trips (Trip 1: $W = 3.45$, $p = 0.0029$, rank-biserial $r = 0.95$, Trip 2: $W = 4.68$, $p = 2.84 \times 10^{-6}$, $r = 0.32$, Trip 3: $W = 2.38$, $p = 0.018$, $r = 0.17$), as seen in Figure 3. These findings indicate that social familiarity enhances interpersonal heart rate synchrony even when controlling for physical proximity, suggesting that pre-existing relationships facilitate physiological alignment.

Fig. 3: **Comparison of ISC-HR between familiar and unfamiliar pairs across three trips.** Mean ISC-HR values are shown for familiar pairs (participants who knew each other prior to the trip) and unfamiliar pairs (participants with no prior relationship), averaged across all group settings. Across all three trips, familiar pairs exhibited significantly higher ISC-HR than unfamiliar pairs, indicating that social proximity—defined by pre-existing social familiarity—enhances physiological synchrony beyond the effect of physical co-presence. Significant differences are denoted with asterisks ($* : p < 0.05$, $** : p < 0.01$, $*** : p < 0.001$).

2.3 ISC-HR across different social contexts

To examine how different types of interactions relate to ISC-HR, events were first manually labeled based on their social and contextual characteristics. Because activities varied widely in structure and social engagement, they were grouped into three categories reflecting shared behavioral and attentional contexts: close-proximity interactions (e.g., group discussions during games or meals), where participants actively engaged in face-to-face interactions; stimulus-locked interactions (e.g., attending lecture presentations or shows), where everyone jointly attended to an external sensory input; and dispersed interactions (e.g., receptions or co-located activities without shared engagement), where participants were more scattered and interactions were brief (Figure 4).

Across all three trips, significant group-level ISC-HR was observed primarily during close-proximity and stimulus-locked events, but not during dispersed interactions. For example, in Trip 1, ISC-HR was significantly elevated compared to time-misaligned control during *Social Dinner 2*, where participants sat close together and engaged in conversation ($t = 4.8$, $p = 2.9 \times 10^{-4}$, $d = 1.24$), and during *Presentation 1*, where participants jointly attended a science lecture ($W = 9.2$, $p = 1.26 \times 10^{-8}$, $d = 2$). In contrast, no significant ISC-HR was observed for *Casual Reception 1*, where participants were scattered around a rooftop terrace ($t = 1.09$, $p = 0.31$, $d = 0.36$).

A similar pattern was found in Trip 2 and Trip 3, with presentation-based and interactive events in close proximity consistently eliciting high ISC-HR. For instance, in Trip 3, *Presentation 7* (joint attention to a shared stimulus) showed very strong HR synchrony ($t = 11$, $p = 1.02 \times 10^{-8}$, $d = 2.33$), as did *Group Game 2*, where participants engaged in intense group discussions to solve an engineering task ($t = 5.51$, $p = 1.7 \times 10^{-4}$, $d = 1.26$). In contrast, dispersed settings such as *Casual Dining 1–4* and *Casual Reception 1–2*, where participants were seated apart and loosely engaged, exhibited low and nonsignificant ISC-HR levels.

Importantly, these effects were not explained by differences in mean heart rate: across events, mean HR did not correlate with ISC-HR, indicating that synchrony reflects shared temporal fluctuations rather than global arousal levels (see S1. for control analyses). Full statistical details for all events are provided in the Supplementary Material (Figures S3–S5, Tables S1–S3).

Fig. 4: **ISC-HR across different social interaction contexts. (A)** Distribution of mean ISC-HR across participants during three types of interaction: close proximity interaction (e.g., group discussion or games), stimulus-locked interaction (e.g., attending a performance or lecture), and dispersed interaction (e.g., rooftop reception). ISC-HR was highest during close proximity interactions, intermediate during stimulus-locked interactions, and lowest during dispersed interactions. **(B)** Example events corresponding to the three categories. Each dot represents the mean ISC-HR of one participant with all others across 5-minute segments within the event, compared against a time-misaligned shuffled control generated by random pairing of time segments. Significant ISC-HR was observed during close proximity and stimulus-locked events, but not during dispersed interactions. Asterisks denote significant differences (Wilcoxon signed-rank test with Bonferroni correction; $*: p < 0.05$, $**: p < 0.01$, $***: p < 0.001$).

2.4 The effect of Sound environment on ISC-HR

We examined whether exposure to different sound environments, characterized by sound pressure levels (SPL) and signal-to-noise ratios (SNR), was associated with variations in physiological synchrony. To contextualize the variability in acoustic sensory experience across the study, we visualized SPL and SNR over time for each trip. Figure 5a shows moment-to-moment and day-to-day fluctuations in the sound environment participants naturally encountered, with colour-coded background

segments indicating when different social context event types occurred. Moreover, it shows that the sound environments were relatively consistent across the three trips.

SPL and SNR values were estimated using participants' hearing aids and categorized into low, medium, and high levels based on tertiles within participants. ISC-HR was modeled using a linear model including SPL level, SNR level, their interaction, and mean heart rate (HR) as a covariate (mean_ISC $\sim$ SPL_level $\times$ SNR_level + mean_HR). Inclusion of mean HR slightly improved explained variance (R^2 increased from 0.0265 to 0.0286), but the effect was small, indicating that general arousal contributed minimally to ISC-HR.

Examination of the model revealed a significant interaction between SPL and SNR, indicating that the relationship between acoustic clarity and synchrony depends on the overall sound level. The following results describe this interaction in detail (Figure 5b).

Low SPL (quiet environments): ISC-HR peaked at medium SNR, followed by high SNR, and was lowest for low SNR. Pairwise contrasts showed that medium SNR was significantly higher than low SNR ($\hat{\Delta} = -0.0697$, $SE = 0.0106$, $t(5886) = -6.59$, $p < 0.0001$) as well as high SNR ($\hat{\Delta} = -0.0410$, $SE = 0.0109$, $t(5886) = -3.77$, $p = 0.0005$). ISC-HR at high SNR was also significantly higher than low SNR ($\hat{\Delta} = 0.0287$, $SE = 0.0075$, $t(5886) = 3.83$, $p = 0.0004$).

Medium SPL (moderately loud environments): ISC-HR at high SNR was significantly higher than at low or medium SNR ($\hat{\Delta} = 0.02875\text{--}0.03086$, $SE = 0.00831\text{--}0.00903$, $t = 3.416\text{--}3.458$, $p = 0.0016\text{--}0.0019$), while low and medium SNR did not differ ($\hat{\Delta} = 0.00211$, $SE = 0.00832$, $t(5886) = 0.254$, $p = 1.000$).

High SPL (loud environments): ISC-HR values were generally lower across all SNR levels. Only the contrast between high and medium SNR reached significance ($\hat{\Delta} = -0.01818$, $SE = 0.00744$, $t(5886) = -2.443$, $p = 0.0437$); other contrasts were not significant (high–low: $p = 0.2264$; low–medium: $p = 1.000$).

This pattern suggests that low-to-medium SPL combined with sufficient acoustic clarity (medium-to-high SNR) supports stronger physiological alignment, whereas excessively loud environments are linked to reduced synchrony. In particular, the noisiest conditions produced ISC-HR levels comparable to those observed during dispersed interactions, indicating that such environments may coincide with less interactive or socially engaging moments.

(a)

(b)

Fig. 5: **Sound environments and their relationship to ISC-HR. a)** Median sound pressure level (left y-axis) and signal-to-noise ratio (right y-axis) in decibels for each trip (rows) and day (columns) across all participants estimated in 5-min windows. The shaded area around each value represents the interquartile range. Vertical shaded areas represent periods of different social interaction contexts (close proximity interaction = blue, stimulus locked = green, dispersed interaction = light blue). **b)** Predicted ISC-HR values across combinations of sound pressure level (SPL: low, medium, high) and signal-to-noise ratio (SNR: low, medium, high) estimated from a linear model including SPL, SNR, their interaction, and mean HR as a covariate. Points and error bars show estimated marginal means ± 95% CI for each SPL × SNR cell. Grey horizontal lines indicate mean ISC-HR observed in different event contexts (solid = close-proximity interaction, dashed = stimulus-locked, dot-dash = dispersed). Synchrony varied as a function of SPL and SNR: in quiet environments (low SPL), ISC-HR peaked at medium SNR; at moderate SPL, ISC-HR was strongest at high SNR, with no difference between low and medium SNR; in loud environments (high SPL), ISC-HR was generally lower, comparable to levels observed during dispersed interactions.

3 Discussion

In this study, we demonstrate that interpersonal physiological synchrony (IPS), as indexed by inter-subject correlation of heart rates (ISC-HR), emerges robustly in everyday social interactions and is modulated by both social and environmental factors. Across three multi-day trips, heart rate synchrony was consistently higher when participants were in close physical proximity, when they interacted with familiar peers, and during activities that fostered shared attention and engagement, such as group games and attending lectures. In contrast, synchrony was reduced during dispersed interactions where opportunities for reciprocal social exchange or shared sensory input were more limited. Moreover, the acoustic environment further modulated ISC-HR: favorable listening conditions (low-to-medium SPL and medium-to-high SNR) enhanced synchrony, whereas noisy conditions reduced it, suggesting that environmental listening conditions influence the degree of physiological alignment, potentially due to increased cognitive effort and attentional load. Importantly, these effects were replicated across the three studies (conducted across different years). These findings show that IPS is not restricted to high-arousal or controlled laboratory settings, but also emerges spontaneously in naturalistic, everyday settings, making it a potential marker of social engagement under typical, low-intensity social conditions.

IPS is a well-documented phenomenon that occurs when individuals engage in both direct and indirect interaction [5, 26, 41, 42]. However, participant interactions in most of studies investigating IPS have been confined to laboratory settings, potentially limiting the ecological validity of the findings; or high arousal settings, which may drive synchrony through large variations in heart rates [16, 23]. To better understand IPS in naturalistic contexts, we analyzed heart rate (HR) data collected via wearable sensors during three different multi-day real-world trips, capturing both scheduled activities and free-roaming exploration.

Our results demonstrate that people's heart rates synchronize when they are in close physical proximity. GPS-based measures of spatial distance revealed that both group- and dyad-level ISC-HR were higher when participants were in close physical proximity of each other, compared to time-misaligned control analyses. This pattern was robust across all three trips with different participants, suggesting that shared physical space alone can facilitate physiological coupling in real-world environments. However, the result was less robust across trips for the dyad-level close–far comparison. While dyads in Trips 2 and 3 showed higher ISC-HR when they were close together than when they were far apart, the difference was not significant for Trip 1. Notably, the effect sizes for the dyad-level (Trip 1: $d = 0.76$; Trip 2: $d = 0.57$; Trip 3: $d = 0.79$) comparisons were smaller than those observed at the group level (Trip 1: $d = 2.94$; Trip 2: $d = 1.9$; Trip 3: $d = 1.35$). This may be because close proximity at the dyadic level does not necessarily indicate that partners were interacting or processing the same information, whereas proximity at the group level more likely reflected participation in

a shared activity. This weaker effect in Trip 1 may thus be explained by lower effect size and the substantially smaller amount of usable data. Specifically, the number of exact matching pair segments with sufficiently long continuous data for the close versus far conditions was much lower in Trip 1 (1262 vs. 1163) compared with Trips 2 (9542 vs. 8682) and 3 (2904 vs. 3385).

Beyond physical proximity, social familiarity was also found to modulate IPS. Pairs who knew each other prior to the trip exhibited higher ISC-HR than unfamiliar pairs, while controlling for physical proximity, indicating that pre-existing familiarity enhances alignment. This effect may reflect greater shared engagement to common stimuli or enhanced face-to-face interaction among familiar individuals. Interestingly, physiological synchrony was robust enough to capture this effect, despite all participants (familiar and unfamiliar) being in close proximity, and hence exposed to common environmental stimuli.

Examining ISC-HR under different social contexts revealed that people's heart rates tend to synchronize during close-proximity interactions (e.g., group discussions) or when co-attending to common stimuli (e.g., watching a lecture or performance), but not during dispersed interactions (e.g., sitting scattered in a restaurant). The observed group ISC-HR during stimulus-driven events appears to align with previous laboratory research on listening to common narrative stimuli [26] and real-world studies on physiological entrainment to music during live concerts [43]. Our findings thus indicate the context-dependent nature of physiological synchrony, which could be related to differences in participants' levels of attentional engagement. For instance, during collaborative group activities, participants may have felt more motivated to engage in frequent, dynamic verbal exchanges compared to less interactive settings. This heightened reciprocity in interpersonal communication might have contributed to the observed higher degree of alignment in physiological responses.

Further, we examined how variations in the acoustic environment, characterized by SPL and SNR, were associated with the strength of IPS. The interaction between SPL and SNR indicated that the association between SNR and physiological synchrony depended on the overall loudness of the environment. In quieter settings (low SPL), ISC-HR was highest at medium SNR, while at moderate SPL ISC-HR was highest for high SNR. In louder environments (high SPL), ISC-HR was generally lower across all SNR levels, with values comparable to those observed during dispersed interactions. We speculate that this may indicate that very loud settings coincide with less interactive activities, or create conditions that make social engagement more challenging. Importantly, because our data are observational, we cannot determine whether these associations reflect a direct effect of the sound environment on physiological synchrony or whether they are driven by the types of activities that typically occur under different acoustic conditions. One limitation is that we did not measure cognitive effort or listening engagement directly, for example via experience sampling. Thus, while our findings show that SPL and SNR are associated with variations in IPS, the underlying mechanisms remain speculative.

In addition, the current study has demonstrated the broader applicability of ISC-HR as a method for quantifying interpersonal synchrony (IPS) beyond controlled laboratory settings or high arousal events. By utilizing wearable sensor data, we have shown the robustness of ISC-HR as a tool for assessing collective, socially and environmentally modulated physiological states. This approach opens new possibilities for studying social dynamics in real-world environments, potentially serving as a marker for monitoring social engagement during real-world interactions.

While the study's ecological validity is a key strength, it also presents with some inherent limitations. Firstly, there is an inevitable trade-off in data quality when conducting research in real-world settings, as HR data were collected using wristbands instead of the raw cardiac activity data typically obtained from electrocardiograms (ECG) in laboratory environments. The wristbands are less accurate and more susceptible to motion artifacts and variations in activity types. Additionally, optimal GPS tracking using mobile phones requires an unobstructed view of the sky [44], whereas in our case, the participants spent considerable time indoors. These limitations in data quality necessitated extensive pre-processing decisions (cleaning, re-sampling, and interpolation). The scarcity of prior literature addressing these methodological challenges made it particularly difficult to establish standardized methods for handling such data in real-world contexts.

There was also limited control over how and when the participants interacted with each other aside from having the timestamps of planned activities. Consequently, we could not precisely identify the type of interaction each participant was engaged in when in close physical proximity to others. This constraint made it challenging to distinguish between reciprocal interaction and stimulus locking as drivers of IPS, leaving us to make inferences and generalize based on event labels. Future research should specifically examine how different types of interactions — such as conversation, shared attention, or cooperative tasks — affect the strength of ISC-HR, to better understand the mechanisms driving physiological synchrony. Such investigations could help isolate the specific social and cognitive factors that contribute to physiological alignment.

The present findings highlight the potential of IPS as a scalable, ecologically valid index of shared engagement in everyday life. By combining wearable sensing, environmental acoustics, and social context mapping with measures of interpersonal synchrony, our approach provides an index of collective behaviour in real-world environments. Importantly, our results are robust, replicating across three different trips and studies. This framework has broad applicability for quantifying real-world social dynamics. Measures of physiological synchrony can be used to track moment-to-moment fluctuations in collective engagement, to study how social connection emerges and evolves across contexts such as education and collaboration, and to index listening effort and participation in individuals with sensory challenges such as hearing loss. More broadly, combining physiological coupling with environmental and behavioural data offers a pathway toward social sensing - a data-driven approach to

characterizing how humans co-regulate their physiological states within complex social and sensory environments.

4 Methods

Participants submitted their consent agreeing to the use of their anonymized data (i.e., no personal identifiers were available) for research purposes on aggregated levels (i.e. no single-case investigation are performed) and agreed that data could be stored on Oticon A/S-owned secure servers. In addition, participants gave specific consent permitting access to their GPS location and health data from Apple HealthKit. All data collection and storage were conducted in a 'privacy by design' manner in accordance with the General Data Protection Regulation (EU regulation 2016/679). The Danish Capital Region Scientific Ethical Committee found that this study was not considered a health research study, according to the Act on Research Ethics Review of Health Research Projects, and therefore did not require ethical approval, reference number: F-25048245 (https://www.nvk.dk/forsker/naar-du-anmelder/hvilke-projekter-skal-jeg-anmelde).

4.1 Participants

Participants were recruited as a part of three audio explorer challenge competition[37] hosted by Oticon A/S. They entered the competition in teams that consist of 2 to 5 people. The winning teams of the competition were rewarded with a study trip to New York. There were in total 72 participants (60 male, 12 female) across three trips (23 on trip 1, 24 on trip 2 and 25 on trip 3). Participants' age ranged from 19 to 32 years old (M = 23 years, s.d. = 2.09 years).

4.2 Study Protocol and Data Collection

Data were collected during three separate trips to New York City, each spanning four days. During each trip, participants followed a pre-planned schedule of activities in and around New York City, interspersed with designated free time. They were informed that their participation would contribute to the creation of a "sound-stress" map of New York City through the use of wearable sensors. Daily competitions encouraged participants to collect acoustic field data.

The data collected were categorized into three types: acoustic, physiological, and environmental data. Every 20 seconds, the hearing aids (Oticon Opn™ S) recorded estimates of the surrounding sound environment, while wristbands (Garmin VivoSmart 4) measured the instantaneous heart rate at each detected heartbeat using pulse plethysmography. Additionally, mobile phones (iPhone 7) stored GPS location data, and researchers maintained detailed timelines of events attended by the participants. The GPS coordinates and event timelines were cross-referenced to define periods of pair or group interaction by identifying moments when participants were in close proximity.

We selected the time frame of interest from 9:00 AM to 10:00 PM for each day of the trip, as this was when most activities occurred.

4.2.1 Acoustic Data

The hearing aids logged acoustic data, including SPLs and noise floor (NF) estimates. Unlike laboratory settings, where SNR calculations involve well-defined target signals and noise, SNR in real-world settings relies on estimates from the processed acoustic features sensed by hearing aid microphones. These include environmental sounds, mixed speech, and background noise sources [45, 46].

In this study, the hearing aids used low-pass infinite impulse-response filters with a time constant of 63 milliseconds to estimate SPL and NF. SNR was determined as the difference between the A-weighted (adjusted to mimic human perception of sound as we are not equally sensitive to all frequencies) for SPL and the NF estimate. The SPL represents the immediate sound environment, while the NF represents slower-varying background noise, derived using a 1-second attack and 0.5-second release time for dynamic changes [47]. Hearing aid microphones were clipped onto participants' clothing, around the collar area, for consistent data collection.

4.2.2 Heart Rate Data

The Garmin wristbands use data from optical sensors (pulse plethysmographs) to estimate heart rates in beats per minute (BPM). These sensors emit green light through the skin, which is reflected by red blood cells, allowing the wristband to detect changes in blood flow due to heartbeats. The BPM value provides an average heart rate over short intervals (approximately a few seconds), making the readings suitable for long-term tracking, but provide less precise recordings than electrocardiograms (ECG).

ECG, often considered the gold standard in clinical studies, records the electrical activity of the heart by measuring voltage changes on the skin surface, capturing details like P-waves, QRS complexes, and T-waves [48]. In contrast, the optical heart rate monitoring of the Garmin wristbands smoothed out small heart rate variations. This method is sensitive to motion artifacts [49] and the participant's physical activity, which can lead to inconsistent RR-intervals not necessarily caused by ectopic beats. As a result, heart rate variability (HRV) analyses were not possible with this dataset. For more details, see Garmin's official heart rate monitoring guide (link) and accuracy disclaimer (link).

4.3 Pre-Processing

4.3.1 GPS Data

First, location logs with invalid coordinates were filtered out. Points within the Greater New York Area were identified based on predefined latitude (40.0–45.0) and longitude (-75.0 to -72.0) ranges.

Time differences and spatial distances between consecutive points for each participant were calculated using the Haversine formula, enabling the derivation of speed metrics. Movement status was labeled as "moving" or "stationary" based on a speed threshold of 2 m/s, with stationary points retained for further analysis. Significant jumps (distance > 30 m within 20 seconds) were detected and interpolated to minimize data inconsistencies.

4.3.2 Heart Rate Data

The instantaneous HR data were resampled to 10 Hz and interpolated using the Piecewise Cubic Hermite Interpolating Polynomial (PCHIP) method, which yielded the lowest root mean squared error (RMSE) for wearable HR signals [50]. Each event, group period, and pair period was then segmented into 5-minute epochs.

The choice of a 5-minute window was based on systematic evaluation of different epoch lengths (3, 5, and 10 minutes) across the three datasets. Shorter windows were unsuitable due to the low sampling rate of the acoustic measures (three samples per minute), while longer windows risked obscuring short-term interpersonal dynamics and increasing noise. The 5-minute segmentation provided the best balance between temporal resolution, data quality, and participant inclusion—yielding consistent data retention across trips (91.8%, 87.7%, and 92.4%, respectively). This window thus allowed sufficient temporal sensitivity while ensuring comparability and robustness across events.

4.4 Analysis

4.4.1 Proximity Measures

Physical proximity. We assessed physical proximity at two levels: the group level and the pair level.

At the *group level*, we calculated the average interpersonal distance among all participants for each minute. Individuals who were consistently at least 1 km away from others were excluded. Group periods were defined as intervals where the average interpersonal distance among all participants was less than 100 m, with each period lasting a minimum of 30 minutes. This 100 m threshold was empirically motivated by the observed proximity patterns across the three trips: during known close-proximity events (e.g., meals, presentations), the mean group proximity ranged from 0.08–0.15 km, whereas during free-time periods, distances were considerably larger and more variable (0.97–2.45 km). The 100 m cutoff thus reliably distinguished structured group activities from dispersed, unstructured periods, while accounting for the typical spread of participants within shared environments (e.g., lecture halls or outdoor areas).

At the *dyad level*, we calculated the interpersonal distance between all possible participant pairs for each minute. Pairs were classified as being in *close proximity* if their distance was less than 20 m, and as *far apart* if their distance exceeded 1 km. Only periods lasting at least 15 minutes were included. The 20 m threshold was selected based on proximity distributions observed during events

where participants were known to be seated or interacting in restricted spaces (e.g., restaurants, small indoor venues). This threshold also accounts for the inherent horizontal position accuracy of smartphone GPS in urban environments, which typically ranges between 7–13 m for static recordings [44]. In other words, distances below approximately twice the GPS accuracy margin can be interpreted as reflecting genuine close physical proximity rather than measurement noise.

Social proximity Participants who entered the audiology challenge worked together in groups of 2 to 5 individuals. We defined those within the same team as familiar members and those outside the team as unfamiliar members. For the group periods where all participants were defined as physically close, we further labeled interactions based on team membership. Specifically, we categorized interactions within the group as familiar interactions and those involving members of different teams as unfamiliar interactions. This distinction allowed us to explore the relationship between physical and social proximity and assess how team membership influenced patterns of interaction during group settings.

4.4.2 Quantifying Interpersonal Heart Rate Synchrony

Computing ISC-HR Intersubject correlations (or correlated component analysis) have been previously used to measure the alignment of physiological signals across individuals [26, 27]. This method evaluates the degree of similarity between individual's neural activity patterns by computing the linear correlations between their response signals. Recently ISC has been extended to quantify the similarity of participants' heart rate modulations over time when they are stimulus-locked (e.g., listening to the same narrative), enabling the study of dynamic alignment in physiological states [26]. In this study we applied ISC to real-world heart rate data collected using wearable sensors, allowing us to assess the degree of synchrony in heart rate fluctuations between participants during naturalistic interactions and gain a deeper understanding of how physiological alignment unfolds in response to real-world stimuli and social dynamics. This method was chosen because it is simple and robust in capturing similarity in fluctuations between two signals, and easily applicable to real-world long-term recordings.

For each period of interest, we divided each participant's instantaneous heart rate (HR) time series into 5-minute segments. We then calculated ISC for each analysis level as follows:

- **Close Proximity Group Level Analysis:** For each participant's segment of HR time series, we computed the ISC with every other participant's time-aligned segments of HR time series and averaged the ISC values across all segments and with all participants. This resulted in a single ISC value for each individual.

- **Physically Close vs. Far Dyad Level Analysis:** We computed the ISC for pairs of participants who were either in close proximity or far apart. The ISC values for these pairs were averaged across the segments, yielding one ISC value for each pair combination.

- **Familiarity Analysis:** For periods of close physical proximity, we computed ISC between all familiar pairs (participants in the same team) and compared these with ISC values between unfamiliar participants and out-group pairs (each participant with participants from different teams). These ISC values were averaged over the segments to obtain a single ISC value per pair.

- **Event Level Analysis:** For each event, we computed the ISC between each participant and every other participant per segment, averaging the ISC over all segments and participant-to-all-other-participants values. This provided one ISC value for each participant per event.

Time-Misaligned Random Segment Pairing Control To assess the statistical significance of the ISC-HR results for each analysis, we implemented random segment pairing controls, in order to measure baseline heart rate synchrony levels by pairing segments between participants that occur at different times. Specifically:

- **Group-Level Analysis (GPS Proximity and Activity Events):** We randomly paired each participant's instantaneous HR segments with time-misaligned segments of all other participants. The number of pairings was the same as in the original data, i.e., the same number of segments they were found to be in close proximity. We then computed the ISC-HR for these random pairings and compared the results to the observed ISC-HR values from the original data to assess statistical significance [26].

- **Pair-Level Analysis:** For each pair of participants, we randomly paired their time-misaligned HR segments, ensuring that the number of randomly paired segments for each pair was the same as for the time-aligned analysis. The ISC values were averaged over the segments, and these random pairings were compared to the observed ISC-HR values to determine significance.

By comparing the ISC-HR values from the original data with those from the random segment pairing control data, we were able to assess whether the observed ISC-HR was statistically different from ISC-HR values obtained between time-misaligned HR time series.

4.5 Statistical approach

To evaluate interpersonal physiological synchrony (IPS) as indexed by inter-subject correlation of heart rate (ISC-HR), several levels of analysis were conducted, and statistical tests were chosen based on data characteristics and experimental design. Across all analyses, normality of differences was assessed using the Shapiro–Wilk test, and appropriate parametric or non-parametric tests were selected accordingly. Effect sizes and multiple comparison corrections are reported where applicable. All statistical analyses were conducted using R (version 2024.04.1) and Python (version 3.8).

Group-level ISC vs. Time-misaligned shuffled controls

To test whether ISC-HR exceeded chance levels arising from time-misaligned signals, we compared group-level ISC-HR against time-misaligned shuffled controls. Differences were first tested for normality using the Shapiro–Wilk test. If normally distributed, paired-sample t-tests were used; otherwise, Wilcoxon signed-rank tests were applied. Effect sizes were reported as Cohen's d for t-tests and rank-biserial r for Wilcoxon tests.

Dyadic ISC across proximity conditions

At the dyadic level, ISC-HR was computed for each pair under three physical proximity conditions: close (< 20 m), far (> 1 km), and shuffled controls. Normality of residuals was assessed using the Shapiro–Wilk test. Repeated-measures ANOVAs were conducted to evaluate the main effect of proximity. When the assumption of sphericity was violated, Greenhouse–Geisser correction was applied. Post hoc pairwise comparisons were conducted between all proximity conditions, with Bonferroni correction for multiple comparisons. Effect sizes are reported as partial η^2 for ANOVAs and Cohen's d for pairwise contrasts.

Social familiarity ISC-HR

To assess the effect of social familiarity on physiological synchrony, ISC-HR was compared between familiar (participants who knew each other prior to the trip) and unfamiliar pairs, controlling for physical proximity by including only segments where participants were in the same group. Normality of ISC-HR differences was assessed via Shapiro–Wilk tests. When normality assumptions were met, Welch's t-tests were used; otherwise, Wilcoxon rank-sum tests were applied. Effect sizes are reported as Cohen's d for t-tests and rank-biserial r for Wilcoxon tests.

Event-level ISC-HR analyses

To examine ISC-HR across different social contexts, events were categorized as close-proximity interactions, stimulus-locked interactions, and dispersed interactions. For each event, ISC-HR was averaged across all 5-minute segments for all participant pairs. Differences between observed ISC-HR and time-misaligned controls were assessed using paired t-tests if normality was satisfied, or Wilcoxon signed-rank tests otherwise. Effect sizes were reported as Cohen's d.

Modeling the Effects of Sound Environment on ISC-HR

For this analysis, we pooled data from all three trips to examine how the acoustic environment was associated with ISC-HR. For each group period, we calculated the segment-level (based on tertiles) median SPL and SNR values across all participants [51]. These values provided a summary of the sound environment participants experienced in 5-minute intervals when they were together.

Categorization thresholds were as follows: SPL — low: 38–67.49 dB, medium: 67.66–81.86 dB, high: 81.88–95.54 dB; SNR — low: -2.32–5.65, medium: 5.67–7.17, high: 7.19–17.86.

To test whether group-level ISC-HR was associated with the sound environment, we fit a series of linear models predicting the mean ISC per segment across participants. The models were designed to evaluate (1) the additive effects of SPL and SNR, (2) whether the interaction between SPL and SNR further improved model fit, and (3) whether mean heart rate (HR), as a proxy for general arousal, contributed additional explanatory power. Specifically, the models were:

$$\text{Model 1: mean\_ISC} \sim \text{SPL\_level} + \text{SNR\_level}$$

$$\text{Model 2: mean\_ISC} \sim \text{SPL\_level} * \text{SNR\_level}$$

$$\text{Model 3: mean\_ISC} \sim \text{SPL\_level} * \text{SNR\_level} + \text{mean\_HR}$$

Model comparison was performed using ANOVA (1) to assess improvements in fit between nested models. Model quality was also evaluated using Akaike Information Criterion (AIC) and changes in R^2.

Model	Res.Df	RSS	Df	F	Pr($>$F)
Model 1: $\sim$ SPL + SNR	5891	138.13	-	-	-
Model 2: $\sim$ SPL * SNR	5887	137.00	4	12.20	7.03×10^{-10} ***
Model 3: $\sim$ SPL * SNR + mean HR	5886	136.71	1	12.40	0.00043 ***

Table 1: ANOVA table comparing nested models predicting mean ISC-HR. Model 2 shows that including the SPL × SNR interaction significantly improves model fit over additive effects (Model 1). Adding mean HR (Model 3) further improves fit, but explains a smaller proportion of variance.

AIC values for the three models were: Model 1 = -5388.47, Model 2 = -5429.06, Model 3 = -5439.47. The marginal R^2 values were: Model 1 = 0.0185, Model 2 = 0.0265, Model 3 = 0.0286. These results indicate that the interaction between SPL and SNR accounts for additional variance in ISC-HR, whereas mean HR contributes minimally, suggesting that general arousal does not fully explain the observed effects of the sound environment.

Data availability

The dataset underlying this Article is available at https://osf.io/cswt2/overview?view_only=364e00ef2c4e4b17a34334d43cf5a3d8

Code availability

The code for replicating the analyses underlying this Article is available at https://osf.io/cswt2/overview?view_only=364e00ef2c4e4b17a34334d43cf5a3d8

Acknowledgements

This work is supported by the Carlsberg Foundation's Semper Ardens: Accelerate grant, "Self, other, and we: mechanisms and dynamics of self-other integration in social observation and interaction", CF22-1251.

Author Contribution

H.H., J.H.C., and I.K. designed the study. J.H.C. collected the data. H.H. performed data preprocessing and analysis with input from I.K., J.H.C., J.M.S. I.K. and H.H. wrote the manuscript. All authors contributed to interpretation of the results and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Ethics approval and consent to participate

Participants submitted their consent agreeing to the use of their anonymized data (i.e., no personal identifiers were available) for research purposes on aggregated levels (i.e. no single-case investigation are performed) and agreed that data could be stored on Oticon A/S-owned secure servers. In addition, participants gave specific consent permitting access to their GPS location and health data from Apple HealthKit. All data collection and storage were conducted in a 'privacy by design' manner in accordance with the General Data Protection Regulation (EU regulation 2016/679). The Danish Capital Region Scientific Ethical Committee found that this study was not considered a health research study, according to the Act on Research Ethics Review of Health Research Projects, and therefore did not require ethical approval, reference number: F-25048245 (https://www.nvk.dk/forsker/naar-du-anmelder/hvilke-projekter-skal-jeg-anmelde).

References

[1] Schmidt, R.C., O'Brien, B.: Evaluating the dynamics of unintended interpersonal coordination. Ecological Psychology **9**(3), 189–206 (1997)

[2] Schmidt, R.C., Richardson, M.J.: Dynamics of interpersonal coordination, 281–308 (2008)

[3] Koban, L., Ramamoorthy, A., Konvalinka, I.: Why do we fall into sync with others? interpersonal synchronization and the brain's optimization principle. Social Neuroscience **14**(1), 1–9 (2019)

[4] Hoehl, S., Fairhurst, M., Schirmer, A.: Interactional synchrony: signals, mechanisms and benefits. Social cognitive and affective neuroscience **16**(1-2), 5–18 (2021)

[5] Golland, Y., Arzouan, Y., Levit-Binnun, N.: The mere co-presence: Synchronization of autonomic signals and emotional responses across co-present individuals not engaged in direct interaction. PloS one **10**(5), 0125804 (2015)

[6] Mayo, O., Gordon, I.: In and out of synchrony—behavioral and physiological dynamics of dyadic interpersonal coordination. Psychophysiology **57**(6), 13574 (2020)

[7] Stuldreher, I.V., Thammasan, N., Erp, J.B., Brouwer, A.-M.: Physiological synchrony in eeg, electrodermal activity and heart rate reflects shared selective auditory attention. Journal of neural engineering **17**(4), 046028 (2020)

[8] Madsen, J., Parra, L.C.: Cognitive processing of a common stimulus synchronizes brains, hearts, and eyes. PNAS nexus **1**(1), 020 (2022)

[9] Ulzen, N.R., Lamoth, C.J., Daffertshofer, A., Semin, G.R., Beek, P.J.: Characteristics of instructed and uninstructed interpersonal coordination while walking side-by-side. Neuroscience letters **432**(2), 88–93 (2008)

[10] Néda, Z., Ravasz, E., Brechet, Y., Vicsek, T., Barabási, A.-L.: The sound of many hands clapping. Nature **403**(6772), 849–850 (2000)

[11] Feldman, R., Magori-Cohen, R., Galili, G., Singer, M., Louzoun, Y.: Mother and infant coordinate heart rhythms through episodes of interaction synchrony. Infant Behavior and Development **34**(4), 569–577 (2011)

[12] Levenson, R.W., Gottman, J.M.: Marital interaction: physiological linkage and affective exchange. Journal of personality and social psychology **45**(3), 587 (1983)

[13] Müller, V., Lindenberger, U.: Cardiac and respiratory patterns synchronize between persons during choir singing. PloS one **6**(9), 24893 (2011)

[14] Fusaroli, R., Bjørndahl, J.S., Roepstorff, A., Tylén, K.: A heart for interaction: Shared physiological dynamics and behavioral coordination in a collective, creative construction task. Journal of Experimental Psychology: Human Perception and Performance **42**(9), 1297 (2016)

[15] Noy, L., Levit-Binun, N., Golland, Y.: Being in the zone: physiological markers of togetherness in joint improvisation. Frontiers in human neuroscience **9**, 187 (2015)

[16] Konvalinka, I., Xygalatas, D., Bulbulia, J., Schjødt, U., Jegindø, E.-M., Wallot, S., Van Orden, G., Roepstorff, A.: Synchronized arousal between performers and related spectators in a fire-walking ritual. Proceedings of the National Academy of Sciences **108**(20), 8514–8519 (2011)

[17] Marsh, K.L., Richardson, M.J., Schmidt, R.C.: Social connection through joint action and interpersonal coordination. Topics in cognitive science **1**(2), 320–339 (2009)

[18] Wiltermuth, S.S., Heath, C.: Synchrony and cooperation. Psychological science **20**(1), 1–5 (2009)

[19] Cirelli, L.K., Einarson, K.M., Trainor, L.J.: Interpersonal synchrony increases prosocial behavior in infants. Developmental science **17**(6), 1003–1011 (2014)

[20] Launay, J., Tarr, B., Dunbar, R.I.: Synchrony as an adaptive mechanism for large-scale human social bonding. Ethology **122**(10), 779–789 (2016)

[21] Tomashin, A., Gordon, I., Wallot, S.: Interpersonal physiological synchrony predicts group cohesion. Frontiers in human neuroscience **16**, 903407 (2022)

[22] Baranowski-Pinto, G., Profeta, V., Newson, M., Whitehouse, H., Xygalatas, D.: Being in a crowd bonds people via physiological synchrony. Scientific reports **12**(1), 613 (2022)

[23] Andersen, M.M., Rosas, F.E., Schjoedt, U., Mediano, P.A., Price, H., Konvalinka, I., Clasen, M.: Scared together: Heart rate synchrony and social closeness in a high-intensity horror setting. Emotion (2025)

[24] Prochazkova, E., Sjak-Shie, E., Behrens, F., Lindh, D., Kret, M.: Physiological synchrony is associated with attraction in a blind date setting. Nature Human Behaviour **6**(2), 269–278 (2022)

[25] Ardizzi, M., Calbi, M., Tavaglione, S., Umiltà, M.A., Gallese, V.: Audience spontaneous entrainment during the collective enjoyment of live performances: physiological and behavioral measurements. Scientific reports **10**(1), 3813 (2020)

[26] Pérez, P., Madsen, J., Banellis, L., Türker, B., Raimondo, F., Perlbarg, V., Valente, M., Niérat, M.-C., Puybasset, L., Naccache, L., et al.: Conscious processing of narrative stimuli synchronizes heart rate between individuals. Cell Reports **36**(11) (2021)

[27] Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., Malach, R.: Intersubject synchronization of cortical activity during natural vision. science **303**(5664), 1634–1640 (2004)

[28] Dmochowski, J.P., Sajda, P., Dias, J., Parra, L.C.: Correlated components of ongoing eeg point to emotionally laden attention–a possible marker of engagement? Frontiers in human neuroscience **6**, 112 (2012)

[29] Cohen, S.S., Madsen, J., Touchan, G., Robles, D., Lima, S.F., Henin, S., Parra, L.C.: Neural engagement with online educational videos predicts learning performance for individual students. Neurobiology of learning and memory **155**, 60–64 (2018)

[30] Dikker, S., Wan, L., Davidesco, I., Kaggen, L., Oostrik, M., McClintock, J., Rowland, J., Michalareas, G., Van Bavel, J.J., Ding, M., *et al.*: Brain-to-brain synchrony tracks real-world dynamic group interactions in the classroom. Current biology **27**(9), 1375–1380 (2017)

[31] Rai, L.A., Lee, H., Becke, E., Trenado, C., Abad-Hernando, S., Sperling, M., Vidaurre, D., Wald-Fuhrmann, M., Richardson, D.C., Ward, J.A., et al.: Delta-band audience brain synchrony tracks engagement with live and recorded dance. Iscience **28**(7) (2025)

[32] Shteynberg, G., Hirsh, J.B., Apfelbaum, E.P., Larsen, J.T., Galinsky, A.D., Roese, N.J.: Feeling more together: group attention intensifies emotion. Emotion **14**(6), 1102 (2014)

[33] Schiano Lomoriello, A., Sessa, P., Doro, M., Konvalinka, I.: Shared attention amplifies the neural processing of emotional faces. Journal of cognitive neuroscience **34**(6), 917–932 (2022)

[34] Heffernan, E., Coulson, N.S., Henshaw, H., Barry, J.G., Ferguson, M.A.: Understanding the psychosocial experiences of adults with mild-moderate hearing loss: An application of leventhal's self-regulatory model. International Journal of Audiology **55**(sup3), 3–12 (2016)

[35] Davis, H., Schlundt, D., Bonnet, K., Camarata, S., Bess, F.H., Hornsby, B.: Understanding listening-related fatigue: Perspectives of adults with hearing loss. International Journal of Audiology **60**(6), 458–468 (2021)

[36] Hornsby, B.W., Naylor, G., Bess, F.H.: A taxonomy of fatigue concepts and their relation to hearing loss. Ear and hearing **37**, 136–144 (2016)

[37] Demant A/S: Audio Explorers Program. https://www.demant.com/jobs-and-career/audio-explorers. Accessed: 2025-11-22

[38] Pichora-Fuller, M.K., Kramer, S.E., Eckert, M.A., Edwards, B., Hornsby, B.W., Humes, L.E., Lemke, U., Lunner, T., Matthen, M., Mackersie, C.L., *et al.*: Hearing impairment and cognitive energy: The framework for understanding effortful listening (fuel). Ear and hearing **37**, 5–27 (2016)

[39] Herrmann, B., Johnsrude, I.S.: A model of listening engagement (mole). Hearing Research **397**, 108016 (2020)

[40] Peelle, J.E.: Listening effort: How the cognitive consequences of acoustic challenge are reflected in brain and behavior. Ear and Hearing **39**(2), 204–214 (2018) https://doi.org/10.1097/AUD.0000000000000494

[41] Palumbo, R.V., Marraccini, M.E., Weyandt, L.L., Wilder-Smith, O., McGee, H.A., Liu, S., Goodwin, M.S.: Interpersonal autonomic physiology: A systematic review of the literature. Personality

and social psychology review **21**(2), 99–141 (2017)

[42] Behrens, F., Snijdewint, J., Moulder, R., Prochazkova, E., Sjak-Shie, E., Boker, S., Kret, M.: Physiological synchrony is associated with cooperative success in real-life interactions. Scientific reports **10**(1), 19609 (2020)

[43] Tschacher, W., Greenwood, S., Egermann, H., Wald-Fuhrmann, M., Czepiel, A., Tröndle, M., Meier, D.: Physiological synchrony in audiences of live concerts. Psychology of aesthetics, creativity, and the arts **17**(2), 152 (2023)

[44] Merry, K., Bettinger, P.: Smartphone gps accuracy study in an urban environment. PloS one **14**(7), 0219890 (2019)

[45] Bradley, J., Reich, R.D., Norcross, S.: On the combined effects of signal-to-noise ratio and room acoustics on speech intelligibility. The Journal of the Acoustical Society of America **106**(4), 1820–1828 (1999)

[46] Christensen, J.H., Saunders, G.H., Porsbo, M., Pontoppidan, N.H.: The everyday acoustic environment and its association with human heart rate: evidence from real-world data logging with hearing aids and wearables. Royal Society open science **8**(2), 201345 (2021)

[47] Kates, J.M.: Digital Hearing Aids. Plural Publishing, San Diego, CA (2008)

[48] Crawford, J., Doherty, L.: Practical Aspects of ECG Recording. M&K Update Ltd, London, UK (2012)

[49] Bent, B., Goldstein, B.A., Kibbe, W.A., Dunn, J.P.: Investigating sources of inaccuracy in wearable optical heart rate sensors. NPJ digital medicine **3**(1), 18 (2020)

[50] Rodrigues, E., Lima, D., Barbosa, P., Gonzaga, K., Guerra, R.O., Pimentel, M., Barbosa, H., Maciel, Á.: Hrv monitoring using commercial wearable devices as a health indicator for older persons during the pandemic. Sensors **22**(5), 2001 (2022)

[51] Christensen, J.H., Rumley, J., Gil-Carvajal, J.C., Whiston, H., Lough, M., Saunders, G.H.: Predicting individual hearing-aid preference from self-reported listening experiences in daily life. Ear and hearing, 10–1097 (2024)