The human hippocampus is involved in auditory short-term memory trace formation

Xiuyuan Liang[1,4,5,†], Zihao Guo[1,†], Dong Zhang[2], Yiming Zhang[2], Aobo Chen[2], Lanlan Wang[3], Xiang Liu[3], Wang Zhang[2], Yinbao Qi[2], Xiaorui Fei[2], , Ming Wang[1], Huawei Li[6], Lin Chen[1,6,*], Ruobing Qian[2,*]

[1]Auditory Research Laboratory, School of Life Sciences, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230027, China

[2]Department of Neurosurgery, The First Affiliated Hospital, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, China

[3]Department of Neuroelectrophysiology, The First Affiliated Hospital, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, China

[4]National Engineering Research Center of Speech and Language Information Processing, University of Science and Technology of China, Hefei 230027, China

[5]Interdisciplinary Research Center for Linguistic Sciences, School of Humanities and Social Sciences, University of Science and Technology of China, Hefei 230027, China

[6]Affiliated Eye and ENT Hospital of Fudan University, Shanghai 200031, China

[†]These authors contributed equally to this work.

*Correspondence to: Lin Chen

Full address: Auditory Research Laboratory, School of Life Sciences, Division of Life Sciences and Medicine,University of Science and Technology of China, Hefei 230027, China

E-mail: linchen@ustc.edu.cn

*Correspondence to: Ruobing Qian

E-mail: qianruobing@ustc.edu.cn

Full address: Department of Neurosurgery, The First Affiliated Hospital, University of Science and Technology of China, Hefei 230001, China

Abstract

The human hippocampus has been claimed to play an important role in long-term memory, or episodic memory, but its contribution to short-term memory remains debated. Here, we demonstrate human hippocampal involvement in auditory short-term memory (ASTM) trace formation. We used a classic oddball paradigm and intracranial recordings across various human brain areas. High frequency activities evoked by deviant stimuli in subjects indicate the generation of ASTM trace. Analysis of auditory response latencies across the hippocampus, insula, and temporal, parietal, and frontal lobes revealed the early processing of ASTM trace at a pre-attention stage. Granger causality analysis further showed that ASTM trace processing engages hierarchical cortical areas and clarified their interactions. Bottom-up signals flow from the insula and auditory temporal regions to the frontal lobe early on, followed by top-down information from the frontal lobe to the hippocampus, inferior temporal lobe, and insula at a later pre-attentive stage. Critically, Bayesian model selection of dynamic causal modeling revealed that deviant stimuli selectively enhanced the effectiveness of top-down connections to the hippocampus, thereby facilitating ASTM trace formation. These results provide compelling evidence that the human hippocampus contributes to auditory short-term memory at a pre-attention stage and suggests a challenge to entrenched beliefs in the classification of memory systems.

Keywords: auditory short-term memory trace; hippocampus; stereoencephalography (sEEG); high-frequency activity (HFA); Granger causality; dynamic causal modeling.

Introduction

The human hippocampus has been claimed to play a crucial role in human memory system especially long-term memory, also referred to as episodic memory[1], it comes with an equally important caveat that the hippocampus does not contribute to short-term memory[2]. The traditional/conventional view of hippocampal amnesia is that short-term memory is preserved despite the severe impairment in long-term memory[1,3,4]. For several years, a wide range of studies have established the medial temporal lobe, and hippocampus in particular, to be

critically associated with long-term memory processing. For instance, researchers studied

H.M., the well-known amnesic patient who had undergone temporal lobe resection, and

published observations of some types of preserved memory (motor skill and perceptual

learning) in the patient[5,6], as well as types of memory that depended on parts of the brain

other than the medial temporal lobe, in particular the hippocampus, or that did not depend on

conscious awareness were termed implicit (non-declarative) memory[7,8]. Previous view

mentioned that amnesia associated with hippocampal region damage is selective to

declarative memory, the ability to recall facts and events[9], and for H.M. the deficit includes

the complete absence of episodic memory, defined as the ability to remember specific

personal experiences[10]. In addition, many researchers have explored the underlying neural

mechanism of memory in which studies have implied that selective injury to the medial

temporal lobe leads to an isolated deficit in long-term memory[11,12]. Evidence from studies in

patients with amnesia due to hippocampal damage proposed that patients exhibit impairment

of conscious episodic encoding and retrieval[13]. Nevertheless, there is different view reports

that amnesic patients with bilateral lesions involving the hippocampus region suffered from

an impairment of short-term memory along with research develops[14,15]. Moreover, recent

visual study in human supports the view that the hippocampus does indeed have a role to play

in short-term memory[16] and study reported the role of cross-frequency coupling in cortico-

hippocampal activity in sequential auditory short-term memory (ASTM)[17].

Short-term memory describes the mental capacity to consciously hold information, such as

keeping a number in mind. Its neural substrate is the memory trace. For example, the active

representation of that number, through specific neuronal firing patterns in networks involving

the hippocampus and prefrontal cortex, constitutes the short-term memory trace. Thus, the

trace is the persistent neural state that enables the transient cognitive experience of short-term

memory. Short-term memory is a storage system that has an evolutionary survival advantage,

such as paying attention to limited but essential information excluding confounding factors

and ASTM is one of the important sensory-associated subsystems[18]. Short-term memory trace

is the physical neural basis for that function, the temporary, biological activity in our brain

that makes short-term memory possible. Mismatch negativity (MMN), first described about

40 years ago by Näätänen and his colleagues[19], is an index of automatic deviance detection. It

reflects cortical responses to deviant changes in the auditory system[20,21] and has been

regarded as a marker for detecting a deviant stimulus within the ASTM[22]. A variety of studies

have led to the understanding that the MMN is the outcome of an automatic comparison

process between a new, deviant stimulus and the memory trace formed by the sensory
representation of the standard stimuli within the short-term memory[23-26]. The MMN depends
on the presence of ASTM trace in the auditory cortex representing the repetitive aspects of
the preceding auditory events[20]. Thus, ASTM trace formation is indexed by the MMN
responses and deviance detection is mediated by a more stable short-term memory trace[27,28].

In the current study, we employed intracranial recordings, a technique affording high
spatial and temporal resolution, across various human brain regions, combined with a short-
term memory paradigm presented with a contrast pure tone auditory streams to explore
whether the human hippocampus contributes to ASTM trace formation. High frequency
activities (HFA, 70–150 Hz) elicited by the auditory stimuli is a reliable electrophysiological
correlate of underlying averaged spiking activity generated by the thousands of neurons that
are in the immediate vicinity of the recording electrodes[29-32]. This type of activity has been
linked to various cognitive processes, including attention, memory, and sensory processing.
Significant HFA of deviance detection evoked by the subjects indicate the generation of
ASTM trace. In our study, HFA results revealed that hippocampus was involved during
process of ASTM trace. Auditory response latencies recorded from the hippocampus, insula,
temporal lobe, parietal lobe, and frontal lobe showed the early processing of ASTM trace at a
pre-attention stage. Moreover, Granger causality analysis further demonstrated ASTM trace
is processed in hierarchical cortical areas, and detailed interactions of brain regions among
this memory trace process have been clarified/interpreted clearly. Specifically, bottom-up
auditory signals processed by the insula, and auditory regions of the temporal lobe and the
frontal lobe at an early stage, then auditory top-down information transmitted from the frontal
lobe to the hippocampus, the inferior temporal lobe, as well as insula at a relatively late pre-
attention stage of the auditory information processing. Although Granger causality
characterizes the temporal precedence and statistical dependencies of directional information
flow, it does not specify the underlying neurobiological mechanism. To identify the specific
neural mechanism, we employed Dynamic Causal Modeling (DCM). Critically, DCM results
revealed that deviant stimuli selectively enhance top-down connections from the frontal,
insula or temporal to the hippocampus, providing a mechanistic explanation for the
information flow that facilitates ASTM trace formation. In summary, our findings provide
converging evidence that the human hippocampus contributes to ASTM trace formation at a
pre-attentive stage through a specific frontal-hippocampal neural circuit that underlies the

formation of this transient memory trace, suggesting a new perspective that challenges
traditional classifications of memory systems.

Results

High frequency responses to auditory deviants across hippocampal and cortical networks.

To investigate the neural mechanisms underlying short-term memory trace formation, we
employed a classic auditory oddball paradigm combined with intracranial recordings across
multiple human brain regions. The hippocampus, a key structure within the medial temporal
lobe memory system, is critically involved in the integration and relay of auditory
information to downstream perceptual and cognitive regions[33-36]. Thus, for each recording
channel in the hippocampal region, we computed trial-averaged high-frequency activity
responses to both standard and deviant pure tones (Fig. 2). Our results revealed that
hippocampal channels displayed diverse response profiles, including both HFA
enhancements and high-frequency suppression (HFS). Specifically, responses to standard
stimuli were predominantly characterized by negative-going deflections (Fig. 2a, c), whereas
responses to deviant stimuli consistently exhibited positive deflections (Fig. 2e, g). To
identify channels exhibiting significant responses to auditory stimuli, a two-tailed cluster-
based permutation test incorporating threshold-free cluster enhancement (TFCE)[37] was
applied separately to standard and deviant trials for each channel. The test was performed
using 1000 sign-flipping randomizations. Channels with a corrected p-value below 0.05 for
either stimulus type were classified as responsive. Exemplar channels exhibiting statistically
significant responses to standard pure tones are shown in or deviant pure tones are shown in
Fig. 2b and Fig. 2d ($p < 0.05$), while exemplar channels exhibiting statistically significant
responses to deviant pure tones are presented in Fig. 2f and Fig. 2h ($p < 0.05$).

Furthermore, electrode pairs that evoked significant mismatched activity were categorized
according to whether the mismatch activity was enhanced or suppressed. We collected sEEG
data from 17 epilepsy patients (Table 1). To obtain a global description of the brain regions in
deviance detection, we examined the HFA responses to standard versus deviant tones across
intracranial channels. In the HFA analysis, 49 channels responded to tone responses in 10 out
of the 17 patients (Fig. 3), which illustrates the overall response profiles across all channels,

including both HFA and high-frequency suppression (HFS). Specifically, Significant HFS

responses were observed in 8 channels across 6 patients with and 41 channels across 10

subjects with HFA responses ($p < 0.05$, Fig. 3a, b). HFS in response to the stimuli at a latency

of 395 ms were recognized at the middle temporal gyrus (MTG), temporo-occipital brain area,

posterior inferior temporal gyrus, and anterior corona radiata a channel. In addition, 29

channels across 9 patients responded exclusively to deviant stimuli (Fig. 3c) whereas 12

channels across 7 patients responded to both standard and deviant tones (Fig. 3d).

Supplementary Figure 1 showed HFA responses across individual subjects. It is known that

studying HFA with sEEG recordings can provide insights into the neural bases of human

cognition that cannot be derived as easily from the study of BOLD effects with fMRI or from

the analysis of EEG[30]. HFS, a transient energy suppression in the gamma-band (40–150 Hz)

during the processing of sensory stimuli, which was first reported by Lachaux[38] and has close

relationship with attention[39]. We also observed widely distributed high frequency suppression

(HFS)[40], particularly in response to standard stimuli, which generally occurred later than high

frequency activity (HFA). This widespread HFS may reflect an energy saving mechanism in

the brain when processing predictable events. Given the known interaction between early

tone responses and later repetition suppression[41], such suppression may be mediated by

repetition suppression mechanisms.

Spatiotemporal responses of deviance processing

Auditory deviance elicited HFA responses across a distributed cortical network, with

response latencies following a systematic spatial order in individual subjects (Fig. S1). Mean

HFA/HFS response waveforms for key brain regions to standard and deviant pure tones,

along with the deviant-minus-standard difference waveforms were shown for multiple brain

regions (Fig. 4a).   Brain regions were defined as follows: the insula; temporal lobe

(encompassing Heschl's gyrus, superior temporal gyrus, and superior temporal sulcus);

frontal lobe (including the middle frontal gyrus, superior frontal gyrus, anterior cingulate

cortex, and frontal operculum); parietal lobe (comprising the postcentral gyrus, parietal

operculum, and supramarginal gyrus); posterior temporal lobe (the temporo-occipital part of

the middle temporal gyrus); inferior temporal lobe (including the inferior temporal gyrus,

fusiform gyrus, and inferior portion of the middle temporal gyrus); and the hippocampal area

(encompassing the hippocampus, parahippocampal gyrus, and amygdala). Additionally,

analysis of response latencies across these regions revealed a precise spatiotemporal gradient

(Fig. 4b). The earliest responses occurred in the insula and primary auditory cortex (~110

ms), followed by sequential engagement of the frontal and parietal cortices. The latest
responses were observed in the hippocampus and inferior temporal cortex, with latencies
extending up to ~438 ms. This ascending latency pattern demonstrates that deviance
processing propagates from early sensory to higher order associative regions.

Granger causality analysis revealed hierarchical information flow between the hippocampus and various brain regions.

To investigate the neural functional connectivity underlying auditory deviance processing, we
applied linear Granger causality (GC) analysis. The multivariate Granger causality analysis
(MVGC) Matlab toolbox used in the current study was designed for applications in
neuroscience[42], which implements numerical routines for calculating multivariate Granger
causality (MVGC) from time series data, both unconditional and conditional, in the time and
frequency domains[43]. The anatomical locations of these responsive brain regions (regions
detailed above) were mapped onto a common cortical space, as shown in the enlarged image
of the boxed area, with different colors denoting distinct brain clusters. The functional
connectivity is depicted in the heatmap (Fig. 4c), which displays the estimated difference in
Granger causal connection strengths between deviant and standard stimulus-evoked activities.
This GC analysis revealed a bottom-up information flow from the insula and auditory-related
regions to the frontal lobe, followed by the top-down information flow from the frontal lobe
back to the hippocampus, inferior temporal lobe, and insula at the pre-attentive stage (Fig. 4c).
This spatiotemporal profile establishes the hippocampus as a key node receiving feedback
signals within the pre-attentive processing. A magnified inset of the demarcated area is
presented, in which unique colors code for separate functional clusters and the directionality
(bottom-up versus top-down) of their causal influences (Fig. 5).

Dynamic causal modeling revealed stimulus gated enhancement of top-down hippocampal connectivity.

Bayesian Model Selection based on dynamic causal modeling yielded distinct winning
models for each subject, indicating individual variability in the optimal circuit architecture.
For Subject 3, Model 2 was identified as optimal (Fig. 6a, b). The key feature of this model is
that it included a top-down connection from the temporal cortex to the hippocampus, which
was significantly enhanced by deviant stimuli. For Subject 16, Model 7 was selected as the
winning model (Fig. 6c, d). This model is characterized by the inclusion of top-down
connections from both the frontal cortex and the insula to the hippocampus, with both

connections showing significant enhancement during deviant processing. Bayesian parameter estimation of the respective winning models confirmed that deviant stimuli selectively and significantly enhanced the effectiveness of the identified top-down connections to the hippocampus in both subjects. In summary, the DCM analysis identified distinct and subject-specific optimal models supporting ASTM trace formation. Despite individual differences in the precise cortical nodes driving the hippocampus, the core mechanistic finding was consistent: deviant processing is characterized by a selective enhancement of top-down effective connectivity directed at the hippocampus. This result provides a mechanistic neural explanation for the Granger causality findings, demonstrating that the observed top-down information flow is supported by a dynamic, stimulus-dependent increase in synaptic efficacy within direct hippocampal afferent pathways.

Intracranial correlates of the mismatch response.

Intracranial event-related potentials (iERPs) revealed a robust neural correlate of the auditory mismatch response. Significant deviant-minus-standard difference potential, which is an intracranial counterpart to the scalp MMN[44], was observed across 188 channels in all 17 patients (Fig. S2). To characterize the dominant response patterns, we performed principal component analysis (PCA) on the iERP waveforms. This analysis identified four distinct spatial-temporal response profiles (Fig. S3). The majority of responses corresponded to a positive-going mismatch response (MMR): Group 1 (121 channels, 6 patients), Group 2 (41 channels, 6 patients), and Group 4 (8 channels, 4 patients) all exhibited a significant positive MMR ($p < 0.05$). In contrast, a smaller subset of responses in Group 3 (18 channels, 3 patients) showed a significant negative deflection, consistent with the classic mismatch negativity (MMN) pattern ($p < 0.05$). Given that the same neural phenomenon can be recorded either as a positive or a negative iERP[30], these results demonstrate that the neural generators underlying deviance detection are broadly accessible in intracranial recordings and exhibit heterogeneous polarity, with a positive MMR being the predominant signature in our dataset.

Discussion

An ASTM trace constitutes the specific, sustained pattern of neural activity that physically represents and temporarily maintains information, serving as the neurobiological substrate of auditory short-term memory. In the current study, we employed a short-term memory paradigm featuring a contrastive auditory stream of pure tones to investigate the role of the human hippocampus in the formation of auditory short-term memory (ASTM) trace. Combined with the high spatiotemporal resolution approach, intracranial recordings used in this study allowed us to examine the electrophysiological activity across multiple brain regions. Using high frequency activity (HFA, 70–150 Hz) as a reliable correlate of localized neuronal firing, we found that deviance detection elicits significant HFA, indicating ASTM trace generation, and revealed hippocampus is involved in the process. Furthermore, the response latencies (110–438 ms) observed in the hippocampus, insula, temporal, parietal, and the frontal regions suggest that the initial process of the ASTM trace occurs at a pre-attention stage. Functional connectivity obtained from Granger causal analysis based on state-space (SS) model showed that ASTM trace processing engages a hierarchical cortical network, with early bottom-up information flow from the insula and auditory regions to the frontal cortex, followed by later top-down information flow from the frontal cortex to the hippocampus, inferior temporal cortex, and insula. Furthermore, to move beyond this descriptive account and identify the neurobiological mechanism underlying this information flow, we applied dynamic causal modeling. DCM results revealed a convergent neural mechanism: deviant stimuli selectively enhanced the effectiveness of top-down connections directed to the hippocampus, indicating that the hippocampus receives and integrates selectively amplified top-down signals during ASTM trace formation. In summary, these findings provide direct evidence that the human hippocampus contributes to ASTM trace formation at a pre-attention stage via a stimulus enhanced top-down pathway. Our study offers a more comprehensive model of ASTM trace formation and challenges entrenched definition of memory systems.

The hippocampus is involved in pre-attentive auditory short-term memory trace formation.

A conventional view in memory neuroscience posits that the human hippocampus plays a crucial role in long-term, or episodic memory but does not contribute to short-term memory[2]. Consistent with this view, previous research has established that the hippocampus participates in auditory processing, often related to later perceptual and cognitive processes[33-36]. For instance, prior research using electrophysiological techniques revealed that auditory processing engages a distributed network and that the hippocampal contribution to the scalp P300 is spatially restricted to the anterior temporal region[45]. Moreover, hippocampal functional engagement has been characterized in contexts of higher-order cognitive operations, including associative learning like auditory fear conditioning[46] and contextual updating at the end of a cognitive episode[47]. Anatomically and physiologically, the auditory association cortices send afferent inputs to the hippocampus, which in turn projects back to the primary auditory cortex and auditory association areas during auditory perception[48,49].

This entrenched view, however, is increasingly difficult to reconcile with findings that place the hippocampus with rapid, pre-attentive processing circuits. Notably, MEG study showed hippocampus forms a functional circuit with the ventromedial prefrontal cortex (vmPFC) during mismatch detection[50], suggesting its engagement occurs at earlier, more fundamental processing stages than previously theorized. The MMN has been regarded as the index of ASTM trace formation and deviance detection for decades, it is known to reflect a cortical comparison process between deviant stimuli and memory trace of standard stimuli[19-26]. While the scalp-recorded, low-frequency ERP and the low spatial resolution limits its ability to delineate the ASTM trace at the neuronal level. Here, we used sEEG, which provides excellent spatio-temporal resolution and unique access to deep limbic structures like the hippocampus[51], and measured HFA (70□150 Hz) as a more direct index of ASTM trace formation. HFA is a well-established signature of localized cortical firing[31] that offers greater temporal and spatial specificity than the integrated, scalp-recorded MMN for tracking rapid network dynamics[52]. In this study, we found that auditory deviant stimuli elicit widespread high-frequency activity (HFA: 70–150 Hz) with latencies ranging from 110 to 438 ms, indicating that ASTM trace formation occurs at a pre-attentive stage and that the hippocampus is involved in this process. This finding fundamentally challenges the traditional segregation of memory systems. The neural responses of hippocampus at a pre-

attentive stage indicate that it is not exclusive to long-term memory encoding, but is involved in the initial, rapid construction of ASTM trace. We therefore propose that ASTM trace formation is supported by a cortico-hippocampal network. This repositions the hippocampus with a newly defined and foundational role in short-term memory.

Hierarchical information flow during ASTM trace formation.

To clarify the specific mechanism of the hippocampus contributes to ASTM trace formation, we first characterized the spatiotemporal dynamics of information flow using Granger causality (GC) analysis based on a state-space model. Functional connectivity results obtained by GC analysis revealed that ASTM trace processing follows a precise spatiotemporal and hierarchical organization/sequence as follows (Fig. 4c and Fig. 5). An early bottom-up information flows from the insula and auditory-related regions to the frontal lobe, followed by later top-down information flow from the frontal lobe back to the hippocampus, inferior temporal lobe, and insula at the pre-attentive stage. This finding not only aligns with the core predictive coding principle, in which bottom-up signals convey prediction errors and top-down signals convey updated predictions[52-54], but also precisely positions the hippocampus as an integration node within the hierarchical network of ASTM trace formation.

The mechanism of hippocampus contributes to auditory short-term memory trace formation.

To further uncover the underlying mechanism, we applied Dynamic Causal Modeling (DCM) to analyze effective connectivity. Bayesian model selection showed that deviant stimuli selectively enhanced the effectiveness of top-down connections directed to the hippocampus, including temporal cortex to the hippocampus, the frontal cortex and the insula to the hippocampus (Fig. 6a and Fig. 6b). This indicates that hippocampal engagement in ASTM trace formation is not a mere epiphenomenon, but is governed by a stimulus-specific, top-down gain-control mechanism. The early engagement of the insula aligns with its well-established role in auditory processing. Converging evidence indicates that the insula is crucial for basic auditory detection, multisensory integration, as well as in allocating auditory attention and orienting to novel stimuli[55,56]. As shown by prior iEEG work, the insula is involved in automatic auditory processing[57], this aligns with our observation of its early engagement in the bottom-up stream. Additionally, the DCM based regulatory network features a direct top-down connection from the insula to the hippocampus. It also

incorporates bidirectional regulation between the frontal and temporal cortices and a direct
frontal-hippocampal pathway, together forming a multi-nodal, cascaded regulatory
architecture. It is also important to note that the parietal region participates in the hierarchical
circuit supporting ASTM trace formation, albeit via frontal and temporal relays to the
hippocampus rather than through direct connections (Fig. 6a, T model). This finding is
consistent with the view that STM has been considered to be the preserve of frontoparietal
networks. Thus, we suggest that the hippocampus does not operate in isolation from classic
STM system but is organized within an extended fronto-parietal-temporal-hippocampal
network. This helps understand the apparent paradox between the hippocampus's role in
long-term memory and its involvement in pre-attentive deviance detection. Hippocampus acts
as a hub that is recruited at the early stage of memory formation, binding significant sensory
events into ASTM under top-down control.

In summary, by combining the directional and temporal network dynamics revealed by GC
analysis with the connection-specific regulatory mechanism identified by DCM, this study
provides converging evidence that the human hippocampus contributes to ASTM trace
formation during a pre-attentive stage via a stimulus-modulated top-down pathway. These
findings challenge the classical dichotomy of memory systems and underscore the
hippocampus's active role in the early construction of short-term memory traces.

Materials and methods

Ethics statement

This study was approved by the Research Ethics Committee of the First Affiliated Hospital, University of Science and Technology of China (2024KY271). All participants provided written informed consent prior to taking part in the study. Neither the study procedures nor the analytical methods were preregistered in a time-stamped institutional registry before the research was conducted.

Participants

Stereoelectroencephalography (sEEG) data were obtained from 17 patients (6 males, 11 females) with drug-resistant epilepsy who were being evaluated as potential candidates for resective surgery. Data collection took place at the First Affiliated Hospital, University of Science and Technology of China. As part of their presurgical assessment, each patient underwent invasive sEEG monitoring. The selection of regions for electrode implantation was determined solely by clinical criteria. All participants were implanted with intracranial depth sEEG electrodes (0.8 mm in diameter, with a 3.5 mm center-to-center interelectrode distance) to help localize the epileptogenic zone. Every participant was right-handed, had normal cognitive function, and normal hearing. Written informed consent was obtained from all subjects prior to enrollment. The experimental protocol received approval from the Institutional Review Board of the University of Science and Technology of China and the First Affiliated Hospital.

Task paradigm

Stimuli in the current study consisted of pure tones generated using Audition 3.0 (Adobe Systems Inc., Mountain View, CA, USA). The auditory contrast was formed by a sequence of 300 ms pure tones, with a 500 Hz tone serving as the frequent standard stimulus and a 700 Hz tone as the infrequent deviant stimulus (**Figure 1**). All stimuli were delivered in random order via headphones using E-Prime (Psychology Software Tools, Inc., USA). Participants were instructed to ignore the auditory stimuli and instead focus on watching a muted movie of their choice with subtitles.

Data acquisition

For each participant, structural MRI scans were acquired prior to electrode implantation, and post-implantation CT scans were obtained afterward. sEEG signals were recorded using a 128/256-channel Neurofax EEG-1200c amplifier/digitizer system (Nihon Kohden, Japan) via implanted depth electrodes. Recording settings included a high-pass filter with a 0.01 Hz cut off, a 50 Hz notch filter, and a sampling rate of either 2000 Hz or 5000 Hz. Electrodes located on the inner surface of the skull served as the ground and reference.

Preprocessing and Electrode localization

Our sEEG data was imported to MATLAB using FieldTrip toolbox[58] and resampled to 1000 Hz. Event onset markers were read from pre-allocated trigger channels. The data were then re-referenced to the mean of the mastoid electrodes, and unused channels were removed. Segments containing epileptiform activity or abnormal fluctuations were manually rejected. The electrode localization was based on the FieldTrip protocol[59] with an adaptation for Windows, in which post-implantation CT images was co-registered to pre-implantation MRI images using SPM 12[60]. The images were subsequently normalized to the MNI-152 template space using SPM12[61], and electrode positions were determined according to the FieldTrip pipeline.

High frequency activity (HFA) analysis

Line noises at 50 Hz, 100 Hz, and 150 Hz was removed using a notch filter. The continuous data were segmented into 900 ms epochs (□200 ms to 700 ms). Trials were rejected if the amplitude exceeded 5 SD standard deviations from the mean for more than 25 consecutive ms, or if the power spectral density exceeded 5 standard deviations for more than 6 consecutive Hz[62]. Eligible trials were re-referenced to a neighboring channel for HFA analysis. HFA signals from –100 ms to 600 ms were extracted for further processing. Preprocessed data were band-pass filtered into eight 10-Hz-wide bands ranging from 70 to 150 Hz using a zero-phase Hamming windowed FIR filters (order 138)[63]. The envelope of each narrowband signal was extracted by taking the absolute value of the analytic signal from the Hilbert transform, and each amplitude time series was divided by its mean value[64]. For channels exhibiting a significant linear trend (R-square > 0.3), the trend was removed. Each channel's baseline period was removed and normalized by dividing the standard deviants of all trials of the baseline period. Auditory stimulus-evoked neural activity exhibits both

enhancements and suppressions, which are spatiotemporally distinct and originate from separate neural mechanisms. Previous studies on auditory cortical responses have consistently demonstrated that the brain displays both increased and decreased activity following sound stimulation, with these opposing responses being generated by different neural processes. It is therefore methodologically necessary, both mathematically and neurobiologically, to model and analyze these responses separately. Notably, stimulus-induced suppression shows characteristic spatiotemporal patterns that are significantly different from those of enhancement, further supporting their generation by distinct neural circuits. Evidence from auditory response research confirms that enhancement and suppression reflect dissociable neurophysiological processes. Thus, independent treatment of these responses is essential for both quantitative and neurobiological interpretations.

Tone response and deviance detection

To identify channels with significant responses to auditory stimuli, a two-tailed cluster-based permutation test with threshold-free cluster enhancement[37] and 1000 sign-flipping randomizations[65] was applied separately to standard and deviant trials for each channel. Channels with a corrected p-value below 0.05 either stimulus type were considered responsive. Channels displaying abnormal waveforms or epileptic activity were subsequently excluded through manual review. Then, in order to assess whether a channel was involved in deviance detection, a two-tailed cluster-based permutation test with threshold-free cluster enhancement[37] and 1000 randomizations was applied to compare trial categories (standard vs. deviant) for each channel. Channels with a corrected p-value less than 0.05 were considered significant for deviance detection. Additional manual rejection was applied to exclude channels with aberrant signals or association with epileptic activity. The latency of high-frequency activity (HFA) responses was estimated using a fractional area method[66]. The procedure began by identifying the time point of maximum amplitude within the response window as the initial seed point. The significant cluster was expanded bilaterally from this seed until the amplitude dropped below 20% of the seed value or reached the epoch boundaries. The latency was defined as the center of mass of the resulting suprathreshold cluster.

Event-related potential analysis and clustering

Event-related potentials (ERP) were derived from the same trials used in the high-frequency activity (HFA) analysis, prior to re-referencing to neighboring channels. The ERP signals

were filtered with a zero-phase FIR filter (order 18, 1–20Hz), segmented into 700 ms epoches

(–100 ms to 600 ms relative to stimulus onset) and baseline-corrected by subtracting the

mean amplitude during the –100 ms to 0 ms interval. For channels exhibiting a significant

linear trend (R-square > 0.3), detrending was applied. Similar to the HFA analysis, a two-

tailed cluster-based permutation test incorporating threshold-free cluster enhancement

(TFCE) was conducted for each electrode shaft to identify channels with significant

responses to deviant stimuli. Using principal component analysis (PCA) in line with

Blenkmann et al.[62], we identified groups of event-related potential (ERP) responses

exhibiting consistent temporal patterns across channels and subjects. This approach also

enabled the mitigation of polarity reversal effects. The analysis proceeded as follows: first, a

difference waveform was computed by subtracting the standard stimulus response from the

deviant stimulus response. PCA was then applied to this difference signal, and the first four

components capturing the highest channel weights were retained. Channels were grouped

based on their highest component weight, those predominately weighted on the first

component were clustered together, followed by those associated with the second component,

and so forth for each subsequent component.

Granger Causality Analysis

Wiener-Granger causality (G-causality, GC) is a statistical notion of causality applicable to

time series data, whereby cause precedes, and helps with prediction[67,68], which has been

developed to a powerful method for identifying information transfer or directed functional

("causal") connectivity in neural time series data[69,70]. It is defined in both time and frequency

domains, and allows for the conditioning out of common causal influences. Originally

developed in the context of econometric theory, it has since achieved broad application in the

neurosciences and beyond[71]. G-causality says that a variable X "G-causes" another variable

Y if the past of X contains information that helps predict the future of Y, over and above the

information already in the past of Y itself (and in the past of other "conditioning" variables Z).

In general terms, when this condition is satisfied one can say that there is "information flow"

from X to Y. This is justified because G-causality is an approximation to transfer entropy (the

approximation is exact for Gaussian variables), which is a directed version of Shannon's

mutual information, a very general way of characterizing the statistical dependency or shared

information between two variables[72]. The multivariate Granger causality analysis (MVGC)

Matlab toolbox used in the current study was designed for applications in neuroscience[42],

which implements numerical routines for calculating multivariate Granger causality (MVGC)

from time series data, both unconditional and conditional, in the time and frequency

domains[43]. Since it has been recognized that a moving average (MA) component in the data

presents a serious confound to Granger causal analysis, as routinely performed via

autoregressive (AR) modeling[42]. Previous studies have solved this problem by demonstrating

that Granger causality may be calculated simply and efficiently from the parameters of a

state-space (SS) model[68]. Because SS models are equivalent to autoregressive moving

average models, Granger causality estimated in this fashion is not degraded by the presence

of a MA component. State-space Granger causal inference stands to significantly enhance our

ability to identify and understand causal interactions and information flow in the current

study[73]. We used the MVGC (Multivariate Granger Causality) Matlab toolbox for calculating

multivariate Granger causality from time series data[42]. Autoregressive moving average

(ARMA) process or, equivalently, finite order linear state-space (SS) have been applied in

this study[73]. We selected a time window of interest spanning –100 ms to 600 ms to perform

identical analyses. The model order was estimated using Akaike information criterion. The

significance of difference of G-causality between standard and deviant conditions was

estimated using a permutation test.

Dynamic causal modeling

To test whether the top-down information flow identified by Granger causality analysis

originates from direct neural connections that are specifically modulated by deviant stimuli,

and to elucidate the underlying neural mechanisms, we conducted Dynamic Causal Modeling

(DCM). Participants for DCM analysis were selected based on stereotactic

electroencephalography (sEEG) electrode coverage and neural response characteristics. Only

participants who exhibited a clear hippocampal mismatch response along with significant

activation in multiple other relevant brain regions (e.g., temporal, frontal, and insular cortices)

were included. Two participants (Subjects 3 and 16) met all criteria and were analyzed. DCM

is a Bayesian framework that employs biophysically constrained neural mass models. Neural

sources are modeled as local field potential generators, allowing for inference about the

mechanisms underlying evoked responses and quantification of how coupling between

sources is modulated by experimental stimuli. Based on prior evidence and the activation

patterns observed in this study, network nodes were defined for each participant. For Subject

3, nodes included the hippocampus, insula, and temporal, frontal, and parietal cortices. For

Subject 16, nodes included the hippocampus, insula, and temporal and frontal cortices. In line

with prior studies suggesting that optimal models of deviance processing incorporate both

bottom-up and top-down interactions, our model space included bidirectional connections

between the hippocampus and cortical regions. A model space of eight competing models

was constructed for each participant. Model 1 served as a baseline (null) model containing no

connections to or from the hippocampus[74-76]. Models 2 through 8 were generated by

systematically adding bidirectional connections between the hippocampus and the other

cortical nodes. Neural activity within a 0–600 ms post-stimulus time window was modeled

using a biophysically and spatially constrained local field potential forward model. Finally,

Bayesian model selection (BMS) was employed to compare the generative models. BMS

evaluates the free-energy bound on the log model evidence, $\ln p(y|m)$, which represents the

probability of the observed data y under each model m. This metric penalizes model

complexity to prevent overfitting. The model with the highest model evidence within the

tested set is identified as the "winning" model.

Data availability

522   The processed datasets generated and/or analysed in the current study are available from the

523   corresponding author upon reasonable request.

Authorship contribution statement

525   X.L., L.C., and R.Q. designed research; X.L. collected sEEG data; R.Q. and X.F. did the

526   electrode implantation surgery for drug-resistant epilepsy patients, Y.Q., D.Z., Y.Z., A.C.,

527   L.W., X.L. and W.Z. and provided fMRI and CT files, Z.G. and X.L. analyzed data; X.L. and

528   L.C. wrote the paper.

Funding

530   This study was supported by the National Natural Science Foundation of China (Grant

531   GG2070000587) and University of Science and Technology of China

532   (GrantWK3460000008).

Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Bird CM, Burgess N. The hippocampus and memory: insights from spatial processing. *Nature reviews neuroscience* **9**, 182-194 (2008).

2. Husain M. The human hippocampus contributes to short-term memory. *Brain : a journal of neurology* **147**, 2593-2594 (2024).

3. Warrington EK. Amnesia and the distinction between long-and short-term memory. *Journal of verbal learning and verbal behavior* **9**, 176-189 (1970).

4. Cave CB, Squire LR. Intact verbal and nonverbal short-term memory following damage to the human hippocampus. *Hippocampus* **2**, 151-163 (1992).

5. Corkin S. Acquisition of Motor Skill after Bilateral Medial Temporal-Lobe Excision. *Neuropsychologia* **6**, 255-& (1968).

6. Moscovitch M. Recovered Consciousness - a Hypothesis Concerning Modularity and Episodic Memory. *Journal of Clinical and Experimental Neuropsychology* **17**, 276-290 (1995).

7. Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychological review* **99**, 195-231 (1992).

8. Schacter DL. Implicit Memory: History and Current Status. *Journal of Experimental Psychology Learning Memory & Cognition* **13**, 501-518 (1987).

9. Cohen NJ, Squire LR. Preserved learning and retention of pattern-analyzing skill in amnesia: Dissociation of knowing how and knowing that. *Science* **210**, 207-210 (1980).

10. Steinvorth S, Levine B, Corkin S. Medial temporal lobe structures are needed to re-experience remote autobiographical memories: evidence from HM and WR. *Neuropsychologia* **43**, 479-496 (2005).

11. Scoville WB, & Milner, B. . Loss of recent memory after bilateral hippocampallesions. *J Neurol Neurosurg Psychiat* **20**, (1957).

12. Reber PJ. The neural basis of implicit learning and memory: a review of neuropsychological and neuroimaging research. *Neuropsychologia* **51**, 2026-2042 (2013).

13. Tulving E. Episodic memory: From mind to brain. *Annual review of psychology* **53**, 1-25 (2002).

14. Hannula DE, Tranel D, Cohen NJ. The long and the short of it: relational memory impairments in amnesia, even at short lags. *Journal of Neuroscience* **26**, 8352-8359 (2006).

15. Olson IR, Page K, Moore KS, Chatterjee A, Verfaellie M. Working memory for
conjunctions relies on the medial temporal lobe. *Journal of Neuroscience* **26**, 4596-
4601 (2006).

16. Daume J*, et al.* Control of working memory by phase–amplitude coupling of human
hippocampal neurons. *Nature* **629**, 393-401 (2024).

17. Griffiths TD*, et al.* Cross-frequency coupling in cortico-hippocampal networks
supports the maintenance of sequential auditory information in short-term memory.
PLOS Biology **22**,　(2024).

18. Cascella M, Al Khalili Y. Short-term memory impairment. In: *StatPearls [Internet]*).
StatPearls Publishing (2024).

19. Näätänen R, Gaillard AWK, Mäntysalo S. Early Selective-Antention Effect on
Evoked Potential Reinterpreted. *Acta Psychologica* **42**, 313-329 (1978).

20. Näätänen RN, Paavilainen P, Rinne T, Alho K. The mismatch negativity (MMN) in
basic research of central auditory processing: a review. *Clinical neurophysiology :*
official journal of the International Federation of Clinical Neurophysiology **118**,
2544-2590 (2007).

21. Partanen E, Kujala T, Naatanen R, Liitola A, Sambeth A, Huotilainen M. Learning-
induced neural plasticity of speech processing before birth. *Proceedings of the*
National Academy of Sciences of the United States of America **110**, 15145-15150
(2013).

22. Colin C, Radeau M, Soquet A, Demolin D, Colin F, Deltenre P. Mismatch negativity
evoked by the McGurk-MacDonald effect: a phonetic representation within short-
term memory. *Clinical Neurophysiology* **113**, 495-506 (2002).

23. Novak GP, Ritter W, Vaughan Jr HG, Wiznitzer ML. Differentiation of negative
event-related potentials in an auditory discrimination task. *Electroencephalography*
and clinical neurophysiology **75**, 255-275 (1990).

24. Näätänen R. The role of attention in auditory information processing as revealed by
event-related potentials and other brain measures of cognitive function. *Behavioral*
and brain sciences **13**, 201-233 (1990).

25. Cowan N, Winkler I, Teder W, Näätänen R. Memory prerequisites of mismatch
negativity in the auditory event-related potential (ERP). *Journal of experimental*
psychology: Learning, Memory, and Cognition **19**, 909 (1993).

26. Näätänen R, Alho K. Mismatch negativity-a unique measure of sensory processing in
audition. *International Journal of Neuroscience* **80**, 317-337 (1995).

27. Teichert T, Jedema H, Zhen Z, Gurnsey K. Complementary functional profiles of
mismatch responses mediated by adaptation and deviance detection point to two
distinct auditory short-term memory systems. *Journal of neurophysiology* **134**, 314-
336 (2025).

28. Tuomainen OT. Auditory short-term memory trace formation for nonspeech and
speech in SLI and dyslexia as indexed by the N100 and mismatch negativity
electrophysiological responses. *Neuroreport* **26**, 374-379 (2015).

29. Ray S, Maunsell JH. Different origins of gamma rhythm and high-gamma activity in
macaque visual cortex. *PLoS Biol* **9**, e1000610 (2011).

30. Lachaux JP, Axmacher N, Mormann F, Halgren E, Crone NE. High-frequency neural
activity and human cognition: past, present and possible future of intracranial EEG
research. *Prog Neurobiol* **98**, 279-301 (2012).

31. Rich EL, Wallis JD. Spatiotemporal dynamics of information encoding revealed in
orbitofrontal high-gamma. *Nature communications* **8**, 1139 (2017).

32. Watson BO, Ding M, Buzsaki G. Temporal coupling of field potentials and action
potentials in the neocortex. *The European journal of neuroscience* **48**, 2482-2497
(2018).

33. Bouton ME, Bolles RC. Conditioned Fear Assessed by Freezing and by the
Suppression of 3 Different Baselines. *Anim Learn Behav* **8**, 429-434 (1980).

34. Liberman T, Velluti RA, Pedemonte M. Temporal correlation between auditory
neurons and the hippocampal theta rhythm induced by novel stimulations in awake
guinea pigs. *Brain research* **1298**, 69-77 (2009).

35. Graham KS, Barense MD, Lee ACH. Going beyond LTM in the MTL: A synthesis of
neuropsychological and neuroimaging findings on the role of the medial temporal
lobe in memory and perception. *Neuropsychologia* **48**, 831-853 (2010).

36. Lee ACH, Yeung LK, Barense MD. The hippocampus and visual perception.
Frontiers in human neuroscience **6**, (2012).

37. Smith SM, Nichols TE. Threshold-free cluster enhancement: Addressing problems of
smoothing, threshold dependence and localisation in cluster inference. *NeuroImage*
44, 83-98 (2009).

38. Lachaux JP, *et al.* The many faces of the gamma band response to complex visual
stimuli. *NeuroImage* **25**, 491-501 (2005).

39. Lachaux JP, *et al.* Silence is golden: transient neural deactivation in the prefrontal
cortex during attentive reading. *Cerebral cortex* **18**, 443-450 (2008).

40. Lachaux J-P, Axmacher N, Mormann F, Halgren E, Crone NE. High-frequency neural
activity and human cognition: Past, present and possible future of intracranial EEG
research. *Progress in neurobiology* **98**, 279-301 (2012).

41. Eckert D, *et al.* Distinct interacting cortical networks for stimulus-response and
repetition-suppression. *Communications biology* **5**, 909 (2022).

42. Barnett L, Seth AK. The MVGC multivariate Granger causality toolbox: a new
approach to Granger-causal inference. *J Neurosci Methods* **223**, 50-68 (2014).

43. Seth AK. A MATLAB toolbox for Granger causal connectivity analysis. *J Neurosci*
Methods **186**, 262-273 (2010).

44. Ishishita Y, *et al.* Deviance detection is the dominant component of auditory
contextual processing in the lateral superior temporal gyrus: A human ECoG study.
Human brain mapping **40**, 1184-1194 (2019).

45. Nishitani N, *et al.* The role of the hippocampus in auditory processing studied by
event-related electric potentials and magnetic fields in epilepsy patients before and
after temporal lobectomy. *Brain : a journal of neurology* **122 (Pt 4)**, 687-707 (1999).

46. Xiao C, Liu Y, Xu J, Gan X, Xiao Z. Septal and Hippocampal Neurons Contribute to
Auditory Relay and Fear Conditioning. *Frontiers in cellular neuroscience* **12**, 102
(2018).

47. Fonken YM, Kam JWY, Knight RT. A differential role for human hippocampus in
novelty and contextual processing: Implications for P300. *Psychophysiology* **57**,
(2019).

48. Mehta AD*, et al.* Seizure propagation in a patient with musicogenic epilepsy. *Epilepsy*
& Behavior **14**, 421-424 (2009).

49. Chatani H*, et al.* Neuromagnetic evidence for hippocampal modulation of auditory
processing. *NeuroImage* **124**, 256-266 (2016).

50. Garrido MI, Barnes GR, Kumaran D, Maguire EA, Dolan RJ. Ventromedial prefrontal
cortex drives hippocampal theta oscillations induced by mismatch computations.
NeuroImage **120**, 362-370 (2015).

51. Peedicail JS*, et al.* Outcomes of stereoelectroencephalography exploration at an
epilepsy surgery center. *Acta Neurologica Scandinavica* **141**, 463-472 (2020).

52. Durschmid S*, et al.* Hierarchy of prediction errors for auditory events in human
temporal and frontal cortex. *Proceedings of the National Academy of Sciences of the*
United States of America **113**, 6755-6760 (2016).

53. Hsu YF, Le Bars S, Hamalainen JA, Waszak F. Distinctive Representation of
Mispredicted and Unpredicted Prediction Errors in Human Electroencephalography.
The Journal of neuroscience : the official journal of the Society for Neuroscience **35**,
14653-14660 (2015).

54. Recasens M, Gross J, Uhlhaas PJ. Low-Frequency Oscillatory Correlates of Auditory
Predictive Processing in Cortical-Subcortical Networks: A MEG-Study. *Sci Rep-Uk* **8**,
(2018).

55. Bamiou DE, Musiek FE, Luxon LM. The insula (Island of Reil) and its role in
auditory processing. Literature review. *Brain research Brain research reviews* **42**,
143-154 (2003).

56. Downar J, Crawley AP, Mikulis DJ, Davis KD. A multimodal cortical network for the
detection of changes in the sensory environment. *Nature neuroscience* **3**, 277-283
(2000).

57. Citherlet D*, et al.* Spatiotemporal dynamics of auditory information processing in the
insular cortex: an intracranial EEG study using an oddball paradigm. *Brain Struct*
Funct **225**, 1537-1559 (2020).

58. Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput*
Intell Neurosci **2011**, 156869 (2011).

59. Stolk A*, et al.* Integrated analysis of anatomical and electrophysiological human
intracranial data. *Nat Protoc* **13**, 1699-1723 (2018).

60. Studholme C, Hill DLG, Hawkes DJ. An overlap invariant entropy measure of 3D
medical image alignment. *Pattern Recognition* **32**, 71-86 (1999).

61. Ashburner J, Friston KJ. Unified segmentation. *NeuroImage* **26**, 839-851 (2005).

62. Blenkmann AO*, et al.* Auditory deviance detection in the human insula: An
intracranial EEG study. *Cortex* **121**, 189-200 (2019).

63. Mesgarani N, Cheung C, Johnson K, Chang EF. Phonetic Feature Encoding in Human
Superior Temporal Gyrus. *Science* **343**, 1006-1010 (2014).

64. Norman Y, Yeagle EM, Harel M, Mehta AD, Malach R. Neuronal baseline shifts
underlying boundary setting during free recall. *Nature Communications* **8**, 1301
(2017).

65. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation
inference for the general linear model. *NeuroImage* **92**, 381-397 (2014).

66. Luck SJ. Event-related potentials from the user's perspective. *Nature neuroscience* **9**,
463 (2006).

67. Granger CWJ. Investigating causal relations by econometric models and cross-
spectral methods. *Econometrica* **37**, 424-438 (1969).

68. Seth AK, Barrett AB, Barnett L. Granger Causality Analysis in Neuroscience and
Neuroimaging. *Journal of Neuroscience* **35**, 3293-3297 (2015).

69. Bressler SL, Seth AK. Wiener-Granger causality: a well established methodology.
NeuroImage **58**, 323-329 (2011).

70. Schreiber T. Measuring information transfer. *Physical review letters* **85**, 461 (2000).

71. Barnett L, Seth AK. The MVGC multivariate Granger causality toolbox: A new
approach to Granger-causal inference. *Journal of Neuroscience Methods* **223**, 50-68
(2014).

72. Barnett L, Barrett AB, Seth AK. Granger causality and transfer entropy are equivalent
for Gaussian variables. *Phys Rev Lett* **103**, 238701 (2009).

73. Barnett L, Seth AK. Granger causality for state-space models. *Physical review E,*
Statistical, nonlinear, and soft matter physics **91**, 040101 (2015).

74. Allen M*, et al.* Anterior insula coordinates hierarchical processing of tactile mismatch
responses. *NeuroImage* **127**, 34-43 (2016).

75. Oestreich LKL, Randeniya R, Garrido MI. Auditory white matter pathways are
associated with effective connectivity of auditory prediction errors within a fronto-
temporal network. *NeuroImage* **195**, 454-462 (2019).

76. Schlossmacher I*, et al.* Differential effects of prediction error and adaptation along the
auditory cortical hierarchy during deviance processing. *NeuroImage* **259**, 119445
(2022).

Figure 1

a

b

n=29, s=12

a High Frequency Suppression (HFS)

b High Frequency Activity (HFA)

c HFA with Standard Non-responsive

d HFA with Standard Responsive

Figure 4

Figure 6