The Human Hippocampus Is Involved in Auditory

Short-Term Memory Trace Formation

Xiuyuan Liang[1,4,5,†], Zihao Guo[1,†], Dong Zhang[2], Yiming Zhang[2], Aobo Chen[2], Lanlan Wang[3], Xiang Liu[3], Wang Zhang[2], Yinbao Qi[2], Xiaorui Fei[2], , Ming Wang[1], Huawei Li[6], Lin Chen[1,6,*], Ruobing Qian[2,*]

[1]Auditory Research Laboratory, School of Life Sciences, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230027, China

[2]Department of Neurosurgery, The First Affiliated Hospital, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, China

[3]Department of Neuroelectrophysiology, The First Affiliated Hospital, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230001, China

[4]National Engineering Research Center of Speech and Language Information Processing, University of Science and Technology of China, Hefei 230027, China

[5]Interdisciplinary Research Center for Linguistic Sciences, School of Humanities and Social Sciences, University of Science and Technology of China, Hefei 230027, China

[6]Affiliated Eye and ENT Hospital of Fudan University, Shanghai 200031, China

[†]These authors contributed equally to this work.

*Correspondence to: Lin Chen

Full address: Auditory Research Laboratory, School of Life Sciences, Division of Life Sciences and Medicine,University of Science and Technology of China, Hefei 230027, China

E-mail: linchen@ustc.edu.cn

*Correspondence to: Ruobing Qian

E-mail: qianruobing@ustc.edu.cn

Full address: Department of Neurosurgery, The First Affiliated Hospital, University of Science and Technology of China, Hefei 230001, China

Abstract

The human hippocampus has been claimed to play an important role in long-term memory, or episodic memory. It has long been argued whether the hippocampus does contribute to short-term memory. Here, we demonstrate human hippocampus has been involved in auditory short-term memory (ASTM) trace formation. We used a classic oddball paradigm and intracranial recordings across various human brain areas. High frequency activities evoked by deviant stimuli in subjects indicate the generation of ASTM trace. Auditory response latencies recorded from the hippocampus, insula, temporal lobe, parietal lobe, and frontal lobe showed the early processing of ASTM trace at a pre-attention stage. Moreover, Granger causality analyses showed ASTM trace is processed in hierarchical cortical areas and interactions among this memory process have been clarified. Specifically, bottom-up auditory signals processed by the insula, and auditory regions of the temporal lobe and the frontal lobe at an early stage, then auditory top-down information transmitted from the frontal lobe to the hippocampus, the inferior temporal lobe, as well as insula at a relatively late pre-attention stage of the auditory information processing. These results provide evidence that the human hippocampus contributes to the short-term memory of auditory perception at a pre-attention stage and suggests a challenge to entrenched beliefs in the classification/definition of memory systems.

Keywords: auditory short-term memory trace; hippocampus; stereoencephalography (sEEG); high-frequency activity (HFA)

Introduction

The human hippocampus has been claimed to play a crucial role in human memory system especially long-term memory, also referred to as episodic memory[1], it comes with an equally important caveat that the hippocampus does not contribute to short-term memory[2]. The traditional/conventional view of hippocampal amnesia is that short-term memory is preserved despite the severe impairment in long-term memory[1,3,4]. For several years, a wide range of studies have established the medial temporal lobe, and hippocampus in particular, to be critically associated with long-term memory processing. For instance, researchers studied H.M., the well-known amnesic patient who had undergone temporal lobe resection, and published observations of some types of preserved memory (motor skill and perceptual learning) in the patient[5,6], as well as types of memory that depended on parts of the brain

other than the medial temporal lobe, in particular the hippocampus, or that did not depend on conscious awareness were termed implicit (non-declarative) memory[7,8]. Previous view mentioned that amnesia associated with hippocampal region damage is selective to declarative memory, the ability to recall facts and events[9], and for H.M. the deficit includes the complete absence of episodic memory, defined as the ability to remember specific personal experiences[10]. In addition, many researchers have explored the underlying neural mechanism of memory in which studies have implied that selective injury to the medial temporal lobe leads to an isolated deficit in long-term memory[11,12]. Evidence from studies in patients with amnesia due to hippocampal damage proposed that patients exhibit impairment of conscious episodic encoding and retrieval[13]. Nevertheless, there is different view reports that amnesic patients with bilateral lesions involving the hippocampus region suffered from an impairment of short-term memory along with research develops[14,15]. Moreover, recent visual study in human supports the view that the hippocampus does indeed have a role to play in short-term memory[16] and study reported the role of cross-frequency coupling in cortico-hippocampal activity in sequential auditory short-term memory (ASTM)[17].

Short-term memory is a storage system that has an evolutionary survival advantage, such as paying attention to limited but essential information excluding confounding factors and ASTM is one of the important sensory-associated subsystems[18]. Short-term memory trace is the physical neural basis for that function—the temporary, biological activity in our brain that makes short-term memory possible. Mismatch negativity (MMN), the index of automatic deviance detection, reflects the cortical responses to deviant changes occurred in the auditory system[19,20], has been regarded as the detection of a deviant stimulus within the ASTM[21], was first described about 40 years ago by Näätänen and his colleagues[22]. A various of studies leading to believe that the MMN is the outcome of an automatic comparison process between a new, deviant stimulus and the memory trace formed by the sensory representation of the standard stimuli within the short-term memory[23-26]. The MMN depends on the presence of a ASTM trace in the auditory cortex representing the repetitive aspects of the preceding auditory events[19]. Thus, ASTM trace formation is indexed by the MMN responses and deviance detection is mediated by a more stable short-term memory trace[27,28]. In the current study, we used a short-term memory paradigm presented with a contrast auditory stream created by different pure tones and intracranial recordings, a high spatial and temporal resolution electrophysiological study across various human brain regions to explore whether the human hippocampus contributes to/is involved in ASTM trace formation process.

High frequency activities (HFA, 70–150 Hz) elicited by the auditory stimuli is a reliable

electrophysiological correlate of underlying averaged spiking activity generated by the

thousands of neurons that are in the immediate vicinity of the recording electrodes[29-32]. This

type of activity has been linked to various cognitive processes, including attention, memory,

and sensory processing. Significant HFA of deviance detection evoked by the subjects

indicate the generation of ASTM trace. In our study, HFA results revealed that hippocampus

was involved during process of ASTM trace. Auditory response latencies recorded from the

hippocampus, insula, temporal lobe, parietal lobe, and frontal lobe showed the early

processing of ASTM trace at a pre-attention stage. Moreover, Granger causality analysis

further demonstrated ASTM trace is processed in hierarchical cortical areas, and detailed

interactions of brain regions among this memory trace process have been clarified/interpreted

clearly. Specifically, bottom-up auditory signals processed by the insula, and auditory regions

of the temporal lobe and the frontal lobe at an early stage, then auditory top-down

information transmitted from the frontal lobe to the hippocampus, the inferior temporal lobe,

as well as insula at a relatively late pre-attention stage of the auditory information processing.

To sum up, these results provide evidence that the human hippocampus contributes to the

ASTM trace of auditory perception at a pre-attention stage. Accompanied by the specific

hierarchical network of deviance processing obtained from Granger causal analysis, the

current study presents a more comprehensive and detailed understanding the formation of

ASTM trace, suggests a challenge to entrenched beliefs in the classification/definition of

memory systems.

Materials and methods

Ethics statement

This study was approved by the Research Ethics Committee of the First Affiliated Hospital, University of Science and Technology of China. All participants provided written informed consent prior to taking part in the study. Neither the study procedures nor the analytical methods were preregistered in a time-stamped institutional registry before the research was conducted.

Participants

Stereoelectroencephalography (sEEG) data were obtained from 17 patients (6 males, 11 females) with drug-resistant epilepsy who were being evaluated as potential candidates for resective surgery. Data collection took place at the First Affiliated Hospital, University of Science and Technology of China. As part of their presurgical assessment, each patient underwent invasive sEEG monitoring. The selection of regions for electrode implantation was determined solely by clinical criteria. All participants were implanted with intracranial depth sEEG electrodes (0.8 mm in diameter, with a 3.5 mm center-to-center interelectrode distance) to help localize the epileptogenic zone. Every participant was right-handed, had normal cognitive function, and normal hearing. Written informed consent was obtained from all subjects prior to enrollment. The experimental protocol received approval from the Institutional Review Board of the University of Science and Technology of China and the First Affiliated Hospital.

Task paradigm

Stimuli in the current study consisted of pure tones generated using Audition 3.0 (Adobe Systems Inc., Mountain View, CA, USA). The auditory contrast was formed by a sequence of 300 ms pure tones, with a 500 Hz tone serving as the frequent standard stimulus and a 700 Hz tone as the infrequent deviant stimulus (**Figure 1**). All stimuli were delivered in random order via headphones using E-Prime (Psychology Software Tools, Inc., USA). Participants were instructed to ignore the auditory stimuli and instead focus on watching a muted movie of their choice with subtitles.

Data acquisition

For each participant, structural MRI scans were acquired prior to electrode implantation, and post-implantation CT scans were obtained afterward. sEEG signals were recorded using a 128/256-channel Neurofax EEG-1200c amplifier/digitizer system (Nihon Kohden, Japan) via implanted depth electrodes. Recording settings included a high-pass filter with a 0.01 Hz cut off, a 50 Hz notch filter, and a sampling rate of either 2000 Hz or 5000 Hz. Electrodes located on the inner surface of the skull served as the ground and reference.

Preprocessing and Electrode localization

Our sEEG data was imported to MATLAB using FieldTrip toolbox[33] and resampled to 1000 Hz. Event onset markers were read from pre-allocated trigger channels. The data were then re-referenced to the mean of the mastoid electrodes, and unused channels were removed. Segments containing epileptiform activity or abnormal fluctuations were manually rejected. The electrode localization was based on the FieldTrip protocol[34] with an adaptation for Windows, in which post-implantation CT images was co-registered to pre-implantation MRI images using SPM 12[35]. The images were subsequently normalized to the MNI-152 template space using SPM12[36], and electrode positions were determined according to the FieldTrip pipeline.

High frequency activity (HFA) analysis

Line noises at 50 Hz, 100 Hz, and 150 Hz was removed using a notch filter. The continuous data were segmented into 900 ms epochs ($\square$200 ms to 700 ms). Trials were rejected if the amplitude exceeded 5 SD standard deviations from the mean for more than 25 consecutive ms, or if the power spectral density exceeded 5 standard deviations for more than 6 consecutive Hz[37]. Eligible trials were re-referenced to a neighboring channel for HFA analysis. HFA signals from –100 ms to 600 ms were extracted for further processing. Preprocessed data were band-pass filtered into eight 10-Hz-wide bands ranging from 70 to 150 Hz using a zero-phase Hamming windowed FIR filters (order 138)[38]. The envelope of each narrowband signal was extracted by taking the absolute value of the analytic signal from the Hilbert transform, and each amplitude time series was divided by its mean value[39]. For channels exhibiting a significant linear trend (R-square > 0.3), the trend was removed. Each channel's baseline period was removed and normalized by dividing the standard deviants of all trials of the baseline period.

Tone response and deviance detection

To identify channels with significant responses to auditory stimuli, a two-tailed cluster-based permutation test with threshold-free cluster enhancement[40] and 1000 sign-flipping randomizations[41] was applied separately to standard and deviant trials for each channel. Channels with a corrected p-value below 0.05 either stimulus type were considered responsive. Channels displaying abnormal waveforms or epileptic activity were subsequently excluded through manual review. Then, in order to assess whether a channel was involved in deviance detection, a two-tailed cluster-based permutation test with threshold-free cluster enhancement[40] and 1000 randomizations was applied to compare trial categories (standard vs. deviant) for each channel. Channels with a corrected p-value less than 0.05 were considered significant for deviance detection. Additional manual rejection was applied to exclude channels with aberrant signals or association with epileptic activity. The latency of high-frequency activity (HFA) responses was estimated using a fractional area method[42]. The procedure began by identifying the time point of maximum amplitude within the response window as the initial seed point. The significant cluster was expanded bilaterally from this seed until the amplitude dropped below 20% of the seed value or reached the epoch boundaries. The latency was defined as the center of mass of the resulting suprathreshold cluster.

Event-related potential analysis and clustering

Event-related potentials (ERP) were derived from the same trials used in the high-frequency activity (HFA) analysis, prior to re-referencing to neighboring channels. The ERP signals were filtered with a zero-phase FIR filter (order 18, 1–20Hz), segmented into 700 ms epochs (–100 ms to 600 ms relative to stimulus onset) and baseline-corrected by subtracting the mean amplitude during the –100 ms to 0 ms interval. For channels exhibiting a significant linear trend (R-square > 0.3), detrending was applied. Similar to the HFA analysis, a two-tailed cluster-based permutation test incorporating threshold-free cluster enhancement (TFCE) was conducted for each electrode shaft to identify channels with significant responses to deviant stimuli. Using principal component analysis (PCA) in line with Blenkmann et al.[37], we identified groups of event-related potential (ERP) responses exhibiting consistent temporal patterns across channels and subjects. This approach also enabled the mitigation of polarity reversal effects. The analysis proceeded as follows: first, a difference waveform was computed by subtracting the standard stimulus response from the

204   deviant stimulus response. PCA was then applied to this difference signal, and the first four

205   components capturing the highest channel weights were retained. Channels were grouped

206   based on their highest component weight, those predominately weighted on the first

207   component were clustered together, followed by those associated with the second component,

208   and so forth for each subsequent component.

Granger Causality Analysis

Wiener-Granger causality (G-causality, GC) is a statistical notion of causality applicable to

time series data, whereby cause precedes, and helps with prediction[43,44], which has been

developed to a powerful method for identifying information transfer or directed functional

("causal") connectivity in neural time series data[45,46]. It is defined in both time and frequency

domains, and allows for the conditioning out of common causal influences. Originally

developed in the context of econometric theory, it has since achieved broad application in the

neurosciences and beyond[47]. G-causality says that a variable X "G-causes" another variable

Y if the past of X contains information that helps predict the future of Y, over and above the

information already in the past of Y itself (and in the past of other "conditioning" variables

Z). In general terms, when this condition is satisfied one can say that there is "information

flow" from X to Y. This is justified because G-causality is an approximation to transfer

entropy (the approximation is exact for Gaussian variables), which is a directed version of

Shannon's mutual information, a very general way of characterizing the statistical

dependency or shared information between two variables[48]. The multivariate Granger

causality analysis (MVGC) Matlab toolbox used in the current study was designed for

applications in neuroscience[49], which implements numerical routines for calculating

multivariate Granger causality (MVGC) from time series data, both unconditional and

conditional, in the time and frequency domains[50]. Since it has been recognized that a moving

average (MA) component in the data presents a serious confound to Granger causal analysis,

as routinely performed via autoregressive (AR) modeling[49]. Previous studies have solved this

problem by demonstrating that Granger causality may be calculated simply and efficiently

from the parameters of a state-space (SS) model[44]. Because SS models are equivalent to

autoregressive moving average models, Granger causality estimated in this fashion is not

degraded by the presence of a MA component. State-space Granger causal inference stands to

significantly enhance our ability to identify and understand causal interactions and

information flow in the current study[51]. We used the MVGC (Multivariate Granger Causality)

Matlab toolbox for calculating multivariate Granger causality from time series

data[49]. Autoregressive moving average (ARMA) process or, equivalently, finite order linear
state-space (SS) have been applied in this study[51]. We selected a time window of interest
spanning –100 ms to 600 ms to perform identical analyses. The model order was estimated
using Akaike information criterion. The significance of difference of G-causality between
standard and deviant conditions was estimated using a permutation test.

Results

The hippocampus exhibited a significant high-frequency activity (HFA) response during auditory deviance detection.

In the current study, we employed a classic auditory oddball paradigm combined with intracranial recordings across multiple human brain regions to investigate the neural mechanisms underlying short-term memory trace formation. The hippocampus, a key structure within the medial temporal lobe memory system, is critically involved in the integration and relay of auditory information to downstream perceptual and cognitive regions[52-55]. Thus, for each recording channel in the hippocampal region, we computed trial-averaged high-frequency activity responses to both standard and deviant pure tones (Fig. 2).

Our results revealed that hippocampal channels displayed diverse response profiles, including both HFA enhancements and high-frequency suppression (HFS). Specifically, responses to standard stimuli were predominantly characterized by negative-going deflections (Fig. 2a and Fig. 2c), whereas responses to deviant stimuli consistently exhibited positive deflections (Fig. 2e and Fig. 2g). To identify channels exhibiting significant responses to auditory stimuli, a two-tailed cluster-based permutation test incorporating threshold-free cluster enhancement (TFCE)[40] was applied separately to standard and deviant trials for each channel. The test was performed using 1000 sign-flipping randomizations. Channels with a corrected p-value below 0.05 for either stimulus type were classified as responsive. Exemplar channels exhibiting statistically significant responses to standard pure tones are shown in or deviant pure tones are shown in Fig. 2b and Fig. 2d ($p < 0.05$), while exemplar channels exhibiting statistically significant responses to deviant pure tones are presented in Fig. 2f and Fig. 2h ($p < 0.05$).

Classifications of High-Frequency Responses to Pure Tones.

Furthermore, electrode pairs that evoked significant mismatched activity were categorized

according to whether the mismatch activity was enhanced or suppressed. We collected sEEG

data from 17 epilepsy patients (Table 1). To obtain a global description of the brain regions in

deviance detection, we examined the HFA responses to standard versus deviant tones across

intracranial channels. In the HFA analysis, 49 channels responded to tone responses in 10 out

of the 17 patients (Fig. 3), which illustrates the overall response profiles across all channels,

including both HFA and high-frequency suppression (HFS). Specifically, Significant HFS

responses were observed in 8 channels across 6 patients with and 41 channels across 10

subjects with HFA responses ($p < 0.05$, Fig. 3a, b). HFS in response to the stimuli at a

latency of 395 ms were recognized at the middle temporal gyrus (MTG), temporo-occipital

brain area, posterior inferior temporal gyrus, and anterior corona radiata a channel. In

addition, 29 channels across 9 patients responded exclusively to deviant stimuli (Fig. 3c)

whereas 12 channels across 7 patients responded to both standard and deviant tones (Fig. 3d).

It is known that studying HFA with sEEG recordings can provide insights into the neural

bases of human cognition that cannot be derived as easily from the study of BOLD effects

with fMRI or from the analysis of EEG[30]. HFS, a transient energy suppression in the gamma-

band (40−150 Hz) during the processing of sensory stimuli, which was first reported by

Lachaux[56] and has close relationship with attention[57]. We also observed widely distributed

high frequency suppression (HFS)[58], particularly in response to standard stimuli, which

generally occurred later than high frequency activity (HFA). This widespread HFS may

reflect an energy saving mechanism in the brain when processing predictable events. Given

the known interaction between early tone responses and later repetition suppression[59], such

suppression may be mediated by repetition suppression mechanisms. Supplementary Figure 1

showed HFA responses across individual patients.

291 Various brain regions showed significant HFA during short-term memory
292 trace formation.

293   We investigated the spatiotemporal dynamics and effective connectivity of high-frequency

294   activity (HFA) responses across key brain regions during the processing of auditory deviant

295   stimuli (Fig S1). Mean HFA/HFS responses to standard and deviant pure tones, along with

296   the deviant-minus-standard difference waveforms were shown for multiple brain regions

297   (Fig. 4a). These brain regions included the insula; temporal lobe (encompassing Heschl's

gyrus, superior temporal gyrus, and superior temporal sulcus); frontal lobe (including the middle frontal gyrus, superior frontal gyrus, anterior cingulate cortex, and frontal operculum); parietal lobe (comprising the postcentral gyrus, parietal operculum, and supramarginal gyrus); posterior temporal lobe (the temporo-occipital part of the middle temporal gyrus); inferior temporal lobe (including the inferior temporal gyrus, fusiform gyrus, and inferior portion of the middle temporal gyrus); and the hippocampal area (encompassing the hippocampus, parahippocampal gyrus, and amygdala). Moreover, A scatter plot illustrates the response latencies of HFA/HFS from electrode pairs in the aforementioned regions. The data reveal a systematic, ascending trend in response latencies across these different brain areas (Fig. 4b), suggesting a sequential propagation of deviance processing. Brain regions included the insula, temporal lobe (i.e., Heschl's gyrus, STG, superior temporal sulcus, STS), frontal lobe (inferior frontal gyrus middle frontal gyrus, anterior (para) cingulate cortex, frontal operculum), parietal lobe (parietal lobe postcentral gyrus, central opercular cortex, parietal operculum, supramarginal gyrus), posterior temporal lobe (temporooccipital division) hippocampus, and the inferior temporal lobe (fusiform gyrus, inferior temporal gyrus, middle temporal gyrus) were all involved during the processing of deviance detection, and the latencies of auditory mismatch HFA of these brain areas ranged from 110 ms to 438 ms (Fig. 4b).

Granger causality analysis showed connectivity of the hippocampus with various brain regions.

Next, we investigated the neural connectivity underlying auditory deviance processing by applying linear Granger causality analysis. To effectively visualize the complex network, electrode pairs were first merged according to their respective brain regions. The analyzed regions included the insula; temporal lobe (Heschl's gyrus, superior temporal gyrus, superior temporal sulcus); frontal lobe (middle frontal gyrus, superior frontal gyrus, anterior cingulate cortex, frontal operculum); parietal lobe (postcentral gyrus, parietal operculum, supramarginal gyrus); posterior temporal lobe (temporo-occipital part of the middle temporal gyrus); inferior temporal lobe (inferior temporal gyrus, fusiform gyrus, inferior portion of the middle temporal gyrus); and the hippocampal area (hippocampus, parahippocampal gyrus, amygdala). The anatomical locations of these regions were mapped onto a common cortical space, as shown in the enlarged image of the boxed area, with different colors denoting

distinct brain clusters. The resulting connectivity is depicted in the heatmap (Fig. 4c), which

displays the estimated difference in Granger causal connection strengths between deviant and

standard stimulus-evoked activities. Each cell represents the strength of the causal influence

from the row-specified region to the column-specified region. Gray indicates connections

with no significant difference between conditions, whereas colored cells represent pathways

where the deviant stimulus elicited a significant change in effective connectivity. The brain

regions implicated by Granger causality analysis were rendered on a standardized cortical

surface. A magnified inset of the demarcated area is presented, in which unique colors code

for separate functional clusters and the directionality (bottom-up versus top-down) of their

causal influences (Fig. 5).

Intracranial event-related potentials

Mean event-related potential (ERP) responses to standard and deviant pure tones, as well as

the deviant-minus-standard difference waveforms were analyzed. Fig S2 showed ERP

responses across individual patients. To ensure waveform comparability, only channels

exhibiting significant responses to both conditions within each analytical dimension were

included in the averages and subsequent plots. The induced activity in the high gamma band

as mismatch response (MMR), an intracranial EEG counterpart to scalp mismatch negativity

(MMN)[60]. Given that the same neural phenomenon can be recorded either as a positive or a

negative intracranial ERP (iERP)[30]. An iERP equivalent to the scalp mismatch response were

shown in Fig S3. And iERP effects were present in 188 channels across 17 patients. Cause

we grouped the iERP responses from different channels according to the highest weights of

the PCA decomposition. Figure 4 shows the iERP responses to deviant and standard tones for

the four groups of channels. Differences in Group 1 ($n = 121$, $s = 6$), group 2 ($n = 41$, $s = 6$)

and group 4 ($n = 8$, $s = 4$) showed significant MMR ($P < 0.05$), and differences in group 3 (n

$= 18$, $s = 3$) showed significant MMN ($P < 0.05$).

Discussion

In the current study, we employed a short-term memory paradigm featuring a contrastive

auditory stream of pure tones to investigate the role of the human hippocampus in the

formation of auditory short-term memory (ASTM) trace. Combined with high spatiotemporal

resolution approach, intracranial recordings used in this study allowed us to examine the
electrophysiological activity across multiple brain regions. High-frequency activity (HFA,
70–150 Hz), a reliable electrophysiological correlate of localized neuronal firing, served as
our primary metric. The significant HFA elicited during deviance detection indicates the
generation of ASTM trace. Our results revealed that the hippocampus is involved in this
process. Furthermore, the response latencies observed in the hippocampus, insula, temporal,
parietal, and the frontal regions suggest that the initial process of the ASTM trace occurs at a
pre-attention stage. Furthermore, Granger causality analysis demonstrate that the ASTM trace
is processed through a hierarchical cortical network. Specifically, bottom-up auditory signals
were initially processed by the insula and auditory regions of the temporal and frontal regions.
This was followed by the top-down information flow from the frontal region to the
hippocampus, inferior temporal lobe, and insula at a later pre-attention stage. In summary,
these findings provide direct evidence that the human hippocampus contributes to ASTM
trace formation at a pre-attention stage. Coupled with the specific hierarchical network for
deviance processing identified through Granger causality, our study offers a more
comprehensive model of ASTM trace formation and challenges entrenched definition of
memory systems.

The hippocampus is involved in processing of auditory short-term memory trace formation.

The hippocampus is well known to be involved in the medial temporal lobe memory system
and also has been thought to play a critical role in relaying auditory information to later
perceptual and cognitive processes[52-55]. In terms of auditory perception, the hippocampus
receives afferent inputs form the auditory association cortices, and in turn, it projects back to
the primary auditory cortex and auditory association areas[61,62]. MRI and MEG recordings
combined with equivalent current dipole (ECD) analysis revealed electrophysiological
signatures of hippocampal involvement in auditory processing[62]. The noise responses of
hippocampal neurons in awake mice have identified a population of neurons that can be
associated with auditory fear conditioning[63]. Previous study used ERPs and event-related
magnetic fields (ERFs) to investigate the relationship between the hippocampus and the
event-related responses in auditory information processing[64]. Results have suggested that the
auditory processing is carried out by the combination of multiple structures including the

mesial temporal area, the superior temporal area and the inferior parietal area on both

hemispheres, and that the generator (hippocampus) in the mesial temporal area contributes to

the scalp-recorded P300 only at the corresponding anterior temporal region[64]. MEG and

effective connectivity analyses (dynamic causal modeling) showed that the hippocampus

and ventromedial prefrontal cortex (vmPFC) work as a functional circuit during mismatch

detection, suggesting that the vmPFC drives the hippocampus during the generation and

processing of mismatch signals[65]. Previous intracranial and lesion research proposed that

hippocampus is involved with novelty processing in a fronto-hippocampal network and the

hippocampus plays an active role in contextual updating at the end of the cognitive episode[66].

In terms of auditory perception, the hippocampus receives afferent inputs form the auditory

association cortices, and in turn, it projects back to the primary auditory cortex and auditory

association areas[61,62]. MRI and MEG recordings combined with equivalent current dipole

(ECD) analysis revealed electrophysiological signatures of hippocampal involvement in

auditory processing[62]. The noise responses of hippocampal neurons in awake mice have

identified a population of neurons that can be associated with auditory fear conditioning[63].

Previous study used ERPs and event-related magnetic fields (ERFs) to investigate the

relationship between the hippocampus and the event-related responses in auditory

information processing[64]. Results have suggested that the auditory processing is carried out

by the combination of multiple structures including the mesial temporal area, the superior

temporal area and the inferior parietal area on both hemispheres, and that the generator

(hippocampus) in the mesial temporal area contributes to the scalp-recorded P300 only at the

corresponding anterior temporal region[64].

Auditory short-term memory trace is processed in hierarchical cortical areas.

415
416

Predictive coding theories posit that neural networks learn statistical regularities in the

environment for comparison with actual outcomes, signaling a prediction error (PE) when

sensory deviation occurs[67]. The predictive coding model of perception proposes that neuronal

responses are modulated by the amount of sensory input that the internal prediction cannot

account for (i.e., prediction error) and postulates that perception entails the top-down

propagation of prediction and the bottom-up propagation of prediction error[68]. Within a

predictive coding framework, MMN and novelty-related ERP signatures are interpreted as

prediction-error responses that signal the mismatch between sensory inputs and internal
models[69]. The MMN is considered the classic PE signal elicited during passive listening to
deviant sounds interrupting the context provided by a sequence of repeated standard
stimuli[70]. Auditory prediction errors, i.e. the mismatch between predicted, forthcoming
auditory sensations and actual sensory input, trigger the detection of surprising auditory
events in the environment (Friston, 2005). Previous studies used DCM to investigate the
effective connectivity of auditory mismatch responses and demonstrated a three-level
hierarchical network comprised of connections from bilateral primary auditory cortex (A1) to
the planum temporale (PT), and from the PT to the right inferior frontal gyrus (IFG) (Garrido
et al., 2007, 2008). This network included intrinsic (within source) connections in A1 and
extrinsic (between-sources), forward and backward, connections between A1, PT and IFG.
The signaling of top-down (feedback signals) predictions and bottom-up (feedforward signal)
prediction errors conveyed across hierarchical regions[71].

The formation of the ASTM trace is not an isolated function of a single region but arises from
coordinated activity within a hierarchical cortical network. Our Granger causality analysis,
based on a state-space model, delineated the specific directional information flows
underlying this process. The network operates through a dynamic interplay of bottom-up and
top-down processing. Bottom-up auditory signals from the deviant stimulus are initially
processed by the insula and auditory regions within the temporal and frontal lobes.
Subsequently, at a later pre-attention stage, a top-down information flow emerges from the
frontal region back to the hippocampus, inferior temporal lobe, and insula. This connectivity
pattern positions the hippocampus within a regulated circuit. The hippocampus appears to be
driven by higher-order frontal regions during the processing of mismatch signals. This is
consistent with previous effective connectivity analyses (e.g., Dynamic Causal Modeling)
showing that structures like the ventromedial prefrontal cortex (vmPFC) drive the
hippocampus during mismatch detection[65]. Furthermore, the insula acts as a critical hub in
this network. Evidence from intracranial EEG (iEEG) indicates its involvement in automatic
deviance detection[37,72] and its role in coordinating hierarchical processing across sensory and
salience networks[73].

In summary, the ASTM trace is processed through a precisely positioned hierarchical
network. The initial, pre-attentive stages involve feedforward signaling of prediction error
from lower sensory areas. This is followed by feedback interactions, where frontal regions
modulate activity in the hippocampus and other temporal areas, facilitating the updating of

the short-term auditory model or context[66]. This comprehensive network model, revealed

through direct intracranial recordings and connectivity analysis, offers a detailed framework

for understanding how distributed brain regions, including the hippocampus, interact to form

and maintain auditory short-term memory traces.

References

1. Bird CM, Burgess N. The hippocampus and memory: insights from spatial processing. *Nature reviews neuroscience* **9**, 182-194 (2008).

2. Husain M. The human hippocampus contributes to short-term memory. *Brain : a journal of neurology* **147**, 2593-2594 (2024).

3. Warrington EK. Amnesia and the distinction between long-and short-term memory. *Journal of verbal learning and verbal behavior* **9**, 176-189 (1970).

4. Cave CB, Squire LR. Intact verbal and nonverbal short-term memory following damage to the human hippocampus. *Hippocampus* **2**, 151-163 (1992).

5. Corkin S. Acquisition of Motor Skill after Bilateral Medial Temporal-Lobe Excision. *Neuropsychologia* **6**, 255-& (1968).

6. Moscovitch M. Recovered Consciousness - a Hypothesis Concerning Modularity and Episodic Memory. *Journal of Clinical and Experimental Neuropsychology* **17**, 276-290 (1995).

7. Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychological review* **99**, 195-231 (1992).

8. Schacter DL. Implicit Memory: History and Current Status. *Journal of Experimental Psychology Learning Memory & Cognition* **13**, 501-518 (1987).

9. Cohen NJ, Squire LR. Preserved learning and retention of pattern-analyzing skill in amnesia: Dissociation of knowing how and knowing that. *Science* **210**, 207-210 (1980).

10. Steinvorth S, Levine B, Corkin S. Medial temporal lobe structures are needed to re-experience remote autobiographical memories: evidence from HM and WR. *Neuropsychologia* **43**, 479-496 (2005).

11. Scoville WB, & Milner, B. . Loss of recent memory after bilateral hippocampallesions. *J Neurol Neurosurg Psychiat* **20**, (1957).

12. Reber PJ. The neural basis of implicit learning and memory: a review of neuropsychological and neuroimaging research. *Neuropsychologia* **51**, 2026-2042 (2013).

13. Tulving E. Episodic memory: From mind to brain. *Annual review of psychology* **53**, 1-25 (2002).

14. Hannula DE, Tranel D, Cohen NJ. The long and the short of it: relational memory impairments in amnesia, even at short lags. *Journal of Neuroscience* **26**, 8352-8359 (2006).

15. Olson IR, Page K, Moore KS, Chatterjee A, Verfaellie M. Working memory for conjunctions relies on the medial temporal lobe. *Journal of Neuroscience* **26**, 4596-4601 (2006).

16. Daume J*, et al.* Control of working memory by phase–amplitude coupling of human hippocampal neurons. *Nature* **629**, 393-401 (2024).

501   17.   Griffiths TD*, et al.* Cross-frequency coupling in cortico-hippocampal networks
502   supports the maintenance of sequential auditory information in short-term memory.
503   *PLOS Biology* **22**, (2024).

504   18.   Cascella M, Al Khalili Y. Short-term memory impairment. In: *StatPearls [Internet]*).
505   StatPearls Publishing (2024).

506   19.   Näätänen RN, Paavilainen P, Rinne T, Alho K. The mismatch negativity (MMN) in
507   basic research of central auditory processing: a review. *Clinical neurophysiology :*
508   *official journal of the International Federation of Clinical Neurophysiology* **118**,
509   2544-2590 (2007).

510   20.   Partanen E, Kujala T, Naatanen R, Liitola A, Sambeth A, Huotilainen M. Learning-
511   induced neural plasticity of speech processing before birth. *Proceedings of the*
512   *National Academy of Sciences of the United States of America* **110**, 15145-15150
513   (2013).

514   21.   Colin C, Radeau M, Soquet A, Demolin D, Colin F, Deltenre P. Mismatch negativity
515   evoked by the McGurk-MacDonald effect: a phonetic representation within short-
516   term memory. *Clinical Neurophysiology* **113**, 495-506 (2002).

517   22.   Näätänen R, Gaillard AWK, Mäntysalo S. Early Selective-Antention Effect on
518   Evoked Potential Reinterpreted. *Acta Psychologica* **42**, 313-329 (1978).

519   23.   Novak GP, Ritter W, Vaughan Jr HG, Wiznitzer ML. Differentiation of negative
520   event-related potentials in an auditory discrimination task. *Electroencephalography*
521   *and clinical neurophysiology* **75**, 255-275 (1990).

522   24.   Näätänen R. The role of attention in auditory information processing as revealed by
523   event-related potentials and other brain measures of cognitive function. *Behavioral*
524   *and brain sciences* **13**, 201-233 (1990).

525   25.   Cowan N, Winkler I, Teder W, Näätänen R. Memory prerequisites of mismatch
526   negativity in the auditory event-related potential (ERP). *Journal of experimental*
527   *psychology: Learning, Memory, and Cognition* **19**, 909 (1993).

528   26.   Näätänen R, Alho K. Mismatch negativity-a unique measure of sensory processing in
529   audition. *International Journal of Neuroscience* **80**, 317-337 (1995).

530   27.   Teichert T, Jedema H, Zhen Z, Gurnsey K. Complementary functional profiles of
531   mismatch responses mediated by adaptation and deviance detection point to two
532   distinct auditory short-term memory systems. *Journal of neurophysiology* **134**, 314-
533   336 (2025).

534   28.   Tuomainen OT. Auditory short-term memory trace formation for nonspeech and
535   speech in SLI and dyslexia as indexed by the N100 and mismatch negativity
536   electrophysiological responses. *Neuroreport* **26**, 374-379 (2015).

537   29.   Ray S, Maunsell JH. Different origins of gamma rhythm and high-gamma activity in
538   macaque visual cortex. *PLoS Biol* **9**, e1000610 (2011).

539   30.   Lachaux JP, Axmacher N, Mormann F, Halgren E, Crone NE. High-frequency neural
540   activity and human cognition: past, present and possible future of intracranial EEG
541   research. *Prog Neurobiol* **98**, 279-301 (2012).

542   31.   Rich EL, Wallis JD. Spatiotemporal dynamics of information encoding revealed in
543   orbitofrontal high-gamma. *Nature communications* **8**, 1139 (2017).

32. Watson BO, Ding M, Buzsaki G. Temporal coupling of field potentials and action
potentials in the neocortex. *The European journal of neuroscience* **48**, 2482-2497
(2018).

33. Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput*
Intell Neurosci **2011**, 156869 (2011).

34. Stolk A*, et al.* Integrated analysis of anatomical and electrophysiological human
intracranial data. *Nat Protoc* **13**, 1699-1723 (2018).

35. Studholme C, Hill DLG, Hawkes DJ. An overlap invariant entropy measure of 3D
medical image alignment. *Pattern Recognition* **32**, 71-86 (1999).

36. Ashburner J, Friston KJ. Unified segmentation. *NeuroImage* **26**, 839-851 (2005).

37. Blenkmann AO*, et al.* Auditory deviance detection in the human insula: An
intracranial EEG study. *Cortex* **121**, 189-200 (2019).

38. Mesgarani N, Cheung C, Johnson K, Chang EF. Phonetic Feature Encoding in Human
Superior Temporal Gyrus. *Science* **343**, 1006-1010 (2014).

39. Norman Y, Yeagle EM, Harel M, Mehta AD, Malach R. Neuronal baseline shifts
underlying boundary setting during free recall. *Nature Communications* **8**, 1301
(2017).

40. Smith SM, Nichols TE. Threshold-free cluster enhancement: Addressing problems of
smoothing, threshold dependence and localisation in cluster inference. *NeuroImage*
44, 83-98 (2009).

41. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation
inference for the general linear model. *NeuroImage* **92**, 381-397 (2014).

42. Luck SJ. Event-related potentials from the user's perspective. *Nature neuroscience* **9**,
463 (2006).

43. Granger CWJ. Investigating causal relations by econometric models and cross-
spectral methods. *Econometrica* **37**, 424-438 (1969).

44. Seth AK, Barrett AB, Barnett L. Granger Causality Analysis in Neuroscience and
Neuroimaging. *Journal of Neuroscience* **35**, 3293-3297 (2015).

45. Bressler SL, Seth AK. Wiener-Granger causality: a well established methodology.
NeuroImage **58**, 323-329 (2011).

46. Schreiber T. Measuring information transfer. *Physical review letters* **85**, 461 (2000).

47. Barnett L, Seth AK. The MVGC multivariate Granger causality toolbox: A new
approach to Granger-causal inference. *Journal of Neuroscience Methods* **223**, 50-68
(2014).

48. Barnett L, Barrett AB, Seth AK. Granger causality and transfer entropy are equivalent
for Gaussian variables. *Phys Rev Lett* **103**, 238701 (2009).

49. Barnett L, Seth AK. The MVGC multivariate Granger causality toolbox: a new
approach to Granger-causal inference. *J Neurosci Methods* **223**, 50-68 (2014).

50. Seth AK. A MATLAB toolbox for Granger causal connectivity analysis. *J Neurosci*
Methods **186**, 262-273 (2010).

51.   Barnett L, Seth AK. Granger causality for state-space models. *Physical review E,*
Statistical, nonlinear, and soft matter physics **91**, 040101 (2015).

52.   Bouton ME, Bolles RC. Conditioned Fear Assessed by Freezing and by the
Suppression of 3 Different Baselines. *Anim Learn Behav* **8**, 429-434 (1980).

53.   Liberman T, Velluti RA, Pedemonte M. Temporal correlation between auditory
neurons and the hippocampal theta rhythm induced by novel stimulations in awake
guinea pigs. *Brain research* **1298**, 69-77 (2009).

54.   Graham KS, Barense MD, Lee ACH. Going beyond LTM in the MTL: A synthesis of
neuropsychological and neuroimaging findings on the role of the medial temporal
lobe in memory and perception. *Neuropsychologia* **48**, 831-853 (2010).

55.   Lee ACH, Yeung LK, Barense MD. The hippocampus and visual perception.
Frontiers in human neuroscience **6**, (2012).

56.   Lachaux JP*, et al.* The many faces of the gamma band response to complex visual
stimuli. *NeuroImage* **25**, 491-501 (2005).

57.   Lachaux JP*, et al.* Silence is golden: transient neural deactivation in the prefrontal
cortex during attentive reading. *Cerebral cortex* **18**, 443-450 (2008).

58.   Lachaux J-P, Axmacher N, Mormann F, Halgren E, Crone NE. High-frequency neural
activity and human cognition: Past, present and possible future of intracranial EEG
research. *Progress in neurobiology* **98**, 279-301 (2012).

59.   Eckert D*, et al.* Distinct interacting cortical networks for stimulus-response and
repetition-suppression. *Communications biology* **5**, 909 (2022).

60.   Ishishita Y*, et al.* Deviance detection is the dominant component of auditory
contextual processing in the lateral superior temporal gyrus: A human ECoG study.
Human brain mapping **40**, 1184-1194 (2019).

61.   Mehta AD*, et al.* Seizure propagation in a patient with musicogenic epilepsy. *Epilepsy*
& Behavior **14**, 421-424 (2009).

62.   Chatani H*, et al.* Neuromagnetic evidence for hippocampal modulation of auditory
processing. *NeuroImage* **124**, 256-266 (2016).

63.   Xiao C, Liu Y, Xu J, Gan X, Xiao Z. Septal and Hippocampal Neurons Contribute to
Auditory Relay and Fear Conditioning. *Frontiers in cellular neuroscience* **12**, 102
(2018).

64.   Nishitani N*, et al.* The role of the hippocampus in auditory processing studied by
event-related electric potentials and magnetic fields in epilepsy patients before and
after temporal lobectomy. *Brain : a journal of neurology* **122 (Pt 4)**, 687-707 (1999).

65.   Garrido MI, Barnes GR, Kumaran D, Maguire EA, Dolan RJ. Ventromedial
prefrontal cortex drives hippocampal theta oscillations induced by mismatch
computations. *NeuroImage* **120**, 362-370 (2015).

66.   Fonken YM, Kam JWY, Knight RT. A differential role for human hippocampus in
novelty and contextual processing: Implications for P300. *Psychophysiology* **57**,
(2019).

67.   Durschmid S*, et al.* Hierarchy of prediction errors for auditory events in human
temporal and frontal cortex. *Proceedings of the National Academy of Sciences of the*
United States of America **113**, 6755-6760 (2016).

68. Hsu YF, Le Bars S, Hamalainen JA, Waszak F. Distinctive Representation of
Mispredicted and Unpredicted Prediction Errors in Human Electroencephalography.
The Journal of neuroscience : the official journal of the Society for Neuroscience **35**,
14653-14660 (2015).

69. Recasens M, Gross J, Uhlhaas PJ. Low-Frequency Oscillatory Correlates of Auditory
Predictive Processing in Cortical-Subcortical Networks: A MEG-Study. *Sci Rep-Uk* **8**,
(2018).

70. Naatanen R, Gaillard AW, Mantysalo S. Early selective-attention effect on evoked
potential reinterpreted. *Acta Psychol (Amst)* **42**, 313-329 (1978).

71. Oestreich LKL, Randeniya R, Garrido MI. Auditory white matter pathways are
associated with effective connectivity of auditory prediction errors within a fronto-
temporal network. *NeuroImage* **195**, 454-462 (2019).

72. Zhang Y*, et al.* The Roles of Subdivisions of Human Insula in Emotion Perception
and Auditory Processing. *Cerebral cortex* **29**, 517-528 (2019).

73. Allen M*, et al.* Anterior insula coordinates hierarchical processing of tactile mismatch
responses. *NeuroImage* **127**, 34-43 (2016).

Figure legends

**Fig. 1 Experimental design and channel localizations. Experiment design and channel
localizations. a** Auditory oddball paradigm: Stimuli consisted of 300 ms pure tones
(including 10 ms fade-in and fade-out periods). The standard tone (500 Hz) accounted for
90% of trials, while the deviant tone (700 Hz) comprised the remaining 10%. **b** 3D
reconstruction of the brain with implanted electrodes showing significant responses. To
improve clarity, non-responsive electrode shafts are omitted. Small black dots represent
channels without significant responses; large colored dots indicate channels with significant
responses, with each color corresponding to a different subject.

Fig. 2 Hippocampal high-frequency activity (HFA) responses to pure tones.

a Averaged positive HFA responses to elicited by standard stimuli. **b** Heatmap of a single
exemplar channel from the group presented in **a**. **c** Averaged negative HFA responses (or
high-frequency suppression) elicited by standard stimuli, **d** Heatmap of a single exemplar
channel from the group presented in **c**. **e** Averaged positive HFA responses to elicited by
deviant stimuli. **f** Heatmap of a single exemplar channel from the group presented in **e**. **g**
Averaged negative HFA responses (or high-frequency suppression) elicited by deviant
stimuli. **h** Heatmap of a single exemplar channel from the group presented in **g**. In panels **a**,

663 **c**, **e**, **g**, blue lines denote responses to standard trials; red lines represent responses to deviant

664 trials. Shaded areas indicate mean ± SEM. n indicates the number of channels averaged, and s

665 indicates the number of subjects. In panels **b**, **d**, **f**, **h**, yellow traces represent positive HFA

666 responses; blue traces indicate negative HFA responses high-frequency suppression (or high-

667 frequency suppression). Solid black vertical line indicates the onset of the stimulus, dashed

668 lines indicate the time windows exhibiting statistically significant responses to auditory

669 stimuli ($p < 0.05$).

671 **Fig. 3 Classification of mean HFA responses to pure tones.** Responses were averaged

672 across channels from different subjects showing significant differences between the deviant

673 and standard conditions. Mean HFA and high frequency suppression (HFS) responses are

674 shown for standard (blue dashed line) and deviant tones (red dashed line), along with deviant-

675 minus-standard difference waveforms (green solid line). **a** Mean HFS responses to standard

676 and deviant pure tones. **b** Mean HFA responses to standard and deviant pure tones. **c** Mean

677 HFA responses with no response to standard pure tones. **d** Mean HFA responses with

678 significant responses to standard pure tones ($p < 0.05$). Shadowed areas depict mean ± SEM,

679 n indicates the number of channels averaged, and s indicates the number of subjects.

681 **Fig. 4 Latencies of HFA responses in different brain regions and Granger causality**

682 **analysis of brain connectivity within different brain regions. a** Mean HFA/HFS responses

683 to standard (blue dash line) and deviant pure tones (red dash line), as well as deviant-minus-

684 standard difference waveforms (green solid line), are plotted for the insula, temporal lobe

685 (including Heschl's gyrus, superior temporal gyrus, superior temporal sulcus), frontal lobe

686 (including the middle frontal gyrus, superior frontal gyrus, anterior cingulate cortex, frontal

687 operculum), parietal lobe (including postcentral gyrus, parietal operculum, supramaginal

688 gyrus), posterior temporal lobe (including temporo-occipital partition in the middle temporal

689 gyrus), inferior temporal lobe (including inferior temporal gyrus, fusiform gyrus, and inferior

690 of middle temporal gyrus) and the hippocampus area (including hippocampus,

691 parahippocampal gyrus, amygdala). n indicates the number of channels averaged, and s

692 indicates the number of subjects. **b** The scatter plot shows the response latencies of HFA/HFS

693 in the electrode pairs from the brain regions mentioned above. The blue line indicates the

694 ascending trend of the latencies of different brain areas. **c** The heatmap shows the

connectivity between these brain regions. Estimates of the difference in Granger causal
connection strengths between deviant and standard stimulus-evoked activity across brain
regions. Each position in the plot indicates the strength of connections corresponding from
the row to the column. Gray indicates no significant difference, and color indicates
significant connection within brain areas. Specific brain regions calculated included in this
figure are shown as follows. The insula; temporal lobe: including Heschl's gyrus, superior
temporal gyrus, superior temporal sulcus; frontal lobe: including the middle frontal gyrus,
superior frontal gyrus, anterior cingulate cortex, frontal operculum; parietal lobe: including
postcentral gyrus, parietal operculum, supramaginal gyrus; posterior temporal lobe: including
temporo-occipital part in the middle temporal gyrus: inferior temporal lobe: including inferior
temporal gyrus, fusiform gyrus, and inferior of middle temporal gyrus; and the hippocampus
area: including hippocampus, parahippocampal gyrus, amygdala.

Fig. 5 Granger causality-based connectivity across brain regions.

Anatomical locations of regions identified by Granger causality analysis are projected onto a
common cortical surface. The enlarged inset highlights the boxed area. Distinct colors
represent different functional clusters and the directionality of connections (bottom-up vs.
top-down) within these regions.

Supplementary Fig. 1 HFA responses to different pure tones of Sub 3. HFA of tone-
responsive channels evoked by standard (blue lines) and deviant (red lines) pure tones are
plotted for sub 3, shadowed areas depict mean ± SEM. The latencies of HFA at each channel
were shown in each plot. Significant effects are marked at the top image, gray indicates there
is no significant difference between response to standard and deviant stimuli, and color
indicates significant deviance detection.

Supplementary Fig. 2 Mean ERP responses to different pure tones. ERP responses were
averaged across channels from different subjects showing significant differences between the
deviant and standard conditions. Only channels that showed significant responses to both
conditions within each dimension of analysis were included in the plots in order to obtain
comparable waveforms. **a** Responses for Group 1. **b** Responses for Group 2. **c** Responses for
Group 3. **d** Responses for Group 4. Mean ERP responses to standard (blue dash line) and

deviant pure tones (red dash line), as well as deviant-minus-standard difference waveforms

(green solid line), are plotted for subjects. Shadowed areas depict mean ± SEM, n indicates

the number of channels averaged, and s indicates the number of subjects.

Supplementary Fig. 3 Mean ERP responses to different pure tones. ERP responses were

averaged across channels from different subjects showing significant differences between the

deviant and standard conditions. Only channels that showed significant responses to both

conditions within each dimension of analysis were included in the plots in order to obtain

comparable waveforms. (A) Responses for Group 1. (B) Responses for Group 2. (C)

Responses for Group 3. d Responses for Group 4. Mean ERP responses to standard (blue

dash line) and deviant pure tones (red dash line), as well as deviant-minus-standard

difference waveforms (green solid line), are plotted for subjects. Shadowed areas depict mean

± SEM, n indicates the number of channels averaged, and s indicates the number of subjects.

Figure 1

a

b

n=29, s=12

Figure 3

Figure 4

Supplementary Figure 1

HFA Normalized Amplitude (Z)

Time (ms)

Non-Responsive

Responsive

Negative

Standard
Deviant

ERP Amplitude (μV)

Group 1

Group 2

Group 3

Group 4

Standard
Deviant

Time (ms)