Biochemical Regulation of Brain Kynurenic Acid Synthesis and Inhibition in Rats is Sensitive to the Time of Day

Courtney J. Wright[1,2], Silas A. Buck[3], Snezana Milosavljevic[1,2],
Ashley M. Lewis[1], Nathan T.J. Wagner[1], Ana Pocivavsek*[1,2,3]

[1]Department of Pharmacology, Physiology, and Neuroscience, University of South Carolina School of Medicine, Columbia, SC, USA
[2]Carolina Autism and Neurodevelopment Research Center, University of South Carolina, Columbia, SC, USA
[3]Maryland Psychiatric Research Center, Department of Psychiatry, University of Maryland School of Medicine, Baltimore, MD, USA

*Corresponding Author:
Ana Pocivavsek, PhD
Pharmacology, Physiology, and Neuroscience
USC School of Medicine
Columbia, SC 29209
Fax: 803-216-3500
Email: ana.pocivavsek@uscmed.sc.edu

Lay Summary: Changes in brain neurochemistry, including elevations in the tryptophan metabolite kynurenic acid (KYNA), are common in brain disorders that present with sleep disturbances and cognitive deficits as symptoms. KYNA interferes with neurotransmission critical for cognition and sleep, so therapeutic strategies to reduce brain KYNA are being pursued. As recent literature has highlighted the impact of time on brain metabolism and drug efficacy, we, for the first time, explored the hypothesis

that KYNA formation and synthesis inhibition change throughout the day. We measured extracellular KYNA levels in the brains of male and female rats and stimulated KYNA synthesis with physiological challenges (exogenous kynurenine, the KYNA bioprecursor, or acute sleep deprivation) and/or inhibited KYNA synthesis pharmacologically (PF-04859989, an inhibitor of the KYNA-synthesizing enzyme kynurenine aminotransferase II (KAT II)). KYNA formation, KAT II enzyme activity, and the effectiveness of PF-04859989 varied throughout the day, highlighting time as a key factor modulating KYNA brain metabolism. PF-04859989 reduced KYNA levels under most conditions. Our findings suggest that the timing of KYNA-targeted treatments should be carefully considered in therapeutic strategies to improve cognition and sleep in brain disorders.

Abstract:
Neurochemical imbalances, including elevations of the tryptophan metabolite kynurenic acid (KYNA), an endogenous antagonist of glutamatergic and cholinergic receptors, are linked to cognitive and sleep disturbances in psychiatric and neurocognitive disorders. Therapeutic strategies to reduce brain KYNA by inhibiting kynurenine aminotransferase II (KAT II) are under investigation. However, few studies consider time as a biological variable, despite recent evidence that the time of day can affect brain metabolism and drug effectiveness. Therefore, we explore the hypothesis that KYNA formation and synthesis inhibition change throughout the day. Using rats of both sexes, we measured basal KYNA levels and the effects of kynurenine (100 mg/kg, i.p.), to stimulate *de novo* KYNA, and/or PF-04859989 (KAT II inhibitor, 30 mg/kg, s.c.), at the beginning of light or dark phases. Microdialysis was used to assess extracellular KYNA in the dorsal hippocampus, and *ex vivo* assays evaluated KAT enzyme activity in separate animals. Additionally, we examined KYNA levels and the effects of PF-04859989 during acute sleep deprivation in male rats. PF-04859989 significantly decreased stimulated KYNA synthesis in both sexes and basal KYNA in males. In the dark phase, compared with the light phase, male rats treated with kynurenine and/or PF-04859989 showed higher KYNA levels and greater KYNA synthesis inhibition, respectively. *In vitro*, KAT II activity was similarly higher, and PF-04859989 was more effective, in the dark than in the light phase. Sleep deprivation increased extracellular KYNA levels, and PF-04859989 prevented this increase. Overall, our findings highlight the need to consider time-dependent factors when developing therapies impacting KYNA synthesis.

Keywords: psychiatric illness, circadian, kynurenine pathway, hippocampus, microdialysis, sleep deprivation, chronotherapy, Zeitgeber, dark phase, kynurenic acid

Introduction

Neurochemical imbalances are regularly observed in individuals with neuropsychiatric illnesses, neurocognitive disorders, and systemic infections, and are causally linked to pervasive symptoms like sleep disturbances and cognitive dysfunction in these conditions. Yet, current treatments do not effectively alleviate such symptoms for patients. Therefore, understanding the mechanisms underlying disrupted neurochemistry provides critical insight into developing new and effective therapeutics to address unresolved symptoms in patients.

Brain elevations in kynurenic acid (KYNA), a tryptophan metabolite of the kynurenine pathway (**Figure 1**), are observed in psychiatric patients with schizophrenia (SZ) and bipolar disorder (BPD),[1-5] neurology patients with neurocognitive dysfunction,[6-11] and individuals with acute inflammation in the central nervous system.[12-15] Mechanistic clinical and preclinical studies show that central KYNA levels increase in response to psychological stress,[16, 17] infection,[18-20] and sleep deprivation.[21, 22] KYNA acts as an endogenous antagonist of the α7 nicotinic

Figure 1: Schematic representation of the kynurenine pathway (KP) of tryptophan degradation, whereby tryptophan is degraded to kynurenine by tryptophan dioxygenase (TDO) or indoleamine dioxygenase (IDO). Subsequently, kynurenine is degraded to 3-hydroxykynurenine (3-HK) or kynurenic acid (KYNA) by kynurenine 3-monooxygenase (KMO) or kynurenine aminotransferase II (KAT II), respectively. PF-04859989 (blue) is an irreversible KAT II inhibitor.

acetylcholine (α7nACh) and N-methyl-d-aspartate (NMDA) glutamate receptors, thereby impacting neurochemical systems that play major roles in modulating cognitive processes and sleep homeostasis.[23, 24] Elevated KYNA levels in key brain regions, such as the hippocampus, which is critical for learning and sleep-dependent memory consolidation, may be causally related to cognitive and sleep disturbances in these conditions.[25-27] Preclinical studies support this association, as elevations in KYNA disrupt sleep architecture and hippocampal-dependent cognition, whereas reductions in KYNA improve these behaviors in rats.[28-32]

Treatment strategies to reduce brain KYNA levels are being pursued therapeutically (clinicaltrials.gov NCT0401355, NCT06225115).[33, 34] Translational studies have utilized pharmacological tools that inhibit the main KYNA-synthesizing enzyme in the brain, kynurenine aminotransferase II (KAT II), to successfully reduce brain KYNA, enhance cognitive function, and promote sleep behavior both at baseline and following physiological challenges.[31-33, 35-41] Yet, our understanding of the influence of circadian time on neuropharmacology as it relates to modulators of the kynurenine pathway remains limited.

A critical limitation in many studies is the omission of 'time' as a variable in the study design, despite recent literature highlighting the significant impact of time of day on brain metabolism and drug efficacy.[42, 43] Clinical work has shown conspicuous time-of-day differences in peripheral tryptophan metabolism and kynurenine pathway metabolite levels,[44-52] yet no studies have closely evaluated

changes in the brain across 24 hours. We currently use a rat model to fill gaps in our understanding of brain-specific kynurenine pathway metabolism and KYNA synthesis across the day.

Rodents experience 'active' and 'rest' periods throughout the day. In laboratory settings, when the time of day can be manipulated by controlling the timing of lights on and off, rodent studies are conducted under 12 h light cycles defined by Zeitgeber time (ZT). ZT 0 corresponds to lights on and ZT 12 to lights off. As nocturnal species, rodents spend most of their time asleep when the lights are on (ZT 0-12, "light phase") and most of their time awake when the lights are off (ZT 12-24, "dark phase"). We have shown that KYNA levels and KYNA-synthesizing enzyme activities in male and female rats change across the light and dark phases.[53, 54] It is, however, unclear whether central nervous system KYNA synthesis and pharmacological KAT II inhibition are affected by time of day, or how physiological challenges, such as enhanced availability of the bioprecursor kynurenine or acute sleep deprivation (SleepDep), affect this system. Understanding whether time of day affects brain KYNA synthesis can enhance our ability to develop chronotherapeutic approaches, and our present study provides the first steps in this direction. We presently determined *in vivo* and *ex vivo* changes in basal KYNA synthesis across the light and dark cycles and investigated the efficacy of KAT II inhibition with the irreversible, brain-penetrable KAT II inhibitor PF-04859989[35, 36] in both sexes of rats. Lastly, we explored KYNA synthesis after homeostatic challenge with SleepDep at the beginning of the light phase, demonstrating a rapid increase in extracellular KYNA that is inhibited by PF-04859989 under these conditions. Taken together, our study highlights ways that KYNA synthesis and inhibition are sensitive to the time of day.

Results

In males, the time of day influences both the effects of KAT II inhibition by PF-04859989 and the synthesis of brain KYNA from exogenous kynurenine

Since previous *in vivo* microdialysis studies on kynurenine pathway neurochemistry have been conducted primarily in male rats, we first examined the time-dependent *de novo* synthesis of KYNA in the dorsal hippocampus of male rats. Peripheral kynurenine challenge (100 mg/kg, to stimulate *de novo* KYNA synthesis in the brain) was administered at either ZT 0, ZT 6, ZT 12, or ZT 18, and extracellular KYNA levels were measured in the dorsal hippocampus. Time of injection significantly influenced the amount of KYNA produced ($F_{3, 23}$ = 4.503, $p < 0.05$), and we determined that kynurenine challenge at ZT 12, the beginning of the dark phase, produced twice as much KYNA in the hippocampus as peripheral kynurenine treatment at ZT 0 ($p < 0.01$), the start of the light phase (**Supplemental Figure 1**). We therefore conducted subsequent microdialysis studies to compare treatments at ZT 0 to ZT 12. Furthermore, as our previous studies have noted sex-specific findings in brain KYNA,[21, 54] we designed the present study to analyze the data separately by sex.

To compare the effects of pharmacologically manipulating KYNA synthesis, either through a kynurenine challenge (as above) or KAT II inhibition using the systemically active, brain penetrant, and irreversible KAT II inhibitor PF-04859989 (30 mg/kg; dose selected based on prior publications[31, 32, 35, 36]), at the start of the light phase (ZT 0) or dark phase (ZT 12), we included the following treatment groups: i) vehicle, ii) PF-04859989, iii) kynurenine, iv) PF-04859989 + kynurenine. Rats received PF-04859989 30 min prior to a kynurenine challenge at ZT 0 or ZT 12 (**Figure 2A**). We first assessed the impact of time on PF-04859989 treatment by comparing PF-04859989 and vehicle conditions. Time of day ("ZT"; light phase: $F_{2.779, 39.18}$ = 3.532, $p < 0.05$), PF-04859989 treatment (light phase: $F_{1, 16}$ = 4.494, $p = 0.05$; dark phase: $F_{1, 15}$ = 4.735, $p < 0.05$) and the interaction between time (ZT) and PF-04859989 treatment (dark phase: $F_{1.953, 24.41}$ = 4.772, $p < 0.05$) significantly influenced extracellular KYNA levels in the male dorsal hippocampus (**Figure 2B**). PF-04859989 treatment, irrespective of phase, consistently reduced KYNA levels over time compared to vehicle. Of note, KYNA levels decreased across the light

phase and increased across the dark phase under vehicle conditions. Area under the curve (AUC) analysis comparing light and dark phases revealed significant effects of phase ($F_{1, 38} = 4.585$, $p < 0.05$; saline post-hoc: $p = 0.0675$) and PF-04859989 treatment ($F_{1, 31} = 6.814$, $p < 0.05$; light phase post-hoc: $p = 0.1177$, dark phase post-hoc: $p < 0.01$) (**Figure 2C**). PF-04859989 reduced KYNA by 41-47%.

Next, we assessed whether KYNA synthesis from exogenous kynurenine (kynurenine challenge) and the effects of PF-04859989 pretreatment on this response are influenced by time of day. Kynurenine challenge transiently elevated KYNA levels in both the light and dark phases, and pretreatment with PF-04859989 significantly blunted this response (**Figure 2D**). When assessing each phase independently and comparing treatment groups across time (ZT), we determined significant ZT x treatment interactions (light phase: $F_{4.174, 45.91} = 10.06$, $p < 0.0001$; dark phase: $F_{6.680, 68.75} = 22.23$, $p < 0.0001$), as well as main effects of ZT (light phase: $F_{2.087, 45.91} = 26.29$, $p < 0.0001$; dark phase: $F_{2, 23} = 51.01$, $p < 0.0001$) and treatment (light phase: $F_{2, 24} = 17.22$, $p < 0.0001$; dark phase: $F_{3.340, 68.75} = 30.96$, $p < 0.0001$). AUC analysis revealed a significant phase x treatment interaction ($F_{2, 44} = 11.68$, $p < 0.0001$) and main effects of both phase ($F_{1, 44} = 8.761$, $p < 0.001$) and treatment ($F_{2, 44} = 56.46$, $p < 0.0001$) on KYNA levels (**Figure 2E**). Kynurenine challenge significantly elevated KYNA levels by 8-fold ($p < 0.0001$) during the light phase and 11.4-fold ($p < 0.0001$) during the dark phase relative to vehicle treatment. Notably, kynurenine challenge during the dark phase produced significantly greater *de novo* extracellular KYNA than during the light phase ($p < 0.0001$). Regardless of phase, PF-04859989 pretreatment significantly attenuated KYNA levels compared to kynurenine alone (light phase: $p < 0.05$, dark phase: $p < 0.0001$). However, when PF-04859989 was given prior to the dark phase, the reduction in *de novo* KYNA production following kynurenine challenge was more pronounced than when the KAT II inhibitor was given before the light phase.

In females, the ability of the KAT II inhibitor PF-04859989 to block kynurenine-induced KYNA synthesis in the brain is significantly influenced by the time of day

To expand our understanding of the impact of biological sex on KP metabolism, we conducted a microdialysis study in female rats (**Figure 3A**). To our knowledge, no prior studies have evaluated whether PF-04859989 can reduce basal extracellular KYNA levels in female rats. Surprisingly, our results demonstrate that the current dose of PF-04859989 fails to reduce basal KYNA levels in the dorsal hippocampus of female rats in either the light or dark phases (**Figure 3B**). Vehicle-treated or PF-04859989-treated females showed no differences in KYNA levels between the light and dark phases in our AUC analysis (**Figure 3C**). Additionally, KYNA levels in females were not significantly influenced by the estrous cycle (**Supplemental Figure 2**).

In females, kynurenine challenge transiently elevated KYNA levels, and pretreatment with PF-04859989 blunted this response regardless of phase (**Figure 3D**). KYNA levels were influenced by interactions between time (ZT) and treatment (light phase: $F_{3.070, 32.62} = 4.148$, $p < 0.05$; dark phase: $F_{5.889, 55.45} = 12.65$, $p < 0.0001$), as well as main effects of ZT (light phase: $F_{1.535, 32.62} = 15.26$, $p < 0.0001$; dark phase: $F_{2.944, 55.45} = 19.55$, $p < 0.0001$) and treatment (light phase: $F_{2, 22} = 6.043$, $p < 0.01$; dark phase: $F_{2, 19} = 31.30$, $p < 0.0001$). AUC analysis revealed that treatment ($F_{2, 40} = 36.14$, $p < 0.0001$), but not phase, affected kynurenine-induced KYNA synthesis and its inhibition by PF-04859989 in females (**Figure 3E**). Kynurenine challenge significantly increased KYNA 6.5-fold during the light phase ($p < 0.0001$) and 6.7-fold during the dark phase ($p < 0.0001$) compared to vehicle treatment. PF-04859989 pretreatment was more effective at preventing kynurenine-induced KYNA synthesis in the dark phase (PF-04859989 + kynurenine vs vehicle: not significant, $p > 0.05$) than in the light phase (PF-04859989 + kynurenine vs vehicle : $p < 0.01$).

Figure 2: In male rats, compared to light phase, dark phase kynurenine challenge (100 mg/kg, i.p) stimulates more extracellular kynurenic acid (KYNA) in the hippocampus, and pre-treatment with PF-04859989 (30 mg/kg, s.c.) inhibits KYNA levels to a greater extent. **(A)** Experimental timeline. Male rats were peripherally injected with kynurenine to stimulate *de novo* KYNA at the beginning of the light phase (Zeitgeber time (ZT) 0) or dark phase (ZT 12). Injection times are denoted by red arrows. PF-04859989, a systemically active brain-penetrant kynurenine aminotransferase (KAT II) inhibitor, was administered to inhibit KYNA synthesis 30 min prior (ZT 23.5 or ZT 11.5). Injection times are denoted by blue arrows. Extracellular KYNA collected via *in vivo* microdialysis in the dorsal hippocampus was evaluated in 30-minute fractions for the duration of the experiment. **(B)** Extracellular KYNA following vehicle or PF-04859989 treatment. **(C)** Area under the curve (AUC) of extracellular KYNA following vehicle or PF-04859989 treatment. **(D)** Extracellular KYNA following kynurenine challenge ± PF-04859989 pretreatment. **(E)** AUC of extracellular KYNA following kynurenine challenge ± PF-04859989 pretreatment. All data are mean ± SEM. Two-way RM ANOVA separated by phase (panels B/D) or two-way ANOVA (panels C/E): ^ $p < 0.05$, ^^ $p < 0.01$, ^^^^ $p < 0.0001$; Dunnett's post hoc test to vehicle (panel D) or Fisher's LSD post hoc test (panels B/C/E): *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. N = 7 - 10 per group.

Figure 3: In female rats, light and dark phase kynurenine challenge (100 mg/kg, i.p.) equivocally stimulates extracellular kynurenic acid (KYNA) in the hippocampus, yet PF-04859989 pretreatment (30 mg/kg, s.c.) inhibits KYNA levels to a greater extent in the dark phase. **(A)** Experimental timeline. Female rats were peripherally injected with kynurenine to stimulate *de novo* KYNA at the beginning of the light phase (Zeitgeber time (ZT) 0) or dark phase (ZT 12). Injection times are denoted by red arrows. PF-04859989, a systemically active brain-penetrant kynurenine aminotransferase (KAT II) inhibitor, was administered to inhibit KYNA synthesis 30 min prior (ZT 23.5 or ZT 11.5). Injection times are denoted by blue arrows. Extracellular KYNA collected via *in vivo* microdialysis in the dorsal hippocampus was evaluated in 30-minute fractions for the duration of the experiment. **(B)** Extracellular KYNA following vehicle or PF-04859989 treatment. **(C)** Area under the curve (AUC) of extracellular KYNA following vehicle or PF-04859989 treatment. **(D)** Extracellular KYNA following kynurenine challenge ± PF-04859989 pretreatment. **(E)** AUC of extracellular KYNA following kynurenine challenge ± PF-04859989 pretreatment. All data are mean ± SEM. Two-way RM ANOVA separated by phase (panels B/D) or two-way ANOVA (panels C/E): ^ $p < 0.05$, ^^ $p < 0.01$, ^^^^ $p < 0.0001$; Dunnett's post hoc test to vehicle (panel D) or Fisher's LSD post hoc test (panels B/C/E): *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. N = 6 – 9 per group.

Brain KAT II enzyme activity exhibits diurnal variation

We next sought to determine whether time-dependent alterations in KAT II activity underlie the previously observed differences in extracellular KYNA. To this end, we conducted *ex vivo* KAT enzyme activity assays in male and female brain homogenates collected every 3 hours for 24 hours (**Figure 4A**). We focused on the KAT II isoform, the predominant KYNA-synthesizing enzyme in the brain,[55] though we also conducted KAT I enzyme activity assays (**Supplemental Figure 3**). ZT significantly affected KAT II enzyme activity in both male ($F_{7, 24}$ = 6.428, $p < 0.001$; **Figure 4B**) and female ($F_{7, 24}$ = 5.495, $p < 0.001$; **Figure 4C**) samples, demonstrating broad variation in KAT II activity across the day.

We next compared KAT II activity at ZT 3 and ZT 15, the first tissues collected after the light and dark phase transitions, respectively, separately by sex. In males, KAT II enzyme activity at ZT 15 was significantly higher than at ZT 3 when assayed with 2 μM kynurenine (**Figure 4D,** $p < 0.05$). In contrast, no difference in KAT II activity was observed between these time points in females (**Figure 4E**).

We also assessed the ability of PF-04859989 to inhibit *ex vivo* KAT II activity at these time points in a dose-response manner. At all tested concentrations, PF-04859989 significantly reduced KAT II enzyme activity compared to the active control in males and females (**Supplemental Figure 4**). In males, both concentration ($F_{1.690, 8.451}$ = 831.9, $p < 0.0001$) and ZT ($F_{1, 5}$ = 10.62, $p < 0.05$) significantly influenced the percent inhibition of KAT II activity by PF-04859989 (**Figure 4F**). At 1 μM and 100 μM concentrations, PF-04859989 produced significantly greater inhibition of KAT II activity at ZT 15 compared to ZT 3 (1μM: $p < 0.05$, 100 μM: $p < 0.01$). In females, no differences in inhibition were observed between ZT 3 and ZT 15, although the expected concentration-dependent effect of PF-04859989 ($F_{1.822, 10.93}$ = 1176, $p < 0.0001$) was evident (**Figure 4G**).

Figure 4: Kynurenine aminotransferase II (KAT II) enzyme activity changes across the time of day. In males, KAT II enzyme activity and its inhibition by PF-04859989 are significantly altered between Zeitgeber time (ZT) 3 and ZT 15. (A) Experimental approach. Brain samples were collected from male and female rats at 3-hour intervals for *ex vivo* enzyme activity assays, which were assessed by measuring kynurenic acid (KYNA) production. **(B)** Male forebrain KAT II enzyme activity. **(C)** Female forebrain KAT II enzyme activity. **(D)** Male forebrain KAT II enzyme activity at ZT 3 and ZT 15. **(E)** Female forebrain KAT II enzyme activity at ZT 3 and ZT 15. **(F)** Male percent of KAT II inhibition with PF-04859989 at ZT 3 and ZT 15. **(G)** Female percent of KAT II inhibition with PF-04859989 at ZT 3 and ZT 15. All data are mean ± SEM. Two-way RM ANOVA (panels B/C/F/G): ^ $p < 0.05$, ^^ $p < 0.01$, ^^^ $p < 0.001$, ^^^^ $p < 0.0001$; Mann-Whitney tests (panels D/E) or Fisher's LSD post-hoc (panels F/G): *$p < 0.05$, **$p < 0.01$. N = 3 – 8 per group.

PF-04859989, KAT II inhibitor, prevents elevated extracellular KYNA induced by acute sleep deprivation (SleepDep) in males.

In addition to exogenous kynurenine, we investigated whether KAT II inhibition could prevent KYNA accumulation induced by a physiological challenge, such as acute SleepDep via gentle handling, which has previously been shown to stimulate brain KYNA synthesis in male but not female rats.[21] Thereby, we investigated whether PF-04859989 (30 mg/kg) could prevent SleepDep-induced elevations in KYNA levels in males (**Figure 5A**). SleepDep was restricted to the beginning of the light phase (ZT 0 - 6), when rats typically obtain most of their sleep, and sleep pressure is highest. Monitoring of extracellular KYNA was extended through ZT 12 to capture recovery from SleepDep.

Our analysis revealed that SleepDep ($F_{1,47}$ = 6.894, $p < 0.05$) and PF-04859989 ($F_{1,47}$ = 8.455, $p < 0.01$) significantly affected extracellular KYNA levels in the hippocampus (**Figure 5B**). Acute SleepDep elevated extracellular KYNA levels throughout most of the light phase compared with control, whereas PF-04859989 treatment prevented this increase. KYNA levels of sleep-deprived males treated with PF-04859989 were not significantly different from those of control (*ad libitum* sleep) vehicle-treated or control PF-04859989-treated males. With AUC analysis, an interaction between SleepDep and PF-04859989 ($F_{1,46}$ = 5.632, $p < 0.05$) and a main effect of PF-04859989 ($F_{1,46}$ = 10.15, $p < 0.01$) influenced KYNA levels (**Figure 5C**). Our post-hoc analysis confirmed that SleepDep elevated KYNA levels (control vs SleepDep: $p < 0.001$) and that pretreatment with PF-04859989 effectively blocked this increase (PF-04859989 + SleepDep vs SleepDep: $p < 0.001$).

Figure 5: Accumulation of extracellular KYNA with sleep deprivation (SleepDep) is prevented with PF-04859989 pretreatment in males. (A) Experimental timeline. Male rats were sleep-deprived from Zeitgeber time (ZT) 0 to ZT 6, as denoted by the green arrows, using a manual, gentle bar-sweeping approach. PF-04859989 (30 mg/kg, s.c.), a systemically active kynurenine aminotransferase II (KAT II) inhibitor, was administered 30 min before SleepDep (ZT 23.5) to prevent KYNA synthesis. Injection time is denoted by a blue arrow. For the duration of the experiment, microdialysis was performed in the dorsal hippocampus to assess extracellular KYNA levels. **(B)** Extracellular KYNA levels are shown as a percent change from baseline following control, PF-04859989, SleepDep, and SleepDep with PF-04859989. **(C)** Area under the curve (AUC) of extracellular KYNA. Three-way ANOVA (panel B) or two-way ANOVA (panel C): ^ $p < 0.05$, ^^ $p < 0.01$, ^^^ $p < 0.001$; Fisher's LSD post-hoc test: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. N = 8 – 17 per group.

Discussion

Our current study found that time of day significantly affected *de novo* KYNA production and the inhibition of KYNA synthesis by PF-04859989 in the rat brain. These time-of-day effects were more pronounced in males than in females; however, in both sexes, the activity of the KYNA-synthesizing enzymes KAT I and KAT II varied across the day. PF-04859989 treatment could ameliorate SleepDep-induced elevations of KYNA in male rats.

Extracellular hippocampal KYNA levels decreased across the light phase and increased across the dark phase in male rats, consistent with previously reported literature indicating that tryptophan and its metabolites exhibit circadian fluctuations both centrally and peripherally.[44-49, 52-54] Both tryptophan and kynurenine can cross the blood-brain barrier, and peripheral levels readily influence brain kynurenine pathway metabolism.[9] After stimulating KYNA levels via exogenous peripheral kynurenine administration, we presently consider brain-specific mechanisms underlying the diurnal changes in extracellular KYNA levels in the hippocampus.

The changes in extracellular KYNA we observed could be due to differences in efflux or transporter activity between the light and dark phases. Efflux of KYNA from the brain occurs via organic anion transporters, an active transport mechanism,[27, 56] and studies suggest that brain efflux is enhanced during sleep.[57, 58] Clearance mechanisms during sleep may thereby contribute to the reduction in extracellular KYNA levels that we observed across the light phase when animals were left undisturbed. In contrast, reduced brain efflux during waking may contribute to the increased KYNA levels we observed across the dark phase and in SleepDep experiments. As KYNA does not readily cross the blood-brain barrier (BBB) and is primarily synthesized from plasma kynurenine, which enters the brain via the large neutral amino acid transporter,[59, 60] circadian changes in transporter function or BBB permeability could influence brain substrate availability and, therefore, KYNA metabolism.[61-63] We were unable to detect extracellular kynurenine levels in our samples, so further investigation is warranted to directly test this hypothesis. Additionally, regulation of kynurenine and KYNA transfer into or out of the astrocytes via sodium-independent processes or organic anion transporters,[56, 59] respectively, could also contribute to these phase-dependent changes. It is also important to note that our current findings are based on extracellular microdialysis measurements; whole-brain or tissue KYNA levels may exhibit slightly different dynamics across the circadian cycle, reflecting distinct pools and regulatory mechanisms between intracellular and extracellular compartments.[56]

We also observed variation in KAT I and KAT II enzyme activity across the light and dark phases. *In vivo* findings showed a marked increase in KYNA formation during the dark phase compared to the light phase in males, whereas females did not exhibit this phase-dependent difference. E*x vivo* assays revealed greater KAT II activity at ZT 15 than at ZT 3 in male rat brains, accompanied by a corresponding decrease in KAT I activity. In females, KAT II activity did not differ substantially between these time points. Because KAT II contributes approximately 60% of *de novo* KYNA synthesis,[55, 64] we interpret that increased KAT II during the dark phase drives the elevated KYNA formation observed in males. The distinct biochemical properties of KAT I and KAT II, including differences in substrate preferences, pH optimum, utilization of α-keto acids, competing endogenous substrates, and enzymatic functions, may underlie their divergent circadian patterns.[27] Additionally, regulators of kynurenine pathway enzymes, such as steroids, growth factors, and immune signaling molecules, fluctuate with time of day[65, 66] and may further modulate brain kynurenine metabolism, though this remains to be tested in the brain.[67, 68]

Time-dependent changes in cellular conditions may further contribute to the circadian pattern of KYNA synthesis. Variations in glucose availability, oxidative stress, and reactive oxygen species (ROS) production across the light-dark cycle could influence both enzymatic and non-enzymatic KYNA

formation.[69] Enzymatic KYNA metabolism is highly dependent on glucose availability,[70] which varies as a function of sleep-wake states and, therefore, ZT.[69, 71] Kynurenine can also be converted to KYNA through non-enzymatic oxidative pathways mediated by ROS.[72] These reactive species are known to accumulate during sleep deprivation[73] and reach peak levels during the dark phase,[74] coinciding with the period of elevated KYNA levels observed in our study.

When we inhibited KAT II with PF-04859989, we observed a more pronounced blockade of *de novo* KYNA production after kynurenine challenge in the dark than in the light phase, suggesting that the drug or enzyme itself exhibits time-of-day-dependent properties. The *in vivo* findings could be attributed to increased brain drug transfer or cofactor availability during the dark phase.[58, 75] Yet *ex vivo* enzyme assays, in which drug and cofactor concentrations are controlled, also supported the notion that PF-04859989 inhibition of KYNA synthesis was greater in tissue collected during the dark phase than during the light phase. Therefore, we speculate that the drug's ability to covalently bind to the essential pyridoxal 5'-phosphate (PLP) cofactor in the KAT II active site may differ across time points.[76] Together, these findings suggest that dosing PF-04859989 before the active period may be most effective in reducing brain KYNA levels, yet further biochemical and functional studies are warranted to better understand the behavioral implications and, thereby, any time-of-day differences that may govern specific behaviors, such as sleep or cognition.

Differences in *de novo* KYNA production, particularly the effectiveness of PF-04859989, in female rats deserve further discussion. In females, PF-04859989 administered before a kynurenine challenge significantly reduced KYNA levels across phases; however, it failed to reduce basal extracellular KYNA levels. This lack of effect was not explained by differences in baseline KYNA between sexes, fluctuations in KYNA across estrous phases, or any deficiency in PF-04859989's ability to inhibit KAT II *ex vivo* in females. Estrogen and its phosphate and sulfur ester forms have been described as reversible inhibitors of KAT II[77, 78] and may compete with PF-04859989 for access to the enzyme's active site,[77] thereby limiting its inhibitory efficacy. It is therefore possible that higher doses of PF-04859989 might overcome this competition in females; however, increasing the dose could compromise KAT II selectivity.[36]

In addition to direct enzymatic effects, sex-dependent differences in compound and metabolite transport may contribute to these findings.[79] PF-04859989 crosses the BBB, however, BBB permeability is itself influenced by biological sex and sex hormones.[80, 81] Estrogen has been shown to decrease BBB permeability in reproductively intact female rats,[82] which could limit drug penetration into the brain during periods of elevated estrogen. Moreover, the transporters responsible for KYNA movement across cellular membranes, organic anion transporters (OATs),[56] exhibit sex differences in expression and activity in the kidneys,[83, 84] wherein renal cortical OAT1 and OAT3 expression is enhanced by androgens and suppressed by estrogens, suggesting a broader hormonal regulation that could extend to the brain and warrants further investigation in regard to the transport of KYNA.[85] Additionally, although time-of-day-dependent changes in these transports have not yet been described, such regulation could influence the extent to which KYNA is released into or cleared from the extracellular space. Together, these factors may interact to shape the sex-specific and potentially circadian dynamics of KYNA metabolism and pharmacological inhibition.

Interestingly, in females, KYNA synthesis from kynurenine was comparable between the light and dark phases, consistent with the absence of time-of-day differences in *ex vivo* KAT II enzyme activity. In addition to enzymatic conversion by KATs, KYNA can be synthesized through oxidative mechanisms.[72] Males often express higher ROS and oxidative stress than females, potentially due to estrogen's antioxidant and neuroprotective roles.[86, 87] Estrogen also acts as a regulator of circadian rhythmicity by modulating clock gene expression,[88] which may, in turn, influence KAT II expression and

activity in a time-of-day-dependent manner. Together, these factors may contribute to the absence of a clear circadian pattern in KYNA production observed in females.

Clinically, elevated KYNA levels,[1-14] along with sleep disruptions[89-91] and cognitive impairments,[92-94] are well-documented in neuropsychiatric and neurocognitive disorders, as well as in chronic infections. Dysfunction of the hippocampus, a brain region crucial for learning, memory, and sleep regulation, may underlie these symptoms. Because elevated KYNA has been implicated in cognitive and sleep abnormalities, [9, 25-27]our microdialysis studies conducted in the dorsal hippocampus provide translationally relevant insight into mechanisms that may contribute to clinical dysfunction.

Inhibiting KYNA synthesis is a promising therapeutic strategy for mitigating cognitive and sleep disturbances associated with psychiatric disease.[33] Preclinical evidence shows that KAT II inhibitors, including PF-04859989, reduce brain KYNA levels and improve cognition in male rats.[37-39] PF-04859989 has been shown to prevent stimulant-induced deficits in auditory gating, working memory, spatial memory, and sustained attention[36]; to attenuate KYNA-mediated impairments in fear discrimination after stress exposure[40]; and to facilitate long-term potentiation in mice with elevated KYNA.[41] Although these findings strongly support the compound's pro-cognitive potential, prior studies have largely focused on males. Further, considering that both cognitive performance and KYNA metabolism exhibit circadian variation, future behavioral studies of KAT II inhibition should incorporate time of day and biological sex as key experimental variables. Moreover, PF-04859989, administered at the onset of either the light or dark phase, enhances sleep in both sexes,[31] suggesting that KAT II inhibition may represent an effective chronotherapeutic approach to improve sleep quality.

In summary, our study demonstrates that the effects of PF-04859989 on KYNA synthesis are influenced by both time of day and biological sex. These data highlight the importance of circadian timing and sex-specific factors when evaluating KAT II inhibitors as potential treatments for disorders characterized by dysregulated KYNA metabolism.

Methods
Animals
Adult male and female Wistar rats (200 - 300g) were obtained from Charles River Laboratories. Animals received standard rodent chow *ad libitum* and were given at least one week to acclimate to the animal facility before being used in any experiment. Animals were kept on a 12/12-hour light/dark cycle, where lights on corresponded to Zeitgeber time (ZT) 0 and lights off to ZT 12. All animals were housed in a temperature-controlled facility fully accredited by the American Association for the Accreditation of Laboratory Animal Care. All protocols were approved by the Institutional Animal Care and Use Committee at the University of South Carolina School of Medicine and were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Chemicals
L-Kynurenine sulfate ("kynurenine," purity: 99.4%) was obtained from Sai Advantium (Hyderabad, India). PF-04959989 was obtained from WuXi AppTec (Shanghai, China). All other chemicals were obtained at the highest commercially available purity from various suppliers. PF-04859989 and kynurenine were prepared for injections daily by dissolving 10 mg/mL in ultrapure water or phosphate-buffered saline, respectively. For kynurenine, the pH was adjusted to 7.2-7.6 with 0.1 N sodium hydroxide. Vehicle injections were ultrapure water (s.c.) or saline (i.p.).

Experiment 1: PF-04859989 and Kynurenine Challenge Microdialysis

Surgery: Animals were placed on a stereotaxic frame (Stoelting Co. Wood Dale IL, USA) under isoflurane anesthesia (2 - 5%) to implant a guide cannula (1.0 mm outer diameter, SciPro Inc., Sanborn, NY, USA) over the dorsal hippocampus (AP: -3.4 mm; LM: ±2.3 mm; DV: -1.5 mm from bregma). Carprofen (5 mg/kg, s.c.) was given for analgesic support. Two surgical screws were inserted into 0.5 mm burr holes on the skull, and acrylic dental cement secured each cannula in place. Animals were given 24-48 hours of recovery before experiments.

Intrahippocampal Microdialysis: On the day of experiments, at ZT 22 or ZT 10, a microdialysis probe (2 mm PES membrane/14 mm shaft/6kD, SciPro Inc.) was inserted through the guide cannula and Ringer solution (147 mM NaCl, 4 mM KCl, 1.4 mM $CaCl_2$) was perfused at a flow rate of 2.5 μL/minute with a microperfusion pump (Harvard Apparatus, Holliston, MA, USA). After 30 minutes, the flow rate was reduced to 1.0 μL/minute for the remainder of the experiment. Samples were collected in 30-minute fractions in a refrigerated autosampler (CMA Microdialysis AB - Harvard Apparatus Inc., Holliston, MA, USA). After one hour of baseline collection, animals received either PF-04859989 (30 mg/kg, s.c.) or vehicle (s.c.) injections at ZT 23.5 (prior to the start of light phase) or ZT 11.5 (prior to the start of dark phase). Animals then received either kynurenine (100 mg/kg, i.p.) or vehicle (i.p.) injections at ZT 0 or ZT 12. Collection continued in 30-minute fractions for 5 hours post-injection. Dialysate samples were stored at -80˚C until biochemical analyses.

Placement Verification: At the end of the microdialysis experiment, brains were removed from euthanized animals and drop-fixed in 10% formalin solution to verify the correct placement of the guide cannula and probe. Brains were moved to 20% sucrose, then 40 μm coronal sections obtained on a cryostat (Microm HM 560, Leica BioSystems, Nussloch, Germany) containing the dorsal hippocampus were collected on gelatin-coated slides. Slides were stained with Neutral Red and evaluated under a light microscope (Laxco, SeBa Pro 4) to confirm probe placement.

Experiment 2: KAT Enzyme Activity Assays

Tissue Collection: Animals were euthanized via CO_2 asphyxiation every three hours across the 24-hour light/dark cycle, beginning at ZT 0. Brains were promptly removed, snap frozen on dry ice, and stored at -80°C.

KAT I/KAT II Assays: On the day of the assays, forebrains were thawed on wet ice and homogenized (1:5 w/v) in ultrapure water. A portion of the homogenate was diluted 1:10 with homogenization buffer (5 mM Tris acetate buffer pH 8.0 containing 50 μM pyridoxal 5'-phosphate (P5P) and 10 mM β-mercaptoethanol) and dialyzed against the homogenization buffer overnight at 4°C. The remaining 1:5 homogenate was stored at -80°C for later use. The next day, 80 μL of dialyzed homogenate (1:20 final dilution) was combined with 20 μL ultrapure water, and 100 μL reaction cocktail (KAT I: 2 μM or 100 μM kynurenine, 80 μM P5P, 80 μM pyruvate, and 150 mM AMP buffer pH 9.5; KAT II: 2 μM or 100 μM kynurenine, 80 μM P5P, 80 μM pyruvate, and 150 mM Tris acetate buffer pH 7.4) and incubated in a 37°C water bath for 2 hours. Blank controls were assessed by adding 20 μL of 1 mM aminooxyacetate, a general KAT inhibitor,[95] to the reaction instead of water. For assays with the KAT II inhibitor PF-04859989, 20 μL of water was replaced with 20 μL of PF-04859989 at the following concentrations: 100 nM, 1 μM, 10 μM, 100 μM. Following incubation, the tubes were placed on ice, and the reactions were stopped with 50 μL of 6% perchloric acid. The tubes were vortexed and centrifuged (14,000 RPM, 10

minutes), and the supernatant was collected. Samples were stored at -80°C until HPLC analysis for KYNA levels. Protein was evaluated in the homogenate using the Lowry method.[96]

Experiment 3: Sleep Deprivation (SleepDep) Microdialysis
Sleep deprivation (SleepDep) was conducted during microdialysis in a separate cohort of animals. Male rats received PF-04859989 (30 mg/kg, s.c.) or vehicle (ultrapure water, s.c.) at ZT 23.5 and were subjected to SleepDep from ZT 0 to ZT 6 for a total of 6 hours. Microdialysis chambers were fitted with custom acrylic arms, manually moved by the experimenter to gently sweep across the bottom of the cage and induce wakefulness in a modified gentle handling approach. When the experimenter observed a complete loss of movement or closed eyelids, the bar was manually swept. Microdialysis samples were collected in 30-minute fractions until ZT 6 and in 1-hour fractions from ZT 6 to ZT 12. Dialysate samples were stored at -80°C until biochemical analyses. Upon completion of the experiment, brains were collected and processed for probe placement, as described above.

HPLC Analysis of Kynurenic Acid (KYNA)
Twenty µL of microdialysate, diluted 1:2 in ultrapure water, or supernatant collected from the enzyme assays (diluted 1:10 for 2 µM kynurenine assays or 1:100 for 100 µM kynurenine assays in ultrapure water) were analyzed with a Reprosil-Pur C18 column (4 × 150 mm; Dr. MaischGmbh, Ammerbuch, Germany) using a 50 mM sodium acetate mobile phase (pH adjusted to 6.20 with glacial acetic acid) containing 5% acetonitrile at a flow rate of 0.5 mL/minute. KYNA (excitation 344 nm, emission 398 nm) was detected fluorometrically in the eluate (Waters Alliance, 2475 fluorescence detector, Bedford MA) with a post-column addition of 500 mM zinc acetate delivered at a flow rate of 0.1 mL/minute. Data was analyzed using Empower 3 software (Waters). Microdialysis data were not corrected for recovery from the microdialysis probe.

Statistics
Our studies investigating kynurenine pathway metabolism and extracellular KYNA in rats have noted sex-specific findings.[21, 53, 54] As such, we designed the present study to analyze the data separately by sex. Data normality was evaluated using the Shapiro–Wilk test and visually inspected with Q-Q plots to confirm an approximately bell-shaped distribution and the absence of outliers. All statistical analyses were performed using GraphPad Prism 10.5.0 software (GraphPad Software, La Jolla, CA, USA) with statistical significance defined as $p < 0.05$.

Experiment 1: PF-04859989 microdialysis data (figures 2B/3B) were analyzed separately by phase (light phase beginning at ZT 0 or dark phase beginning at ZT 12) and sex (male or female) using two-way repeated measures (RM) ANOVAs with ZT as a within-subject factor and treatment (vehicle or PF-04859989) as a between-subject factor. Significant main effects were followed up with Fisher's LSD post hoc tests. Area under the curve analysis (Figures 2C/3C), calculated with data points (as above) relative to Y = 0, was analyzed separately by sex using two-way ANOVAs with phase (light or dark) and treatment (vehicle or PF-04859989) as between-subject factors. For male AUC, any subjects with undetectable KYNA levels by ZT 5 or ZT 17 were not included for AUC analysis (1 light phase and 1 dark phase PF-04859989-treated animals). Significant main effects were followed up with Fisher's LSD post hoc tests. Kynurenine challenge microdialysis data (Figures 2D/3D) were analyzed for all collected data points separately by phase (light or dark) and sex (male or female), using two-way RM ANOVAs with ZT as a within-subject factor and treatment (vehicle, kynurenine, or PF-04859989 + kynurenine) as a between-subject factor. Significant main effects were followed up with Dunnett's post hoc tests to

vehicle. Area under the curve (Figures 2E/3E), calculated with all measured data points relative to Y=0, was analyzed separately by sex using two-way ANOVAs with phase (light or dark) and treatment (vehicle, kynurenine, or PF-04859989 + kynurenine) as between-subject factors. Significant main effects were followed up with Fisher's LSD post hoc tests. Female microdialysis data grouped by estrous phase (Supplemental Figure 2) were analyzed with light and dark phase data combined, as no significant phase effects were observed in the female data.

Experiment 2: KAT II activity data (Figure 4B/4C) were analyzed by two-way RM ANOVAs with kynurenine concentration as a within-subject factor and ZT as a between-subject factor. Enzyme activity data comparing ZT 3 and ZT 15 (Figure 4D/4E) were analyzed with Mann-Whitney tests at each concentration. PF-04859989 KAT II inhibition data (Figure 4F/4G) were analyzed by two-way RM ANOVAs with PF-04859989 concentration as a within-subject factor and ZT as a between-subject factor. Significant main effects of ZT were followed up with Fisher's LSD post-hoc tests.

Experiment 3: Microdialysis data from SleepDep experiments were calculated as a percentage of basal KYNA and analyzed with a three-way RM ANOVA with ZT as a within-subject factor and SleepDep and PF-04859989 as between-subject factors. Significant main effects of SleepDep or PF-04859989 were followed up with two-way RM ANOVAs separated by SleepDep or PF-04859989 and Fisher's LSD post-hoc tests compared to control. The area under the microdialysis data curve, calculated using all measured data points relative to Y = 100, was analyzed using a two-way ANOVA with SleepDep and PF-04859989 as between-subject factors, followed by Fisher's LSD post hoc tests.

Safety: This work is not associated with any unexpected, new, or significant hazards or risks.

Abbreviations: schizophrenia (SZ), bipolar disorder (BPD), $\alpha 7$ nicotinic acetylcholine (α7nACh), N-methyl-d-aspartate (NMDA), kynurenic acid (KYNA), sleep deprivation (SleepDep), Zeitgeber time (ZT), kynurenine aminotransferase (KAT), reactive oxygen species (ROS), organic anion transporter (OAT), blood-brain barrier (BBB), pyridoxal 5'-phosphate (PLP).

Author Contributions: **Courtney J. Wright**: Investigation, methodology, formal analysis, visualization, writing – original draft, writing – review and editing. **Silas A. Buck**: Investigation, methodology, formal analysis, writing – review and editing. **Snezana Milosavljevic**: Investigation, formal analysis, writing – review and editing. **Ashley M. Lewis**: Investigation, methodology, formal analysis. **Nathan T.J. Wagner**: Investigation, methodology, formal analysis, visualization, writing – original draft. **Ana Pocivavsek**: Investigation, methodology, formal analysis, visualization, writing – original draft, writing – review and editing, conceptualization, funding acquisition, project administration, supervision, resources.

Funding Sources: This work was supported by National Institutes of Health Grant Nos. R01 HL174802, R21 AG080335, and P50 MH103222.

Conflict of Interest: The authors declare no conflicts of interest.

Data Availability: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Bibliography

(1) Erhardt, S.; Blennow, K.; Nordin, C.; Skogh, E.; Lindström, L. H.; Engberg, G. Kynurenic acid levels are elevated in the cerebrospinal fluid of patients with schizophrenia. *Neuroscience Letters* **2001**, *313* (1-2), 96-98. DOI: 10.1016/s0304-3940(01)02242-x.

(2) Schwarcz, R.; Rassoulpour, A.; Wu, H.-Q.; Medoff, D.; Tamminga, C. A.; Roberts, R. C. Increased cortical kynurenate content in schizophrenia. *Biological Psychiatry* **2001**, *50* (7), 521-530. DOI: 10.1016/s0006-3223(01)01078-2.

(3) Miller, C. L.; Llenos, I. C.; Dulay, J. R.; Weis, S. Upregulation of the initiating step of the kynurenine pathway in postmortem anterior cingulate cortex from individuals with schizophrenia and bipolar disorder. *Brain Research* **2006**, *1073-1074*, 25-37. DOI: 10.1016/j.brainres.2005.12.056.

(4) Olsson, S. K.; Samuelsson, M.; Saetre, P.; Lindström, L.; Jönsson, E. G.; Nordin, C.; Engberg, G.; Erhardt, S.; Landén, M. Elevated levels of kynurenic acid in the cerebrospinal fluid of patients with bipolar disorder. *Journal of Psychiatry and Neuroscience* **2010**, *35* (3), 195-199. DOI: 10.1503/jpn.090180.

(5) Sellgren, C. M.; Gracias, J.; Jungholm, O.; Perlis, R. H.; Engberg, G.; Schwieler, L.; Landen, M.; Erhardt, S. Peripheral and central levels of kynurenic acid in bipolar disorder subjects and healthy controls. *Transl Psychiatry* **2019**, *9* (1), 37. DOI: 10.1038/s41398-019-0378-9.

(6) Sinz, E. H.; Kochanek, P. M.; Heyes, M. P.; Wisniewski, S. R.; Bell, M. J.; Clark, R. S.; DeKosky, S. T.; Blight, A. R.; Marion, D. W. Quinolinic acid is increased in CSF and associated with mortality after traumatic brain injury in humans. *J Cereb Blood Flow Metab* **1998**, *18* (6), 610-615. DOI: 10.1097/00004647-199806000-00002.

(7) Baran, H.; Jellinger, K.; Deecke, L. Kynurenine metabolism in Alzheimer's disease. *J Neural Transm (Vienna)* **1999**, *106* (2), 165-181. DOI: 10.1007/s007020050149.

(8) Ting, K. K.; Brew, B.; Guillemin, G. The involvement of astrocytes and kynurenine pathway in Alzheimer's disease. *Neurotoxicity Research* **2007**, *12* (4), 247-262. DOI: 10.1007/bf03033908.

(9) Schwarcz, R.; Bruno, J. P.; Muchowski, P. J.; Wu, H. Q. Kynurenines in the mammalian brain: when physiology meets pathology. *Nat Rev Neurosci* **2012**, *13* (7), 465-477. DOI: 10.1038/nrn3257.

(10) Yan, E. B.; Frugier, T.; Lim, C. K.; Heng, B.; Sundaram, G.; Tan, M.; Rosenfeld, J. V.; Walker, D. W.; Guillemin, G. J.; Morganti-Kossmann, M. C. Activation of the kynurenine pathway and increased production of the excitotoxin quinolinic acid following traumatic brain injury in humans. *J Neuroinflammation* **2015**, *12*, 110. DOI: 10.1186/s12974-015-0328-2.

(11) González-Sánchez, M.; Jiménez, J.; Narváez, A.; Antequera, D.; Llamas-Velasco, S.; Martín, A. H.; Arjona, J. A. M.; Munain, A. L.; Bisa, A. L.; Marco, M. P.; et al. Kynurenic Acid Levels are Increased in the CSF of Alzheimer's Disease Patients. *Biomolecules* **2020**, *10* (4). DOI: 10.3390/biom10040571.

(12) Atlas, A.; Gisslén, M.; Nordin, C.; Lindström, L.; Schwieler, L. Acute psychotic symptoms in HIV-1 infected patients are associated with increased levels of kynurenic acid in cerebrospinal fluid. *Brain Behav Immun* **2007**, *21* (1), 86-91. DOI: 10.1016/j.bbi.2006.02.005.

(13) Atlas, A.; Franzen-Röhl, E.; Söderlund, J.; Jönsson, E. G.; Samuelsson, M.; Schwieler, L.; Sköldenberg, B.; Engberg, G. Sustained elevation of kynurenic Acid in the cerebrospinal fluid of patients with herpes simplex virus type 1 encephalitis. *Int J Tryptophan Res* **2013**, *6*, 89-96. DOI: 10.4137/IJTR.S13256.

(14) Cihan, M.; Doğan, Ö.; Ceran Serdar, C.; Altunçekiç Yıldırım, A.; Kurt, C.; Serdar, M. A. Kynurenine pathway in Coronavirus disease (COVID-19): Potential role in prognosis. *J Clin Lab Anal* **2022**, *36* (3), e24257. DOI: 10.1002/jcla.24257.

(15) Li, X.; Edén, A.; Malwade, S.; Cunningham, J. L.; Bergquist, J.; Weidenfors, J. A.; Sellgren, C. M.; Engberg, G.; Piehl, F.; Gisslen, M.; et al. Central and peripheral kynurenine pathway metabolites in COVID-19: Implications for neurological and immunological responses. *Brain Behav Immun* **2025**, *124*, 163-176. DOI: 10.1016/j.bbi.2024.11.031.

(16) Notarangelo, F. M.; Schwarcz, R. Restraint Stress during Pregnancy Rapidly Raises Kynurenic Acid Levels in Mouse Placenta and Fetal Brain. *Dev Neurosci* **2016**, *38* (6), 458-468. DOI: 10.1159/000455228.

(17) Dugan, A. M.; Parrott, J. M.; Redus, L.; Hensler, J. G.; O'Connor, J. C. Low-Level Stress Induces Production of Neuroprotective Factors in Wild-Type but Not BDNF+/- Mice: Interleukin-10 and Kynurenic Acid. *Int J Neuropsychopharmacol* **2015**, *19* (3), pyv089. DOI: 10.1093/ijnp/pyv089.

(18) Larsson, M. K.; Faka, A.; Bhat, M.; Imbeault, S.; Goiny, M.; Orhan, F.; Oliveros, A.; Ståhl, S.; Liu, X. C.; Choi, D. S.; et al. Repeated LPS Injection Induces Distinct Changes in the Kynurenine Pathway in Mice. *Neurochem Res* **2016**, *41* (9), 2243-2255. DOI: 10.1007/s11064-016-1939-4.

(19) Zheng, Y. R.; Tufvesson-Alm, M.; Trepci, A.; Imbeault, S.; Li, X. Q.; Schwieler, L.; Engberg, G.; Erhardt, S. Dual administration of lipopolysaccharide induces behavioural changes in rats relevant to psychotic disorders. *Acta Neuropsychiatr* **2023**, *37*, e18. DOI: 10.1017/neu.2023.40.

(20) Raison, C. L.; Dantzer, R.; Kelley, K. W.; Lawson, M. A.; Woolwine, B. J.; Vogt, G.; Spivey, J. R.; Saito, K.; Miller, A. H. CSF concentrations of brain tryptophan and kynurenines during immune stimulation with IFN-alpha: relationship to CNS immune responses and depression. *Mol Psychiatry* **2010**, *15* (4), 393-403. DOI: 10.1038/mp.2009.116.

(21) Baratta, A. M.; Buck, S. A.; Buchla, A. D.; Fabian, C. B.; Chen, S.; Mong, J. A.; Pocivavsek, A. Sex Differences in Hippocampal Memory and Kynurenic Acid Formation Following Acute Sleep Deprivation in Rats. *Sci Rep* **2018**, *8* (1), 6963. DOI: 10.1038/s41598-018-25288-w.

(22) Baratta, A. M.; Kanyuch, N. R.; Cole, C. A.; Valafar, H.; Deslauriers, J.; Pocivavsek, A. Acute sleep deprivation during pregnancy in rats: Rapid elevation of placental and fetal inflammation and kynurenic acid. *Neurobiol Stress* **2020**, *12*, 100204. DOI: 10.1016/j.ynstr.2019.100204.

(23) Birch, P. J.; Grossman, C. J.; Hayes, A. G. Kynurenic acid antagonises responses to NMDA via an action at the strychnine-insensitive glycine receptor. *Eur J Pharmacol* **1988**, *154* (1), 85-87. DOI: 10.1016/0014-2999(88)90367-6.

(24) Hilmas, C.; Pereira, E. F.; Alkondon, M.; Rassoulpour, A.; Schwarcz, R.; Albuquerque, E. X. The brain metabolite kynurenic acid inhibits alpha7 nicotinic receptor activity and increases non-alpha7 nicotinic receptor expression: physiopathological implications. *J Neurosci* **2001**, *21* (19), 7463-7473. DOI: 10.1523/JNEUROSCI.21-19-07463.2001.

(25) Pocivavsek, A.; Rowland, L. M. Basic Neuroscience Illuminates Causal Relationship Between Sleep and Memory: Translating to Schizophrenia. *Schizophr Bull* **2018**, *44* (1), 7-14. DOI: 10.1093/schbul/sbx151.

(26) Sapienza, J.; Spangaro, M.; Guillemin, G. J.; Comai, S.; Bosia, M. Importance of the dysregulation of the kynurenine pathway on cognition in schizophrenia: a systematic review of clinical studies. *Eur Arch Psychiatry Clin Neurosci* **2023**, *273* (6), 1317-1328. DOI: 10.1007/s00406-022-01519-0.

(27) Pocivavsek, A.; Schwarcz, R.; Erhardt, S. Neuroactive Kynurenines as Pharmacological Targets: New Experimental Tools and Exciting Therapeutic Opportunities. *Pharmacol Rev* **2024**, *76* (6), 978-1008. DOI: 10.1124/pharmrev.124.000239.

(28) Chess, A. C.; Landers, A. M.; Bucci, D. J. L-kynurenine treatment alters contextual fear conditioning and context discrimination but not cue-specific fear conditioning. *Behav Brain Res* **2009**, *201* (2), 325-331. DOI: 10.1016/j.bbr.2009.03.013.

(29) Pocivavsek, A.; Wu, H. Q.; Potter, M. C.; Elmer, G. I.; Pellicciari, R.; Schwarcz, R. Fluctuations in endogenous kynurenic acid control hippocampal glutamate and memory. *Neuropsychopharmacology* **2011**, *36* (11), 2357-2367. DOI: 10.1038/npp.2011.127.

(30) Pocivavsek, A.; Baratta, A. M.; Mong, J. A.; Viechweg, S. S. Acute Kynurenine Challenge Disrupts Sleep-Wake Architecture and Impairs Contextual Memory in Adult Rats. *Sleep* **2017**, *40* (11). DOI: 10.1093/sleep/zsx141.

(31) Milosavljevic, S.; Smith, A. K.; Wright, C. J.; Valafar, H.; Pocivavsek, A. Kynurenine aminotransferase II inhibition promotes sleep and rescues impairments induced by neurodevelopmental insult. *Transl Psychiatry* **2023**, *13* (1), 106. DOI: 10.1038/s41398-023-02399-1.

(32) Rentschler, K. M.; Milosavljevic, S.; Baratta, A. M.; Wright, C. J.; Piroli, M. V.; Tentor, Z.; Valafar, H.; O'Reilly, C.; Pocivavsek, A. Reducing brain kynurenic acid synthesis precludes kynurenine-induced sleep disturbances. *J Sleep Res* **2023**, e14038. DOI: 10.1111/jsr.14038.

(33) Pocivavsek, A.; Erhardt, S. Kynurenic acid: translational perspectives of a therapeutically targetable gliotransmitter. *Neuropsychopharmacology* **2024**, *49* (1), 307-308. DOI: 10.1038/s41386-023-01681-6.

(34) Fukuwatari, T. Possibility of Amino Acid Treatment to Prevent the Psychiatric Disorders via Modulation of the Production of Tryptophan Metabolite Kynurenic Acid. *Nutrients* **2020**, *12* (5). DOI: 10.3390/nu12051403.

(35) Koshy Cherian, A.; Gritton, H.; Johnson, D. E.; Young, D.; Kozak, R.; Sarter, M. A systemically-available kynurenine aminotransferase II (KAT II) inhibitor restores nicotine-evoked glutamatergic activity in the cortex of rats. *Neuropharmacology* **2014**, *82*, 41-48. DOI: 10.1016/j.neuropharm.2014.03.004.

(36) Kozak, R.; Campbell, B. M.; Strick, C. A.; Horner, W.; Hoffmann, W. E.; Kiss, T.; Chapin, D. S.; McGinnis, D.; Abbott, A. L.; Roberts, B. M.; et al. Reduction of brain kynurenic acid improves cognitive function. *J Neurosci* **2014**, *34* (32), 10592-10602. DOI: 10.1523/JNEUROSCI.1107-14.2014.

(37) Wu, H. Q.; Okuyama, M.; Kajii, Y.; Pocivavsek, A.; Bruno, J. P.; Schwarcz, R. Targeting kynurenine aminotransferase II in psychiatric diseases: promising effects of an orally active enzyme inhibitor. *Schizophr Bull* **2014**, *40 Suppl 2* (Suppl 2), S152-158. DOI: 10.1093/schbul/sbt157.

(38) Bortz, D. M.; Wu, H. Q.; Schwarcz, R.; Bruno, J. P. Oral administration of a specific kynurenic acid synthesis (KAT II) inhibitor attenuates evoked glutamate release in rat prefrontal cortex. *Neuropharmacology* **2017**, *121*, 69-78. DOI: 10.1016/j.neuropharm.2017.04.023.

(39) Pocivavsek, A.; Elmer, G. I.; Schwarcz, R. Inhibition of kynurenine aminotransferase II attenuates hippocampus-dependent memory deficit in adult rats treated prenatally with kynurenine. *Hippocampus* **2019**, *29* (2), 73-77. DOI: 10.1002/hipo.23040.

(40) Klausing, A. D.; Fukuwatari, T.; Bucci, D. J.; Schwarcz, R. Stress-induced impairment in fear discrimination is causally related to increased kynurenic acid formation in the prefrontal cortex. *Psychopharmacology (Berl)* **2020**, *237* (7), 1931-1941. DOI: 10.1007/s00213-020-05507-x.

(41) Imbeault, S.; Gubert Olivé, M.; Jungholm, O.; Erhardt, S.; Wigström, H.; Engberg, G.; Jardemark, K. Blockade of KAT II Facilitates LTP in Kynurenine 3-Monooxygenase Depleted Mice. *Int J Tryptophan Res* **2021**, *14*, 11786469211041368. DOI: 10.1177/11786469211041368.

(42) Ruben, M. D.; Smith, D. F.; FitzGerald, G. A.; Hogenesch, J. B. Dosing time matters. *Science* **2019**, *365* (6453), 547-549. DOI: 10.1126/science.aax7621.

(43) Ayyar, V. S.; Sukumaran, S. Circadian rhythms: influence on physiology, pharmacology, and therapeutic interventions. *Journal of Pharmacokinetics and Pharmacodynamics* **2021**, *48* (3), 321-338. DOI: 10.1007/s10928-021-09751-2.

(44) Rapoport, M. I.; Beisel, W. R. Circadian periodicity of tryptophan metabolism. *J Clin Invest* **1968**, *47* (4), 934-939. DOI: 10.1172/JCI105785.

(45) Munõz-Hoyos, A.; Molina-Carballo, A.; Rodríguez-Cabezas, T.; Uberos-Fernández, J.; Ruiz-Cosano, C.; Acuña-Castroviejo, D. Relationships between methoxyindole and kynurenine pathway metabolites in plasma and urine in children suffering from febrile and epileptic seizures. *Clin Endocrinol (Oxf)* **1997**, *47* (6), 667-677. DOI: 10.1046/j.1365-2265.1997.2991136.x.

(46) Muñoz-Hoyos, A.; Molina-Carballo, A.; Macías, M.; Rodríguez-Cabezas, T.; Martín-Medina, E.; Narbona-López, E.; Valenzuela-Ruiz, A.; Acuña-Castroviejo, D. Comparison between tryptophan methoxyindole and kynurenine metabolic pathways in normal and preterm neonates and in neonates with acute fetal distress. *Eur J Endocrinol* **1998**, *139* (1), 89-95. DOI: 10.1530/eje.0.1390089.

(47) de Jong, W. H.; Smit, R.; Bakker, S. J.; de Vries, E. G.; Kema, I. P. Plasma tryptophan, kynurenine and 3-hydroxykynurenine measurement using automated on-line solid-phase extraction HPLC-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* **2009**, *877* (7), 603-609. DOI: 10.1016/j.jchromb.2009.01.015.

(48) Coggan, S. E.; Smythe, G. A.; Bilgin, A.; Grant, R. S. Age and circadian influences on picolinic acid concentrations in human cerebrospinal fluid. *Journal of Neurochemistry* **2009**, *108* (5), 1220-1225. DOI: 10.1111/j.1471-4159.2009.05868.x.

(49) Molina-Carballo, A.; Cubero-Millán, I.; Fernández-López, L.; Checa-Ros, A.; Machado-Casas, I.; Jerez-Calero, A.; Blanca-Jover, E.; Cantarero-Malagón, A.-M.; Uberos, J.; Muñoz-Hoyos, A.

Methylphenidate ameliorates the homeostatic balance between levels of kynurenines in ADHD children. *Psychiatry Research* **2021**, *303*, 114060. DOI: 10.1016/j.psychres.2021.114060.

(50) Moravcová, S.; Spišská, V.; Pačesová, D.; Hrubcová, L.; Kubištová, A.; Novotný, J.; Bendová, Z. Circadian control of kynurenine pathway enzymes in the rat pineal gland, liver, and heart and tissue- and enzyme-specific responses to lipopolysaccharide. *Arch Biochem Biophys* **2022**, *722*, 109213. DOI: 10.1016/j.abb.2022.109213.

(51) Stincardini, C.; Pariano, M.; D'Onofrio, F.; Renga, G.; Orecchini, E.; Orabona, C.; Nunzi, E.; Gargaro, M.; Fallarino, F.; Chun, S. K.; et al. The circadian control of tryptophan metabolism regulates the host response to pulmonary fungal infections. *PNAS Nexus* **2023**, *2* (3), pgad036. DOI: 10.1093/pnasnexus/pgad036.

(52) Johanns, M.; Haas, J. T.; Raverdy, V.; Vandel, J.; Chevalier-Dubois, J.; Guille, L.; Derudas, B.; Legendre, B.; Caiazzo, R.; Verkindt, H.; et al. Time-of-day-dependent variation of the human liver transcriptome and metabolome is disrupted in MASLD. *JHEP Rep* **2024**, *6* (1), 100948. DOI: 10.1016/j.jhepr.2023.100948.

(53) Rentschler, K. M.; Baratta, A. M.; Ditty, A. L.; Wagner, N. T. J.; Wright, C. J.; Milosavljevic, S.; Mong, J. A.; Pocivavsek, A. Prenatal Kynurenine Elevation Elicits Sex-Dependent Changes in Sleep and Arousal During Adulthood: Implications for Psychotic Disorders. *Schizophr Bull* **2021**, *47* (5), 1320-1330. DOI: 10.1093/schbul/sbab029.

(54) Wright, C. J.; Rentschler, K. M.; Wagner, N. T. J.; Lewis, A. M.; Beggiato, S.; Pocivavsek, A. Time of Day-Dependent Alterations in Hippocampal Kynurenic Acid, Glutamate, and GABA in Adult Rats Exposed to Elevated Kynurenic Acid During Neurodevelopment. *Front Psychiatry* **2021**, *12*, 734984. DOI: 10.3389/fpsyt.2021.734984.

(55) Guidetti, P.; Hoffman, G. E.; Melendez-Ferro, M.; Albuquerque, E. X.; Schwarcz, R. Astrocytic localization of kynurenine aminotransferase II in the rat brain visualized by immunocytochemistry. *Glia* **2007**, *55* (1), 78-92. DOI: 10.1002/glia.20432.

(56) Schwarcz, R.; Stone, T. W. The kynurenine pathway and the brain: Challenges, controversies and promises. *Neuropharmacology* **2017**, *112* (Pt B), 237-247. DOI: 10.1016/j.neuropharm.2016.08.003.

(57) Xie, L.; Kang, H.; Xu, Q.; Chen, M. J.; Liao, Y.; Thiyagarajan, M.; O'Donnell, J.; Christensen, D. J.; Nicholson, C.; Iliff, J. J.; et al. Sleep drives metabolite clearance from the adult brain. *Science* **2013**, *342* (6156), 373-377. DOI: 10.1126/science.1241224.

(58) Zhang, S. L.; Lahens, N. F.; Yue, Z.; Arnold, D. M.; Pakstis, P. P.; Schwarz, J. E.; Sehgal, A. A circadian clock regulates efflux by the blood-brain barrier in mice and human cells. *Nat Commun* **2021**, *12* (1), 617. DOI: 10.1038/s41467-020-20795-9.

(59) Segawa, H.; Fukasawa, Y.; Miyamoto, K.; Takeda, E.; Endou, H.; Kanai, Y. Identification and functional characterization of a Na+-independent neutral amino acid transporter with broad substrate selectivity. *J Biol Chem* **1999**, *274* (28), 19745-19751. DOI: 10.1074/jbc.274.28.19745.

(60) Sekine, A.; Kuroki, Y.; Urata, T.; Mori, N.; Fukuwatari, T. Inhibition of Large Neutral Amino Acid Transporters Suppresses Kynurenic Acid Production Via Inhibition of Kynurenine Uptake in Rodent Brain. *Neurochem Res* **2016**, *41* (9), 2256-2266. DOI: 10.1007/s11064-016-1940-y.

(61) Cuddapah, V. A.; Zhang, S. L.; Sehgal, A. Regulation of the Blood-Brain Barrier by Circadian Rhythms and Sleep. *Trends Neurosci* **2019**, *42* (7), 500-510. DOI: 10.1016/j.tins.2019.05.001.

(62) Pulido, R. S.; Munji, R. N.; Chan, T. C.; Quirk, C. R.; Weiner, G. A.; Weger, B. D.; Rossi, M. J.; Elmsaouri, S.; Malfavon, M.; Deng, A.; et al. Neuronal Activity Regulates Blood-Brain Barrier Efflux Transport through Endothelial Circadian Genes. *Neuron* **2020**, *108* (5), 937-952.e937. DOI: 10.1016/j.neuron.2020.09.002.

(63) Kim, M.; Keep, R. F.; Zhang, S. L. Circadian Rhythms of the Blood-Brain Barrier and Drug Delivery. *Circ Res* **2024**, *134* (6), 727-747. DOI: 10.1161/CIRCRESAHA.123.323521.

(64) Guillemin, G. J.; Kerr, S. J.; Smythe, G. A.; Smith, D. G.; Kapoor, V.; Armati, P. J.; Croitoru, J.; Brew, B. J. Kynurenine pathway metabolism in human astrocytes: a paradox for neuronal protection. *J Neurochem* **2001**, *78* (4), 842-853. DOI: 10.1046/j.1471-4159.2001.00498.x.

(65) Oster, H.; Challet, E.; Ott, V.; Arvat, E.; de Kloet, E. R.; Dijk, D. J.; Lightman, S.; Vgontzas, A.; Van Cauter, E. The Functional and Clinical Significance of the 24-Hour Rhythm of Circulating Glucocorticoids. *Endocr Rev* **2017**, *38* (1), 3-45. DOI: 10.1210/er.2015-1080.

(66) Chen, R.; Routh, B. N.; Gaudet, A. D.; Fonken, L. K. Circadian Regulation of the Neuroimmune Environment Across the Lifespan: From Brain Development to Aging. *J Biol Rhythms* **2023**, *38* (5), 419-446. DOI: 10.1177/07487304231178950.

(67) Haroon, E.; Raison, C. L.; Miller, A. H. Psychoneuroimmunology meets neuropsychopharmacology: translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology* **2012**, *37* (1), 137-162. DOI: 10.1038/npp.2011.205.

(68) Mithaiwala, M. N.; Santana-Coelho, D.; Porter, G. A.; O'Connor, J. C. Neuroinflammation and the Kynurenine Pathway in CNS Disease: Molecular Mechanisms and Therapeutic Implications. *Cells* **2021**, *10* (6). DOI: 10.3390/cells10061548.

(69) Dash, M. B.; Bellesi, M.; Tononi, G.; Cirelli, C. Sleep/wake dependent changes in cortical glucose concentrations. *J Neurochem* **2013**, *124* (1), 79-89. DOI: 10.1111/jnc.12063.

(70) Gramsbergen, J. B.; Hodgkins, P. S.; Rassoulpour, A.; Turski, W. A.; Guidetti, P.; Schwarcz, R. Brain-specific modulation of kynurenic acid synthesis in the rat. *J Neurochem* **1997**, *69* (1), 290-298. DOI: 10.1046/j.1471-4159.1997.69010290.x.

(71) Naylor, E.; Aillon, D. V.; Barrett, B. S.; Wilson, G. S.; Johnson, D. A.; Harmon, H. P.; Gabbert, S.; Petillo, P. A. Lactate as a biomarker for sleep. *Sleep* **2012**, *35* (9), 1209-1222. DOI: 10.5665/sleep.2072.

(72) Blanco Ayala, T.; Lugo Huitrón, R.; Carmona Aparicio, L.; Ramírez Ortega, D.; González Esquivel, D.; Pedraza Chaverrí, J.; Pérez de la Cruz, G.; Ríos, C.; Schwarcz, R.; Pérez de la Cruz, V. Alternative kynurenic acid synthesis routes studied in the rat cerebellum. *Front Cell Neurosci* **2015**, *9*, 178. DOI: 10.3389/fncel.2015.00178.

(73) Tian, Y.; Liu, X.; Liu, D. Physiological roles of reactive oxygen species in sleep homeostasis. *Free Radic Biol Med* **2025**, *242*, 220-236. DOI: 10.1016/j.freeradbiomed.2025.10.273.

(74) Logan, R. W.; McClung, C. A. Rhythms of life: circadian disruption and brain disorders across the lifespan. *Nat Rev Neurosci* **2019**, *20* (1), 49-65. DOI: 10.1038/s41583-018-0088-y.

(75) Walker, W. H.; Sprowls, S. A.; Bumgarner, J. R.; Liu, J. A.; Meléndez-Fernández, O. H.; Walton, J. C.; Lockman, P. R.; DeVries, A. C.; Nelson, R. J. Circadian Influences on Chemotherapy Efficacy in a Mouse Model of Brain Metastases of Breast Cancer. *Front Oncol* **2021**, *11*, 752331. DOI: 10.3389/fonc.2021.752331.

(76) Dounay, A. B.; Anderson, M.; Bechle, B. M.; Campbell, B. M.; Claffey, M. M.; Evdokimov, A.; Evrard, E.; Fonseca, K. R.; Gan, X.; Ghosh, S.; et al. Discovery of Brain-Penetrant, Irreversible Kynurenine Aminotransferase II Inhibitors for Schizophrenia. *ACS Med Chem Lett* **2012**, *3* (3), 187-192. DOI: 10.1021/ml200204m.

(77) Mason, M.; Manning, B. Effects of steroid conjugates on availability of pyridoxal phosphate for kynureninase and kynurenine aminotransferase activity. *Am J Clin Nutr* **1971**, *24* (7), 786-791. DOI: 10.1093/ajcn/24.7.786.

(78) Jayawickrama, G. S.; Nematollahi, A.; Sun, G.; Gorrell, M. D.; Church, W. B. Inhibition of human kynurenine aminotransferase isozymes by estrogen and its derivatives. *Sci Rep* **2017**, *7* (1), 17559. DOI: 10.1038/s41598-017-17979-7.

(79) Morris, M. E.; Lee, H. J.; Predko, L. M. Gender differences in the membrane transport of endogenous and exogenous compounds. *Pharmacol Rev* **2003**, *55* (2), 229-240. DOI: 10.1124/pr.55.2.1.

(80) Sohrabji, F. Guarding the blood-brain barrier: a role for estrogen in the etiology of neurodegenerative disease. *Gene Expr* **2007**, *13* (6), 311-319. DOI: 10.3727/000000006781510723.

(81) Dalla, C.; Pavlidi, P.; Sakelliadou, D. G.; Grammatikopoulou, T.; Kokras, N. Sex Differences in Blood-Brain Barrier Transport of Psychotropic Drugs. *Front Behav Neurosci* **2022**, *16*, 844916. DOI: 10.3389/fnbeh.2022.844916.

(82) Bake, S.; Sohrabji, F. 17beta-estradiol differentially regulates blood-brain barrier permeability in young and aging female rats. *Endocrinology* **2004**, *145* (12), 5471-5475. DOI: 10.1210/en.2004-0984.

(83) Ljubojevic, M.; Herak-Kramberger, C. M.; Hagos, Y.; Bahn, A.; Endou, H.; Burckhardt, G.; Sabolic, I. Rat renal cortical OAT1 and OAT3 exhibit gender differences determined by both androgen stimulation and estrogen inhibition. *Am J Physiol Renal Physiol* **2004**, *287* (1), F124-138. DOI: 10.1152/ajprenal.00029.2004.

(84) Breljak, D.; Brzica, H.; Sweet, D. H.; Anzai, N.; Sabolic, I. Sex-dependent expression of Oat3 (Slc22a8) and Oat1 (Slc22a6) proteins in murine kidneys. *Am J Physiol Renal Physiol* **2013**, *304* (8), F1114-1126. DOI: 10.1152/ajprenal.00201.2012.

(85) Brzica, H.; Abdullahi, W.; Reilly, B. G.; Ronaldson, P. T. Sex-specific differences in organic anion transporting polypeptide 1a4 (Oatp1a4) functional expression at the blood-brain barrier in Sprague-Dawley rats. *Fluids Barriers CNS* **2018**, *15* (1), 25. DOI: 10.1186/s12987-018-0110-9.

(86) Kander, M. C.; Cui, Y.; Liu, Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med* **2017**, *21* (5), 1024-1032. DOI: 10.1111/jcmm.13038.

(87) Martínez de Toda, I.; González-Sánchez, M.; Díaz-Del Cerro, E.; Valera, G.; Carracedo, J.; Guerra-Pérez, N. Sex differences in markers of oxidation and inflammation. Implications for ageing. *Mech Ageing Dev* **2023**, *211*, 111797. DOI: 10.1016/j.mad.2023.111797.

(88) Alvord, V. M.; Kantra, E. J.; Pendergast, J. S. Estrogens and the circadian system. *Semin Cell Dev Biol* **2022**, *126*, 56-65. DOI: 10.1016/j.semcdb.2021.04.010.

(89) Winkelman, J. W.; Lecea, L. Sleep and neuropsychiatric illness. *Neuropsychopharmacology* **2020**, *45* (1), 1-2. DOI: 10.1038/s41386-019-0514-5.

(90) Zhang, Y.; Ren, R.; Yang, L.; Zhang, H.; Shi, Y.; Okhravi, H. R.; Vitiello, M. V.; Sanford, L. D.; Tang, X. Sleep in Alzheimer's disease: a systematic review and meta-analysis of polysomnographic findings. *Transl Psychiatry* **2022**, *12* (1), 136. DOI: 10.1038/s41398-022-01897-y.

(91) Ferrarelli, F. Sleep disturbance in schizophrenia spectrum disorders: more than just a symptom? *Int Clin Psychopharmacol* **2023**, *38* (3), 187-188. DOI: 10.1097/YIC.0000000000000467.

(92) McCutcheon, R. A.; Keefe, R. S. E.; McGuire, P. K. Cognitive impairment in schizophrenia: aetiology, pathophysiology, and treatment. *Mol Psychiatry* **2023**, *28* (5), 1902-1918. DOI: 10.1038/s41380-023-01949-9.

(93) Hampshire, A.; Elliott, P. Cognition and Memory after Covid-19 in a Large Community Sample. Reply. *N Engl J Med* **2024**, *390* (21), 2035. DOI: 10.1056/NEJMc2403996.

(94) Lee, M.; Cullen, A. E.; Matheson, G. J.; Lu, Z. A.; Bergen, S. E.; Sellgren, C. M.; Erhardt, S.; Fatouros-Bergman, H.; Cervenka, S. Cognition before and after psychosis onset: A naturalistic study of change, heterogeneity, and prognosis. *Schizophr Res Cogn* **2025**, *42*, 100387. DOI: 10.1016/j.scog.2025.100387.

(95) John, R. A.; Charteris, A. The reaction of amino-oxyacetate with pyridoxal phosphate-dependent enzymes. *Biochem J* **1978**, *171* (3), 771-779. DOI: 10.1042/bj1710771.

(96) LOWRY, O. H.; ROSEBROUGH, N. J.; FARR, A. L.; RANDALL, R. J. Protein measurement with the Folin phenol reagent. *J Biol Chem* **1951**, *193* (1), 265-275.

TITLE: Biochemical Regulation of Brain Kynurenic Acid Synthesis and Inhibition in Rats is Sensitive to the Time of Day

Courtney J. Wright[1,2], Silas A. Buck[3], Snezana Milosavljevic[1,2], Ashley M. Lewis[1], Nathan T.J. Wagner[1], Ana Pocivavsek*[1,2,3]

[1]Department of Pharmacology, Physiology, and Neuroscience, University of South Carolina School of Medicine, Columbia, SC, USA
[2]Carolina Autism and Neurodevelopment Center, University of South Carolina, Columbia, SC, USA
[3]Maryland Psychiatric Research Center, Department of Psychiatry, University of Maryland School of Medicine, Baltimore, MD, USA

Corresponding Author:
Ana Pocivavsek, PhD
Pharmacology, Physiology, and Neuroscience
USC School of Medicine
Columbia, SC 29209
Fax: 803-216-3500
Email: ana.pocivavsek@uscmed.sc.edu

Supplemental Figure 1: Time of day differentially affects KYNA production following an acute kynurenine challenge, and maximal *de novo* KYNA synthesis occurs at the start of the dark phase. Kynurenine (100 mg/kg, i.p.) was administered at Zeitgeber time (ZT) 0, ZT 6, ZT 12, and ZT 18 in adult male Wistar rats. Extracellular KYNA, collected via in vivo microdialysis, was evaluated in the dorsal hippocampus for the duration of the experiment. **(A)** Extracellular KYNA (two-way RM ANOVA, sample number x time of injection: $F_{9.820, 71.47}$ = 3.250, ^^ p < 0.01, sample number: $F_{3.273, 71.47}$ = 55.29, ^^^^ p < 0.0001; time of injection: $F_{3, 23}$ = 5.346, ^^ p < 0.01). **(B)** Area under the curve (AUC) of extracellular KYNA (one-way ANOVA, $F_{3, 23}$ = 4.503, ^ p < 0.05). Data are mean ± SEM and Fisher's LSD post-hoc test: ** p < 0.01. N = 5 - 10 per group.

Supplemental Figure 2: Estrous cycle stage does not impact basal KYNA, response to kynurenine challenge, or PF-04859989 treatment in females. (A) Extracellular KYNA in females (light and dark phase data combined) were evaluated in females with high circulating estrogen (**proestrus/estrus, P/E**) upon kynurenine or PF-04859989 + kynurenine treatment (two-way RM ANOVA, treatment x ZT: $F_{12, 137} = 7.899$, ^^^^ $p < 0.0001$, treatment: $F_{1, 12} = 12.38$, ^^ $p < 0.01$, ZT: $F_{2.133, 24.36} = 24.81$, ^^^^ $p < 0.0001$). **(B)** Extracellular KYNA in females (light and dark phase data combined) were evaluated in females with low circulating estrogen (**metestrus/diestrus, M/D**) upon kynurenine or PF-04859989 + kynurenine treatment (two-way ANOVA, treatment x ZT: $F_{2.586}$, 34.70 = 2.803, $p = 0.0616$, ZT: $F_{2.586, 34.70} = 20.49$, ^^^^ $p < 0.0001$, treatment: $F_{1,14} = 4.403$, $p = 0.0545$). **(C)** Area under the curve (AUC) of extracellular KYNA in females treated with kynurenine or PF-04859989 + kynurenine, separated by P/E or M/D (two-way ANOVA, treatment: $F_{1, 26} = 16.11$, ^^^ $p < 0.001$). **(D)** Extracellular KYNA levels in P/E females with vehicle or PF-04859989 alone (two-way RM ANOVA, ZT: $F_{4.957, 62.79} = 2.497$, ^ $p < 0.05$). **(E)** Extracellular KYNA levels in M/D females treated with vehicle or PF-04859989 alone (two-way RM ANOVA). **(F)** AUC of extracellular KYNA in females treated with vehicle or PF-04859989 alone, separated by P/E or M/D (two-way ANOVA). All data are mean ± SEM. Fisher's LSD post-hoc: *$p < 0.05$, **$p < 0.01$. N = 5 - 8 per group.

Supplemental Figure 3: Kynurenine aminotransferase I (KAT I) enzyme activity changes across Zeitgeber time (ZT). (A) Experimental approach. Brain samples were collected from male and female rats at 3-hour intervals for *ex vivo* enzyme activity assays, KAT I and KAT II, which were assessed by measuring kynurenic acid (KYNA) production. **(B)** Forebrain KAT I enzyme activity assay, 2 μM kynurenine (two-way ANOVA, ZT x sex: $F_{7, 48}$ = 4.252, p < 0.01; ZT: $F_{7, 48}$ = 10.79, p < 0.0001). **(C)** Forebrain KAT I enzyme activity assay, 100 μM kynurenine (two-way ANOVA, ZT x sex: $F_{7, 48}$ = 5.199, p < 0.001, ZT: $F_{7, 48}$ = 30.68, p < 0.0001, sex: $F_{1, 48}$ = 4.337, p < 0.05). All data are mean ± SEM. N = 3 – 4 per group.

Supplemental Figure 4: PF-04859989 significantly inhibits kynurenine aminotransferase (KAT) II enzyme activity in a dose-dependent manner. (A) Male forebrain percent activity KAT II with increasing concentrations of PF-04859989 (two-way ANOVA, concentration: $F_{2.065, 10.33}$ = 1516, ^^^^ $p < 0.0001$, ZT: $F_{1, 5}$ = 10.62, ^ $p < 0.05$). **(B)** Female forebrain percent activity KAT II with increasing concentrations of PF-04859989 (two-way ANOVA, concentration: $F_{1.978, 11.87}$ = 2221, ^^^^ $p < 0.0001$). All data are mean ± SEM and Dunnett's post-hoc comparison to 0 nM PF-04859989. Post-hoc significance denotes the least significant ZT 3 or ZT 15 comparison between 0 nM and subsequent PF-04859989 concentrations: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. N = 3 – 4 per group.