μ-Opioid Receptor Superagonists Engage a Sodium-Bound Active State

Author Names: *Alexander Powell[1], Nicholas W. Griggs[1], Jorge Iniguez-Lluhi[1], John R. Traynor[1,2]*

*denotes 1st author

Author Address: [1]Department of Pharmacology, Edward F. Domino Research Center, Medical School, University of Michigan, Ann Arbor, MI 48109, United States

[2]Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, Ann Arbor, MI 48109, United States

Corresponding Author: John R. Traynor (jtraynor@umich.edu | 734-647-7479)

1st Author Email: powelaj@umich.edu

Abstract

The conformational selection model of G-protein coupled receptor (GPCR) activation suggests that, by binding to a high affinity site, an agonist will shift receptor equilibrium in favor of active state (R*) conformations that recruit heterotrimeric G proteins over inactive state (R) conformations. Agonist binding to the μ-opioid receptor is highly sensitive to sodium ions which stabilize an inactive receptor state. Higher efficacy opioid agonists, such as DAMGO and fentanyl, are sensitive to Na^+ compared to lower efficacy ligands at the mu-opioid receptor (μOR). However, the binding of the highly potent oripavine agonists etorphine and dihydroetorphine are less sensitive to Na^+ than expected such that the prevailing models fail to explain their pharmacology. In an attempt to explain this discrepancy, experiments were performed to evaluate the binding properties and G protein activation of the highly potent agonists carfentanil, BU72, etorphine, etonitazene and similarly potent opioid peptidomimetics in comparison to the standard agonists DAMGO, fentanyl, and morphine in the presence or absence of Na^+ or K^+. In agreement with prior studies on etorphine, several of the superagonists showed minimal affinity shifts and retained high potency in both ionic conditions, whereas conventional agonists DAMGO, fentanyl and morphine displayed enhanced

binding and signaling only in K^+. These functional parameters were used to determine an intrinsic efficacy

value, determined as $Log(\frac{Max}{EC50})$. Comparison of affinity shifts with intrinsic efficacy gives a negative

correlation in which superagonists with the highest intrinsic efficacy are least sensitive to Na^+. These data

suggest that select μ-opioid receptor superagonists have high affinity for the Na^+ bound receptor (R) and

shift these species into active receptor conformations (R*) that efficiently couple to G proteins.

Significance Statement

The prevailing theory of conformational selection is that high efficacy ligands, such as fentanyl and

DAMGO, are more sensitive to Na^+ and guanine nucleotide compared to lower efficacy agonists. Here,

instead of a positive correlation and what has previously thought to be true, there is a negative correlation

in which the ligands with the highest intrinsic efficacy (colloquially termed as superagonists) are the least

sensitive to Na^+ and guanine nucleotide and thus show a smaller affinity shift. Thus, demonstrating that

these highly potent ligands are capable of engaging a Na^+-bound receptor that may then efficiently couple

to G protein. This work has broad implications for GPCR pharmacology – challenging foundational models

of class A GPCR activation and providing framework for designing next-generation GPCR therapeutics

that retain high affinity and potency under physiological ionic conditions.

Key words: Opioid, GPCR, Mu-opioid Receptor, Intrinsic Efficacy, Affinity, Conformation Selection

Theory

Abbreviations: CHO, Chinese hamster ovary; DAMGO, [D-Ala^2, NMe-Phe^4, Gly-ol^5]-enkephalin; GDP,

Guanosine 5'-diphosphate; GPCR, G protein-coupled receptor; GTPγ^{35}S, (Guanosine-5'-O-(γ-thio)

triphosphate-[^{35}S]); μOR, mu-opioid receptor.

Introduction

Mu-opioid receptors (μOR) are G protein-coupled receptor (GPCR) proteins that activate heterotrimeric G proteins of the $G\alpha_{i/o}$ family. The mu-opioid receptors, like other GPCRs, exist in an equilibrium of various conformational states that can be grouped into ensembles of inactive (R) and active states (R*); the R* state interacting with the G protein α-subunit. Intrinsic efficacy describes the ability of a ligand to shift the equilibrium from inactive-state conformations to active-state conformations. This parameter distinguishes ligands that prefer the "low affinity" R state (inverse agonists) from those that prefer a "high affinity" R* state (partial agonists and full agonists) and from those that display no preference (neutral antagonists). Intrinsic efficacy, in conjunction with binding affinity and pharmacokinetic properties, are the driving factors that determine the pharmacological activity of a drug for a particular receptor system.[1]

Stimulation of GTPγ[35]S binding to G proteins is a sensitive indicator of the active R* state and may be used as a predictive measure of the intrinsic efficacy of a ligand.[2] GTPγ[35]S binding assays allow for the determination of potency (EC_{50}) and relative efficacy as that is maximal stimulation by a ligand compared to a standard agonist value. Determination of a $Log\left(\frac{Max}{EC50}\right)$ from these data is a convenient way to quantify the degree of agonism for a ligand.[3] One advantage of using these assays to assess ligand efficacy is that partial agonists are aptly discriminated from full agonists, since G protein activation is directly downstream of ligand-receptor binding and is less susceptible to signal amplification. On the other hand, and like other functional assays that rely on measurement of signal transduction events downstream of the receptor, results are dependent on experimental conditions.[4,5]

Agonist-mediated activation of opioid receptors is negatively modulated by the binding of Na^+ ions.[6,7,8] These ions bind deep in the receptor below the orthosteric site and form ionic interactions with transmembrane domains 2, 3, and 7, holding the receptor in an inactive set of conformations.[9] Thus, agonists and Na^+ ions work in opposition; agonists shift the equilibrium toward active R* ensembles

whereas sodium ions shift the equilibrium toward inactive R ensembles.[9,10,11] This negative cooperativity

has been long known at μOR, in which the addition of Na^+ and guanine nucleotide, to uncouple G proteins,

decreases the binding affinity (K_i) of opioid agonists.[6,12] Higher efficacy opioid agonists show a greater

decrease in binding affinity upon the addition of Na^+ and guanine nucleotide compared to lower efficacy

agonists and antagonists display no shift in affinity.[6,13] There appears to be a positive association between

the decrease in the K_i of an agonist by Na^+ (i.e. sodium sensitivity) and the intrinsic efficacy of an opioid.

However, two very potent opioid agonists, etorphine and dihydroetorphine, demonstrate significantly

reduced sodium sensitivity which questions the validity of the observed positive association between

sodium sensitivity and intrinsic efficacy.[13]

From first principles, the effects of sodium on μOR activation cannot be treated as uniform or universal.

Sodium may stabilize multiple inactive conformations, but its allosteric influence should be considered

probe-dependent and vary with the specific agonist or allosteric modulator engaged. Different ligands

may couple to the receptor in distinct ways that either disrupt or accommodate sodium binding. For

superagonists, it is possible that receptor activation does not require complete displacement of the sodium

ion; instead, these ligands may stabilize sodium-bound intermediate receptor ensembles that are still

competent to recruit G proteins. Thus, receptor activation may proceed through several parallel

trajectories, with some involving sodium displacement and others proceeding through Na^+-bound "active-

like" states that facilitate G protein coupling. These observations suggest that sodium's role in GPCR

activation should not be considered strictly inhibitory, but rather as part of an ensemble of conformations

whose functional relevance depends on the ligand's mode of allosteric engagement.

Here, we report the determination of intrinsic efficacy for a series potent opioids, including carfentanil,

BU72, etorphine and etonitazene, together with a series of highly potent opioid peptidomimetics and

compare them with standard opioids morphine, DAMGO and fentanyl (**Figure 1**). By evaluating the affinity

of these ligands in the absence and presence of Na^+ and calculating the difference in affinity between the

two conditions (i.e. "affinity shift"), we confirm that for highly potent opioid agonists, reduced Na^+ sensitivity correlates significantly with higher intrinsic efficacy. The prevailing model would predict the compounds to be neutral antagonists or low efficacy agonists, as they demonstrate little affinity shifts and recognize different affinity states in a near equal manner. Instead, we develop a model that shows how highly potent agonists bind to a low-affinity state of μOR, even in the presence of Na^+ ions, and shift the receptor into a high-affinity G protein-coupled state, explaining the exceptional potency of these compounds. Present results suggest that a simple conformational selection model of agonist receptor interaction controlled by Na^+ ions may not be adequate to explain the receptor kinetics of highly potent opioid agonists.

Methods

Materials and Reagents

We have previously reported the synthesis and pharmacological evaluation of novel opioid peptidomimetic compounds.[14,15,16,17,18,19,20] AAH8 was synthesized by Aubrie A. Harland, AFN42 was synthesized by Anthony F. Nastase, AMB46 and AMB47 were synthesized by Aaron M. Bender.[14,15,20] All chemicals, unless otherwise specified, were purchased from Sigma Aldrich. Radiolabeled [^{3}H]-diprenorphine (DPN) and radiolabeled guanosine-5′-O-(3-[^{35}S] thio)-triphosphate (GTPγ^{35}S) were purchased from PerkinElmer Life Sciences. All tissue culture media, penicillin-streptomycin, geneticin (G148), trypsin, and fetal bovine serum were purchased from Gibco LifeSciences.

Cell Lines

The tissue culture and maintenance of C6 rat glioma cells stably transfected with rat μOR was performed as previously described.[21] C6 cells were grown to confluence at 37 °C in 5 % CO_2 in Dulbecco's modified Eagle media (DMEM) containing 10 % fetal bovine serum (FBS), 5 % penicillin/streptomycin (P/S), and 400 ug/mL G418. CHO cells stably expressing wild-type human μOR were grown to confluence at 37 °C in 5 % CO_2 in Dulbecco's modified Eagle media (DMEM) containing 10 % fetal bovine serum (FBS), 1 % penicillin/streptomycin (P/S), and 400 ug/mL G418.

Cell Membrane Preparation

Cell membrane homogenates from C6 and CHO cells were prepared by washing confluent cells three times with phosphate buffered saline (pH 7.4), after which they were then detached using harvesting buffer (20 mM HEPES, 150 mM NaCl, 0.68 mM EDTA, pH7.4) and pelleted by centrifugation at 200 x g for 3 min at room temperature. The pellet was resuspended in ice-cold 50 mM Tris buffer (pH 7.4) and homogenized using a Tissue Tearor (Biospec Products, Inc.). The cell lysate was centrifuged at 20,000 x g at 4 °C for 20 minutes, after which the pellet was then resuspended on ice cold 50 mM Tris (pH7.4), homogenized, and centrifuged once more at 20000 x g at 4 C for 20 minutes. The final pellet was resuspended in 50 mM Tris (pH 7.4) using a glass dounce homogenizer and then dispensed into aliquots and stored at -80 °C. The final protein concentration was determined by performing a BCA protein assay (Thermo Scientific Pierce) using bovine serum albumin as the standard.

Competition Radioligand Binding Assays

^{3}H-diprenorphine competition binding assays were performed in 500μL reactions in 96-well plates using the cell membrane homogenates described above. Briefly, cell membranes [5-20 μg] were incubated in 50 mM Tris buffer containing 0.2 nM ^{3}H-Diprenorphine, and various concentrations of opioid compound for 90 minutes in a shaking water bath at room temperature to reach equilibrium. Nonspecific binding was determined using 10 μM of the opioid antagonist, naloxone. The reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue harvester (Brandel, Gaithersburg, MD) and washed with 50 mM Tris buffer. Filters were dried and following the addition of EcoLume scintillation cocktail, bound radioactivity was quantified in a MicroBeta 2450 Liquid Scintillation and Luminescence Counter (PerkinElmer).

Radioligand binding assays in Na$^+$ or K$^+$ containing buffers were performed in 200 μL reactions in 96-well plates using the cell membrane homogenates described above, Briefly, cell membranes [5-20 μg] were incubated in Na$^+$ or K$^+$ containing buffers (50 mM Tris, 100 mM NaCl or 100 mM KCl, 5 mM MgCl2, 1 mM EDTA, pH 7.4) or (50 mM Tris, 300 mM NaCl, 5 mM MgCl2, 1 mM EDTA, pH 7.4)

containing 10 μM cold GTPγS, and various concentrations of opioid compound for 90 minutes in a

shaking water bath at room temperature. Nonspecific binding was determined using 10 μM naloxone. The

reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue harvester

and washed with Na^+ containing buffer, and the filters were processed as described above.

Stimulation of GTPγ^{35}S Binding Assays

Stimulation of GTPγ^{35}S binding experiments were performed in 200 μL reactions in 96-well plates using

cell membrane homogenates prepared as described above. Briefly, cell membranes [10-20 μg] were

incubated in either Na^+ or K^+ containing buffer [50 mM Tris, 100 mM NaCl or 100mM KCl, 5 mM

MgCl2, 1 mM EDTA, pH 7.4] containing 0.1 nM guanosine-5′-O-(3-[^{35}S]thio)triphosphate (GTPγ^{35}S)

and 30 μM guanosine 5'-diphosphate (GDP) and various concentrations of opioid compound or

peptidomimetic for 1 hour in a shaking water bath at 25 °C. Basal stimulation of GTPγ^{35}S binding was

determined by incubation in the absence of any ligand and maximal stimulation of was defined by using

10 μM of the standard mu opioid receptor agonist, DAMGO ([D-Ala^2,N-MePhe^4,Gly-ol]-enkephalin).

The reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue

harvester and washed with Na^+ containing buffer, and filters were processed as described above.

Results

GTPγ^{35}S Binding in Sodium

As evaluated by concentration response curves of GTPγ^{35}S binding in C6 μOR membranes, etorphine, BU72, carfentanil and etonitazene are very potent with EC$_{50}$ values in the low nanomolar and sub-nanomolar range (**Table 1 and Figure 2**). In addition, we have previously described a series of highly potent peptidomimetics that bind to μOR with exceptional affinity.[20,22] They are at least 70 times more potent than the standard opioid peptide DAMGO, fentanyl, or morphine with potencies that are similar to that of the highly potent opioid agonists, etorphine, BU72, carfentanil and etonitazene. All compounds demonstrated a high relative efficacy, as evaluated by maximal stimulation of GTPγ^{35}S binding. The fractional maximal response (i.e. "Max") compared to DAMGO (Max = 1) was nearly equivalent for all ligands, except for carfentanil and AFN42, which demonstrate supramaximal μOR stimulation.

Binding Affinities in 50 mM Tris, 100 mM Na$^+$ and Guanine Nucleotide, or 300 mM Na$^+$ and Guanine Nucleotide

It has been known for decades that physiologically relevant Na$^+$ concentrations decrease the binding affinity of agonists to μOR.[23] Radioligand competition binding assays were performed using C6 μOR cell membranes in the absence or presence of Na$^+$ ions and guanine nucleotide to induce R and R* receptor states. Affinity shifts were then determined (**Table 1 and Figure 2**). In 50 mM Tris, the superagonists carfentanil, BU72, etorphine and etonitazene bound with roughly 10-fold higher affinity than the classical agonists DAMGO, morphine and fentanyl. The peptidomimetics also exhibited binding affinities similar to etorphine, BU72, carfentanil and etonitazene, with AAH8 displaying the highest affinity.

μOR binding affinities of DAMGO, fentanyl, and morphine decreased in the presence of 100 mM Na$^+$ and guanine nucleotide, producing affinity shifts of 131, 52 and 77-fold, respectively (**Table 1**). Of the superagonists, etorphine and etonitazene displayed the greatest affinity shifts of 24 and 33-fold,

respectively. Carfentanil, AAH8 and AMB47 all displayed 10-fold decreases in affinity, whereas BU72,

AFN42 and AMB47 displayed little shift.

Since the superagonists in question displayed a less than expected affinity shift in 100 mM Na^+ and guanine

nucleotide, we utilized a supraphysiological concentration of Na^+ to determine whether the affinity shift for

each superagonist was saturated. I (**Table** 1). In the presence of 300 mM Na^+ and guanine nucleotide, an

affinity shift of 1,240-fold was observed for DAMGO, whereas fentanyl and morphine displayed smaller

shifts of 105 and 84-fold, respectively. Of the highly potent agonists, only etonitazene produced an affinity

shift greater than 100-fold. Etorphine, AAH8 and AMB46 had comparable affinity shifts ranging from 29

to 67-fold. Carfentanil and AMB47 displayed smaller shifts of 14 and 15-fold, respectively. Two

compounds exhibited shifts of less than 10-fold, these ligands being BU72 (7-fold) and AFN42 (3-fold).

Effect of Ionic Composition on GTPγ^{35}S Binding and Ligand Affinity

Since we compared Na^+ to Tris, we needed to evaluate the role of ionic composition in functional receptor

activation with GTPγ^{35}S assays either in the presence of 100 mM KCl or 100 mM NaCl. The larger K^+ ion

does not fit well into the allosteric Na^+ binding pocket of MOR and so would not hold the receptor in an

inactive conformation. Subsequently, the standard opioid agonists like DAMGO exhibited 4-fold greater

potency in 100 mM KCl relative to 100 mM NaCl (11 nM (8.4 – 16 nM) vs. 42 nM (39 – 48 nM)), consistent

with previous evidence of Na^+ stabilizing inactive receptor (R) conformations that prevent G protein

signaling (**Figure 3**).[6,7,9] However, the two superagonists tested – AFN42 and carfentanil – demonstrate

increased potency (AFN42 in NaCl: 0.058 nM (0.043 – 0.081 nM) vs. in KCl: 0.25 nM (0.19 – 0.35 nM)) or

near equivalent potency (carfentanil in NaCl: 0.37 nM (0.28 – 0.49 nM) vs in KCl (0.19 nM (0.15 – 0.25

nM) in the presence of Na^+ relative to K^+. This suggests that Na^+ does not universally prevent agonist

activity but may instead promote a distinct active Na^+-occupied receptor state that may then still efficiently

couple to G proteins when bound to these highly potent ligands.

To evaluate the contribution of Na^+ ions on to ligand-receptor interactions, we performed radioligand

binding assays in the presence of 100 mM KCl and guanine nucleotide. Our conventional opioid agonists,

DAMGO and morphine, exhibited higher affinity in 100 mM KCl, or a smaller rightward shift in binding

affinity compared to 100 mM NaCl (**Table 2**). This is consistent with reduced binding affinity due to Na^+-

mediated stabilization of inactive (R) receptor states. In contrast, AFN42 displayed a greater rightward shift

in 100 mM KCl relative to 100 mM NaCl (1.68 nM vs. 0.07 nM), and this effect is still seen in 300 mM

NaCl conditions (**Table 2**). Supporting this narrative, Carfentanil displayed no statistically significant

change in affinity between 100 mM KCl, 100 mM NaCl and 300 mM NaCl conditions. These binding data

suggest that Na^+ is not inhibitory to binding – rather it may enhance binding affinity in the case of AFN42,

further supporting the existence of a Na^+-bound receptor state that may then efficiently couple to G proteins

when stabilized by these superagonists.

Radioligand Binding in CHO-µOR Cells with 50 mM Tris or 100 mM Na^+ and Guanine Nucleotide

Additional radioligand binding experiments were performed in CHO-µOR cell membranes in the absence

or presence of 100 mM Na^+ and guanine nucleotide to confirm the affinity shifts were cell type/environment

specific (**Table 3**). Of the three ligands tested – DAMGO, BU72 and carfentanil all displayed sub-

nanomolar binding affinities in 50 mM Tris. In the presence of 100 mM Na^+ and guanine nucleotide, only

DAMGO showed a large decrease in binding affinity, resulting in a shift of 810-fold. Both BU72 and

carfentanil exhibited smaller shifts of 3 and 35-fold, respectively. The differences in affinity shifts observed

between ligands and cell types could be due to differences between the human and rat µOR, as there are

subtle variations in amino acid sequences which have been shown can lead to differences in ligand binding

affinities.[24] Additionally, their cellular environments (hamster ovary versus rat glioma cell), level of

receptor expression and access to different accessory proteins on the membranes may also be of influence.

Nevertheless, the same pattern can be observed between cell types.

Correlating Intrinsic Efficacy with Affinity Shifts in NaCl Conditions

The level of agonism was then determined for each opioid agonist by calculating the $Log \left(\frac{Max}{EC50}\right)$ from

GTPγ^{35}S binding assays performed in 100 mM NaCl (as described in Kenakin, 2017).[3] The value for

AFN42 (10.26) is much greater than that of DAMGO (7.38). Since these parameters are on a logarithmic

scale, the difference in agonism between AFN42 and DAMGO (10.26 – 7.38 = 2.88) represents a 760-fold

difference. Since AFN42 and DAMGO are full agonists ($Max \geq 1$), as well as the other three highly potent

ligands, the difference in agonism is attributable to the difference in EC_{50}. The agonism value for each

compound was then plotted in comparison to the affinity shift due to Na$^+$ and guanine nucleotide (**Figure**

4). This correlation plot shows that agonism and affinity shift are negatively correlated to both the 100 mM

Na$^+$ conditions ($r^2 = 0.63$, p-value < 0.003) and the 300 mM Na$^+$ conditions ($r^2 = 0.45$, p-value $= 0.02$). This

correlation confirms that these experimental methods are both predictive measures of intrinsic efficacy.

Moreover, etorphine, BU72, carfentanil, etonitazene, and the opioid peptidomimetics demonstrate a

remarkable degree of agonism and a significantly decreased sensitivity to Na$^+$ (100 mM or 300 mM) and

free nucleotide (10 μM cold GTPγS). AFN42 was the most potent compound examined with roughly 1,000

to 3,000-fold higher potency than DAMGO, fentanyl, and morphine. AFN42 also is the least sensitive

opioid compound to Na$^+$ and guanine nucleotide, and AFN42 has the same affinity for the inactive R state

and active R* state receptor.

Discussion

In this study, we investigate the potency and binding affinity of a series of highly potent agonists at μOR

using GTPγ^{35}S assays, and radioligand competition binding assays in the presence of Na$^+$ and guanine

nucleotide. Correlation plots utilizing our functional data and affinity shifts revealed that these

superagonists are far less sensitive to Na$^+$ and guanine nucleotide than predicted, supporting an inadequacy

of the conformational selection model to explain their pharmacology. The conformational selection model

of receptor activation suggests that, by binding to high affinity states of the receptor, an agonist will shift

the equilibrium in favor of an active state (R*) conformations. Although this model predicts the action of

most types of agonists (e.g. inverse agonists, partial agonists, antagonists), the current data shows it falls

short in explaining the mechanism of action of extremely potent opioid agonists. It is generally accepted

that agonists have a greater affinity for R* compared to R and are sensitive to Na^+ and guanine nucleotide,

because these act as negative allosteric modulators and stabilize the low affinity R conformational state.[8]

In contrast to this idea, the very potent and high efficacy opioid agonists appear to be much less sensitive,

even insensitive to Na^+ or may even prefer the Na^+-bound state. Indeed, these compounds bind with a

similar high affinity to both active (R*) and inactive state (R) conformations. Binding assays performed in

Na^+-free conditions reinforce this difference in behavior seen between conventional agonists and

superagonists. DAMGO and morphine bind more tightly in 100 mM KCl conditions, consistent with the

absence of Na^+-mediated allosteric inhibition. However, AFN42 exhibits a paradoxical decrease in affinity

in KCl, with greater binding affinity in 100 mM NaCl and little to no further shift in 300 mM NaCl

conditions. This suggests that Na^+ occupancy of its allosteric site may even favor the binding conformation

of superagonists. The opposing shifts in KCl buffer between ligand classes reveal that the influence of

sodium ions on receptor conformation, and therefore ligand affinity, is not universal and ligand-dependent,

further supporting a model in which superagonists access a Na^+-bound receptor that is not accessible to

conventional agonists.

Additionally, our functional data suggests that Na^+ does not exhibit a uniform inhibitory effect across all

agonists. While the conventional opioid agonists such as fentanyl show reduced potency in Na^+ relative to

K^+ conditions, consistent with the established models of Na^+-mediated negative allosteric modulation,

superagonists like AFN42 display positive cooperativity with Na^+ in GTPγ^{35}S assays, maintaining or even

enhancing their potency in Na^+ conditions. These findings provide the functional support for the existence

of ligands that interact with a Na^+-bound state of μOR and drive it to couple to G proteins; where

superagonists not only tolerate Na^+ binding to its conserved allosteric site but may actively exploit it to

promote receptor activation. These findings challenge the traditional two-state model of GPCR activation

and suggest that the role of Na^+ as a negative allosteric modulator is ligand-dependent. Together with our

binding data, these results support a model in which superagonists stabilize a distinct active-like conformation that is permissive to Na^+ binding.

The β_1AR is the only currently known GPCR to not be modulated by Na^+ ions.[26] Na^+ interacts with a highly conserved residue, $Asp^{2.50}$, which is found in 98% of GPCRs, including the β_1AR and opioid receptor family.[27,28] Mutating this residue causes different functional outcomes depending on the GPCR. In the rat μOR, a D114(2.50) N mutation reduces binding affinities, potencies and efficacies of both DAMGO and morphine, as well as the binding affinity of the peptide antagonist, CTAP.[29] Similarly, for the delta-opioid receptor (δOR), the binding affinities of peptide and non-peptide agonists are reduced, while antagonists bind equally well to both mutated and wild-type receptors.[30] Additional mutagenesis studies of this residue have shown classical δOR antagonists transformed into potent beta-arrestin-biased agonists – suggesting an important role in mediating the R-R* equilibriums.[31] Similar experiments have been performed in a variety of Class A GPCRs and lead to a variety of different pharmacological effects, including reduced downstream signaling and receptor internalization, increases in agonist affinity with no change in antagonist affinity, no change in agonist affinity, and changes in antagonist affinity while abolishing G protein signaling.[32,33,34,35] However, mutating this residue at the β_1AR does not lead to significant differences in agonist binding affinity - even in the presence of Na^+ - suggesting the activity of this receptor is not modulated by Na^+.[26] The β_1AR is the only known Class A GPCR to not show modulation by Na^+ ions. Further functional, kinetic, and structural studies will need to be performed on additional GPCRs to determine whether superagonist insensitivity to Na^+ and guanine nucleotide is an opioid receptor phenomenon only (cite MOR/DOR insensitivity here) and whether specific ligand-receptor interactions mediate this phenomenon.[36]

The structure of the allosteric Na^+ site has been finely tuned to delicately balance the transition between R and R* through stabilizing GPCRs in their ligand-free state. Our results seem to indicate that superagonists of μOR with high intrinsic efficacy may negate this physiological role of Na^+ through additionally binding to a Na^+-bound receptor state, and facilitating it's transition into an active receptor state that is able to

efficiently couple to heterotrimeric G proteins. The implication of our findings is that a decrease in affinity

due to the presence of Na^+ is not a reliable predictor of efficacy when dealing with very potent compounds

and these observations are in contrast with previously accepted ideas.[6,12] It will be intriguing if this is the

reason why it is more difficult for naloxone to reverse respiratory depression caused by a highly potent

agonist like carfentanil.[37] Additionally, understanding the mechanisms and specific receptor-ligand

interactions that mediate Na^+ insensitivity may allow for the design of more potent GPCR agonists *in vivo*,

where there is an abundance of Na^+ and guanine nucleotide. It is important to note that this phenomenon

does not occur with existing endogenous opioid peptides, but exclusively with high affinity and intrinsic

efficacy ligands as observed so far.

The influence of sodium on GPCR activation should not be viewed as a uniform or as an obligatory

inhibitory mechanism. Sodium's allosteric effects appear to be probe-dependent, varying with the chemical

nature and coupling behavior of the bound ligand. Different agonists, or in this case superagonists, may

stabilize unique Na^+-bound intermediate or active-like receptor states that remain competent for G protein

engagement. In this context, superagonists may exploit sodium binding rather than oppose it, therefore

promoting receptor activation through Na^+-bound pathways that conventional agonists cannot access. Thus,

sodium's role in μOR activation may encompass multiple parallel trajectories, some requiring ion

displacement and others proceeding through stabilized Na^+-bound intermediates that facilitate G protein

coupling, as seen with superagonists like AFN42.

This work has broad implications for GPCR pharmacology – challenging foundational models of class A

GPCR activation and providing framework for designing next-generation GPCR therapeutics that retain

high affinity and potency under physiological ionic conditions. There is a significant negative correlation

between sodium sensitivity and intrinsic efficacy. The prevailing theory is that high efficacy ligands, such

as fentanyl and morphine, are more sensitive to Na^+ and guanine nucleotide compared to low efficacy

ligands. Here, instead of a positive correlation and what was previously thought to be true, there is a negative

correlation in which the ligands with the highest intrinsic efficacy are the least sensitive to Na^+ and guanine

nucleotide and thus show a smaller affinity shift.

C6 μ Cells	GTPγ^{35}S Binding			^{3}H-Diprenorphine Binding				
Compound	EC$_{50}$ (nM)	Max Stimulation (% of DAMGO, fractional)	Agonism $\text{Log} \frac{\text{Max}}{\text{EC50 (M)}}$	K$_i$ (nM) 50 mM Tris	K$_i$ (nM) + 100mM Na$^+$ +10μM GTPγS	Affinity Shift $\frac{\text{Ki 100mM Na +}}{\text{Ki 50mM Tris}}$	K$_i$ (nM) + 300mM Na$^+$ +10μM GTPγS	Affinity Shift $\frac{\text{Ki 300mM Na +}}{\text{Ki 50mM Tris}}$
DAMGO	42.2 (39.0-47.9)	1	7.38	4.71 (3.63-6.13)	618 (421-929)	131	5840 (4040-8470)	1240
Fentanyl	233 (178-308)	0.76 (0.72-0.80)	6.51	8.33 (5.74-12.3)	433 (239-780)	52.0	873 (488-1590)	105
Morphine	195 (148-260)	0.60 (0.57-0.63)	6.49	3.32 (2.47-4.42)	254 (169-377)	76.5	280 (155-485)	84.3
Diprenorphine	-	-	-	0.12 (0.094-0.16)	-	-	0.10 (0.077-0.13)	0.83
Etorphine	0.79 (0.42-1.39)	0.96 (0.89-1.04)	9.08	0.40 (0.24-0.66)	9.66 (6.02-15.4)	24.2	26.6 (19.7-35.6)	66.5
BU72	0.35 (0.24-0.51)	1.02 (0.96-1.08)	9.46	0.20 (0.14-0.29)	0.50 (0.33-0.77)	2.5	1.48 (0.96-2.33)	7.4
Carfentanil	0.37 (0.28-0.49)	1.06 (1.01-1.11)	9.46	0.17 (0.13-0.23)	2.05 (1.37-3.03)	12.1	2.36 (1.63-3.40)	13.9
Etonitazene	0.63 (0.38-1.03)	1.0 (0.92-1.08)	9.20	0.18 (0.14-0.22)	5.94 (4.19-8.46)	33	29.8 (19.3-45.1)	166
AAH8	0.64 (0.43-0.95)	0.9 (0.85-0.95)	9.15	0.058 (0.043-0.079)	0.50 (0.37-0.68)	8.6	1.70 (1.25-2.31)	29.3
AFN42	0.058 (0.043-0.081)	1.06 (1.01-1.11)	10.26	0.093 (0.072-0.12)	0.067 (0.054-0.083)	0.72	0.27 (0.19-0.36)	2.9
AMB46	0.86 (0.53-1.37)	0.96 (0.89-1.03)	9.04	0.077 (0.052-0.11)	0.91 (0.60-1.42)	11.8	3.39 (2.17-5.23)	44
AMB47	0.59 (0.37-0.92)	0.98 (0.91-1.05)	9.22	0.15 (0.097-0.28)	0.32 (0.27-0.38)	2.1	2.29 (1.65-3.15)	15.3

Table 1: Stimulation of GTPγ^{35}S Binding and Inhibition of ^{3}H-Diprenorphine Binding by Opioid Ligands in C6 μ Cells. Stimulation of GTPγ^{35}S binding at the μOR was performed using membranes from C6 cells with various concentrations of opioid agonists. Maximal stimulation of GTPγ^{35}S binding was determined by 10 μM of the reference agonist, DAMGO. Effective concentrations (EC$_{50}$) and the fractional maximal response (Max Stimulation) were determined by nonlinear regression analysis of concentration-response curves from GTPγ^{35}S binding assays. Agonism (Log [Max/EC50]) was calculated using EC$_{50}$ and max stimulation values for each opioid agonist. ^{3}H-Diprenorphine binding assays were performed using 0.2 – 0.5 nM of the radioligand in the following three buffers: **1.** 50 mM Tris buffer, **2.** 50 mM Tris, 100 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM GTPγS, or **3.** 50 mM Tris, 300 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM GTPγS. The affinity shift was determined for 100 mM Na$^+$ and 300 mM Na$^+$ conditions. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate or quadruplicate (n ≥ 3).

Figure 1: Chemical Structures of Opioid Agonists and Peptidomimetics. (A) DAMGO, (B) Fentanyl, (C) Morphine, (D) Etorphine, (E) BU72, (F) Carfentanil, (G) Etonitazene, (H) AAH8, (I) AFN42, (J) AMB46, and (K) AMB47

Figure 2: Stimulation of GTPγ³⁵S Binding and Inhibition of ³H-Diprenorphine Binding Concentration-Response Curves in C6 μ Cells. (A) GTPγ³⁵S binding assays were performed using membranes from C6 cells with various concentrations of opioid agonists. Maximal stimulation of GTPγ³⁵S binding was produced by 10 μM of the reference agonist, DAMGO. (B) Radioligand competition binding assays were performed in the following buffers: 50 mM Tris (circles), 50 mM Tris, 100 mM NaCl, 5 mM MgCl₂ 1 mM EDTA, 10 μM GTPγS (squares), or 50 mM Tris, 300 mM NaCl, 5 mM MgCl₂ 1 mM EDTA, 10 μM GTPγS (triangles). These assays used 20 μg membrane protein from C6 cells, 0.2 – 0.5 nM ³H-Diprenorphine, and various concentrations of the following opioid ligands: Diprenorphine, DAMGO, Fentanyl, Etorphine, BU72, Carfentanil, Etonitazene, AAH8, AFN42, AMB46, AMB47. All experiments were performed in duplicate or quadruplicate (n ≥ 3).

Figure 3: Stimulation of GTPγ^{35}S Binding in 100 mM NaCl and KCl buffer. Stimulation of GTPγ^{35}S binding at the µOR was performed using membranes from C6 cells with increasing concentrations of opioid agonists. Maximal stimulation of GTPγ^{35}S binding was determined by 10 µM of the reference agonist, DAMGO. Effective concentrations (EC$_{50}$) were determined by nonlinear regression analysis of concentration-response curves from GTPγ^{35}S binding assays. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate (n ≥ 3).

C6 μ Cells	^{3}H-Diprenorphine Binding		
Compound	K_i (nM) 50 mM Tris	K_i (nM) + 100mM K^+ +10μM GTPγS	Affinity Shift $\frac{\text{Ki 100mM K +}}{\text{Ki 50mM Tris}}$
DAMGO	4.71 (3.63-6.13)	111 (66.8-187)	23.6
Fentanyl	8.33 (5.74-12.3)	218 (140-344)	26.2
Morphine	3.32 (2.47-4.42)	150 (93.9-246)	45.2
Diprenorphine	0.12 (0.094-0.16)	1.96 (1.29-2.94)	16.3
Carfentanil	0.17 (0.13-0.23)	1.19 (0.81-1.76)	7
AFN42	0.093 (0.072-0.12)	1.68 (1.02-2.86)	18.1

Table 2. Inhibition of ^{3}H-Diprenorphine Binding in 100 mM KCl. Radioligand competition binding assays were performed in 50 mM Tris, 100 mM KCl, 5 mM MgCl$_2$ 1 mM EDTA, 10 μM GTPγS. These assays used 20 μg membrane protein from C6 cells, 0.2 – 0.5 nM ^{3}H-Diprenorphine, and various concentrations of opioid ligand. Experiments were performed in duplicate or quadruplicate (n ≥ 3) and are presented as mean with 95% confidence intervals.

CHO μ Cells	^{3}H-Diprenorphine Binding		
Compound	**K_i (nM)** 50 mM Tris	**K_i (nM)** + 100mM Na$^+$ +10μM GTPγS	**Affinity Shift** $\dfrac{\text{Ki 100mM Na}}{\text{Ki 50mM Tris}}$
DAMGO	0.65 (0.40-1.05)	529 (293-946)	814
BU72	0.27 (0.18-0.39)	0.73 (0.54-1.01)	2.7
Carfentanil	0.06 (0.035-0.10)	2.09 (1.56-2.79)	34.8

Table 3: Inhibition of ^{3}H-Diprenorphine Binding by Opioid Ligands in CHO μ Cells. ^{3}H-Diprenorphine binding assays were performed using 0.2 – 0.5 nM of the radioligand in the following two buffers: **1.** 50 mM Tris buffer, and **2.** 50 mM Tris, 100 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM cold GTPγS. The affinity shift was determined for 100 mM Na$^+$ conditions. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate (n ≥ 3).

Figure 4: Intrinsic Efficacy Plots for C6 μ. Agonism (Log [Max/EC$_{50}$]) was determined from stimulation of GTPγ^{35}S binding assays and Affinity Shift (Log [Ki, 100 mM Na$^+$ / K$_i$ 50 mM Tris] in **(A)** or Log [Ki, 300 mM Na$^+$ / K$_i$ 50 mM Tris] in **(B)** was determined from ^{3}H-Diprenorphine competition binding assays. Data was plotted for each of the following ligands: DAMGO, Fentanyl, Morphine, Etorphine, BU72, Carfentanil, Etonitazene, AFN42, AAH8, AMB46, and AMB47. r^2 = 0.63, p** = 0.003 **(A)**; r^2 = 0.45, p* = 0.02 **(B)**

References

1. Ehlert, F. J. (2018). Analysis of biased agonism. Progress in Molecular Biology and Translational Science, 63–104. https://doi.org/10.1016/bs.pmbts.2018.08.001

2. Traynor, J. R., & Nahorski, S. R. (1995). Modulation by mu-opioid agonists of guanosine-5'-O-(3-[35S]thio)triphosphate binding to membranes from human neuroblastoma SH-SY5Y cells. Molecular Pharmacology, 47(4), 848–854. https://doi.org/10.1016/s0026-895x(25)08634-1

3. Kenakin, T. (2017). A scale of agonism and allosteric modulation for assessment of selectivity, bias, and receptor mutation. *Carolina Digital Repository (University of North Carolina at Chapel Hill)*. https://doi.org/10.17615/hhsf-de46

4. Kenakin, T., & Christopoulos, A. (2011). Analytical pharmacology: the impact of numbers on pharmacology. Trends in Pharmacological Sciences, 32(4), 189–196. https://doi.org/10.1016/j.tips.2011.01.002

5. Herenbrink, C. K., Sykes, D. A., Donthamsetti, P., Canals, M., Coudrat, T., Shonberg, J., Scammells, P. J., Capuano, B., Sexton, P. M., Charlton, S. J., Javitch, J. A., Christopoulos, A., & Lane, J. R. (2016). The role of kinetic context in apparent biased agonism at GPCRs. Nature Communications, 7(1). https://doi.org/10.1038/ncomms10842

6. Pert, C. B., Pasternak, G., & Snyder, S. H. (1973). Opiate agonists and antagonists discriminated by receptor binding in brain. *Science*, *182*(4119), 1359–1361. https://doi.org/10.1126/science.182.4119.1359

7. Puttfarcken, P., Werling, L. L., Brown, S. R., Cote, T. E., & Cox, B. M. (1986). Sodium regulation of agonist binding at opioid receptors. I. Effects of sodium replacement on binding at mu- and delta-type receptors in 7315c and NG108-15 cells and cell membranes. *PubMed*, *30*(2), 81–89. https://doi.org/10.1016/s0026-895x(25)10340-4

8. Livingston, K. E. and J. R. Traynor (2014). Disruption of the Na+ ion binding site as a mechanism for positive allosteric modulation of the mu-opioid receptor. Proceedings of the National Academy of Science **111**(51): 18369-18374. https://doi.org/10.1073/pnas.1415013111

9. Katritch, V., G. Fenalti, E. E. Abola, B. L. Roth, V. Cherezov and R. C. Stevens (2014). Allosteric sodium in class A GPCR signaling. Trends in biochemical sciences **39**(5): 233-244. https://doi.org/10.1016/j.tibs.2014.03.002

10. Liu, W., E. Chun, A. A. Thompson, P. Chubukov, F. Xu, V. Katritch, G. W. Han, C. B. Roth, L. H. Heitman, A. P. Ijzerman, V. Cherezov and R. C. Stevens (2012). Structural basis for allosteric regulation of GPCRs by sodium ions. Science (New York, N.Y.) **337**(6091): 232-236. https://doi.org/10.1126/science.1219218

11. White, K. L., M. T. Eddy, Z.-G. Gao, G. W. Han, T. Lian, A. Deary, N. Patel, K. A. Jacobson, V. Katritch and R. C. Stevens (2018). Structural Connection between Activation Microswitch and Allosteric Sodium Site in GPCR Signaling. Structure **26**(2): 259-269.e255. https://doi.org/10.1016/j.str.2017.12.013

12. Childers, S. R., & Snyder, S. H. (1980). Differential regulation by guanine nucleotides of opiate agonist and antagonist receptor interactions. Journal of Neurochemistry, 34(3), 583–593. https://doi.org/10.1111/j.1471-4159.1980.tb11184.x

13. Lee, K. O., Akil, H., Woods, J. H., & Traynor, J. R. (1999). Differential binding properties of oripavines at cloned μ- and δ-opioid receptors. European Journal of Pharmacology, 378(3), 323–330. https://doi.org/10.1016/s0014-2999(99)00460-4

14. Bender, A. M., N. W. Griggs, J. P. Anand, J. R. Traynor, E. M. Jutkiewicz and H. I. Mosberg (2015). Asymmetric Synthesis and in Vitro and in Vivo Activity of Tetrahydroquinolines Featuring a Diverse Set of Polar Substitutions at the 6 Position as Mixed-Efficacy mu Opioid Receptor/delta Opioid Receptor Ligands. ACS Chem Neurosci **6**(8): 1428-1435. https://doi.org/10.1021/acschemneuro.5b00100

15. Harland, A. A., L. Yeomans, N. W. Griggs, J. P. Anand, I. D. Pogozheva, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2015). Further Optimization and Evaluation of Bioavailable, Mixed-Efficacy mu-Opioid Receptor (MOR) Agonists/delta-Opioid Receptor (DOR) Antagonists: Balancing MOR and DOR Affinities. J Med Chem **58**(22): 8952-8969. https://doi.org/10.1021/acs.jmedchem.5b01270

16. Harland, A. A., A. M. Bender, N. W. Griggs, C. Gao, J. P. Anand, I. D. Pogozheva, J. R. Traynor, E. M. Jutkiewicz and H. I. Mosberg (2016). Effects of N-Substitutions on the Tetrahydroquinoline (THQ) Core of

Mixed-Efficacy mu-Opioid Receptor (MOR)/delta-Opioid Receptor (DOR) Ligands. J Med Chem **59**(10): 4985-4998. https://doi.org/10.1021/acs.jmedchem.6b00308

17. Nastase, A. F., N. W. Griggs, J. P. Anand, T. J. Fernandez, A. A. Harland, T. J. Trask, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2018). Synthesis and Pharmacological Evaluation of Novel C-8 Substituted Tetrahydroquinolines as Balanced-Affinity Mu/Delta Opioid Ligands for the Treatment of Pain. ACS Chem Neurosci **9**(7): 1840-1848. https://doi.org/10.1021/acschemneuro.8b00139

18. Montgomery, D., J. P. Anand, M. A. Baber, J. J. Twarozynski, J. G. Hartman, L. J. Delong, J. R. Traynor and H. I. Mosberg (2019). Structure-Activity Relationships of 7-Substituted Dimethyltyrosine-Tetrahydroisoquinoline Opioid Peptidomimetics. Molecules **24**(23). https://doi.org/10.3390/molecules24234302

19. Montgomery, D., J. P. Anand, N. W. Griggs, T. J. Fernandez, J. G. Hartman, A. A. Sanchez-Santiago, I. D. Pogozheva, J. R. Traynor and H. I. Mosberg (2019). Novel Dimethyltyrosine-Tetrahydroisoquinoline Peptidomimetics with Aromatic Tetrahydroisoquinoline Substitutions Show in Vitro Kappa and Mu Opioid Receptor Agonism. ACS Chem Neurosci **10**(8): 3682-3689. https://doi.org/10.1021/acschemneuro.9b00250

20. Nastase, A. F., J. P. Anand, A. M. Bender, D. Montgomery, N. W. Griggs, T. J. Fernandez, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2019). Dual Pharmacophores Explored via Structure-Activity Relationship (SAR) Matrix: Insights into Potent, Bifunctional Opioid Ligand Design. J Med Chem **62**(8): 4193-4203. https://doi.org/10.1021/acs.jmedchem.9b00378

21. Clark, M. J., Harrison, C., Zhong, H., Neubig, R. R., & Traynor, J. R. (2003). Endogenous RGS Protein Action Modulates μ-Opioid Signaling through Gαo. Journal of Biological Chemistry, 278(11), 9418–9425. https://doi.org/10.1074/jbc.m208885200

22. Anand, J. P., Kochan, K. E., Nastase, A. F., Montgomery, D., Griggs, N. W., Traynor, J. R., Mosberg, H. I., & Jutkiewicz, E. M. (2018). In vivo effects of μ-opioid receptor agonist/δ-opioid receptor antagonist peptidomimetics following acute and repeated administration. *British Journal of Pharmacology*, *175*(11), 2013–2027. https://doi.org/10.1111/bph.14148

23. Pasternak, G. W., & Pan, Y. (2013). MU Opioids and their Receptors: Evolution of a Concept. *Pharmacological Reviews*, *65*(4), 1257–1317. https://doi.org/10.1124/pr.112.007138

24. Rothman, R. B., Xu, H., Wang, J. B., Partilla, J. S., Kayakiri, H., Rice, K. C., & Uhl, G. R. (1994). The ligand-selectivity of cloned human and rat opiate mu receptors studied with [125I]IOXY-AGO. *Regulatory Peptides*, *54*(1), 249–250. https://doi.org/10.1002/syn.890210109

25. Malcolm, N. J., Palkovic, B., Sprague, D. J., Calkins, M. M., Lanham, J. K., Halberstadt, A. L., Stucke, A. G., & McCorvy, J. D. (2023). Mu-opioid receptor selective superagonists produce prolonged respiratory depression. *iScience*, *26*(7), 107121. https://doi.org/10.1016/j.isci.2023.107121

26. Miller-Gallacher, J. L., Nehmé, R., Warne, T., Edwards, P. C., Schertler, G. F. X., Leslie, A. G. W., & Tate, C. G. (2014). The 2.1 Å Resolution Structure of Cyanopindolol-Bound β1-Adrenoceptor Identifies an Intramembrane Na+ Ion that Stabilises the Ligand-Free Receptor. *PLoS ONE*, *9*(3), e92727. https://doi.org/10.1371/journal.pone.0092727

27. Parker, M. S., Wong, Y. Y., & Parker, S. L. (2008). An ion-responsive motif in the second transmembrane segment of rhodopsin-like receptors. *Amino Acids*, *35*(1), 1–15. https://doi.org/10.1007/s00726-008-0637-6

28. Shang, Y., LeRouzic, V., Schneider, S., Bisignano, P., Pasternak, G. W., & Filizola, M. (2014). Mechanistic Insights into the Allosteric Modulation of Opioid Receptors by Sodium Ions. *Biochemistry*, *53*(31), 5140–5149. https://doi.org/10.1021/bi5006915

29. Xu, W., Ozdener, F., Li, J., Chen, C., De Riel, J., Weinstein, H., & Liu-Chen, L. (1999). Functional role of the spatial proximity of Asp114(2.50) in TMH 2 and Asn332(7.49) in TMH 7 of the μ opioid receptor. *FEBS Letters*, *447*(2–3), 318–324. https://doi.org/10.1016/s0014-5793(99)00316-6

30. Kong, H., Raynor, K., Yasuda, K., Moe, S. T., Portoghese, P. S., Bell, G. I., & Reisine, T. (1993). A single residue, aspartic acid 95, in the delta opioid receptor specifies selective high affinity agonist binding. *Journal of Biological Chemistry*, *268*(31), 23055–23058. https://doi.org/10.1016/s0021-9258(19)49424-6

31. Fenalti, G., Giguere, P. M., Katritch, V., Huang, X., Thompson, A. A., Cherezov, V., Roth, B. L., & Stevens, R. C. (2014). Molecular control of δ-opioid receptor signalling. *Nature*, *506*(7487), 191–196. https://doi.org/10.1038/nature12944

32. Proulx, C. D., Holleran, B. J., Boucard, A. A., Escher, E., Guillemette, G., & Leduc, R. (2008). Mutational analysis of the conserved ASP2.50and ERY motif reveals signaling bias of the urotensin II receptor. *Molecular Pharmacology*, *74*(3), 552–561. https://doi.org/10.1124/mol.108.045054

33. H Barbhaiya, McClain R, A Ijzerman, Rivkees SA. Site-directed mutagenesis of the human A1 adenosine receptor: influences of acidic and hydroxy residues in the first four transmembrane domains on ligand binding (1996). *Molecular Pharmacology*. 50(6):1635-1642. doi:https://doi.org/10.1016/s0026-895x(25)09624-5

34. Gao, Z., Kim, S., Gross, A. S., Chen, A., Blaustein, J. B., & Jacobson, K. A. (2003). Identification of essential residues involved in the allosteric modulation of the human A3Adenosine receptor. *Molecular Pharmacology*, *63*(5), 1021–1031. https://doi.org/10.1124/mol.63.5.1021

35. Bihoreau, C., Monnot, C., Davies, E., Teutsch, B., Bernstein, K. E., Corvol, P., & Clauser, E. (1993). Mutation of Asp74 of the rat angiotensin II receptor confers changes in antagonist affinities and abolishes G-protein coupling. *Proceedings of the National Academy of Sciences*, *90*(11), 5133–5137. https://doi.org/10.1073/pnas.90.11.5133

36. Childers, S. R., Fleming, L. M., Selley, D. E., McNutt, R. W., & Chang, K. J. (1993). BW373U86: a nonpeptidic delta-opioid agonist with novel receptor-G protein-mediated actions in rat brain membranes and neuroblastoma cells. *Molecular Pharmacology*, *44*(4), 827–834. https://doi.org/10.1016/s0026-895x(25)13269-0

37. Flynn, S. M., & France, C. P. (2025). Comparison of ventilatory depression by carfentanil, fentanyl, and heroin, and reversal by opioid receptor antagonists in rats. *Neuropharmacology*, 110601. https://doi.org/10.1016/j.neuropharm.2025.110601