# μ-Opioid Receptor Superagonists Engage Sodium-Bound Receptor States

**Author Names:** *Alexander Powell[1], Nicholas W. Griggs[1], Jorge Iniguez-Lluhi[1], Henry Mosberg[2], John R. Traynor[1,2]*

**Author Address:** [1]Department of Pharmacology, Edward F. Domino Research Center, Medical School, University of Michigan, Ann Arbor, MI 48109, United States

[2]Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, Ann Arbor, MI 48109, United States

**Corresponding Author:** John R. Traynor (jtraynor@umich.edu | 734-647-7479)

**Abstract**

The simple two-state conformational selection model of G-protein coupled receptor (GPCR) activation suggests that, by binding to a high affinity site, an agonist will shift receptor equilibrium in favor of active state (R*) conformations that recruit heterotrimeric G proteins over inactive state (R) conformations. Agonist binding to the μ-opioid receptor is highly sensitive to $Na^+$ ions which stabilize an inactive receptor state. Higher efficacy opioid agonists, such as DAMGO and fentanyl, are sensitive to $Na^+$ compared to lower efficacy ligands at the μ-opioid receptor. However, the binding of the highly potent oripavine agonists etorphine and dihydroetorphine are less sensitive to $Na^+$ than expected such that the prevailing models fail to explain their pharmacology. To explain this discrepancy, experiments were performed to evaluate the binding properties and G protein activation of the highly potent agonists carfentanil, BU72, etorphine, etonitazene and similarly potent opioid peptidomimetics in comparison to the standard agonists DAMGO, fentanyl, and morphine in the presence or absence of $Na^+$ or $K^+$ ions. Several of the superagonists retained high affinity and potency in both ionic conditions, whereas DAMGO, fentanyl and morphine displayed enhanced binding and signaling in $K^+$, compared to $Na^+$ ions. These functional parameters were used to determine an intrinsic efficacy value, determined as $Log(\frac{Max}{EC50})$. Comparison of affinity shifts with intrinsic

efficacy afforded a negative correlation in which superagonists with the highest intrinsic efficacy are least

sensitive to Na$^+$. These data suggest that select μ-opioid receptor superagonists have high affinity for the

Na$^+$ bound receptor states (R) and shift these species into active receptor conformations (R*) that efficiently

couple to G proteins.

**Significance Statement**

The simple theory of conformational selection suggests the binding affinities of high efficacy μ-opioid

receptor ligands, such as fentanyl and DAMGO, are more sensitive to Na$^+$ and guanine nucleotide which

stabilize inactive receptor states than lower efficacy agonists and antagonists. Here, we show that ligands

with high intrinsic efficacy (superagonists) are much less sensitive to Na$^+$ and guanine nucleotide. This

work demonstrates that  highly potent ligands can engage a low affinity Na$^+$-bound receptor state that may

then convert to a receptor species that efficiently couples to G protein – i.e. a conformational induction.

**Key words:** Opioid, GPCR, Mu-opioid Receptor, Intrinsic Efficacy, Affinity, Conformation Selection

Theory

**Abbreviations:** CHO, Chinese hamster ovary; DAMGO, [D-Ala$^2$, NMe-Phe$^4$, Gly-ol$^5$]-enkephalin; GDP,

Guanosine 5'-diphosphate; GPCR, G protein-coupled receptor; GTPγ$^{35}$S, (Guanosine-5'-O-(γ-thio)

triphosphate-[$^{35}$S]); μOR, mu-opioid receptor.

## Introduction

Mu-opioid receptors (μOR) are G protein-coupled receptor (GPCR) proteins that activate heterotrimeric G proteins of the $G\alpha_{i/o}$ family. The mu-opioid receptors, like other GPCRs, exist in an equilibrium of various conformational states that can be grouped into ensembles of inactive (R) and active states (R*); the R* state interacting with the G protein α-subunit. Intrinsic efficacy describes the ability of a ligand to shift the equilibrium from inactive-state conformations to active-state conformations. This parameter distinguishes ligands that prefer the "low affinity" R state (inverse agonists) from those that prefer a "high affinity" R* state (partial agonists and full agonists) and from those that display no preference (neutral antagonists). Intrinsic efficacy, in conjunction with binding affinity and pharmacokinetic properties, are the driving factors that determine the pharmacological activity of a drug for a particular receptor system.[1]

Stimulation of $GTP\gamma^{35}S$ binding to G proteins is a sensitive indicator of the active R* state and may be used as a predictive measure of the intrinsic efficacy of a ligand.[2] $GTP\gamma^{35}S$ binding assays allow for the determination of potency ($EC_{50}$) and relative efficacy as maximal stimulation by a ligand compared to a standard agonist. Determination of a $Log\left(\frac{Max}{EC50}\right)$ from these data is a convenient way to quantify the degree of ligand agonism.[3] One advantage of using the $GTP\gamma^{35}S$ binding assay to assess ligand efficacy is that partial agonists are aptly discriminated from full agonists, since G protein activation is directly downstream of ligand-receptor binding and is less susceptible to signal amplification. On the other hand, and like other functional assays that rely on measurement of signal transduction events downstream of the receptor, results are dependent on experimental conditions.[4,5]

Agonist-mediated activation of the μOR is negatively modulated by the binding of $Na^+$ ions.[6,7,8] These ions bind deep in the receptor below the orthosteric site and form ionic interactions with transmembrane domains 2, 3, and 7, holding the receptor in an inactive set of conformations.[9] Thus, agonists and $Na^+$ ions work in opposition; agonists shift the equilibrium toward active R* ensembles whereas $Na^+$ ions

shift the equilibrium toward inactive R ensembles.[9,10,11] This negative cooperativity has been long known

at μOR, in which the addition of $Na^+$ ions with guanine nucleotide to uncouple G proteins, decreases the

binding affinity ($K_i$) of opioid agonists.[6,12] Higher efficacy opioid agonists show a greater decrease in

binding affinity upon the addition of $Na^+$ and guanine nucleotide compared to lower efficacy agonists and

antagonists display no shift in affinity.[6,13] There appears to be a positive association between the decrease

in the $K_i$ of an agonist by $Na^+$ (i.e. sodium sensitivity) and the intrinsic efficacy of an opioid. However, two

very potent opioid agonists, etorphine and dihydroetorphine, demonstrate significantly reduced sodium

sensitivity which questions the validity of the observed positive association between sodium sensitivity and

intrinsic efficacy at μOR.[13]

The effects of sodium on μOR activation cannot be treated as uniform or universal. Sodium may stabilize

multiple inactive conformations, but its allosteric influence is probe-dependent and varies with the

specific agonist engaged, based largely on agonist efficacy. Different ligands couple to the receptor in

distinct ways that either disrupt or accommodate sodium binding.[13,14] For superagonists, it is possible that

receptor engagement does not require displacement of the $Na^+$ ion; instead, these ligands may interact

with $Na^+$-bound states leading to displacement of $Na^+$ and receptor activation, i.e. conformational

induction or induced fit. Alternatively, superagonists bind and stabilize $Na^+$-bound receptor ensembles

that are still competent to recruit G protein. Thus, receptor activation may proceed through several parallel

trajectories, with some involving sodium displacement and others proceeding through $Na^+$-bound "active-

like" states that still facilitate G protein coupling. These observations suggest that the role of $Na^+$ in μOR

activation should not be considered strictly inhibitory but instead may include $Na^+$-bound states that

remain competent to interact with high-efficacy agonists.

Here, we report the determination of intrinsic efficacy for a series of potent opioids including carfentanil,

BU72, etorphine and etonitazene, together with a series of highly potent opioid peptidomimetics, and

compare them with standard opioids morphine, DAMGO and fentanyl (**Figure 1**). By evaluating the affinity

of these ligands in the absence and presence of $Na^+$ and guanine nucleotide and calculating the difference

in affinity between the two conditions (i.e. "affinity shift"), we show that for highly potent opioid agonists,

reduced $Na^+$ sensitivity correlates significantly with higher intrinsic efficacy. The model of conformational

selection would predict the compounds to be neutral antagonists or low efficacy agonists, as they

demonstrate small affinity shifts and recognize different affinity states (inactive and active) in a near equal

manner. Instead, we suggest a conformational induction model in which highly potent agonists bind to a

low-affinity state of μOR in the presence of $Na^+$ ions and shift the receptor into a high-affinity state capable

of recruiting G protein.

**Methods**

***Materials and Reagents***

We have previously reported the synthesis and pharmacological evaluation of novel opioid

peptidomimetic compounds.[15,16,17,18,19,20,21] AAH8 (compound 14a) was synthesized by Aubrie A.

Harland[16], AFN42 (compound 4d) was synthesized by Anthony F. Nastase[21], AMB46 (compound 10n)

and AMB47(Compound 10m) were synthesized by Aaron M. Bender.[15] All chemicals, unless otherwise

specified, were purchased from Sigma Aldrich. Radiolabeled $[^3H]$-diprenorphine (DPN) and radiolabeled

guanosine-5′-O-(3-$[^{35}S]$ thio)-triphosphate (GTPγ$^{35}$S) were purchased from PerkinElmer Life Sciences.

All tissue culture media, penicillin-streptomycin, geneticin (G148), trypsin, and fetal bovine serum were

purchased from Gibco LifeSciences.

***Cell Lines***

The tissue culture and maintenance of C6 rat glioma cells stably transfected with rat μOR was performed

as previously described.[21] C6 cells were grown to confluence at 37 °C in 5 % $CO_2$ in Dulbecco's modified

Eagle media (DMEM) containing 10 % fetal bovine serum (FBS), 5 % penicillin/streptomycin (P/S), and

400 ug/mL G418. CHO cells stably expressing wild-type human μOR were grown to confluence at 37 °C

in 5 % $CO_2$ in Dulbecco's modified Eagle media (DMEM) containing 10 % fetal bovine serum (FBS),

1 % penicillin/streptomycin (P/S), and 400 ug/mL G418.

### Cell Membrane Preparation

Cell membrane homogenates from C6 and CHO cells were prepared by washing confluent cells three times with phosphate buffered saline (pH 7.4), after which they were then detached using harvesting buffer (20 mM HEPES, 150 mM NaCl, 0.68 mM EDTA, pH7.4) and pelleted by centrifugation at 200 x g for 3 min at room temperature. The pellet was resuspended in ice-cold 50 mM Tris buffer (pH 7.4) and homogenized using a Tissue Tearor (Biospec Products, Inc.). The cell lysate was centrifuged at 20,000 x g at 4 °C for 20 minutes, after which the pellet was then resuspended on ice cold 50 mM Tris (pH7.4), homogenized, and centrifuged once more at 20000 x g at 4 C for 20 minutes. The final pellet was resuspended in 50 mM Tris (pH 7.4) using a glass dounce homogenizer and then dispensed into aliquots and stored at -80 °C. The final protein concentration was determined by performing a BCA protein assay (Thermo Scientific Pierce) using bovine serum albumin as the standard.

### Competition Radioligand Binding Assays

$^3$H-diprenorphine competition binding assays were performed in 500μL reactions in 96-well plates using the cell membrane homogenates described above. Briefly, cell membranes [5-20 μg] were incubated in 50 mM Tris buffer containing 0.2 nM $^3$H-Diprenorphine, and various concentrations of opioid compound for 90 minutes in a shaking water bath at room temperature to reach equilibrium. Nonspecific binding was determined using 10 μM of the opioid antagonist, naloxone. The reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue harvester (Brandel, Gaithersburg, MD) and washed with 50 mM Tris buffer. Filters were dried and following the addition of EcoLume scintillation cocktail, bound radioactivity was quantified in a MicroBeta 2450 Liquid Scintillation and Luminescence Counter (PerkinElmer).

Radioligand binding assays in $Na^+$ or $K^+$ containing buffers were performed in 200 μL reactions in 96-well plates using the cell membrane homogenates described above, Briefly, cell membranes [5-20 μg] were incubated in $Na^+$ or $K^+$ containing buffers (50 mM Tris, 100 mM NaCl or 100 mM KCl, 5 mM MgCl2, 1 mM EDTA, pH 7.4) or (50 mM Tris, 300 mM NaCl, 5 mM MgCl2, 1 mM EDTA, pH 7.4)

containing 10 μM cold GTPγS, and various concentrations of opioid compound for 90 minutes in a

shaking water bath at room temperature. Nonspecific binding was determined using 10 μM naloxone. The

reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue harvester

and washed with $Na^+$ containing buffer, and the filters were processed as described above.

***Stimulation of GTPγ$^{35}$S Binding Assays***

Stimulation of GTPγ$^{35}$S binding experiments were performed in 200 μL reactions in 96-well plates using

cell membrane homogenates prepared as described above. Briefly, cell membranes [10-20 μg] were

incubated in either $Na^+$ or $K^+$ containing buffer [50 mM Tris, 100 mM NaCl or 100mM KCl, 5 mM

MgCl2, 1 mM EDTA, pH 7.4] containing 0.1 nM guanosine-5′-O-(3-[$^{35}$S]thio)triphosphate (GTPγ$^{35}$S)

and 30 μM guanosine 5'-diphosphate (GDP) and various concentrations of opioid compound or

peptidomimetic for 1 hour in a shaking water bath at 25 °C. Basal stimulation of GTPγ$^{35}$S binding was

determined by incubation in the absence of any ligand and maximal stimulation of was defined by using

10 μM of the standard mu opioid receptor agonist, DAMGO ([D-Ala$^2$,N-MePhe$^4$,Gly-ol]-enkephalin).

The reaction was terminated by vacuum filtration onto glass microfiber GF/C filters using a tissue

harvester and washed with $Na^+$ containing buffer, and filters were processed as described above.

## Results

### GTPγ$^{35}$S Binding

As evaluated by concentration response curves of GTPγ$^{35}$S binding in C6 μOR membranes, etorphine, BU72, carfentanil and etonitazene were seen to be very potent agonists with EC$_{50}$ values in the low nanomolar and sub-nanomolar range (**Table 1 and Figure 2**). In addition, we have previously described a series of highly potent peptidomimetics that bind to μOR with exceptional affinity.[15,16,21] These compounds (H-K in **Figure 1**) are at least 70 times more potent than the standard opioid peptide DAMGO, fentanyl, or morphine with potencies that are similar to carfentanil and etonitazene. All compounds demonstrated high relative efficacy, similar to DAMGO, as evaluated by maximal stimulation of GTPγ$^{35}$S binding. The fractional maximal response (i.e. "Max") compared to DAMGO (Max = 1) was equivalent for all ligands.

### Agonist Binding Affinities

It has been known for decades that physiologically relevant Na$^+$ ion concentrations decrease the binding affinity of agonists to μOR.[22] Radioligand competition binding assays were performed using membranes from C6 cells expressing the rat μOR in the presence or absence of Na$^+$ ions and guanine nucleotide (GTPγS) to induce R and R* receptor states, respectively. Difference in binding affinities between the two conditions were given as affinity shifts (**Table 1 and Figure 2**). In 50 mM Tris, the superagonists carfentanil, BU72, etorphine and etonitazene bound with approximately 10-fold higher affinity than the classical agonists DAMGO, morphine and fentanyl. The peptidomimetics exhibited similar binding affinities, with AAH8 displaying the highest affinity.

Binding affinities of DAMGO, fentanyl, and morphine to μOR decreased in the presence of 100 mM Na$^+$ and GTPγS, producing affinity shifts of 131, 52 and 77-fold, respectively (**Table 1**). Of the superagonists, etorphine and etonitazene displayed the greatest affinity shifts of 24 and 33-fold, respectively. Carfentanil,

AAH8 and AMB47 all displayed 10-fold decreases in affinity, whereas BU72, AFN42 and AMB47

displayed much smaller shifts or no shift.

Since the superagonists displayed a less than expected affinity shift in the presence of 100 mM $Na^+$ and

GTPγS, we utilized a supraphysiological concentration of $Na^+$ to determine whether the affinity shift for

each superagonist was saturated (**Table 1**). In the presence of 300 mM $Na^+$ ions and GTPγS, an affinity

shift of 1,240-fold was observed for DAMGO, whereas fentanyl and morphine displayed smaller shifts of

105 and 84-fold, respectively. Of the highly potent agonists, only etonitazene produced an affinity shift

greater than 100-fold.  Etorphine, AAH8 and AMB46 had comparable affinity shifts ranging from 29 to 67-

fold. Carfentanil and AMB47 displayed smaller shifts of 14 and 15-fold, respectively. Two compounds

exhibited shifts of less than 10-fold, these ligands being BU72 (7-fold) and AFN42 (3-fold).

**Comparison of the Effect of $Na^+$ and $K^+$ Ions on Agonist-Mediated GTPγ$^{35}$S Binding and Ligand**

**Affinity**

Since we compared agonist binding in $Na^+$ and GTPγS to binding in 50 mM Tris-HCl buffer alone, we

needed to evaluate the role of ionic strength on the binding and functional measures. To do this, we replaced

NaCl with the same concentration of KCl and examined two superagonists, carfentanil and AFN42. The

conserved class A GPCR allosteric sodium site is defined by a tight coordination network involving $Asp^{2.50}$

and multiple other polar residues. This geometry selectively accommodates $Na^+$ and is not observed with

larger monovalent cations, like $K^+$, in solved crystal structures.[23] Therefore, the larger $K^+$ ion would not be

expected to efficiently hold the receptor in an inactive conformation. Consequently, in the GTPγ$^{35}$S assay

the standard opioid agonist DAMGO exhibited 4-fold greater potency in 100 mM KCl relative to 100 mM

NaCl (11 nM (8.4 – 16 nM) vs. 42 nM (39 – 48 nM) respectively), consistent with previous evidence that

$Na^+$ stabilizes an inactive receptor (R) conformations and so inhibits G protein coupling (**Figure 3**).[6,7,9] In

support, diprenorphine exhibited partial agonist activity in the 100 mM KCl condition. However, the

superagonist tested AFN42 demonstrated increased potency in the presence of $Na^+$ compared to $K^+$ (AFN42

in NaCl: 0.058 nM (0.043 – 0.081 nM) vs. in KCl: 0.25 nM (0.19 – 0.35 nM) and carfentanil near equivalent

potency (carfentanil in NaCl: 0.37 nM (0.28 – 0.49 nM) vs in KCl (0.19 nM (0.15 – 0.25 nM). This suggests

that $Na^+$ does not universally prevent agonist activity unlike previously thought to be true.

To evaluate the contribution of $Na^+$ ions and GTPγS on ligand-receptor interactions, we compared the above

results with radioligand binding assays in the presence of 100 mM KCl and GTPγS. These latter conditions

lead to a smaller rightward shift in affinity, consistent with the inability of $K^+$ ions to replace $Na^+$ ions and

stabilize a low affinity state (**Table 2**). In contrast, AFN42 displayed a greater rightward shift in 100 mM

KCl (18.1-fold) than it did in the presence of 100 mM NaCl (0.7-fold) or even 300 mM NaCl (2.9-fold)

(**Table 2**). Similarly, carfentanil displayed no significant change in affinity when examined in 100 mM KCl

($K_i$: 1.2 nM), 100 mM NaCl ($K_i$: 2.1 nM), or 300 mM NaCl ($K_i$: 2.4 nM).

These data suggest that $Na^+$ is not always inhibitory to agonist binding and indicates superagonists are

capable of binding to a $Na^+$-bound receptor state. It may then convert this receptor species into a $Na^+$-free

state that can then efficiently couple to G proteins.

**Agonist Binding in CHO Cells Expressing Human μOR**

Since the above data was obtained in C6 glioma cells expressing the rat μOR, we performed experiments

in CHO cells expressing human μOR (CHO-μOR). Radioligand binding experiments were performed in

CHO-μOR cell membranes in the absence or presence of 100 mM $Na^+$ ions and GTPγS to confirm that the

affinity shifts observed were not specific to the rat μOR or dependent on the cell type/receptor environment

(**Table 3**). Three ligands were examined – DAMGO, BU72 and carfentanil. In the presence of 100 mM $Na^+$

and GTPγS, DAMGO showed a substantial decrease in binding affinity, resulting in a shift of 810-fold,

larger than the 131-fold shift observed in the C6 μOR cells. Both BU72 and carfentanil exhibited smaller

shifts of 3 and 35-fold, respectively, although these shifts were greater than observed in the C6 cells. The

differences in affinity shifts observed between ligands and cell types could be due to differences between

the human and rat μOR, as there are subtle variations in amino acid sequences which have been shown can

lead to differences in ligand binding affinities.[24] Additionally, their cellular environments (hamster ovary

versus rat glioma cell) and access to different accessory proteins on the membranes may also be of influence.

Nevertheless, the same pattern is observed across the two species of μOR and cell types.

**Correlating Intrinsic Efficacy with Affinity Shifts in NaCl Conditions**

The level of agonism was determined for each opioid agonist by calculating the $Log\ (\frac{Max}{EC50})$ from GTPγ$^{35}$S

binding assays performed in 100 mM NaCl (as described in Kenakin, 2017).[3] The value for AFN42 (10.26)

is much greater than that of DAMGO (7.38). Since these parameters are on a logarithmic scale, the

difference in agonism between AFN42 and DAMGO (10.26 – 7.38 = 2.88) is a 760-fold difference. Since

the superagonists tested give the same maximal response the difference in agonism is attributable to the

difference in $EC_{50}$. The agonism value for each compound was then plotted in comparison to the affinity

shift seen in the presence of Na$^+$ ions and GTPγS (**Figure 4**). This plot shows that for full agonists in the

GTPγ$^{35}$S assay, agonism and affinity shift are negatively correlated in both the 100 mM Na$^+$/GTPγS

conditions ($r^2 = 0.63$, p-value < 0.003) and the 300 mM Na$^+$/GTPγS conditions ($r^2 = 0.45$, p-value = 0.02).

Thus, both the level of agonism and Na$^+$-dependent affinity shifts serve as complementary and predictive

readouts of superagonism at the receptor.

**Discussion**

In this study, we investigate the potency and binding affinity of a series of highly potent agonists

(superagonists) at μOR using GTPγ$^{35}$S assays, and radioligand competition binding assays in the absence

or presence of Na$^+$ ions and guanine nucleotide (GTPγS). Na$^+$ ions act as negative allosteric modulators of

the receptor, stabilizing the inactive R resting state. Similarly, guanine nucleotides prevent the G protein

from stabilizing the active R* state by binding to the Gα subunit of the G protein heterotrimer. Agonists

preferentially bind to the active R* receptor state, exhibiting higher affinity for R* than for R. Under the

conformational selection model, this binding shifts the receptor population or equilibrium toward the active

R* state. However, our data comparing differences in agonist affinities in two ionic conditions with the

GTP$\gamma^{35}$S functional data revealed that superagonists may be far less sensitive to Na$^+$ ions and guanine

nucleotide than predicted. Although the conformational selection model predicts the action of most types

of ligands (such as endogenous agonists and agonists with similar kinetic properties, inverse agonists,

partial agonists and antagonists), our data suggests it falls short in explaining the action of extremely potent

opioid agonists, which appear able to function through a conformation induced or induced-fit model,

binding to an inactive Na$^+$-bound R states and converting these to active R* states.

Binding assays performed in conditions in which Na$^+$ ions were replaced with K$^+$ ions to maintain

osmolarity highlighted the difference in behavior seen between superagonists and less potent agonists.

DAMGO and morphine bind more tightly to μOR in the presence of 100 mM KCl and GTPγS, consistent

with the absence of Na$^+$-mediated allosteric inhibition. However, AFN42 exhibits a paradoxical increase in

binding affinity in the presence of 100 mM or even 300 mM Na$^+$ ions and GTPγS compared to its affinity

in 100 mM K$^+$ ions with GTPγS. This suggests that Na$^+$ occupancy of its allosteric site on μOR may even

favor the binding of some superagonists, supporting a model in which these agonists access a Na$^+$-bound

receptor conformation that is far less accessible to conventional, less potent agonists. Thus, Na$^+$ ions do not

exhibit a uniform inhibitory effect across all agonists, rather the role of Na$^+$ as a negative allosteric

modulator can be in part dependent on agonist affinity. In support of this, at the delta-opioid receptor (δOR),

binding of the highly potent agonist BW373U86 has been reported as insensitive to Na$^+$ ions and guanine

nucleotide, compared to a peptide agonist.[25] However, high-affinity superagonists are not universally

agnostic to the effects of Na$^+$ ions as seen in the aforementioned study with the peptide agonist DSLET and

in our own study with etonitazene. It may reflect a specific interaction, or a coordinated interaction network,

that mediates this Na$^+$ insensitivity.

A recent structural and computational study of µOR partial, full and superagonists demonstrate that G

protein activation proceeds through a series of intermediate conformational states that are separated by

energetic barriers, and that the differences in agonist efficacy arise from the kinetics of traversing these

steps rather than from distinct activation pathways.[26] Superagonists increase receptor dynamics, allowing

for a substantially more rapid transition towards active ternary complexes relative to full and partial

agonists, where partial agonists stabilize more rigid intermediate conformations that struggle to overcome

these energy barriers needed to initiate fully active complexes. Consistent with this, molecular dynamics

simulations initiated from the inactive µOR demonstrate that the superagonist lofentanil can drive

spontaneous transitions into an active receptor-G protein complex in a subset of trajectories, further

indicating that superagonists can bind to inactive receptors and markedly reduce the energetic barrier

required for activation.[26]

The $Na^+$ ion binding site on class A GPCRs comprising of $Asp^{2.50}$, $Asn^{1.50}$, $Ser^{3.39}$, $Asn^{7.45}$, $Ser^{7.46}$, $Asn^{7.49}$

and $Tyr^{7.53}$ is highly conserved across GPCRs.[9] For example, $Asp^{2.50}$, is found in 90% of non-olfactory class

A GPCRs, including the opioid receptor family.[9,27,28] Mutating $Asp^{2.50}$ causes different functional outcomes

depending on the GPCR. For example, with the rat µOR, a D114(2.50) N mutation reduces binding

affinities, potencies and efficacies of both DAMGO and morphine, as well as the binding affinity of the

peptide antagonist, CTAP.[29] Mutagenesis studies of this residue in the closely related δOR transform

antagonists into potent beta-arrestin-biased agonists, suggesting an important role in mediating the R-R*

equilibria.[31] Similar experiments have been performed in a range of Class A GPCRs and lead to a variety

of different pharmacological effects, including reduced downstream signaling and receptor internalization,

increases in agonist affinity with no change in antagonist affinity, no change in agonist affinity, and changes

in antagonist affinity while abolishing G protein signaling.[31-34] However, these effects are not universal.

Thus, mutating this residue at the $\beta_1AR$ does not lead to significant differences in agonist binding affinity -

even in the presence of $Na^+$ - suggesting the activity of this receptor is not modulated by $Na^+$.[35] However

consideration should be taken, as the reported experiments were done in insect Sf9 cell membranes and

using only one agonist, isoprenaline. Further functional, kinetic, and structural studies will need to be performed on additional GPCRs to determine whether insensitivity to $Na^+$ ions and guanine nucleotide with highly potent agonists is a general phenomenon.

The structure of the allosteric $Na^+$ site has been finely tuned to delicately balance the transition between R and R* through stabilizing GPCRs in their ligand-free state. Our results indicate that superagonists of µOR with high intrinsic efficacy may negate this physiological role of $Na^+$ by binding to a $Na^+$-bound receptor state, and facilitating it's transition into an active receptor state that is able to efficiently couple to heterotrimeric G proteins. Alternatively, $Na^+$ ions and guanine nucleotides by promoting the R form of the receptor speed up agonist dissociation. It is possible that very potent agonists hold the receptor in a high affinity active ternary complex due to their slow rates of dissociation from the receptor and markedly slow the ability of $Na^+$ and guanine nucleotides to influence the state of the receptor. Certainly, it is known that carfentanil[36] and BU72[37] dissociate very slowly from the µOR, as does BWU373U86 at δOR[25], and a slow dissociation rate may also explain why it is more difficult for naloxone to reverse respiratory depression caused by highly potent agonists like carfentanil.[38] Furthermore, the rate of dissociation of carfentanil is much slower than that of etonitazene which could explain their different sensitivities to $Na^+$ even when considering their equivalent affinities and functional activity at µOR in this study. Whatever the mechanism, the implication is that a decrease in binding affinity due to the presence of $Na^+$ is not a reliable predictor of efficacy when dealing with very potent compounds in contrast with previous findings.[6,12] Understanding the mechanisms and specific receptor-ligand interactions that mediate $Na^+$ insensitivity may allow for the design of more potent GPCR agonists that are active *in vivo*, in the presence of $Na^+$ and guanine nucleotide. It is important to note that this phenomenon does not occur with existing endogenous opioid peptides, but exclusively with high affinity and intrinsic efficacy ligands.

In conclusion, the influence of $Na^+$ ions and guanine nucleotides on GPCR activation should not be viewed as a uniform or obligatory inhibitory mechanism. The allosteric effect of $Na^+$ ions appears to be probe-

dependent, varying with the chemical scaffold of the bound agonist. Different agonists, or in the case

reported here, superagonists can bind to low affinity receptor and lead to the collapse of the $Na^+$ binding

pocket to generate active states. Alternatively, superagonists may hold the receptor in an active

conformation due to their slow dissociation rates or could even stabilize unique $Na^+$-bound intermediate or

active-like receptor states that remain competent for G protein engagement. In this context, superagonists

may exploit $Na^+$ binding rather than oppose it, thereby promoting receptor activation through $Na^+$-bound

states that conventional agonists alone cannot access. It is possible that the role of $Na^+$ in μOR activation

may encompass multiple parallel trajectories, some requiring ion displacement and others proceeding

through stabilized $Na^+$-bound intermediates that facilitate G protein coupling.

| C6 μ Cells | GTPγ$^{35}$S Binding | | | $^3$H-Diprenorphine Binding | | | | |
|---|---|---|---|---|---|---|---|---|
| Compound | EC$_{50}$ (nM) | Max Stimulation (% of DAMGO, fractional) | Agonism Log $\frac{Max}{EC50\ (M)}$ | K$_i$ (nM) 50 mM Tris | K$_i$ (nM) + 100mM Na$^+$ +10μM GTPγS | Affinity Shift $\frac{Ki\ 100mM\ Na\ +}{Ki\ 50mM\ Tris}$ | K$_i$ (nM) + 300mM Na$^+$ +10μM GTPγS | Affinity Shift $\frac{Ki\ 300mM\ Na\ +}{Ki\ 50mM\ Tris}$ |
| DAMGO | 42.2 (39.0-47.9) | 1 | 7.38 | 4.71 (3.63-6.13) | 618 (421-929) | 131 | 5840 (4040-8470) | 1240 |
| Fentanyl | 233 (178-308) | 0.76 (0.72-0.80) | 6.51 | 8.33 (5.74-12.3) | 433 (239-780) | 52.0 | 873 (488-1590) | 105 |
| Morphine | 195 (148-260) | 0.60 (0.57-0.63) | 6.49 | 3.32 (2.47-4.42) | 254 (169-377) | 76.5 | 280 (155-485) | 84.3 |
| Diprenorphine | - | - | - | 0.12 (0.094-0.16) | - | - | 0.10 (0.077-0.13) | 0.83 |
| Etorphine | 0.79 (0.42-1.39) | 0.96 (0.89-1.04) | 9.08 | 0.40 (0.24-0.66) | 9.66 (6.02-15.4) | 24.2 | 26.6 (19.7-35.6) | 66.5 |
| BU72 | 0.35 (0.24-0.51) | 1.02 (0.96-1.08) | 9.46 | 0.20 (0.14-0.29) | 0.50 (0.33-0.77) | 2.5 | 1.48 (0.96-2.33) | 7.4 |
| Carfentanil | 0.37 (0.28-0.49) | 1.06 (1.01-1.11) | 9.46 | 0.17 (0.13-0.23) | 2.05 (1.37-3.03) | 12.1 | 2.36 (1.63-3.40) | 13.9 |
| Etonitazene | 0.63 (0.38-1.03) | 1.0 (0.92-1.08) | 9.20 | 0.18 (0.14-0.22) | 5.94 (4.19-8.46) | 33 | 29.8 (19.3-45.1) | 166 |
| AAH8 | 0.64 (0.43-0.95) | 0.9 (0.85-0.95) | 9.15 | 0.058 (0.043-0.079) | 0.50 (0.37-0.68) | 8.6 | 1.70 (1.25-2.31) | 29.3 |
| AFN42 | 0.058 (0.043-0.081) | 1.06 (1.01-1.11) | 10.26 | 0.093 (0.072-0.12) | 0.067 (0.054-0.083) | 0.72 | 0.27 (0.19-0.36) | 2.9 |
| AMB46 | 0.86 (0.53-1.37) | 0.96 (0.89-1.03) | 9.04 | 0.077 (0.052-0.11) | 0.91 (0.60-1.42) | 11.8 | 3.39 (2.17-5.23) | 44 |
| AMB47 | 0.59 (0.37-0.92) | 0.98 (0.91-1.05) | 9.22 | 0.15 (0.097-0.28) | 0.32 (0.27-0.38) | 2.1 | 2.29 (1.65-3.15) | 15.3 |

**Table 1: Stimulation of GTPγ$^{35}$S Binding and Inhibition of $^{3}$H-Diprenorphine Binding by Opioid Ligands in C6 μ Cells.** Stimulation of GTPγ$^{35}$S binding at the μOR was performed using membranes from C6 cells with various concentrations of opioid agonists. Maximal stimulation of GTPγ$^{35}$S binding was determined by 10 μM of the reference agonist, DAMGO. Effective concentrations (EC$_{50}$) and the fractional maximal response (Max Stimulation) were determined by nonlinear regression analysis of concentration-response curves from GTPγ$^{35}$S binding assays. Agonism (Log [Max/EC50]) was calculated using EC$_{50}$ and max stimulation values for each opioid agonist. $^{3}$H-Diprenorphine binding assays were performed using 0.2 – 0.5 nM of the radioligand in the following three buffers: **1.** 50 mM Tris buffer, **2.** 50 mM Tris, 100 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM GTPγS, or **3.** 50 mM Tris, 300 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM GTPγS. The affinity shift was determined for 100 mM Na$^{+}$ and 300 mM Na$^{+}$ conditions. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate or quadruplicate (n ≥ 3).

**Figure 1: Chemical Structures of Opioid Agonists and Peptidomimetics.** (A) DAMGO, (B) Fentanyl, (C) Morphine, (D) Etorphine, (E) BU72, (F) Carfentanil, (G) Etonitazene, (H) AAH8, (I) AFN42, (J) AMB46, and (K) AMB47

**Figure 2: Stimulation of GTPγ³⁵S Binding and Inhibition of ³H-Diprenorphine Binding Concentration-Response Curves in C6 µ Cells.** (A) GTPγ³⁵S binding assays were performed using membranes from C6 cells with various concentrations of opioid agonists. Maximal stimulation of GTPγ³⁵S binding was produced by 10 µM of the reference agonist, DAMGO. (B) Radioligand competition binding assays were performed in the following buffers: 50 mM Tris (circles), 50 mM Tris, 100 mM NaCl, 5 mM MgCl₂ 1 mM EDTA, 10 µM GTPγS (squares), or 50 mM Tris, 300 mM NaCl, 5 mM MgCl₂ 1 mM EDTA, 10 µM GTPγS (triangles). These assays used 20 µg membrane protein from C6 cells, 0.2 – 0.5 nM ³H-Diprenorphine, and various concentrations of the following opioid ligands: Diprenorphine, DAMGO, Fentanyl, Etorphine, BU72, Carfentanil, Etonitazene, AAH8, AFN42, AMB46, AMB47. All experiments were performed in duplicate or quadruplicate (n ≥ 3).

**Figure 3: Stimulation of GTPγ³⁵S Binding in 100 mM NaCl and KCl buffer.** Stimulation of GTPγ³⁵S binding at the μOR was performed using membranes from C6 cells with increasing concentrations of opioid agonists. Maximal stimulation of GTPγ³⁵S binding was determined by 10 μM of the reference agonist, DAMGO. Effective concentrations (EC$_{50}$) were determined by nonlinear regression analysis of concentration-response curves from GTPγ³⁵S binding assays. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate (n ≥ 3).

| C6 $\mu$ Cells | ³H-Diprenorphine Binding | | |
|---|---|---|---|
| Compound | $K_i$ (nM) 50 mM Tris | $K_i$ (nM) + 100mM $K^+$ +10$\mu$M GTP$\gamma$S | Affinity Shift $\dfrac{Ki\ 100mM\ K +}{Ki\ 50mM\ Tris}$ |
| DAMGO | 4.71 (3.63-6.13) | 111 (66.8-187) | 23.6 |
| Fentanyl | 8.33 (5.74-12.3) | 218 (140-344) | 26.2 |
| Morphine | 3.32 (2.47-4.42) | 150 (93.9-246) | 45.2 |
| Diprenorphine | 0.12 (0.094-0.16) | 1.96 (1.29-2.94) | 16.3 |
| Carfentanil | 0.17 (0.13-0.23) | 1.19 (0.81-1.76) | 7 |
| AFN42 | 0.093 (0.072-0.12) | 1.68 (1.02-2.86) | 18.1 |

**Table 2. Inhibition of ³H-Diprenorphine Binding in 100 mM KCl.** Radioligand competition binding assays were performed in 50 mM Tris, 100 mM KCl, 5 mM $MgCl_2$ 1 mM EDTA, 10 $\mu$M GTP$\gamma$S. These assays used 20 $\mu$g membrane protein from C6 cells, 0.2 – 0.5 nM ³H-Diprenorphine, and various concentrations of opioid ligand. Experiments were performed in duplicate or quadruplicate (n $\geq$ 3) and are presented as mean with 95% confidence intervals.

| CHO μ Cells | $^3$H-Diprenorphine Binding | | |
|---|---|---|---|
| **Compound** | **$K_i$ (nM)** 50 mM Tris | **$K_i$ (nM)** + 100mM Na$^+$ +10μM GTPγS | **Affinity Shift** $\dfrac{\text{Ki 100mM Na +}}{\text{Ki 50mM Tris}}$ |
| DAMGO | 0.65 (0.40-1.05) | 529 (293-946) | 814 |
| BU72 | 0.27 (0.18-0.39) | 0.73 (0.54-1.01) | 2.7 |
| Carfentanil | 0.06 (0.035-0.10) | 2.09 (1.56-2.79) | 34.8 |

**Table 3: Inhibition of $^3$H-Diprenorphine Binding by Opioid Ligands in CHO μ Cells.** $^3$H-Diprenorphine binding assays were performed using 0.2 – 0.5 nM of the radioligand in the following two buffers: **1.** 50 mM Tris buffer, and **2.** 50 mM Tris, 100 mM NaCl, 5 mM MgCl$_2$ 1 mM EDTA, and 10 μM cold GTPγS. The affinity shift was determined for 100 mM Na$^+$ conditions. Data are represented as mean plus 95% confidence intervals and all experiments were performed in duplicate (n ≥ 3).

**Figure 4: Intrinsic Efficacy Plots for C6 μ.** Agonism (Log [Max/EC$_{50}$]) was determined from stimulation of GTPγ$^{35}$S binding assays and Affinity Shift (Log [Ki, 100 mM Na$^+$ / K$_i$ 50 mM Tris] in **(A)** or Log [Ki, 300 mM Na$^+$ / K$_i$ 50 mM Tris] in **(B)** was determined from $^3$H-Diprenorphine competition binding assays. Data was plotted for each of the following ligands: DAMGO, Fentanyl, Morphine, Etorphine, BU72, Carfentanil, Etonitazene, AFN42, AAH8, AMB46, and AMB47. r$^2$ = 0.63, p$^{**}$ = 0.003 **(A)**; r$^2$ = 0.45, p$^*$ = 0.02 **(B)**

# References

1. Ehlert, F. J. (2018). Analysis of biased agonism. Progress in Molecular Biology and Translational Science, 63–104. https://doi.org/10.1016/bs.pmbts.2018.08.001

2. Traynor, J. R., & Nahorski, S. R. (1995). Modulation by mu-opioid agonists of guanosine-5'-O-(3-[35S]thio)triphosphate binding to membranes from human neuroblastoma SH-SY5Y cells. Molecular Pharmacology, 47(4), 848–854. https://doi.org/10.1016/s0026-895x(25)08634-1

3. Kenakin, T. (2017). A scale of agonism and allosteric modulation for assessment of selectivity, bias, and receptor mutation. *Carolina Digital Repository (University of North Carolina at Chapel Hill)*. https://doi.org/10.17615/hhsf-de46

4. Kenakin, T., & Christopoulos, A. (2011). Analytical pharmacology: the impact of numbers on pharmacology. Trends in Pharmacological Sciences, 32(4), 189–196. https://doi.org/10.1016/j.tips.2011.01.002

5. Herenbrink, C. K., Sykes, D. A., Donthamsetti, P., Canals, M., Coudrat, T., Shonberg, J., Scammells, P. J., Capuano, B., Sexton, P. M., Charlton, S. J., Javitch, J. A., Christopoulos, A., & Lane, J. R. (2016). The role of kinetic context in apparent biased agonism at GPCRs. Nature Communications, 7(1). https://doi.org/10.1038/ncomms10842

6. Pert, C. B., Pasternak, G., & Snyder, S. H. (1973). Opiate agonists and antagonists discriminated by receptor binding in brain. *Science*, *182*(4119), 1359–1361. https://doi.org/10.1126/science.182.4119.1359

7. Puttfarcken, P., Werling, L. L., Brown, S. R., Cote, T. E., & Cox, B. M. (1986). Sodium regulation of agonist binding at opioid receptors. I. Effects of sodium replacement on binding at mu- and delta-type receptors in 7315c and NG108-15 cells and cell membranes. *PubMed*, *30*(2), 81–89. https://doi.org/10.1016/s0026-895x(25)10340-4

8. Livingston, K. E. and J. R. Traynor (2014). Disruption of the Na+ ion binding site as a mechanism for positive allosteric modulation of the mu-opioid receptor. Proceedings of the National Academy of Science **111**(51): 18369-18374. https://doi.org/10.1073/pnas.1415013111

9. Katritch, V., G. Fenalti, E. E. Abola, B. L. Roth, V. Cherezov and R. C. Stevens (2014). Allosteric sodium in class A GPCR signaling. Trends in biochemical sciences **39**(5): 233-244. https://doi.org/10.1016/j.tibs.2014.03.002

10. Liu, W., E. Chun, A. A. Thompson, P. Chubukov, F. Xu, V. Katritch, G. W. Han, C. B. Roth, L. H. Heitman, A. P. Ijzerman, V. Cherezov and R. C. Stevens (2012). Structural basis for allosteric regulation of GPCRs by sodium ions. Science (New York, N.Y.) **337**(6091): 232-236. https://doi.org/10.1126/science.1219218

11. White, K. L., M. T. Eddy, Z.-G. Gao, G. W. Han, T. Lian, A. Deary, N. Patel, K. A. Jacobson, V. Katritch and R. C. Stevens (2018). Structural Connection between Activation Microswitch and Allosteric Sodium Site in GPCR Signaling. Structure **26**(2): 259-269.e255. https://doi.org/10.1016/j.str.2017.12.013

12. Childers, S. R., & Snyder, S. H. (1980). Differential regulation by guanine nucleotides of opiate agonist and antagonist receptor interactions. Journal of Neurochemistry, 34(3), 583–593. https://doi.org/10.1111/j.1471-4159.1980.tb11184.x

13. Lee, K. O., Akil, H., Woods, J. H., & Traynor, J. R. (1999). Differential binding properties of oripavines at cloned μ- and δ-opioid receptors. European Journal of Pharmacology, 378(3), 323–330. https://doi.org/10.1016/s0014-2999(99)00460-4

14. Faouzi, A., Wang, H., Zaidi, S. A., DiBerto, J. F., Che, T., Qu, Q., Robertson, M. J., Madasu, M. K., Daibani, A. E., Varga, B. R., Zhang, T., Ruiz, C., Liu, S., Xu, J., Appourchaux, K., Slocum, S. T., Eans, S. O., Cameron, M. D., Al-Hasani, R., . . . Majumdar, S. (2022). Structure-based design of bitopic ligands for the μ-opioid receptor. *Nature*, *613*(7945), 767–774. https://doi.org/10.1038/s41586-022-05588-y

15. Bender, A. M., N. W. Griggs, J. P. Anand, J. R. Traynor, E. M. Jutkiewicz and H. I. Mosberg (2015). Asymmetric Synthesis and in Vitro and in Vivo Activity of Tetrahydroquinolines Featuring a Diverse Set of Polar Substitutions at the 6 Position as Mixed-Efficacy mu Opioid Receptor/delta Opioid Receptor Ligands. ACS Chem Neurosci **6**(8): 1428-1435. https://doi.org/10.1021/acschemneuro.5b00100

16. Harland, A. A., L. Yeomans, N. W. Griggs, J. P. Anand, I. D. Pogozheva, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2015). Further Optimization and Evaluation of Bioavailable, Mixed-Efficacy mu-Opioid Receptor (MOR) Agonists/delta-Opioid Receptor (DOR) Antagonists: Balancing MOR and DOR Affinities. J Med Chem **58**(22): 8952-8969. https://doi.org/10.1021/acs.jmedchem.5b01270

17. Harland, A. A., A. M. Bender, N. W. Griggs, C. Gao, J. P. Anand, I. D. Pogozheva, J. R. Traynor, E. M. Jutkiewicz and H. I. Mosberg (2016). Effects of N-Substitutions on the Tetrahydroquinoline (THQ) Core of Mixed-Efficacy mu-Opioid Receptor (MOR)/delta-Opioid Receptor (DOR) Ligands. J Med Chem **59**(10): 4985-4998. https://doi.org/10.1021/acs.jmedchem.6b00308

18. Nastase, A. F., N. W. Griggs, J. P. Anand, T. J. Fernandez, A. A. Harland, T. J. Trask, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2018). Synthesis and Pharmacological Evaluation of Novel C-8 Substituted Tetrahydroquinolines as Balanced-Affinity Mu/Delta Opioid Ligands for the Treatment of Pain. ACS Chem Neurosci **9**(7): 1840-1848. https://doi.org/10.1021/acschemneuro.8b00139

19. Montgomery, D., J. P. Anand, M. A. Baber, J. J. Twarozynski, J. G. Hartman, L. J. Delong, J. R. Traynor and H. I. Mosberg (2019). Structure-Activity Relationships of 7-Substituted Dimethyltyrosine-Tetrahydroisoquinoline Opioid Peptidomimetics. Molecules **24**(23). https://doi.org/10.3390/molecules24234302

20. Montgomery, D., J. P. Anand, N. W. Griggs, T. J. Fernandez, J. G. Hartman, A. A. Sanchez-Santiago, I. D. Pogozheva, J. R. Traynor and H. I. Mosberg (2019). Novel Dimethyltyrosine-Tetrahydroisoquinoline Peptidomimetics with Aromatic Tetrahydroisoquinoline Substitutions Show in Vitro Kappa and Mu Opioid Receptor Agonism. ACS Chem Neurosci **10**(8): 3682-3689. https://doi.org/10.1021/acschemneuro.9b00250

21. Nastase, A. F., J. P. Anand, A. M. Bender, D. Montgomery, N. W. Griggs, T. J. Fernandez, E. M. Jutkiewicz, J. R. Traynor and H. I. Mosberg (2019). Dual Pharmacophores Explored via Structure-Activity Relationship (SAR) Matrix: Insights into Potent, Bifunctional Opioid Ligand Design. J Med Chem **62**(8): 4193-4203. https://doi.org/10.1021/acs.jmedchem.9b00378

22. Pasternak, G. W., & Pan, Y. (2013). MU Opioids and their Receptors: Evolution of a Concept. *Pharmacological Reviews*, *65*(4), 1257–1317. https://doi.org/10.1124/pr.112.007138

23. Katritch, V., Fenalti, G., Abola, E. E., Roth, B. L., Cherezov, V., & Stevens, R. C. (2014). Allosteric sodium in class A GPCR signaling. *Trends in Biochemical Sciences*, *39*(5), 233–244. https://doi.org/10.1016/j.tibs.2014.03.002

24. Rothman, R. B., Xu, H., Wang, J. B., Partilla, J. S., Kayakiri, H., Rice, K. C., & Uhl, G. R. (1994). The ligand-selectivity of cloned human and rat opiate mu receptors studied with [125I]IOXY-AGO. *Regulatory Peptides*, *54*(1), 249–250. https://doi.org/10.1002/syn.890210109

25. Childers, S. R., Fleming, L. M., Selley, D. E., McNutt, R. W., & Chang, K. J. (1993). BW373U86: a nonpeptidic delta-opioid agonist with novel receptor-G protein-mediated actions in rat brain membranes and neuroblastoma cells. *Molecular Pharmacology*, *44*(4), 827–834. https://doi.org/10.1016/s0026-895x(25)13269-0

26. Robertson, M. J., Modak, A., Papasergi-Scott, M. M., Hu, M., Peroto, M. C., Varga, B. R., Majumdar, S., Kalathur, R., Blanchard, S. C., & Skiniotis, G. (2025). Non-equilibrium snapshots of ligand efficacy at the μ-opioid receptor. *Nature*. https://doi.org/10.1038/s41586-025-10056-4

27. Parker, M. S., Wong, Y. Y., & Parker, S. L. (2008). An ion-responsive motif in the second transmembrane segment of rhodopsin-like receptors. *Amino Acids*, *35*(1), 1–15. https://doi.org/10.1007/s00726-008-0637-6

28. Shang, Y., LeRouzic, V., Schneider, S., Bisignano, P., Pasternak, G. W., & Filizola, M. (2014). Mechanistic Insights into the Allosteric Modulation of Opioid Receptors by Sodium Ions. *Biochemistry*, *53*(31), 5140–5149. https://doi.org/10.1021/bi5006915

29. Xu, W., Ozdener, F., Li, J., Chen, C., De Riel, J., Weinstein, H., & Liu-Chen, L. (1999). Functional role of the spatial proximity of Asp114(2.50) in TMH 2 and Asn332(7.49) in TMH 7 of the μ opioid receptor. *FEBS Letters*, *447*(2–3), 318–324. https://doi.org/10.1016/s0014-5793(99)00316-6

30. Fenalti, G., Giguere, P. M., Katritch, V., Huang, X., Thompson, A. A., Cherezov, V., Roth, B. L., & Stevens, R. C. (2014). Molecular control of δ-opioid receptor signalling. *Nature*, *506*(7487), 191–196. https://doi.org/10.1038/nature12944

31. Proulx, C. D., Holleran, B. J., Boucard, A. A., Escher, E., Guillemette, G., & Leduc, R. (2008). Mutational analysis of the conserved ASP2.50and ERY motif reveals signaling bias of the urotensin II receptor. *Molecular Pharmacology*, *74*(3), 552–561. https://doi.org/10.1124/mol.108.045054

32. H Barbhaiya, McClain R, A Ijzerman, Rivkees SA. Site-directed mutagenesis of the human A1 adenosine receptor: influences of acidic and hydroxy residues in the first four transmembrane domains on ligand binding (1996). *Molecular Pharmacology*. 50(6):1635-1642. doi:https://doi.org/10.1016/s0026-895x(25)09624-5

33. Gao, Z., Kim, S., Gross, A. S., Chen, A., Blaustein, J. B., & Jacobson, K. A. (2003). Identification of essential residues involved in the allosteric modulation of the human A3Adenosine receptor. *Molecular Pharmacology*, *63*(5), 1021–1031. https://doi.org/10.1124/mol.63.5.1021

34. Bihoreau, C., Monnot, C., Davies, E., Teutsch, B., Bernstein, K. E., Corvol, P., & Clauser, E. (1993). Mutation of Asp74 of the rat angiotensin II receptor confers changes in antagonist affinities and abolishes G-protein coupling. *Proceedings of the National Academy of Sciences*, *90*(11), 5133–5137. https://doi.org/10.1073/pnas.90.11.5133

35. Miller-Gallacher, J. L., Nehmé, R., Warne, T., Edwards, P. C., Schertler, G. F. X., Leslie, A. G. W., & Tate, C. G. (2014). The 2.1 Å Resolution Structure of Cyanopindolol-Bound β1-Adrenoceptor Identifies an Intramembrane Na+ Ion that Stabilises the Ligand-Free Receptor. *PLoS ONE*, *9*(3), e92727. https://doi.org/10.1371/journal.pone.0092727

36. Alhosan, N., Cavallo, D., Santiago, M., Kelly, E., & Henderson, G. (2024). Slow dissociation kinetics of fentanyls and nitazenes correlates with reduced sensitivity to naloxone reversal at the µ-opioid receptor. *British Journal of Pharmacology*, *182*(4), 969–987. https://doi.org/10.1111/bph.17376

37. Huang, W., Manglik, A., Venkatakrishnan, A. J., Laeremans, T., Feinberg, E. N., Sanborn, A. L., Kato, H. E., Livingston, K. E., Thorsen, T. S., Kling, R. C., Granier, S., Gmeiner, P., Husbands, S. M., Traynor, J. R., Weis, W. I., Steyaert, J., Dror, R. O., & Kobilka, B. K. (2020). Author

Correction: Structural insights into μ-opioid receptor activation. *Nature*, *584*(7820), E16.

https://doi.org/10.1038/s41586-020-2542-z

38. Flynn, S. M., & France, C. P. (2025). Comparison of ventilatory depression by carfentanil, fentanyl, and heroin, and reversal by opioid receptor antagonists in rats. *Neuropharmacology*, 110601. https://doi.org/10.1016/j.neuropharm.2025.110601