Smooth muscle LRRC8A knockout reduces $O_2^{\cdot-}$ influx, inflammation, senescence and atherosclerosis

Sourav Panja[1*], Hyehun Choi[1*], Hong N. Nguyen[1], Julia Russolillo[1], Sergey Dikalov[2], Ryan J. Stark[1] and Fred S. Lamb[1]

[1]Department of Pediatrics, Division of Critical Care, Vanderbilt University Medical Center, Nashville, TN 37232

[2]Department of Internal Medicine, Division of Clinical Pharmacology, Vanderbilt University Medical Center, Nashville, TN 37232

* These authors contributed equally to this work.

Running title: LRRC8A KO in atherosclerosis

Word count:

Number of figures:

Corresponding author:
Fred S. Lamb MD, PhD
Vanderbilt University Medical Center
Department of Pediatrics
2215 Garland Avenue, Light Hall-1055D
Nashville, TN 37232-3122
Phone: +1 (615) 936-1301 Fax: +1 (615) 936-3467
fred.s.lamb@vanderbilt.edu

ABSTRACT

Background: Leucine Rich Repeat Containing 8A (LRRC8A) anion channels (VRACs) associate with NADPH oxidase 1 (Nox1) and support extracellular superoxide ($O_2^{\cdot-}$) production, inflammation, and contractility in vascular smooth muscle cells (VSMCs). We proposed previously that VRACs also support influx of $O_2^{\cdot-}$ to produce localized cytoplasmic signaling.

Methods: We assayed $O_2^{\cdot-}$ influx and guided by changes in mRNA expression (RNAseq) tested multiple phenotypes of cultured LRRC8A knockout (KO) VSMCs. Aortic atherosclerotic burden and inflammation, and mesenteric vascular reactivity were compared between wild type (WT) apolipoprotein E null (ApoE$^{-/-}$) mice and VSMC-specific LRRC8A KO, ApoE$^{-/-}$ mice.

Results: KO cells were less permeable to extracellular $O_2^{\cdot-}$, produced less mitochondrial $O_2^{\cdot-}$, and experienced less oxidant stress (GSH/GSSG, lipid peroxidation, Nrf2 activity) than WT. RNAseq and reporter assays demonstrated reduced pro-inflammatory (NF-κB and Hif1α) transcription. KO cells also under-expressed multiple senescence markers and had longer telomeres. Inflammation causes a metabolic shift from oxidative phosphorylation (OCR) to glycolysis (ECAR). Both pathways were less active in KO cells, as was expression of glycolytic enzymes. Mitochondrial membrane potential was lower in KO cells, but ADP/ATP and NADP$^+$/NADPH were unaltered, suggestive of lower energy demand. Consistent with this, proliferation and migration were both reduced in KO cells. Protein-protein interaction analysis of RNAseq (PPI Hub) identified the Epidermal Growth Factor Receptor (EGFR) signaling pathway. Activating phosphorylation of EGFR (Y1068) as well as downstream AKT (S473 and T308) were reduced. Following 15 weeks of exposure to a high fat diet (42%) VSMC-specific LRRC8A$^{-/-}$ and 8A$^{+/-}$, ApoE$^{-/-}$ mice had less atherosclerotic lesion area, and reduced aortic senescence (β-Gal), inflammation (ICAM, VCAM) and proliferation (PCNA) marker expression. Mesenteric artery vasomotor function was also preserved compared to WT, ApoE$^{-/-}$ controls, and the abundance of MYPT1 and CPI17 was lower

in KO vessels. Uptake of oxidized LDL (OxLDL) was significantly reduced in KO VSMCs and similarly impaired by VRAC block in WT cells.

Conclusions: Loss of LRRC8A reduces $O_2^{\cdot-}$ influx, oxidative stress, inflammation and senescence, lowers energy demand, and in the setting of hypercholesterolemia impairs uptake of oxidized LDL and abrogates atherosclerosis. VRAC inhibitors warrant further investigation as therapeutic vascular anti-inflammatory agents.

KEYWORDS

Leucine Rich Repeat Containing 8, vascular smooth muscle, inflammation, oxidative stress, senescence, atherosclerosis

INTRODUCTION

LRRC8 family genes (LRRC8A-E) encode volume-regulated anion channels (VRACs), originally identified by their activation in response to hypotonicity and contribution to a regulatory volume decrease. LRRC8A is obligatory for the trafficking of all VRACs to the plasma membrane[1], where they have also evolved to serve other important roles[2-5]. These channels are regulated by NADPH oxidase (Nox)-derived oxidants which are produced in response to cell swelling, and to signaling molecules including angiotensin II, TNFα and growth factors in vascular smooth muscle cells (VSMCs)[6,7]. VRACs support proliferation in multiple cell types including VSMCs[8,9], and the magnitude of their current varies with the cell cycle[10,11], being more active in proliferative cells[12]. TNFα activates VRACs[13,14] and NF-κB leading to VSMC proliferation[15]. All of these signals require production of extracellular superoxide ($O_2^{\cdot-}$) by Nox1 which physically-associates with LRRC8A, 8C and 8D, and Nox1 activity depends upon VRAC function[16,17].

VRACs are linked to redox balance[18] not only by their ability to both impact oxidant-dependent signaling[16,17,19], their activity is modulated by oxidants[16,20]. We previously proposed that VRACs also provide a mechanism by which newly created extracellular $O_2^{\cdot-}$ can enter the cytoplasm to drive localized redox reactions (See[19] for review). This concept was supported by several indirect observations. TNFα receptor endocytosis and c-Jun N-terminal kinase activation were impaired by extracellular superoxide dismutase (SOD, degrades $O_2^{\cdot-}$ to produce H_2O_2), while catalase (degrades H_2O_2 to $H_2O + O_2$) did not affect these signals[17]. Furthermore, activation of apoptosis signaling-related kinase 1 (ASK1) by TNFα, which depends upon oxidation of cytoplasmic thioredoxin, is Nox1-dependent and potentiated by knockdown of extracellular SOD[19]. In addition, TNFα-induced RhoA activation and sulfenylation (oxidation) of

the cytoplasmic LRRC8A-associated scaffolding protein Myosin Phosphatase Rho Interacting Protein (MPRIP), are both LRRC8A-dependent[21]. Recently this hypothesis was supported by direct evidence for LRRC8A-dependent $O_2^{\cdot-}$ influx in neurons, and disruption of VRAC function was protective against oxidative injury induced by NMDA or transient ischemia[22].

Noxes and the extracellular or endosomal $O_2^{\cdot-}$ that they produce play a fundamental role in vascular inflammation[23,24]. $O_2^{\cdot-}$ is the conjugate base of $HO_2^{\cdot}$ with a very short half-life and is rapidly converted to H_2O_2 either spontaneously or enzymatically by superoxide dismutase. It can also react with nitric oxide to form peroxynitrite[25], with metals to create more reactive oxidants (e.g. $OH^{\cdot}$), or it can induce membrane lipid peroxidation or directly oxidize proteins[26]. Access to the cytoplasm would allow these reactions to occur in the direct vicinity of signaling proteins. Thus, association of Nox1 with LRRC8A may allow even small amounts of $O_2^{\cdot-}$ to target proteins that bind to the cytoplasmic leucine rich domains of the channel[21]. In this scenario the short half-life and limited diffusion distance of $O_2^{\cdot-}$ may limit off-target oxidation.

Atherosclerosis is a chronic vascular inflammatory disease linked to hypercholesterolemia-induced oxidative stress[27]. It underlies the pathogenesis of coronary artery disease and stroke, the top two causes of death worldwide. VSMCs play a critical pathophysiological role as lineage tracing has revealed that the majority of the fat-containing foam cells in the vascular wall are of smooth muscle origin[28]. Atherosclerotic VSMCs are inflamed[29], under oxidative stress[30], have incurred telomere damage[31], are more frequently senescent[32], and utilize glycolysis as a preferred energy source[33]. These changes are driven by activation of scavenger receptors such as LOX-1 and CD36 that take up oxidized LDL and drive inflammation, oxidant stress, and senescence[34].

Here we provide direct evidence that VRACs promote $O_2^{\cdot-}$ access to the cytoplasm. We then explore how the loss of these channels in VSMCs impacts gene expression, oxidative stress, mitochondrial function, metabolism, proliferation and signaling pathways related to inflammation, senescence and longevity. Compared to wild type (WT) controls, cultured LRRC8A knockout (KO) cells are under less oxidative stress, utilize less energy, and are resistant to stimuli that drive vascular disease. This motivated assessment of how losing these channels, specifically in VSMCs, affected the development of atherosclerosis and resistance artery dysfunction in hypercholesterolemic mice lacking the ApoE receptor ($ApoE^{-/-}$). Consistent with the *in vitro* characteristics of cultured KO VSMCs, smooth muscle LRRC8A KO mice, and to a significant degree heterozygous animals (HET), were protected from disease.

METHODS

A detailed description of all materials and methods is provided in the Data Supplement.

Animal Studies and Ethics Statement

All animal procedures were approved by the Institutional Animal Care and Use Committee of Vanderbilt University Medical Center and complied with the ARRIVE guidelines. Vascular smooth muscle cell (VSMC)-specific *Lrrc8a* knockout (KO) mice were generated by crossing *Lrrc8a*^{fl/fl} mice with transgenic mice expressing Cre recombinase under the SM22α promoter. To induce atherosclerosis, these mice were crossed onto an Apolipoprotein E-null (*ApoE*^{−/−}) background and fed a high-fat diet (42% Fat, TD88137, Harlan Teklad) for 15 weeks after weaning.

Cell Culture and In Vitro Assays

Primary aortic VSMCs were isolated from male KO and WT littermates as previously described[35] and used for experiments between passages 6 and 15. Cellular redox status was assessed in dye-loaded cells by confocal microscopy (for DHE and ROSstar 550 imaging) and flow cytometry (for DHE, ROSstar 550, MitoSOX, and TMRE), measuring superoxide influx, mitochondrial/cytosolic superoxide, mitochondrial membrane potential, membrane lipid peroxidation (MDA assay), and the reduced-to-oxidized glutathione (GSH/GSSG) ratio. Metabolic flux was analyzed using a Seahorse XF Analyzer to determine the extracellular acidification rate (ECAR) and oxygen consumption rate (OCR). VSMC proliferation, migration, and senescence were quantified by MTT, scratch-wound healing assays, and senescence-associated β-galactosidase activity, respectively. Uptake of fluorescently labeled Dil-OxLDL (ThermoFisher) was quantified by confocal microscopy using ImageJ software.

Molecular and Signaling Analyses

Global transcriptomic changes were profiled by bulk RNA sequencing (Illumina NovaSeq), with differential expression analysis performed in Partek Flow using the DESeq2 algorithm. Key protein expression and phosphorylation status of signaling intermediates (e.g., pEGFR, pAKT) were validated by Western blotting using an infrared imaging system (LI-COR Odyssey). Transcription factor activity (NF-κB, Hif-1α, Nrf2) was quantified using lentiviral luciferase reporters, with activity normalized for infection efficiency. Relative telomere length was determined by quantitative PCR.

Statistical Analysis

Data are presented as mean ± SEM. Statistical analyses were performed using GraphPad Prism 9. Comparisons were made using a one sample t- test, unpaired Student's t-test or one-way

ANOVA with Tukey's post hoc test, as appropriate. A (*$P<0.05$, **$P<0.01$, ***$P<0.001$) was considered statistically significant.

RESULTS

Superoxide flux

To detect influx of $O_2^{\cdot-}$, unstimulated WT and LRRC8A KO HEK293 cells were loaded with DHE (5 µM.) for 30 min, washed 3 times and then exposed to $O_2^{\cdot-}$ for an additional 30 min under culture conditions (McCoy's 5A medium) (Fig. 1A). Oxidized (2-OH)-ethidium, a highly specific reaction product of ethidium and $O_2^{\cdot-}$, was quantified by HPLC and normalized to total protein (pmol/mg protein). Ethidium was oxidized by extracellular $O_2^{\cdot-}$ in WT but not LRRC8A KO cells (Fig. 1A). To control the nature and activity of the VRACs present in the plasma membrane, HCT116 cells lacking all LRRC8 isoforms (LRRC8$^{-/-}$) were made to express LRRC8A, LRRC8C or LRRC8D with C-terminal GFP tags. Under isotonic conditions homomeric LRRC8A channels are minimally active[36], even in the presence of a C-terminal tag[37]. However, when LRRC8C or 8D channels are given the intracellular loop of LRRC8A they are trafficked to the plasma membrane[38], and incorporation of a C-terminal GFP tag renders them active under isotonic conditions[16,37]. DHE loaded cells were exposed to extracellular $O_2^{\cdot-}$ (Fig. 1B), then fixed and imaged by confocal microscopy (Fig. 1C). Transfection efficiency was less than 100%, allowing comparison of the DHE signal between expressing (GFP positive) and non-expressing (GFP negative) cells in single microscopic fields. Expression of LRRC8A-GFP or LRRC8D-GFP had no visually discernable effect on DHE fluorescence in either resting or $O_2^{\cdot-}$-exposed cells, while the presence of LRRC8C-GFP cell-specifically increased the subjective DHE signal after $O_2^{\cdot-}$ treatment. To quantify this association, flow cytometry was used to

compare DHE signal intensity under control conditions and after exposure to $O_2^{\cdot-}$ (Fig 1C). Resting LRRC8C-GFP and LRRC8D-GFP expressing cells both had a higher average resting DHE fluorescence than those expressing inactive LRRC8A-GFP and this difference was larger in cells expressing LRRC8C. Following exposure to $O_2^{\cdot-}$ the DHE signal increased in all cells. This increase did not differ between LRRC8A-GFP and LRRC8D-GFP expressors but was significantly greater in cells expressing LRRC8C-GFP. This suggests that LRRC8C supports both resting endogenous superoxide production and $O_2^{\cdot-}$ influx better than LRRC8A or LRRC8D.

Oxidative Stress

NADPH oxidase activity can trigger mitochondrial $O_2^{\cdot-}$ production in a feed-forward cycle of ROS production[39]. We hypothesized that reduced Nox activity and $O_2^{\cdot-}$ influx in LRRC8A KO VSMCs[17] would reduce mitochondrial ROS production. Confocal microscopy confirmed that ROSstar (cytoplasmic + mitochondria) intensity appeared to be markedly lower in KO cells compared to WT, indicating reduced overall ROS levels. Therefore MitoSOX (mitochondria-specific) and ROSstar oxidant signals were quantitatively compared by flow cytometry, and both were significantly reduced in KO compared to WT cells (Fig. 1E).

Membrane lipid peroxidation occurs under oxidant stress. This was quantified under resting conditions and after TNFα (10 ng/ml) exposure. Resting levels were lower but not significantly so (P = 0.06, N = 4) in KO cells but TNFα stimulation increased MDA levels only in WT cells (Fig. 1F). The ratio of reduced to oxidized glutathione (GSH/GSSG) was significantly higher in KO cells (Fig. 1G), indicative of a more reduced intracellular environment. As an integrated assessment of oxidative stress, Nrf2 transcriptional activity was measured using a luciferase reporter. Nrf2 activity was significantly lower in KO VSMCs (Fig. 1H), consistent with a

reduced need for antioxidant protection. Protein products of Nrf2 target genes, including extracellular superoxide dismutase (SOD3) and heme oxygenase (HO-1) were reduced in KO cells (Suppl. Fig. 1A), as was the expression of mRNA (qPCR) for other Nrf2 target genes including glutathione peroxidase 3 (*Gpx3*), glutathione S-transferase alpha 3 (*Gsta3*), and heme oxygenase-1 (*Hmox1*) (Suppl. Fig. 1B). Expression of the membrane components of the Nox1 complex (Nox1, p22phox) were unchanged in LRRC8A KO VSMCs. However, the abundance of the cytosolic activator Noxa1 subunit was significantly lower in KO cells. Of note, VSMC-specific KO of Noxa1 protected against both neointimal hyperplasia following endovascular injury and atheroma formation in ApoE$^{-/-}$ mice[40]. These data, and many others were normalized to tubulin which was not differentially expressed in KO cells (Suppl. Fig. 1D).

Transcriptomic Analysis and HIF-1α Signaling

Bulk RNA sequencing was performed on primary cultured aortic VSMCs from 3 pairs of WT and KO littermates (Suppl. Fig. 2A). Principal component analysis showed clear segregation of WT and KO samples and a heatmap of global gene expression patterns indicated distinct clustering by genotype (Suppl. Fig. 2 B and C). Differential expression analysis using DESeq2 identified 573 genes revealed significant expression changes between KO and WT VSMCs. Among these, 206 genes were upregulated, and 367 genes downregulated in KO cells (p ≤ 0.05 and Fold Change ≥ 1.5). A volcano plot is provided in Suppl. Fig. 2D.

To interpret the impact of these transcriptional changes we performed pathway enrichment analysis on the differentially expressed genes (DEGs) using KEGG pathways via Enrichr. Fig. 2A presents the top eight affected KEGG pathways based on all DEGs. Genes showing increased expression are in red and decreased genes are in green with bar height indicating the combined

score/enrichment. Statistically significant pathways were most frequently associated with; 1) metabolism (lipolysis, glutathione metabolism), 2) inflammation (Hif-1α and MAPK signaling), 3) stress response (oxidative stress resistance), and 4) muscle contraction (Fig. 2B, left panel). Key pathways were primarily driven by genes that were downregulated in KO cells (green). When analysis was confined to the 367 downregulated genes the most affected functional categories were inflammatory/immune pathways, metabolic processes, stress-response pathways, muscle contraction and the response to hypoxia (Fig. 2B, left panel). The top KEGG pathways enriched exclusively among downregulated genes confirmed an anti-inflammatory signature (Fig. 2B, right panel). This aligns with our previous findings that LRRC8A knockdown impaired NF-κB signaling[16,17]. Numerous NF-κB target genes and cytokine signaling components were expressed at significantly lower levels in KO VSMCs at baseline, including receptors for interleukin-6 (Il6ra) and interleukin-1β (Il1r1). Read counts for strongly repressed genes in key inflammatory pathways are provided in Suppl. Fig. 3. Categorical analysis of upregulated transcripts is shown in Suppl. Fig. 4. where KEGG analysis reveals enrichment of Rap1 signaling. This small GTPase regulates proliferation and migration of VSMCs[41]. Collectively, the RNAseq data indicate that loss of LRRC8A induces an anti-inflammatory transcriptional profile that is coupled to significant alterations in metabolism and stress-related processes. Accordingly, an analysis of gene-disease association for the downregulated gene set using the DisGeNET database revealed a link to vascular inflammation associated with atherosclerosis and myocardial infarction (Suppl. Fig. 2E).

Hif-1α signaling was the most significantly altered pathway so this was directly assessed using a hypoxia response element (HRE)-luciferase reporter. Activity was ~3X higher in resting WT cells. Hypoxia (1% O_2, 24 h) and chemical induction with $CoCl_2$ induced a similar fold

change activation in all cells, but WT activity remained significantly greater than KO under all conditions (Fig. 2C). Analysis of the RNA-seq data for a panel of canonical Hif-1α target genes revealed consistent downregulation in KO cells (Fig. 2D). An exception was Rps4l, a known target of HIF-1α-mediated repression, with a read count that was elevated over 60-fold in KO VSMCs (Fig. 2D). This gene has both long-noncoding and peptide-encoding functions, both of which impair S6 kinase activity and reduce VSMC proliferation and VSMC Rps4l expression is protective in pulmonary hypertension[42,43]. To further explore the regulation of Rps4l qPCR was performed under control conditions and after exposure to hypoxia or $CoCl_2$. Expression decreased appropriately but remained markedly higher in KO cells under all conditions (Fig. 2E). Many of the genes that are regulated by Hif-1α control glycolysis which is activated under hypoxic conditions. We therefore measured expression of two critical glycolytic genes; hexokinase 2 (Hk2) and phosphofructokinase (Pfkp) (Suppl. Fig. 5). Hk2 mRNA was reduced under control conditions in KO cells, but Pfkp was not. Both genes displayed impaired transcriptional responses to hypoxia in KO cells.

Metabolism, Proliferation and Migration

Since transcriptomic analysis suggested downregulation of glycolysis in KO cells, we assessed the abundance of Hk2, pyruvate kinase M2 (Pkm), and 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase-3 (Pfkfb3) and found significant decreases in all three in KO cells. The latter is a master regulator of glycolytic flux in VSMCs and is upregulated in intimal hyperplasia[44] (Fig. 3A). Therefore, we assessed the metabolic activity of WT and KO VSMCs using an XF extracellular flux analyzer. Extracellular acidification rate (ECAR) revealed a significantly lower glucose-stimulated glycolytic rate in KO cells but no difference in maximal glycolytic capacity or glycolytic reserve (Fig. 3B, left panel; Suppl. Fig. 6, left panel).

Mitochondrial oxidative phosphorylation (OCR) was also suppressed in KO cells. Significant reductions were observed in all major parameters, including basal respiration, ATP-linked respiration, maximal respiratory capacity and non-mitochondrial respiration (Fig. 3B, right panel; Suppl. Fig. 6, right panel). This was not associated with a change in proton leak, indicating that ATP production was not differentially uncoupled. OCR was previously reported to be reduced in skeletal muscle myotubes lacking LRRC8A[45]. We explored whether reduced OCR was associated with a change in mitochondrial membrane potential ($\Delta\Psi$m) as assessed by TMRE staining (Fig. 3C, left panel). This was significantly reduced in KO cells, indicating a reduced proton gradient. VDAC1 abundance correlates with mitochondrial mass and was not altered in KO cells (Fig. 3C, bottom panel). To discern if a reduced $\Delta\Psi$m reflects mitochondrial dysfunction or lower energy demand we assessed ATP and availability of reducing equivalents as reflected by ADP/ATP and $NADP^+$/NADPH respectively, and both were unaltered (Fig. 3C, middle and right panel). Maintenance of energy homeostasis in the setting of lower energy production points to lower demand. Proliferation increases this demand and a proliferative VSMC phenotype is associated with upregulation of glycolysis. MTT assays document a decrease in proliferation that was confirmed by western blotting for Proliferating Cell Nuclear Antigen (PCNA) which was also significantly reduced in KO cells (Fig. 3D). Movement is another important energy demand, and the migratory rate of KO cells was reduced both at rest and following TNFα stimulation in a scratch assay (Fig. 3E). Collectively, these results indicate that loss of LRRC8A in VSMCs is associated with a reduced metabolic rate.

Epidermal Growth Factor Signaling

Protein-Protein Interaction (PPI) hub analysis of the RNAseq data identified significantly enriched hubs as visualized by heatmap clustering (Figure 4A). The top hits were two receptor

tyrosine kinases, the epidermal growth factor receptor (EGFR) and the type 1 insulin-like growth factor receptor (IGF1R), that both share critical functional features with the insulin receptor. Differentially expressed genes associated with the EGFR hub included key adaptor and regulatory proteins (e.g., Grb10, Sh3kbp1, Ptpre, Socs3), multiple ligands and growth factors (e.g., Hbegf, Ereg, Hgf, Vegfc), cell adhesion molecules (e.g., Icam1, Ncam1, Tnc) and downstream signaling effectors (Fig. 4A). Guided by this finding we explored the functional status of EGFR signaling. Under resting conditions, the abundance of EGFR protein was unchanged but phosphorylation at Y1068 (pEGFR), an autophosphorylation site reflecting activation, was significantly reduced in KO cells (Fig. 4B). Insulin, EGFR and IGF1R all activate the PI3K/AKT cascade to promote glycolysis, cell growth and survival, and to inhibit lipolysis (the second most altered pathway based on RNAseq (Fig. 2A). EGFR activates PI3 kinase to create phosphinositol (3,4,5)-triphosphate (PIP3) which coordinates PIP3-Dependent Kinase 1 (PDK1), Mechanistic Target of Rapamycin Complex 2 (mTORC2), and AKT at the plasma membrane. PDK1 and mTORC2 phosphorylate AKT at T308 and S473 respectively to cause activation. There was a significant increase in total AKT protein in KO cells and pAKT was reduced at both sites. Exposure to EGF (50ng/ml) increased pEGFR but the difference between WT and KO only remained statistically significant at T308 pointing to greater impact on PDK1 (Fig. 4C). In view of the known LRRC8A-dependence of AKT activation by insulin we also examined the abundance of Insulin Receptor Substrate 1 (IRS1), the primary docking protein for the insulin receptor, EGFR and IGF1R. Western blotting confirmed what was suggested by RNAseq, a significant increase in IRS1 protein in KO cells (Fig. 4D).

Inflammation and Senescence

RNAseq pointed to altered TNFα/NF-κB signaling, as we reported previously following siRNA knockdown of LRRC8A[16,17,29]. We used an NF-κB-luciferase reporter to confirm that both basal and TNFα-stimulated NF-κB activity were reduced >10-fold in KO compared to WT VSMCs (Fig. 5A). However, KO cells remained responsive to TNFα based on fold change from a much lower baseline.

Reduced oxidative stress, a low metabolic rate, and less inflammation should have significant phenotypic consequences. We next assessed mRNA expression for known markers of contractile vs. secretory VSMCs (Suppl. Table 1). There was a significant reduction in only 3 of 23 markers of a contractile phenotype (Itga1, Myh11, Lmod1), and no consistent upregulation of synthetic markers (Col1a1 up, Ereg down). There were no indications of phenotypic transformation (fibroblast, fat, bone, etc.), nor changes in autophagy or apoptosis. However, there was decreased expression of 38% (15/39) of established markers of senescence including cell cycle inhibitors (Cdkn1a, Cdkn2b) and the lysosomal enzyme β-galactosidase (β-Gal, encoded by Glb1). Western blot analysis showed reduced β-Gal protein, which correlated with lower SA-β-galactosidase activity in KO cells (Fig. 5B and C). Sustained stress drives senescence partly through telomere attrition. We therefore measured relative telomere length by qPCR. LRRC8A KO VSMCs had significantly longer telomeres than WT (Fig. 5D), suggestive of enhanced genomic stability and greater replicative capacity.

Signaling changes in cultured VSMCs are summarized in Fig. 5E. Loss of LRRC8A impairs redox-dependent activation of the EGFR, reducing PIP3 production and AKT activation. This impairs Hif-1α and NF-κB-driven inflammation and reduces senescence. This anti-inflammatory and anti-senescent phenotype in cultured cells provided the impetus to assess the impact of VSMC-specific LRRC8A KO in hypercholesterolemia *in vivo*.

Atherosclerosis

LRRC8A floxed mice were bred into an ApoE[-/-], VSMC-specific Cre (SM-22α) background to create hypercholesterolemic, VSMC-specific LRRC8A knockout (KO) mice. Serum cholesterol and triglycerides did not differ between WT, HET (LRRC8A[+/-]), and KO mice (Suppl. Fig. 7A and B). After 15 weeks on an high fat diet Oil Red-O staining was performed on aortae to visualize atherosclerotic plaque coverage. LRRC8A KO and HET aortae from both male and female mice showed a significant reduction in lesion area compared to WT (Fig. 6A), HET lesion area was not significantly different from KO.

LRRC8A protein was dramatically reduced in aortae from KO mice but remained detectable due to protein from non-VSMC cells in the vascular wall. LRRC8A in aortae from male HET mice was intermediate between WT and KO, significantly lower than WT. A similar pattern was observed for markers of vascular inflammation (ICAM, VCAM), senescence (β-Gal), and proliferation (PCNA) but these changes were only statistically significant in KO vessels (Fig. 6B).

Uptake of oxidized LDL (OxLDL) occurs through scavenger receptors including LOX-1 and CD36 and both pathways trigger ROS production by NADPH oxidases[46,47]. The abundance of LOX-1 and CD36 are increased by their own pro-inflammatory activity[45,46] but did not differ between WT and KO aortae, while the expression of LDLR was increased in KO aortae from male mice. To assess the function of these receptors we compared uptake of Dil-labelled OxLDL (10 µg/ml, 6 hrs) using confocal microscopy. Uptake was reduced in KO compared to WT cells (Fig. 6C). To test the role of VRAC channel function, the impact of three non-structurally related VRAC inhibitors (Carbenoxolone; CBX, DCPIB, Tamoxifen; TAM) on OxLDL uptake by was

assessed in WT cells. All three attenuated uptake similarly to that of KO cells. In contrast, uptake of non-oxidized LDL occurs via the LDL receptor (LDLR) which does not activate ROS production. Accumulation of Dil-labeled LDL did not differ between WT and KO cells (Fig. 6C).

Vascular function

Atherosclerosis enhances vasoconstriction[48] and impairs both endothelium-dependent and -independent relaxation[49]. Mesenteric arterial vasomotor function in ApoE$^{-/-}$ mice was assessed via wire myography. In LRRC8A WT male and female animals, maximal contraction in response to both norepinephrine (NE, mixed α1-, α2- and β2-adrenergic agonist) and phenylephrine (selective α1) were enhanced compared to HET or KO mice (Fig. 7A). Endothelium-dependent relaxation to acetylcholine (ACh) was better in female but not male LRRC8A KO and HET rings compared to WT. Endothelium-independent relaxation to sodium nitroprusside (SNP) was significantly increased only in male LRRC8A KO rings.

Mesenteric artery injury induced by a prolonged (2 day) *in vitro* exposure to TNFα was mitigated by loss of LRRC8A and enhanced relaxation was shown to be associated with reduced RhoA and Rho kinase (ROCK) activation[21]. ROCK phosphorylates myosin phosphatase targeting subunit 1 (MYPT1), the regulatory subunit of myosin light chain phosphatase (MLCP) causing inhibition which promotes vasoconstriction and impairs vasorelaxation. CPI17 can be activated by protein kinase C or ROCK and also inhibits MLCP[50,51]. Western blot analysis of mesenteric artery beds from LRRC8A WT and KO, ApoE$^{-/-}$ mice demonstrated that the abundance of ROCK was not altered, but MYPT1 expression was significantly decreased in male

KO mesenteries. CPI17 protein expression was reduced in KO mice of both sexes compared to WT, while expression of MLC was not altered. Downregulation of MYPT1 and CPI17 in KO vessels may contribute to decreased vasoconstriction and enhanced relaxation. Thus, both plaque progression and vascular dysfunction in ApoE$^{-/-}$ mice is LRRC8A-dependent.

DISCUSSION

LRRC8 VRACs support Nox1 activity[16,17]. We now show that the functional status of LRRC8A affects oxidation of cytoplasmic DHE by extracellular $O_2^{\cdot-}$, consistent with our prior proposal that these anion channels support influx of $O_2^{\cdot-}$. This focused experimentation on endpoints linked to redox signaling/stress and this was further guided by changes in gene expression in KO cells. We observed an increase in GSH/GSSG, and reduced mitochondrial $O_2^{\cdot-}$ production, lipid peroxidation, Nrf2 activity and expression of antioxidant enzymes. Gene expression reflected decreased inflammatory signaling (NF-κB, Hif1α) and reduced glycolysis. KO cells produced less ATP via both ECAR and OCR and mitochondrial membrane potential was reduced, but neither ATP nor NADPH availability were altered. These changes, in the setting of lower EGFR and AKT activity, reductions in proliferation, motility and senescence, and longer telomeres collectively point to low energy demand in KO VSMCs. We hypothesized that these changes would alter vascular inflammation and oxidative stress in response to hypercholesterolemia. Accordingly, VSMC-specific LRRC8A KO mice were protected from both atheroma formation and increased contractility in an $ApoE^{-/-}$ environment. Oxidized but not reduced LDL uptake was impaired in KO cells, and LRRC8A channel blockers impaired OxLDL uptake in WT cells to a similar degree as seen in LRRC8A KO cells. These data suggest for the first time that LRRC8A channels and ROS support endocytosis of OxLDL.

Superoxide flux and oxidative stress

The ability of extracellular SOD3 to modulate Nox1-dependent activation of intracellular ASK1[19], and LRRC8A-dependence of MPRIP oxidation[21] suggested that extracellular $O_2^{\cdot-}$ might reach the cytoplasm via these channels[52]. We proposed that influx is promoted by the local

membrane depolarization produced by electron current flow through physically-associated Nox1 protein[19,29]. Here we sought to directly link $O_2^{\cdot-}$ flux to VRAC activity. Unfortunately, influx of $O_2^{\cdot-}$ produced by Nox1 activation cannot be compared because loss or inhibition of VRACs reduces Nox1 activity[17], and Nox1 activation also increases mitochondrial $O_2^{\cdot-}$ production[53,54]. Therefore, we utilized a controlled enzymatic source of extracellular $O_2^{\cdot-}$. Unstimulated WT HEK293 cells took up more $O_2^{\cdot-}$ than LRRC8A KO HEK293 cells, even though endogenous VRAC activity is low in this setting. We therefore heterologously expressed LRRC8 homomers of known, variable activity[16]. Constitutively active LRRC8C homomers increased the oxidation of cytoplasmic DHE in $O_2^{\cdot-}$-exposed cells significantly more than did expression of inactive LRRC8A, or active LRRC8D homomers. This supports the concept that Nox1-dependent redox signaling is best supported by LRRC8A/C channels[16] and is consistent with 8C best supporting small anion flux.

Within the Nox1 complex, the abundance of the integral membrane components Nox1 and p22phox were unaltered in LRRC8A KO cells, but cytoplasmic Noxa1 was reduced. This may reflect reduced inflammation as Noxa1 expression is induced by TNFα and increases in ApoE$^{-/-}$ vessels[55]. Nox1-dependent $O_2^{\cdot-}$ can also amplify ROS production by mitochondria via so-called "ROS-induced ROS production"[56], the mechanism for which is poorly understood. Thus, reduced Nox1 activity and impaired $O_2^{\cdot-}$ influx may limit mitochondrial $O_2^{\cdot-}$ production (Fig. 3C). The combination of these changes is consistent with less membrane lipid peroxidation, an increased GHS/GSSG ratio, reduced Nrf2 activity, and altered expression of SOD3 and HO-1.

Gene Expression

The dominant categories of altered gene expression were metabolism and inflammation, and the most downregulated pathway was Hif1-α which senses hypoxia and drives a metabolic

switch to glycolysis. The inflammation associated with atherosclerosis[57] and pulmonary

hypertension[58] activates Hif1-α and drives aerobic glycolysis (Warburg effect). Hif1-α can also

be activated by growth factors and cytokines via PI3K/AKT/mTOR and Ras/MAPK signaling,

both of which are triggered by EGFR activation, and impaired by loss of LRRC8A. Hif-1α

activity can be modified by multiple factors[59] including FOXO3a[60] which interferes with its

transcriptional activity. However, FOXO3a activity was impaired in LRRC8A KO VSMCs[61]

making this unlikely to account for reduced Hif1-α activity. RPS4l is a long non-coding RNA

that also encodes a peptide, and both mechanisms interfere with RPS6 function to reduce

proliferation. RPS4l is repressed by Hif1-α and consistent with this was over-expressed >20X in

KO cells. This may be of physiological significance as knockdown of RPSl4 in VSMCs

promoted pulmonary vascular inflammation and pulmonary hypertension, while overexpression

was vasculoprotective[42,43].

Glutathione provides electrons to reduce oxidized proteins, and under oxidative stress

regeneration of reduced glutathione (GSH) requires significant energy as NADPH created via the

pentose phosphate shunt[62]. Thus, lowers redox stress reduce energy demand. Lower expression

of GPX3 (Suppl. Fig. 3), a secreted enzyme that acts extracellularly, suggests that extracellular

oxidant stress is also lower in KO VSMCs. Rap1 signaling was the top pathway identified by

analysis of upregulated genes (Suppl. Fig. 4). This small GTPase promotes proliferation and

migration of VSMCs[41] but mice lacking Rap1b were hypertensive, with increased vascular

responsiveness to multiple vasoconstrictors[63]. Rap1 regulates ERK signaling[41] which is impaired

by SOD3 and by siLRRC8A[17], therefore increases in gene expression related to Rap1 signaling

may be compensatory. Based on downregulated genes, the DisGenNET gene-disease association

network identified coronary artery disease, myocardial infarction and atherosclerosis (Suppl. Fig 2).

Metabolism, proliferation and migration

An important clue to the metabolic state of LRRC8A KO VSMCs is reduced expression of Glut1 (Slc2a1, 2.3X, $p < 0.0001$, Suppl. Fig.3) which mediates insulin-independent glucose uptake. Insulin-sensitive Glut4 was not detected. Consistently, key glycolytic enzymes (Hk2, Pfkfb3 and Pkm) and metabolic activity were reduced in KO cells. In many cases, changes in ECAR are compensated by increased OCR but this was not the case in LRRC8A KO cells. Energy availability is the primary regulator of OCR. Based on an unaltered ADP/ATP and mitochondrial mass despite a reduced membrane potential, lower OCR is unlikely to reflect mitochondrial dysfunction. Low energy production in KO cells is more likely related to reduced energy requirements for reduction of oxidized proteins and lipids, and to support inflammation, proliferation and migration.

Lipolysis was the second most significantly altered pathway identified by KEGG analysis (Fig. 2A). Lipases are activated by cAMP to release free fatty acids (FFAs) from stored triglycerides to provide substrate for beta-oxidation. AKT activation by insulin blocks this process by phosphorylating and activating phosphodiesterase 3B (Pde3b, 11.3X increased, $p < 0.01$) which degrades cAMP. Expression of genes that promote cAMP production such as Ptgs2 (COX2) and Adcy8 (Adenylate cyclase 8) were reduced, but so was Ptger3, the prostaglandin E2 receptor which inhibits adenylate cyclase. Because FFAs are potent pro-inflammatory agents and promote atherosclerosis[64] additional investigation into the status of lipolysis and fatty acid oxidation in LRRC8A KO cells is warranted.

Signaling

AKT activation by insulin is impaired in LRRC8A KO adipocytes[65] and skeletal muscle cells[45], and PIP3 kinase and AKT activation by angiotensin II is disrupted in KO VSMCs[66]. Our PPI analysis identified EGFR as the top protein association hub. The insulin receptor and EGFR tyrosine kinases share key associated proteins (e.g., Irs1, Grb2), signaling pathways (PI3K/Akt and Ras/Erk)[67], and Nox-dependance[68]. LRRC8A also associates with Grb2[69], Nox1[17], and Nox2[70] and oxidants from Nox1 promote phosphorylation and transactivation of the EGFR[71]. EGFR phosphorylation at Y1068 reflects receptor activation and was reduced in KO cells both at rest and in response to EGF. Activated EGFR associates with IRS1 which activates PIP3 kinase, producing the PIP3 that drives membrane association of AKT, mTORC2 and PDK1, promoting activating AKT activation by phosphorylation at S473 and T308. Reduced phosphorylation at these sites is therefore consistent with reduced PIP3 production due to failed EGRF activation in the setting of reduced ROS.

Inflammation and Senescence

The RNAseq data supported our prior observations of suppressed NF-κB activation by TNFα in cells treated with siRNA targeting VRACs[16,17]. This was confirmed using a luciferase reporter (Fig. 3C and 5A). Reduced NF-κB and Hif1-α activity is consistent with less inflammation, and delayed senescence - a state characterized by growth arrest and the senescence-associated secretory phenotype (SASP)[72]. It occurs with aging and telomere shortening[31] (replicative senescence), and with oxidant stress[32,72,73] (premature senescence) and is associated with hypertension, atherosclerosis, and cardiovascular risk[74]. SASP genes were consistently under-expressed in KO VSMCs (Suppl. Table 1). β-gal, a classic marker of

senescence, had reduced mRNA (Glb1, -1.72, p = 0.0008), protein expression (Fig. 5B), and activity (Fig. 5C) in KO cells. Reduced inflammation may explain the longer telomeres of KO VSMCs. Consistent with these *in vitro* findings, VSMC-specific LRRC8A KO mice exposed to AngII infusion were protected from senescence as reflected by reduced aortic βGal[75]. This was again seen in the setting of atherosclerosis (Fig. 6C). Surprisingly, selective KO of LRRC8A in endothelial cells was associated with premature senescence and vascular aging[76]. It will be critical to determine how LRRC8A-dependent events differ in endothelial cells, and whether these endothelial changes are linked to altered redox signaling.

Atherosclerosis

The contribution of VSMCs to atheroma formation requires VSMC phenotypic switching from contractile to secretory and migration to the intima with subsequent transformation into LDL-accumulating foam cells[28]. Disease progression is Nox1-dependent[77] and characterized by oxidative stress, and increased glycolysis, proliferation, migration, and senescence[74]. OxLDL promotes atherogenesis by activating LOX-1 and CD36 scavenger receptors[47,78] that can trigger ROS production by both Nox1 and mitochondria, activate NF-κB, and enhance their own expression[34,79] creating a maladaptive positive feedback loop. Both KO, and very importantly LRRC8A HET mice, were resistant to hypercholesterolemia-induced aortic atheroma formation. Expression of LRRC8A was correlated with the abundance of markers of inflammation including ICAM and VCAM. LRRC8A mRNA expression was decreased in atherosclerotic aortae from humans[16]. In the same dataset, cytokine-cytokine receptor interactions was the KEGG pathway most associated with the downregulated genes (p < 5.8 x 10^{-7}), while OCR and ECAR-related genes (Metabolic Pathways p = 1.5 x 10^{-43}) accounted for ~12% of all of those that were

upregulated[80]. Thus, LRRC8A downregulation may represent a critical compensatory change by which VSMCs try to maintain function in a cytokine-rich, high energy demand environment.

LOX-1 and CD36 abundance were not altered in aortae from KO mice, but uptake of OxLDL was significantly reduced in cultured KO VSMCs. Thus, resistance to atherosclerosis in KO mice is likely to be related to impaired function of these receptors, as well as to other Nox1-dependent inflammatory signals that drive the process. A connection to LRRC8A was supported by the finding that three structurally unrelated VRAC inhibitors reduced OxLDL uptake in WT VSMC to a degree similar to that seen in KO cells. This demonstrates that LRRC8A function, and potentially the ROS produced in response to LOX-1 or CD36 activation, is required for receptor endocytosis and OxLDL uptake. Importantly, LRRC8A KO did not alter uptake of non-oxidized LDL by the LDLR, which does not activate Nox1.

We previously assessed vascular reactivity in mesenteric arteries from VSMC-specific LRRC8A KO mice[21,35]. In freshly isolated vessels contractile responses to KCl and PE were unaltered while relaxation to both ACh and SNP were augmented. In the pro-inflammatory setting of hypercholesterolemia, contractility to PE and NE were increased in WT but not in KO or HET vessels. Protected dilation to ACh was again seen in female but not in male mice. This is in accordance with prior demonstration of impaired responses to ACh only in female ApoE$^{-/-}$ mice[81,82]. We attributed vasomotor changes in VSMC-specific LRRC8A KO mice to reduced RhoA activity and hypothesized that this was due to the loss of activating effects of $O_2^{\cdot-}$ on the small GTPase positioned in proximity to oxidants in a multi-protein complex organized by MPRIP. Here we assessed expression of key proteins in this complex and found reduced

expression of MYPT1 and CPI17, both of which are capable of impairing activity of MLCP and increasing calcium sensitivity.

Summary

Impaired Akt activation by insulin has been well documented in LRRC8A KO cells[65]. We now report similar defects in EGFR signaling, suggesting a shared defect in protein tyrosine kinase signaling, perhaps attributable to loss of oxidant influx. Thus, reduced AKT activity is most likely related to impaired PI3K activation[66] which reduces PDK1 and mTORC2 activation of AKT. The integrated downstream effects of these changes include impaired Hif1- α and NF-κB activation, reduced inflammation, proliferation migration all of which limit energy demand, reducing both ECAR and OCR.

These signaling defects are protective of VSMCs in the pro-inflammatory environment of hypercholesterolemia. Agonist-triggered cytoplasmic oxidant signals may be required for endocytosis of the LOX-1 and CD36 scavenger receptors as they are for TNFR1[17]. LRRC8A-dependance of OxLDL uptake is a key novel finding that will require detailed investigation to determine how VRACs and ROS contribute to this process. This also provides an exciting potential target for selective inhibition of ROS-dependent pathophysiology.

ACKNOWLEDGMENTS

We thank Benjamin Crawford for their expert assistance with flow cytometry data acquisition.

Sources of funding

This work was supported by NIH GM138191 to RJS and HL160975 and DK132948 to FSL.

FOOTNOTE

Nonstandard Abbreviations and Acronyms

ACh	Acetylcholine
ApoE	Apolipoprotein E
ApoE$^{-/-}$	Apolipoprotein E–knockout
ASK1	Apoptosis signal–regulating kinase 1
β-Gal	B-galactosidase
CBX	Carbenoxolone
CPI-17	PKC-potentiated inhibitor protein of PP1 (17 kda)
DHE	Dihydroethidium
ECAR	Extracellular acidification rate
EGFR	Epidermal growth factor receptor
FACS	Fluorescence-activated cell sorting
HCT116	Human colorectal carcinoma cell line HCT-116
HET	Heterozygous
HIF-1α	Hypoxia-inducible factor-1α
HK2	Hexokinase 2
HO-1	Heme oxygenase-1
HRE	Hypoxia response element
ICAM	Intercellular adhesion molecule-1
IGF1R	Insulin-like growth factor-1 receptor
IRS1	Insulin receptor substrate-1
KO	Knockout
LDL	Low-density lipoprotein
LDLR	Low-density lipoprotein receptor
LOX-1	Lectin-like oxidized LDL receptor-1
LRRC8A/C/D	Leucine-rich repeat–containing 8A/8C/8D
MDA	Malondialdehyde
MLC	Myosin light chain
MLCP	Myosin light chain phosphatase
MYPT1	Myosin phosphatase target subunit-1

NE	Norepinephrine
Nox1	NADPH oxidase 1
Nrf2	Nuclear factor erythroid 2–related factor 2
OCR	Oxygen consumption rate
OxLDL	Oxidized low-density lipoprotein
PCNA	Proliferating cell nuclear antigen
Pdk1	3-phosphoinositide-dependent protein kinase-1
Pfkfb3	6-phosphofructo-2-kinase/fructose-2
PIP$_3$	Phosphatidylinositol 3,4,5-trisphosphate
PKM	Pyruvate kinase M
PPI	Protein–protein interaction
Rap1	Ras-associated protein 1
RNAseq	RNA sequencing
ROS	Reactive oxygen species
ROSstar550	Cell-permeant fluorescent ROS probe
Rps4l	Ribosomal protein S4-like
ROCK	Rho-associated coiled-coil containing protein kinase
SA-β-gal	Senescence-associated β-galactosidase
SASP	Senescence-associated secretory phenotype
SNP	Sodium nitroprusside
SOD3	Extracellular superoxide dismutase
TAM	Tamoxifen
TMRE	Tetramethylrhodamine ethyl ester
TNFα	Tumor necrosis factor-α
VDAC1	Voltage-dependent anion channel 1
VCAM	Vascular cell adhesion molecule-1
VRAC	Volume-regulated anion channel
VSMC	Vascular smooth muscle cell
WT	Wild type
ΔΨm	Mitochondrial membrane potential

CONFLICT OF INTEREST

None declared.

REFERENCES

1. Voss FK, Ullrich F, Munch J, Lazarow K, Lutter D, Mah N, Andrade-Navarro MA, von Kries JP, Stauber T, Jentsch TJ. Identification of LRRC8 heteromers as an essential component of the volume-regulated anion channel VRAC. *Science*. 2014;344:634–638. doi: 10.1126/science.1252826

2. Chen L, Konig B, Liu T, Pervaiz S, Razzaque YS, Stauber T. More than just a pressure release valve: physiological roles of volume-regulated LRRC8 anion channels. *Biol Chem*. 2019. doi: 10.1515/hsz-2019-0189

3. Syeda R, Qiu Z, Dubin AE, Murthy SE, Florendo MN, Mason DE, Mathur J, Cahalan SM, Peters EC, Montal M, et al. LRRC8 Proteins Form Volume-Regulated Anion Channels that Sense Ionic Strength. *Cell*. 2016;164:499–511. doi: 10.1016/j.cell.2015.12.031

4. Okada Y. Physiology of the volume-sensitive/regulatory anion channel VSOR/VRAC. Part 1: from its discovery and phenotype characterization to the molecular entity identification. *J Physiol Sci*. 2024;74:3. doi: 10.1186/s12576-023-00897-x

5. Okada Y. Physiology of the volume-sensitive/regulatory anion channel VSOR/VRAC: part 2: its activation mechanisms and essential roles in organic signal release. *J Physiol Sci*. 2024;74:34. doi: 10.1186/s12576-024-00926-3

6. Ren Z, Raucci FJ, Jr., Browe DM, Baumgarten CM. Regulation of swelling-activated Cl(-) current by angiotensin II signalling and NADPH oxidase in rabbit ventricle. *Cardiovasc Res*. 2008;77:73–80.

7. Varela D, Simon F, Riveros A, Jorgensen F, Stutzin A. NAD(P)H oxidase-derived H(2)O(2) signals chloride channel activation in cell volume regulation and cell proliferation. *J Biol Chem*. 2004;279:13301–13304.

8. Liang W, Huang L, Zhao D, He JZ, Sharma P, Liu J, Gramolini AO, Ward ME, Cho HC, Backx PH. Swelling-activated Cl- currents and intracellular CLC-3 are involved in proliferation of human pulmonary artery smooth muscle cells. *Journal of hypertension*. 2014;32:318–330. doi: 10.1097/HJH.0000000000000013

9. Qian JS, Pang RP, Zhu KS, Liu DY, Li ZR, Deng CY, Wang SM. Static pressure promotes rat aortic smooth muscle cell proliferation via upregulation of volume-regulated chloride channel. *Cell Physiol Biochem*. 2009;24:461–470.

10. Wang L, Chen L, Zhu L, Rawle M, Nie S, Zhang J, Ping Z, Kangrong C, Jacob TJ. Regulatory volume decrease is actively modulated during the cell cycle. *J Cell Physiol*. 2002;193:110–119.

11. Chen L, Wang L, Zhu L, Nie S, Zhang J, Zhong P, Cai B, Luo H, Jacob TJ. Cell cycle-dependent expression of volume-activated chloride currents in nasopharyngeal carcinoma cells. *Am J Physiol Cell Physiol*. 2002;283:C1313–1323.

12. Yin Z, Tong Y, Zhu H, Watsky MA. ClC-3 is required for LPA-activated Cl- current activity and fibroblast-to-myofibroblast differentiation. *Am J Physiol Cell Physiol*. 2008;294:C535–542.

13. Matsuda JJ, Filali MS, Volk KA, Lamb FS. Activation of IClswell by TNFalpha requires ClC-3-dependent ROS production. *FASEB J*. 2008;22:937.918.

14. Yang H, Huang LY, Zeng DY, Huang EW, Liang SJ, Tang YB, Su YX, Tao J, Shang F, Wu QQ, et al. Decrease of intracellular chloride concentration promotes endothelial cell inflammation by activating nuclear factor-kappaB pathway. *Hypertension*. 2012;60:1287–1293. doi: 10.1161/HYPERTENSIONAHA.112.198648

15. Selzman CH, Shames BD, McIntyre RC, Jr., Banerjee A, Harken AH. The NFkappaB inhibitory peptide, IkappaBalpha, prevents human vascular smooth muscle proliferation. *Ann Thorac Surg*. 1999;67:1227–1231; discussion 1231–1222.

16.   Choi H, Rohrbough JC, Nguyen HN, Dikalova A, Lamb FS. Oxidant-resistant LRRC8A/C anion channels support superoxide production by NADPH oxidase 1. *J Physiol*. 2021;599:3013–3036. doi: 10.1113/JP281577

17.   Choi H, Ettinger N, Rohrbough J, Dikalova A, Nguyen HN, Lamb FS. LRRC8A channels support TNFalpha-induced superoxide production by Nox1 which is required for receptor endocytosis. *Free Radic Biol Med*. 2016;101:413–423. doi: 10.1016/j.freeradbiomed.2016.11.003

18.   Friard J, Laurain A, Rubera I, Duranton C. LRRC8/VRAC Channels and the Redox Balance: A Complex Relationship. *Cell Physiol Biochem*. 2021;55:106–118. doi: 10.33594/000000341

19.   Lamb FS, Choi H, Miller MR, Stark RJ. Vascular Inflammation and Smooth Muscle Contractility: The Role of Nox1-Derived Superoxide and LRRC8 Anion Channels. *Hypertension*. 2024. doi: 10.1161/HYPERTENSIONAHA.123.19434

20.   Bertelli S, Zuccolini P, Gavazzo P, Pusch M. Molecular determinants underlying volume-regulated anion channel subunit-dependent oxidation sensitivity. *J Physiol*. 2022;600:3965–3982. doi: 10.1113/JP283321

21.   Choi H, Miller MR, Nguyen HN, Rohrbough JC, Koch SR, Boatwright N, Yarboro MT, Sah R, McDonald WH, Reese JJ, et al. LRRC8A anion channels modulate vascular reactivity via association with myosin phosphatase rho interacting protein. *FASEB J*. 2023;37:e23028. doi: 10.1096/fj.202300561R

22.   Harris K, Won SJ, Uruk G, Mai N, Ogut D, Zhao Y, Xie L, Baxter P, Everaerts K, Sah R, et al. Excitotoxic neuronal death requires superoxide entry into neurons through volume-regulated anion channels. *Sci Adv*. 2025;11:eadw0424. doi: 10.1126/sciadv.adw0424

23.   Konior A, Schramm A, Czesnikiewicz-Guzik M, Guzik TJ. NADPH oxidases in vascular pathology. *Antioxid Redox Signal*. 2014;20:2794–2814. doi: 10.1089/ars.2013.5607

24.   Rivera J, Sobey CG, Walduck AK, Drummond GR. Nox isoforms in vascular pathophysiology: insights from transgenic and knockout mouse models. *Redox Rep*. 2010;15:50–63. doi: 10.1179/174329210X12650506623401

25.   Radi R. Peroxynitrite, a stealthy biological oxidant. *J Biol Chem*. 2013;288:26464–26472. doi: 10.1074/jbc.R113.472936

26.   Andres CMC, Perez de la Lastra JM, Andres Juan C, Plou FJ, Perez-Lebena E. Superoxide Anion Chemistry-Its Role at the Core of the Innate Immunity. *International journal of molecular sciences*. 2023;24. doi: 10.3390/ijms24031841

27.   Myszko M, Bychowski J, Skrzydlewska E, Luczaj W. The Dual Role of Oxidative Stress in Atherosclerosis and Coronary Artery Disease: Pathological Mechanisms and Diagnostic Potential. *Antioxidants (Basel)*. 2025;14. doi: 10.3390/antiox14030275

28.   Basatemur GL, Jorgensen HF, Clarke MCH, Bennett MR, Mallat Z. Vascular smooth muscle cells in atherosclerosis. *Nat Rev Cardiol*. 2019;16:727–744. doi: 10.1038/s41569-019-0227-9

29.   Lamb FS, Choi H, Miller MR, Stark RJ. TNFalpha and Reactive Oxygen Signaling in Vascular Smooth Muscle Cells in Hypertension and Atherosclerosis. *American journal of hypertension*. 2020. doi: 10.1093/ajh/hpaa089

30.   Siti HN, Kamisah Y, Kamsiah J. The role of oxidative stress, antioxidants and vascular inflammation in cardiovascular disease (a review). *Vascul Pharmacol*. 2015;71:40–56. doi: 10.1016/j.vph.2015.03.005

31.   Uryga AK, Grootaert MOJ, Garrido AM, Oc S, Foote K, Chappell J, Finigan A, Rossiello F, d'Adda di Fagagna F, Aravani D, et al. Telomere damage promotes vascular smooth muscle cell senescence and immune cell recruitment after vessel injury. *Commun Biol*. 2021;4:611. doi: 10.1038/s42003-021-02123-z

32.   Matthews C, Gorenne I, Scott S, Figg N, Kirkpatrick P, Ritchie A, Goddard M, Bennett M. Vascular smooth muscle cells undergo telomere-based senescence in human atherosclerosis: effects of

telomerase and oxidative stress. *Circ Res*. 2006;99:156–164. doi: 10.1161/01.RES.0000233315.38086.bc

33.   Shi J, Yang Y, Cheng A, Xu G, He F. Metabolism of vascular smooth muscle cells in vascular diseases. *Am J Physiol Heart Circ Physiol*. 2020;319:H613–H631. doi: 10.1152/ajpheart.00220.2020

34.   Zingg JM, Vlad A, Ricciarelli R. Oxidized LDLs as Signaling Molecules. *Antioxidants (Basel)*. 2021;10. doi: 10.3390/antiox10081184

35.   Choi H, Panja S, Nguyen HN, Stark RJ, Lamb FS. Smooth Muscle LRRC8A Knockout Preserves Vascular Function in Ang II Hypertension. *Hypertension*. 2025. doi: 10.1161/HYPERTENSIONAHA.125.25222

36.   Yamada T, Figueroa EE, Denton JS, Strange K. LRRC8A homohexameric channels poorly recapitulate VRAC regulation and pharmacology. *Am J Physiol Cell Physiol*. 2021;320:C293–C303. doi: 10.1152/ajpcell.00454.2020

37.   Gaitan-Penas H, Gradogna A, Laparra-Cuervo L, Solsona C, Fernandez-Duenas V, Barrallo-Gimeno A, Ciruela F, Lakadamyali M, Pusch M, Estevez R. Investigation of LRRC8-Mediated Volume-Regulated Anion Currents in Xenopus Oocytes. *Biophys J*. 2016;111:1429–1443. doi: 10.1016/j.bpj.2016.08.030

38.   Yamada T, Strange K. Intracellular and extracellular loops of LRRC8 are essential for volume-regulated anion channel function. *J Gen Physiol*. 2018. doi: 10.1085/jgp.201812016

39.   Dikalov S. Cross talk between mitochondria and NADPH oxidases. *Free Radic Biol Med*. 2011;51:1289–1301. doi: 10.1016/j.freeradbiomed.2011.06.033

40.   Vendrov AE, Stevenson MD, Lozhkin A, Hayami T, Holland NA, Yang X, Moss N, Pan H, Wickline SA, Stockand JD, et al. Renal NOXA1/NOX1 Signaling Regulates Epithelial Sodium Channel and Sodium Retention in Angiotensin II-induced Hypertension. *Antioxidants & Redox Signaling*. 2021;36:550–566. doi: 10.1089/ars.2021.0047

41.   Li Q, Teng Y, Wang J, Yu M, Li Y, Zheng H. Rap1 promotes proliferation and migration of vascular smooth muscle cell via the ERK pathway. *Pathol Res Pract*. 2018;214:1045–1050. doi: 10.1016/j.prp.2018.04.007

42.   Li Y, Zhang J, Sun H, Chen Y, Li W, Yu X, Zhao X, Zhang L, Yang J, Xin W, et al. lnc-Rps4l-encoded peptide RPS4XL regulates RPS6 phosphorylation and inhibits the proliferation of PASMCs caused by hypoxia. *Mol Ther*. 2021;29:1411–1424. doi: 10.1016/j.ymthe.2021.01.005

43.   Liu Y, Zhang H, Li Y, Yan L, Du W, Wang S, Zheng X, Zhang M, Zhang J, Qi J, et al. Long Noncoding RNA Rps4l Mediates the Proliferation of Hypoxic Pulmonary Artery Smooth Muscle Cells. *Hypertension*. 2020;76:1124–1133. doi: 10.1161/HYPERTENSIONAHA.120.14644

44.   Pan H, Wu Z, Gao Y, Yao W, Feng G, Wang H. The relevance of resveratrol in ameliorating carotid atherosclerosis through glycolysis. *BMC Cardiovasc Disord*. 2025;25:301. doi: 10.1186/s12872-025-04735-3

45.   Kumar A, Xie L, Ta CM, Hinton AJ, Gunasekar SK, Minerath RA, Shen K, Maurer JM, Grueter CE, Abel ED, et al. SWELL1 regulates skeletal muscle cell size, intracellular signalling, adiposity and glucose metabolism. *Elife*. 2020;9. doi: 10.7554/eLife.58941

46.   Mitra S, Goyal T, Mehta JL. Oxidized LDL, LOX-1 and atherosclerosis. *Cardiovasc Drugs Ther*. 2011;25:419–429. doi: 10.1007/s10557-011-6341-5

47.   Yang M, Silverstein RL. CD36 signaling in vascular redox stress. *Free Radic Biol Med*. 2019;136:159–171. doi: 10.1016/j.freeradbiomed.2019.02.021

48.   Leal MA, Balarini CM, Dias AT, Porto ML, Gava AL, Pereira TM, Meyrelles SS, Vasquez EC. Mechanisms of enhanced vasoconstriction in the mouse model of atherosclerosis: the beneficial effects of sildenafil. *Curr Pharm Biotechnol*. 2015;16:517–530. doi: 10.2174/1389201016061504071113458

49. Mian MO, Idris-Khodja N, Li MW, Leibowitz A, Paradis P, Rautureau Y, Schiffrin EL. Preservation of endothelium-dependent relaxation in atherosclerotic mice with endothelium-restricted endothelin-1 overexpression. *J Pharmacol Exp Ther*. 2013;347:30–37. doi: 10.1124/jpet.113.206532

50. Nunes KP, Webb RC. New insights into RhoA/Rho-kinase signaling: a key regulator of vascular contraction. *Small GTPases*. 2021;12:458–469. doi: 10.1080/21541248.2020.1822721

51. Eto M, Kitazawa T, Matsuzawa F, Aikawa S, Kirkbride JA, Isozumi N, Nishimura Y, Brautigan DL, Ohki SY. Phosphorylation-induced conformational switching of CPI-17 produces a potent myosin phosphatase inhibitor. *Structure*. 2007;15:1591–1602. doi: 10.1016/j.str.2007.10.014

52. Choi H, Stark RJ, Raja BS, Dikalova A, Lamb FS. Apoptosis signal-regulating kinase 1 activation by Nox1-derived oxidants is required for TNFalpha receptor endocytosis. *Am J Physiol Heart Circ Physiol*. 2019;316:H1528–H1537. doi: 10.1152/ajpheart.00741.2018

53. Sylvester AL, Zhang DX, Ran S, Zinkevich NS. Inhibiting NADPH Oxidases to Target Vascular and Other Pathologies: An Update on Recent Experimental and Clinical Studies. *Biomolecules*. 2022;12. doi: 10.3390/biom12060823

54. Ward JPT. From Physiological Redox Signalling to Oxidant Stress. *Adv Exp Med Biol*. 2017;967:335–342. doi: 10.1007/978-3-319-63245-2_21

55. Niu XL, Madamanchi NR, Vendrov AE, Tchivilev I, Rojas M, Madamanchi C, Brandes RP, Krause KH, Humphries J, Smith A, et al. Nox activator 1: a potential target for modulation of vascular reactive oxygen species in atherosclerotic arteries. *Circulation*. 2010;121:549–559. doi: 10.1161/CIRCULATIONAHA.109.908319

56. Daiber A, Di Lisa F, Oelze M, Kroller-Schon S, Steven S, Schulz E, Munzel T. Crosstalk of mitochondria with NADPH oxidase via reactive oxygen and nitrogen species signalling and its role for vascular function. *Br J Pharmacol*. 2017;174:1670–1689. doi: 10.1111/bph.13403

57. Jain T, Nikolopoulou EA, Xu Q, Qu A. Hypoxia inducible factor as a therapeutic target for atherosclerosis. *Pharmacol Ther*. 2018;183:22–33. doi: 10.1016/j.pharmthera.2017.09.003

58. Liu M, Galli G, Wang Y, Fan Q, Wang Z, Wang X, Xiao W. Novel Therapeutic Targets for Hypoxia-Related Cardiovascular Diseases: The Role of HIF-1. *Front Physiol*. 2020;11:774. doi: 10.3389/fphys.2020.00774

59. Koyasu S, Kobayashi M, Goto Y, Hiraoka M, Harada H. Regulatory mechanisms of hypoxia-inducible factor 1 activity: Two decades of knowledge. *Cancer science*. 2018;109:560–571. doi: 10.1111/cas.13483

60. Emerling BM, Weinberg F, Liu JL, Mak TW, Chandel NS. PTEN regulates p300-dependent hypoxia-inducible factor 1 transcriptional activity through Forkhead transcription factor 3a (FOXO3a). *Proc Natl Acad Sci U S A*. 2008;105:2622–2627. doi: 10.1073/pnas.0706790105

61. Lu FT, Huang CC, Lai WY, Yang GY, Liang ZJ, Zhang ZY, Chokshi T, Guo KM, Tang YB, Chen Y, et al. Vascular smooth muscle-specific LRRC8A knockout ameliorates angiotensin II-induced cerebrovascular remodeling by inhibiting the WNK1/FOXO3a/MMP signaling pathway. *Acta Pharmacol Sin*. 2024;45:1848–1860. doi: 10.1038/s41401-024-01280-1

62. TeSlaa T, Ralser M, Fan J, Rabinowitz JD. The pentose phosphate pathway in health and disease. *Nat Metab*. 2023;5:1275–1289. doi: 10.1038/s42255-023-00863-2

63. Lakshmikanthan S, Zieba BJ, Ge ZD, Momotani K, Zheng X, Lund H, Artamonov MV, Maas JE, Szabo A, Zhang DX, et al. Rap1b in smooth muscle and endothelium is required for maintenance of vascular tone and normal blood pressure. *Arterioscler Thromb Vasc Biol*. 2014;34:1486–1494. doi: 10.1161/ATVBAHA.114.303678

64. Khoi CS, Lin TY, Chiang CK. Targeting Insulin Resistance, Reactive Oxygen Species, Inflammation, Programmed Cell Death, ER Stress, and Mitochondrial Dysfunction for the Therapeutic

Prevention of Free Fatty Acid-Induced Vascular Endothelial Lipotoxicity. *Antioxidants (Basel)*. 2024;13. doi: 10.3390/antiox13121486

65. Zhang Y, Xie L, Gunasekar SK, Tong D, Mishra A, Gibson WJ, Wang C, Fidler T, Marthaler B, Klingelhutz A, et al. SWELL1 is a regulator of adipocyte size, insulin signalling and glucose homeostasis. *Nat Cell Biol*. 2017;19:504–517. doi: 10.1038/ncb3514

66. Lu J, Xu F, Zhang J. Inhibition of angiotensin II-induced cerebrovascular smooth muscle cell proliferation by LRRC8A downregulation through suppressing PI3K/AKT activation. *Hum Cell*. 2019;32:316–325. doi: 10.1007/s13577-019-00260-6

67. Borisov N, Aksamitiene E, Kiyatkin A, Legewie S, Berkhout J, Maiwald T, Kaimachnikov NP, Timmer J, Hoek JB, Kholodenko BN. Systems-level interactions between insulin-EGF networks amplify mitogenic signaling. *Mol Syst Biol*. 2009;5:256. doi: 10.1038/msb.2009.19

68. Henriquez-Olguin C, Boronat S, Cabello-Verrugio C, Jaimovich E, Hidalgo E, Jensen TE. The Emerging Roles of Nicotinamide Adenine Dinucleotide Phosphate Oxidase 2 in Skeletal Muscle Redox Signaling and Metabolism. *Antioxid Redox Signal*. 2019;31:1371–1410. doi: 10.1089/ars.2018.7678

69. Alghanem AF, Abello J, Maurer JM, Kumar A, Ta CM, Gunasekar SK, Fatima U, Kang C, Xie L, Adeola O, et al. The SWELL1-LRRC8 complex regulates endothelial AKT-eNOS signaling and vascular function. *Elife*. 2021;10. doi: 10.7554/eLife.61313

70. Huo C, Liu Y, Li X, Xu R, Jia X, Hou L, Wang X. LRRC8A contributes to angiotensin II-induced cardiac hypertrophy by interacting with NADPH oxidases via the C-terminal leucine-rich repeat domain. *Free Radic Biol Med*. 2021;165:191–202. doi: 10.1016/j.freeradbiomed.2021.01.022

71. Jagadeesha DK, Takapoo M, Banfi B, Bhalla RC, Miller FJ, Jr. Nox1 transactivation of epidermal growth factor receptor promotes N-cadherin shedding and smooth muscle cell migration. *Cardiovasc Res*. 2012;93:406–413. doi: 10.1093/cvr/cvr308

72. Chi C, Li DJ, Jiang YJ, Tong J, Fu H, Wu YH, Shen FM. Vascular smooth muscle cell senescence and age-related diseases: State of the art. *Biochim Biophys Acta Mol Basis Dis*. 2019;1865:1810–1821. doi: 10.1016/j.bbadis.2018.08.015

73. Herbert KE, Mistry Y, Hastings R, Poolman T, Niklason L, Williams B. Angiotensin II-mediated oxidative DNA damage accelerates cellular senescence in cultured human vascular smooth muscle cells via telomere-dependent and independent pathways. *Circ Res*. 2008;102:201–208. doi: 10.1161/CIRCRESAHA.107.158626

74. Picos A, Seoane N, Campos-Toimil M, Vina D. Vascular senescence and aging: mechanisms, clinical implications, and therapeutic prospects. *Biogerontology*. 2025;26:118. doi: 10.1007/s10522-025-10256-5

75. Choi H, Panja S, Nguyen HN, Stark RJ, Lamb FS. Smooth muscle LRRC8A knockout preserves vascular function in AngII hypertension. *bioRxiv*. 2025. doi: 10.1101/2025.05.08.652978

76. Pang J, Pan L, Shen W, Yi G, Kong X, Zhang H, Feng B, Sun H, Chen L, Li G, et al. Endothelial LRRC8A delays vascular ageing in natural and accelerated ageing mouse models. *Cardiovasc Res*. 2025. doi: 10.1093/cvr/cvaf212

77. Lee MY, San Martin A, Mehta PK, Dikalova AE, Garrido AM, Datla SR, Lyons E, Krause KH, Banfi B, Lambeth JD, et al. Mechanisms of vascular smooth muscle NADPH oxidase 1 (Nox1) contribution to injury-induced neointimal formation. *Arterioscler Thromb Vasc Biol*. 2009;29:480–487.

78. Goyal T, Mitra S, Khaidakov M, Wang X, Singla S, Ding Z, Liu S, Mehta JL. Current Concepts of the Role of Oxidized LDL Receptors in Atherosclerosis. *Curr Atheroscler Rep*. 2012. doi: 10.1007/s11883-012-0228-1

79. Mulumba M, Le C, Schelsohn E, Namkung Y, Laporte SA, Febbraio M, Servant MJ, Chemtob S, Lubell WD, Marleau S, et al. Selective Azapeptide CD36 Ligand MPE-298 Regulates oxLDL-LOX-1-

Mediated Inflammation and Mitochondrial Oxidative Stress in Macrophages. *Cells*. 2025;14. doi: 10.3390/cells14050385

80. Biros E, Gabel G, Moran CS, Schreurs C, Lindeman JH, Walker PJ, Nataatmadja M, West M, Holdt LM, Hinterseher I, et al. Differential gene expression in human abdominal aortic aneurysm and aortic occlusive disease. *Oncotarget*. 2015;6:12984–12996. doi: 10.18632/oncotarget.3848

81. Arruda RM, Peotta VA, Meyrelles SS, Vasquez EC. Evaluation of vascular function in apolipoprotein E knockout mice with angiotensin-dependent renovascular hypertension. *Hypertension*. 2005;46:932–936. doi: 10.1161/01.HYP.0000182154.61862.52

82. Cola MS, Gava AL, Meyrelles SS, Vasquez EC. Endothelial dysfunction of resistance vessels in female apolipoprotein E-deficient mice. *Lipids in Health and Disease*. 2010;9:51. doi: 10.1186/1476-511X-9-51

FIGURE LEGENDS

Figure 1. LRRC8 channels support superoxide influx and knockout reduces oxidative stress.
A, WT and LRRC8A KO HEK293 cells were loaded with dihydroethidium (DHE), washed, and exposed to extracellular $O_2^{\cdot-}$ generated by hypoxanthine/xanthine oxidase (HX–XO). 2-OH-ethidium was quantified by HPLC and normalized to protein. $O_2^{\cdot-}$ increased 2-OH-ethidium in WT but not KO cells (mean ± SEM; n = 3 independent experiments; two-way ANOVA). **B,** Protocol for experiments in (**C)** and (**D)**: GFP-tagged LRRC8 proteins were expressed in DHE-loaded HCT116 cells lacking all other LRRC8 proteins and exposed to $O_2^{\cdot-}$. **C,** Confocal LRRC8A-GFP (top) or LRRC8C-GFP (bottom); GFP tag (left), DHE (middle) and merged images (right). Cells that express LRRC8C but not LRRC8A have higher DHE fluorescence. **D,** *Left:* Flow-cytometric quantification of DHE fluorescence in LRRC8A-, LRRC8D-, or LRRC8C-GFP–expressing cells under control and $O_2^{\cdot-}$-exposed conditions (2-way ANOVA); *Right:* overlaid histograms for LRRC8C-GFP cells demonstrating increased fluorescence in cells that express LRRC8C-GFP and an additional increase in cells exposed to $O_2^{\cdot-}$. **E,** ROS imaging in WT and KO primary aortic VSMCs using ROSstar550 or MitoSOX; *Left*: confocal images of cells exposed to ROSstar550. *Right*: Quantification of ROSstar and Mitosox signals by flow cytometry. Violin plots show single-cell fluorescence (dye-positive gate, all cells from 3 independent experiments, Welch's *t* test). **F,** Lipid peroxidation (malondialdehyde adducts) at baseline and after TNFα. The difference at baseline was not significant (P = 0.0125) but TNFα increased peroxidation only in WT cells (n = 3 - 4). **G,** Reduced/oxidized glutathione (GSH/GSSG) under resting conditions; KO had a significantly more reduced ratio (n = 3). **H,** Nrf2 reporter activity under resting conditions is lower in KO VSMCs (n = 4). All bars show mean ± SEM and dots represent independent replicates. *P<0.05, and ***P<0.001.

Figure 2. Effects of LRRC8A KO on mRNA expression and Hif1α signaling in VSMCs. A, Most altered KEGG pathways (combined score) with increased (red) and decreased (green) genes noted. **B,** Categories of downregulated genes and associated pathways. **C.** Hif-1α reporter (HRE-luciferase) activity in WT and KO VSMCs under normoxia, hypoxia (1% O_2, 24 h), or $CoCl_2$ (300 μM, 6 h) stimulation; activity normalized to protein and corrected for reporter infection efficiency (GFP/tubulin). (normoxia, one-sample t test; hypoxia and $CoCl_2$, Mann–Whitney test; n = 3). **D,** RNA-seq log_2 fold change (KO vs WT) for canonical HIF-1α target genes. **E,** qPCR validation of Rps4l expression under the same conditions as in (**C**) normalized to WT in normoxia (normoxia, one-sample t test; hypoxia and $CoCl_2$, Mann–Whitney test; n = 6 for control and n = 3 for hypoxia and $CoCl_2$). Bars are mean ± SEM, dots represent replicates. *P<0.05, ***P<0.001.

Figure 3. Metabolism and energy balance in LRRC8A KO VSMCs. A, Abundance of HK2, PFKFB3, PKM, and LRRC8A in primary aortic VSMCs (WT, KO) normalized to α-tubulin (unpaired t test; $n = 4$ independent lysates per genotype). **B,** Seahorse glycolytic stress test (ECAR; injections: glucose, oligomycin, 2-deoxyglucose) and mitochondrial stress test (OCR; injections: oligomycin, FCCP, antimycin/rotenone) n = 4. **C,** Mitochondrial membrane potential comparison by TMRE fluorescence/flow cytometry (violin plots of single-cell fluorescence, dye-positive gate) (pooled cells from $n = 3$ independent experiments, unpaired t test). ADP/ATP and $NADP^+$/NADPH ($n = 3$-4, Welch's t test), and VDAC1 abundance as an estimate of mitochondrial biomass. (n = 5, unpaired t test). **D,** Proliferation assays: MTT over 72 h (n = 6, One-way ANOVA**),** and PCNA immunoblotting. (unpaired t test; $n = 4$). **E,** Migration by scratch-wound (IncuCyte) under control conditions and with TNFα (10 ng/mL); average wound width (μm). (One-way ANOVA $n = 3$ with 6 wells/condition). Bars/plots show mean ± SEM, dots are independent replicates. *P<0.05, **P<0.01, ***P<0.001).

Figure 4. EGFR-centered network and EGF–AKT signaling in LRRC8A-deficient VSMCs.
A, Top hits from Protein-Protein Interaction (PPI) hub analysis of RNAseq (KO vs WT) with associated partner proteins (bottom dot map). **B,** Immunoblotting for pT1068 and normalized to total EGFR or α-tubulin. For control lanes, n = 5, one-sample t test; for EGF-treated comparisons n = 3, unpaired t test. **C,** The same lysates were probed for pAKT (T308), pAKT (S473), or total AKT and normalized to total AKT or α-tubulin. For control n = 6, one-sample t test; EGF-treated WT vs. KO, n = 3, unpaired t test. **D,** IRS1 protein normalized to α-tubulin. (n = 6, unpaired t test) Bars/plots show mean ± SEM with individual biological replicates; significance is indicated on the panels (*$P<0.05$, **$P<0.01$, ***$P<0.001$).

Figure 5. Inflammation, senescence, telomere length and a working model for EGFR signaling in LRRC8A KO VSMCs. A, NF-κB reporter activity in WT and KO VSMCs under control conditions and after TNFα (10 ng/mL, 6 h) normalized to protein and corrected for infection efficiency (GFP/tubulin) and expressed as fold change from control KO cells, (control samples, one-sample t test; for TNF treated samples Mann-Whitney test; n ≥ 9). **B,** β-galactosidase (β-Gal) protein abundance in WT and KO VSMCs normalized to α-tubulin (n = 3, unpaired t test). **C,** β-Gal enzymatic activity (fluorescence/μg protein) in WT and KO VSMCs grown in either low (*left*) or normal (*right*) serum (n = 6, One-way ANOVA). **D,** Relative telomere length (qPCR T/S ratio, arbitrary units) in WT and KO VSMCs (n = 12, unpaired t test). **E,** Schematic model linking Nox1/LRRC8A/$O_2^{\cdot-}$ and EGFR/AKT signaling to NF-κB and HIF-1α pathways and effects on ECAR/OCR, proliferation, migration, senescence, and longevity. Means ± SEM. *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Figure 6. Reduces aortic atherosclerosis, inflammation, proliferation and senescence in VSMC-specific LRRC8A KO mice is associated with impaired VSMC oxidized-LDL uptake.

A, Aortic atherosclerotic lesions. Representative en-face Oil Red O–stained aortae from a WT and a KO mouse, and quantification of percent lipid-stained area relative to total aortic area in male and female ApoE$^{-/-}$;SM22α-Cre LRRC8A genotypes (WT, HET, KO) mice (One-way ANOVA). **B,** Lipoprotein uptake in primary VSMCs. Confocal quantification of reduced DiI-oxLDL uptake (10 µg/mL, 4 h) in KO VSMCs, or with pharmacologic VRAC inhibition in WT cells using carbenoxolone (CBX), DCPIB, or tamoxifen (TAM). Uptake of non-oxidized DiI-LDL was not altered in KO cells (Mann Whitney test, one way ANOVA). **C,** Aortic protein expression. Immunoblotting of aortic lysates from male ApoE-/-, LRRC8A WT, HET or KO mice normalized to GAPDH for LRRC8A, β-Gal, VCAM, ICAM, PCNA, LOX-1, CD36, and LDLR with densitometric quantification. Each point represents one animal. Bars are mean ± SEM. *p < 0.05.

Figure 8. Mesenteric vasomotor function in ApoE$^{-/-}$ mice lacking VSMC LRRC8A. A, Cumulative concentration-response curves for contraction to phenylephrine (PE) and norepinephrine (NE), and relaxation to acetylcholine (ACh, endothelium-dependent) and sodium nitroprusside (SNP, endothelium-independent) in mesenteric arteries from female and male LRRC8A WT, HET or KO, ApoE$^{-/-}$ mice. Black asterisks indicate comparisons of maximal responses between WT vs HET or KO. Red asterisks indicate EC$_{50}$ comparisons (ACh: WT vs HET or KO; SNP: WT or HET vs KO, male n =6-7, female male n =5-6), error bars are mean ± SEM. **B,** Representative immunoblots from isolated mesenteric artery beds (WT or KO, ApoE$^{-/-}$) normalized to GAPDH. Bars are mean ± SEM normalized to WT. *p < 0.05 (male n = 4 - 8, female n = 3)

A

2OH-Ethidium/protein (pmol/mg)

Wild Type: Control, O₂˙⁻ (*)
LRRC8A KO: Control, O₂˙⁻

B

Hypoxanthine + Xanthine Oxidase → $O_2^{\cdot-}$

$H_2O_2 + Catalase \rightarrow O_2 + H_2O$

$O_2^{\cdot-}$ $O_2^{\cdot-}$ $O_2^{\cdot-}$

LRRC8A LRRC8C LRRC8D

$DHE + O_2^{\cdot-} \rightarrow 2OH\text{-ethidium}$

C

LRRC8A-GFP   DHE   Merge

LRRC8C-GFP   DHE   Merge

D

Relative DHE Fluorescence

Con / O₂˙⁻ (LRRC8A), Con / O₂˙⁻ (LRRC8D), Con / O₂˙⁻ (LRRC8C)

Relative Fluorescence Intensity

GFP Neg
GFP Pos
GFP Pos + (HX-XO)

LRRC8C-GFP

E Cellular and Mitochondrial ROS

WT

KO

Relative Fluorescence Intensity (a.u)   ***
WT   KO
ROSstar550

Relative Fluorescence Intensity (a.u)   ***
WT   KO
Mitosox

F Lipid Peroxides

Fold Change

WT (open)
KO (gray)

Control   TNFα (*)

G GSH/GSSG

GSH/GSSG   *

WT   KO

H Nrf2 Activity

Fold Change   ***

WT   KO

A — PPI Hub Proteins

Term	P-value	Adjusted P-value	Combined Score
EGFR	0.00047	0.0685	16.33
IGF1R	0.00049	0.0685	21.14
JAK2	0.00068	0.0685	22.36
ITGB1	0.00122	0.0924	21.48
MAPK3	0.00286	0.173	12.51

row min   row max

id | EGFR | IGF1R | JAK2 | ITGB1 | MAPK3 | id

PTPRE
GRB10
SH3KBP1
IRS1
PLCG2
NCAM1
VEGFC
FGF2
NEDD9
BRAF
AR
GJA1
ESR1
PGR
DOK1
SOCS2
SOCS3
FES
OSMR
LIFR

B — EGF Signaling

WT KO WT KO
EGF (50ng/ml) - - + +

pEGFR (T1068)   175 KD
EGFR   175 KD
Tubulin   52 KD

pEGFR / EGFR   **   *   EGF - - + +
pEGFR / Tubulin   **   *   EGF - - + +
EGFR / Tubulin   EGF - - + +

C

WT KO WT WT KO KO
EGF (50ng/ml) - - + + + +

pAKT (T^{308})   60 KD
pAKT (S^{473})   60 KD
AKT   60 KD
Tubulin   52 KD

pAKT (T^{308}) / AKT   **   *   EGF - - + +
pAKT (T^{308}) / Tubulin   **   **   EGF - - + +
pAKT (S^{473}) / AKT   ***   EGF - - + +
pAKT (S^{473}) / Tubulin   ***   EGF - - + +
AKT / Tubulin   *   EGF - - + +

D

WT KO
IRS1   180 KD
Tubulin   52 KD

IRS1 / Tubulin   *   WT KO