Semi-automated image analysis of root architecture and early root development in faba bean and white clover and genomic estimation of breeding values and correlations

Istvan Nagy[1,*,†,©], Peter Skov Kristensen[2,†], Elesandro Bornhofen[1], Marta Malinowska[2], Linda Kærgaard Nielsen[3], Andrea Schiemann[4], Niels Rolund[2], Stig Uggerhøj Andersen[5], and Torben Asp[1]

[1]Center for Quantitative Genetics and Genomics, Aarhus University, Forsøgsvej 1, DK-4200 Slagelse, Denmark

[2]DLF Seeds A/S, Ny Østergade 9, DK-4000 Roskilde, Denmark

[3]Sejet Plant Breeding, Nørremarksvej 67, DK-8700 Horsens, Denmark

[4]Nordic Seed, Kornmarken 1, DK-8464 Galten, Denmark

[5]Department of Molecular Biology and Genetics, Aarhus University, Universitetsbyen 83, DK-8000 Aarhus, Denmark

[†]These authors contributed equally

[*]Correspondence: Istvan.Nagy@qgg.au.dk

Abstract

Protein-rich leguminous plants, such as faba bean and white clover are prospectively interesting crops in the North-European countries for reducing dependence on soybean import. Significant expansion of the production area of leguminous crops is challenged by the sub-optimal climatic conditions in this region, especially by the increasing probability of year-to-year fluctuation of extreme weather conditions due to global climate changes. To overcome these challenges, development of new climate-resilient varieties suitable for growing under Northern-European conditions are needed. Root architecture and early root development, as well as the availability of efficient root phenotyping technologies are crucial factors of advancing in breeding of adequate varieties. We report a study of a simple and affordable screening technology of early root development using rhizoboxes in connection with semi-automated image analysis and provide a conceptual pipeline for estimation of Genomic Estimated Breeding Values (GEBVs) and correlating greenhouse- and field phenotype data. Based on bivariate models, high genetic correlation could be detected between total root length values recorded in greenhouse rhizobox experiments and field grain yield in faba bean. In white clover, moderately positive genetic correlation between estimated breeding values of rhizobox-detected total root length and field yield could be identified. Our results suggest that phenotyping and selection of early root development components could potentially be useful in breeding programs to increase the genetic gain for field yield.

Keywords

Root System Architecture, early root development, rhizoboxes, image analysis, multivariate mixed models, Genomic Estimated Breeding Values, faba bean, white clover

Introduction

Agriculture and food industry of Northern European countries, including Denmark are greatly depending on import of crop products providing protein resources - mainly soybean and processed soybean products. To reduce dependence on soybean import, there is an increasing interest in the Nordic countries in expanding local production of protein-rich leguminous crops, like faba bean (*Vicia faba* L.) and white clover (*Trifolium repens* L.).

Expansion of the area of leguminous crops is beneficial for the environmental-friendly agriculture as well: Owing the ability of biological nitrogen fixation, legumes reduce the input of chemical fertilizers.[1] Legumes fit well into the crop rotation, alternating with cereals, tubers or industrial crops, because the spacious legume root system remediates soil structure by creating stable biopores in the subsoil, improving water infiltration, and root penetration for subsequent crops.[2] In addition, the deep root system of legumes contribute to long-term soil carbon storage by stabilizing organic carbon below the frequently disturbed topsoil.[3]

The most limiting factors of expansion of leguminous protein crop production in the Nordic region are the sub-optimal climatic conditions, particularly the increasing probability of year-to-year fluctuation of extreme weather conditions driven by global climate changes. Spring drought episodes can critically restrict water availability during reproductive stages, while water abundance in cold winters may lead to temporary soil saturation and subsoil hypoxia, affecting root function and resource uptake.[4,5] These challenges underline the need of developing new climate-resilient varieties suitable for growing under Northern-European conditions. Root architecture and early root development, together with the availability of efficient root phenotyping technologies are therefore important factors for advancing breeding of adequate varieties.

Root system architecture (RSA) increasingly attracts the interest of scientist in the era of climate change. RSA comprises the shape and spatial arrangement of the root system within the soil[6] and is determined by genetic- as well as by environmental factors, like soil temperature and supply of water and nutrients, while roots exhibit a high level of phenotypic plasticity in response of environmental conditions. Evidently, root architecture is a key factor of efficient nutrient and water uptake, and even a 'second green revolution' has been envisioned that deploys crops with improved below ground traits.[7,8] The development of new plant varieties with improved root phenotypes requires advances in the characterization of root development and morphology, as well as in the relationships between genetic factors and RSA. As being a complex, polygenic trait, studying of RSA should be conducted by parameterization of its elements, such as root depth, total root length, root angle, root thickness, root density and root surface area.[9,10]

A wide range of root phenotyping technologies have been developed and applied during the last decades, ranging from excavating root crowns on the fields ("showelomics"[11]) to semi-field based micro-rhizothron facilities with automated image acquisition.[12] Non-destructive 3D root phenotyping using tomographic techniques, like X-ray computed tomography, magnetic resonance imaging (MRI) and positron emission tomography (PET) also become achievable during the last decades (refer[8] for a review).Soil-filled slab chambers with transparent lids (rhizoboxes) offer a reasonable and space-efficient approach for studying early root development under controlled conditions.

Investigations on early root development are of fundamental importance, as a vigorous early root system is essential for healthy plant growth and establishment, especially under sub-optimal conditions,[13] and recent quantitative genetics studies in perennial ryegrass demonstrated that early root development traits can act as proxies for field yield across multiple years and locations.[14] Earlier studies identified correlations between early root traits and crop productivity and/or nutrient- and water use efficiency in other species, like maize[15,16] and rice[17] as well. Such relations have been confirmed by the co-occurrence of quantitative trait loci (QTLs) between early root traits and yield components.[16,18,19]

The decreasing genotyping costs and the availability of high-quality genomic references in most important crop species provide opportunities for breeders to obtain abundant genotyping information at affordable costs and to utilize large-scale genotype data for genomic predictions along with phenotpype data collected by multiple technologies across multiple years and locations. However, for the successful implementation of such efforts, the development of appropriate statistical models and methodologies is necessary. For polygenic quantitative traits, like RSA components, genomic selection offers opportunities for breeders to increase the rate of genetic gain, by reducing the time needed to complete a breeding cycle, increasing the selection intensity and by utilizing within-family variation that can be captured using molecular markers.[20] Genomic selection is a form of Marker Assisted Selection that simultaneously estimates all locus, haplotype or marker effects across the entire genome to calculate Genomic Estimated Breeding Values (GEBVs[21]). Multivariate models are valuable for integrating multiple correlated traits into the breeding process and estimation of genetic correlations between traits and can potentially improve the accuracy of Genomic Estimated Breeding Values.[22–24]

To facilitate quantitative genetics studies for correlating early root development traits and yield components, we conducted greenhouse rhizobox experiments using faba bean and white clover genotypes, including standard varieties and breeding lines. The specific objectives of the study were to quantify early RSA traits using image-based phenotyping, assess genetic variation and heritabilities using genomic data, and evaluate genetic correlations between early RSA traits and agronomic performance to determine their potential for indirect selection in breeding programs.

Materials and methods

Plant material

For faba bean, 180 lines from the breeding program of the Danish breeding companies Nordic Seeds (75 lines) and Sejet (75 lines), as well as 10 variety standards and 20 core lines belonging to the IMFABA Consortium (https://projects.au.dk/fabagenome/genomics-data) were used in the study. For white clover, a panel consisting of 174 lines selected from 20 commercial varieties with diverse agronomic characters was used. Initial white clover plant material was maintained in the greenhouse as clonally propagated stocks in 17cm pots.

Rhizobox setup and growing conditions

Plants were grown in custom-made plastic rhizoboxes (internal dimensions: 36x18x2.5cm). Boxes were filled with 1.8L substrate (a mix of turf, local topsoil and sand) and supplemented with 300ml water. In case of faba bean, two seeds were sown in each box. For white clover, 8 to 12 cm long cuttings of young lateral shoots were collected from the middle part of the pots. It was attempted that cuttings have a nodus above the cutting site and fully developed leaflets below the shoot tip. During processing, cuttings were kept between wet filter paper sheets. Boxes were placed in the greenhouse on racks that ensured to incline them at 60° to facilitate root growth along the visible side. During growing, the transparent plexiglass front side of the boxes was covered with a black foil. For both plant species, experiments were

conducted in three subsequent batches in randomized blocks throughout a 12 month period in the same greenhouse compartment equipped with an automatic shadowing system during summer and additional lighting and heating (16/8h day/night at 21/18 °C) during winter. In each replicate, plants were grown for 25 to 28 days, by applying manual watering with equal amounts (100 to 200ml) of tap water in 5 to 10 days periods, depending on humidity and outer temperature.

Phenotype data collection

Root imaging

On the final day of growing period, in each batch experiment root images were taken by a horizontal office scanner (Epson Perfection V700) with a modified lid. Root imaging was conducted by the RootPainter software.[25] The analysis procedure involved training with manual annotation of clear foreground and background regions, and corrective annotations to refine the segmentation model. The final model was used to automatically segment the entire root dataset. In addition, segmented root images were exported and further processed by the RhizoVision software[26] to quantify and record the following root phenotype components: total root length, number of root tips, number of branching points and branching frequency, root diameter and perimeter, root volume and surface area.

White clover leaf phenotyping

For leaf phenotyping, in each genotype, three fully developed trifoliate leaves were collected from the source plants maintained in the greenhouse. Separated leaflets (9 leaflets for each genotype) were fixed on white paper sheets by slight glueing their back side and scanned in full-color mode at the highest possible resolution. The scanned images were used for digital image analysis to determine leaf size, leaf shape and leaf color. Leaf size (average surface area of leaflets) and leaf shape parameters (circularity and solidity) were determined by ImageJ (v1.54g).[27] For leaf color analysis, 20x20 pixel squares were cut out from the middle part of the leaflet images and saved to separate files using the image editing software Inkscape.[28] Average pixel RGB values were calculated from the 400px image files by the *convert* tool of the ImageMagick software[29] using a single-line command including the following formula:

$$RGB = convert\ [400px\_image\_file]\ \text{-}resize\ 1x1\backslash!$$
$$\text{-}format\ "\%[fx:int(255*r+.5)], \%[fx:int(255*g+.5)], \%[fx:int(255*b+.5)]" \tag{1}$$

For each line, RGB values were converted to perceived brightness values using the following formula (Luma conversion for perceived brightness, in which the green component has the main effect):

$$Y = 0.2126*R + 0.7152*G + 0.0722*B \tag{2}$$

Faba bean field phenotype data

Faba bean breeding lines and variety standards were tested in replicated multi-year field trials in 3 years (2022, 2023, 2024) at 3 locations (Dyngby, Svinø and Gamborg, Denmark) in alpha-lattice design with 3 replicates. Plant height, Thousand Kernel Weight (TKW), grain yield and grain protein content were recorded, resulting in ca. 4000 observations in total for each trait.

White clover field phenotype data

A two year field experiment was conducted in Haldrup, Denmark, with 12.5 m^2 plots in two replicates.
Fresh and dry biomass yield was recorded after four cuts in the first year and five cuts in the second year.

White clover greenhouse biomass data

Image-based biomass measurements were carried out on a panel of white clover lines overlapping with
the panels of rhizobox- and field experiments. For each line, 10 individual plants were monitored using
stationary cameras to estimate cumulated biomass after 10 days of growth.[30]

Genetic variant data

Faba bean variant data

Variant data based on pseudo-chromosome references of the latest version of the *Vicia faba* genome
sequences (Hedin/2 v.2),[31] containing 122,291 bi-allelic SNPs was provided by the IMFABA consortium
(https://projects.au.dk/fabagenome/genomics-data). Of the 178 lines used for the rhizobox experiments,
157 were represented in the VCF file. Based on the variant data stored in the VCF file, three-dimensional
Prinicipal Component Analysis (PCA) plots were generated using VCF2PCACluster.[32]

White clover variant data

Genetic variant data based on Genotyping by Sequencing (GBS) short-read sequences mapped onto
the *Trifolium repens* v.4.0 genome references.[33] Of the 195 genotypes represented in the original variant
discovery project, 167 were common with the line set used in the root phenotyping experiments, therefore
variant data for the not matching lines were removed from the VCF file and genomic analysis studies
was further on restricted to these 167 lines). The VCF file was further on filtered for a minimum Minor
Allele Frequency of 3%, maximum number of missing data of 30% across lines and a minimum Read
Depth of 5, keeping bi-allelic variants on 280,265 loci.

Genomic analysis conception

Calculating variance components and heritabilities

GBLUP (Genomic Best Linear Unbiased Prediction) models were used in order to partition environmental,
spatial, and genetic effects from each other for the different traits. Here, genomic relationships between
lines based on SNP markers were included in order to estimate breeding values (additive genetic effects)
for each trait, heritabilities (i.e. the ratio between genetic variance and total phenotypic variance), and
to estimate the genetic correlations between traits.[34]

Variance components of all models were estimated by restricted maximum likelihood using the soft-
ware package DMU for analyzing Multivariate Mixed Models (provided by the Center of Quantitative
Genetis and Genomics, Aarhus University[35]).

Heritabilities at field plot or rhizobox level were calculated as narrow sense using the following formula: 181

$$h^2 = \frac{\sigma_g^2}{\sigma_p^2} \tag{3}$$

and as broad sense using the following formula: 182

$$H^2 = \frac{\sigma_g^2 + \sigma_i^2}{\sigma_p^2} \tag{4}$$

where σ_p^2 is the sum of all estimated variance components within each of the models, σ_g^2 and σ_i^2 are 183 variances of additive genetics and of line effects, respectively. 184

Correlations between traits were calculated as Pearson correlations between estimated additive genomic 186 breeding values for the respective traits. 187

Correlated response to selection 188

The genetic response to selection per generation of a trait can be calculated using the breeder's equation: 189

$$R = ih\sigma_A \tag{5}$$

where i is selection intensity, h is the square root of the narrow-sense heritability, σ_A is the genetic 190 standard deviation of the trait (i.e. the square root of the genetic variance). The genetic gain of a trait 191 can be improved by incorporating information from another genetically correlated trait. To determine 192 how much genetic gain can be achieved for a primary trait (Y), when selection is made based on a 193 secondary trait (X), the correlated response to selection can be calculated as: 194

$$CR_Y = ih_X\gamma_A\sigma_{AY} = ih_Xh_Y\gamma_A\sigma_{PY} \tag{6}$$

where $h_Xh_Y\gamma_A$ is the co-heritability of the two traits X and Y, γ_A is the genetic correlation between 195 traits X and Y, the genetic and phenotypic standard deviations of trait Y are σ_{AY},σ_{AP} and σ_{PY} 196 respectively, according to Falconer and Mackay.[34] 197

Thus, the genetic gain of trait Y, when selecting indirectly based on trait X, will better in cases 198 where the heritability of trait X is high, where the genetic correlation between the traits is high, or where 199 the selection intensity can be increased. 200

Yield is often an important breeding goal, but it is relatively expensive and time-consuming to get 201 phenotypic observations for yield, since varieties ideally should be tested in replicated field trials in 202 several locations and several years due to potential spatial variation within fields and due to Genotype 203 by Environment (GxE) interactions across different environments. Therefore, yield cannot directly be 204 measured very accurately early in breeding programs, when only one or few plants are available per 205 variety. However, if other traits can be identified that are genetically correlated to yield and that can be 206 phenotyped more easily and/or in shorter time, e.g. in controlled conditions, using few seeds or plants 207 per variety, then such traits would be useful to implement in early selection stages of breeding programs 208 in order to increase the genetic gain of yield. 209

Dominance and epistatic genomic relationships were not explicitly included in the models, since large data sets are required to estimate such non-additive genetic effects. Instead, the line variance was included in order to capture non-additive genetic variance as well as any additive genetic variance that was not captured by the SNP markers used in the genomic relationship matrix. The line variance can only be estimated for the traits, where phenotypic observations are available for more than one replicate, since the line variance will be confounded with the residual variance if only one observation is available per line.

Verifying genetic correlations

For entry means of two correlated traits (for example total root length *vs.* grain yield), a simple bivariate Bayesian model was fitted using the R package *brms*.[36] Each response included a random intercept for entry, with residual correlations fixed to zero. Models were run for 10,000 iterations (1,000 warm-up; thinning = 5; one chain)

Models

Faba bean models

Yield data from field trials was modelled by:

$$y_1 = X_1 b_1 + Z_1 t + Z_2 g_1 + Z_3 i_1 + e_1 \tag{7}$$

where $\boldsymbol{y}_1$ is a vector of phenotypic observations, $\boldsymbol{X}_1$ is a design matrix for fixed effects, $\boldsymbol{b}_1$ is a vector of effects of year-location-replicate, $\boldsymbol{Z}_1$, $\boldsymbol{Z}_2$ and $\boldsymbol{Z}_3$ are design matrices for random effects, $\boldsymbol{t}$ is a vector of effects of blocks within year-location-replicate with $\boldsymbol{t} \sim N(0, \boldsymbol{I}_1 \sigma_t^2)$, σ^2_t is the variance of block effects, $\boldsymbol{g}_1$ is a vector of additive genomic breeding values with $\boldsymbol{g}_1 \sim N(0, \boldsymbol{G}_1 \sigma_{g_1}^2)$, $\boldsymbol{G}_1$ is a genomic relationship matrix calculated using method one proposed by VanRaden,[37] $\sigma_{g_1}^2$ is the additive genetic variance, $\boldsymbol{i}_1$ is a vector of line effects with $\boldsymbol{i}_1 \sim N(0, \boldsymbol{I}_2 \sigma_{i_1}^2)$, $\sigma_{i_1}^2$ is the variance of line effects, and $\boldsymbol{e}_1$ is a vector of residual effects with $\boldsymbol{e}_1 \sim N(0, \boldsymbol{I}_3 \sigma_{e_1}^2)$, $\sigma_{e_1}^2$ is the variance of residual effects, and $\boldsymbol{I}_n$ are identity matrices.

Root length data from rhizoboxes was modelled by:

$$y_2 = X_2 b_2 + cw + Z_4 g_2 + Z_5 i_2 + e_2 \tag{8}$$

where $\boldsymbol{y}_2$ is a vector of phenotypic observations, $\boldsymbol{X}_2$ is a design matrix for fixed effects, $\boldsymbol{b}_2$ is a vector of effects of experiment, $\boldsymbol{c}$ is a fixed regression coefficient for observed root lengths regressed on thousand seed weights, and $\boldsymbol{w}$ is a vector of thousand seed weights of seeds sown in rhizoboxes, $\boldsymbol{Z}_4$ and $\boldsymbol{Z}_5$ are design matrices for random effects, $\boldsymbol{g}_2$ is a vector of additive genomic breeding values with $\boldsymbol{g}_2 \sim N(0, \boldsymbol{G}_1 \sigma_{g_2}^2)$, $\sigma_{g_2}^2$ is the additive genetic variance, $\boldsymbol{i}_2$ is a vector of line effects with $\boldsymbol{i}_2 \sim N(0, \boldsymbol{I}_4 \sigma_{i_2}^2)$, $\sigma_{i_2}^2$ is the variance of line effects, and $\boldsymbol{e}_2$ is a vector of residual effects with $\boldsymbol{e}_2 \sim N(0, \boldsymbol{I}_5 \sigma_{e_2}^2)$, $\sigma_{e_2}^2$ is the variance of residual effects, and $\boldsymbol{I}_n$ are identity matrices.

Combining the two models above, a bivariate model was used for estimating genetic covariances between field yield and rhizobox root length traits:

$$\begin{bmatrix} \boldsymbol{g}_1 \\ \boldsymbol{g}_2 \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{g1}^2 & \sigma_{g_1g_2} \\ \sigma_{g1g2} & \sigma_{g2}^2 \end{bmatrix} \otimes \boldsymbol{G}_1 \right), \begin{bmatrix} \boldsymbol{i}_1 \\ \boldsymbol{i}_2 \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{i1}^2 & \sigma_{i_1i_2} \\ \sigma_{i1i2} & \sigma_{i2}^2 \end{bmatrix} \right), \begin{bmatrix} \boldsymbol{e}_1 \\ \boldsymbol{e}_2 \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} \sigma_{e1}^2 & 0 \\ 0 & \sigma_{e2}^2 \end{bmatrix} \right)$$
(9)

The remaining random effects were assumed normally and independently distributed with zero means and variances as described above.

The genetic correlation between traits, $\gamma_{g_1g_2}$, can be calculated from the estimated genetic variances and co-variances:

$$\gamma_{g_1g_2} = \frac{\sigma_{g_1g_2}}{\sqrt{\sigma_{g1}^2 * \sigma_{g2}^2}}$$
(10)

Clover models

Yield data from field trials was modelled by:

$$y_3 = X_3b_3 + Z_6p + Z_7g_3 + e_3$$
(11)

where $\boldsymbol{y}_3$ is a vector of phenotypic observations, $\boldsymbol{X}_3$ is a design matrix for fixed effects, $\boldsymbol{b}_3$ is a vector of effects of cut number, $\boldsymbol{Z}_6$ and $\boldsymbol{Z}_7$ are design matrices for random effects, $\boldsymbol{p}$ is a vector of effects of plot with $\boldsymbol{p} \sim N(0, \boldsymbol{I}_6\sigma_p^2)$, σ_p^2 is the variance of plot effects, $\boldsymbol{g}_1$ is a vector of additive genomic breeding values with $\boldsymbol{g}_3 \sim N(0, \boldsymbol{G}_2\sigma_{g3}^2)$, $\boldsymbol{G}_2$ is a genomic relationship matrix, σ_{g3}^2 is the additive genetic variance, and $\boldsymbol{e}_3$ is a vector of residual effects with $\boldsymbol{e}_3 \sim N(0, \boldsymbol{I}_7\sigma_{e3}^2)$, σ_{e3}^2 is the variance of residual effects, and $\boldsymbol{I}_n$ are identity matrices.

Data from greenhouse was modelled by:

$$y_4 = X_4b_4 + Z_8g_4 + e_4$$
(12)

where $\boldsymbol{y}_4$ is a vector of phenotypic observations, $\boldsymbol{X}_4$ is a design matrix for fixed effects, $\boldsymbol{b}_4$ is a vector of effects of experiment, $\boldsymbol{Z}_8$ is a design matrix for random effects, $\boldsymbol{g}_4$ is a vector of additive genomic breeding values with $\boldsymbol{g}_4 \sim N(0, \boldsymbol{G}_2\sigma_{g4}^2)$, σ_{g4}^2 is the additive genetic variance, and $\boldsymbol{e}_4$ is a vector of residual effects with is the variance of residual effects with $\boldsymbol{e}_4 \sim N(0, \boldsymbol{I}_8\sigma_{e4}^2)$, σ_{e4}^2 is the variance of residual effects, and $\boldsymbol{I}_n$ identity matrices.

Results and discussion

Rhizobox experiments

Rhizobox setups differed between the two species: faba bean seeds were sown directly, while white clover plants were raised from shoot cuttings. The latter approach proved more challenging, as the greenhouse stock plants showed some inherent variability, which affected the uniformity of starting material. In addition, the three experimental batches were conducted sequentially over a full year in the same greenhouse compartment. Seasonal differences certainly influenced the consistency of data

obtained from different experiment batches. Roots were scanned when they reached about two-thirds of the rhizobox hight: after 12 to 15 days in case of faba bean, and 15 to 20 days in case of white clover (Figure 1). RootPainter provided a reliable root image segmentation across images. Image-based quantification resulted in approximately normal distributions for most parameters (Figure 2). Among the parameters extracted, total root length (TRL) was consistently the most robust trait, with lower sensitivity to background noise or overlapping roots. In contrast, vertical measurements, such as root diameter or perimeter were more prone to errors, as overlapping side roots and image artefacts interfered with accurate detection. These issues underline that TRL is an appropriate RSA descriptor for downstream analyses, while caution is needed when interpreting vertical or diameter-based traits.

Genetic variant data, diversity and population structure

In case of both species, high-quality reference genomes and large sets of genetic variant data were available for the lines included in the study (bi-allelic SNPs on 122,291 loci for faba bean and on 280,265 loci for white clover). In faba bean, the core plant material included 75-75 breeding lines from two Danish breeding companies. According to the PCA plot based on genetic variant data is showing that the majority of these faba breeding lines are separating to distinct clusters according to company origin, suggesting that the breeding programs of the companies is predominantly based in different germ plasm resources. resources (Figure 3). For white clover, a panel consisting of 174 lines selected from 20 commercial varieties (4 to 10 lines for each variety) was used. In both species, the available genetic variant data enabled to build robust and reliable Genomic Relationship Matrices for genomic estimation of breeding values and correlations (Figure 4).

Genomic estimations and correlations

Faba bean

The narrow-sense heritability at plot level for faba bean grain yield was 0.18, and the broad-sense heritability was 0.23, which indicates that the genetic control of yield is largely due to additive genetic effects and to a smaller extent due to non-additive genetic effects. It is likely, that with larger datasets it would be possible to determine whether such differences are due to non-additive effects or non-captured genomic regions.

On the other hand, the narrow-sense and broad-sense heritabilities for total root length of 0.07 and 0.13, respectively, indicate that additive and non-additive genetic effects were approximately equally important for root length, and that root length could also be strongly affected by other factors, such as environmental effects or GxE interactions.

Based on the bivariate model for grain yield and total root length, the estimated additive genetic correlation between the two traits was 0.83 and the estimated correlation of the line effects was 0.53. So, these two traits seem to be highly genetically correlated, albeit at a quite high level of standard errors in both cases, 0.47 and 0.45 respectively (Table 1, Figure 5). Nevertheless, distribution analyses confirmed the robustness of the estimated genetic correlation between greenhouse total root length (TRL) values and field grain yield: A histogram (Figure 7a) reveals a strongly positive correlation, with the posterior

Table 1: Genomic estimations and heritabilities - Faba bean

	Variance component				Heritability	
	Block	**Id$_g$**	**Id$_l$**	**e**	**h^2**	**H^2**
Grain yield						
Estimate (*Standard error*)	15.2 (*1.1*)	8.8 (*2.3*)	2.6 (*1.4*)	23.1 (*0.6*)	0.18	0.23
Protein content						
Estimate (*Standard error*)	0.15 (*0.03*)	1.35 (*0.26*)	0.16 (*0.13*)	1.33 (*0.04*)	0.45	0.50
Total root length						
Estimate (*Standard error*)	-	510.1 (*473.8*)	405.0 (*529.1*)	6356.3 (*516.3*)	0.07	0.13

distribution peaking near 0.8 and the majority of the probability mass falling between 0.3 and 1.0. As the 95% Bayesian Credible Interval (CI) does not cover zero, there is strong evidence that the genetic correlation is positive, distinct from zero, and statistically credible, and a trace plot (Figure 7b) confirms that the model has successfully converged with well-mixed and stationary sampling. This suggests that the genetic mechanisms driving early root length are reliably linked to seed yield performance in field conditions, indicating that phenotyping and selection for early root development components, particularly total root length could potentially be useful in breeding programs to increase the genetic gain for grain yield.

White clover

We found a positive correlation of 0.17 between GEBVs of total root length (TRL), measured in rhizoboxes and field yield, although the heritability of TRL was very low, which can be explained by the restricted availabilty of field phenotype data, as only a few lines were phenotyped for all traits in two replicates, and therefore the estimates of the variances, breeding values, and correlations could not be very accurate.

On the other hand, phenotyping for leaf parameters could be carried out in the greenhouse for all lines, for which genetic variant data was available. Moderately positive genetic correlations of 0.26 between estimated breeding values of yield from field cuts and of both leaf size and of leaf solidity were identified. For leaf size and leaf solidity, moderate to high narrow-sense heritabilities of 0.46 and 0.63, respectively, were estimated. Both traits had higher narrow-sense heritabilities than 0.22 that was estimated for field yield, and they could therefore be suitable indicator traits in breeding programs for improving yield by indirect selection. The leaf traits can be measured on few, young plants grown in controlled greenhouse conditions, and selection based on these traits might thereby be used at low costs in early stages of breeding programs to increase the genetic gain of yield. Furthermore, phenotyping such traits on selection candidates may improve genomic prediction accuracies for yield in cases where larger training sets are available to use for multivariate models (Table 2, Figure 6).

Table 2: Genomic estimations and heritabilities - White clover

	Heritability[a]	KGGM	TRL	LS	Lsize	iSize_GEBV[b]	gpdCor_GEBV[c]
Drymatter, kg (KGGM)	0.22	1.00	0.17	0.26	0.26	0.21	-0.13
Total root length (TRL)	0.02	0.17	1.00	0.08	0.11	-0.06	0.13
Leaf solidity (LS)	0.63	0.26	0.08	1.00	0.61	0.27	-0.07
Leaf size (Lsize)	0.46	0.26	0.11	0.61	1.00	0.40	-0.01
iSize_GEBV	0.83	0.21	-0.06	0.27	0.40	1.00	-0.06
gpdCor_GEBV	0.24	-0.13	0.13	-0.07	-0.01	-0.06	1.00

[a]Narrow sense
[b]Image-based biomass measurements calculated from pixel counts in the first 10 days of growth
[c]Dry weight per plant corrected for the full effect of iSize

Conclusions

Field performance of crop varieties, in terms of biomass- or grain yield is significantly affected by genotype-by-environment (GxE) interactions, which complicates the selection of genotypes consistently outperforming under variable environmental conditions. Field trials, as standard tools for crop yield performance estimations, are costly, time consuming and laborious, and even with a sufficient number of replicates, plot experiments are often subjected to representation bias due to soil inhomogenity and various technical- and management issues. Incorporating quantitative phenotype traits that can be assessed under controlled environmental conditions in early developmental phases, while positively correlating with field yield components, can improve the predictability of field performance of crops in different stages of breeding programs.In accordance with earlier findings, our results confirmed that early root development traits can be used as prognostic indicators for field yield across different environments and locations: Both in faba bean as well as in white clover we found significant genetic correlations between Genomic Estimated Breeding Values (GEBVs) of total root length assessed in rhizoboxes and harvested yield in field plot experiments.These findings highlights the value of early root development traits in enhancing genetic gain in breeding programs. Currently, the accessibility to state-of-the-art sequencing technologies and high quality genomic resources effectively support efficient and accurate genomic prediction in all important crop species, but the availability of robust and reliable field phenotype data, well representing the genetic variations targeted by the study, often remains a bottleneck in such projects. Another important aspects of genomic prediction are the availability, development and implementation of appropriate statistical models and methodologies. Our results demonstrated that multi- and bivariate models along with the conceptual pipeline provided in the present case study are well applicable for correlating early root development traits and field yield components. Consequently, these results suggest that early root development traits are of high predictive potential in connection with field yield components, especially if field phehotype data are available to a limited extent.

Acknowledgements 350

This work was supported by Promilleafgiftsfonden for Landbrug grant Cultivation of protein crops with 350
a low environmental and climate footprint for the future climate (41647 Klimaprotein). Faba bean 351
genotyping- and field experiment data were generated in the frame of the IMFABA project, supported 352
by The European Union's Horizon 2020 Programme for Research & Innovation in the ERA-NET Cofund 353
SusCrop project ProFaba (Grant no. 771134). White clover genotying- and greenhouse biomass data 354
were generated in the frame of the NCHAIN project, funded by the Innovation Fund Denmark (Grant 355
no. 4105-00007A). 356

Author contribution 357

Conceptualization: TA, Methodology: IN, MM and PSK, Investigations: IN, LKN, AS and NR, Data 358
analysis: IN, PSK and EB, Writing: IN, PSK and MM, Funding acquisition: TA and SUA 359

Figures

Figure 1: Greenhouse rhizobox experiments and root image processing
  (a) Faba bean seedlings in rhizoboxes
  (b) White clover explants in rhizoboxes
  (c) Scanned root image
  (d) Segmentation image made by RootPainter
  (e) Segmentation image exported to analysis by RhizoVision

Figure 2: Distribution of quantitative root parameter values in three faba bean rhizobox batch experiments

Figure 3: 3D clustered PCA results of Faba bean breeding lines based on bi-allelic SNP markers. **SJ** and **NS** are for the Danish breeding companies Sejet and Nordic Seed

Figure 4: Genomic realationship matrices based on bi-allalic SNP markers.
(a) Faba bean lines, Blue bars: **Nordic Seed** lines, Red bars: **Sejet** lines
(b) White clover lines

Figure 5: Correlations between Genomic Estimated Breeding Values in faba bean breeding lines.
TRL: Total root length

Figure 6: Correlations between Genomic Estimated Breeding Values in white clover breeding lines. **KGGM**: Green matter, kg, **TRL**: Total root length, **LS**: Leaf solidity

Figure 7: Posterior distribution and trace plot of the genetic correlation between total root length (**TRL**) and grain yield in faba bean.

(a) Histogram of posterior samples for the correlation parameter between the random intercepts of the two traits

(b) Trace plot (MCMC chain), indicating the sampling behavior over 3,000 iterations

References

1. Heslop, A.D., Ford, J.L., Jahufer, Z., and Hofmann, R.W. (2025). From root to shoot: Morphological evaluation of an international collection of red clover (*Trifolium pratense* L.) populations. Journal of Agronomy and Crop Science *211*. doi: `10.1111/jac.70055`.

2. Behrend, D., Athmann, M., Han, E., Küpper, P.M., Perkons, U., Bauke, S.L., Köpke, U., Kautz, T., Gaiser, T., and Seidel, S.J. (2025). Do biopores created by perennial fodder crops improve the

growth of subsequent annual crops? A synthesis of multiple field experiments. Field Crops Research *322*, 109687. doi: `10.1016/j.fcr.2024.109687`.

3. Kell, D.B. (2011). Breeding crop plants with deep roots: their role in sustainable carbon, nutrient and water sequestration. Annals of Botany *108*, 407–418. doi: `10.1093/aob/mcr175`.

4. Odone, A., Popovic, O., and Thorup-Kristensen, K. (2023). Deep roots: implications for nitrogen uptake and drought tolerance among winter wheat cultivars. Plant and Soil *500*, 13–32. doi: `10.1007/s11104-023-06255-5`.

5. Ober, E.S., Alahmad, S., Cockram, J., Forestan, C., Hickey, L.T., Kant, J., Maccaferri, M., Marr, E., Milner, M., Pinto, F., Rambla, C., Reynolds, M., Salvi, S., Sciara, G., Snowdon, R.J., Thomelin, P., Tuberosa, R., Uauy, C., Voss-Fels, K.P., Wallington, E., and Watt, M. (2021). Wheat root systems as a breeding target for climate resilience. Theoretical and Applied Genetics *134*, 1645–1662. doi: `10.1007/s00122-021-03819-w`.

6. Rogers, E.D., and Benfey, P.N. (2015). Regulation of plant root system architecture: implications for crop advancement. Current Opinion in Biotechnology *32*, 93–98. doi: `10.1016/j.copbio.2014.11.015`.

7. Lynch, J.P. (2007). Roots of the second green revolution. Australian Journal of Botany *55*, 493–512. doi: `10.1071/bt06118`.

8. Atkinson, J.A., Pound, M.P., Bennett, M.J., and Wells, D.M. (2019). Uncovering the hidden half of plants using new advances in root phenotyping. Current Opinion in Biotechnology *55*, 1–8. doi: `10.1016/j.copbio.2018.06.002`.

9. Lynch, J. (1995). Root architecture and plant productivity. Plant Physiology *109*, 7–13. doi: `10.1104/pp.109.1.7`.

10. Maqbool, S., Hassan, M.A., Xia, X., York, L.M., Rasheed, A., and He, Z. (2022). Root system architecture in cereals: progress, challenges and perspective. The Plant Journal *110*, 23–42. doi: `10.1111/tpj.15669`.

11. Trachsel, S., Kaeppler, S.M., Brown, K.M., and Lynch, J.P. (2010). Shovelomics: high throughput phenotyping of maize (*Zea mays* L.) root architecture in the field. Plant and Soil *341*, 75–87. doi: `10.1007/s11104-010-0623-8`.

12. Svane, S.F., Jensen, C.S., and Thorup-Kristensen, K. (2019). Construction of a large-scale semi-field facility to study genotypic differences in deep root growth and resources acquisition. Plant Methods *15*. doi: `10.1186/s13007-019-0409-9`.

13. Wang, Y., Thorup-Kristensen, K., Jensen, L.S., and Magid, J. (2016). Vigorous root growth is a better indicator of early nutrient uptake than root hair traits in spring wheat grown under low fertility. Frontiers in Plant Science *7*. doi: `10.3389/fpls.2016.00865`.

14. Malinowska, M., Kristensen, P.S., Nielsen, B., Fè, D., Ruud, A.K., Lenk, I., Greve, M., and Asp, T. (2025). The value of early root development traits in breeding programs for biomass yield in perennial ryegrass *Lolium perenne* L.). Theoretical and Applied Genetics *138*. doi: `10.1007/s00122-024-04797-5`.

15. Tuberosa, R., Sanguineti, M.C., Landi, P., Michela Giuliani, M., Salvi, S., and Conti, S. (2002). Identification of QTLs for root characteristics in maize grown in hydroponics and analysis of their overlap with QTLs for grain yield in the field at two water regimes. Plant Molecular Biology *48*, 697–712. doi: `10.1023/a:1014897607670`.

16. Li, P., Chen, F., Cai, H., Liu, J., Pan, Q., Liu, Z., Gu, R., Mi, G., Zhang, F., and Yuan, L. (2015). A genetic relationship between nitrogen use efficiency and seedling root traits in maize as revealed by QTL analysis. Journal of Experimental Botany *66*, 3175–3188. doi: `10.1093/jxb/erv127`.

17. Uga, Y., Sugimoto, K., Ogawa, S., Rane, J., Ishitani, M., Hara, N., Kitomi, Y., Inukai, Y., Ono, K., Kanno, N., Inoue, H., Takehisa, H., Motoyama, R., Nagamura, Y., Wu, J., Matsumoto, T., Takai, T., Okuno, K., and Yano, M. (2013). Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. Nature Genetics *45*, 1097–1102. doi: `10.1038/ng.2725`.

18. Atkinson, J.A., Wingen, L.U., Griffiths, M., Pound, M.P., Gaju, O., Foulkes, M.J., Le Gouis, J., Griffiths, S., Bennett, M.J., King, J., and Wells, D.M. (2015). Phenotyping pipeline reveals major seedling root growth QTL in hexaploid wheat. Journal of Experimental Botany *66*, 2283–2292. doi: `10.1093/jxb/erv006`.

19. Jia, Z., Liu, Y., Gruber, B.D., Neumann, K., Kilian, B., Graner, A., and von Wirén, N. (2019). Genetic dissection of root system architectural traits in spring barley. Frontiers in Plant Science *10*. doi: `10.3389/fpls.2019.00400`.

20. Heslot, N., Jannink, J., and Sorrells, M.E. (2015). Perspectives for genomic selection applications and research in plants. Crop Science *55*, 1–12. doi: `10.2135/cropsci2014.03.0249`.

21. Meuwissen, T.H.E., Hayes, B.J., and Goddard, M.E. (2001). Prediction of total genetic value using genome-wide dense marker maps. Genetics *157*, 1819–1829. doi: `10.1093/genetics/157.4.1819`.

22. Thompson, R., and Meyer, K. (1986). A review of theoretical aspects in the estimation of breeding values for multi-trait selection. Livestock Production Science *15*, 299–313. doi: `10.1016/0301-6226(86)90071-0`.

23. Habier, D., Fernando, R.L., and Dekkers, J.C.M. (2007). The impact of genetic relationship information on genome-assisted breeding values. Genetics *177*, 2389–2397. doi: `10.1534/genetics.107.081190`.

24. Jia, Y., and Jannink, J.L. (2012). Multiple-trait genomic selection methods increase genetic value prediction accuracy. Genetics *192*, 1513–1522. doi: `10.1534/genetics.112.144246`.

25. Smith, A.G., Han, E., Petersen, J., Olsen, N.A.F., Giese, C., Athmann, M., Dresbøll, D.B., and Thorup-Kristensen, K. (2022). RootPainter: deep learning segmentation of biological images with corrective annotation. New Phytologist *236*, 774–791. doi: `10.1111/nph.18387`.

26. Seethepalli, A., Guo, H., Liu, X., Griffiths, M., Almtarfi, H., Li, Z., Liu, S., Zare, A., Fritschi, F.B., Blancaflor, E.B., Ma, X.F., and York, L.M. (2020). Rhizovision crown: An integrated hardware and software platform for root crown phenotyping. Plant Phenomics *2020*, 3074916. doi: `10.34133/2020/3074916`.

27. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH image to ImageJ: 25 years of image analysis. Nature Methods *9*, 671–675. doi: `10.1038/nmeth.2089`.

28. Inkscape. Inkscape project (2020). Tech. Rep. Retrieved from https://inkscape.org (2020).

29. ImageMagick. ImageMagick Studio LLC. Tech. Rep. Retrieved from https://imagemagick.org (2024).

30. Moeskjær, S., Skovbjerg, C.K., Tausen, M., Wind, R., Roulund, N., Janss, L., and Andersen, S.U. (2021). Major effect loci for plant size before onset of nitrogen fixation allow accurate prediction of yield in white clover. Theoretical and Applied Genetics *135*, 125–143. doi: `10.1007/s00122-021-03955-3`.

31. Jayakodi, M., Zhang, H., Windhorst, A., Bornhofen, E., Tulpova, Z., Novak, P., Macas, J., Simkova, H., Nadzieja, M., Kim, J.M., Cram, D., Cao, Y., Konkin, D., Sass, O., Welna, G., Himmelbach, A., Mascher, M., Link, W., Kwon, S.J., Andersen, S., and Yang, T.J. (2024). Allelic variation at a single locus distinguishes spring and winter faba beans. Research Square. doi: `10.21203/rs.3.rs-5356723/v1`.

32. He, W., Xu, L., Wang, J., Yue, Z., Jing, Y., Tai, S., Yang, J., and Fang, X. (2024). VCF2PCACluster: a simple, fast and memory-efficient tool for principal component analysis of tens of millions of SNPs. BMC Bioinformatics *25*. doi: `10.1186/s12859-024-05770-1`.

33. Griffiths, A.G., Moraga, R., Tausen, M., Gupta, V., Bilton, T.P., Campbell, M.A., Ashby, R., Nagy, I., Khan, A., Larking, A., Anderson, C., Franzmayr, B., Hancock, K., Scott, A., Ellison, N.W., Cox, M.P., Asp, T., Mailund, T., Schierup, M.H., and Andersen, S.U. (2019). Breaking free: The genomics of allopolyploidy-facilitated niche expansion in white clover. The Plant Cell *31*, 1466–1487. doi: `10.1105/tpc.18.00606`.

34. Falconer, D.S., and Mackay, T. (2009). Introduction to quantitative genetics. Pearson, Prentice Hall. ISBN 9780582243026.

35. Madsen, P., and Jensen, J. A user's guide to DMU. a package for analysing multivariate mixed models. Version 6, release 5.6. Tech. Rep. Available from https://dmu.ghpc.au.dk (2024).

36. Bürkner, P.C. (2021). Bayesian Item Response Modeling in R with brms and Stan. Journal of Statistical Software *100.* doi: `10.18637/jss.v100.i05`.

37. VanRaden, P. (2008). Efficient methods to compute genomic predictions. Journal of Dairy Science *91*, 4414–4423. doi: `10.3168/jds.2007-0980`.