Rapid adaptation follows experimental assisted gene flow in subset of annual monkeyflower populations

Authors: Donna M. Hinrichs[1], Courtney M. Patterson[1], Andrea Turcu[1], Stacy D. Holt Jr.[1], Simon G. Innes[1], Nicholas J. Kooyers[1,2]

Affiliations:

[1]Department of Biology, University of Louisiana, Lafayette; Lafayette, LA 70503 USA

[2]Corresponding author. Email: Nicholas.kooyers@louisiana.edu, Phone : +1(337)-482-1258, Address : 410 E. St. Mary Blvd.; Billeaud Hall, Rm 108; Lafayette, LA 70503 USA

ORCID and Email Addresses

DMH: 0009-0009-9100-1204 - donnasaurrus@gmail.com

CMP: 0000-0002-8304-5276 - patterson.courtneym@gmail.com

AT: 0000-0002-1899-4969 - andrea.turcu1@louisiana.edu

SDH: 0000-0001-7684-024X - stacy.holt1@louisiana.edu

SGI: 0000-0003-3859-1400 - simon.innes1@louisiana.edu

NJK: 0000-0003-3398-7377 - Nicholas.kooyers@louisiana.edu

Short Title : Rapid adaptation follows assisted gene flow

Author Contributions

N.J.K. conceived and designed the experiments. DMH, CMP, AT, SDH, SGI, and NJK collected data and performed the experiments. DMH and NJK analyzed the data. DMH and NJK wrote the original draft of the manuscript, and all authors reviewed and edited the manuscript. All authors approved of the final submission.

Data Availability Statement

Datasets containing field fitness data and phenotypic data from the resurrection experiment will be deposited on Dryad. Sequence data will be uploaded to NCBI SRA as .fastq files.

Word Counts

  Abstract: 200
  Main Text: 4476
  Main Text + Methods: 6360
  Number of Figures and Tables: 4 figures, 0 tables
  Supplemental Material: 9 figures, 14 tables, Extended Methods

Keywords: Evolutionary Rescue, Genetic Rescue, Introgression, Adaptation, Conservation Genomics, Restoration, Applied Management, Assisted Migration, Seed Sourcing, *Mimulus guttatus* (syn. *Erythranthe guttata*)

Abstract

Assisted gene flow, the human-facilitated movement of species within its range, has the

potential to increase genetic diversity and facilitate adaptation for climate-threatened

populations. However, concerns regarding outbreeding depression and gene swamping have

limited the application of assisted gene flow, and experimental tests have been largely confined

to simulations and laboratories. We conducted a landscape-scale manipulation, planting seeds

and seedlings from historically hotter and drier source populations into climate-threatened

populations of the common yellow monkeyflower (*Mimulus guttatus*) and tracked change in

genomes, phenotypes, and fitness. Within three generations, source alleles introgressed into

half of the target populations. Assisted gene flow increased fitness in one experimental block of

populations and slightly decreased fitness in a second block. Fitness increases were associated

with flowering earlier and producing fewer trichomes. Greater fitness changes occurred in

populations where seeds rather than seedlings were introduced. Both the amount of

introgression and fitness changes were heterogeneous between years, suggesting that

temporal fluctuations in climate likely impact initial introgression and success of assisted gene

flow. Initial introgression from source populations was low across the genome, limiting concerns

of gene swamping. Our results are generally consistent with theoretical models, and provide

cautious optimism for an often-maligned conservation strategy.

Introduction

Increasing patterns of maladaptation and declining population sizes indicate that there is a

substantial threat of climate change to many organisms [1–5]. The potential risk of species

extinction [4,6–9] and loss of key ecosystem services [10–14] suggest that conservation action

is necessary with stipulation that actions will not further endanger species. An increasingly

popular intervention is assisted gene flow (AGF) - the facilitated movement of individuals within

a species range to increase the overall genetic diversity of a target population or facilitate

adaptation of a target population to changing conditions [15]. Simulation and experimental

evolution studies suggest that AGF could be highly successful at increasing the long-term

fitness of populations [16,17], with the caveat that AGF is largely untested in natural populations

[18]. Based on these potential benefits, assisted gene flow has been conducted in several plant

and animal populations that are at risk of extirpation [18–21].

Although the potential of AGF is well-recognized, it is still highly controversial within the

conservation community due to both perceived risks and lack of experimental data within natural

populations. Two common perceived risks are outbreeding depression and gene swamping [22–

24]. Introducing novel alleles into a population via outcrossing and subsequent recombination

could add potentially maladapted alleles into a locally adapted population or break-up co-

adaptive gene complexes [25–32]. Adding non-local alleles that are much more adaptive than

local alleles could cause the loss of local genetic diversity [33,34]. Theoretical models of

evolutionary rescue suggest that some maladaptation is likely to occur following introduction of

new variation [35–38], but evidence of outbreeding depression is mixed with most research

emphasizing that the effects of outbreeding are temporary [30,39,40]. Instead, simulation

studies suggest that the benefits of AGF can be rapid (<10 generations) when fitness traits are

controlled by few large-effect loci [41]. Landscape-scale tests of AGF in natural populations are

necessary to evaluate potential benefits and risks.

Because AGF has historically been reserved for highly threatened species, there has been

limited opportunity to dissect the consequences of AGF through controlled experiments (but see

[18,42,43]. Several important hypotheses need to be evaluated. First, the impact of AGF is

predicted to be proportional to the number of individuals introduced within natural populations

and the life stage of the introduced organisms [21,41]. For instance, in plant populations, it may

be more useful to introduce seedlings rather than seeds, as seedlings do not need to meet local

germination requirements. Alternatively, introducing seeds may buffer against extreme years

and allow germination within more benign conditions [44]. Second, determining the amount and

location of introgression of non-local alleles into target population genomes is key for assessing

the potential for gene swamping, outbreeding depression, and hybrid vigor. Third, environmental

heterogeneity between different target populations or years may or may not create variation in

the success of AGF [43]. Both the level of introgression and intensity of selection could fluctuate

between years. Finally, evaluating the potential for non-local alleles to move via gene flow

between geographically proximate populations is necessary for predicting off target effects (e.g.

similar to transgenes into wild populations; [45]). Evaluating the uncertainty and understanding

the mechanisms underlying success or failure of AGF provides more confidence for undertaking

AGF for threatened or endangered species.

Here, we evaluate short-term fitness consequences after conducting a landscape-scale assisted

gene flow experiment in climate-threatened subalpine populations of the common yellow

monkeyflower, *Mimulus guttatus* (syn. *Erythranthe guttata*). While *M. guttatus* is widespread

across western North America and invasive in the U.K. and New Zealand [46,47], annual

populations in the central Oregon Cascades are arguably climate-threatened [44,48,49]. Annual

populations of *M. guttatus* are characterized by highly variable growing seasons that depend on

ephemeral water availability – growing seasons begin with winter/spring rain or snowmelt and

end with terminal droughts [50]. Growing seasons are particular short and variable in annual

populations in the central Oregon Cascades, and climate change has both shifted growing

seasons earlier and caused more frequent and severe heat waves within these seeps. These

populations now exhibit an adaptation lag, where populations from warmer and drier climates

with earlier growing seasons (i.e. populations from the foothills of the Sierra Nevada in

California) have higher fitness than native Oregon populations [48,51]. Additionally, short heat

waves in the last decade have had severe demographic impacts on these populations (50.4%

average mortality in one heat wave), with some populations exhibiting near or complete

mortality [44,49]. Despite the threatened nature of these particular populations, annual *M.*

guttatus continues thrives in other parts of its large range and is not in danger of extinction.

These past experiments, along with the substantial genomic resources of this model system,

create an ideal scenario to evaluate the impact of assisted gene flow within threatened

populations.

We leverage twelve central Oregon Cascade populations with long-term demographic data to

conduct our landscape-scale AGF experiment (Fig. 1, Table S1). Populations were clustered

geographically into four blocks of three populations with all populations contained within a

25km^2 region (Blocks A-D below). Baseline data collection began in 2018 with yearly surveys of

phenotypes and fecundity in each population. The assisted gene flow experiment began in 2021

with the introduction of individuals from a cross (F$_1$) between two maternal lines, one from a low

elevation population in the Sierra Nevada (BEL) and one from a high elevation population in the

Sierra Nevada (SHL). These populations were chosen because they had high relative fitness in

a nearby Oregon common garden over multiple years [44,48,51]. Within each of our

experimental blocks, we assigned populations to one of three experimental treatments: control,

seeds, or seedlings. In the control populations, we dispersed 200 seeds native to that

population. These seeds were pooled from as many different maternal lines as possible from

the baseline collections (2018-2020) described above. In the seed treatment populations, we

dispersed 200 seeds of the F_1 source individuals uniformly across the population. In the

seedling treatment populations, we planted 19-20 seedlings of the F_1 source individuals along a

transect extending across the center of target populations. We use field surveys and collections

before and after assisted gene flow to evaluate the fitness consequences and underlying

genetic and phenotypic mechanisms of AGF.

Results

Fitness consequences of assisted gene flow

Assisted gene flow was associated with modest differences in fitness across the target

populations, but the effects differed in direction and magnitude when evaluating different fitness

metrics and different life stages of the introduced individuals. Floral abundance increased by

25% post-AGF in populations where California (CA) seeds were introduced and 13.2% post-

AGF in populations where CA seedlings were introduced, but only increased by 4.6% in control

populations, relative to pre-AGF levels (AGF:treatment, X^2 = 9.58, p = 0.008; Fig. 2A, Table S2-

3). Seed production did not change post-AGF in target populations relative to control

populations after conducting assisted gene flow (AGF:treatment, X^2 = 3.0, p = 0.22). Instead,

there were moderate increases in seed production on average across all populations post-AGF,

relative to pre-AGF production (28-43%; Fig. 2B). Introduction of novel alleles may also be

expected to increase variation in fitness within target populations. Standard deviation in floral

abundance increased by 26.4% post-AGF in populations where CA seeds were dispersed and

by 3.1% in populations where CA seedlings were planted (Fig. S2A). The standard deviation in

seed production decreased by 19.3% in populations where CA seeds were dispersed and

increased by 43.0% in populations where CA seedlings were planted (Fig. S2B). These results

suggest that assisted gene flow is either weakly effective across all populations or only effective

in some of the target populations.

Target populations exhibited substantial heterogeneity in their responses to AGF. Assisted gene

flow-related increases in flower abundance were primarily driven by two of the four experimental

blocks (Block A and Block C; Fig. 2C, Table S4-5). Conducting AGF via seeds increased floral

abundance post-AGF by 62.7% (Block A) and 24.8% (Block C), and conducting AGF via

seedlings increased floral abundance by 39.7% (Block A) and 23.7% (Block C) (Block A:

AGF:treatment, $X^2 = 66.8$, p = 3.1 x 10^{-15}; Block C: AGF:treatment, $X^2 = 8.4$, p = 0.014). Similar

trends were observed in these experimental blocks regarding seed production (Block A:

AGF:treatment, $X^2 = 13.4$, p = 0.001; Block C: AGF:treatment, $X^2 = 19.1$, p = 6.8 x 10^{-5}; Fig.

2D). However, AGF resulted in decreases in both floral abundance and seed production post-

AGF in a third block relative to control populations (Block D; floral abundance: AGF:treatment,

$X^2 = 11.2$, p = 0.004, seed production: AGF:treatment, $X^2 = 12.5$, p = 0.002). Differences in

responses across populations likely depends both on stochastic (e.g., gene flow between plants

within populations) and deterministic factors (e.g., seed inviability and the magnitude of early

season selection on hybrid individuals).

We observed changes in both the magnitude and direction of fitness responses to AGF in the

years following introduction (Fig. S3-S5, Table S6-7). There was a statistically significant

increase in floral abundance in AGF treatments in the year following AGF implementation (2022:

AGF:treatment, X^2 = 9.9, p = 0.007) but not in 2023. Seed production exhibited similar

heterogeneity between years with greater increases in populations receiving AGF in 2022 and

increases in control populations in 2023. There were also differences between populations in

the degree of temporal heterogeneity in fitness. For instance, in Block A, introducing CA seeds

increased floral abundance most prominently in 2022, but introducing seedlings into the paired

population increased floral abundance more prominently in 2023. These results highlight that

temporal heterogeneity in selection will have a strong impact on AGF results, but also that some

populations exhibit consistent increases in fitness across multiple years.

Genetic evidence for introgression into AGF-targeted populations.

To validate whether the fitness consequences above were consistent with introgression of

California alleles into target populations, we grew seeds from each year and population in the

UL-Lafayette greenhouse and sequenced whole genomes to an average of 10.9x coverage for

872 individuals. Each population and year combination was represented by 5-27 individuals

(Total N = 872; mean = 14.53; SD = 5.2; Table S7). We also sequenced 7 F_1 lines and 20 plants

from the low elevation source population. We examined two lines of evidence with this dataset

for evidence of introgression: introgression of private alleles from California and whole genome

shifts toward California-like ancestry.

We find evidence of introgression of private alleles into two of the four experimental blocks. In

the experimental block with the greatest fitness increases after conducting AGF (Block A), we

observed that AGF increased the number of CA private alleles that were successfully

incorporated relative to the control population (treatment: X^2 = 616.7, p = 2.2 x 10^{-16}; Fig. 3A,

Table S8). Likewise, there is an increase in the number of CA private alleles in the experimental

block with the greatest fitness decreases after conducting assisted gene flow (Block D:

treatment: $X^2 = 76.151$, p = 2.2 x 10^{-16}). However, there is no evidence of introgression from

private alleles in the other two experiment blocks – including Block C that had increased fitness

in AGF treatment relative to control populations. The lack of evidence for genome-wide

introgression does not necessarily mean that there were not source alleles introduced, but does

indicate that introgression was, at best, rare. In populations with evidence of introgression from

source populations, there was no obvious pattern in the distribution of introgression in private

alleles across the genome and all private alleles were found at relatively low frequencies in both

2021 and 2022 (Fig. 3B, Fig. S6). Notably, introgression was higher in 2021 than in 2022 and

different private alleles introgressed in 2021 and 2022 as only 5.86% of loci have CA alleles

present in both years (Table S9). These results confirm that introgression of CA alleles did

occur in multiple experimental blocks, and non-local alleles remain at relatively low frequencies

in the first years following introduction.

Introgression of CA alleles should also cause AGF target populations to become more

genetically similar to CA parents and F_1 lines (Fig. S7). Genetic principal component analysis

conducted with 4-fold degenerate sites suggest that there is substantial genetic divergence

between source populations (CA parents/F_1) and native Oregon populations, represented by

PC1 (48.5% of variation). Only one experimental block (Block A) has evidence of a shift toward

CA/F_1 plants along PC1 (AGF:treatment, $X^2 = 18.3$, p = 0.0001). Both introducing seeds and

introducing seedlings were effective in increasing CA ancestry in this block. However, the

magnitude of the shift toward CA ancestry is small and all plants are much closer to Oregon

ancestry than California ancestry. This suggests that very low genome-wide levels of

introgression can create substantial fitness impacts.

Phenotype mechanisms underlying AGF fitness responses

Changes in fitness due to the introgression of CA alleles suggests that there should be some

underlying change in phenotype that links genetic variation to population performance. We

conducted a greenhouse resurrection experiment growing maternal lines collected from all

years (2018-2022; Table S10) and populations to determine how phenological, morphological,

and physiological phenotypes changed in populations that received AGF relative to control

populations. Both flowering time and number of trichomes decreased across populations with

introduced CA F_1 individuals relative to control populations (flowering time: $X^2 = 6.74$, p = 0.03;

trichomes: $X^2 = 6.8$, p = 0.03; Fig. 4AB, Fig. S8-9, Table S11). AGF-driven phenotypic

differences were stronger for populations receiving seeds rather than seedlings. Despite this

general pattern, there was heterogeneity in phenotype responses across the four experimental

blocks. In the population block with evidence of introgression and fitness increases (Block A),

flowering time decreased by 6.8 and 3.1 days on average in treatments introducing seeds and

seedlings compared to an increase of 0.88 days in the control treatment (Fig. 4C). Trichomes

decrease in a similar fashion (Fig. 4D). However, in the experimental block with evidence of

introgression and decreased fitness (Block D), flowering time increased by 4.3 and 0.8 days on

average in treatments introducing seeds and seedlings (Fig. 4E). Trichomes were not

significantly altered in this block (Fig. 4F). These results indicate that phenotypic evolution

occurred in the same target populations that had introgression of CA alleles and changes in

fitness.

We hypothesize that phenotypic evolution in AGF-target population populations should be in the

direction of source California populations. Previous greenhouse common garden studies

suggest both California parent populations flower more quickly on average than any of our 12

central Oregon focal populations (30.7-35 mean days following germination; Kooyers et al.

2015) and have high levels of variation in trichome production (range: 6-250 trichomes; Kooyers

et al. 2015). There are individuals in every focal Oregon population that flower as quickly as

California populations. However, populations in Block A were 3 of the 4 slowest flowering

central Oregon populations prior to AGF and evolve earlier flowering in the direction of the

California parent lines. Populations in block D, particularly BMP, had relatively faster times to

flowering prior to AGF that were more closely resembled California source populations. Thus,

evolution of later flowering within this block would not have been predicted, but could be

generated if introgression broke up co-adapted combinations of alleles. We cannot make the

same predictions for the evolution in trichome number as the source populations are too

variable. Taken together, these results suggest that the introgression of California alleles altered

flowering time and created variation in fitness following AGF.

Discussion

We find that assisted gene flow has the potential to increase fitness of target populations within

only a few generations. Theoretical models simulating genetic or evolutionary rescue indicate

that introduction of foreign alleles often lead to initial maladaptation due to outbreeding

depression followed by increases in fitness as adaptive alleles are selected and maladaptive

alleles purged from the population [36,37]. We find two blocks that had evidence of

introgression and that also exhibited changes in fitness post AGF. Both target populations in

one block with introgression exhibited slight evidence of maladaptation or outbreeding

depression relative to control populations in the first three generations, consistent with

theoretical expectations (Fig. 2). Surveys over the next decade will determine whether

introgression eventually increases population fitness in this initially maladapted block.

However, target populations in the other experimental block with introgression had substantial

fitness gains within these first generations (Fig. 2). There are multiple non-mutually exclusive

hypotheses explaining this rapid fitness increase. One explanation for fitness gains is hybrid

vigor or heterosis, where outbreeding leads to increased heterozygosity and an increase in

fitness [52]. However, since allele frequencies associated with the introgression of private

alleles are low (Fig. 3), this is unlikely to be the driving mechanism. Also unlikely is a scenario

where the influx of genetic diversity rescues the population from inbreeding or high genetic load

as target populations have high levels of genetic diversity [53]. The most likely possibility is that

target populations were poorly adapted to their local conditions and AGF introduced novel

genetic variants that shifted phenotype means and allowed transgressive phenotypes or novel

phenotypic combinations to emerge. While our greenhouse work establishes post-AGF shifts

toward earlier flowering and lower trichomes relative to control populations, field common

gardens with manipulations to key climate variables (e.g. growing season length) are necessary

to dissect whether the observed phenotypic evolution could underlie our observed fitness gains.

One potential caveat to our results is that fitness differences pre- and post-AGF could be

associated with intrinsic differences between the individual populations within a block, rather the

AGF treatments. Indeed, if a control population were to have declining fitness following AGF

due to any random reason, both treatment populations would appear to have enhanced fitness.

We believe our results are robust, particularly for increase in fitness associated with AGF in

Block A as well as the decrease in fitness in Block D, for three reasons. First, we control for

intrinsic differences among population (and years) within our models using random effects.

Second, populations *within* blocks are geographically paired and nearly indistinguishable in

population structure analysis (with the exception of low levels of differentiation in Block D).

Finally, the same target populations that exhibit fitness differences in certain years also exhibit

phenotypic evolution and show the strongest signals of introgression (Block A &D). Thus,

although our results are necessarily correlative because of our landscape-scale experimental

design, these results strongly suggest that AGF caused introgression, produced phenotypic

evolution, and altered fitness of focal populations.

Climatic variation impacts introgression and fitness gains associated with AGF

Our results provide key insights into the expectations for assisted gene flow within complex

environments. We find variation in the success of assisted gene flow across both space and

time (Fig. 2, Fig S4). Our finding that two experimental blocks had evidence of introgression

from assisted gene flow while two other blocks did not could be explained by both deterministic

and stochastic factors. Lack of introgression could stem from stochastic factors. For instance,

relatively few seeds or seedlings were introduced and introgression is dependent on pollination

between native and introduced individuals.

However, success of introgression and associated fitness differences were likely impacted by

the climatic differences between populations. While populations within each experimental block

were located at the same elevation, experimental blocks were located over an elevation

gradient of ~500m – creating substantial differences in growing season timing and both abiotic

and biotic conditions during the growing season. The lowest elevation experimental block was

the block with evidence of introgression, phenotypic evolution and fitness gains. These

populations have an earlier growing season start date and a longer growth season than higher

elevation populations – conditions that are most similar to the historical conditions with the

source California populations [50]. Notably, low elevation populations also had more delayed

times to flowering than high elevation populations at the beginning of the experiment. An

infusion of novel alleles promoting earlier flowering, such as those from the early flowering

California source plants, would be most likely to be effective in these populations and could be

detrimental in target populations that already flower earlier than the source populations, i.e., are

already better adapted to changing conditions. Conditions for germination of seeds may also

have differed between populations and this is likely a key barrier for the success of assisted

gene flow [54]. Our results suggest, broadly, that the success of AGF is dependent on whether

local populations are already well adapted to the environmental conditions during the

translocation years.

The difference in introgression and fitness in different years following AGF also suggests that

climate abnormalities in the initial years of AGF act as an important filter for AGF success.

Patterns of introgression differ substantially between years. Most introgression took place during

the year that seeds and seedling were introduced (2021). This should be expected as seeds

germinate readily in *M. guttatus* and we planted a number of seedlings that were ready to flower

(Fig. S1). However, there was also a substantial heat dome event during the 2021 growing

season [55] that had devastating consequences on many target populations [49] and prevented

many of the source seedlings from producing seed (Table S12). We also expected evidence of

introgression in subsequent years from seeds that did not germinate the first year and from

seeds which were generated as hybrids between source and target populations. There was far

more limited evidence of introgression of private alleles relative to control populations in 2022

(Fig. 3). It is notable that we judge levels of introgression relative to control populations as the

likely pollinators (bees) are capable of flying between control and treatment populations within

an experimental block (1-2km; 55). Indeed, alleles that we inferred as private to source

populations appear in every control population (Fig. 3). This is primarily due to our inference of

private alleles as we likely did not detect some rare alleles in Oregon in the sequencing of 2018-

20 Oregon populations. However, tracking source population alleles that escape treatment

populations is important for determining off-target impacts of AGF [57].

Limited introgression of private alleles genome-wide does not exclude the possibility that there

may be introgression in key genomic regions that could have impacted phenotypes. On the

contrary, some populations with relatively limited introgression of private alleles in 2022 had

strongly increased floral abundance relative to control populations in the same year (e.g. Block

A seeds treatment). Future studies scanning the genome for allele frequency change will detect

the key regions underlying shifts in fitness. However, temporal variance in relative fitness in

AGF populations relative to control populations likely indicates that temporal variation in climate

plays a key role in the outcome of AGF. High elevation *M. guttatus* populations in the central

Oregon Cascades are known for strong fluctuating selection maintaining high genetic and

phenotypic diversity within populations [53,58–60]. Notably, one of the key phenotypes

undergoing fluctuating selection is flowering time with year-to-year variation driven by

differences in growing season start date and duration [59]. An important consideration for AGF

is that introduced alleles may be adaptive in some years but not in others. For instance, we

worry that introduced alleles that increase in frequency during the relatively benign years (i.e.

2022 growing season) may be maladaptive and cause populations to be less resilient during

future heatwaves. Moreover, strong fluctuations during the first years after AGF could either

enhance or prevent source population alleles from incorporating into target populations

depending on whether the fluctuation makes the environment more similar to the historic source

environment. The relative impacts of fluctuating selection during AGF require additional

theoretical work.

Assessing key choices for implementing AGF

Our results suggest that the logistics of assisted gene flow are key to the end result. Current recommendations suggest introducing 5-20% of the target population census size from source populations [21,41]. Our experiment introduced ~ 0.5 - 2 % of the census size depending on the population and whether we introduced seeds or seedlings. Population census sizes can differ dramatically between years, with limited individuals flowering during years with early growing season start dates and abnormally hot conditions [44,49]. This decision likely limited the initial amount of introgression in target populations. Notably, even with a limited number of individuals introduced, we observed both introgression and fitness impacts within half of our target populations. Despite this success, we suggest introducing higher numbers of individuals to ensure more rapid incorporation into the genome (i.e., 10-20% of census size).

Another key choice is deciding how to introduce source populations. Our experiment examines a single introduction of seeds vs. seedlings. In experimental blocks with clear evidence of introgression, both seed and seedling treatments had effects, but the introduction of seeds increased introgression and led to larger magnitude fitness effects. The introduction of seeds was likely more effective for three reasons: first, we introduced more seeds than planted seedlings; second, 77% of seedlings died following introduction due to a substantial heatwave and the rest produced limited fruits (1.17 ± 0.99 SD fruits); third, seeds may not have germinated within the first year and could have provided a reservoir capable of introducing alleles in multiple growing seasons. Thus, we do not observe the potential downside of introducing seeds – that they must overcome seed dormancy requirements and novel establishment conditions associated with non-native sites [54]. Our results suggest that single introductions are less likely to be effective than multiple introductions and that introducing seeds may have longer lasting effects in unpredictable climates.

Our experiment also included a non-standard choice for applied assisted gene flow efforts. We

introduced germplasm resulting from a cross of two maternal lines sourced from two

populations. Our choice was justified by the success of these two lines and populations in a

nearby common garden experiment [44,48], evidence of inbreeding depression in nearby

natural populations [61], and the need for a genetically-tractable system. Yet, most species with

the need for AGF will not have the large history of evolutionary ecology and genetics studies

that our study system possesses. Source populations for AGF are typically selected via climate

transfer zones or seedlot selection tools based on historical climates and contemporary

conditions [62]. Experimental and simulation studies are needed to better assess how to choose

source populations – phenotypic and genomic data could provide rapid assessment of potential

source populations that could complement or even replace climate data. In the absence of data,

including both native and climate-adapted germplasm in introductions either via seed mixes or

through hybridization of populations can allow natural selection to incorporate ecologically-

important variants.

Conclusions

Our study provides cautious support for utilizing assisted gene flow to preserve natural

populations experiencing the detrimental effects of climate change, consistent with the growing

recommendations for conducting AGF in a variety of maladapted populations [16,17,63,64]. We

find evidence of the beginning stages of evolutionary rescue despite relatively limited genome-

wide patterns of introgression, limiting potential concerns regarding outbreeding depression or

gene swamping. However, attention to organismal biology of the threatened species and

logistical details is clearly key to the success or failure of AGF and we encourage responsible

evidence-based decision making.

Materials and Methods

We conducted a landscape-scale manipulative experiment involving 12 target populations within

a 25km^2 region of the Central Oregon Cascade Mountains. Populations were geographically

clustered into four groups of three populations ('Blocks') with 0.5 - 2km between populations

within blocks (Fig. 1). Elevations were similar within blocks but varied between blocks (1087-

1567 m; Table S1). We selected two source populations from the foothills of the Sierra Nevada

in central California (Fig. 1; Table S1). These populations were chosen because they had higher

fitness than the native populations from the Oregon Cascades in nearby common garden

experiments [48,65]. A single maternal line from the two source populations (BEL, SHL) were

crossed together in bulk to produce the F_1 seeds used for AGF. In each block, populations were

designated as control, seed or seedling treatment. In the control treatment populations, 200

seeds previously collected from the native population were uniformly sown throughout the

population (20+ maternal lines). In the seed treatment populations, we dispersed 200 F_1 source

seeds uniformly across the population. In the seedling treatment populations, we planted 19-20

14-day old seedlings grown from the F_1 seeds along a transect extending across the center of

target populations. The difference in sample size reflects germination and establishment likely

provide a substantial demographic filter within these populations. Treatments occurred as soon

as possible after snowmelt to correspond to population phenology. For seedling growth

conditions prior to planting, see extended methods. We recorded survival of translocated

seedlings weekly, and counted number of fruits produced after senescence in order to assess

the potential for introgression of new alleles in these populations; only 23% of seedlings

survived to flower, and those individuals produced an average of 1.17 ± 0.99 SD fruits (Table

S11). However, we do note that a single fruit can contain hundreds of seeds, and natural

populations also had very few individuals surviving to reproduction and producing seeds during

these years [49].

Evaluating AGF impacts on fitness within natural populations

We assessed fitness at the end of the growing season in each population in each year from

2018-2023 (previously reported in [44,49]. Plants were collected along two 7.5 m transects

spanning the population. From 2018-2020, one individual was collected every 15 cm along each

transect, totaling 100 individuals/population. From 2021-2023, one individual was collected

every 30 cm along each transect, totaling 50 individuals/population. We counted all flowers and

seeds for each individual as proxies of fitness. We note that assessing population demography

is methodically challenging within this species. Central Oregon populations exhibit sweepstakes

reproduction where an individual seed has a low chance of producing seeds in the next

generation. Specifically, seeds must land in ephemerally wet patches within an extremely

heterogeneous seepy meadow. Thus, fecundity (i.e., increasing the number of chances at

reproductive success) is likely strongly associated with population growth rate, but our work

cannot assess whether AGF increases population growth rates above replacement level.

We evaluated whether AGF increased fecundity of the target populations within a general linear

mixed model framework (LMM). We conducted univariate LMM to assess the effects of AGF

(pre/post), Treatment (control/seeds/seedlings), and the interaction between AGF:Treatment on

both metrics of fitness (flower abundance and seed production). Year and population were

random effects in each model. Floral abundance and seed production were log+1-transformed

to better fit model assumptions. To determine whether assisted gene flow increases the

variance in fitness, we conduced similar univariate LMMs (without population as a random

factor) using the standard deviation of floral abundance and seed production between

individuals within a population as response variables. Standard deviation of flower abundance

was log+1-transformed. Analyses were performed using the *lme4* v 1.1-35.5 package in R v

4.4.1 [66]. Statistical significance of LMM was assessed via ANOVA with type III sum of squares

using the Anova() function in the *car* v 3.1-2 package [67]. We note that interpretation of models

differed for each fitness variable as flower abundances could logistically not be affected by AGF

in 2021 as plants sampled along transects were almost certainly native Oregon plants (i.e., not

from seed germinated from source populations); thus, floral abundance in 2021 is considered

pre-AGF in models. However, seed production could have been impacted in 2021 through

pollen transfer from introduced CA seeds/seedlings and can be considered post-AGF in models.

To better dissect the impacts of AGF in the experimental manipulation, we evaluated variation

among blocks and between post-AGF years. We examined differences among different

experimental blocks by subsetting the data into the four blocks and repeating the above

univariate LMMs for each block. Because each treatment was only represented by one

population, population was removed as a random factor for these models. We assessed

heterogeneity in AGF-related fitness variation across post-AGF years by subsetting data into

each post-AGF year and retaining all baseline years. In these models, year was removed and

population was included as a random variable.

Validating AGF introgression via genomic analysis.

We germinated seeds from 24-30 maternal lines/population ~2 months after collection each

year in the UL-Lafayette Greenhouse (*Refresher Generation Growth Conditions* in extended

methods). We also included 7 individuals from one parental population (BEL) and 20 F1

individuals grown in the 2021 common garden. Seeds from the other parent line (SHL) did not

germinate. One individual from each maternal line had buds or leaves collected for genetics and

this individual was selfed for use in the resurrection experiment described below. DNA was

extracted using a modified CTAB extraction [68] and diluted to 3 ng/μL for library construction.

Libraries for whole genome sequencing were constructed using a customized dual-index

Nextera library construction protocol [69]. Each population and year combination was

represented by 5-27 individuals (Total N = 872; mean = 14.53; SD = 5.2; Table S7). Poor fitness

in the field that limited collections as well as reduced germination led to the reduced sample size

of some populations in some years. Post-barcoding, we pooled either 48 or 96 individual

equimolar libraries and sequenced each pool on a single lane on either a HiSeqX or NovoseqX

Plus platforms (150bp paired end reads; Novogene, Davis, CA). Batch effects due to either

sequencing platform or lane were minimal.

We constructed a custom bioinformatics pipeline for our population genomics study

(https://github.com/donnasaurrus/AGF). Initial read quality was evaluated using fastQC [70].

High-quality reads were aligned a *Mimulus guttatus* reference genome from a nearby population

(IM62 v3: https://phytozome-next.jgi.doe.gov/info/Mguttatusvar_IM62_v3_1) utilizing bwa-mem

with default parameters [71] and converted to bam files using SAMtools [72]. Read groups and

duplicates were marked using PicardTools, and SNPs were called using bcftools [73]. We

filtered for variant quality, strand bias, and mapping quality using GATK [QD < 2.0 ll FS > 60.0 ll

MQ < 40.0 ll MQRankSum < -12.5] [74]. We used VCFtools [75] to filter retaining biallelic SNPs

with minimum quality score of 500, and removing any SNPs with >25% missing data. This SNP

dataset included 18,118,076 SNPs. Mean coverage and error rates of our sequence data (.bam

files) were assessed using Qualimap [76,77]. Mean coverage depth was 10.9x (SD = 7.7), and

mean error rate was 2.7% (SD = 0.39%) (Table S13).

We first evaluated introgression via AGF into Oregon populations using private alleles. We

filtered the SNP dataset to only include sites where both reference and alternative alleles were

at a frequency > 0.3 in BEL and F_1 individuals *and* were monomorphic in all populations in

Oregon prior to conducting AGF (2018-2020). We note that these alleles may not necessarily be

truly private as some rare alleles in Oregon populations may escape detection in the three

baseline years, either due to the presence of a seed bank or a finite sample size. This dataset

consisted of 15,691 SNPs distributed genome-wide. We evaluated whether there was

introgression in target populations that received AGF from source populations compared to

control populations. We transformed allele frequencies into a binomial variable, where loci with

a CA private allele frequency of 0 were coded as 0, and any non-zero allele frequencies were

coded as 1. We conducted a generalized linear mixed model with a binomial distribution and a

logit link function using the *glmmTMB* v 1.1.9 package in R v 4.4.1 [78], with treatment (levels:

Control, Seed, Seedling) included as a fixed factor, and with year and population included as

random variables. To assess differential responses between blocks, we repeated this model,

subsetting data into individual experimental blocks and removing population as a random

variable. Statistical significance was assessed as in above GLMs for fitness.

We evaluated the extent that introgression shifted genome-wide genetic diversity of Oregon

populations toward California source populations using a principal component analysis that

included Oregon populations from all years as well as individuals from California parental and F_1

lines. We filtered SNPs to include only 4-fold degenerate sites and only used individuals with <

20% missing data (N = 803). This dataset included 1,056,971 SNPs. A PCA was conducted

using --pca in *plink* v1.9 [79]. PC1 separated Oregon from California populations (Fig. S7). We

examined shifts in PC1 toward CA populations following AGF using univariate GLMs identical

those used above for fitness except with PC1 and PC2 as response variables. In these models,

AGF (pre/post), treatment (control/seeds/seedlings), and the interaction between

AGF:Tntroduction were fixed effects while population and year were included as random effects.

Identifying phenotypic shifts following AGF

We conducted a greenhouse resurrection experiment to examine how phenotypes shift

following AGF. We germinated seed from 988 maternal lines stemming from every population

and year from 2018 - 2022 (mean ± SD: 17.0 ± 6.1 lines per population per year, Table S9).

These seeds were selfed lines from the refresher generation(s) described above. Germination

and growth conditions are described in extended methods (*Resurrection Experiment Growth*

Conditions). We surveyed for germination and flowering daily and calculated flowering time as

the number of days between germination and opening of the first flower. At flowering, we

identified the node of the first flower, counted the number of leaves and branches, and

measured plant height, corolla width, corolla length, corolla height, and stem diameter. We also

collected the larger single 2nd true leaf to measure leaf area, specific leaf area, number of

trichomes, and relative water content. Further details in phenotypes can be found in extended

methods. There were strong correlations among several of the phenotypes (Table S14) so we

choose to only include a subset of phenotypes in analyses including flowering time, plant height,

leaf number, corolla width and number of trichomes.

We used univariate GLMMs to evaluate potential shifts in phenotypes following AGF treatment.

As in above GLMMs, AGF (pre/post), Treatment (control/seeds/seedlings), and the interaction

between AGF:Treatment were fixed effects while population and year were included as random

effects. Distributions, links, and transformations were chosen to maximize model fit. The GLMM

for flowering time had a Poisson distribution with a log link. All other variables were log-

transformed and had Gaussian distributions with identity links. Statistical significance of fixed

factors was assessed via ANOVA with type III sum of squares using the Anova() function in the

car v3.1-2 R package [67]. Changes in phenotype due to AGF would create significant

AGF:Treatment interactions with shifts likely toward CA phenotypic means. To better examine

how phenotypic change was occurring within experimental blocks, we subset the dataset by

block and ran identical models for each block without population as a random factor.

Acknowledgements

The authors thank Josh FitzPatrick, Braden Doucet, Luke Rabalais, Elissa Harb, Jace Segura,

Gabbie Dietel, Lana Gaspard, Noah Richard, Emily Bollich, Haley St. Martin, Laura McDonald,

Anna Scharnagl, and Benjamin LeBlanc who helped with field, greenhouse and lab work

associated with this project. This work was logistically supported by H.J. Andrews Experimental

Forest and permitted through the Willamette Forest District of the United States Forest Service.

Funding for this research came from University of Louisiana, Lafayette and NSF grants DEB-

2045643 and IOS-2222466 to N.J.K.

References

1. Exposito-Alonso M, Booker TR, Czech L, Gillespie L, Hateley S, Kyriazis CC, et al. Genetic diversity loss in the Anthropocene. Science. 2022;377: 1431–1435. doi:10.1126/science.abn5642

2. Calvin K, Dasgupta D, Krinner G, Mukherji A, Thorne PW, Trisos C, et al. Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. First. Geneva, Switzerland: Intergovernmental Panel on Climate Change (IPCC); 2023 July. doi:10.59327/IPCC/AR6-9789291691647

3. Anderson JT, DeMarche ML, Denney DA, Breckheimer I, Santangelo J, Wadgymar SM. Adaptation and gene flow are insufficient to rescue a montane plant under climate change. Science. 2025;388: 525–531. doi:10.1126/science.adr1010

4. Campbell DR. Early snowmelt projected to cause population decline in a subalpine plant. Proc Natl Acad Sci USA. 2019;116: 12901–12906. doi:10.1073/pnas.1820096116

5. Kooyers NJ, Anderson JT, Angert AL, Avolio ML, Campbell DR, Exposito-Alonso M, et al. Responses to climate change – insights and limitations from herbaceous plant model species. New Phytologist. 2025;248: 461–493. doi:10.1111/nph.70468

6. Krushelnycky PD, Loope LL, Giambelluca TW, Starr F, Starr K, Drake DR, et al. Climate-associated population declines reverse recovery and threaten future of an iconic high-elevation plant. Global Change Biology. 2013;19: 911–922. doi:10.1111/gcb.12111

7. Reed PB, Bridgham SD, Pfeifer-Meister LE, DeMarche ML, Johnson BR, Roy BA, et al. Climate warming threatens the persistence of a community of disturbance-adapted native annual plants. Ecology. 2021;102: e03464. doi:10.1002/ecy.3464

8. Sheth SN, Angert AL. Demographic compensation does not rescue populations at a trailing range edge. Proc Natl Acad Sci USA. 2018;115: 2413–2418. doi:10.1073/pnas.1715899115

9. Iknayan KJ, Beissinger SR. Collapse of a desert bird community over the past century driven by climate change. Proc Natl Acad Sci USA. 2018;115: 8597–8602. doi:10.1073/pnas.1805123115

10. Smith MD, La Pierre KJ, Collins SL, Knapp AK, Gross KL, Barrett JE, et al. Global environmental change and the nature of aboveground net primary productivity responses: insights from long-term experiments. Oecologia. 2015;177: 935–947. doi:10.1007/s00442-015-3230-9

11. Bartley TJ, McCann KS, Bieg C, Cazelles K, Granados M, Guzzo MM, et al. Food web rewiring in a changing world. Nat Ecol Evol. 2019;3: 345–354. doi:10.1038/s41559-018-0772-3

12. Vahsen ML, Kleiner HS, Kodak H, Summers JL, Vahsen WL, Blum MJ, et al. Complex eco-evolutionary responses of a foundational coastal marsh plant to global change. New Phytologist. 2023;240: 2121–2136. doi:10.1111/nph.19117

13.   Zoller L, Bennett J, Knight TM. Plant–pollinator network change across a century in the
subarctic. Nat Ecol Evol. 2023;7: 102–112. doi:10.1038/s41559-022-01928-3

14.   Raubenheimer SL, Zheng L, Stefanski A, Reich PB. Climate-change-driven shifts in C_3 and
C_4 grass distributions and leaf traits could lead to changes in community-level flammability.
American J of Botany. 2025; e70081. doi:10.1002/ajb2.70081

15.   Aitken SN, Whitlock MC. Assisted Gene Flow to Facilitate Local Adaptation to Climate
Change. Annual Review of Ecology, Evolution, and Systematics. 2013;44: 367–388.
doi:10.1146/annurev-ecolsys-110512-135747

16.   Browne L, Wright JW, Fitz-Gibbon S, Gugger PF, Sork VL. Adaptational lag to temperature
in valley oak (Quercus lobata) can be mitigated by genome-informed assisted gene flow.
Proceedings of the National Academy of Sciences. 2019;116: 25179–25185.
doi:10.1073/pnas.1908771116

17.   Grummer JA, Booker TR, Matthey-Doret R, Nietlisbach P, Thomaz AT, Whitlock MC. The
immediate costs and long-term benefits of assisted gene flow in large populations.
Conservation Biology. 2022;36: e13911. doi:10.1111/cobi.13911

18.   Pregler KC, Obedzinski M, Gilbert-Horvath EA, White B, Carlson SM, Garza JC. Assisted
gene flow from outcrossing shows the potential for genetic rescue in an endangered
salmon population. Conservation Letters. 2023;16: e12934. doi:10.1111/conl.12934

19.   Fredrickson RJ, Siminski P, Woolf M, Hedrick PW. Genetic rescue and inbreeding
depression in Mexican wolves. Proceedings of the Royal Society B: Biological Sciences.
2007;274: 2365–2371. doi:10.1098/rspb.2007.0785

20.   Hagedorn M, Page CA, O'Neil KL, Flores DM, Tichy L, Conn T, et al. Assisted gene flow
using cryopreserved sperm in critically endangered coral. Proceedings of the National
Academy of Sciences. 2021;118: e2110559118. doi:10.1073/pnas.2110559118

21.   Hedrick PW. Gene Flow and Genetic Restoration: The Florida Panther as a Case Study.
Conservation Biology. 1995;9: 996–1007.

22.   Weeks AR, Sgro CM, Young AG, Frankham R, Mitchell NJ, Miller KA, et al. Assessing the
benefits and risks of translocations in changing environments: a genetic perspective:
Translocations in changing environments. Evolutionary Applications. 2011;4: 709–725.
doi:10.1111/j.1752-4571.2011.00192.x

23.   Todesco M, Pascual MA, Owens GL, Ostevik KL, Moyers BT, Hübner S, et al.
Hybridization and extinction. Evolutionary Applications. 2016;9: 892–908.
doi:10.1111/eva.12367

24.   Ricciardi A, Simberloff D. Assisted colonization is not a viable conservation strategy.
Trends in Ecology & Evolution. 2009;24: 248–253. doi:10.1016/j.tree.2008.12.006

25.   Edmands S. Between a rock and a hard place: evaluating the relative risks of inbreeding
and outbreeding for conservation and management. Molecular Ecology. 2007;16: 463–
475. doi:10.1111/j.1365-294X.2006.03148.x

26.   Frank A, Howe GT, Sperisen C, Brang P, Clair JBSt, Schmatz DR, et al. Risk of genetic
671   maladaptation due to climate change in three major European tree species. Global Change
672   Biology. 2017;23: 5358–5371. doi:10.1111/gcb.13802

673   27.   Price MV, Waser NM. Pollen dispersal and optimal outcrossing in Delphinium nelsoni.
674   Nature. 1979;277: 294–297. doi:10.1038/277294a0

675   28.   Ronce O, Kirkpatrick M. When sources become sinks: Migrational meltdown in
676   heterogeneous habitats. Evolution. 2001;55: 1520–1531. doi:10.1111/j.0014-
677   3820.2001.tb00672.x

678   29.   Tallmon DA, Luikart G, Waples RS. The alluring simplicity and complex reality of genetic
679   rescue. Trends in Ecology & Evolution. 2004;19: 489–496. doi:10.1016/j.tree.2004.07.003

680   30.   Waller DM. Genetic rescue: a safe or risky bet? Molecular Ecology. 2015;24: 2595–2597.
681   doi:10.1111/mec.13220

682   31.   Weeks AR, Sgro CM, Young AG, Frankham R, Mitchell NJ, Miller KA, et al. Assessing the
683   benefits and risks of translocations in changing environments: a genetic perspective.
684   Evolutionary Applications. 2011;4: 709–725. doi:10.1111/j.1752-4571.2011.00192.x

685   32.   Waser NM, Price MV. Crossing-distance effects in *Delphinium nelsonii* : Outbreeding and
686   inbreeding depression in progeny fitness. Evolution. 1994;48: 842–852.
687   doi:10.1111/j.1558-5646.1994.tb01366.x

688   33.   Barton NH. The effects of linkage and density-dependent regulation on gene flow.
689   Heredity. 1986;57: 415–426. doi:10.1038/hdy.1986.142

690   34.   Lenormand T. Gene flow and the limits to natural selection. Trends in Ecology & Evolution.
691   2002;17: 183–189. doi:10.1016/S0169-5347(02)02497-7

692   35.   Gomulkiewicz R, Holt RD. When does evolution by natural selection prevent extinction?
693   Evolution. 1995;49: 201–207. doi:10.1111/j.1558-5646.1995.tb05971.x

694   36.   Orr HA, Unckless RL. The population genetics of evolutionary rescue. Pritchard JK, editor.
695   PLoS Genet. 2014;10: e1004551. doi:10.1371/journal.pgen.1004551

696   37.   Carlson SM, Cunningham CJ, Westley PAH. Evolutionary rescue in a changing world.
697   Trends in Ecology & Evolution. 2014;29: 521–530. doi:10.1016/j.tree.2014.06.005

698   38.   Bell G. Evolutionary Rescue. Annu Rev Ecol Evol Syst. 2017;48: 605–627.
699   doi:10.1146/annurev-ecolsys-110316-023011

700   39.   Frankham R, Ballou JD, Eldridge MDB, Lacy RC, Ralls K, Dudash MR, et al. Predicting the
701   probability of outbreeding depression. Conservation Biology. 2011;25: 465–475.
702   doi:10.1111/j.1523-1739.2011.01662.x

703   40.   Frankham R. Genetic rescue of small inbred populations: Meta-analysis reveals large and
704   consistent benefits of gene flow. Molecular Ecology. 2015;24: 2610–2618.
705   doi:10.1111/mec.13139

41. Grummer JA, Booker TR, Matthey-Doret R, Nietlisbach P, Thomaz AT, Whitlock MC. The
immediate costs and long-term benefits of assisted gene flow in large populations.
Conservation Biology. 2022;36: e13911. doi:10.1111/cobi.13911

42. Pain R. Evolutionary Impacts of Assisted Gene Flow: Fitness Consequences of
Hybridization Along a Geo-Climatic Gradient in an Annual Prairie Legume. ProQuest
Dissertations and Theses. Ph.D., University of Minnesota. 2024. Available:
https://ezproxyprod.ucs.louisiana.edu:2443/login?url=https://www.proquest.com/dissertatio
ns-theses/evolutionary-impacts-assisted-gene-flow-fitness/docview/2967058057/se-
2?accountid=14566

43. Wadgymar SM, Cumming MN, Weis AE. The success of assisted colonization and
assisted gene flow depends on phenology. Global Change Biology. 2015;21: 3786–3799.
doi:10.1111/gcb.12988

44. McDonald LM, Scharnagl A, Turcu AK, Patterson CM, Kooyers NJ. Demographic
consequences of an extreme heat wave are mitigated by spatial heterogeneity in an
annual monkeyflower. Ecology and Evolution. 2023;13: e10397. doi:10.1002/ece3.10397

45. Stewart CN, Halfhill MD, Warwick SI. Transgene introgression from genetically modified
crops to their wild relatives. Nat Rev Genet. 2003;4: 806–817. doi:10.1038/nrg1179

46. Wu CA, Lowry DB, Cooley AM, Wright KM, Lee YW, Willis JH. *Mimulus* is an emerging
model system for the integration of ecological and genomic studies. Heredity. 2008;100:
220–230.

47. Vallejo-Marín M, Friedman J, Twyford AD, Lepais O, Ickert-Bond SM, Streisfeld MA, et al.
Population genomic and historical analysis suggests a global invasion by bridgehead
processes in *Mimulus guttatus*. Commun Biol. 2021;4: 327. doi:10.1038/s42003-021-
01795-x

48. Kooyers NJ, Colicchio JM, Greenlee AB, Patterson E, Handloser NT, Blackman BK.
Lagging adaptation to climate supersedes local adaptation to herbivory in an annual
monkeyflower. American Naturalist. 2019;194: 541–557. doi:10.1086/702312

49. Kooyers NJ, Genung MA, Innes SG, Turcu AK, Hinrichs DM, LeBlanc BJ, et al. Heat
waves decrease fitness and alter maternal provisioning in natural populations of *Mimulus
guttatus*. American Journal of Botany. 2025;112: e70087. doi:10.1002/ajb2.70087

50. Kooyers NJ, Greenlee AB, Colicchio JM, Oh M, Blackman BK. Replicate altitudinal clines
reveal that evolutionary flexibility underlies adaptation to drought stress in annual *Mimulus
guttatus*. New Phytologist. 2015;206: 152–165. doi:10.1111/nph.13153

51. Scharnagl A, Genung MA, Holeski LM, Kooyers NJ. Life history and chemical defense
interact to drive patterns of local adaptation in an annual monkeyflower. Evolution.
2023;77: 370–383. doi:https://doi.org/10.1093/evolut/qpac048

52. Szűcs M, Melbourne BA, Tuff T, Weiss-Lehman C, Hufbauer RA. Genetic and
demographic founder effects have long-term fitness consequences for colonising
populations. Angert A, editor. Ecology Letters. 2017;20: 436–444. doi:10.1111/ele.12743

53.   Puzey JR, Willis JH, Kelly JK. Population structure and local selection yield high genomic
variation in Mimulus guttatus. Molecular Ecology. 2017;26: 519–535.
doi:https://doi.org/10.1111/mec.13922

54.   Kildisheva OA, Dixon KW, Silveira FAO, Chapman T, Di Sacco A, Mondoni A, et al.
Dormancy and germination: making every seed count in restoration. Restoration Ecology.
2020;28. doi:10.1111/rec.13140

55.   White RH, Anderson S, Booth JF, Braich G, Draeger C, Fei C, et al. The unprecedented
Pacific Northwest heatwave of June 2021. Nat Commun. 2023;14: 727.
doi:10.1038/s41467-023-36289-3

56.   Osborne JL, Martin AP, Carreck NL, Swain JL, Knight ME, Goulson D, et al. Bumblebee
flight distances in relation to the forage landscape. Journal of Animal Ecology. 2008;77:
406–415. doi:10.1111/j.1365-2656.2007.01333.x

57.   Pasquet RS, Peltier A, Hufford MB, Oudin E, Saulnier J, Paul L, et al. Long-distance pollen
flow assessment through evaluation of pollinator foraging range suggests transgene
escape distances. Proceedings of the National Academy of Sciences. 2008;105: 13456–
13461. doi:10.1073/pnas.0806040105

58.   Mojica JP, Lee YW, Willis JH, Kelly JK. Spatially and temporally varying selection on
intrapopulation quantitative trait loci for a life history trade-off in *Mimulus guttatus*.
Molecular Ecology. 2012;21: 3718–3728. doi:10.1111/j.1365-294X.2012.05662.x

59.   Troth A, Puzey JR, Kim RS, Willis JH, Kelly JK. Selective trade-offs maintain alleles
underpinning complex trait variation in plants. Science. 2018;361: 475–478.
doi:DOI:%2010.1126/science.aat576

60.   Kelly JK. The genomic scale of fluctuating selection in a natural plant population. Evolution
Letters. 2022;6: 506–521. doi:10.1002/evl3.308

61.   Willis JH. Effects of different levels of inbreeding on fitness components in *Mimulus
guttatus*. Evolution. 1993;47: 864–876. doi:10.1111/j.1558-5646.1993.tb01240.x

62.   St.Clair JB, Richardson BA, Stevenson-Molnar N, Howe GT, Bower AD, Erickson VJ, et al.
Seedlot Selection Tool and Climate-Smart Restoration Tool: Web-based tools for sourcing
seed adapted to future climates. Ecosphere. 2022;13: e4089. doi:10.1002/ecs2.4089

63.   Aitken SN, Bemmels JB. Time to get moving: assisted gene flow of forest trees.
Evolutionary Applications. 2016;9: 271–290. doi:10.1111/eva.12293

64.   Pavlova A, Beheregaray LB, Coleman R, Gilligan D, Harrisson KA, Ingram BA, et al.
Severe consequences of habitat fragmentation on genetic diversity of an endangered
Australian freshwater fish: A call for assisted gene flow. Evolutionary Applications.
2017;10: 531–550. doi:10.1111/eva.12484

65.   McDonald LM, Scharnagl A, Turcu AK, Patterson CM, Kooyers NJ. Demographic
consequences of an extreme heat wave are mitigated by spatial heterogeneity in an
annual monkeyflower. Ecology and Evolution. 2023;13: e10397. doi:10.1002/ece3.10397

66. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using lme4.
arXiv; 2014. Available: http://arxiv.org/abs/1406.5823

67. Fox J, Friendly M, Weisberg S. Hypothesis Tests for Multivariate Linear Models Using the
car Package. The R Journal. 2013;5: 39. doi:10.32614/RJ-2013-004

68. Kelly AJ, Willis JH. Polymorphic microsatellite loci in Mimulus guttatus and related species.
Molecular Ecology. 1998;7: 769–774. doi:10.1046/j.1365-294x.1998.00328.x

69. Therkildsen NO, Palumbi SR. Practical low-coverage genomewide sequencing of
hundreds of individually barcoded samples for population and evolutionary genomics in
nonmodel species. Molecular Ecology Resources. 2017;17: 194–208. doi:10.1111/1755-
0998.12593

70. Andrews S. FastQC: a quality control tool for high throughput sequence data. Babraham
Bioinformatics, Babraham Institute, Cambridge, United Kingdom; 2010.

71. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

72. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.
doi:10.1093/bioinformatics/btp352

73. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27:
2987–2993. doi:10.1093/bioinformatics/btr509

74. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Res. 2010;20: 1297–1303. doi:10.1101/gr.107524.110

75. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
doi:10.1093/bioinformatics/btr330

76. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, et al.
Qualimap: evaluating next-generation sequencing alignment data. Bioinformatics. 2012;28:
2678–2679. doi:10.1093/bioinformatics/bts503

77. Okonechnikov K, Conesa A, García-Alcalde F. Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data. Bioinformatics. 2016;32: 292–294.
doi:10.1093/bioinformatics/btv566

78. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al.
glmmTMB balances speed and flexibility among packages for zero-inflated generalized
linear mixed modeling. The R journal. 2017;9: 378–400. doi:10.32614/RJ-2017-066

79.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: A
tool set for whole-genome association and population-based linkage analyses. The
American Journal of Human Genetics. 2007;81: 559–575. doi:10.1086/519795

Figures

823
Fig 1: Experimental design and variation in fitness from 2018-2023 across 12 central
Oregon Populations. **A.** Map of study area and locations of source populations (BEL and SHL)
and target populations for assisted gene flow. Raster depicts elevation with warmer value
indicative of higher elevations. Colors represent different experiment blocks and shapes
represent treatment (circle = local seeds, triangle = source seedlings, square = source seeds).
B. Experimental design for assisted gene flow manipulation into each of the four experimental
blocks. Each block consists of three populations, one that receives seed from a cross between
source populations (F_1 seed), one that receives seedlings grown from the same cross, and a
third that receives seed from the native population. Assisted gene flow occurred in 2021.

834
Fig. 2: Assessing changes in fitness following assisted gene flow. **A-B**. Boxplots depict changes in floral abundance and seed production following assisted gene flow across treatments: control (native seed introduced), source seeds introduced (CA F_1), or source seedling introduced (CA F_1). Each point is an individual. Pre-AGF (green boxes) includes 2018-2021 growing seasons for floral abundance and 2018-2020 growing seasons for seed production. **C-D.** Changes in floral abundance and seed production parsed into the four experimental blocks. Summary statistics and p-values correspond to the AGF:Treatment term within the GLMMs. Border colors on the subpanels correspond to the location of the experimental block on the map in Fig. 1. Edges of boxes in boxplots correspond to the interquartile range, the line inside boxes to medians and the whiskers extend to the most extreme points within 1.5 × IQR from box hinges.

847
Fig. 3 Patterns of introgression into target populations following assisted gene flow. **A**.
Number of loci with evidence of introgression of CA private alleles for three experimental blocks.
In each subpanel, left, center and right populations correspond to control, seedling and seed
population treatments, respectively. Total number of potential loci with California private alleles
was 15,691. **B**. Allele frequencies of California private alleles across a representative
chromosome (chromosome 2 in the RRM population – a population with source seeds
introduced). Allele frequencies may be higher in 2022 because there was greater sample size
(27 vs. 14) in 2021. **C.** Variation in genetic principal component 1 before and after assisted gene
flow within three experimental blocks. California populations are distinct from Oregon
populations with greater values of PC1 (Fig. S7). Edges of boxes in boxplots correspond to the
interquartile range, the line inside boxes to medians and the whiskers extend to the most
extreme points within 1.5 × IQR from box hinges.

Fig. 4 Shifts in flowering time and number of trichomes accompany assisted gene flow.
A-B. Violin plots depict variation in flowering time and number of trichomes before (green) and after (gold) assisted gene flow. Four populations are pooled within each treatment. **C-F.** Variation in flowering time and number of trichomes from the two experimental blocks with genomic evidence of introgression and fitness impacts of AGF. Fitness was elevated following AFG in block A, but lower in the block D (Fig. 2). Border color of panels corresponds to experiment block color in Fig.1. Points and whiskers inside violin plots correspond to mean and standard error, respectively. Summary statistics and p-values correspond to the AGF:Treatment term within the GLMMs.

Extended Data
Extended Methods

Fig. S1: Seedling transplant into a central Oregon Population (FIR) at different developmental stages.
Fig. S2: Changes in variation in fitness following assisted gene flow.
Fig. S3: Variation in floral abundance and seed production across six growing seasons and twelve populations
Fig. S4: Relative changes in floral abundance and seed production across years following assisted gene flow.
Fig. S5: Temporal variation in fitness during the assisted gene flow experiment.
Fig. S6: Limited introgression of California private alleles within target populations.
Fig. S7: Populations structure across source and target monkeyflower populations.
Fig. S8: Variation between pre- and post-assisted gene flow in flowering time within a common garden by experimental block
Fig. S9: Variation between pre- and post-assisted gene flow in number of trichomes within a common garden by experimental block.

Supplemental Tables
Table S1: Population geographic locations and treatments
Table S2: Model summaries of fitness effects of assisted gene flow within Oregon populations
Table S3: Estimated marginal means and raw means for of fitness effects of assisted gene flow across Oregon populations
Table S4: ANOVA summary of fitness effects of assisted gene flow within each block
Table S5: Estimated marginal means and raw means for of fitness effects of assisted gene flow within individual blocks
Table S6: ANOVA summary of fitness effects in individual years following assisted gene flow
Table S7: Sampling summary for genomic analysis
Table S8: Generalized linear model summaries for private allele analysis
Table S9: Counts of private alleles across target Oregon populations and years
Table S10: Sampling summary from the resurrection experiment
Table S11: Generalized linear modeling results from the resurrection experiment
Table S12: Source seedling performance in target Oregon populations
Table S13: Population summary of coverage and mean depth for genomic analysis
Table S14: Phenotypic correlations in the resurrection experiment