Distinct denitrification phenotypes in closely related bacteria: clues to understanding variations in nitrite accumulation among *Stutzerimonas* strains

Martin Menestreau[1], Daniel A. Milligan[1], Louise B. Sennett[1], Linda Bergaust[1], Lars R. Bakken[1], Gary Rowley[2], Morten Kjos[1], James P. Shapleigh[3] and Åsa Frostegård[1]

[1] Faculty of Chemistry, Biotechnology, and Food Sciences, Norwegian University of Life Sciences, Norwegian University of Life Sciences, Ås, Norway

[2] School of Biological Sciences, University of East Anglia, Norwich, United Kingdom

[3] Department of Microbiology, Cornell University, Ithaca, NY, USA

Correspondance: Åsa Frostegård (asa.frostegard@nmbu.no)

Funding: This work was supported by the Research Council of Norway, project no. 325770 (awarded to Å.F.).

Running title: Nitrite accumulation in *Stutzerimonas* strains

Keywords: Denitrification phenotypes, Nitrite accumulation, *Stutzerimonas* strains

Author details:

Martin Menestreau, e-mail: martin.menestreau@nmbu.no; ORCID id: 0000-0001-5060-5903

Daniel A. Milligan, e-mail: daniel.a.milligan@gmail.com; ORCID id: 0009-0002-3792-1308

Louise B. Sennett, e-mail : louise.sennett@nmbu.no; ORCID id: 0000-0001-5445-9712

Linda Bergaust, e-mail: linda.bergaust@nmbu.no; ORCID id: 0000-0001-5100-6611

Lars R. Bakken, e-mail: lars.bakken@nmbu.no; ORCID id: 0000-0002-7293-8108

Gary Rowley, e-mail: g.rowley@uea.ac.uk; ORCID id: 0000-0002-5421-4333

Morten Kjos, e-mail: morten.kjos@nmbu.no; ORCID id: 0000-0003-4448-9082

James P. Shapleigh, e-mail: jps2@cornell.edu; ORCID id: 0000-0002-8628-3966

Åsa Frostegård, e-mail: asa.frostegard@nmbu.no; ORCID id: 0000-0002-4754-5994

Abstract

Nitrite (NO_2^-) is a key denitrification intermediate, formed from nitrate (NO_3^-). Transient NO_2^- accumulation varies among denitrifiers, yet the underlying causes remain poorly understood, despite its potential toxicity and role in NO and N_2O emissions. We profiled eighteen related *Stutzerimonas* strains, including the model *S. perfectomarina* ZoBell, and identified three phenotypic clusters (full, partial and low nitrite accumulators; FNA, PNA and LNA) based on the fraction of NO_3^--N transiently accumulated as NO_2^-. LNA strains lack or express the membrane-bound nitrate reductase (NarG) late, relying on periplasmic NapA for NO_3^- reduction, possibly explaining their balanced NO_2^- production/reduction. FNA and PNA strains possess NapA and NarG but differ in their nitrite reductase (NirS) clades. Delayed *nirS* transcription as FNA strains transition to NO_3^- respiration likely accounts for some NO_2^- accumulation. However, addition of NO_3^- halted NO_2^- reduction in FNA strains, suggesting additional metabolic control. This may require the cytochromes NirTB, which are only found in FNA strains. The regulator DnrE was also unique to NO_2^--accumulators, likely having a role in finetuning NO_2^- regulation. Our findings reveal diverse NO_2^--handling phenotypes among denitrifiers and provide insights for optimizing wastewater nitrogen removal and soil bioaugmentation strategies to mitigate N_2O emissions.

Introduction

Denitrification is one of the main pathways driving the global nitrogen cycle. Complete denitrification is the four-step reduction of nitrate (NO_3^-) to dinitrogen (N_2) via the intermediate products nitrite (NO_2^-), nitric oxide (NO) and nitrous oxide (N_2O). In this process, the denitrification enzymes Nar, Nir, Nor and Nos are responsible for reduction of NO_3^-, NO_2^-, NO and N_2O, respectively (Berks et al. 1995; Zumft 1997). A wide variety of facultative anaerobic organisms, including bacteria, archaea and fungi, can perform all or some of the reduction steps associated with denitrification (Shapleigh 2013; Roothans et al. 2025). A shared trait amongst all denitrifiers is that O_2 is the preferred electron acceptor, but if the O_2 concentration decreases, denitrifiers use a variety of regulatory mechanisms to switch from O_2 to N-oxides as electron acceptors (Spiro 2007; Bakken et al. 2012), leading to large variations in their transient accumulation of denitrification

intermediates (Lycus et al. 2017). Current knowledge of how denitrifiers control the transition from aerobic

respiration to denitrification is largely based on studies of a few model organisms capable of complete

denitrification of NO_3^- to N_2. These studies have resulted in a relatively clear, though likely incomplete,

understanding of transcriptional and post-transcriptional mechanisms influencing denitrification

phenotypes (Spiro, 2012; Vaccaro et al. 2016; Gaimster et al. 2018).

One knowledge gap is the understanding of mechanisms controlling NO_2^- accumulation in

denitrifying bacteria (Liu et al. 2013). Nitrite is a central molecule in the nitrogen cycle that can be produced

and consumed in many redox processes, and its reduction is the defining reaction of denitrification (Kuypers

et al. 2018). In its protonated form, HNO_2, it can diffuse into cells and cause damage itself or through the

formation of toxic, reactive nitrogen species (Chislett et al. 2022). Moreover, accumulation of NO_2^- in the

environment can cause increased emissions of the greenhouse gas N_2O by impairing N_2O reduction

(Highton et al. 2022) or by producing N_2O through reactions with ferrous iron or organic matter compounds

(Van Cleemput and Samater 1995). Paradoxically, in some systems, NO_2^- accumulation is desirable. One

example is denitratation in wastewater treatment plants, where denitrification is used instead of ammonia

oxidation to provide nitrite for the anammox process (Zhuang et al. 2022).

The best-studied denitrifying bacteria, *Paracoccus denitrificans*, *Pseudomonas aeruginosa* and

Pseudomonas stutzeri ZoBell, all transiently accumulate some NO_2^- during anaerobic growth on NO_3^-

(Carlson and Ingraham 1983). However, studies extending beyond these model organisms have revealed

substantial variations in NO_2^- accumulation among diverse taxa (Lycus et al. 2017). A comparison of

Thauera strains (Liu et al. 2013) showed that some exhibited a rapid and complete onset (RPO) of all four

steps of the denitrification pathway, resulting in little to no NO_2^- accumulation. Others displayed a

progressive onset (PO) in which all available NO_3^- was reduced to NO_2^- prior to any reduction of NO_2^- took

place, leading to NO_2^- accumulation. A shared feature of the RPO strains was that they carried the

periplasmic NapA for dissimilatory NO_3^- reduction but lacked the membrane-bound NarG, while the PO

strains carried both NapA and NarG. All strains induced NO_3^- reduction as O_2 approached depletion, but

the regulatory dynamics varied. In the RPO strains, transcription of the *nirS* and *nosZ* genes - which encode

Nir and Nos, respectively – was initiated concurrently with Nar. In contrast, the NO_2^- accumulating PO

strains delayed transcription of these genes until all available NO_3^- had been reduced to NO_2^-. This led to

the tentative suggestion that a NO_3^- responsive mechanism might regulate *nir*-transcription in the NO_2^-

accumulating strains, although no direct evidence for this was found (Liu et al. 2013).

The present study was initiated to clarify how regulatory mechanisms drive variations in

denitrification phenotypes, and how these can differ even among closely related bacteria, with a particular

focus on NO_2^- accumulation. We compared eighteen strains within the former *Pseudomonas stutzeri* species

cluster, now reclassified as the genus *Stutzerimonas* (Lalucat et al. 2022). This complex contains

metabolically versatile organisms that are widespread in various environments (Lalucat et al. 2006). We

combined genome analyses with determinations of high-resolution denitrification kinetics and transcription

patterns of denitrification related genes during and after transition from aerobic to anaerobic respiration.

Using deletion mutations, we investigated whether the Crp/Fnr-like regulator DnrE, known to affect

transcriptional regulation in *S. stutzeri* (Härtig et al. 1999), influences NO_2^- accumulation in a closely related

strain. Moreover, we explored the potential role of sRNAs, which can act as both activators and repressors

of transcription and translation and have been shown to influence denitrification via control of NirR in *P.*

denitrificans (Gaimster et al. 2018; Moeller et al. 2021). We also considered whether electron competition

between electron pathways to the different denitrification reductases (Cole 2025; Richardson 2025) could

explain differences in NO_2^- accumulation among the *Stutzerimonas* strains. Electrons are delivered from

the TCA cycle via dehydrogenases, such as NADH or FDH dehydrogenase, to the quinone/quinol pool,

which serves as a key branching point in the electron transport chain. In complete denitrifiers, electrons are

then directed either to NO_3^- reductases (NapA, NarG) or to the bc_1 complex, a second branching point, from

which electrons are passed to Nir, Nor, and NosZ via cytochromes or azurins/pseudoazurins. Competition

occurs primarily at these branching points (Chen and Strous, 2013; Mania et al. 2020). In this study, we

specifically investigated the electron competition between NO_3^- and NO_2^- reduction pathways.

Our results categorize the eighteen strains into three distinct phenotypic groups showing full, partial or low NO_2^- accumulation (FNA, PNA and LNA). A major difference among these phenotypes was the presence and regulation of dissimilatory NO_3^- reductases (Nar/Nap), with Nar either absent or showing delayed transcription in LNA strains. Among the NO_2^--accumulating strains (FNA and PNA), differences were associated with *nir* gene cluster arrangement and NirS clade affiliation. Additional differences among phenotypic groups were observed in the transcriptional patterns of denitrification reductase genes and in the metabolic control of NO_3^- and NO_2^- reduction pathways. These findings indicate that denitrification phenotypes cannot be predicted solely from gene content but are instead governed by a multilayered regulatory network operating at both transcriptional and post-transcriptional levels. Since NO_2^- is a central molecule in the biochemical N-cycle and is utilized by several biological processes, this knowledge is important for developing biotechnological approaches, such as using denitrifiers in combination with anammox bacteria for improved wastewater treatment technologies, or engineering the soil microbiome in agriculture to reduce greenhouse gas emissions.

Materials and Methods

Strains and growth conditions

Eighteen bacterial strains, previously classified as belonging to the *P. stutzeri* species complex (Lalucat et al. 2006; Li et al. 2022), were selected for comparison of their denitrification phenotypes. The strains were obtained from various culture collections. A recent taxonomic revision of the genus *Pseudomonas* reclassified these strains into several species of the novel genus *Stutzerimonas* (Lalucat et al. 2022; Gomila et al. 2022). A list of the strains, including their origin of isolation and culture collection accession numbers, is found in Table S1.

All strains were incubated at 20°C in 120 mL serum vials containing 50 mL Sistrom's mineral medium (Sistrom 1960; Bergaust et al. 2010) supplemented with 2 mM KNO_3 or KNO_2, added as filter sterilized solutions. Vials prepared for cultivation under limited O_2 conditions were sealed with gas tight

butyl rubber septa and aluminum caps (Matriks AS, Norway), and air was evacuated from the 70 mL

headspace by six repetitions of applying vacuum for 360 s and He overpressure for 30 s. The overpressure

was released prior to the injection of the desired amount of O_2 and N_2O. Near-equilibrium between

headspace and the liquid phase was maintained by vigorous stirring at 600 rpm using a Teflon-coated

triangular magnetic bar. The stirring also prevented cell aggregation, ensuring that the cells were evenly

dispersed and experienced the intended gas concentrations. At the start of each experiment, the vials were

inoculated with 1 mL of aerobically growing bacterial cultures to an initial OD_{600} of 0.05-0.10. The cultures

had been raised from frozen stocks and pre-cultured with vigorous stirring under fully oxic conditions for

at least 4 generations to minimize the amounts of previously synthesized denitrification enzymes in the

cells (Mania et al. 2020; Gao et al. 2021).

Gas kinetics and chemical analyses

Headspace gases were monitored using a robotized incubation system as described in Molstad et al. (2007)

and Molstad et al. (2016). Briefly, gas samples were taken automatically by an autosampler (Agilent GC

Sampler 80) through the rubber septa of the vials using a peristaltic pump. The samples were then pumped

into a gas chromatograph (Agilent 7890A) for analysis of O_2, N_2O, N_2 and CO_2 and into a

chemifluorescence NO analyzer (Model 200A, Advanced Pollution Instrumentation). A thermal

conductivity detector was used to measure O_2 and N_2, while an electron capture detector was used to

measure N_2O. To maintain a slight overpressure in the vials, each 1 mL gas sample was replaced with a

slightly larger volume of He at near-atmospheric pressure. Three replicate vials were measured for each

treatment unless otherwise stated (n=3).

To determine NO_2^- concentrations, samples were collected from a parallel set of vials to avoid

sampling losses and changes in the headspace-to-liquid ratio. The samples were collected manually at

regular intervals through the rubber septum using a sterile syringe. Concentrations of NO_2^- were determined

immediately after each sampling, using the methodology described in Lim et al. (2018), which is based on

the method by Braman and Hendric (1989) and Cox (1980). Briefly, 10 µl liquid sample was injected into

a purging device containing a reducing agent (NaI) in 50% acetic acid at room temperature and the NO

produced was measured with a chemifluorescence NOx analyzer (Sievers NOA 280i). Samples from three

replicate vials were measured (n=3). Calculations based on cubic spline interpolation allowed estimation

of values between sampling times.

Statistical analyses

Statistical analyses were performed using the base statistics software in RStudio. In brief, one-way analysis

of variance (ANOVA) tests were performed to assess whether there were significant differences in the

levels of denitrification intermediates produced by the different NO_2^- accumulation phenotypes and by the

various overexpression mutants. When differences occurred ($p = 0.05$), mean pairwise comparisons were

performed using Tukey's test (HSD) for the post hoc analysis of the ANOVA models. Normality was tested

using homoscedasticity plots and the Shapiro-Wilks test of the residuals ($p > 0.05$).

Transcription analysis

Samples for RNA extraction (approximately 5×10^8 cells per sample) were taken at intervals from the liquid

phase. Cells were pelleted at 10,000 g for 7 minutes at 4°C and stabilized using RNAprotect (Bacteria

Reagent, Qiagen) according to the manufacturer's protocol. Total RNA was extracted using the RNeasy

Mini Kit (Quiagen). RNA concentrations were measured using the Qubit RNA BR Assay Kit (Life

Technologies). Genomic DNA was removed with Turbo DNase (Life Technologies) and genomic DNA

contamination was ruled out by SYBR green qPCR amplification using primers for *nosZ* (Table S2) prior

to reverse transcription. The assays were run on an Applied Biosystems StepOnePlus system, and samples

with values < 5 *nosZ* copies per reaction were deemed gDNA free (Lim et al. 2016). Reverse transcription

was then carried out with Maxima First Strand cDNA Synthesis Kit (Thermo Scientific), the product was

diluted, and gene transcripts were quantified by digital droplet PCR (ddPCR). A ddPCR reaction mix was

prepared (QX200 ddPCR EvaGreen supermix; Bio-Rad). Oil droplet suspensions were prepared using a

QX200 droplet generator (Bio-Rad), and PCR reactions were performed using the temperature program:

enzyme activation at 95 °C for 5 min; 40 cycles of denaturation at 95 °C for 30 s and combined

annealing/extension at 60 °C for 1 min; followed by signal stabilization at 4 °C for 5 min and a final step at 90 °C for 5 min. PCR products were quantified using a QX200 droplet reader (Bio-Rad) and the data was analyzed using QuantaSoft Analysis Pro 1.0.596 (Bio-Rad). All primers used are listed in Table S2.

Deletion of *narG* and *dnrE* in *Stutzerimonas decontaminans* strain 19SMN4

The pACRISPR/pCasPA system developed by Chen et al. (2018) for genome editing in various *Pseudomonas* species was used to construct the two deletion mutants, Δ*narG* and Δ*dnrE*, in *Stutzerimonas decontaminans* 19SMN4. pACRISPR plasmids specific to the targeted gene were constructed to contain a spacer targeting the specific gene designed using the tool CRISPOR (Concordet and Haeussler 2018) and a repair template consisting of 1000-bp flanking sequences, upstream and the downstream of the gene. The pCasPA was electroporated into *S. decontaminans* strain 19SMN4 following the protocol of Choi et al. (2006), but with electroporation parameters optimized for this strain (2500V, 200Ω, 25µF, 2 mm cuvette). Colonies containing pCasPA were selected after incubation on LB agar in the presence of 50 µg mL^{-1} tetracycline. Before the second electroporation for introduction of the gene specific plasmid pACRISPR, the strain containing the pCasPA plasmid was induced for 2 hours with 2 mg mL^{-1} L-arabinose, followed by electroporation under the same conditions as the first electroporation. Transformants were then selected on 50 µg mL^{-1} tetracycline and 150 µg mL^{-1} carbenicillin plates. The deletions were validated by PCR (for primers see Table S2) and Sanger sequencing. Colonies of successfully mutated cells were restreaked one to three times on LB agar supplemented with 10% w/v sucrose to cure the plasmids from the cells.

Stutzerimonas perfectomarina ZoBell sRNA overexpression mutants

Small RNAs (sRNAs) were overexpressed in *Stutzerimonas perfectomarina* ZoBell using the arabinose-inducible vector pL2020 (Sommer et al. 2017). The pL2020 plasmid was linearized by digestion with NdeI and HindIII. sRNAs were amplified from *S. perfectomarina* ZoBell genomic DNA using overlapping PCR. The resulting PCR fragments were assembled with the digested vector using the NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs) and transformed into *E. coli* DH5α competent cells. Transformants were selected on LB agar plates containing 30 µg mL^{-1} chloramphenicol. Positive clones

205   were screened by colony PCR and verified by Sanger sequencing. Constructed plasmids were subsequently

206   electroporated into the *S. perfectomarina* ZoBell host strain using an adapted protocol (Choi et al. 2006)

207   with optimized electroporation parameters as described above. The growth of the sRNA overexpression

208   mutants was assessed in LB medium supplemented with 30 µg mL^{-1} chloramphenicol to maintain plasmid

209   selection.

Genome sequencing and assembly

211   DNA was extracted with the QIAmp DNA Mini Kit (Qiagen, USA) and sent to the Norwegian Sequencing

212   Centre (University of Oslo) for TruSeq Nano preparation and Illumina MiSeq sequencing. Reads were

213   trimmed using Nesoni clip (https://github.com/Victorian-Bioinformatics-Consortium/nesoni) and contigs

214   were assembled using Velvet (Zerbino and Birney 2008). Sequence annotation was carried out on the RAST

215   server (Aziz et al. 2008). NCBI accession numbers are listed in Table S1.

Gene sequence analyses and phylogenetic determinations

Sequences coding for housekeeping genes assigned to strains in the *P. stutzeri* species complex, as well as

regulatory and structural genes encoding relevant nitrogen cycle processes, were retrieved from the NCBI

databases. For protein-coding genes, the match sequences that represented uninterrupted open reading

frames (ORFs) were identified, and gene synteny was compared for strains for which extensive operon data

was available. The phylogenetic relationship of the strains was determined based on a multilocus sequence

analysis (MLSA) of the genes encoding the small subunit RNA (16S rRNA), gyrase B (*gyrB*) and RNA

polymerase (*rpoD*). The sequences were complete, except in the case of 16S rRNA genes where the first

30 residues were omitted, as they were not available for strain DSM 50238. Genes were aligned using

MUSCLE v. 3.8.31 (Edgar 2004) and examined before Jukes Cantor distance estimation and maximum-

likelihood tree construction, using PhyMl 3.1 (Guindon et al. 2010). Amino acid-based enzyme phylogenies

were calculated using Clustal Omega (Sievers et al. 2011) and phylogenetic trees were constructed using

the Whelan and Goldman model in PhyMl.

Results

Nitrite accumulation phenotypes and phylogeny

Analyses of the denitrification phenotypes revealed significant differences in transient NO_2^- accumulation among the strains (Table 1). When incubated under denitrifying conditions with initially 100 µmol vial^{-1} of NO_3^- (2 mM), the full nitrite accumulator (FNA) strains reduced ≥89 % of the provided NO_3^- to NO_2^- before any NO_2^- reduction took place (Figure 1a and Figure S1). The partial nitrite accumulator (PNA) strains (Figure 1b and Figure S2) exhibited transient NO_2^- accumulation, amounting to about half of the provided NO_3^- (43.7- 62.2 µmol vial^{-1} of NO_2^-). The low nitrite accumulator (LNA) phenotype (Figure 1c and Figure S3) showed no or little NO_2^- accumulation (0.2-1.6 µmol vial^{-1}), except for strains 24a75 and 4C29, which transiently accumulated 11.3 and 17.1 µmol NO_2^- vial^{-1}, respectively.

A phylogenetic tree was constructed, incorporating the eighteen *Stutzerimonas* strains selected for analysis of denitrification phenotypes, along with additional related strains. Each strain´s classification is marked in the phylogenetic tree with red (FNA), yellow (PNA) or blue (LNA) (Figure 2). The tree was based on multilocus sequence analysis (MLSA) of two complete housekeeping genes (*gyrB* and *rpoD*) and nearly complete 16S rRNA gene sequences. Only limited congruence was observed between phylogeny and the NO_2^- accumulation phenotype: FNA and PNA strains clustered together and were often intertwined, whereas LNA strains were distributed across multiple, widely separated branches. The neighbor joining tree based on kmer clustering of a more extensive collection of genomes from *Stutzerimonas* strains (Figure S4) revealed a similar pattern, also consistent with other recently published phylogenetic studies (Li et al. 2022; Gomila et al. 2022).

Selected genetic features related to denitrification

An analysis of denitrification related gene clusters was performed to identify possible relationships with the NO_2^- accumulation phenotypes (Table 1). All eighteen strains had genetic potential to perform complete denitrification. They all carry the *napEDABC* gene cluster encoding periplasmic nitrate reductase (Nap). Nap also was the dominant NO_3^- reductase in the clustered set of genomes (Figure S4). Two strains in the LNA group (DNSP21 and 28a3) rely only on Nap for dissimilatory NO_3^- reduction, since they lack the

cytoplasmic nitrate reductase Nar, encoded by the *nar* genes. The *narGHJIKMXL* cluster was found in all

other strains. A notable difference in *nar*-related genes among the phenotype groups was the presence of

the *dnrE* gene, which was found in all FNA and PNA strains but was absent in all LNA strains. DnrE is

located within the *nar* gene cluster. It belongs to the group dissimilatory nitrate respiration regulators

(DNR), a subgroup of transcriptional regulators related to the Fnr/Crp family (Trunk et al. 2010; Ebert et

al. 2017), and has a potential role in regulating the *narG* gene expression (Härtig et al. 1999).

All analyzed strains carried the *nirS* gene encoding the cytochrome cd_1 nitrite reductase (Zumft

1997). The *nirS* gene is part of a large gene cluster encoding various enzymes involved in the synthesis and

function of NirS (Jüngst et al. 1991; Philippot 2002). The FNA and PNA strains have identical *nirS* gene

cluster configurations, apart from the presence of *nirTB* in the FNA strains, while the *nirS* gene cluster in

the LNA strains is more diverse (Figure S5). Based on their amino acid sequences, the NirS reductases

have been divided into several clades. (Jones et al. 2008; Pold et al. 2024). Following this classification,

the NirS type of the FNA organisms in the present study belongs to clade 1b, while that of the PNA and

LNA strains belong to clade 1a (Table 1), except the LNA strain JM300 which carries the NirS clade 1b

(Figure S5). A common trait for all analyzed (Table 1) or clustered (Figure S4) *Stutzerimonas* strains with

clade 1b NirS is the consistent presence of the genes *nirT* and *nirB*. NirT and NirB are tetraheme and diheme

cytochromes, respectively, with a putative role in electron transfer and possibly also in the catalytic

activation of the cd_1 cytochrome (Härtig and Zumft 1999; Zajicek et al. 2004). The *nirTB* genes are absent

in all clade 1a strains.

Of the 314 genomes in the clustered tree (Figure S4), 111 carried the *nirK* gene, which encodes the

copper-containing nitrite reductase NirK. Among the analyzed strains, seven possessed both *nirK* and *nirS*

(Table 1). Notably, *nirK* was present in strains across all three NO_2^- accumulator groups. Previously, *nirK*

was divided into two main clades based on copper-binding histidine motifs: TRPHL for clade I; and

SSFH(V/I/P) for clade II (Jones et al. 2008; Declayre et al. 2016). A recent study (Pold et al. 2024) proposed

a more detailed classification, reassigning *Stutzerimonas* strains with clade I NirK to clade 1a, and those

with clade II to clade 1e. In the analyzed *Stutzerimonas* strains carrying the *nirK* clade I gene (1a), the gene

is consistently followed by *nirV*. This was observed in two of the strains in the present study, both belonging

to the LNA phenotype group (Table 1). Half of the FNA strains analyzed in this study, along with one PNA

strain, possess *nirK* clade II (1e), consistently followed by the *norCBQD* gene cluster.

The nitric oxide reductase cNor, which was present in all eighteen strains, is encoded by the

norCBD operon located in the denitrification island together with the *nir* and *nos* clusters (Zumft 1997;

Vollack et al. 1998). The *norCBQD* cluster present downstream of the *nirK* clade II gene in some of the

FNA and PNA strains may be used under conditions when *nirK* is expressed, given their proximity. The

gene cluster *nosRZDFYL* encoding the nitrous oxide reductase (NosZ) clade I was also found in the

denitrification island in all the studied *Stutzerimonas* strains. The *nor* and *nos* gene cluster arrangements

were identical in all strains (Table 1).

Denitrification kinetics and transcription analysis

Electron flow rates to the different reductases were calculated from denitrification kinetics measured for all

eighteen strains during and after the switch from aerobic respiration to denitrification when incubated with

1% initial O_2 and 2 mM NO_3^- (100 μmol vial^{-1}). All strains could perform the complete denitrification

process, reducing NO_3^- to N_2. Examples for some representative strains are shown in Figures 1 and 3, while

Figures S1, S2 and S3 show the denitrification kinetics for the remainder. Common to all strains,

irrespective of which NO_2^- accumulation group they belonged to, was that NO_3^- reduction started when the

O_2 had decreased to 10-15 μmol vial^{-1} (corresponding to 0.6-0.9 μM in the liquid). The FNA strains reduced

all the provided NO_3^- to NO_2^- before any NO_2^- reduction took place, as shown by the lack of intermediate

gas products (Figure 1a and Figure S1). The start of NO_2^- reduction coincided with the peak of NO_2^-

accumulation and was accompanied by nearly simultaneous production of NO, N_2O and N_2. The sequential

reduction of NO_3^- and NO_2^- in the FNA strains is also reflected in the electron flow kinetics (Figure 1d and

Figure S1), with nearly no overlap between the curve showing the electron flow rate to NO_3^- ($V_{eNar/Nap}$, red

shading) and that to NO_2^- (V_{eNir}; purple shading) and a clear separation between O_2 reduction (V_{eO2}; blue

shading) and NO_2^- reduction (V_{eNir}; purple shading). The electron flow for the FNA strains typically showed

a prominent dip when the cultures shifted from NO_3^- reduction to NO_2^- reduction (Figure 1d, Figure 3e and

Figure S2), suggesting that only a fraction of the cells respired NO_2^- as found for *Paracoccus denitrificans*

(Lycus et al. 2018).

The PNA and LNA strains reduced NO_3^- and NO_2^- simultaneously, as indicated by the production

of NO and N_2O, followed by the appearance of N_2 (Figure 1b-c, Figure S2 and Figure S3). The strains

CCUG 11256 (PNA) and 28a3 (LNA) were similar in their denitrification phenotypes (Figure 1b-c), except

for the transient accumulation of NO_2^- by the PNA strain. The simultaneous reduction of NO_3^- and NO_2^-

was reflected in the overlap between the areas under the curves for $V_{eNar/Nap}$ (red shading) and V_{eNir} (purple

shading) (Figure 2e-f, Figure S2 and Figure S3). Production of N_2 was detected somewhat later, when O_2

was near depletion. For both the PNA and LNA groups, the electron flow indicated smooth transitions

between the different electron acceptors with no distinct dips and thus an overlap between the curves for

$V_{eNar/Nap}$ and Ve_{Nir}.

All three NO_2^- accumulation phenotypes showed little transient accumulation of NO and N_2O

(Table 1), suggesting a strong regulatory control of the production and consumption of these gaseous

denitrification intermediates. When incubated with 1% initial O_2 and 2 mM initial concentration of NO_3^-,

the maximum net NO production was 9.1–70.5 nmol vial^{-1} with no significant ($p = 0.598$) difference

between the phenotype groups in terms of maximum NO accumulation. The N_2O accumulation remained

low for all strains, ranging between 0.3 and 89 nmol vial^{-1}, and was significantly lower for FNA compared

to LNA ($p = 0.030$), while PNA exhibited intermediate levels, showing no significant difference from either

FNA or LNA ($p = 0.840$ and 0.217).

Transcription of the denitrification reductase genes *narG, napA*, *nirS*, *norB* and *nosZ* was monitored

in the strains ZoBell (FNA) and JM300 (LNA) during and after the transition from aerobic respiration to

denitrification, along with the denitrification kinetics (Figure 3). In addition, transcription of the *nirK* gene,

which is lacking in strain ZoBell (Table 1), was monitored in strain JM300. Strain ZoBell reduced all NO_3^-

(100 µmol vial^{-1}) to NO_2^- before any significant NO_2^- reduction took place (Figure 3a), confirming its

classification into the FNA group. This was also reflected in the electron flow kinetics which showed that

electrons were channeled to Nir only once the electron flow to Nar stalled due to NO_3^- depletion (Figure

3e). Transcription of *napA* and *nosZ* in strain ZoBell (Figure 3c) was detected already at 0 hpi (hours post

inoculation). Increased transcription was observed for all the monitored genes at 10-13 hpi when less than

10 µmol of O_2 remained in the vial, with an initial transcription peak for most genes at 18 hpi. Although

transcripts for genes involved in all four denitrification steps were detected during this initial peak, it

resulted in only minimal reduction activity, except for NO_3^- reduction. A second pulse of transcription

occurred once all NO_3^- had been reduced and the NO_2^- concentration reached its maximum. During this

second event, the maximal levels of *norB* and *nirS* (inset) transcription were observed at 30-35 hpi. There

was also a notable increase in *nosZ* transcription, although not surpassing what was measured in the initial

phase. These peaks corresponded to the nearly simultaneous reduction of NO_2^-, NO and N_2O (Figure 3a).

Transcription was sustained until all intermediates were fully reduced to N_2.

The LNA strain JM300 (Figure 3b, d, f) reduced NO_3^- to gases, via NO_2^-, without any detectable

net accumulation of NO_2^-. NO and N_2O became detectable when the O_2 concentration was approximately

20 µmol vial^{-1}, corresponding to1.2 µM in the liquid (at 7.5 hpi), consistent with previous findings for this

strain (Wittorf et al. 2018). Concentrations of these gases peaked around 11 hpi, immediately followed by

N_2 production. The simultaneous reduction of the different electron acceptors is seen in the gradual increase

in total electron flow rates in JM300 as well as in the other LNA strains (Figure 3f and Figure S3). Also

notable is the absence of the significant dip in the electron flow rates that was observed in most of the FNA

strains during the shift from NO_3^- to NO_2^- reduction (Figure 3e and Figure S1). The overlapping $V_{eNar/Nap}$

and V_{eNir} curves (Figure 3f) reflect the simultaneous reduction of NO_3^- and NO_2^-, also observed for the other

strains characterized as LNA (Figure 2c, f and Figure S3). An important finding was that *narG* transcription

was delayed by five hours compared to the other genes, indicating that NO_3^- reduction was carried out by

the periplasmic nitrate reductase Nap during the early phase of denitrification. A slight delay in the

transcription of *nirS*, *nirK*, and *norB* was also observed compared to *napA* and *nosZ*. Like this LNA strain

(JM300), delayed *narG* transcription was seen in the LNA strain DSM 50238 (Figure S6), which exhibited

a transcriptional profile similar to JM300, although it lacks *nirK*.

Effect of *narG* and *dnrE* deletions on nitrite accumulation

The wild-type PNA strain 19SMN4 carries *narG*, which is either absent (Table 1) or transcribed with a

delay (Figure 3 and Figure S6) in the LNA strains, as well as *dnrE*, which is absent in all the LNA strains

(Table 1). We hypothesized that deletion of either gene would shift strain 19SMN4 from the PNA to the

LNA phenotype. Specifically, we predicted that NO_3^- reduction rate would slow down in the *narG* mutant,

as it would rely solely on the periplasmic NO_3^- reductase Nap for dissimilatory NO_3^- reduction, resulting in

decreased NO_2^- accumulation. Furthermore, given that all LNA strains lack the *dnrE* gene (Table 1), we

hypothesized that the absence of this transcriptional regulator would reduce or delay the transcription of

narG in the presence of NO_3^-, compared to the wild-type. This would similarly result in reduced NO_2^-

accumulation producing a phenotype resembling that of the LNA strains. To test these hypotheses, we

generated deletion *narG* and *dnrE* mutants in the PNA strain 19SMN4.

A comparison of the denitrification phenotypes of the wild-type and the two mutants showed that

the mutants remained complete denitrifiers, as all provided NO_3^- was fully reduced to N_2 (Figure 4). Both

the wild-type and the mutants transitioned to denitrification, indicated by the accumulation of low amounts

of NO (16.5-43.2 nmol vial^{-1}) and N_2O (11.0-34.9 nmol vial^{-1}), when O_2 fell below 7.5 µmol vial^{-1}(0.45 µM

in the liquid). The two mutants and the wild-type strain showed, however, different denitrification

phenotypes regarding their NO_2^- accumulation and the time it took to complete denitrification (seen as an

N_2 plateau). The wild-type strain accumulated approximately 50% of the provided 100 µmol NO_3^- vial^{-1} as

NO_2^- (62.8 µmol vial^{-1}), supporting its classification as PNA, determined in a separate experiment (Figure

S2) where this strain accumulated 44.0 µmol vial^{-1} NO_2^- . In contrast to the wild-type, the $\Delta narG$ mutant

accumulated <1.3 µmol vial^{-1} NO_2^- (Figure 4). It had thus obtained an LNA phenotype similar to the strains

Stutzerimonas frequens DNSP21 and *S. stutzeri* 28a3 (Figure 1c and Figure S3a), both of which lack *narG*

and thus use only Nap for dissimilatory NO_3^- reduction (Table 1). As expected, deleting *narG* slowed

denitrification so that the reduction of the provided NO_3^- to N_2 took almost three times longer than for the

wild-type (78 h *vs* 28 h). Knocking out the *dnrE* gene had a less severe effect on the time it took to complete

denitrification (38 h). However, and contrary to our expectations, the $\Delta dnrE$ mutant showed increased

accumulation of NO_2^- compared to the wild-type, reaching 80.1 ± 4.99 µmol vial^{-1}. The NO_2^- accumulation

rates were however almost equal with 5.5 ± 0.7 µmol vial^{-1} h^{-1} for the wild-type and 5.5 ± 0.4 µmol vial^{-1} h$^-$

^{1} for the $\Delta dnrE$ mutant.

Competition for electrons between the NO_3^- and NO_2^- reduction pathways

To investigate whether competition for electrons between the NO_3^- and NO_2^- reduction pathways influences

NO_2^- accumulation, we added NO_3^- to actively NO_2^- reducing cultures. Three strains were studied (ZoBell,

FNA; 19SMNA, PNA and JM300, LNA), each representing one of the phenotype groups (Figure 5a-c).

The cultures were initially incubated under the denitrifying conditions described above to induce the

development of the denitrification proteome, after which the cells were transferred to new anoxic vials

(initial OD_{600} 0.3) with medium containing 100 µmol NO_2^- vial^{-1} (corresponding to 2 mM NO_2^- in the

medium) (t=0). When approximately one third of the initial NO_2^- had been reduced, 100 µmol vial^{-1} of NO_3^-

(corresponding to 2 mM NO_3^- in the medium) was added to the vial. For strain ZoBell (Figure 5a), the

addition of NO_3^- to the actively NO_2^- respiring culture led to an immediate shift to NO_3^- respiration and a

complete halt of the NO_2^- reduction until all the added NO_3^- had been reduced. This shift is also seen from

the plummeting of the NO concentration and the plateauing of N_2. Once all the NO_3^- had been reduced,

NO_2^- respiration resumed along with the downstream denitrification steps as seen from the decrease of NO_2^-

and the concomitant accumulation of NO (<50 nmol vial^{-1}). N_2 increased until it reached a plateau at

approximately 200 µmol N_2-N vial^{-1}, reflecting the complete reduction of the added NO_3^- and NO_2^-.

In a similar experiment, strain 19SMN4 (PNA) reduced NO_3^- and NO_2^- simultaneously following

supplementation with NO_3^-, and continuously reduced NO_2^- further to N_2, as seen by the constant increase

of N_2 and lack of an intermittent N_2 plateau (Figure 5b). NO_3^- was apparently reduced faster than NO_2^-,

resulting in a transient accumulation of NO_2^- from 75 to 123 µmol vial^{-1}(thus about half of the added NO_3^-

) until all the available NO_3^- was depleted. The LNA strain JM300 (Figure 5c) continued to reduce NO_2^-

despite the addition of NO_3^-, with no accumulation of NO_2^-. The NO_2^- reduction proceeded without any

apparent reduction in rate until all NO_3^- and NO_2^- had been consumed, seen as N_2-N reaching a plateau. As

observed with strain ZoBell, the accumulation of NO peaked at $\leq$ 50 nmol vial^{-1} in experiments with strains

19SMN4 or JM300 (Figure 5b-c).

Analysis of a possible role of sRNAs in regulating nitrite accumulation

We investigated potential regulatory mechanisms underlying the nitrite accumulation phenotype by

examining the role of small RNAs (sRNAs). Inclusion of selected sRNAs was based on their previous

association to either regulation of the broader nitrogen cycle, to denitrification specifically, to anaerobic

regulators, and to regulators located within denitrification gene clusters in *Stutzerimonas* sp. (Lin et al.

2015; Zhan et al. 2016; Wang et al. 2022). With these criteria, eight putative sRNAs (crcZ, nfiS, P11, P15,

pda200, phrS, prrF1, rsmY) were selected for investigation and overexpression strains constructed.

Overexpression has been used to assign function to sRNAs from a wide range of bacterial species and

associated processes (Massé and Gottesman 2002; Gaimster et al. 2019). We assessed the NO_2^-

accumulation under anaerobic conditions for the sRNA-overexpressing strains (Figure S8). During

anaerobic growth, NO_2^- accumulation peaked between 83.2 $\pm$ 7.7 and 104.2 $\pm$ 0.4 μmol vial^{-1}. However,

pairwise Tukey HSD tests revealed no statistically significant differences in NO_2^- accumulation between

any of the overexpression strains and the control ($p > 0.05$).

Discussion

Nitrite accumulation can have significant environmental consequences, as it plays a central role in the

emissions of N_2O, NO and HONO from various environments (Bhattarai et al. 2018; Frostegård et al. 2022;

Song et al. 2023), and can also hamper N removal efficiency in wastewater plants (Zhou et al. 2011). Despite

its importance, variation in patterns of transient NO_2^- accumulation among denitrifying organisms has

received relatively little attention, and mechanisms underlying this phenotype remain poorly understood.

Our study of eighteen closely related strains revealed a greater diversity in denitrification phenotypes within

the genus *Stutzerimonas* than previously recognized, with strains exhibiting full, partial, or low NO_2^--

accumulating phenotypes. We identified multiple factors that either individually or in concert contribute to

varying levels of transient NO_2^- accumulation in *Stutzerimonas*, including genetic configurations as well as

regulatory mechanisms operating at both the transcriptional and post-transcriptional/metabolic levels. The

phenotype grouping showed partial congruence with phylogeny (Figure 2): FNA and PNA were

phylogenetically intertwined, and in some cases, even strains of the same species belonged to different

phenotype groups. The LNA strains were mostly phylogenetically separated from the two other phenotype

groups, with one exception (DNSP21). Genome clustering supported these observations, with the exception

that strains DNSP21 and 24a75 showed proximity (Figure S4). The limited congruence between phylogeny

or genome clustering and NO_2^- accumulation phenotype is consistent with findings from the genus *Thauera*

(Liu et al. 2013) and underscores the need for caution against inferring regulatory phenotypes based solely

on taxonomic relatedness.

The genome survey of denitrification-related genes (Table 1) did not conclusively explain the three

phenotype groups, as variations in the gene clusters encoding both dissimilatory NO_3^- and NO_2^- reduction

were not consistently associated with specific phenotypes. In contrast, the clusters for NO and N_2O

reduction were arranged consistently across all three groups. A previous investigation of bacteria in the

genus *Thauera* (Liu et al. 2013) suggested that *nar* plays a key role in NO_2^- accumulation, since strains

carrying both the *nar* and *nap* clusters accumulated NO_2^-, while those with only the *nap* cluster did not. In

line with this, all *Stutzerimonas* strains classified as FNA and PNA in the present study carried the *nar*

genes. However, unlike the *Thauera* strains, the *nar* cluster was also found in all but two of the LNA strains.

This suggests that additional mechanisms controlling the expression of denitrification related genes are

involved in determining the NO_2^- accumulation levels.

To investigate whether transcriptional control mechanisms could explain why the LNA strains

possessing both *nar* and *nap* accumulated little or no NO_2^-, we measured transcriptional activity in two such

strains. Strain JM300 showed significantly delayed *narG* transcription by approximately 4-6 h compared

to other denitrification related genes (Figure 3d), implying that it relied only on Nap for NO_3^- reduction

during this period. This slower NO_3^- reduction allowed Nir to maintain pace, preventing NO_2^- accumulation.

Interestingly, strain JM300 carries the NO_2^- reductase gene *nirK* in addition to *nirS*, and it cannot be

excluded that this contributed to the lack of NO_2^- accumulation. To explore this further, we performed a

transcription analysis of the LNA strain DSM 50238, which carries both *napA* and *narG* but only *nirS* for

dissimilary NO_2^- reduction (thus lacking *nirK*). These strains showed the same temporal pattern of

transcription as JM300, with a significant delay in *narG* transcription (Figure S5), although copy numbers

were lower for DSM 50238, for which real-time PCR was used, compared to the ddPCR analysis used for

JM300. Collectively, these results reinforce the role of NarG as a primary factor in determining whether a

denitrifier accumulates NO_2^-. This is further supported by the results from the *narG* knock-out mutant,

which shifts PNA strain 19SMN4 to a LNA phenotype (Figure 4).

While the presence and activity of Nar appear to be significant factors in determining whether a

strain accumulates NO_2^-, they do not conclusively explain the difference between the phenotype groups. A

primary distinction between the FNA group and the PNA/LNA groups appears to lie in their transcriptional

control of *nirS* (Figure 3). The FNA phenotype was characterized by a progressive onset of denitrification,

whereas the LNA phenotype showed a rapid and complete onset, similar to reports on *Thauera* strains (Liu

et al. 2013). In that study, the FNA strain *Thauera terpenica* showed a low transcription peak of *nirS* shortly

before O_2 depletion, but the main transcription was delayed and coincided with the peak of NO_2^-

accumulation and depletion of NO_3^-. This precise timing of *nirS* transcription led to the tentative suggestion

that NO_3^- interacted with one of the regulatory proteins involved in *nirS* transcription, possibly a repressor

protein, though no evidence was provided to support such a mechanism. The present, more detailed analysis

revealed a similar transcription pattern in *S. perfectomarina* strain ZoBell. Taken together, the findings

suggest that temporal control of transcription plays a major role in the near complete NO_2^- accumulation in

FNA strains, although other control mechanisms are likely at play as well.

The FNA strains showed an almost complete inhibition of NO_2^- reduction while NO_3^- was present,

whereas the PNA strains reduced NO_2^- in the presence of NO_3^-, albeit not at a rate sufficient to immediately

reduce the produced NO_2^- as effectively as the LNA strains (Figure 5a-c). This suggests that, in both PNA

and LNA organisms, electron flow can occur simultaneously to Nar/Nap and Nir, enabling some Nir activity

even when NO_3^- is available. The observed difference in NO_2^- accumulation between FNA and PNA strains

is unlikely to be explained by the electron pathway to NarG, via NarHI, being a stronger competitor for

electrons than the bc_1 complex in FNA organisms relative to PNA organisms. One notable difference

between these two groups is the clade of NirS, with FNA strains possessing clade 1b while PNA strains

have clade a. This distinction is accompanied by the presence of the genes *nirT* and *nirB* in the clade 1b

cluster, but not in the clade 1a cluster (Table 1). These genes encode tetraheme and diheme *c*-type

cytochromes, respectively. NirT is a member of the NapC/NirT family of cytochromes, which are

characterized by their electron transfer capacity. In addition, NirT may catalytically activate cytochrome

cd₁, potentially serving as an essential activating partner for NirS (Zajicek et al. 2004). Notably, NirT is

needed for functional NO_2^- reduction in *S. perfectomarina* strain ZoBell (Jüngst et al. 1991; Härtig and

Zumft 1999). The NirT and NirB proteins can apparently form a functional heterodimer, as fused NirTB

has been found in other NO_2^- accumulating bacteria, such as *Halomonas* sp. 2A, which was detected in a

denitritation bioreactor where denitrifying bacteria supplied NO_2^- for anaerobic ammonium oxidation (Li

et al. 2018). The involvement of NirTB in NO_2^- reduction is supported by the location of their genes

immediately downstream of *nirS*. However, despite these indications of function, the extent to which NirT

and NirB contribute to the near-complete inhibition of NirS activity by NO_3^- in FNA strains remains

unresolved.

Another difference between the phenotype groups was that the *dnrE* gene was present in all FNA

and PNA strains but absent in all LNA strains (Table 1), making it a candidate gene for involvement in NO_2^-

accumulation. DnrE belongs to the FNR/CRP family of transcriptional regulators, which sense

environmental signals such as O_2 and NO. We hypothesized that knocking out *dnrE* in an FNA or PNA

strain would decrease or delay the transcription of *narG*, thereby slowing NO_3^- reduction and reducing NO_2^-

accumulation. Contrary to our expectations, the NO_2^- accumulation rate of the PNA strain 19SMN4 deletion

mutant remained equal to the wild-type (Figure 4), excluding DnrE from playing a direct regulatory role in

NO_2^- accumulation. The *dnrE* gene is located within the *nar* gene cluster, alongside *narXL*. NarXL is a two-

component sensor–regulator system that senses NO_3^- and NO_2^- with a higher affinity for NO_3^-, which

directly controls the expression of *narG* and *dnrE* (Härtig et al. 1999). Consequently, in the *dnrE* knockout

mutant, *narG* expression was maintained, possibly through NarXL, despite the absence of *dnrE*, which

makes it challenging to elucidate the specific role of *dnrE* in regulating NO_2^- accumulation. Despite high

homology among the Dnr regulatory proteins DnrE, DnrS and DnrD (>40% identity and >90% coverage),

their functions are distinct: DnrS senses O_2, DnrD senses NO, while the role of DnrE remains unknown.

The regulation of *dnrE* appears to be conserved in all FNA and PNA strains, as indicated by a highly

conserved upstream intergenic region containing two putative NarL binding sites, a ribosome binding site

(RBS), and two promoters (Figure S9). This conservation suggests a similar regulatory mechanism

controlling *dnrE* expression. Consistent with Vollack et al. (1999), no FNR-box was detected upstream of

dnrE, suggesting that the main anaerobic regulator, Anr, which senses O_2, may not directly regulate *dnrE*.

Nevertheless, *dnrE* transcription increases in the presence of both NO_3^- and O_2, indicating that another

oxygen-sensitive factor may trigger its expression. This suggests that DnrE is involved in a complex,

multilayered regulatory network (Spiro 2012).

In our search for additional regulatory mechanisms influencing denitrification, we examined the

potential role of small RNAs (sRNAs) in modulating NO_2^- accumulation in the FNA strain *S.*

perfectomarina ZoBell. Candidate sRNAs were selected based on their conservation across *Pseudomonas*

species, their genomic proximity to denitrification gene clusters, and previous associations with nitrogen

metabolism. Although several candidates were successfully overexpressed under anaerobic conditions, no

significant changes in NO_2^- accumulation were observed, suggesting that none of the selected sRNAs

directly regulates denitrification under the tested conditions although mutational analyses would be needed

to fully rule them out. These findings highlight the limitations of a candidate-based approach and emphasize

the need for broader transcriptomic analyses. In particular, *de novo* RNA-seq analyses, comparing closely

related *Stutzerimona*s strains exhibiting different NO_2^- accumulation phenotypes, could provide valuable

insight into the denitrification specific transcriptional landscape.

Taken together, the results enhance the understanding of how variations in regulatory phenotypes

may influence NO_2^- accumulation across various environments and highlight how organisms optimize

energy savings by temporarily blocking steps in the denitrification pathway, albeit at the cost of providing

substrates to competing populations. Importantly, these findings have significant implications for various

biotechnological approaches. FNA strains can supply NO_2^- for denitratation in anammox wastewater

treatment (Zhuang et al. 2022). NNA strains, on the other hand, may be candidates for engineering soil

microbiomes for N_2O mitigation (Hiis et al. 2024), since they maintain low levels of all denitrification

intermediates.

Conclusions

Our study demonstrates substantial variation in denitrification phenotypes among closely

related *Stutzerimonas* strains, driven by a complex interplay of genetic, transcriptional, and post-

transcriptional mechanisms that influence NO_2^- accumulation. We found that even organisms with a

complete set of denitrification genes can act as partial denitrifiers by temporarily constraining specific steps,

as seen in the FNA strains. This likely conserves energy, providing an advantage in environments with

fluctuating O_2 levels where full denitrification is only occasionally required. Similar to classical "partial

denitrifiers", which lack certain genes and therefore share metabolic tasks (Roothans et al. 2025), FNA-like

organisms rely on community members capable of consuming NO_2^-, including other denitrifiers, DNRA

organisms (performing dissimilatory NO_3^-/NO_2^- reduction to ammonia), or NO_2^- oxidizers if O_2 is available,

to keep NO_2^- concentrations non-toxic. This collaboration helps prevent the buildup of NO_2^- and the

formation of environmentally harmful byproducts such as NO and N_2O, including those from abiotic NO_2^-

reactions (Lim et al. 2018). While the presence and timing of *narG* transcription were identified as key

determinants of transient NO_2^- accumulation, other factors—including the type of *nirS* and associated *nirT/nirB* genes, as well as competition for electrons between Nar and Nir pathways—also play important roles. Our results highlight the limitations of predicting NO_2^- accumulation potential based solely on taxonomy or gene presence. Overall, these findings underscore the complexity of denitrification regulation and emphasize the need for integrative approaches combining microbial physiology, biochemistry, and transcriptomics to link genotype to denitrification phenotypes. A deeper understanding of these mechanisms is essential for managing NO_2^- and related intermediates in both environmental and engineered systems, such as wastewater treatment, and for supporting efforts to mitigate greenhouse gas emissions.

Author contributions

Martin Menestreau: investigations, formal analysis, visualization, data interpretation, writing – original draft, writing – review and editing. **Daniel A. Milligan:** preliminary investigations (gas kinetics, transcription, genome analyses), writing – early draft. **Louise B. Sennett:** data analysis (statistics), writing- - editing. **Linda Bergaust:** data interpretation, writing – review and editing. **Lars R. Bakken** data interpretation, writing – review and editing. **Gary Rowley:** supervision of sRNA work, writing – review and editing. **Morten Kjos:** supervision of CRISPR analyses, writing – review and editing. **James P. Shapleigh:** data analysis (phylogeny), data interpretation, validation, writing – review and editing. **Åsa Frostegård:** conceptualization, funding acquisition, supervision, project administration, data interpretation, validation, writing – original draft, writing – review and editing. All authors contributed to the article and approved the submitted version.

Acknowledgements

This work was supported by the Research Council of Norway, project no. 325770 (awarded to Å.F.).

Conflicts of interest

The authors declare no conflicts of interest

References

Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., … Zagnitko, O. (2008). The
RAST server: Rapid annotations using subsystems technology. *BMC Genomics, 9*(1), 75.
https://doi.org/10.1186/1471-2164-9-75

Bakken, L. R., Bergaust, L., Liu, B., & Frostegård, Å. (2012). Regulation of denitrification at the cellular
level: A clue to the understanding of N_2O emissions from soils. *Philosophical Transactions of the Royal*
Society B: Biological Sciences, 367(1593), 1226–1234. https://doi.org/10.1098/rstb.2011.0321

Bergaust, L., Mao, Y., Bakken, L. R., & Frostegård, Å. (2010). Denitrification response patterns during
the transition to anoxic respiration and posttranscriptional effects of suboptimal pH on nitrogen oxide
reductase in *Paracoccus denitrificans*. *Applied and Environmental Microbiology, 76*(19), 6387–6396.
https://doi.org/10.1128/AEM.00608-10

Berks, B. C., Ferguson, S. J., Moir, J. W., & Richardson, D. J. (1995). Enzymes and associated electron
transport systems that catalyze the respiratory reduction of nitrogen oxides and oxyanions. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics, 1232, 97–173. https://doi.org/10.1016/0005-2728(95)00092-5

Bhattarai, H. R., Virkajärvi, P., Yli-Pirilä, P., & Maljanen, M. (2018). Emissions of atmospherically
important nitrous acid (HONO) gas from northern grassland soil increases in the presence of nitrite (NO_2^-
). *Agriculture, Ecosystems & Environment, 256*, 194–199. https://doi.org/10.1016/j.agee.2018.01.017

Braman, R. S., & Hendrix, S. A. (1989). Nanogram nitrite and nitrate determination in environmental and
biological materials by vanadium(III) reduction with chemiluminescence detection. *Analytical Chemistry,*
61(24), 2715–2718. https://doi.org/10.1021/ac00199a007

Carlson, C. A., & Ingraham, J. L. (1983). Comparison of denitrification by *Pseudomonas stutzeri,*
Pseudomonas aeruginosa, and *Paracoccus denitrificans*. *Applied and Environmental Microbiology,*
45(4), 1247–1253. https://doi.org/10.1128/aem.45.4.1247-1253.1983

Chen, J., & Strous, M. (2013). Denitrification and aerobic respiration, hybrid electron transport chains
and co-evolution. *Biochimica et Biophysica Acta (BBA) - Bioenergetics, 1827*(2), 136–144.
https://doi.org/10.1016/j.bbabio.2012.10.002

Chen, W., Zhang, Y., Zhang, Y., Pi, Y., Gu, T., Song, L., Wang, Y., & Ji, Q. (2018). CRISPR/Cas9-based
genome editing in *Pseudomonas aeruginosa* and cytidine deaminase-mediated base editing in
Pseudomonas species. *iScience, 6*, 222–231. https://doi.org/10.1016/j.isci.2018.07.024

Chislett, M., Guo, J., Bond, P. L., Wang, Y., Donose, B. C., & Yuan, Z. (2022). Reactive nitrogen species
from free nitrous acid (FNA) cause cell lysis. *Water Research*, *217*, 118401.
https://doi.org/10.1016/j.watres.2022.118401

Choi, K.-H., Kumar, A., & Schweizer, H. P. (2006). A 10-min method for preparation of highly
electrocompetent *Pseudomonas aeruginosa* cells: Application for DNA fragment transfer between

chromosomes and plasmid transformation. *Journal of Microbiological Methods, 64*(3), 391–397. https://doi.org/10.1016/j.mimet.2005.06.001

Cole, J. A. (2025). Nitrate reduction for survival in a nanomolar world, not the millimolar world of a laboratory. In *Advances in Microbial Physiology* (Vol. 87, pp. 79–117). Elsevier. https://doi.org/10.1016/bs.ampbs.2025.07.005

Concordet, J.-P., & Haeussler, M. (2018). CRISPOR: Intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Research, 46*(W1), W242–W245. https://doi.org/10.1093/nar/gky354

Cox, R. D. (1980). Determination of nitrate and nitrite at the parts per billion level by chemiluminescence. *Analytical Chemistry, 52*(2), 332–335. https://doi.org/10.1021/ac50052a028

Decleyre, H., Tytgat, B., & Anne, W. (2016). Highly diverse *nirK* genes comprise two major clades that harbour ammonium-producing denitrifiers. *BMC Genomics, 17*(1), 155. https://doi.org/10.1186/s12864-016-2465-0

Ebert, M., Laaß, S., Thürmer, A., Roselius, L., Eckweiler, D., Daniel, R., Härtig, E., & Jahn, D. (2017). FnrL and three Dnr regulators are used for the metabolic adaptation to low oxygen tension in *Dinoroseobacter shibae*. *Frontiers in Microbiology, 8*, 642. https://doi.org/10.3389/fmicb.2017.00642

Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research, 32*(5), 1792–1797. https://doi.org/10.1093/nar/gkh340

Frostegård, Å., Vick, S. H. W., Lim, N. Y. N., Bakken, L. R., & Shapleigh, J. P. (2022). Linking meta-omics to the kinetics of denitrification intermediates reveals pH-dependent causes of N_2O emissions and nitrite accumulation in soil. *The ISME Journal, 16*(1), 26–37. https://doi.org/10.1038/s41396-021-01045-2

Gaimster, H., Alston, M., Richardson, D. J., Gates, A. J., & Rowley, G. (2018). Transcriptional and environmental control of bacterial denitrification and N_2O emissions. *FEMS Microbiology Letters, 365*(5). https://doi.org/10.1093/femsle/fnx277

Gaimster, H., Hews, C. L., Griffiths, R., Soriano-Laguna, M. J., Alston, M., Richardson, D. J., Gates, A. J., & Rowley, G. (2019). A central small RNA regulatory circuit controlling bacterial denitrification and N_2O emissions. *mBio, 10*(4), e01165-19. https://doi.org/10.1128/mBio.01165-19

Gao, Y., Mania, D., Mousavi, S. A., Lycus, P., Arntzen, M. Ø., Woliy, K., Lindström, K., Shapleigh, J. P., Bakken, L. R., & Frostegård, Å. (2021). Competition for electrons favors N_2O reduction in denitrifying *Bradyrhizobium* isolates. *Environmental Microbiology, 23*(4), 2244–2259. https://doi.org/10.1111/1462-2920.15404

Gomila, M., Mulet, M., García-Valdés, E., & Lalucat, J. (2022). Genome-based taxonomy of the genus *Stutzerimonas* and proposal of *S. frequens* sp. nov. and *S. degradans* sp. nov. and emended descriptions of *S. perfectomarina* and *S. chloritidismutans*. *Microorganisms, 10*(7), 1363. https://doi.org/10.3390/microorganisms10071363

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology, 59*(3), 307–321. https://doi.org/10.1093/sysbio/syq010

Härtig, E., Schiek, U., Vollack, K.-U., & Zumft, W. G. (1999). Nitrate and nitrite control of respiratory nitrate reduction in denitrifying *Pseudomonas stutzeri* by a two-component regulatory system homologous to NarXL of *Escherichia coli*. *Journal of Bacteriology, 181*(12), 3658–3665. https://doi.org/10.1128/JB.181.12.3658-3665.1999

Härtig, E., & Zumft, W. G. (1999). Kinetics of *nirS* expression (cytochrome cd1 nitrite reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to denitrification: Evidence for a denitrification-specific nitrate- and nitrite-responsive regulatory system. *Journal of Bacteriology, 181*(1), 161–166. https://doi.org/10.1128/JB.181.1.161-166.1999

Highton, M. P., Bakken, L. R., Dörsch, P., Molstad, L., & Morales, S. E. (2022). Nitrite accumulation and impairment of N$_2$O reduction explains contrasting soil denitrification phenotypes. *Soil Biology and Biochemistry, 166*, 108529. https://doi.org/10.1016/j.soilbio.2021.108529

Hiis, E. G., Vick, S. H. W., Molstad, L., Røsdal, K., Jonassen, K. R., Winiwarter, W., & Bakken, L. R. (2024). Unlocking bacterial potential to reduce farmland N$_2$O emissions. *Nature*, *630*(8016), 421–428. https://doi.org/10.1038/s41586-024-07464-3

Jones, C. M., Stres, B., Rosenquist, M., & Hallin, S. (2008). Phylogenetic analysis of nitrite, nitric oxide, and nitrous oxide respiratory enzymes reveal a complex evolutionary history for denitrification. *Molecular Biology and Evolution, 25*(9), 1955–1966. https://doi.org/10.1093/molbev/msn146

Jüngst, A., Wakabayashi, S., Matsubara, H., & Zumft, W. G. (1991). The *nirSTBM* region coding for cytochrome cd1-dependent nitrite respiration of *Pseudomonas stutzeri* consists of a cluster of mono-, di-, and tetraheme proteins. *FEBS Letters, 279*(2), 205–209. https://doi.org/10.1016/0014-5793(91)80150-2

Kuypers, M. M. M., Marchant, H. K., & Kartal, B. (2018). The microbial nitrogen-cycling network. *Nature Reviews Microbiology, 16*(5), 263–276. https://doi.org/10.1038/nrmicro.2018.9

Lalucat, J., Bennasar, A., Bosch, R., García-Valdés, E., & Palleroni, N. J. (2006). Biology of *Pseudomonas stutzeri*. *Microbiology and Molecular Biology Reviews, 70*(2), 510–547. https://doi.org/10.1128/MMBR.00047-05

Lalucat, J., Gomila, M., Mulet, M., Zaruma, A., & García-Valdés, E. (2022). Past, present and future of the boundaries of the *Pseudomonas* genus: Proposal of *Stutzerimonas* gen. nov. *Systematic and Applied Microbiology, 45*(1), 126289. https://doi.org/10.1016/j.syapm.2021.126289

Li, W., Li, H., Liu, Y., Zheng, P., & Shapleigh, J. P. (2018). Salinity-aided selection of progressive onset denitrifiers as a means of providing nitrite for anammox. *Environmental Science & Technology, 52*(18), 10665–10672. https://doi.org/10.1021/acs.est.8b02314

Li, X., Yang, Z., Wang, Z., Li, W., Zhang, G., & Yan, H. (2022). Comparative genomics of *Pseudomonas stutzeri* complex: Taxonomic assignments and genetic diversity. *Frontiers in Microbiology, 12*, 755874. https://doi.org/10.3389/fmicb.2021.755874

Lim, N. Y. N., Frostegård, Å., & Bakken, L. R. (2018). Nitrite kinetics during anoxia: The role of abiotic reactions versus microbial reduction. *Soil Biology and Biochemistry, 119*, 203–209. https://doi.org/10.1016/j.soilbio.2018.01.006

Lim, N. Y. N., Roco, C. A., & Frostegård, Å. (2016). Transparent DNA/RNA co-extraction workflow protocol suitable for inhibitor-rich environmental samples that focuses on complete DNA removal for transcriptomic analyses. *Frontiers in Microbiology, 7*, 1588. https://doi.org/10.3389/fmicb.2016.01588

Lin, M., Yan, Y., Lu, W., Zhan, Y., Zhang, Y., & Elmerich, C. (2015). Regulatory coupling of nitrogen and
carbon metabolism in nitrogen-fixing *Pseudomonas stutzeri* A1501. In F. J. De Bruijn (Ed.), *Biological*
nitrogen fixation (1st ed., pp. 109–120). Wiley. https://doi.org/10.1002/9781119053095.ch10

Liu, B., Mao, Y., Bergaust, L., Bakken, L. R., & Frostegård, Å. (2013). Strains in the genus *Thauera*
exhibit remarkably different denitrification regulatory phenotypes. *Environmental Microbiology, 15*(10),
2816–2828. https://doi.org/10.1111/1462-2920.12142

Lycus, P., Lovise Bøthun, K., Bergaust, L., Peele Shapleigh, J., Reier Bakken, L., & Frostegård, Å.
(2017). Phenotypic and genotypic richness of denitrifiers revealed by a novel isolation strategy. *The ISME*
Journal, 11(10), 2219–2232. https://doi.org/10.1038/ismej.2017.82

Lycus, P., Soriano-Laguna, M. J., Kjos, M., Richardson, D. J., Gates, A. J., Milligan, D. A., Frostegård,
Å., Bergaust, L., & Bakken, L. R. (2018). A bet-hedging strategy for denitrifying bacteria curtails their
release of N_2O. *Proceedings of the National Academy of Sciences, 115*(46), 11820–11825.
https://doi.org/10.1073/pnas.1805000115

Mania, D., Woliy, K., Degefu, T., & Frostegård, Å. (2020). A common mechanism for efficient N_2O
reduction in diverse isolates of nodule-forming bradyrhizobia. *Environmental Microbiology, 22*(1), 17–31.
https://doi.org/10.1111/1462-2920.14731

Massé, E., & Gottesman, S. (2002). A small RNA regulates the expression of genes involved in iron
metabolism in *Escherichia coli*. *Proceedings of the National Academy of Sciences, 99*(7), 4620–4625.
https://doi.org/10.1073/pnas.032066599

Moeller, S., Payá, G., Bonete, M.-J., Gates, A. J., Richardson, D. J., Esclapez, J., & Rowley, G. (2021).
Microbial small RNAs – The missing link in the nitrogen cycle? *Frontiers in Environmental Science, 9*,
660055. https://doi.org/10.3389/fenvs.2021.660055

Molstad, L., Dörsch, P., & Bakken, L. R. (2007). Robotized incubation system for monitoring gases (O_2,
NO, N_2O, N_2) in denitrifying cultures. *Journal of Microbiological Methods, 71*(3), 202–211.
https://doi.org/10.1016/j.mimet.2007.08.011

Molstad, L., Dörsch, P., & Bakken, L. R. (2016). Improved robotized incubation system for gas kinetics
in batch cultures. Unpublished. https://doi.org/10.13140/RG.2.2.30688.07680

Philippot, L. (2002). Denitrifying genes in bacterial and archaeal genomes. *Biochimica et Biophysica Acta*
(BBA) - Gene Structure and Expression, 1577(3), 355–376. https://doi.org/10.1016/S0167-
4781(02)00420-7

Pold, G., Bonilla-Rosso, G., Saghaï, A., Strous, M., Jones, C. M., & Hallin, S. (2024). Phylogenetics and
environmental distribution of nitric oxide-forming nitrite reductases reveal their distinct functional and
ecological roles. *ISME Communications, 4*(1), ycae020. https://doi.org/10.1093/ismeco/ycae020

Richardson, D. (2025). Life on oxidized nitrogen: The biochemistry and physiology of prokaryotic life
supported by the nitrate-nitrite redox couple. In *Advances in Microbial Physiology* (Vol. 87, pp. 11–77).
Elsevier. https://doi.org/10.1016/bs.ampbs.2025.07.004

Roothans, N., Van Loosdrecht, M. C. M., & Laureni, M. (2025). Metabolic labour division trade-offs in
denitrifying microbiomes. *The ISME Journal, 19*(1), wraf020. https://doi.org/10.1093/ismejo/wraf020

Shapleigh, J. P. (2013). Denitrifying prokaryotes. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes* (pp. 405–425). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-30141-4_71

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J. D., & Higgins, D. G. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology, 7*(1), 539. https://doi.org/10.1038/msb.2011.75

Sistrom, W. R. (1960). A requirement for sodium in the growth of *Rhodopseudomonas spheroides*. *Journal of General Microbiology, 22*(3), 778–785. https://doi.org/10.1099/00221287-22-3-778

Sommer, M., Xie, H., & Michel, H. (2017). *Pseudomonas stutzeri* as an alternative host for membrane proteins. *Microbial Cell Factories, 16*(1), 157. https://doi.org/10.1186/s12934-017-0771-0

Song, M., Zhao, X., Liu, P., Mu, J., He, G., Zhang, C., Tong, S., Xue, C., Zhao, X., Ge, M., & Mu, Y. (2023). Atmospheric NOx oxidation as major sources for nitrous acid (HONO). *NPJ Climate and Atmospheric Science, 6*(1), 30. https://doi.org/10.1038/s41612-023-00357-8

Spiro, S. (2007). Regulators of bacterial responses to nitric oxide. *FEMS Microbiology Reviews, 31*(2), 193–211. https://doi.org/10.1111/j.1574-6976.2006.00061.x

Spiro, S. (2012). Nitrous oxide production and consumption: Regulation of gene expression by gas-sensitive transcription factors. *Philosophical Transactions of the Royal Society B: Biological Sciences, 367*(1593), 1213–1225. https://doi.org/10.1098/rstb.2011.0309

Trunk, K., Benkert, B., Quäck, N., Münch, R., Scheer, M., Garbe, J., Jänsch, L., Trost, M., Wehland, J., Buer, J., Jahn, M., Schobert, M., & Jahn, D. (2010). Anaerobic adaptation in *Pseudomonas aeruginosa*: Definition of the Anr and Dnr regulons. *Environmental Microbiology, 12*(6), 1719–1733. https://doi.org/10.1111/j.1462-2920.2010.02252.x

Vaccaro, B. J., Thorgersen, M. P., Lancaster, W. A., Price, M. N., Wetmore, K. M., Poole, F. L., Deutschbauer, A., Arkin, A. P., & Adams, M. W. W. (2016). Determining roles of accessory genes in denitrification by mutant fitness analyses. *Applied and Environmental Microbiology, 82*(1), 51–61. https://doi.org/10.1128/AEM.02602-15

Van Cleemput, O., & Samater, A. H. (1995). Nitrite in soils: Accumulation and role in the formation of gaseous N compounds. *Fertilizer Research, 45*(1), 81–89. https://doi.org/10.1007/BF00749884

Vollack, K., Härtig, E., Körner, H., & Zumft, W. G. (1999). Multiple transcription factors of the FNR family in denitrifying Pseudomonas stutzeri: Characterization of four fnr-like genes, regulatory responses and cognate metabolic processes. Molecular Microbiology, 31(6), 1681–1694. https://doi.org/10.1046/j.1365-2958.1999.01302.x

Vollack, K.-U., Xie, J., Härtig, E., Römling, U., & Zumft, W. G. (1998). Localization of denitrification genes on the chromosomal map of Pseudomonas aeruginosa. Microbiology, 144(2), 441–448. https://doi.org/10.1099/00221287-144-2-441

Wang, M., Liang, Y., Li, F., Shen, S., Huang, X., & Sun, Y. (2022). Enhancement of biological denitrification by the addition of novel sRNA Pda200 under antibiotic pressure. Bioresource Technology, 365, 128113. https://doi.org/10.1016/j.biortech.2022.128113

Wittorf, L., Jones, C. M., Bonilla-Rosso, G., & Hallin, S. (2018). Expression of nirK and nirS genes in
two strains of Pseudomonas stutzeri harboring both types of NO-forming nitrite reductases. Research in
Microbiology, 169(6), 343–347. https://doi.org/10.1016/j.resmic.2018.04.010

Zajicek, R. S., Allen, J. W. A., Cartron, M. L., Richardson, D. J., & Ferguson, S. J. (2004). Paracoccus
pantotrophus NapC can reductively activate cytochrome cd1 nitrite reductase. FEBS Letters, 565(1–3),
48–52. https://doi.org/10.1016/j.febslet.2004.03.072

Zerbino, D. R., & Birney, E. (2008). Velvet: Algorithms for de novo short read assembly using de Bruijn
graphs. Genome Research, 18(5), 821–829. https://doi.org/10.1101/gr.074492.107

Zhan, Y., Yan, Y., Deng, Z., Chen, M., Lu, W., Lu, C., Shang, L., Yang, Z., Zhang, W., Wang, W., Li, Y.,
Ke, Q., Lu, J., Xu, Y., Zhang, L., Xie, Z., Cheng, Q., Elmerich, C., & Lin, M. (2016). The novel
regulatory ncRNA, NfiS, optimizes nitrogen fixation via base pairing with the nitrogenase gene nifK
mRNA in Pseudomonas stutzeri A1501. Proceedings of the National Academy of Sciences, 113(30).
https://doi.org/10.1073/pnas.1604514113

Zhou, Y., Oehmen, A., Lim, M., Vadivelu, V., & Ng, W. J. (2011). The role of nitrite and free nitrous acid
(FNA) in wastewater treatment plants. Water Research, 45(15), 4672–4682.
https://doi.org/10.1016/j.watres.2011.06.025

Zhuang, J.-L., Sun, X., Zhao, W.-Q., Zhang, X., Zhou, J.-J., Ni, B.-J., Liu, Y.-D., Shapleigh, J. P., & Li,
W. (2022). The anammox-coupled partial-denitrification process in an integrated granular sludge and
fixed-biofilm reactor developed for mainstream wastewater treatment: Performance and community
structure. Water Research, 210, 117964. https://doi.org/10.1016/j.watres.2021.117964

Zumft, W. G. (1997). Cell biology and molecular basis of denitrification. Microbiology and Molecular
Biology Reviews, 61(4), 533–616. https://doi.org/10.1128/mmbr.61.4.533-616.1997

Figure 1. Denitrification kinetics (a–c) and electron flow rates to reductases (d–f) during and after the transition from aerobic respiration to denitrification. Strains were incubated at 20 °C in 120 mL medical vials containing 50 mL Sistrom's mineral medium, supplemented with KNO_3 to an initial concentration of 2 mM (100 µmol vial^{-1}), with He and 1% O_2 in the headspace. The panels show the results for three strains representing the FNA, PNA, and LNA groups. The upper panels show the measured NO_2^- and gases and the lower panels show the calculated rate of electron flow to O_2 (blue), NO_3^- (red), and NO_2^- (purple), with the total electron flow shown as a dashed line. All graphs show mean values from three independent replicates (n = 3); error bars represent standard deviation.

Figure 2. Maximum likelihood phylogenetic tree of selected *Stutzerimonas* strains and their NO₂⁻ accumulation phenotype. The three NO_2^- accumulation groups are indicated by different colors in the phylogenetic tree. Red: Full NO_2^- accumulators (FNA); reduced almost all available NO_3^- before reducing any NO_2^-. Yellow: Partial NO_2^- accumulators (PNA); transient accumulation of NO_2^- reaching approximately 40-60% of the amount of added NO_3^- Blue: Low NO_2^- accumulators (LNA); 0-20% of the reduced NO_3^- was transiently accumulated as NO_2^-. The phylogenetic analysis was a multilocus sequence analysis (MLSA) based on concatenated sequences of partial 16S rRNA and full *gyrB* and *rpoD* gene sequences. In addition to the strains analyzed for denitrification phenotypes, some other strains were included for which complete genome sequence data were publicly available. Only branches with bootstrap values ≥ 700 (of 1000) are shown. The strains were until recently included in a species complex encompassing the former *Pseudomonas stutzeri* and closely related strains (Li et al. 2022) but were recently circumscribed into several species of the novel genus *Stutzerimonas* (Laculat et al. 2022; Gomila et al. 2022). A list of the strains, including their origin and culture collection numbers, is provided in Table S1. A more detailed tree based on genome clustering, including all presently available *Stutzerimonas* genomes, is found in Figure S4.

Figure 3. Denitrification kinetics, gene transcription and electron flow in the FNA strain ZoBell and the LNA strain JM300, switching from aerobic to anaerobic respiration. The panels show measured NO_2^- and gases (a, b), the denitrification gene transcription (c, d), and electron flow rates to reductases (e, f) during and after the transition from aerobic respiration to denitrification. Strains were incubated at 20 °C in 120 mL medical vials containing 50 mL Sistrom's mineral medium, supplemented with KNO_3 to an initial concentration of 2 mM (100 µmol vial^{-1}), with He and 1% O_2 in the headspace. Graphs show mean values from six independent replicates (n = 6); error bars represent standard deviation.

Figure 4. Effect of deleting *narG* and *dnrE* on the phenotype of the PNA strain *Stutzerimonas decontaminans* 19SMN4. The panels show the measured NO_2^- and gases during and after the transition from aerobic respiration to denitrification for the wild type (a), *narG* deletion mutant (b) and *dnrE* deletion mutant (c). Strains were incubated at 20 °C in 120 mL medical vials containing 50 mL Sistrom's mineral medium, supplemented with KNO_3 to an initial concentration of 2 mM (100 µmol vial^{-1}), with He and 1% O_2 in the headspace. Graphs show mean values from three independent replicates (n = 3); error bars represent standard deviation.

Figure 5. Competition for electrons between NO$_3^-$ and NO$_2^-$ reduction pathways. Results for three *Stutzerimonas* strains, each representing one of the three nitrite accumulation phenotypes (a–c). Experiments were performed in biological triplicates (n = 3). The timing of NO$_3^-$ addition varied between replicate vials; therefore, one representative replicate is shown here, while the other replicates, which displayed very similar patterns, are presented in Fig. S7. Cultures were pre-grown anoxically (100% He headspace) in Sistrom's medium with NO$_3^-$ to induce full denitrification proteomes, then transferred at t = 0 to fresh anoxic vials containing only 2 mM NO$_2^-$ (100 µmol vial^{-1}). When approximately one-third of the NO$_2^-$ was reduced, 100 µmol of NO$_3^-$ (corresponding to 2 mM in the medium) was added (indicated by the arrow).

Group	Strain	napABCDE	narGHJIKMXL	dnrE	NirS clade	nirJENY	nirTB	NirK clade	norBCD	nosRZDFYL	Max NO$_2^-$		Max NO		Max N$_2$O	
FNA	CCUG 36651	+	+	+	Clade 1b	downstream	+	clade II	+	+	96.5 ± 0.6		39.3 ± 0.3		22.1 ± 1.0	
	DSM 50227	+	+	+	Clade 1b	downstream	+	clade II	+	+	101.4 ± 5.7		62.5 ± 15.5		7.7 ± 1.7	
	ATCC 17591	+	+	+	Clade 1b	downstream	+	clade II	+	+	92.9 ± 0.4		26.5 ± 4.2		15.8 ± 2.2	
	ZoBell	+	+	+	Clade 1b	downstream	+	−	+	+	108.8 ± 0.25	a	70.5 ± 5.0	abc	11.8 ± 1.2	a
	24a13	+	+	+	Clade 1b	downstream	+	clade II	+	+	71.7 ± 5.7		32.1 ± 2.8		19.7 ± 1.3	
	KC	+	+	+	Clade 1b	downstream	+	−	+	+	96.7 ± 0.2		25.0 ± 0.6		13.7 ± 0.2	
	ST27MN3	+	+	+	Clade 1b	downstream	+	−	+	+	101.5 ± 6.0		20.1 ± 3.4		0.3 ± 0.1	
	CMT.9.A	+	+	+	Clade 1b	downstream	+	−	+	+	89.1 ± 1.6		27.5 ± 2.7		16.1 ± 1.4	
PNA	AN10	+	+	+	Clade 1a	downstream	−	clade II	+	+	62.2 ± 4.8		25.6 ± 5.0		24.9 ± 1.8	
	19SMN4	+	+	+	Clade 1a	downstream	−	−	+	+	44.0 ± 2.5	b	29.3 ± 1.7	abc	25.6 ± 3.0	ab
	CCUG 11256	+	+	+	Clade 1a	downstream	−	−	+	+	43.7 ± 3.3		20.4 ± 1.6		18.0 ± 1.2	
LNA	DNSP21	+	−	−	Clade 1a	downstream	−	−	+	+	1.2 ± 0.01		9.1 ± 0.6		6.5 ± 0.3	
	24a75	+	+	−	Clade 1a	downstream	−	−	+	+	11.3 ± 0.7		47.4 ± 4.3		83.7 ± 44.6	
	DSM 50238	+	+	−	Clade 1a	downstream	−	−	+	+	0.01 ± 0.01		13.9 ± 0.7		56.9 ± 13.6	
	DSM 18231	+	+	−	Clade 1a	upstream	−	clade I	+	+	0.2 ± 0.07	c	43.1 ± 4.2	abc	21.4 ± 5.6	bc
	4C29	+	+	−	Clade 1a	upstream	−	−	+	+	17.1 ± 6.7		41.2 ± 3.0		88.9 ± 26.5	
	28a3	+	−	−	Clade 1a	upstream	−	−	+	+	1.38 ± 0.01		28.4 ± 8.5		52.6 ± 27.5	
	JM300	+	+	−	Clade 1b	upstream	+	clade I	+	+	1.6 ± 0.04		38.6 ± 13.8		67.0 ± 25.7	

Column group headers:
- napABCDE: Periplasmic nitrate red. (NO$_3^-$ to NO$_2^-$)
- narGHJIKMXL, dnrE: Membrane-bound nitrate red. (NO$_3^-$ to NO$_2^-$)
- NirS clade / nirJENY / nirTB: Cytochrome cd$_1$-nitrite red. (NO$_2^-$ to NO)
- NirK clade: Copper nitrite red. (NO$_2^-$ to NO)
- norBCD: c-Nitric oxide red. (NO to N$_2$O)
- nosRZDFYL: Nitrous oxide red. clade I (N$_2$O to N$_2$)
- Max NO$_2^-$, Max NO, Max N$_2$O: Maximum Denitrification intermediates accumulation (NO$_2^-$ µmol vial^{-1}, NO and N$_2$O nmol vial^{-1})

Table 1. Denitrification-related genetic features and maximum accumulation of the intermediate products of 18 *Stutzerimonas* strains. The strains were characterized and grouped as full nitrite accumulators (FNA), partial nitrite accumulators (PNA) or low nitrite accumulators (LNA). The two nitrite reductases NirS and NirK were classified into different clades according to Jones et al. (2008). The genetic organization of the *nirS* related clusters *nirMCFDLGH*, located directly downstream and *nirQOP*, positioned directly upstream of *nirS*, was the same in all strains. These are therefore not mentioned in the table. The *nirJENY* cluster was found either upstream or downstream of *nirS*, depending on the strain. All *nirK* clade I genes were accompanied by the accessory gene *nirV*, located directly downstream of the *nirK* gene, while all *nirK* clade II genes were followed by an extra *norCBQD* gene cluster, not shown in the table. Statistical significance for the maximum denitrification intermediates accumulation was assessed using one-way ANOVAs (p = 0.05). Post hoc multiple comparisons of means were conducted using Tukey´s Honest Significant Difference (HSD; p = 0.05) tests. Statistical differences are indicated where the groups sharing the same letter are not statistically different.