Title: Process for Standardizing and Assessing the Parameters Governing MS2 Virus-Like

Particle Reassembly around Nucleic Acid Cargo

Authors: Daniel de Castro Assumpcao[1], Emma S. Vinokour[2], Madeline M. Mills[1], Shiqi Liang[3],

Carolyn E. Mills[4], Aline Carvalho da Costa[5], Nolan W. Kennedy[1], Danielle Tullman-Ercek[1,*]

Affiliations: [1]Department of Chemical and Biological Engineering, Northwestern University,

Evanston, Illinois, USA

[2]Weinberg College of Arts and Sciences, Northwestern University, Evanston, Illinois, USA

[3]Interdisciplinary Biological Sciences Program, Northwestern University, Evanston,

Illinois, USA

[4]Department of Bioengineering, University of California, Santa Barbara, California, USA

[5]School of Chemical Engineering, University of Campinas, Campinas, Sao Paulo, Brazil

* Corresponding author: 2145 Sheridan Road, Silverman 3619, Evanston, IL 60208;

ercek@northwestern.edu. **Fax:** 847-491-3728. **Phone:** 847-491-7043.

Running title: Reassembly of MS2 Protein Nanoparticles

Pages: 28

Supplementary material pages: 12

Figures: 4

Supplementary material description:

Supplemental Information 1 summarizes literature on MS2 VLP disassembly/reassembly

conditions. Supplemental Information 2 compares linearity across protein quantification

methods. Supplemental Information 3 provides supporting calculations for statistical design of

experiments using reassembly yield as the response.

Keywords: Capsid assembly, design of experiments, self-assembly

Abbreviations:

ANOVA: Analysis of Variance

BCA: Bicinchoninic Acid Assay

CP: Coat Protein

DOE: Design of Experiments

HPLC: High Performance Liquid Chromatography

SDS-PAGE: Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis

SEC: Size Exclusion Chromatography

TEM: Transmission Electron Microscopy

TMAO: Trimethylamine N-Oxide

UV-Vis: Ultraviolet-Visible light spectrum

VLP: Virus-Like Particle

Abstract:

MS2 virus-like particles (VLPs) are widely used as protein nanocages for cargo encapsulation, yet *in vitro* disassembly–reassembly protocols remain poorly standardized, and reassembly yields are reported inconsistently. As a result, the same experiments reported in literature produce widely divergent yields, limiting reproducibility and cross-study comparability. Here, we introduce a cargo-specific, quantitative framework for standardized MS2 VLP reassembly yield determination. We evaluate commonly used disassembly and post-disassembly processing methods and identify practical trade-offs between protein recovery, accessibility, and reproducibility. Reassembly yield is quantified using size exclusion chromatography calibrated against purified VLP standards, enabling robust, cargo-specific yield measurement. Using this framework, we apply a full factorial design of experiments to quantify the individual and combined effects of coat protein concentration, ionic strength, buffer pH, and molecular crowding on reassembly yield. The resulting statistical model explains more than 99% of the explainable variance and its linear fit to the experimental data indicates that optimal reassembly conditions extend beyond those tested to date. Protein concentration and ionic strength dominate reassembly yield, whereas pH and osmolyte concentration contribute more modestly within the tested ranges. Finally, we propose practical guidelines for standardized MS2 VLP disassembly, reassembly, and yield reporting, defining a transferable operating envelope for MS2 VLP reconstruction. While demonstrated here using a single nucleic acid cargo (tr-DNA), the framework is readily extensible to alternative cargos and coat protein variants.

INTRODUCTION

The MS2 virus-like particle (VLP) is a protein-based nanoparticle derived from the

capsid of the male-specific bacteriophage MS2 (*Emesvirus zinderi*) [1]. Composed solely of 180

identical copies of a 129-amino acid coat protein (CP), the MS2 VLP assembles into a 27 nm

icosahedral structure that lacks viral RNA and is non-infectious [2,3]. These protein

nanoparticles can be readily produced in *Escherichia coli* using plasmid-based expression

systems, where the expressed CPs spontaneously self-assemble around RNA molecules,

including their encoding mRNA during translation [4]. Owing to their simplicity, symmetry, and

genetic tractability, MS2 VLPs have become a widely used model in virology and a versatile

platform in nanotechnology. They have been extensively investigated for applications in

imaging, diagnostics, vaccination, and targeted drug delivery [5–7]. In therapeutic delivery, a

common strategy involves disassembling the capsid to remove encapsulated bacterial RNA

("native" cargo), followed by reassembly in a buffered solution containing the cargo of interest

resulting in a protein-based nanoparticle loaded with a desired molecule [8–17]. Despite the

widespread use of this approach, experimental protocols remain highly heterogeneous, and

researchers seldom report standardized metrics for evaluating reassembly efficiency. As a result,

nominally identical MS2 disassembly–reassembly experiments have reported widely divergent

yields, and a standardized method for MS2 disassembly/reassembly, as well as reproducible and

quantitative reassembly yield estimation, have yet to be formalized.

Some alternative strategies avoid disassembly altogether, relying instead on *in vivo*

expression systems to alter the cargo [18], *in vivo* cargo conjugation to the interior of the VLP

[19], or cargo replacement without disassembling the VLP [20]. Nevertheless, the predominant

in vitro protocol for cargo loading involves four main steps: (1) disassembling the MS2 VLP

using acidic conditions to generate free coat protein, (2) removing the encapsulated native

mRNA, (3) adjusting the pH of the coat protein solution, and (4) incubating the coat proteins

with the target cargo under reassembly conditions (**Figure 1**). Despite the widespread use of this

general framework, the specific conditions used in each step vary widely across studies. For

example, incubation time and acid concentration during disassembly differ significantly (e.g. 30

min to 2 h), and reassembly conditions, such as buffer composition, pH, and the inclusion of

crowding agents or salts, also lack standardization. Moreover, few publications clearly define,

quantify, or report reassembly yield in a consistent manner, making it difficult to compare

reassembly efficiencies across different protocols or draw conclusions from the existing

literature (see **Supplemental Information 1** for an extensive comparison between experimental

conditions across literature).

*Figure 1: **Overall process for encapsulating a cargo of interest in MS2 VLPs**. Biologically produced and purified MS2 VLPs*
are first incubated under acidic conditions to induce capsid disassembly and precipitate nucleic acids natively packed during
VLP expression. The precipitated nucleic acids are removed by centrifugation, and the resulting coat protein solution is buffer
exchanged to eliminate excess acid and adjust the pH. Finally, the coat proteins are incubated with the cargo of interest in
buffer, promoting spontaneous self-assembly and encapsulation of cargo with total or local negative charges on its structure.

97   In this study, we systematically investigate each step of the *in vitro* cargo-loading process

for MS2 VLPs, with a focus on the protein-level parameters and solution conditions that

influence disassembly and reassembly. First, we examine how variations in acid incubation time

affect the removal of native RNA. We then discuss alternative methods for adjusting the pH of

the coat protein solution post-disassembly. Next, we evaluate current approaches for calculating

reassembly yield and propose a quantitative, reproducible metric designed to enable cross-study

comparability. Finally, we implement a full factorial design of experiments (DOE) framework to

simultaneously assess the effects of buffer pH, ionic strength, protein concentration, and

molecular crowding on reassembly efficiency. Through this comprehensive analysis, we define a

method for MS2 VLP disassembly–reassembly, highlight the strengths and limitations of

commonly used methods and propose standardized conditions for MS2 VLP disassembly and

reassembly. While this study focuses on a single nucleic acid cargo (tr-DNA), the framework

introduced here is designed to be extensible to alternative cargos and coat protein variants,

providing a transferable foundation for systematic and reproducible MS2 VLP engineering.

RESULTS AND DISCUSSION

Incubation time with acetic acid impacts native cargo removal during disassembly

The first step in loading non-native cargo into MS2 VLPs involves disassembling the

capsid and removing encapsulated native RNA using glacial acetic acid. Here, the acid serves

two purposes: to disassemble the VLP and to precipitate its native nucleic acid cargo. Most

published protocols employ a 2:1 volumetric ratio of glacial acetic acid to VLP solutions with

VLPs typically at 10 mg/mL [8,10–17,21]. However, some studies report using a 3:1 ratio [9]

or lower VLP concentrations (e.g., 4 mg/mL) [16]. Among the reported parameters, incubation

time varies the most significantly across literature, from 30 min [8,10,12,14–17] to 2 h [9] (See

Supplemental Information 1 for a comprehensive comparison). To identify optimal conditions

for VLP disassembly, we considered the three primary goals of this step: (1) complete

dissociation of the capsid into coat protein subunits, (2) efficient precipitation and removal of the

encapsulated native RNA, and (3) maximum possible coat protein concentration after

disassembly. This last goal (maximum coat protein concentration) enables the choice of a wider

range of initial coat protein concentrations in reassembly reactions.

To determine whether the lowest reported acetic acid-to-VLP ratio (2:1 v/v) and the

highest reported VLP concentration (10 mg/mL) are sufficient for complete capsid disassembly,

we analyzed samples before acid treatment and after acid treatment followed by centrifugation

using size-exclusion chromatography (SEC) and transmission electron microscopy (TEM)

(**Figure 2A–B**). The shift in SEC retention time from 8.1 min (intact capsid) to 14.0 min

(disassembled protein) and the absence of intact particles in TEM micrographs indicate that these

conditions fully disassemble the capsids. We confirmed protein integrity preservation after

disassembly with sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE)

(**Figure 2C**).

135
*Figure 2: **Characterization of MS2 VLP disassembly following acid treatment**. A) Size exclusion chromatography (SEC) traces*
showing a shift in retention time before (labelled "VLP") and after (labelled "CP") incubation with acetic acid, indicating
capsid disassembly. A concurrent decrease in A absorbance reflects the removal of native nucleic acid via precipitation
and centrifugation. B) Transmission electron microscopy (TEM) images of VLPs before and after acid incubation, demonstrating
the transition from intact capsids to irregular protein aggregates. C) SDS-PAGE of samples before and after disassembly in
acetic acid indicating that protein does not degrade during disassembly D) A/A absorbance ratio of the protein
solution as a function of incubation time, showing that RNA removal improves with longer incubation and reaches a plateau after
~90 min.

In addition to disassembling the capsid, the low pH generated by glacial acetic acid also

promotes spontaneous precipitation of the native RNA packaged within MS2 VLPs. Efficient

removal of this RNA is a critical step in the disassembly–reassembly process, as residual

negatively charged nucleic acids can trigger unintended, premature self-assembly during pH

neutralization, prior to the introduction of the desired cargo. A simple diagnostic metric to

confirm the removal of nucleic acids from VLP is the <u>u</u>ltra<u>v</u>iolet-<u>vis</u>ible spectroscopy (UV-Vis)

absorbance profiles, particularly ratio between absorbances at 260 nm and 280 nm. Pure RNA

has an A_{260}/A_{280} of approximately 2.1 and pure proteins have an A_{260}/A_{280} of approximately 0.6.

These values are heuristic with widespread use in the field, rather than a theoretical value and

thus, may change according to the molecule composition. For MS2, we observe a drop in the

A_{260}/A_{280} ratio of samples pre-disassembly (A_{260}/A_{280} ~1.65) and post-disassembly followed by

centrifugation (A_{260}/A_{280} ~0.65), suggesting removal of the native cargo (**Figure 2A**). As noted

by Garmann *et al.* [11] , the clarified coat protein solution should ideally have an A_{260}/A_{280} ratio

below 0.67 — preferably closer to 0.60 — to confirm effective removal of RNA.

Nucleic acid precipitation is driven by its protonation under acidic conditions, which

decreases its solubility and enables nucleic acid separation from protein through standard solid–

liquid separation techniques. While a single step centrifugation is used by all the consulted

references, there is no consensus in the operational conditions (**Supplemental Information 1**).

We tested only one condition, in which centrifugation for 15 min at $17,000 \times g$ and 4 °C was

sufficient to pellet nucleic acids and achieve an $A_{260}/A_{280} = 0.65$. Centrifugation followed by

filtration could also be employed if the filter material is compatible with glacial acetic acid.

Chromatography should be avoided as passing insoluble material through a column can cause

clogging and reduce column lifespan.

To assess the impact of VLP-acetic acid incubation time on RNA removal, we measured

A_{260}/A_{280} values following disassembly of 10 mg/mL VLPs in a 2:1 acetic acid solution across

varying incubation durations. Our results show that a 90-min incubation consistently produces

coat protein solutions with A_{260}/A_{280} ratios below 0.65, suggesting efficient RNA removal and

optimal conditions for downstream reassembly (**Figure 2D**). Since not all laboratories have

access to identical equipment, acid incubation time and centrifugation operational conditions

should be tailored to specific experimental setups to ensure that this ratio approaches the ideal

value of 0.60. It is also important to note that very high protein concentrations (>30 mg/mL) may

lead to coat protein loss due to aggregation and should be tested prior to establishing internal

protocols.

Once the capsid is disassembled and the native nucleic acid is removed, the next step is to

eliminate the excess acetic acid. Common methods reported in the literature include desalting

columns [8,12,14,15,17], dialysis [9,13,16], and centrifugal filtration [10] (See **Supplemental**

Information 1 for operational details), each with distinct advantages and limitations. Centrifugal

filtration is the fastest and most cost-effective approach but is also labor-intensive and can

promote premature protein aggregation, especially if local protein concentrations become high

near the filter membrane. Desalting columns offer an alternative approach and can be operated

via either high-performance or gravity flow liquid chromatography. While high-performance

liquid chromatography allows for higher precision, it requires capital equipment that may not be

accessible in all settings. Gravity-based desalting is simple and accessible but can lead to protein

loss due to residual acidity, as some collected fractions may retain low pH values due to the non-

automated nature of the technique and thus should be excluded from reassembly steps. Dialysis

offers the highest protein recovery at a reassembly-compatible pH, is straightforward to perform,

and is inexpensive; however, it is the most time-consuming method. Overall, each of these

approaches is acceptable for acetic acid removal, and the optimal choice depends on laboratory

constraints such as available resources, equipment, and desired throughput. In this work, we

adopted dialysis to maximize protein recovery and ensure precise pH control of the coat protein

solution prior to reassembly.

Another critical variable in this buffer exchange step is the choice of buffer. An effective

buffer should maintain the coat proteins in a disassembled and soluble state, prevent

precipitation, and avoid introducing components that interfere with downstream reassembly.

Most studies recommend acetic acid solutions ranging from 1 mM (pH ~4.4) to 20 mM (pH

~3.7) (See **Supplemental Information 1** for additional buffer compositions). For example,

Ashley *et al.* [9] employed a 10 mM acetic acid buffer supplemented with 50 mM sodium

chloride. While sodium chloride can enhance protein solubility, its effect on disassembled coat

proteins remains unclear. Excess ionic strength may promote premature capsid reassembly in the

absence of cargo, thereby reducing encapsulation efficiency. For this reason, we use a 1 mM

acetic acid solution with no salts as a dialysis buffer. The impact of salt concentration on

reassembly efficiency is addressed in detail below.

Methods for determining reassembly yield and cargo impact on quantification

Before reassembly conditions can be optimized, reliable and standardized methods for

quantifying reassembly yield must be established. Because reassembly rarely reaches 100%

efficiency, the final mixture inevitably contains both assembled VLPs and unassembled coat

protein (CP). Quantification of reassembly yield therefore requires two steps: (1) separation of

assembled VLPs from free CP, and (2) determination of the protein concentrations in the VLP

fraction and the total solution before the reassembly reaction. Among studies that report yield of

reassembly, the most common approaches for this quantification are colorimetric protein assays

such as bicinchoninic acid (BCA) assay; densitometric analysis of protein bands resolved by

SDS–PAGE; and absorbance-based techniques such as UV–vis spectroscopy, coupled or not to

liquid chromatography (*i.e.* SEC) instruments (see **Supplemental Information 1** for methods

details).

The BCA assay is a colorimetric, protein-specific method that quantifies protein

concentration based on the reduction of $Cu^{2\square}$ to $Cu^{\square}$ by peptide bonds and thus not affected by

the presence of nucleic acids. BCA assay works well for quantifying total protein. However, it is

unsuitable for determining true reassembly yield if VLP reassembly is done around another

encapsulated protein because it cannot distinguish between the two. We also considered

densitometric analysis of SDS–PAGE bands, a semi-quantitative approach that correlates band

intensity with protein concentration. Like BCA, this method is also not confounded by nucleic

acids and can therefore provide informative estimates of reassembly yield when protein loading

in the gel is carefully controlled. However, its utility is limited if the cargo is a protein of similar

molecular weight to the MS2 coat protein. SEC appears particularly advantageous for VLPs

because it simultaneously separates assembled from unassembled particles while providing a

measurement (A_{280} peak area) that correlates to capsid concentration, seemingly condensing two

processes into a single procedure. However, as is true for all absorbance-based methods, the

absorbance at 280 nm is influenced by cargo composition. We demonstrated this by

reassembling MS2 around either luciferase RNA (~1900 nt) or a short tr-DNA (19 nt). Following

nuclease digestion of unencapsulated cargo and SEC purification, both samples were brought to

equal protein concentration as determined by BCA assay. When analyzed by SEC (**Figure 3A**),

the luciferase RNA sample (A_{260}/A_{280} = 1.45) produced a stronger A_{280} signal than the tr-DNA

sample (A_{260}/A_{280} = 1.09), despite identical protein concentrations, illustrating how cargo

composition can confound absorbance-based yield quantification. In the case of this procedure,

estimating the yield by comparing the absorbance of unassembled capsid with reassembled

capsid would overestimate yield. Therefore, to establish a reliable correlation between A_{280} peak

area and protein concentration, we recommend using the reassembled product itself as a

calibrant, described in detail below.

Figure 3: **Standardized, cargo-based reassembly yield calculation method**. *A) SEC chromatograms normalized by A280 of two samples containing different nucleic acid-based cargos (varying in amount and nature) but with identical protein concentrations as measured by BCA. The resulting differences in absorbance highlight the interference caused by cargo. B) Schematic of the proposed method for quantifying VLP reassembly yield. The concentrations of the purified VLP (1) and the disassembled capsid (2) are determined using the BCA assay. The reassembly reaction product is analyzed by SEC (3). Additional reassembly reactions are performed to generate a standard curve relating SEC peak area to concentration (4). This standard is then used to convert the SEC peak area of the experimental reassembly product into concentration, enabling calculation of reassembly yield.*

To continue to evaluate which protein concentration method is the most suitable, we assessed the sensitivity and linear dynamic range of BCA and SDS-PAGE methods. We also included UV-Vis based methods, either coupled to SEC or not, and using either the coupled A_{280}

peak area or a NanoDrop^{TM} spectrophotometer, respectively. To this end, we expressed and

purified MS2 VLPs to characterize with each method. We normalized the VLP-containing

solution to $A_{280} = 1$, prepared dilutions spanning A_{280} values from 0.1 to 1.0, and measured each

dilution using all four methods (**Supplemental Information 2**). Among them, NanoDrop^{TM}

UV–vis spectroscopy, SEC, and BCA exhibited lower coefficients of variation between technical

replicates compared to SDS-PAGE densitometry (average coefficient of variance [CV] across

nonzero datapoints: $CV_{UV–Vis} = 0.02$, $CV_{SEC} = 0.008$, $CV_{BCA} = 0.03$, $CV_{Densitometry} = 0.1$).

Moreover, the relationship between expected A_{280} values and observed measurements revealed

that densitometry of SDS–PAGE dilutions had a lower coefficient of determination and higher

normalized root mean square error than the other methods ($R^2_{Densitometry} = 0.886$, $R^2_{BCA} = 0.996$,

$R^2_{UV–Vis} = 0.999$, $R^2_{SEC} = 0.999$ | $RMSE_{Densitometry} = 0.122$, $RMSE_{BCA} = 0.024$, $RMSE_{UV–Vis} =$

0.012, $RMSE_{SEC} = 0.011$). These findings reinforce that densitometry-based methods suffer from

intrinsically higher variability and a narrower linear dynamic range than absorbance-based

techniques [22], rendering the common practice of single-point calibration (i.e., using a single

band of known concentration to proportionally estimate the concentration of another band)

unreliable. Nevertheless, accuracy limitations due to poor linearity can be mitigated by using a

multi-point standard curve generated from purified coat protein and a high–dynamic-range gel

imager. Finally, this dynamic range analysis indicates that the BCA assay does not span as broad

a dynamic range as SEC coupled with A_{280} peak area. However, it remains the only tested

method capable of reporting absolute protein concentration without interference from nucleic

acids or other small molecules, making it a valuable complement to relative quantification

approaches.

Given our findings, we propose a standardized and reproducible method for quantifying

reassembly efficiency (**Figure 3B**) combining both separation of assembled and disassembled

VLPs and yield quantification while addressing variability in how reassembly yield is reported in

the literature. This approach allows for the calculation of either the reassembly yield alone (*Yr*)

or the overall disassembly–reassembly yield (*Yg*), depending on which parameters are measured.

The core of this method involves generating a custom VLP standard specific to the VLP–

cargo system under study. To do this, approximately six parallel reassembly reactions should be

prepared using known amounts of coat protein and cargo to generate a study-specific VLP-cargo

standard. After reassembly, these reactions should be pooled and subjected to centrifugation

using a 100 kDa spin filter to remove unassembled components. The resulting purified sample

represents fully reassembled VLPs that serve as the reference standard. This is then analyzed by

both absorbance at 280 nm (in this case, measured by SEC) and BCA assay to determine protein

concentration. From these two measurements, a conversion factor (*cf*) is calculated by correlating

standard protein concentration (C_S) determined by BCA with the A_{280} peak area (A_S) measured

by SEC (Equation 1):

$$(Equation\ 1)\ \boldsymbol{cf} = \frac{\boldsymbol{C_S}}{\boldsymbol{A_S}}$$

290   Because SEC has a broad linear dynamic range, a single calibration point—run in

291   triplicate—is sufficient. However, for added robustness, performing a short dilution series and

292   fitting a linear regression is recommended. This calibration allows the conversion of SEC A_{280}

293   peak areas from any experimental reassembly sample (*A*) into capsid concentration (*C*) (Equation

294   2):

$$(Equation\ 2)\ \boldsymbol{C} = \boldsymbol{cf} \times \boldsymbol{A}$$

Since the standard and the reassembled samples share the same coat protein and cargo

composition, the calibration factor derived from this standard is directly applicable. Once the

capsid concentration of the reassembled sample (*C*) is determined, the reassembly yield (*Y_r*) can

be calculated using the concentration of disassembled coat protein used in the reassembly

reaction (*C_CP*), the volume of disassembled coat protein used in the reassembly (*V_CP*), the volume

of the reassembly reaction (*V_reassembly*), and the volume injected in the chromatography system

(V_injection) – Equation 3:

$$(Equation\ 3)\ \boldsymbol{Y_r} = \frac{C \times V_{injection}}{C_{CP} \times V_{CP\ used\ in\ reassembly}} \times \frac{V_{reassembly}}{V_{injection}} = \frac{\boldsymbol{cf} \times \boldsymbol{A} \times \boldsymbol{V_{reassembly}}}{\boldsymbol{C_{CP}} \times \boldsymbol{V_{CP\ used\ in\ reassembly}}}$$

It is also important to consider the overall process efficiency, material losses during

handling, and the practical feasibility of the method. To this end, we define the global

disassembly–reassembly yield (*Yg*) metric, which quantifies the fraction of protein VLPs

recovered relative to the initial amount subjected to disassembly. To calculate *Yg* (Equations 4

and 5), the following parameters are required: the concentration of VLPs prior to disassembly

(*C_VLP*), the volume of VLP solution subjected to disassembly (*V_VLP disassembled*), and a correction

factor to account for any volume changes introduced during post-disassembly purification (V_CP

stock after disassembly and V_CP *used in reassembly*):

$$(Equation\ 4)\ \boldsymbol{Y_g} = \frac{C \times V_{injection} \times \dfrac{V_{reassembly}}{V_{injection}} \times \dfrac{V_{CP\ stock\ after\ desalting}}{V_{CP\ used\ in\ reassembly}}}{C_{VLP} \times V_{VLP\ disassembled}}$$

$$(Equation\ 5)\ \boldsymbol{Y_g} = \frac{cf \times A \times V_{reassembly} \times \dfrac{V_{CP\ stock\ after\ desalting}}{V_{CP\ used\ in\ reassembly}}}{C_{VLP} \times V_{VLP\ disassembled}}$$

This standardized framework enables objective, reproducible, and cargo-specific

quantification of VLP reassembly, improving the comparability of results across experiments and

studies. With this method in hand, it becomes possible to systematically investigate how

different variables influence the final reassembly yield.

Impact of protein, crowding agent, sodium chloride concentration and buffer pH on VLP

reassembly yield

Among the steps in the disassembly/reassembly process, the conditions used for

reassembly vary the most. Coat protein concentration ranges from 0.05 µM of coat protein

monomer [23] to 146 µM (2 mg/mL) [17]. The buffer pH ranges from 5.8 [23] to 8.5 [9], but is

not always reported. The sodium content in the buffer ranges from 0 [9,13,21] to 100 mM

[11,12], and some research groups add trimethylamine N-oxide (TMAO) at varying

concentrations as a crowding agent [8,12]. The reassembly incubation temperature varies from 4

°C [8] to 48 °C [23]. Reported incubation time for the reassembly reaction varies from 10 min

[10] to 51 h [16]. A summary of reported reassembly conditions and the corresponding

publications can be found in **Supplemental Information 1.**

To understand the impact of these variables and their interactions on reassembly yield,

we estimated the effect (quantitative influence that an independent variable has on a response) by

performing a full factorial 2^4 design of experiments (DOE) with a central point, in which we

combinatorially tested four factors (variables) with each one assuming two levels (values). The

factors and levels we adopted in the combinatorial design of experiments are coat protein

monomer molarity [10 µM, 30 µM], sodium chloride molarity [0 mM, 200 mM], TMAO

molarity [0 mM, 250 mM], and buffer pH [5,7]. In addition to estimating the effects, we

statistically modeled yield as a function of the tested experimental conditions to predict which

conditions maximize yield. The proposed model is linear and accounts for all main effects of the

independent variables (x_1, x_2, x_3, x_4) as well as their two- and three-way interactions.

$$\hat{y} = b_0 + x_1 b_1 + x_2 b_2 + x_3 b_3 + x_4 b_4 + x_1 x_2 b_{12} + x_1 x_3 b_{13} + x_1 x_4 b_{14} + x_2 x_3 b_{23} + x_2 x_4 b_{24}$$

$$+ x_3 x_4 b_{34} + x_1 x_2 x_3 b_{123} + x_1 x_2 x_4 b_{124} + x_1 x_3 x_4 b_{134} + x_2 x_3 x_4 b_{234}$$

335  The modulus of the coefficients *b* indicates the intensity of the effect, and their signal

336 indicates whether they have a positive or negative effect on yield. All the tested combinatorial

337 conditions necessary for estimating the effects and generating the statistical model are

338 represented in the experimental design matrix (**Supplemental Information 3**)

339  With the experimental results of the tested 17 conditions, we estimated the effects, their

340 statistical significance (**Figure 4A**), the model parameters (**Table S3-5**), and standardized

341 absolute effect (**Figure 4B**). Yield was quantified using the methodology developed in this

342 study. Because the VLP standard for yield estimation was generated under the central-point

343 condition (condition 17), yields for conditions with higher packing efficiency are likely

344 overestimated and those with lower packing efficiency underestimated. However, this systematic

345 bias is consistent across conditions and does not affect the interpretation of the comparative

346 results. The interaction parameters protein-TMAO (*x1x2*), protein-TMAO-pH (*x1x2x4*), protein-

347 NaCl-pH (*x1x3x4*) and TMAO-NaCl-pH (*x2x3x4*) are not statistically significant with a

348 significance level $\alpha = 0.05$ and were thus removed from the proposed model. We tested the

349 model significance by performing an ANOVA, testing the null hypothesis H_0 that all regression

350 coefficients (except *b₀*) are 0. The ratio between the regression mean square and the residual

351 mean square ($QM_R = 0.254802$, $QM_r = 0.000563$, $QM_R/QM_r = 452.93$) was higher than a critical

352 F-value ($v_1 = 10$, $v_2 = 40$, $\alpha = 0.05$, $Fv_1, v_2 = 2.077$), and the ratio of the mean square for lack of fit

353 and the mean square for pure error ($QM_{pe} = 0.000514$, $QM_{lof} = 0.000838$, $QM_{pe}/QM_{lof} = 1.632$)

354 was below a critical F-value ($v_1 = 6$, $v_2 = 34$, $\alpha = 0.05$, $Fv_1, v_2 = 2.380$). The results of the F-tests

in conjunction with a model with coefficient of determination $R^2 = 0.9912$ suggest that the model

is statistically significant and adequately fits the data without significant lack of fit (**Figure 4C**).

*Figure 4: **Determinants of Reassembly Efficiency.** A) Reassembly yield for each individual replicate of each tested condition. B)*
Pareto chart of standardized effects showing the influence of individual factors and their interactions on yield. The vertical
dashed line represents the critical t-value for statistical significance. C) Experimental versus predicted yield from the fitted
model. Each point represents one replicate. The dashed line corresponds to the ideal fit (y = x), indicating strong agreement
between experimental data and model predictions. D) Marginal means plot indicating the impact of protein molarity, sodium
chloride   molarity,   and   pH   and   their   interactions   on   reassembly   yield

The effect of protein concentration on the VLP reassembly yield is interconnected with

other variables. While the main effect of protein concentration is one of the strongest in the

model (0.232 ± 0.007, p-value = 3 ×10^{-27}), it also had a strong negative interaction effect with

sodium chloride (-0.1189 ± 0.007, p-value = 3 ×10^{-15}) and a weaker negative interaction effect

with pH (-0.0410 ± 0.007, p-value = 3 ×10^{-6}). These interaction effects indicate that the effect of

protein concentration is pronounced in certain conditions (*e.g.* pH 5, sodium chloride = 0 mM),

but minimal in others (*e.g.* pH 7, sodium chloride = 200 mM) (**Figure 4D**). Conversely, sodium

chloride had a consistently negative effect on reassembly, suggesting that 200 mM sodium

chloride is too high of a concentration for efficient reassembly. We hypothesize that this is

because elevated ionic strength disrupts the electrostatic interactions between coat proteins that

are necessary for proper nucleation and propagation of capsid structures [24]. Interestingly, the

magnitude of the interaction between protein concentration and sodium chloride concentration

exceeds the independent effects of pH and TMAO, highlighting their comparatively greater

impact on yield.

The effects of pH and the crowding agent TMAO on reassembly yield are more subtle.

Within the ranges tested, these parameters produced a milder influence on reassembly yield,

suggesting the reassembly process tolerates moderate pH shifts and osmolyte levels without

negatively impacting reassembly yield, within the tested range. This finding complements the

understanding of TMAO's impact in MS2 VLP reassembly reported by our own research group.

Glasgow *et al.* [12] reported that increasing TMAO concentration from 0 to 250 mM in

reassembly reactions with 30 µM coat protein monomer, 3:5 protein monomer to cargo (tr-DNA)

molar ratio, and 100 mM of sodium chloride increased the reassembly from approximately 10%

to 50%. For the same coat protein monomer concentration (30 µM), sodium chloride

concentration (100 mM), and pH (7), but for a coat protein monomer to tr-DNA cargo ratio of

2:1, our model predicts a reassembly yield variation from 34% to 38%. This difference suggests

that the interaction between TMAO concentration and cargo concentration or protein to cargo

ratio has a significant impact on reassembly yield. It is important to note that the method utilized

to estimate yield in both studies is different, highlighting the necessity of the standardized

method for measuring reassembly yield developed earlier in this manuscript.

Another important outcome of the model is the absence of curvature in the response

surface, which indicates that the optimal conditions for reassembly are not within the range of

conditions tested here. If an optimal set of conditions had been captured within the experimental

space, we would expect to see a nonlinear model with maxima. Instead, the linear behavior

observed suggests that the best conditions lie outside of the ranges reported in the literature and

explored to date. While it would be possible to work with higher protein and TMAO

concentrations, lowering the pH below that used in this work could result in buffer conditions

close to those in which the reassembled capsid is not stable. While lowering the pH may be

prohibitive, the DOE results imply that the field has not yet converged on optimal reassembly

parameters, and that established conditions represent only partial, rather than maximal,

efficiency. Another important observation is that these experiments were performed using tr-

DNA as a cargo and physicochemical interactions between the tested conditions and other cargos

may vary drastically. Therefore, extrapolating the results here to other MS2 VLP-cargo systems

should be done carefully and additional studies with an expanded set of variables is necessary to

determine optimal conditions. Importantly, while each new cargo may require its own

optimization, the present findings highlight that not all parameters will need to be reinvestigated

from the underlying basics. Overall, although perfect reassembly conditions remain elusive, we

identify key parameters that help guide future efforts, and the quantitative yield framework

introduced here finally provides a foundation for systematic, cargo-specific optimization going

forward.

Based on the experimental results presented here, we propose a standardized workflow

for MS2 VLP disassembly and reassembly with nucleic acid cargo to improve reproducibility

and comparability across studies. Purified MS2 VLPs should be disassembled at high protein

concentration (10 mg/mL) using a 2:1 (v/v) ratio of glacial acetic acid for 90 min at 4 °C,

followed by removal of precipitated native RNA by centrifugation ($\geq$17,000 $\times$ g, 15 min, 4 °C)

and verification of RNA clearance by $A_{260}/A_{280} \leq 0.65$. Excess acid should then be removed via

dialysis against 1 mM acetic acid without added salt. We recommend reassembling the VLP in a

buffer at pH 5 with no sodium chloride and 250 mM of TMAO. The initial concentration of Coat

Protein in a reassembly reaction should be close to 50 µM and the reaction should be carried at 4

°C for 48 h. Finally, reassembly yield should be quantified using cargo-specific SEC calibration

curves anchored to absolute protein concentration measured by BCA. Adoption of this workflow

provides a practical, experimentally grounded baseline for MS2 VLP reconstruction and enables

systematic extension to alternative cargos and assembly conditions.

CONCLUSION

In this study, we dissected the *in vitro* disassembly–reassembly process of MS2 VLPs to

identify parameters that influence efficiency and reproducibility. We show that using a 2:1 acetic

acid-to-VLP ratio at high protein concentrations with a 90-min incubation provides robust

disassembly and RNA clearance. We also highlight the importance of selecting appropriate

methods for buffer exchange, as trade-offs exist between protein recovery, accessibility, and

processing time. Our comparison of quantification methods for reassembly yield reveals that

electrophoresis and spectroscopy alone are insufficient for accurate determination; instead, we

propose a standardized, cargo-specific calibration framework using HPLC-SEC coupled with

BCA. Through factorial DOE analysis, we demonstrate that protein concentration is the

dominant factor driving reassembly efficiency, whereas sodium chloride has a strongly negative

effect, and pH and TMAO contribute more modestly. Notably, the absence of curvature in our

response surfaces suggests that the optimal conditions for reassembly have not yet been defined

within the literature and lie beyond the ranges typically reported.

These findings underscore two key points: (i) methodological variation has hindered

progress toward reproducible, high-yield encapsulation, and (ii) the field would benefit from

adopting standardized conditions and yield reporting practices. More broadly, while individual

reassembly parameters have been explored empirically in prior studies, the field has largely

lacked a quantitative approach capable of evaluating multiple variables and their interactions

simultaneously. Although design of experiments (DOE) methodology has existed for decades,

their application remains uncommon in the VLP assembly literature, where most studies vary

one factor at a time. By implementing a statistically rigorous full factorial DOE, we show that

reassembly yield is dominated by protein concentration and ionic strength, with comparatively

modest contributions from osmolyte concentration under the tested conditions. Importantly,

this analysis reveals interaction effects that are not accessible through one-variable-at-a-time

approaches. These findings emphasize that a DOE-based approach is necessary to understand

the cumulative and interacting effects of multiple physicochemical variables on reassembly

yield.

We also acknowledge the limitations of this work, which include the focus on a single

type of cargo (tr-DNA) and omission of other variables such as temperature, incubation time, or

different cargo-to-coat protein ratios. Numerous studies have demonstrated successful MS2 VLP

reassembly with a range of cargos, including diverse nucleic acids and proteins (summarized in

Supplemental Information 1). However, even among studies employing the same cargos and

reassembly conditions, reported yields vary substantially [12, 15], and in some cases differ by

nearly twofold. This variability highlights the challenges associated with comparing results

across literature and emphasizes the need for standardized experimental and yield-quantification

frameworks. By introducing a cargo-specific, SEC-based calibration strategy anchored to

absolute protein concentration, this work provides a common quantitative baseline that facilitates

reproducibility, cross-study comparability, and rational troubleshooting of reassembly

workflows. Future studies expanding these parameters and testing diverse cargos will be

essential for developing broadly applicable guidelines. To that end, our study provides a

systematic framework to improve reproducibility, comparability, and ultimately the reliability of

MS2 VLPs as a versatile nanotechnology platform. While developed here for MS2 VLPs, this

framework is readily extendable to other VLP systems that rely on disassembly–reassembly

strategies.

MATERIALS AND METHODS

MS2 VLP expression: The strain used in this experiment is the same as that described by

Robinson *et al* [25]. It consists of *E. coli* DH10B cells harboring the MS2 coat protein gene

cloned into a pBAD33 vector. Key features of this plasmid include a chloramphenicol resistance

cassette for selection and an arabinose-inducible pBAD promoter for controlled expression of the

MS2 gene. Cells from a 30% glycerol stock were used to inoculate a 125 mL baffled shake flask

containing 12 mL of autoclaved Lysogeny Broth (LB-Miller formulation) supplemented with

34 μg/mL chloramphenicol. Cultures were grown overnight (14–16 h) at 37 °C with shaking

at 225 rpm. 10 mL of this overnight culture were used to inoculate a 2.8 L baffled shake flask

containing 1 L of autoclaved LB-M medium and 1 mL of a 34 mg/mL chloramphenicol stock

solution. The culture was incubated at 37 °C with shaking at 225 rpm until it reached an optical

density at 600 nm (OD) of ~0.6. At this point, VLP expression was induced by adding

L(+)-arabinose to a final concentration of 0.02% (w/v). The culture was incubated for an

additional 16–20 h at 37 °C and 225 rpm. Cells were harvested by centrifugation at $4,800 \times g$

for 10 min at 4 °C, and the supernatant was discarded. The resulting pellet was placed on ice to

halt further protein expression and resuspended in 50 mL of SEC buffer (10 mM sodium

phosphate, 200 mM sodium chloride, pH 7.2). If not purified immediately, cell suspension was

stored at −80 °C for later use.

MS2 VLP purification: To recover and purify MS2 capsid proteins, the resuspended cell pellets

were lysed using a probe sonicator (Fisher Scientific, Model FB120) set to 50% amplitude for 10

min (2-second pulses followed by 4-second intervals, for a total sonication program duration of

30 min), with samples kept on ice throughout sonication to prevent overheating. Lysates were

clarified by centrifugation at $17,000 \times g$ for 10 min at 4 °C, and the pellet containing cell debris

was discarded. The resulting supernatant was mixed 1:1 (v/v) with saturated ammonium sulfate

aqueous solution and incubated on a vibrating platform at 4 °C for at least 2 h to precipitate

protein. Following incubation, samples were centrifuged again at $17,000 \times g$ for 10 min at 4 °C.

The supernatant was discarded, and the resulting pellet was resuspended in 3 mL of 1× SEC

buffer. To remove residual ammonium sulfate, the sample was dialyzed overnight at 4 °C

using a 7,000 molecular weight cut off (MWCO) (Snakeskin) dialysis membrane against 4 L of

1× SEC buffer. The dialyzed sample was clarified by centrifugation ($21,000 \times g$, 5 min, 4 °C),

followed by filtration through a 0.45 μm syringe filter. For purification, the sample was

subjected to Fast Protein Liquid Chromatography (FPLC) using an ÄKTA Pure system (Cytiva)

equipped with a HiPrep 16/60 Sephacryl S-500 HR column, using 1x SEC buffer as the mobile

phase run at a flow rate of 1 mL/min at room temperature. Up to 5 mL of sample was injected

per run, and MS2 capsid elution was monitored at A_{280}, consistently appearing between 60 and

80 mL of elution volume.

High-Performance Liquid Chromatography (HPLC): VLP reassembly reactions were

assessed using an HPLC system (Agilent 1260 infinity II) and an SEC column (Yara 4000,

Phenomenex). The mobile phase consisted of $1 \times$ SEC buffer (10 mM phosphate, 200 mM NaCl,

pH 7.2). Runs were carried out at room temperature by injecting 100 µL of sample at the time

with a total method length of 25 min. VLPs eluted between 7.7 and 9.0 min.

Transmission Electron Microscopy: 400-mesh copper grids with Formvar/carbon film (EMS

P/N FCF400-Cu-50) were hydrophilized via glow discharge. We applied 10 µL of 150 nM

purified capsid solution for 5 seconds, then blotted with filter paper. A 10 µL aliquot of 1%

(w/v) uranyl acetate was added and immediately wicked away; this step was repeated once. A

final 10 µL of stain was applied, incubated for 4 min, and then blotted. Grids were imaged on

a JEOL 1400 Flash TEM with a Gatan OneView camera.

Sodium Dodecyl Sulfate–Polyacrylamide Gel Electrophoresis (SDS-PAGE): Proteins were

resolved on 12.5% polyacrylamide gels prepared with a standard Laemmli formulation. Samples

were mixed with Laemmli buffer containing 10% β-mercaptoethanol, heated at 95 °C for 5 min,

and loaded alongside a molecular weight ladder (EZ-run, Fisher Scientific). Electrophoresis was

carried out at 150 V for ~1 h in $1\times$ SDS running buffer until the dye front reached the bottom of

the gel. Protein gels were scanned using Azure biosystems 600 and densitometry analysis was

performed utilizing Fiji software (Schindelin et al., 2012).

Bicinchoninic Acid Assay. Protein concentrations were determined using the bicinchoninic acid

(BCA) assay (Pierce BCA Protein Assay Kit, Thermo Scientific) according to the manufacturer's

instructions, using a 96 well plate and BSA standards.

MS2 VLP disassembly and reassembly: Native MS2 VLPs (≤1 week old) were concentrated to

$A_{280} > 25.0$ (approximately 10 mg/mL) using a 100 kDa molecular weight cutoff centrifugal

filter (Amicon Ultra-15, Sigma-Aldrich) spun at 3000 × g in a swinging-bucket rotor.

Disassembly was initiated by adding glacial acetic acid at a 2:1 (v/v) acid-to-capsid ratio,

followed by incubation on ice for 1.5 h. Samples were centrifuged at 17,000 × g for 15 min at 4

°C to pellet residual nucleic acids. The clarified supernatant, containing disassembled capsids,

was dialyzed at 4 °C against 1.5 L of chilled 1 mM acetic acid using a 3.5 kDa molecular weight

cutoff dialysis cassette. Dialysis was performed in two steps: 3 h in fresh buffer, followed by

overnight dialysis. Successful RNA removal was confirmed by an A_{260}/A_{280} absorbance ratio of

~0.6. Reassembly conditions were varied as noted in the results section. In all cases, water,

buffer, salt, crowding agent, cargo, and protein were added at listed concentrations in the order

listed here, and incubated for 48 h at 4 °C. The samples were centrifuged at 17,000 × g for 15

min and supernatant collected for further analysis.

Supplementary material description

Supplemental Information 1 summarizes literature on MS2 VLP disassembly/reassembly

conditions. Supplemental Information 2 compares linearity across protein quantification

methods. Supplemental Information 3 provides supporting calculations for statistical design of

experiments using reassembly yield as the response. Supplemental Information 4 contains raw

data and results of statistical design of experiments using A_{260}/A_{280} as the response.

Acknowledgements

This work was supported by the National Science Foundation (CBET-2043973, to D.T.-E.).

D.C.A. received support from the Chemistry of Life Processes Predoctoral Training Program at

Northwestern University (NIH T32GM149439). M.M.M. was supported in part by the

Northwestern University Graduate School Cluster in Biotechnology, Systems, and Synthetic

Biology, which is affiliated with the Biotechnology Training Program. S.L. was partially

supported by the Molecular Biophysics Training Program at Northwestern University (NIH

5T32GM140995-04). A.C.C. was supported by Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq grant number 303063/2022-0). N.W.K. was supported by the

U.S. Army Research Office (W52P1J-21-9-3023 P00003/CS-21-0701-001).

Author's Contributions (CRediT)

D.C.A.: Conceptualization (lead), methodology (lead), validation (lead), formal analysis (equal),

investigation (equal), resources (supporting), data curation (lead), writing – original draft (lead);

writing – review and edit (supporting), visualization (lead), project administration (equal),

funding acquisition (supporting). **E.S.V.:** validation (supporting), formal analysis (supporting),

investigation (equal), visualization (support). **M.M.M.:** investigation (supporting), data curation

(supporting) **S.L.:** investigation (supporting) **C.E.M.:** writing – review and edit (equal) **A.C.C.:**

formal analysis (equal), data curation (supporting) **N.W.K.:** writing – review and edit (equal),

visualization (supporting) **D.T.-E.:** writing – review and edit (equal), project administration

(equal), funding acquisition (lead).

References

[1] J. Callanan, S.R. Stockdale, E.M. Adriaenssens, J.H. Kuhn, J. Rumnieks, M.J. Pallen, A.N.
Shkoporov, L.A. Draper, R.P. Ross, C. Hill, Leviviricetes: expanding and restructuring the
taxonomy of bacteria-infecting single-stranded RNA viruses, Microb. Genomics 7 (2021)
000686. https://doi.org/10.1099/mgen.0.000686.

[2] R. Golmohammadi, K. Valegård, K. Fridborg, L. Liljas, The Refined Structure of
Bacteriophage MS2 at 2·8 Å Resolution, J. Mol. Biol. 234 (1993) 620–639.
https://doi.org/10.1006/jmbi.1993.1616.
[3] J.H. Strauss, R.L. Sinsheimer, Purification and properties of bacteriophage MS2 and of its
ribonucleic acid, J. Mol. Biol. 7 (1963) 43–54. https://doi.org/10.1016/S0022-
2836(63)80017-0.
[4] E.C. Hartman, C.M. Jakobson, A.H. Favor, M.J. Lobba, E. Álvarez-Benedicto, M.B. Francis, D.
Tullman-Ercek, Quantitative characterization of all single amino acid variants of a viral
capsid-based drug delivery vehicle, Nat. Commun. 9 (2018) 1385.
https://doi.org/10.1038/s41467-018-03783-y.
[5] K.M. Frietze, D.S. Peabody, B. Chackerian, Engineering virus-like particles as vaccine
platforms, Curr. Opin. Virol. 18 (2016) 44–49.
https://doi.org/10.1016/j.coviro.2016.03.001.
[6] Y. Fu, J. Li, A novel delivery platform based on Bacteriophage MS2 virus-like particles, Virus
Res. 211 (2016) 9–16. https://doi.org/10.1016/j.virusres.2015.08.022.
[7] B. Ikwuagwu, D. Tullman-Ercek, Virus-like particles for drug delivery: a review of methods
and applications, Curr. Opin. Biotechnol. 78 (2022) 102785.
https://doi.org/10.1016/j.copbio.2022.102785.
[8] I.L. Aanei, T. Huynh, Y. Seo, M.B. Francis, Vascular Cell Adhesion Molecule-Targeted MS2
Viral Capsids for the Detection of Early-Stage Atherosclerotic Plaques, Bioconjug. Chem. 29
(2018) 2526–2530. https://doi.org/10.1021/acs.bioconjchem.8b00453.
[9] C.E. Ashley, E.C. Carnes, G.K. Phillips, P.N. Durfee, M.D. Buley, C.A. Lino, D.P. Padilla, B.
Phillips, M.B. Carter, C.L. Willman, C.J. Brinker, J. do C. Caldeira, B. Chackerian, W.
Wharton, D.S. Peabody, Cell-Specific Delivery of Diverse Cargos by Bacteriophage MS2
Virus-like Particles, ACS Nano 5 (2011) 5729–5745. https://doi.org/10.1021/nn201397z.
[10] L.A. Williams, A. Neophytou, R.F. Garmann, D. Chakrabarti, V.N. Manoharan, Effect of
coat-protein concentration on the self-assembly of bacteriophage MS2 capsids around
RNA, Nanoscale 16 (2024) 3121–3132. https://doi.org/10.1039/D3NR03292B.
[11] R.F. Garmann, A.M. Goldfain, V.N. Manoharan, Measurements of the self-assembly
kinetics of individual viral capsids around their RNA genome, Proc. Natl. Acad. Sci. 116
(2019) 22485–22490. https://doi.org/10.1073/pnas.1909223116.
[12] J.E. Glasgow, S.L. Capehart, M.B. Francis, D. Tullman-Ercek, Osmolyte-Mediated
Encapsulation of Proteins inside MS2 Viral Capsids, ACS Nano 6 (2012) 8658–8664.
https://doi.org/10.1021/nn302183h.
[13] T. Sugiyama, R.R. Hebert, K.A. Hartman, Ribonucleoprotein complexes formed between
bacteriophage MS2 RNA and MS2 Protein in vitro, J. Mol. Biol. 25 (1967) 455–463.
https://doi.org/10.1016/0022-2836(67)90198-2.
[14] M. Wu, W.L. Brown, P.G. Stockley, Cell-specific delivery of bacteriophage-encapsidated
ricin A chain, Bioconjug. Chem. 6 (1995) 587–595. https://doi.org/10.1021/bc00035a013.
[15] M. Yang, G. Rao, L. Li, L. Qi, C. Ma, H. Zhang, J. Gong, B. Wei, X.-E. Zhang, G. Chen, S. Cao,
F. Li, Transformation of a Viral Capsid from Nanocages to Nanotubes and Then to
Hydrogels: Redirected Self-Assembly and Effects on Immunogenicity, ACS Nano 18 (2024)
13755–13767. https://doi.org/10.1021/acsnano.4c01969.

[16] L. Zhang, Y. Sun, L. Chang, T. Jia, G. Wang, R. Zhang, K. Zhang, J. Li, A novel method to produce armored double-stranded DNA by encapsulation of MS2 viral capsids, Appl. Microbiol. Biotechnol. 99 (2015) 7047–7057. https://doi.org/10.1007/s00253-015-6664-4.

[17] W. Zhou, S. Ma, R. Gao, Y. Tang, H. Zhang, A. Liang, M. Yang, C. Ma, Q. Fan, X.-E. Zhang, F. Li, Assembly of Matryoshka-Type Protein Nanocages for Compartmentalized Oxygen Sensing, Nano Lett. 25 (2025) 4433–4440. https://doi.org/10.1021/acs.nanolett.4c06699.

[18] B.L. Pasloske, C.R. Walkerpeach, R.D. Obermoeller, M. Winkler, D.B. DuBois, Armored RNA technology for production of ribonuclease-resistant viral RNA controls and standards, J. Clin. Microbiol. 36 (1998) 3590–3594. https://doi.org/10.1128/JCM.36.12.3590-3594.1998.

[19] S. Liang, K. Butaney, D. de C. Assumpção, J. Jung, N.W. Kennedy, D. Tullman-Ercek, Synthetic Rewiring of Virus-Like Particles via Circular Permutation Enables Modular Peptide Display and Protein Encapsulation, (2025) 2025.06.13.659512. https://doi.org/10.1101/2025.06.13.659512.

[20] T.N. Szyszka, R. Siddiquee, A. Loustau, L.S.R. Adamson, C. Rennie, T. Huang, R. Young, A. Care, Y.H. Lau, High-Fidelity In Vitro Packaging of Diverse Synthetic Cargo into Encapsulin Protein Cages, Angew. Chem. Int. Ed. 64 (2025) e202422459. https://doi.org/10.1002/anie.202422459.

[21] P.G. Stockley, O. Rolfsson, G.S. Thompson, G. Basnak, S. Francese, N.J. Stonehouse, S.W. Homans, A.E. Ashcroft, A Simple, RNA-Mediated Allosteric Switch Controls the Pathway to Formation of a T=3 Viral Capsid, J. Mol. Biol. 369 (2007) 541–552. https://doi.org/10.1016/j.jmb.2007.03.020.

[22] W. Holzmüller, U. Kulozik, Protein quantification by means of a stain-free SDS-PAGE technology without the need for analytical standards: Verification and validation of the method, J. Food Compos. Anal. 48 (2016) 128–134. https://doi.org/10.1016/j.jfca.2016.03.003.

[23] O. Rolfsson, K. Toropova, V. Morton, S. Francese, G. Basnak, G.S. Thompson, S.W. Homans, A.E. Ashcroft, N.J. Stonehouse, N.A. Ranson, P.G. Stockley, RNA packing specificity and folding during assembly of the bacteriophage MS2, Comput. Math. Methods Med. 9 (2008) 339–349. https://doi.org/10.1080/17486700802168445.

[24] J.D. Perlmutter, M.R. Perkett, M.F. Hagan, Pathways for virus assembly around nucleic acids, J. Mol. Biol. 426 (2014) 3148–3165. https://doi.org/10.1016/j.jmb.2014.07.004.

[25] S.A. Robinson, E.C. Hartman, B.C. Ikwuagwu, M.B. Francis, D. Tullman-Ercek, Engineering a Virus-like Particle to Display Peptide Insertions Using an Apparent Fitness Landscape, Biomacromolecules 21 (2020) 4194–4204. https://doi.org/10.1021/acs.biomac.0c00987.