1 **An Expectation and Maximization Algorithm for Multivariate Genome-wide Association**
2 **Studies (EMmvGWAS)**

3 Chin-Sheng Teng[1], Xuesong Wang[2], Cheng Liu[3], Qishan Wang[3], Yanru Cui[4] and Shizhong Xu[2*]
4
5 [1]Department of Statistics, University of California, Riverside, CA 92521, USA
6 [2]Department of Botany and Plant Sciences, University of California, Riverside, CA 92521, USA
7 [3] Department of Animal Science, College of Animal Sciences, Zhejiang University, 866#
8 Yuhangtang Road, Hangzhou 310058, China
9 [4]College of Agronomy, Hebei Agricultural University, Baoding 071001, China
10
11 *shizhong.xu@ucr.edu
12

Abstract

Genome-wide association studies (GWAS) commonly focus on one quantitative trait at a time,
even when multiple traits are collected. However, joint analysis of multiple traits can increase the
power to detect genetic associations by leveraging trait correlations. Multivariate GWAS is
particularly important for identifying pleiotropic effects and uncovering shared genetic
architecture underlying complex traits. Despite its great potential, multivariate GWAS methods
face substantial computational challenges due to the high dimensionality of polygenic covariance
structures and the cost of scanning genome-wide markers. We present EMmvGWAS, an efficient
computational framework implemented as an R, which dramatically reduces computation time
for multivariate GWAS involving a moderate number of traits. The algorithm estimates the
genetic and environmental covariance matrices from a pure polygenic model using an
Expectation-Maximization (EM) algorithm. To improve computational efficiency during
genome-wide scanning, we use a semi-exact method that fixes the estimated covariance matrices
and treats their ratio, defined as the genetic covariance matrix multiplied by the inverse of the
environmental covariance matrix, as a known constant across all markers. This approach enables
closed-form solutions for marker effects at each locus, dramatically reducing the computation
time without compromising statistical power. We demonstrate the scalability and performance of
our method through both simulation studies and real dataset analyses from rice, mouse and
human populations. The method is implemented in R and is designed to support multivariate
GWAS involving a moderate number of traits, while also accommodating univariate analyses as
a special case. The R package is available at https://github.com/Jason-Teng/EMmvGWAS.

39 **Keywords:** Expectation and maximization algorithm, Genome-wide association studies,
40 Multivariate linear mixed model, Restricted maximum likelihood method.

1. Background

Genome-wide association studies (GWAS) have significantly advanced our understanding of the genetic architecture underlying complex traits by identifying genetic variants associated with quantitative traits (Visscher et al., 2017). Traditionally, GWAS are often conducted on individual traits independently, even though multiple correlated phenotypes are always measured in genetic studies. Analyzing traits individually overlooks valuable trait correlations, thus potentially reducing the statistical power and increasing the type 1 error rate. Multivariate GWAS leverages these correlations by using multivariate linear mixed models, improving power and reducing false positive associations in identifying pleiotropic genetic variants, which affect multiple traits simultaneously (Korte et al., 2012; Porter & O'Reilly, 2017; Turley et al., 2018; Zhou & Stephens, 2014). Such methods are crucial for uncovering shared genetic bases among related traits, providing deeper insights into biological pathways and mechanisms underlying complex diseases and phenotypes (Hackinger & Zeggini, 2017).

Despite their clear advantages, multivariate GWAS methods confront significant computational hurdles, primarily due to the complexity of modeling high-dimensional covariance structures and the computational cost associated with scanning genome-wide markers. Estimating genetic and environmental covariance matrices between traits becomes increasingly demanding as the number of traits grows, often requiring iterative optimization over large parameter spaces. At the same time, genome-wide marker scanning involves evaluating potentially millions of SNPs across all traits, which remains computationally intensive. These two challenges compound in high-dimensional settings, making many existing multivariate GWAS tools impractical for real-world applications involving many traits. Although R is widely used in statistical genetics, there is a lack of efficient R packages for multivariate GWAS, especially for analyses involving many traits.

To address these challenges, we developed EMmvGWAS, a general statistical framework that supports multivariate GWAS with improved computational efficiency. While suitable for small trait sets, EMmvGWAS offers substantial speed improvements as the number of traits grows, making it especially effective for studies involving 6 to 10 traits or more. For variance component estimation, we applied an approach similar to previous methods that utilize eigen decomposition of the genetic relationship matrix (or kinship matrix) and linear transformation to achieve uncorrelated observations. However, we simplify the model into a weighted linear mixed model with a random intercept, allowing for more efficient estimation. Variance components are estimated using an expectation-maximization (EM) algorithm under a pure polygenic model.

For marker scanning, we developed a semi-exact method using an innovative linear transformation framework. Traditional approximate approaches use variance components estimated under the null model, offering faster computation but at the cost of reduced statistical power and lacking a natural extension to multivariate settings (Kang et al., 2010; Zhang et al., 2010). In contrast, exact methods such as GEMMA maintain statistical rigor but become computationally burdensome as the number of traits increases (Zhou & Stephens, 2014). Our

EMmvGWAS algorithm addresses these limitations by using a semi-exact method that fixes the
ratio of variance components while explicitly re-estimating the residual covariance matrix at
each locus. This enables a closed-form Wald test for each marker, eliminating the need for
iterative optimization during marker testing and substantially reduces time complexity, achieving
a practical balance between computational efficiency and statistical power.

To promote accessibility and reproducibility, we implemented the method in R, a widely used
platform in statistical genetics. Our package supports moderate-dimensional multivariate GWAS
with flexible input for phenotypes, covariates, and kinship matrices, enabling scalable
multivariate analyses and deeper insight into the genetic architecture of complex traits.

2. Methods

2.1. Multivariate linear mixed model

The multivariate linear mixed model for p traits measured on n individuals is defined as:

$$y = X\beta + Z_k\gamma_k + \xi + \varepsilon \qquad (1)$$

where the phenotype vector is stacked by individuals, $y = (y_1^\mathrm{T},\ldots,y_n^\mathrm{T})^\mathrm{T}$, and $y_j = (y_{j1},\ldots,y_{jp})^\mathrm{T}$ is

the $p\times 1$ vector of trait values for individual j, $j = 1,\ldots,n$. To test for associations, the model

considers one marker at a time. For the k th marker, the design matrix is $Z_k = z_k \otimes I_p$, where $\otimes$

denotes the Kronecker product, z_k is the n-dimensional vector of genotypes for that specific

marker, and $\gamma_k = (\gamma_{k1},\ldots,\gamma_{kp})^\mathrm{T}$ is the vector of its trait-specific effects. The fixed effects design

matrix $X = 1_n \otimes I_p$ contains trait-specific intercepts and the corresponding vector of fixed effects

is $\beta = (\beta_1,\ldots,\beta_p)^\mathrm{T}$. The random polygenic effects and residual errors are denoted by

$\xi = (\xi_1^\mathrm{T},\ldots,\xi_n^\mathrm{T})^\mathrm{T}$ and $\varepsilon = (\varepsilon_1^\mathrm{T},\ldots,\varepsilon_n^\mathrm{T})^\mathrm{T}$, respectively, where each ξ_j and ε_j is a p-dimensional

vector corresponding to individual j. We assume $\mathrm{var}(\xi) = K \otimes \Sigma_\xi$ and $\mathrm{var}(\varepsilon) = I_n \otimes \Sigma_\varepsilon$, where

Σ_ξ and Σ_ε are $p\times p$ symmetric positive-definite matrices describing trait-level polygenic and

residual covariances, respectively, and K is the known kinship matrix. Consequently, the total

variance of the observed phenotypes is expressed as:

$$\mathrm{var}(y) = V = K \otimes \Sigma_\xi + I_n \otimes \Sigma_\varepsilon \qquad (2)$$

This structured covariance allows the model to capture genetic background effects and

environmental noise across all p traits simultaneously. The inclusion of correlated individuals

substantially increases computational complexity, primarily due to the necessity of repeated

inversions and determinant calculations of the large-scale matrix V during variance component

estimation. Therefore, we propose a decorrelation method to render individuals uncorrelated to

simplify the inversion of covariance matrices.

2.2. Decorrelation of the polygenic effects

Let the eigen-decomposition of the kinship matrix be $K = UDU^T$, where $D = diag\{d_1,\ldots,d_n\}$ is the diagonal matrix holding the ordered eigenvalues and U is the matrix holding the eigenvectors corresponding to the eigenvalues. We define the transformation matrix as $H = D^{-\frac{1}{2}}U^T$. Since y is a stacked vector of phenotypic values across n individuals and p traits, the transformation is applied to all variables in the model using the Kronecker product $H \otimes I_p$. The transformed variables are defined as $y = (H \otimes I_p)y$, $X = (H \otimes I_p)X$, $Z_k = (H \otimes I_p)Z_k$, $\xi = (H \otimes I_p)\xi$, and $\varepsilon = (H \otimes I_p)\varepsilon$. Note that we have used a pseudo code notation to simplify the derivation. Hereafter, all variables in the linear mixed model are eigen-transformed. The phenotypic variance-covariance matrix of the transformed model is given by:

$$\text{var}(y) = V = I_n \otimes \Sigma_\xi + D^{-1} \otimes \Sigma_\varepsilon \tag{3}$$

The matrix V becomes block diagonal with n blocks of dimension $p \times p$. Compared with the original variance structure, the transformed covariance matrix is substantially simplified, as observations from different individuals are uncorrelated. For the j th individual, let $y_j = (y_{j1},\ldots,y_{jp})^T$ denote the vector of transformed phenotypic values for the p traits. The corresponding polygenic effects and residual errors are denoted by ξ_j and ε_j, respectively, where ξ_j and ε_j are $p \times 1$ vectors for $j = 1,\ldots,n$. The transformed model for the j th individual is

$$y_j = X_j\beta + Z_{jk}\gamma_k + \xi_j + \varepsilon_j \tag{4}$$

where X_j and Z_{jk} denote the transformed fixed-effect and marker design matrices corresponding to transformed individual j, respectively. The variance of y_j is

$$\text{Var}(y_j) = V_j = \Sigma_\xi + d_j^{-1}\Sigma_\varepsilon = d_j^{-1}(d_j\Sigma_R + I)\Sigma_\varepsilon = H_j\Sigma_\varepsilon \tag{5}$$

where $H_j = d_j^{-1}(d_j\Sigma_R + I)$, and $\Sigma_R = \Sigma_\xi\Sigma_\varepsilon^{-1}$ is the matrix version of variance ratio.

2.3 The expectation and maximization (EM) algorithm for estimating the covariance matrices

The decorrelation of polygenic effects makes all individuals independent. As a result, the log-likelihood function only involves matrices of dimension $p \times p$ for p traits. We estimate the covariance matrices under the null model, which includes only fixed intercepts and polygenic effects and excludes marker effects. Let $\theta = \{\Sigma_\xi,\Sigma_\varepsilon\}$ be the parameter set, while the β vector is not a vector of parameters in REML (it is a vector of parameter for ML). The restricted log likelihood function is

$$l(\theta) = -\frac{1}{2}\sum_{j=1}^{n}\left(\log|d_j\Sigma_R+I|+\log|d_j^{-1}\Sigma_\varepsilon|\right) - \frac{1}{2}\log\left|\sum_{j=1}^{n}X_j^T(d_j\Sigma_\varepsilon^{-1})(d_j\Sigma_R+I)^{-1}X_j\right|$$

$$-\frac{1}{2}\sum_{j=1}^{n}(y_j-X_j\hat{\beta})^T(d_j\Sigma_\varepsilon^{-1})(d_j\Sigma_R+I)^{-1}(y_j-X_j\hat{\beta}) \tag{6}$$

where

$$\hat{\beta} = \left[\sum_{j=1}^{n}d_jX_j^T(\Sigma_R d_j+I)^{-1}X_j\right]^{-1}\left[\sum_{j=1}^{n}d_jX_j^T(\Sigma_R d_j+I)^{-1}y_j\right] \tag{7}$$

Here, $\hat{\beta}$ is not a parameter vector but a function of θ. The covariance parameters are obtained by maximizing $l(\theta)$. The MIXED procedure PROC MIXED in SAS is readily available to perform the variance component estimations. However, PROC MIXED becomes computationally inefficient and may fail to converge properly due to the large number of parameters. To address these issues, we adopt an Expectation-Maximization (EM) algorithm. The EM algorithm guarantees a monotonic increase in the likelihood and achieves stable convergence.

In the EM algorithm framework, the missing values in the complete-data log likelihood function are the polygenic effects, ξ_j for $j=1,\cdots,n$, where each ξ_j is an $p\times1$ vector. We first update β using

$$\beta^{(t+1)} = \left[\sum_{j=1}^{n}d_jX_j^T(d_j\Sigma_\xi^{(t)}+I)^{-1}X_j\right]^{-1}\left[\sum_{j=1}^{n}d_jX_j^T(d_j\Sigma_\xi^{(t)}+I)^{-1}y_j\right] \tag{8}$$

Given ξ_j for $j=1,\cdots,n$, the polygenic variance and covariance matrix is define as

$$\Sigma_\xi = \frac{1}{n}\sum_{j=1}^{n}\xi_j\xi_j^T \tag{9}$$

When ξ_j is missing, we simply replace them by the conditional expectations to update this matrix,

$$\Sigma_\xi^{(t+1)} = \frac{1}{n}\sum_{j=1}^{n}E(\xi_j\xi_j^T\,|\,y_j) = \frac{1}{n}\sum_{j=1}^{n}\left[E(\xi_j\,|\,y_j)E(\xi_j^T\,|\,y_j)+\mathrm{var}(\xi_{j.}\,|\,y_j)\right] \tag{10}$$

where

$$E(\xi_j\,|\,y_j) = \hat{\xi}_j = d_j\Sigma_\xi^{(t)}(d_j\Sigma_\xi^{(t)}+I)^{-1}(y_j-X_j\beta^{(t)}) \tag{11}$$

and

$$\mathrm{var}(\xi_j\,|\,y_j) = \Sigma_\xi^{(t)}-d_j\Sigma_\xi^{(t)}(d_j\Sigma_\xi^{(t)}+I)^{-1}\Sigma_\xi^{(t)} \tag{12}$$

Finally, the updated residual covariance matrix is

$$\Sigma_\varepsilon^{(t+1)} = \frac{1}{n-1}\sum_{j=1}^{n}d_j(y_j-X_j\beta^{(t)}-\hat{\xi}_j)y_j^T \tag{13}$$

When the iterations converge, we report the two covariance matrices as the estimated parameters
denoted by $\hat{\theta} = \{\hat{\Sigma}_\xi, \hat{\Sigma}_\varepsilon\}$ and the estimated covariance matrix ratio denoted by $\hat{\Sigma}_R = \hat{\Sigma}_\xi \hat{\Sigma}_\varepsilon^{-1}$. This
matrix ratio will be treated as a constant and used in genome-wide scanning for associations of
genome-wide markers with multiple traits.

2.4 Scanning the entire genome for associations

Genome-wide association analysis is carried out by testing markers one at a time. In principle,
marker effects can be incorporated into the model by re-estimating all parameters for each
marker, yielding an exact but computationally expensive approach. In this study, we develop an
efficient strategy that fixes the variance matrix ratio estimated from the null model and scans the
genome marker by marker. This strategy allows closed-form estimation of marker effects and
enables re-estimation of the residual covariance matrix for each marker, thereby substantially
improving computational efficiency while maintaining good statistical performance. With the

variance matrix ratio fixed from the null model, genome-wide association testing is carried out
marker by marker. For marker k, the fixed effects include both the intercepts β and the marker
effects γ_k, which are estimated jointly by

194
$$
\begin{bmatrix} \hat{\beta} \\ \hat{\gamma}_k \end{bmatrix} = \begin{bmatrix} \sum X_j^T H_j^{-1} X_j & \sum X_j^T H_j^{-1} Z_{jk} \\ \sum Z_{jk}^T H_j^{-1} X_j & \sum Z_{jk}^T H_j^{-1} Z_{jk} \end{bmatrix}^{-1} \begin{bmatrix} \sum X_j^T H_j^{-1} y_j \\ \sum Z_{jk}^T H_j^{-1} y_j \end{bmatrix} = \begin{bmatrix} A_{XX} & A_{XZ} \\ A_{ZX} & A_{ZZ} \end{bmatrix}^{-1} \begin{bmatrix} A_{Xy} \\ A_{Zy} \end{bmatrix}
\tag{14}
$$

195　where $A_{XX} = \sum X_j^T H_j^{-1} X_j$, $A_{Xy} = \sum X_j^T H_j^{-1} y_j$, and similarly for the remaining block terms.

197　The intercepts β are nuisance parameters and can be eliminated prior to estimating γ_k via the
198　annihilator matrix (Xu, 2024) so that γ_k are estimated explicitly. Let us also define
199　$A_{yy} = \sum H_j^{-1} y_j y_j^T$. Define the following quantities

200
$$
\begin{aligned}
C_{ZZ} &= A_{ZZ} - A_{ZX} A_{XX}^{-1} A_{XZ} \\
C_{Zy} &= A_{Zy} - A_{ZX} A_{XX}^{-1} A_{Xy} \\
C_{yy} &= A_{yy} - A_{yX} A_{XX}^{-1} A_{Xy}
\end{aligned}
$$

201　The estimated γ_k is

202
$$
\hat{\gamma}_k = C_{ZZ}^{-1} C_{Zy}
\tag{15}
$$

203　with variance

204
$$
\mathrm{var}(\hat{\gamma}_k) = C_{ZZ}^{-1} \hat{\Sigma}_\varepsilon
\tag{16}
$$

205　where

206
$$
\hat{\Sigma}_\varepsilon = C_{yy} - C_{yZ} C_{ZZ}^{-1} C_{Zy}
\tag{17}
$$

207 is the estimated residual covariance matrix and $C_{yZ} = C_{Zy}^T$. The Wald test statistic for testing

208 $H_0 : \gamma_k = 0$ is

209 $$W_k = \hat{\gamma}_k^T \left[\text{var}(\hat{\gamma}_k) \right]^{-1} \hat{\gamma}_k = \hat{\gamma}_k^T \hat{\Sigma}_\varepsilon^{-1} C_{ZZ} \hat{\gamma}_k \qquad (18)$$

210 which follows approximately a Chi-square distribution with p degrees of freedom.

2.5. Dataset

We evaluated the performance and computational efficiency of the new method using three
genetic association datasets from rice, human, and mouse populations. Each dataset includes
genome-wide genotype data and multiple phenotypic traits, enabling a range of multivariate
GWAS applications. The rice dataset consists of 1,495 hybrid rice individuals (Huang et al.,
2015). For the human dataset, we analyzed a subset of 5,000 unrelated white British individuals
from the UK Biobank (Sudlow et al., 2015) cohort for demonstration purposes. The mouse
dataset was obtained from the CFW outbred population study (Parker et al., 2016), available
from the Dryad Digital Repository (doi:10.5061/dryad.2rs41).

For the rice and mouse datasets, we applied standard quality control procedures, including
removal of SNPs with minor allele frequency below 0.01 and individuals or variants with
missing data. For the UK Biobank data, genotype data were filtered using PLINK2 with
thresholds of 0.05 for missing genotype rate and minor allele frequency, along with linkage
disequilibrium pruning (Chang et al., 2015). Detailed information is provided in Supplementary
Table S1. All phenotypes were standardized prior to analysis.

2.6. Simulation analysis

To evaluate the type I error control and statistical power of our method, we performed null
simulations and alternative simulations using real genotype data from the rice and mouse
datasets. We excluded the human dataset from simulations due to its large sample size and dense
marker coverage, which made simulations computationally impractical.

For null simulations, phenotypes were generated under a pure polygenic model without any
marker effects. Specifically, we simulated 100 sets of phenotypes based on the real genetic
relatedness matrix and the estimated genetic and environmental covariance matrices from the
rice and mouse data. We randomly selected 10,000 SNPs from the real genotype data. We then
computed p-values for all combinations of these SNPs and the simulated phenotypes, resulting in
one million tests per dataset. For alternative simulations, we added SNP effects into the
phenotypes to assess empirical power. We first selected 1,000 evenly spaced SNPs from a set of
SNPs that showed no significant association with either trait (P > 0.1 in all univariate and
multivariate tests). Each selected SNP was assigned an effect on the first trait to explain a fixed
proportion of phenotypic variance (PVE). The effect on the second trait was scaled to explain
either 20% or 40% of PVE of the second trait, with effects either in the same or opposite
direction across traits. These synthetic effects were added to the original phenotypes to generate

datasets with known causal SNPs. Power was calculated as the proportion of causal SNPs
detected at the Bonferroni-corrected significance level with initial significance level of 0.05. We
then applied both EMmvGWAS and GEMMA to these simulated datasets to compare their
empirical power under various effect sizes and trait correlation scenarios.

3. Results

3.1. Computational results

To evaluate the computational performance of our method, we presented its theoretical time
complexity in comparison with existing approaches. EMmvGWAS benefits from an explicit
solution, resulting in a reduced overall computational complexity of
$O(n^3 + n^2 p + t_1 np^2 + sn^2 + snp^2)$, where n is the sample size, p is the number of traits, t_1 is the
number of iterations of the EM algorithm, and s is the number of markers. The terms n^3 and
$n^2 d$ arise from eigen decomposition and transformation of the phenotypes. The term $t_1 np^2$
corresponds to the variance component estimation under the null model using the EM algorithm,
while $s(n^2 + np^2)$ reflects the cost of genome-wide marker scanning across multiple traits. In
practice, the number of markers s is typically much larger than both n and p, making the last
term $s(n^2 + np^2)$ the dominant contributor to the overall complexity. The number of fixed effects
per test is omitted in these expressions because its contribution is negligible and can be adjusted
in the phenotypes. In contrast, the exact algorithms such as GEMMA require
$O(n^3 + n^2 p + sn^2 + st_1 np^2 + st_2 np^2)$, where t_1 and t_2 represent the number of optimization
iterations needed for the EM algorithm and the Newton-Raphson algorithm, respectively. As the
number of traits increases, the additional iterative steps in genome marker scanning in GEMMA
lead to substantially higher computational burden.

We evaluated the computational efficiency of EMmvGWAS using real-world datasets from rice,
mouse, and human, comparing it against GEMMA, the gold standard exact algorithm. For each
dataset, we varied the number of traits and recorded the total runtime. As shown in Figure 1, the
runtime of GEMMA increases rapidly with the number of traits. For example, GEMMA required
more than three weeks (554 hours) for 10 traits in rice and nearly 200 hours for 10 traits in the
human dataset. In contrast, our EMmvGWAS maintains a nearly flat runtime profile, requiring
less than 5 hours for the rice and the mouse data. For the human dataset, we applied 10-core
parallel computing to EMmvGWAS, which enabled analysis of up to 10 traits within 40 hours,
compared to nearly 200 hours for GEMMA.

3.2. Consistency of association results

281 We showed that the semi-exact method EMmvGWAS is highly consistent with the exact method
282 implemented in GEMMA when analyzing 10 traits across the rice, the mouse, and the human
283 datasets (Figure 2). The Pearson correlation coefficients were nearly 1 in all cases, indicating that
284 the two approaches produced almost identical test statistics. Figure 3 displays the Manhattan

plots from both methods, which show highly similar peak patterns throughout the genome, further confirming the alignment of their results.

3.3. Simulation study

To illustrate that our method produces P values with proper control of type 1 error and power, we conducted both null and alternative simulations based on real genotype data from the rice and mouse datasets. Under the null model, simulated phenotypes were generated without any marker effects. Quantile–quantile (QQ) plots from the null simulations in both the mouse and the rice data are shown in Figure 4. For both EMmvGWAS and GEMMA, the observed p-values closely followed the expected distribution under the null hypothesis, with only minor deviations in the extreme tail. This indicates that both approaches properly control the type 1 error across species. To evaluate statistical power, we conducted alternative simulations by injecting known genetic effects into selected SNPs with predefined proportions of variance explained. In Figure 5, GEMMA exhibited slightly higher, EMmvGWAS maintained competitive performance with minimal power loss. Importantly, EMmvGWAS achieved this with substantially reduced computational burden, demonstrating its practical advantage for multivariate GWAS applications.

4. Discussion

These results highlight the substantial computational gains offered by EMmvGWAS. While GEMMA scales poorly with the dimensionality of the trait space, the EMmvGWAS framework maintains a nearly constant runtime. This efficiency arises from the closed-form re-estimation of the residual variance, which avoids repeated iterative optimization. The performance advantage is consistent across organisms with diverse sample sizes and genetic architectures, demonstrating the scalability and robustness of EMmvGWAS. The human analysis further demonstrates that the semi-exact framework can efficiently leverage parallel computing, achieving additional reductions in runtime without compromise in accuracy. The efficiency of EMmvGWAS opens the door to comprehensive multivariate GWAS across diverse datasets and trait dimensions.

While exact methods are often expected to produce more significant results, this was not observed in our comparisons. We suspect that GEMMA's internal calibration procedure reduces the apparent significance of its test statistics, leading to closer agreement with the semi-exact method. QQ plots from null simulations confirmed that EMmvGWAS yields well-calibrated p-values that are comparable to GEMMA, with no evidence of systematic inflation or deflation of the type 1 error. This indicates that EMmvGWAS does not compromise type I error control while offering substantial computational efficiency.

For variance component estimation, we employed the expectation-maximization (EM) algorithm under the pure polygenic (null) model. Despite its simplicity, the EM algorithm provided sufficiently accurate estimates and was computationally efficient in our analyses. In contrast, GEMMA uses a more complex procedure that combines parameter-expanded EM algorithm with

Newton–Raphson optimization to accelerate convergence. However, our results demonstrate that
the standard EM approach is already fast in practice and does not compromise the accuracy of
downstream test statistics. This suggests that, for moderate numbers of traits, the EM-based
variance component estimation strikes a favorable balance between simplicity, speed, and
reliability.

Our method is implemented in the R programming language to ensure accessibility and ease of
use within the statistical genetics community. While R is not optimized for high-performance
computing, our implementation still demonstrates competitive efficiency. We encourage future
implementation in compiled languages such as C++, which could offer further improvement in
speed. Additionally, genome-wide marker scanning is inherently parallelizable, and our package
includes built-in support for parallel computing to take advantage of multi-core computing
systems. The framework could also be extended to support GPU acceleration, enabling even
greater scalability for large-scale multivariate GWAS.

Declarations

Ethics approval and consent to participate

This study did not involve the collection of any new human or animal data. All analyses were
conducted using previously published datasets with existing ethical approval, including UK
Biobank data accessed under an approved application. No additional ethics approval or consent
was required for this work.

Consent for publication

Not applicable

Availability of data and materials

The data set of 1,495 hybrid rice varieties is available at the Rice Haplotype Map Project
database (http://www.ncgr.ac.cn/RiceHap4). The CFW mouse datasets can be available at Dryad
Digital Repository (doi:10.5061/dryad.2rs41). UK Biobank (UKB) data can be accessed through
the UKB Research Analysis Platform (RAP) following approval via the UKB access system
(https://www.ukbiobank.ac.uk).

Competing interests

The authors declare that they have no competing interests

Funding

Authors' contributions

SX conceived the proposed method. CT analyzed and interpreted the data and generated the figures in the manuscript. CL and QW provided processed UKB dataset. YC provided the hybrid rice dataset. XW advised on the GEMMA implementation. CT wrote the manuscript. SX supervised the research and edited the manuscript. All authors reviewed the manuscript.

Reference

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*, *4*(1), s13742-13015-10047-13748.

Hackinger, S., & Zeggini, E. (2017). Statistical methods to detect pleiotropy in human complex traits. *Open biology*, *7*(11), 170125.

Huang, X., Yang, S., Gong, J., Zhao, Y., Feng, Q., Gong, H., Li, W., Zhan, Q., Cheng, B., & Xia, J. (2015). Genomic analysis of hybrid rice varieties reveals numerous superior alleles that contribute to heterosis. *Nature communications*, *6*(1), 6258.

Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S.-y., Freimer, N. B., Sabatti, C., & Eskin, E. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nature genetics*, *42*(4), 348-354.

Korte, A., Vilhjálmsson, B. J., Segura, V., Platt, A., Long, Q., & Nordborg, M. (2012). A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature genetics*, *44*(9), 1066-1071.

Parker, C. C., Gopalakrishnan, S., Carbonetto, P., Gonzales, N. M., Leung, E., Park, Y. J., Aryee, E., Davis, J., Blizard, D. A., & Ackert-Bicknell, C. L. (2016). Genome-wide association study of behavioral, physiological and gene expression traits in outbred CFW mice. *Nature genetics*, *48*(8), 919-926.

Porter, H. F., & O'Reilly, P. F. (2017). Multivariate simulation framework reveals performance of multi-trait GWAS methods. *Scientific reports*, *7*(1), 38837.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., & Landray, M. (2015). UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine*, *12*(3), e1001779.

Turley, P., Walters, R. K., Maghzian, O., Okbay, A., Lee, J. J., Fontana, M. A., Nguyen-Viet, T. A., Wedow, R., Zacher, M., & Furlotte, N. A. (2018). Multi-trait analysis of genome-wide association summary statistics using MTAG. *Nature genetics*, *50*(2), 229-237.

Visscher, P. M., Wray, N. R., Zhang, Q., Sklar, P., McCarthy, M. I., Brown, M. A., & Yang, J. (2017). 10 years of GWAS discovery: biology, function, and translation. *The American Journal of Human Genetics*, *101*(1), 5-22.

Xu, S. (2024). Estimating genetic variance contributed by a quantitative trait locus: removing nuisance parameters. *Genetics*, *227*(4), iyae095.

Zhang, Z., Ersoz, E., Lai, C.-Q., Todhunter, R. J., Tiwari, H. K., Gore, M. A., Bradbury, P. J., Yu, J., Arnett, D. K., & Ordovas, J. M. (2010). Mixed linear model approach adapted for genome-wide association studies. *Nature genetics*, *42*(4), 355-360.

Zhou, X., & Stephens, M. (2014). Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature methods*, *11*(4), 407-409.

Figures

Figure 1. Computational runtime of EMmvGWAS and GEMMA across datasets and trait dimensions.

Runtime (in hours) is plotted across the number of traits for the rice, the mouse, and the human datasets. The red
solid line indicates GEMMA and the blue dashed line indicate EMmvGWAS. The rice and mouse analyses were
carried out on 1 core, while the human data analysis was run with 10 cores.

Figure 2. Comparison of −log10(p) values between the EMmvGWAS and GEMMA for 10 traits.

Scatterplots of EMmvGWAS versus GEMMA results are shown for the rice, the mouse, and the human datasets. The
red line represents the fitted regression line, and *r* denotes the Pearson correlation coefficient between methods.

447

Figure 3. Manhattan plots for EMmvGWAS and GEMMA methods across rice, mouse, and human datasets using 10 traits.

Each row shows results for one species (rice, mouse, and human), and each column corresponds to a method (EMmvGWAS on the left and GEMMA on the right). Genome-wide association −log10(p) values are plotted along the genome. The red horizontal line marks the Bonferroni-corrected significance threshold for each species.

453

Figure 4. QQ plots for null simulations in the mouse and the rice data.

Observed versus expected $-\log_{10}(p)$ values are shown for EMmvGWAS (left panels) and GEMMA (right panels). Results for the mouse and rice datasets are displayed in the top and bottom panels, respectively. The red diagonal line represents the null expectation.

Figure 5. Power comparison between EMmvGWAS and GEMMA under alternative simulations using rice data.

Causal SNPs were simulated with varying levels of phenotypic variance explained for the first trait (PVE), and effect sizes on the second trait were set as either 20% or 40% of PVE, with effects in the same (sign +) or opposite (sign −) directions. Power was defined as the proportion of causal SNPs detected at the Bonferroni-corrected significance level.