An Expectation and Maximization Algorithm for Multivariate Genome-wide Association Studies (EMmvGWAS)

Chin-Sheng Teng[1], Xuesong Wang[2], Cheng Liu[3], Qishan Wang[3], Yanru Cui[4] and Shizhong Xu[2*]

[1]Department of Statistics, University of California, Riverside, CA 92521, USA
[2]Department of Botany and Plant Sciences, University of California, Riverside, CA 92521, USA
[3] Department of Animal Science, College of Animal Sciences, Zhejiang University, 866# Yuhangtang Road, Hangzhou 310058, China
[4]College of Agronomy, Hebei Agricultural University, Baoding 071001, China

*shizhong.xu@ucr.edu

Abstract

Genome-wide association studies (GWAS) often analyze one trait at a time, but multivariate GWAS can increase the power by leveraging trait correlations. However, existing methods struggle with high computational demands, especially when analyzing over five traits. We present EMmvGWAS, an efficient multivariate GWAS framework, which dramatically reduces computational time when analyzing a moderate number of traits. EMmvGWAS estimates covariance matrices via an Expectation-Maximization algorithm and accelerates genome-wide scans using a semi-exact method. We validate its accuracy and scalability through simulations and real data from rice, mouse, and human populations. The package is available at https://github.com/Jason-Teng/EMmvGWAS.

Keywords: Expectation and maximization algorithm, Genome-wide association studies, Multivariate linear mixed model, Restricted maximum likelihood method.

1. Background

Genome-wide association studies (GWAS) have significantly advanced our understanding of the genetic architecture underlying complex traits by identifying genetic variants associated with quantitative traits (Visscher et al., 2017). Traditionally, GWAS are often conducted on individual traits independently, even though multiple correlated phenotypes are always measured in genetic studies. Analyzing traits individually overlooks valuable trait correlations, thus potentially reducing the statistical power and increasing the type 1 error rate. Multivariate GWAS leverages these correlations by using multivariate linear mixed models, improving power and reducing false positive associations in identifying pleiotropic genetic variants, which affect multiple traits simultaneously (Korte et al., 2012; Porter & O'Reilly, 2017; Turley et al., 2018; Zhou & Stephens, 2014). Such methods are crucial for uncovering shared genetic bases among related traits, providing deeper insights into biological pathways and mechanisms underlying complex diseases and phenotypes (Hackinger & Zeggini, 2017).

Despite their clear advantages, multivariate GWAS methods confront significant computational hurdles, primarily due to the complexity of modeling high-dimensional covariance structures and the computational cost associated with scanning genome-wide markers. Estimating genetic and environmental covariance matrices between traits becomes increasingly demanding as the number of traits grows, often requiring iterative optimization over large parameter spaces. At the same time, genome-wide marker scanning involves evaluating potentially millions of SNPs across all traits, which remains computationally intensive. These two challenges compound in high-dimensional settings, making many existing multivariate GWAS tools impractical for real-world applications involving many traits. Although R is widely used in statistical genetics, there is a lack of efficient R packages for multivariate GWAS, especially for analyses involving many traits.

To address these challenges, we developed EMmvGWAS, a general statistical framework that supports multivariate GWAS with improved computational efficiency. While suitable for small trait sets, EMmvGWAS offers substantial speed improvements as the number of traits grows, making it especially effective for studies involving 6 to 10 traits or more. For variance component estimation, we applied an approach similar to previous methods that utilize eigen decomposition of the genetic relationship matrix (or kinship matrix) and linear transformation to achieve uncorrelated observations. However, we simplify the model into a weighted linear mixed model with a random intercept, allowing for more efficient estimation. Variance components are estimated using an expectation-maximization (EM) algorithm under a pure polygenic model.

For marker scanning, we developed a semi-exact method using an innovative linear
transformation framework. Traditional approximate approaches use variance components
estimated under the null model, offering faster computation but at the cost of reduced statistical
power and lacking a natural extension to multivariate settings (Kang et al., 2010; Zhang et al.,
2010). In contrast, exact methods such as GEMMA maintain statistical rigor but become
computationally burdensome as the number of traits increases (Zhou & Stephens, 2014). Our
EMmvGWAS algorithm addresses these limitations by using a semi-exact method that fixes the
ratio of variance components while explicitly re-estimating the residual covariance matrix at
each locus. This enables a closed-form Wald test for each marker, eliminating the need for
iterative optimization during marker testing and substantially reduces time complexity, achieving
a practical balance between computational efficiency and statistical power.

To promote accessibility and reproducibility, we implemented the method in R, a widely used
platform in statistical genetics. Our package supports moderate-dimensional multivariate GWAS
with flexible input for phenotypes, covariates, and kinship matrices, enabling scalable
multivariate analyses and deeper insight into the genetic architecture of complex traits.

88   2. Results

89   2.1. Computational results

To evaluate the computational performance of our method, we presented its theoretical time
complexity in comparison with existing approaches. EMmvGWAS benefits from an explicit
solution, resulting in a reduced overall computational complexity of
$O(n^3 + n^2d + t_1nd^2 + sn^2 + snd^2)$, where n is the sample size, d is the number of traits, t_1 is the
number of iterations of the EM algorithm, and s is the number of markers. The terms n^3 and
n^2d arise from eigen decomposition and transformation of the phenotypes. The term t_1nd^2
corresponds to the variance component estimation under the null model using the EM algorithm,
while $s(n^2 + nd^2)$ reflects the cost of genome-wide marker scanning across multiple traits. In
practice, the number of markers s is typically much larger than both n and d, making the last term
$s(n^2 + nd^2)$ the dominant contributor to the overall complexity. The number of fixed effects per
test is omitted in these expressions because its contribution is negligible and can be adjusted in
the phenotypes. In contrast, the exact algorithms such as GEMMA require
$O(n^3 + n^2d + sn^2 + st_1nd^2 + st_2nd^2)$, where t_1 and t_2 represent the number of optimization
iterations needed for the EM algorithm and the Newton-Raphson algorithm, respectively. As the
number of traits increases, the additional iterative steps in genome marker scanning in GEMMA
lead to substantially higher computational burden.

We evaluated the computational efficiency of EMmvGWAS using real-world datasets from rice,
mouse, and human, comparing it against GEMMA, the gold standard exact algorithm. For each
dataset, we varied the number of traits and recorded the total runtime. As shown in Figure 1, the
runtime of GEMMA increases rapidly with the number of traits. For example, GEMMA required
more than three weeks (554 hours) for 10 traits in rice and nearly 200 hours for 10 traits in the
human dataset. In contrast, our EMmvGWAS maintains a nearly flat runtime profile, requiring
less than 5 hours for the rice and the mouse data. For the human dataset, we applied 10-core
parallel computing to EMmvGWAS, which enabled analysis of up to 10 traits within 40 hours,
compared to nearly 200 hours for GEMMA.

Figure 1. Computational runtime of EMmvGWAS and GEMMA across datasets and trait dimensions.

118 Runtime (in hours) is plotted across the number of traits for the rice, the mouse, and the human datasets. The red
119 solid line indicates GEMMA and the blue dashed line indicate EMmvGWAS. The rice and mouse analyses were
120 carried out on 1 core, while the human data analysis was run with 10 cores.

2.2. Consistency of association results

We showed that the semi-exact method EMmvGWAS is highly consistent with the exact method
implemented in GEMMA when analyzing 10 traits across the rice, the mouse, and the human
datasets (Figure 2). The Pearson correlation coefficients were nearly 1 in all cases, indicating that
the two approaches produced almost identical test statistics. Figure 3 displays the Manhattan
plots from both methods, which show highly similar peak patterns throughout the genome,
further confirming the alignment of their results.

Figure 2. Comparison of −log10(p) values between the EMmvGWAS and GEMMA for 10 traits.

132 Scatterplots of EMmvGWAS versus GEMMA results are shown for the rice, the mouse, and the human datasets. The
133 red line represents the fitted regression line, and r denotes the Pearson correlation coefficient between methods.

Figure 3. Manhattan plots for EMmvGWAS and GEMMA methods across rice, mouse, and human datasets using 10 traits.

Each row shows results for one species (rice, mouse, and human), and each column corresponds to a method (EMmvGWAS on the left and GEMMA on the right). Genome-wide association −log10(p) values are plotted along the genome. The red horizontal line marks the Bonferroni-corrected significance threshold for each species.

2.3. Simulation study

To illustrate that our method produces P values with proper control of type 1 error and power, we conducted both null and alternative simulations based on real genotype data from the rice and mouse datasets. Under the null model, simulated phenotypes were generated without any marker effects. Quantile–quantile (QQ) plots from the null simulations in both the mouse and the rice data are shown in Figure 4. For both EMmvGWAS and GEMMA, the observed p-values closely

followed the expected distribution under the null hypothesis, with only minor deviations in the
extreme tail. This indicates that both approaches properly control the type 1 error across species.
To evaluate statistical power, we conducted alternative simulations by injecting known genetic
effects into selected SNPs with predefined proportions of variance explained. In Figure 5,
GEMMA exhibited slightly higher, EMmvGWAS maintained competitive performance with
minimal power loss. Importantly, EMmvGWAS achieved this with substantially reduced
computational burden, demonstrating its practical advantage for multivariate GWAS
applications.

Figure 4. QQ plots for null simulations in the mouse and the rice data.

158 Observed versus expected −log10(p) values are shown for EMmvGWAS (left panels) and GEMMA (right panels).
159 Results for the mouse and rice datasets are displayed in the top and bottom panels, respectively. The red diagonal
160 line represents the null expectation.

Figure 5. Power comparison between EMmvGWAS and GEMMA under alternative simulations using rice data.

Causal SNPs were simulated with varying levels of phenotypic variance explained for the first trait (PVE), and effect sizes on the second trait were set as either 20% or 40% of PVE, with effects in the same (sign +) or opposite (sign −) directions. Power was defined as the proportion of causal SNPs detected at the Bonferroni-corrected significance level.

3. Discussion

These results highlight the substantial computational gains offered by EMmvGWAS. While GEMMA scales poorly with the dimensionality of the trait space, the EMmvGWAS framework maintains a nearly constant runtime. This efficiency arises from the closed-form re-estimation of the residual variance, which avoids repeated iterative optimization. The performance advantage is consistent across organisms with diverse sample sizes and genetic architectures, demonstrating the scalability and robustness of EMmvGWAS. The human analysis further demonstrates that the semi-exact framework can efficiently leverage parallel computing, achieving additional reductions in runtime without compromise in accuracy. The efficiency of EMmvGWAS opens the door to comprehensive multivariate GWAS across diverse datasets and trait dimensions.

181 While exact methods are often expected to produce more significant results, this was not
182 observed in our comparisons. We suspect that GEMMA's internal calibration procedure reduces
183 the apparent significance of its test statistics, leading to closer agreement with the semi-exact
184 method. QQ plots from null simulations confirmed that EMmvGWAS yields well-calibrated p-
185 values that are comparable to GEMMA, with no evidence of systematic inflation or deflation of
186 the type 1 error. This indicates that EMmvGWAS does not compromise type I error control while
187 offering substantial computational efficiency.

189 For variance component estimation, we employed the expectation-maximization (EM) algorithm
190 under the pure polygenic (null) model. Despite its simplicity, the EM algorithm provided
191 sufficiently accurate estimates and was computationally efficient in our analyses. In contrast,
192 GEMMA uses a more complex procedure that combines parameter-expanded EM algorithm with
193 Newton–Raphson optimization to accelerate convergence. However, our results demonstrate that
194 the standard EM approach is already fast in practice and does not compromise the accuracy of
195 downstream test statistics. This suggests that, for moderate numbers of traits, the EM-based
196 variance component estimation strikes a favorable balance between simplicity, speed, and
197 reliability.

199 Our method is implemented in the R programming language to ensure accessibility and ease of
200 use within the statistical genetics community. While R is not optimized for high-performance
201 computing, our implementation still demonstrates competitive efficiency. We encourage future
202 implementation in compiled languages such as C++, which could offer further improvement in
203 speed. Additionally, genome-wide marker scanning is inherently parallelizable, and our package
204 includes built-in support for parallel computing to take advantage of multi-core computing
205 systems. The framework could also be extended to support GPU acceleration, enabling even
206 greater scalability for large-scale multivariate GWAS.

208 **4. Methods**

209 **4.1. Multivariate linear mixed model**

210 Let $y_{\cdot t}$ be an $n \times 1$ vector of the phenotypic values of the tth trait collected from n individuals
211 for $t = 1, \cdots, p$ and p is the number of traits. To intuitively introduce the linear mixed model, we
212 assume $p = 2$ and define the linear mixed model by

213
$$
\begin{bmatrix} y_{\cdot 1} \\ y_{\cdot 2} \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} \xi_{\cdot 1} \\ \xi_{\cdot 2} \end{bmatrix} + \begin{bmatrix} \varepsilon_{\cdot 1} \\ \varepsilon_{\cdot 2} \end{bmatrix} \tag{1}
$$

where $y_{\bullet 1}$ is an $n \times 1$ vector of phenotypic values for the first trait and $y_{\bullet 2}$ is an $n \times 1$ vector of

phenotypic values for the second trait, X_1 and X_2 are unity vectors of length n to capture the

intercepts of the two traits, β_1 and β_2 are the corresponding intercepts, $\xi_{\bullet 1}$ and $\xi_{\bullet 2}$ are $n \times 1$

vectors of polygenic effects of the two traits, and $\varepsilon_{\bullet 1}$ and $\varepsilon_{\bullet 2}$ are the vectors of environmental

effects (residual errors) of the two traits. The polygenic effects are random following assumed

normal distributions, $\xi_{\bullet t} \sim N(0, K\sigma_t^2)$ for $t = 1, 2$, where K is an $n \times n$ kinship matrix inferred

from genome-wide markers. Similarly, the residual errors are also assumed to be normally

distributed $\varepsilon_{\bullet t} \sim N(0, I\phi_t^2)$ for $t = 1, 2$. In addition to the variances, the two traits are genetically

and environmentally correlated with $\mathrm{cov}(\xi_{\bullet 1}, \xi_{\bullet 2}) = K\sigma_{12}$ and $\mathrm{cov}(\varepsilon_{\bullet 1}, \varepsilon_{\bullet 2}) = I\phi_{12}$. Define the

polygenic covariance matrix and the residual covariance matrix by

224
$$\Sigma_{\xi} = \begin{bmatrix} \sigma_1^2 & \sigma_{12} \\ \sigma_{12} & \sigma_2^2 \end{bmatrix} \tag{2}$$

225   and

226
$$\Sigma_{\varepsilon} = \begin{bmatrix} \phi_1^2 & \phi_{12} \\ \phi_{12} & \phi_2^2 \end{bmatrix} \tag{3}$$

227   respectively. The expectation of the joint model for the two traits is

228
$$\mathrm{E}\begin{bmatrix} y_{\bullet 1} \\ y_{\bullet 2} \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} \tag{4}$$

229   The covariance matrix of the model is

230
$$\mathrm{var}\begin{bmatrix} y_{\bullet 1} \\ y_{\bullet 2} \end{bmatrix} = \begin{bmatrix} K\sigma_1^2 & K\sigma_{12} \\ K\sigma_{12} & K\sigma_2^2 \end{bmatrix} + \begin{bmatrix} I\phi_1^2 & I\phi_{12} \\ I\phi_{12} & I\phi_1^2 \end{bmatrix} \tag{5}$$

231   Let

232
$$y = \begin{bmatrix} y_{\bullet 1} \\ y_{\bullet 2} \end{bmatrix}, X = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix}, \beta = \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}, \xi = \begin{bmatrix} \xi_{\bullet 1} \\ \xi_{\bullet 2} \end{bmatrix}, \varepsilon = \begin{bmatrix} \varepsilon_{\bullet 1} \\ \varepsilon_{\bullet 2} \end{bmatrix} \tag{6}$$

233   The joint linear mixed model written in an even more compact matrix notation is

234
$$y = X\beta + \xi + \varepsilon \tag{7}$$

235   The expectation of the model is $\mathrm{E}(y) = X\beta$ and the variance is

236
$$\mathrm{var}(y) = V = \Sigma_{\xi} \otimes K + \Sigma_{\varepsilon} \otimes I \tag{8}$$

237   where I is an $n \times n$ identity matrix and $\otimes$ represents Kronecker matrix product.

4.2. Decorrelation of the polygenic effects

Let us perform eigen decomposition for the kinship matrix, $K = UDU^T$. We then define

$H = D^{-\frac{1}{2}}U^T$ and use it to transform all variables in the model with $y_t = Hy_t$, $X_t = HX_t$,

$\xi_t = H\xi_t$, and $\varepsilon_t = H\varepsilon_t$. Note that we have used a pseudo code notation to simplify the

derivation. Hereafter, all variables in the linear mixed model are eigen-transformed. The

expectation of the model becomes

$$E\begin{bmatrix} y_{\cdot 1} \\ y_{\cdot 2} \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix}\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} \tag{9}$$

and the variance matrix is

$$\mathrm{var}\begin{bmatrix} y_{\cdot 1} \\ y_{\cdot 2} \end{bmatrix} = \begin{bmatrix} \mathrm{var}(\xi_{\cdot 1}) & \mathrm{cov}(\xi_{\cdot 1},\xi_{\cdot 2}) \\ \mathrm{var}(\xi_{\cdot 2},\xi_{\cdot 1}) & \mathrm{var}(\xi_{\cdot 2}) \end{bmatrix} + \begin{bmatrix} \mathrm{var}(\varepsilon_{\cdot 1}) & \mathrm{cov}(\varepsilon_{\cdot 1},\varepsilon_{\cdot 2}) \\ \mathrm{var}(\varepsilon_{\cdot 2},\varepsilon_{\cdot 1}) & \mathrm{var}(\varepsilon_{\cdot 2}) \end{bmatrix} \tag{10}$$

where

$$\mathrm{var}(\xi_{\cdot t}) = HKH^T\sigma_t^2 = D^{-\frac{1}{2}}U^T UDU^T UD^{-\frac{1}{2}}\sigma_t^2 = I\sigma_t^2 \tag{11}$$

and $\mathrm{cov}(\xi_{\cdot 1},\xi_{\cdot 2}) = I\sigma_{12}$ is obtained the same way. The residual covariance matrix is

$$\mathrm{var}(\varepsilon_{\cdot t}) = HH^T\phi_t^2 = D^{-\frac{1}{2}}U^T UD^{-\frac{1}{2}}\phi_t^2 = D^{-1}\phi_t^2 \tag{12}$$

and $\mathrm{cov}(\varepsilon_{\cdot 1},\varepsilon_{\cdot 2}) = D^{-1}\phi_{12}$. Therefore,

$$\mathrm{var}\begin{bmatrix} y_{\cdot 1} \\ y_{\cdot 2} \end{bmatrix} = \begin{bmatrix} I\sigma_1^2 & I\sigma_{12} \\ I\sigma_{12} & I\sigma_2^2 \end{bmatrix} + \begin{bmatrix} D^{-1}\phi_1^2 & D^{-1}\phi_{12} \\ D^{-1}\phi_{12} & D^{-1}\phi_2^2 \end{bmatrix} \tag{13}$$

Let $y = \begin{bmatrix} y_{\cdot 1}^T & y_{\cdot 2}^T \end{bmatrix}^T$, we have

$$\mathrm{var}(y) = V = \Sigma_\xi \otimes I + \Sigma_\varepsilon \otimes D^{-1} \tag{14}$$

Comparing equation (14) with equation (8), we see that the variance matrix of the transformed
model is quite different from the variance matrix of the original model. The advantage of the
transformed model will be obvious after we sort y by individuals, not by traits.

Let y_{jt} be the transformed phenotypic value for trait $t = 1, 2$ ($p = 2$ traits) from individual

$j = 1, 2, 3$ ($n = 3$ individuals). The first subscript indicates the individual ID and the second

subscript indicates the trait ID. The sorted model is

$$
\begin{bmatrix} y_{11} \\ y_{12} \\ y_{21} \\ y_{22} \\ y_{31} \\ y_{32} \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_1 \\ X_2 & 0 \\ 0 & X_2 \\ X_3 & 0 \\ 0 & X_3 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} \xi_{11} \\ \xi_{12} \\ \xi_{21} \\ \xi_{22} \\ \xi_{31} \\ \xi_{32} \end{bmatrix} + \begin{bmatrix} \varepsilon_{11} \\ \varepsilon_{12} \\ \varepsilon_{21} \\ \varepsilon_{22} \\ \varepsilon_{31} \\ \xi_{32} \end{bmatrix}
\tag{15}
$$

where $X_1 = X_2 = X_3 = 1$ are constants capturing the intercepts for the two traits. In matrix notation, the expectation of the model is

$$
E(y) = X\beta
\tag{16}
$$

and the variance matrix is

$$
\text{var}(y) = V = I \otimes \Sigma_\xi + D^{-1} \otimes \Sigma_\varepsilon
\tag{17}
$$

which looks very similar to equation (14) but the two matrices around the Kronecker product sign $\otimes$ swapped positions. This V matrix is block diagonal and inverting this block diagonal matrix of dimension $(np) \times (np)$ can be formulated into inverting n matrices of dimension $p \times p$. Substantial computational time can be saved with this transformation. The covariance structure of the polygene is

$$
\text{var} \begin{bmatrix} \xi_{11} \\ \xi_{12} \\ \xi_{21} \\ \xi_{22} \\ \xi_{31} \\ \xi_{32} \end{bmatrix} = \begin{bmatrix} \sigma_1^2 & \sigma_{12} & 0 & 0 & 0 & 0 \\ \sigma_{12} & \sigma_2^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_1^2 & \sigma_{12} & 0 & 0 \\ 0 & 0 & \sigma_{12} & \sigma_2^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_1^2 & \sigma_{12} \\ 0 & 0 & 0 & 0 & \sigma_{12} & \sigma_2^2 \end{bmatrix}
\tag{18}
$$

The covariance structure of the residuals is

$$
\text{var} \begin{bmatrix} \varepsilon_{11} \\ \varepsilon_{12} \\ \varepsilon_{21} \\ \varepsilon_{22} \\ \varepsilon_{31} \\ \xi_{32} \end{bmatrix} = \begin{bmatrix} d_1^{-1}\phi_1^2 & d_1^{-1}\phi_{12} & 0 & 0 & 0 & 0 \\ d_1^{-1}\phi_{12} & d_1^{-1}\phi_2^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & d_2^{-1}\phi_1^2 & d_2^{-1}\phi_{12} & 0 & 0 \\ 0 & 0 & d_2^{-1}\phi_{12} & d_2^{-1}\phi_2^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & d_3^{-1}\phi_1^2 & d_3^{-1}\phi_{12} \\ 0 & 0 & 0 & 0 & d_3^{-1}\phi_{12} & d_3^{-1}\phi_2^2 \end{bmatrix}
\tag{19}
$$

The two traits from the jth individual $y_{j\cdot} = \begin{bmatrix} y_{j1} & y_{j2} \end{bmatrix}^T$ has a covariance matrix of

$$
\text{var}(y_{j\cdot}) = \text{var}(\xi_{j\cdot}) + \text{var}(\varepsilon_{j\cdot}) = \Sigma_\xi + d_j^{-1}\Sigma_\varepsilon = d_j^{-1}(d_j\Sigma_R + I)\Sigma_\varepsilon
\tag{20}
$$

where $\Sigma_R = \Sigma_\xi \Sigma_\varepsilon^{-1}$ is the matrix version of variance ratio. The V matrix is

$$282 \quad V = \operatorname{var}\begin{bmatrix} y_{1\cdot} \\ y_{2\cdot} \\ y_{3\cdot} \end{bmatrix} = \begin{bmatrix} d_1^{-1}(d_1\Sigma_R + I)\Sigma_\varepsilon & 0 & 0 \\ 0 & d_2^{-1}(d_2\Sigma_R + I)\Sigma_\varepsilon & 0 \\ 0 & 0 & d_3^{-1}(d_3\Sigma_R + I)\Sigma_\varepsilon \end{bmatrix} \quad (21)$$

283   The inverse of matrix V is simply obtained via

$$284 \quad V^{-1} = \begin{bmatrix} d_1(d_1\Sigma_R + I)^{-1}\Sigma_\varepsilon^{-1} & 0 & 0 \\ 0 & d_2(d_2\Sigma_R + I)^{-1}\Sigma_\varepsilon^{-1} & 0 \\ 0 & 0 & d_3(d_3\Sigma_R + I)^{-1}\Sigma_\varepsilon^{-1} \end{bmatrix} \quad (22)$$

285   where V is an $(np) \times (np)$ matrix but $d_1\Sigma_R + I$ and Σ_ε are $p \times p$ matrices. In a typical GWAS

286   population, n may vary from a few hundreds to a few thousands but we cannot imagine the

287   number of traits be $p > 20$ unless we are talking about molecular traits, e.g., transcriptomic

288   traits.

290   4.3 The expectation and maximization (EM) algorithm for estimating the covariance
291   matrices

292   The decorrelation of polygenic effects makes all individuals independent. The log likelihood

293   function only deals with matrices of dimension 2×2 or $p \times p$ for p traits. Let $\theta = \{\Sigma_\xi, \Sigma_\varepsilon\}$ be

294   the parameter set, while the β vector is not a vector of parameters in REML (it is a vector of

295   parameter for ML). The restricted log likelihood function is

$$296 \quad \begin{aligned} l(\theta) = &-\frac{1}{2}\sum_{j=1}^{n}\left(\log|d_j\Sigma_R + I| + \log|d_j^{-1}\Sigma_\varepsilon|\right) - \frac{1}{2}\log\left|\sum_{j=1}^{n}X_j^T(d_j\Sigma_\varepsilon^{-1})(d_j\Sigma_R + I)^{-1}X_j\right| \\ &-\frac{1}{2}\sum_{j=1}^{n}(y_{j\cdot} - X_j\hat{\beta})^T(d_j\Sigma_\varepsilon^{-1})(d_j\Sigma_R + I)^{-1}(y_{j\cdot} - X_j\hat{\beta}) \end{aligned} \quad (23)$$

297   where

$$298 \quad \hat{\beta} = \left[\sum_{j=1}^{n}d_j X_j^T(\Sigma_R d_j + I)^{-1}X_j\right]^{-1}\left[\sum_{j=1}^{n}d_j X_j^T(\Sigma_R d_j + I)^{-1}y_{j\cdot}\right] \quad (24)$$

is not a parameter vector but a function of θ. The covariance parameters are found by

maximization of $L(\theta)$. The MIXED procedure PROC MIXED in SAS is readily available to

perform the variance component estimations. However, PROC MIXED becomes

computationally inefficient and may fail to converge properly because of the large number of

parameters. To address these issues, we adopt an Expectation-Maximization (EM) algorithm. EM

algorithm guarantees a monotonic increase in likelihood and achieves stable convergence.

In the EM algorithm framework, the missing values in the complete-data log likelihood function
are the polygenic effects, ξ_j for $j = 1, \cdots, n$, where each ξ_j is an $p \times 1$ vector. We first update β
using

$$\beta^{(t+1)} = \left[\sum_{j=1}^{n} d_j X_j^T (d_j \Sigma_\xi^{(t)} + I)^{-1} X_j \right]^{-1} \left[\sum_{j=1}^{n} d_j X_j^T (d_j \Sigma_\xi^{(t)} + I)^{-1} y_{j.} \right] \tag{25}$$

310 Given ξ_j for $j = 1, \cdots, n$, the polygenic variance and covariance matrix is define as

$$\Sigma_\xi = \frac{1}{n} \sum_{j=1}^{n} \xi_j \xi_j^T \tag{26}$$

312 When ξ_j is missing, we simply replace them by the conditional expectations to update this
313 matrix,

$$\Sigma_\xi^{(t+1)} = \frac{1}{n} \sum_{j=1}^{n} E(\xi_{j.} \xi_{j.}^T \mid y_{j.}) = \frac{1}{n} \sum_{j=1}^{n} \left[E(\xi_{j.} \mid y_{j.}) E(\xi_{j.}^T \mid y_{j.}) + \text{var}(\xi_{j.} \mid y_{j.}) \right] \tag{27}$$

315 where

$$E(\xi_{j.} \mid y_{j.}) = \hat{\xi}_{j.} = d_j \Sigma_\xi^{(t)} (d_j \Sigma_\xi^{(t)} + I)^{-1} (y_{j.} - X_j \beta^{(t)}) \tag{28}$$

317 and

$$\text{var}(\xi_{j.} \mid y_{j.}) = \Sigma_\xi^{(t)} - d_j \Sigma_\xi^{(t)} (d_j \Sigma_\xi^{(t)} + I)^{-1} \Sigma_\xi^{(t)} \tag{29}$$

319 Finally, the updated residual covariance matrix is

$$\Sigma_\varepsilon^{(t+1)} = \frac{1}{n-1} \sum_{j=1}^{n} d_j (y_{j.} - X_j \beta^{(t)} - \hat{\xi}_{j.}) y_{j.}^T \tag{30}$$

321 When the iterations converge, we report the two covariance matrices as the estimated parameters
322 denoted by $\hat{\theta} = \{ \hat{\Sigma}_\xi, \hat{\Sigma}_\varepsilon \}$ and the estimated covariance matrix ratio denoted by $\hat{\Sigma}_R = \hat{\Sigma}_\xi \hat{\Sigma}_\varepsilon^{-1}$. This
323 matrix ratio will be treated as constants and used in genome scanning for associations of
324 genome-wide markers with multiple traits.

4.4 Scanning the entire genome for associations

The method described so far is under the null model, i.e., the pure polygenic model where
intercepts are the only fixed effects in the model. To scan the genome for associations, we can
take one of two approaches: (1) Add a marker to the null model as an additional fixed effect and
repeat the method for every marker of the genome. There is no approximation involved and thus
it is called the exact method. (2) Take the variance matrix ratio estimated from the null model
$(\hat{\Sigma}_R)$ as a constant matrix and scan the entire genome one marker at a time. The fixed effects,

including the effects of the current marker, and the residual covariance matrix are estimated for

each marker. The model for scanning the markers is

$$
\begin{bmatrix} y_{11} \\ y_{12} \\ y_{21} \\ y_{22} \\ y_{31} \\ y_{32} \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_1 \\ X_2 & 0 \\ 0 & X_2 \\ X_3 & 0 \\ 0 & X_3 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} Z_{1k} & 0 \\ 0 & Z_{1k} \\ Z_{2k} & 0 \\ 0 & Z_{2k} \\ Z_{3k} & 0 \\ 0 & Z_{3k} \end{bmatrix} \begin{bmatrix} \gamma_{1k} \\ \gamma_{2k} \end{bmatrix} + \begin{bmatrix} \xi_{11} \\ \xi_{12} \\ \xi_{21} \\ \xi_{22} \\ \xi_{31} \\ \xi_{32} \end{bmatrix} + \begin{bmatrix} \varepsilon_{11} \\ \varepsilon_{12} \\ \varepsilon_{21} \\ \varepsilon_{22} \\ \varepsilon_{31} \\ \xi_{32} \end{bmatrix}
\tag{31}
$$

where Z_{jk} is the genotype indicator of individual j for marker k and γ_{tk} is the effect of marker k

on trait t for $t = 1, 2$. Define the following variables for individual j,

$$ y_{j\bullet} = \begin{bmatrix} y_{j1} \\ y_{j2} \end{bmatrix}, \ \xi_{j\bullet} = \begin{bmatrix} \xi_{j1} \\ \xi_{j2} \end{bmatrix}, \ \varepsilon_{j\bullet} = \begin{bmatrix} \varepsilon_{j1} \\ \varepsilon_{j2} \end{bmatrix}, \gamma_k = \begin{bmatrix} \gamma_{k1} \\ \gamma_{k2} \end{bmatrix}, \ X_j = \begin{bmatrix} X_j & 0 \\ 0 & X_j \end{bmatrix}, \text{ and } Z_{jk} = \begin{bmatrix} Z_{jk} & 0 \\ 0 & Z_{jk} \end{bmatrix} $$

A more compact expression of equation (31) is

$$ y_{j\bullet} = X_j \beta + Z_{jk} \gamma_k + \xi_{j\bullet} + \varepsilon_{j\bullet} \tag{32} $$

The expectation and variance of $y_{j\bullet}$ is

$$ \mathrm{E}(y_{j\bullet}) = X_j \beta + Z_{jk} \gamma_k \tag{33} $$

and

$$ \mathrm{var}(y_{j\bullet}) = V_j = d_j^{-1}(d_j \Sigma_R + I)\Sigma_\varepsilon = H_j \Sigma_\varepsilon \tag{34} $$

where $H_j = d_j^{-1}(d_j \Sigma_R + I)$ and the variance matrix ratio Σ_R is estimated from the null model

prior to scanning the genome. The fixed effects include both β and γ_k, which are estimated

with

$$ \begin{bmatrix} \hat{\beta} \\ \hat{\gamma}_k \end{bmatrix} = \begin{bmatrix} \sum X_j^T H_j^{-1} X_j & \sum X_j^T H_j^{-1} Z_{jk} \\ \sum Z_{jk}^T H_j^{-1} X_j & \sum Z_{jk}^T H_j^{-1} Z_{jk} \end{bmatrix}^{-1} \begin{bmatrix} \sum X_j^T H_j^{-1} y_{j\bullet} \\ \sum Z_{jk}^T H_j^{-1} y_{j\bullet} \end{bmatrix} = \begin{bmatrix} A_{XX} & A_{XZ} \\ A_{ZX} & A_{ZZ} \end{bmatrix}^{-1} \begin{bmatrix} A_{Xy} \\ A_{Zy} \end{bmatrix} \tag{35} $$

where $A_{Xy} = \sum X_j^T H_j^{-1} y_{j\bullet}$, $A_{XX} = \sum X_j^T H_j^{-1} X_j$ and so on. Let us also define

$$ A_{yy} = \sum H_j^{-1} y_{j\bullet} y_{j\bullet}^T $$

The intercepts β are nuisance parameters that can be removed prior to estimate γ_k via the

annihilator matrix (Xu, 2024) so that γ_k are estimated explicitly. Let us define the following

quantities

$$C_{ZZ} = A_{ZZ} - A_{ZX} A_{XX}^{-1} A_{XZ}$$

354
$$C_{Zy} = A_{Zy} - A_{ZX} A_{XX}^{-1} A_{Xy}$$

$$C_{yy} = A_{yy} - A_{yX} A_{XX}^{-1} A_{Xy}$$

355 The estimated γ_k is

356
$$\hat{\gamma}_k = C_{ZZ}^{-1} C_{Zy} \tag{36}$$

357 with a variance of

358
$$\text{var}(\hat{\gamma}_k) = C_{ZZ}^{-1} \hat{\Sigma}_\varepsilon \tag{37}$$

359 where

360
$$\hat{\Sigma}_\varepsilon = C_{yy} - C_{yZ} C_{ZZ}^{-1} C_{Zy} \tag{38}$$

361 is the estimated residual covariance matrix and $C_{yZ} = C_{Zy}^T$. The Wald test statistic for $H_0 : \gamma_k = 0$

362 is

363
$$W_k = \hat{\gamma}_k^T \left[\text{var}(\hat{\gamma}_k) \right]^{-1} \hat{\gamma}_k = \hat{\gamma}_k^T \Sigma_\varepsilon^{-1} C_{ZZ} \hat{\gamma}_k \tag{39}$$

364 which follows approximately a Chi-square distribution with p degrees of freedom.

4.5. Dataset

We evaluated the performance and computational efficiency of the new method using three
genetic association datasets from rice, human, and mouse populations. Each dataset includes
genome-wide genotype data and multiple phenotypic traits, enabling a range of multivariate
GWAS applications. The rice dataset consists of 1,495 hybrid rice individuals (Huang et al.,
2015). For the human dataset, we analyzed a subset of 5,000 unrelated white British individuals
from the UK Biobank (Sudlow et al., 2015) cohort for demonstration purposes. The mouse
dataset was obtained from the CFW outbred population study (Parker et al., 2016), available
from the Dryad Digital Repository (doi:10.5061/dryad.2rs41).

For the rice and mouse datasets, we applied standard quality control procedures, including
removal of SNPs with minor allele frequency below 0.01 and individuals or variants with
missing data. For the UK Biobank data, genotype data were filtered using PLINK2 with
thresholds of 0.05 for missing genotype rate and minor allele frequency, along with linkage
disequilibrium pruning (Chang et al., 2015). Detailed information is provided in Supplementary
Table S1. All phenotypes were standardized prior to analysis.

4.6. Simulation analysis

To evaluate the type I error control and statistical power of our method, we performed null simulations and alternative simulations using real genotype data from the rice and mouse datasets. We excluded the human dataset from simulations due to its large sample size and dense marker coverage, which made simulations computationally impractical.

For null simulations, phenotypes were generated under a pure polygenic model without any marker effects. Specifically, we simulated 100 sets of phenotypes based on the real genetic relatedness matrix and the estimated genetic and environmental covariance matrices from the rice and mouse data. We randomly selected 10,000 SNPs from the real genotype data. We then computed p-values for all combinations of these SNPs and the simulated phenotypes, resulting in one million tests per dataset. For alternative simulations, we added SNP effects into the phenotypes to assess empirical power. We first selected 1,000 evenly spaced SNPs from a set of SNPs that showed no significant association with either trait ($P > 0.1$ in all univariate and multivariate tests). Each selected SNP was assigned an effect on the first trait to explain a fixed proportion of phenotypic variance (PVE). The effect on the second trait was scaled to explain either 20% or 40% of PVE of the second trait, with effects either in the same or opposite direction across traits. These synthetic effects were added to the original phenotypes to generate datasets with known causal SNPs. Power was calculated as the proportion of causal SNPs detected at the Bonferroni-corrected significance level with initial significance level of 0.05. We then applied both EMmvGWAS and GEMMA to these simulated datasets to compare their empirical power under various effect sizes and trait correlation scenarios.

Declarations

Ethics approval and consent to participate

This study did not involve the collection of any new human or animal data. All analyses were conducted using previously published datasets with existing ethical approval, including UK Biobank data accessed under an approved application. No additional ethics approval or consent was required for this work.

Consent for publication

Not applicable

Availability of data and materials

The data set of 1,495 hybrid rice varieties is available at the Rice Haplotype Map Project database (http://www.ncgr.ac.cn/RiceHap4). The CFW mouse datasets can be available at Dryad Digital Repository (doi:10.5061/dryad.2rs41). UK Biobank (UKB) data can be accessed through the UKB Research Analysis Platform (RAP) following approval via the UKB access system (https://www.ukbiobank.ac.uk).

Competing interests

The authors declare that they have no competing interests

Funding

The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Authors' contributions

SX conceived the proposed method. CT analyzed and interpreted the data and generated the figures in the manuscript. CL and QW provided processed UKB dataset. YC provided the hybrid rice dataset. XW advised on the GEMMA implementation. CT wrote the manuscript. SX supervised the research and edited the manuscript. All authors reviewed the manuscript.

Reference

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*, *4*(1), s13742–13015–10047–13748.

Hackinger, S., & Zeggini, E. (2017). Statistical methods to detect pleiotropy in human complex traits. *Open biology*, *7*(11), 170125.

Huang, X., Yang, S., Gong, J., Zhao, Y., Feng, Q., Gong, H., Li, W., Zhan, Q., Cheng, B., & Xia, J. (2015). Genomic analysis of hybrid rice varieties reveals numerous superior alleles that contribute to heterosis. *Nature communications*, *6*(1), 6258.

Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S.-y., Freimer, N. B., Sabatti, C., & Eskin, E. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nature genetics*, *42*(4), 348–354.

Korte, A., Vilhjálmsson, B. J., Segura, V., Platt, A., Long, Q., & Nordborg, M. (2012). A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature genetics*, *44*(9), 1066–1071.

Parker, C. C., Gopalakrishnan, S., Carbonetto, P., Gonzales, N. M., Leung, E., Park, Y. J., Aryee, E., Davis, J., Blizard, D. A., & Ackert-Bicknell, C. L. (2016). Genome-wide association study of behavioral, physiological and gene expression traits in outbred CFW mice. *Nature genetics*, *48*(8), 919–926.

Porter, H. F., & O'Reilly, P. F. (2017). Multivariate simulation framework reveals performance of multi-trait GWAS methods. *Scientific reports*, *7*(1), 38837.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., & Landray, M. (2015). UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine*, *12*(3), e1001779.

Turley, P., Walters, R. K., Maghzian, O., Okbay, A., Lee, J. J., Fontana, M. A., Nguyen-Viet, T. A., Wedow, R., Zacher, M., & Furlotte, N. A. (2018). Multi-trait analysis of genome-wide association summary statistics using MTAG. *Nature genetics*, *50*(2), 229–237.

Visscher, P. M., Wray, N. R., Zhang, Q., Sklar, P., McCarthy, M. I., Brown, M. A., & Yang, J. (2017). 10 years of GWAS discovery: biology, function, and translation. *The American Journal of Human Genetics*, *101*(1), 5–22.

Xu, S. (2024). Estimating genetic variance contributed by a quantitative trait locus: removing nuisance parameters. *Genetics*, *227*(4), iyae095.

Zhang, Z., Ersoz, E., Lai, C.-Q., Todhunter, R. J., Tiwari, H. K., Gore, M. A., Bradbury, P. J., Yu, J., Arnett, D. K., & Ordovas, J. M. (2010). Mixed linear model approach adapted for genome-wide association studies. *Nature genetics*, *42*(4), 355–360.

Zhou, X., & Stephens, M. (2014). Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature methods*, *11*(4), 407–409.