1   **Reduced vomeronasal receptor diversity and impaired pheromone detection in Tex15**
2   **knockout mice**
3

4   Nader Boutros Ghali[1], Paige Kramer[1], Joshua Danoff[1], Ryan Edwards[1], Hetanshi Patel[1], Nusrath
5   Yusuf[1], Zain Zaidi[1], Kevin Monahan[1]
6
7   [1] Department of Molecular Biology and Biochemistry, Rutgers University, Piscataway, NJ,
8   08904, USA
9
10 Correspondence to be sent to Kevin Monahan, kevin.monahan@rutgers.edu

14
Abstract

The mouse vomeronasal organ (VNO) detects pheromones, which provide vital
information needed to find a mate, detect predators, or alert to the presence of an intruder.
Within the VNO, pheromones are sensed by two families of G-protein Coupled Receptors
(GPCRs) the type 1 and type 2 vomeronasal receptors (V1Rs and V2Rs). However, little is
known about the regulatory mechanisms that control the expression of V1R and V2R genes by
vomeronasal sensory neurons (VSNs). Here we identify a transcriptional repressor, *Testis-*
expressed gene 15 (*Tex15*), that regulates vomeronasal receptor gene choice and is required for
the proper functioning of the vomeronasal system in regulating male behavior. We find that
Tex15 is transiently expressed by VSN precursors prior to strong VR gene expression. The
absence of *Tex15* results in a pervasive change in the expression of V1R and V2R genes, which
results in a less diverse repertoire of VR expressing cells and includes a dramatic reduction in the
expression of specific receptors that have been tied to intermale aggression. Accordingly, *Tex15*
knockout mice exhibit lowered activation in the Accessory Olfactory Bulb (AOB) after exposure
to male odorants, and a loss of stereotyped aggression between male mice. Taken together, these
results show that *Tex15* plays a critical role in pheromone sensing by ensuring that VSNs express
a diverse set of receptor proteins.

Introduction

Introduction

Olfaction plays a critical role in the survival and reproductive success of many terrestrial
animals. The vomeronasal organ (VNO) is the key organ for detecting conspecific and
allospecific pheromones from the environment, which can convey vital information such as
another animal's social position in the hierarchy, the receptivity of a female to mating, and the
presence of unfamiliar intruder animals (Fu et al., 2015; Ibarra-Soria et al., 2014; Itakura et al.,
2022; Leypold et al., 2002; Wysocki, 1979). Within the VNO, vomeronasal sensory neurons
(VSNs) are responsible for detecting chemical signals and transmitting this information to the
anterior olfactory bulb (AOB), from which it is relayed further on to the relevant neuronal
circuits (Itakura et al., 2022). Mice with mutations that affect VSN function exhibit impaired
social behaviour such as a lack of aggression in males or abnormal sexual behaviour in females
(Ibarra-Soria et al., 2014; Itakura et al., 2022; Kondo et al., 2024; Leypold et al., 2002).
In mice, VSNs recognize chemical signals using G-protein coupled receptor proteins
from the type 1 and type 2 families of vomeronasal receptors, known as the V1Rs and the V2Rs.
The V1R genes are expressed in a monogenic and monoallelic fashion by VSNs in the apical
layer of the epithelium, such that each apical VSN expresses only one type of receptor out of the
230 genes in the V1R family (Dietschi et al., 2022; Hills et al., 2024). In contrast, basal layer
VSNs express specific combinations of the 160 V2R genes. The V2R genes are split between
four sub-families: A, B, C, and D. Each basal VSN expresses one A, B, or D family gene
together with one or more genes from the C subfamily, which includes seven members: *Vmn2r1-*
7. The pairing is not random, as individual A, B, D genes tend to be expressed together with
either *Vmn2r1* or with the other C family members. Expression of A, B, and D genes precedes
expression of C genes, so the choice of an A, B, or D gene may instruct subsequent C family
expression (Akiyoshi et al., 2018; Ishii & Mombaerts, 2011). In addition, in rodents a small
number of VSNs express a receptor from a family of immune GPCRs, the formyl peptide
receptors (Fprs). This family expanded in the rodent lineage after an ancestral Fpr was
translocated into a VR gene cluster and gained expression in VSNs (Boillat et al., 2021; Dietschi
et al., 2022).
VSN progenitors commit to expressing a type 1 or type 2 VR gene over VSN
differentiation. VSNs originate from a population of globose basal cells (GBCs) found in the
marginal zones of the VNO (Dietz et al., 2025). These GBCs differentiate into immature VSNs
(iVSNs) that are committed to either the apical (V1R-expressing) or basal (V2R-expressing)
lineage (Dietz et al., 2025; Hills et al., 2024). The mechanism behind this divergence involves
multiple regulatory proteins, with differentiating basal cells expressing combinations of ATF5,
Bcl11b, and Notch signaling factors (Katreddi et al., 2022). Perturbing these regulatory proteins
results in a disorganised VNO with a different distribution of apical and basal VSNs and a
corresponding change in VR expression (Katreddi et al., 2022; Lin et al., 2018; Nakano et al.,
2015). However, beyond the specification of neuronal type, the mechanism that controls the
choice of VR genes for expression remains unknown.

Here we show that a transcriptional repressor known for its role in repressing
transposable elements in the male germ-line, *testis expressed gene 15 (Tex15)*, is expressed by
iVSNs and is required for wild-type patterns of VR gene expression, VNO function, and
stereotyped intermale aggression. We show that differentiating VSN progenitors transiently
express *Tex15* RNA and protein. The peak of *Tex15* transcription coincides with the split
between apical and basal VSNs lineages. Mice that are homozygous for a *Tex15* null allele
exhibit altered expression of VR genes and altered abundances of neurons expressing an
individual VR gene. When exposed to soiled male nesting material, *Tex15* null mice exhibit
reduced c-Fos expression, a maker of neuronal activity, in downstream neurons in the accessory
olfactory bulb. Finally, male *Tex15* null mice fail to attack intruder males in a resident-intruder
paradigm and instead exhibit increased exploratory anogenital sniffing behaviour.

Results

Single-cell RNA-seq has established *Tex15* as a marker of the immediate neuronal
progenitor stage in the olfactory sensory neuron lineage in the main olfactory epithelium (Brann
et al., 2025; Pourmorady et al., 2024), but its expression in the VSN lineage has not been
characterized. To determine whether *Tex15* is expressed at any point during VSN differentiation,
we examined previously published single-cell RNAseq data from mouse VNO (Hills et al.,
2024). This dataset contains eight mice, four collected at postnatal day 56 (P56) and four
collected at postnatal day 14 (P14). Since we were interested in examining the immature VSNs,
we used the P14 samples, which have a higher count of immature VSNs but otherwise resemble
the P56 samples. This cohort has two males and two females. Consistent with prior work, UMAP
analysis generates distinct clusters of cells expressing markers of mature apical and basal VSNs
as well as clusters corresponding to various stages of VSN progenitors (Figure S1A). We
excluded a cluster of atypical VSNs, noted by Hills et al, that express markers of both apical and
basal VSNs, since these cells do not express *Tex15* and their connection to the canonical set of
VSN progenitors is uncertain.
Tex15 expression is restricted to cells in the neuronal lineage (Figure S1B, C). Thus, to
examine *Tex15* expression over VSN differentiation, we used marker genes to identify cell
clusters corresponding to the neuronal lineage and then isolated and reclustered these cells. We
then assigned each cluster to a stage of differentiation based upon the expression of known
markers, namely: *Ascl1* for early progenitors, *Neurod1* for immediate neuronal progenitors,
Gap43 for immature VSNs, *Gnai2* for mature apical VSNs, and *Gnao1* for mature basal VSNs
(Figure 1A). Notably, *Tex15* transcript levels peaks in the INP population coinciding with the
divergence of INP cells into the apical and basal differentiation lineages (Figure 1B, C).
To examine the precise timing of *Tex15* expression, we conducted pseudotime analysis of
the neuronal lineage (Figure 1D). Consistent with the cell stage analysis, *Tex15* transcripts peak
after *Neurod1* and prior to *Gap43* (Figure 1E, F) in both the V1R and V2R lineages. Next, we
examined how *Tex15* aligns with the onset VR gene expression. To accomplish this, we
determined the highest expressed V1R and V2R genes within each cell in the corresponding

lineage. In the basal lineage, we took the additional step of analyzing the C-subfamily V2Rs
separately from the A, B, D subfamily V2Rs. We find that in the apical lineage, *Tex15*
expression peaks just prior to a dramatic rise in V1R gene transcripts (Figure 1G). Similarly, in
the basal lineage, we observed that *Tex15* expression precedes A, B, D subfamily expression,
which in turn precedes, C-subfamily expression (Figure 1H). The temporal separation between
the rise in A, B, D transcript levels and the rise in C-subfamily transcript levels is consistent with
a prior study that observed that expression of A, B and D subfamilies of V2R genes was
detectable by RNA in situ hybridisation (ISH) in immature cells prior to the expression of a C
subfamily gene (Ishii & Mombaerts, 2011). We conclude that *Tex15* expression is tightly
restricted to a time point in the development of VSNs, which precedes VR expression.
We next sought to determine whether TEX15 protein is present in immature VSNs in the
VNO. For these experiments, we used VNO tissue sections postnatal day 8 (P8) mice bearing a
Neurog1-GFP reporter gene (Gong, et. al., 2003), which labels INPs and iVSNs in the marginal
zones. To visualize TEX15 expression, we raised a mouse monoclonal antibody against an
epitope in the central region of TEX15 (Figure S2A), and then used it for immunohistochemistry
with a GFP antibody to enhance the endogenous Neurog1-GFP signal. We observe *Tex15*
immunoreactivity in the nuclei of cells in the marginal zones of the VNO, all of which also
exhibit labeling with Neurog1-GFP (Figure 1I). We also observe immunoreactivity along the
basal membrane below the basal VSN, but this is retained in Tex15 knockout tissue and thus is
likely non-specific (see below). Thus, we conclude that TEX15 protein is abundant in the nuclei
of VSN progenitors found in the marginal zone of the VNO.
To examine the effects of *Tex15* we used a *Tex15* null allele (Tex15KO) in which the
majority of the *Tex15* coding sequence has been replaced by beta galactosidase (Yang et al.,
2008). Homozygous Tex15KO mice and littermate controls were obtained by crossing Tex15 +/-
heterozygous mice. To confirm that the knockout lacks *Tex15* expression in the VNO, we
performed a *Tex15* immunofluorescence staining in postnatal day 2 KO mice and littermate
controls (Figure 2A). As above, nuclear *Tex15* immunoreactivity is present in the marginal
regions of the VNO, although we also observe occasional labeled nuclei throughout the rest of
the VNO in heterozygous and wildtype animals (Figure S2B). The nuclear signal is completely
absent from both regions in Tex15KO tissue. Otherwise, the overall structure and organization of
the VNO appear similar in Tex15KO mice.
Based upon its expression in INP cells, we hypothesized that Tex15 might regulate the
expression of VR genes. To test this, we performed RNAseq on VNO tissue obtained from adult
Tex15KO mice and heterozygous controls. We sequenced poly-A selected RNAseq libraries
from four animals per genotype, and then performed differential gene expression analysis using
DESeq2 (Love et al., 2014). As expected, the samples cluster by genotype when subjected to
principal component analysis (Figure 2B). Differential expression analysis identifies 283 genes
that exhibit a greater than 50% change in expression between KO and Het animals ($p_{adj} < 0.05$
for >50% change in expression) Most differentially expressed genes are V1Rs and V2Rs: 151
out of a total of 283. However, the total expression of V1R and V2R genes is not affected; the

sum of transcript counts from each lineage are similar between Het and KO animals (Figure
S3A). Rather, the expression of individual receptors is significantly different (Figure 2D).
Notably, the changes to V2R expression are restricted to the A, B, and D subfamilies, 102 out of
160 genes of which are differentially expressed in the KO (Figure 2E). None of the seven C-
family V2Rs, which are co-expressed depending on prior A, B, or D choice, are differentially
expressed in the Tex15KO (Figure 2F). Interestingly, we also see reduced expression of more
than half of the FPR genes, with 4 significantly decreased in Tex15KO mice (Figure S3B).
Taken together, these results suggest that Tex15 regulates the choice of a receptor protein
by VSNs, but suggests that it may not affect the downstream processes that influence the
subsequent selection of a C-subfamily V2R. Consistent with this idea, among the non-VR genes
that are differentially expressed in the Tex15KO VNOs are four H2-MV group genes: *H2-M1*,
H2-M9, *H2-M11*, and *H2-M10.4* (Figure S3C). The H2-MV genes are expressed in specific
combinations with VR genes, and the first three of these are specifically co-expressed with
Vmn2r82 and *Vmn2r81 (Devakinandan et al., 2024)*, which are downregulated in the knockout
(Figure S3D).
Overall, the Tex15KO results in higher expression of a few VR genes, whereas many
more exhibit decreased transcript levels. This results in a reduction in the diversity of the VR
gene repertoire (Figure 2G, H). We quantified VR gene diversity by calculating the Shannon
diversity index value for all VR gene expression levels in Tex15KO and control samples. We
observe a significant reduction in these scores, indicating reduced VR diversity in KO mice (t(df)
= 3.9187, p-value = 6.491e-05) (Figure 2I).
VR and FPR genes are arranged in clusters across chromosomes, and this clustered
organization contributes to their regulation (Dietschi et al., 2022). We asked whether the altered
expression of VR genes in Tex15KO animals reflected the chromosome cluster location of each
VR gene. Grouping VR genes by cluster did not reveal any consistent patterns as to which VRs
were upregulated and which were downregulated. Many VR clusters contain both upregulated
and downregulated VR genes, nor are there any individual VR clusters that exhibit uniform
increase or decrease in expression (Figure S3E, F).
Since V1R and V2R A, B, and D genes are expressed monogenically within the
population of VSN, the changes we observed by bulk RNAseq could arise from either a change
in the number of cells expressing each VR genes, reflecting a change in the frequency of VR
choice, or a change in the levels of VR expression within the same number of cells. To test
whether differentially expressed VR genes are expressed by a different number of cells in
Tex15KO animals, we performed RNA in situ hybridization (ISH) to label individual cells
expressing specific VR genes in tissue sections from the VNO. Due to the very high sequence
similarity among VR genes, we relied on a chromogenic ISH strategy (BaseScope) that can
target genes that vary over only short regions, as few as 50 base pairs. For this analysis, we first
sought to identify candidate VR genes that had large differences in raw expression values
(transcripts per million) as well as a large fold change between conditions. Even with a 50 base
sequence length there was only one VR which met our criteria and was not predicted to cross

react with other VR genes, *Vmn1r202* (log2fold change = -3.45, p_{adj}= 4.89e-35). As a positive
control for the BaseScope assay, we also examined the expression of a C family VR, *Vmn2r5*,
that is not differentially expressed (log2fold change = 0.09, p_{adj}= 0.20). We dissected the VNOs
of Tex15KO and littermate controls and counted the nuclei present in the epithelial layer to see
what percentage of nuclei were labeled by each probe. As expected, the *Vmn2r5* probe is widely
expressed in the basal layer of the VNO (Figure S4A) and is similar in Tex15KO and control
tissue sections. In contrast, we observe significantly fewer cells expressing *Vmn1r202* in
Tex15KO mice (F(1,3.63)=11.68, p=0.031, using a mixed linear model with a fixed effect of
genotype and a random effect of subject), in accordance with the transcriptional changes
observed by bulk RNA-seq of Tex15KO and control VNO (Figure 3A). Importantly, we still
observe strongly positive cells in the KO animals (Figure S4B), but at a reduced rate compared
to control tissue, which is consistent with an alteration in the frequency of receptor choice rather
than expression per cell.
Our prior results indicate that Tex15KO VNOs contain a reduced diversity of VSNs.
Since the expression of a diverse repertoire of VRs is thought to facilitate the detection of signals
from conspecifics, we asked whether the reduced VSN diversity in Tex15KO mice might impair
the response to mouse derived odorants. We tested this hypothesis by quantifying the expression
of c-Fos, an immediate early gene that is expressed in response to neuronal activity in the mitral
cells of the accessory olfactory bulb (AOB). These are the first layer of cells downstream of
VSN and serve to process the information encoded in pheromones and relay it to the downstream
targets in the brain (Imamura et al., 2020). To test this, *Tex15* knockout mice and wildtype
controls were placed in a new cage that contained either soiled male nesting material or clean
nesting material for 2 hours (Figure 4A), and then we assayed for c-Fos expression in the AOB.
To facilitate identification of the AOB, these experiments were performed with mice bearing an
OMP-ires-GFP reporter gene, which labels VSN axons located within the glomeruli of the
Accessory Olfactory Bulb (AOB) (Figure 4B). As expected, there is a significant genotype by
condition interaction ($F_{(1,8)}$=6.64, p=0.033) (Figure 4C). Wild type mice exposed to dirty bedding
had significantly higher rates (p = 0.0121) of c-Fos positive nuclei (32.3%± 2.63%) than
wildtype mice exposed to clean bedding (20.3% ± 3.79%). In contrast, *Tex15* knockout mice
exposed to clean bedding and dirty bedding have similar rates of c-Fos expression (17% ±
2.95% vs 18.7% ± 2.36%, p = 0.5697). Moreover, c-Fos expression in the AOBs of Tex15KO
mice exposed to dirty bedding was similar to c-Fos express in the AOBs of wildtype mice
exposed to clean bedding (p= .9357), supporting a failure to detect male pheromones in the KO.
Taken together, these results suggest that the alterations in VR expression rates observed in
Tex15KO mice impair the response of the vomeronasal system to male derived odorants.
The impaired activation of the AOB in response to male odorants suggests that Tex15KO
mice may also exhibit altered behavior in response to male cues. The VNO plays a critical role in
many innate behaviours, such as intermale aggression or identifying the sex of the other mouse
(Leypold et al., 2002). To test VNO function in Tex15KO mice we chose the resident intruder
experiment, as a functioning VNO is key to eliciting the stereotyped aggression from the male

resident (Itakura et al., 2022; Leypold et al., 2002). A male mouse in an established territory will
react aggressively toward an unknown male intruder (Leypold et al., 2002). This assay typically
uses sexually experienced males as the resident, as they exhibit greater responses than sexually
naive males (Koolhaas et al., 2013). However, due to widespread changes in VR receptor
diversity, we chose to use sexually naive males for these experiments, to avoid potentially
confounding effects of *Tex15* on mating behavior.
We ran the resident intruder with three 10-minute trials over the course of three days
while comparing control and *Tex15* knockout naive males interacting with intact intruders that
are age-matched (Figure 5A). An attack latency of 600 seconds indicates that the resident did not
initiate an attack during the trial. However, Tex15KO males exhibited significantly different
behaviour as calculated by Fisher test (p= 0.002997); Tex15KO male residents never initiated an
attack against an intruder at any point during any of the trials (Figure 5B). We noted that the
knockout residents spent more time anogenitally sniffing their intruders when compared to the
control (p=0.048) (Figure 5C).

Discussion

We find that *Tex15* is expressed at a critical time point in VSN development, coinciding with the
divergence between the basal and apical populations and shortly preceding the onset of VR gene
transcription. In mice lacking *Tex15*, the overall expression of VR genes is unchanged, but the
relative abundance of individual VR genes is altered. Most VR genes exhibit reduced expression,
whereas a few are expressed at higher levels, resulting in a reduction in the overall diversity of
VR gene expression. We show that the downregulated genes are a result of lowered frequency of
being chosen. These transcriptional changes are accompanied by reduced AOB activation in
response to soiled male nesting material. We also observe that Tex15KO mice exhibit a lack of
stereotyped intermale aggression and increased ano-genital sniffing behaviour. Taken together,
these findings indicate that *Tex15* has a crucial role in ensuring that VR receptor choice results in
diverse outcomes, generating a VNO in which many subtypes of VSN are represented.
The transient expression of *Tex15* during a precise developmental window, coupled with
the striking alterations in VR gene expression observed in *Tex15* knockout mice, implicates
Tex15 as an important regulator of VR gene choice. The loss of *Tex15* affects all of the main
chemoreceptor receptor families expressed by VSNs, the V1Rs, the subfamily A, B, and D
V2Rs, and the FPRs, implying these receptors share a *Tex15*-dependent choice mechanism.
Conversely, gene regulatory decisions that are downstream of VR choice, such as V2R C family
expression and H2-MV expression, seem to respond to the overall changes in the VR repertoire.
What then is the role of *Tex15* in regulating VR gene choice? In testes, *Tex15* associates
with PIWI family proteins MILI and MIWI2, which mediate piRNA-directed methylation and
silencing of transposable elements. Tex15KO mice exhibit reduced DNA methylation of LINE1
and LTR family transposable elements in testes, which is accompanied by increased transposon
expression (Schöpp et al., 2020; Wang et al., 2005; Yang et al., 2020). Expression of PIWI
family members is very low or absent in the VNO, making it unlikely that the altered VR

expression results from altered piRNA directed gene regulation. However, it is possible that
Tex15 associates with the same gene silencing machinery that carries out the piRNA directed
DNA methylation and heterochromatin deposition on transposable elements. This would
implicate a role for heterochromatin deposition in regulating VR gene choice, similar to what has
been proposed for the olfactory receptor genes (Bashkirova et al., 2023; Magklara et al., 2011).
These results also suggest a possible role for small RNA directed silencing in VR gene
regulation, which has an incompletely understood role in the olfactory system (D. Yang et al.,
2020). Identifying the protein partners of Tex15 in the VNO will be critical for characterizing its
tissue specific mechanism of action.
278   Alternatively, it is possible that the observed effects on VR gene expression are a direct
result of dysregulation of transposable elements in Tex15KO mice. The transcription start sites
(TSSs) of over half of VR genes were found to be upstream of LINE1 elements and, moreover,
these LINE1 elements are transcribed (Pascarella et al., 2014). It may be that this LINE1
transcription influences subsequent VR gene choice, and that alterations in this transcription in
Tex15KO mice results in the observed changes in VR gene expression. In this scenario, variation
in the proximity of transposable elements to the transcriptional start site of VR genes may
account for the variable effect of the Tex15KO on VR gene expression. Future work should
examine the Tex15KO mice for potential changes in transposable element regulation that might
be upstream of altered VR expression.
288   We observe striking defects in the function of the vomeronasal system in Tex15KO mice.
Indeed, the responses to male bedding observed in the accessory olfactory bulb of Tex15KO
mice are comparable to those from wildtype mice housed in a clean environment. It is unclear
whether these differences arise directly from the altered repertoire of expressed VR genes, or
some other defect in VSN signaling in Tex15KO mice. However, this loss of sensitivity to male
odorants aligns with a dramatic reduction in aggressive behavior in Tex15KO male mice in a
resident-intruder behavioural assay. Tex15KO male mice never attacked a male intruder across
any of our trials. This loss of aggressive behaviour was accompanied by a significant increase in
anno-genital sniffing behaviour. We speculate that this increased exploratory behaviour may
reflect a response to deficient signaling through the VNO pathway.
298   We show that *Tex15* expression is restricted to a key stage of VSN differentiation,
corresponding to the split between the V1R and V2R lineages, and directly preceding the onset -
of VR gene expression. As such, *Tex15* could serve as an excellent marker for this crucial stage
of differentiation. A large portion of the literature has focused on this stage of VSN
differentiation (Devakinandan et al., 2024; Enomoto et al., 2011; Katreddi et al., 2022; Lin et al.,
2018). As such, Tex15 may serve as a useful driver for genetic tools designed to target this
population. Thus, our work identifies *Tex15* as both a key regulator of VR choice and VNO
function as well as a useful marker for a key stage in VSN differentiation.
306
307
308

309 **Methods**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
C-Fos Antibody	Abcam	Cat#ab190289
GFP Antibody	Antibodies Inc	Cat#gfp-1020
1H11-A3-5 Tex15 Antibody	Homemade	
IgY (H+L) Goat anti-Chicken, Alexa Fluor® 488, Invitrogen	Fisher	Cat#A11039
Cy™3 AffiniPure Goat Anti-Mouse IgG, Fcγ subclass 1 specific	Jackson Immunoresearch	Cat#115-165-205
IgG (H+L) Highly Cross-Adsorbed Donkey anti-Rabbit, Alexa Fluor® 555, Invitrogen	Fisher	Cat#A31572
Critical commercial assays		
Basescope Duplex Assay	ACD	
Direct-Zol RNA Miniprep	Zymo Research	Cat#R2050
Bulk RNA-seq	Admera Health	
Deposited data		
P14 VNO Single Cell Dataset	Hills Jr. et. al.	https://doi.org/10.7554/eLife.97356.3
Experimental models: Organisms/strains		
Tex15 KO mouse	Yang, et. al., 2008	https://doi.org/10.1083/jcb.200709057
OMP-IRES-GFP	Mombaerts et al., 1996	https://www.jax.org/strain/006667
Neurog1-EGFP	Gong, et. al., 2003	#010618-UCD
Software and algorithms		
R software(4.5.2)	R Core Team 2016	https://www.r -project.org/
RStudio(2024.12.1.563)	Rstudio Team 2020	https://rstudio.com/
Tidyverse(2.0.0)	Wickham H, et. al. (2019)	doi:10.21105/joss.01686

Dplyr(1.2.0)	Wickham H, François R, Henry L, Müller K, Vaughan D (2026)	https://dplyr.tidyverse.org
ggplot2(4.0.2)	Wickham H (2016)	https://ggplot2.tidyverse.org.
rstatix(0.7.3)	Kassambara A (2023)	https://rpkgs.datanovia.com/rstatix/
ggpubr(0.6.3)	Kassambara A (2025)	https://rpkgs.datanovia.com/ggpubr/.
DEseq2 (3.22)	Love, M.I., Huber, W. and Anders, S., 2014	10.1186/s13059-014-0550-8
Monocle3(3)	Cao et al, Nature 2019	https://cole-trapnell-lab.github.io/monocle3
Seurat(5.4.0)	Hao. Y. et al 2023	https://cran.r-project.org/web/packages/Seurat/citation.html
SeuratObject(5.3.0)	Satija R, et. al. 2023	https://satijalab.r-universe.dev/SeuratObject/citation
Vegan(2.7.3)	Oksanen J et. al. 2025	https://vegandevs.github.io/vegan/authors.html
naniar(1.1.0)	Tierney N, Cook D (2023)	https://www.jstatsoft.org/article/view/v105i07
forcats(1.0.1)	Wickham. H. 2025	10.32614/CRAN.package.forcats
Fiji(2.17.0)	Schindelin et al. 2012 DOI: doi:10.1038/nmeth.2019	https://imagej.net/software/fiji/
afex(1.5.1)	Singmann H. et. al. 2016 DOI: 10.32614/CRAN.package.afex	https://CRAN.R-project.org/package=afex
emmeans(2.0.2)	Lenth & Piaskowski 2026	https://rvlenth.github.io/emmeans/authors.html
Trim Galore (0.6.5)	https://doi.org/10.5281/zenodo.5127898	https://github.com/felixkrueger/trimgalore
Cutadapt (5.1)	Martin M. 2011 DOI: 10.14806/ej.17.1.200	https://cutadapt.readthedocs.io/en/stable/
STAR (2.7.11b)	Dobin et al 2013	https://github.com/alexdobin/STAR

	DOI: 10.1093/bioinformatics/bts635	
samtools (1.19.2)	Danecek et al 2021 DOI: 10.1093/gigascience/giab008	https://www.htslib.org/
htslib (1.21)	Bonfield et al 2021 DOI: 10.1093/gigascience/giab007	https://www.htslib.org/

Tex15 Antibody Generation

DNA encoding the Tex15 immunogen was cloned from mouse genomic DNA then inserted into pGex4T-2. The protein was expressed in *E. coli* and purified from the insoluble pellet by solubilizing with Sarkosyl, TX100 and Chaps at a ratio of 2% to 4% to 2.5%. Mice were immunized with 100 ug of protein emulsified in Freund's Adjuvant and were subsequently bled. A mouse was boosted with Tex15 protein in PBS and spleen cells were fused with an NS-1 parental cell line using Roche PEG1500 (Sigma 10783641001). The fused cells were plated in IMDM 20% FCS with HAT selection. Cells were grown for 10 days and then the supernatants from the growing cells were screened for the presence of anti-Tex15 antibody by screening against Tex15 fusion protein in an ELISA. ELISA positives were screened on P5 olfactory epithelia from Tex15KO and heterozygous mice. Clone 1H1 was determined to be positive after screening on tissue blocked with Donkey Fab anti Mouse IgG, 30ug/ml (Jackson ImmunoResearch 715-007-003) for 1 hour at RT or overnight at 4°C prior to staining.

Tex15 immunogen, aa 597-1070:
RDDKNPNEAKEHNTDNINGSEKQDCLANDHFTNIVEMREIKSNTEVEILNSEECFT
FNSFRGKNGKPAETASSESEAVEQRHAPNDQRGLEHLVSSFPEIEGSSVCVASNATKQI
VGTTVLTVSTSLGDHQKDELKEICSSESSDLGLVKHSISECEIDTDKDKLQDFHQLVN
ENSALKTGLGSEIEVDLEHDNASVFQQNMHSQGNDLCEEFELYESLKSRIDWEGLFGS
SYEEIESSSFARREGTDQHSSTECNCVSFCSQDKRELHNPIFLPDLQVTITNLLSLRI
SPTDESLELKDNFYKQVTESTEPETNKEGNASGFGMCSQPSGENSSFSCANKFGNSVQ
ESGDVSKSESSHSSNSSHNTHVDQGSGKPNNDSLSTEPSNVTVMNDKSKCPTKSKPVF
NDTRNKKDMQSRSSKRTLHASSSRGQNIANKDLREHETHEKKRRPTSHGSSDRFSSLS
QGRIKTFSQS

Tex15 Immunofluorescence

Mice under P8 were decapitated and their heads were fixed in 4% PFA in PBS for 30 minutes at room temperature. The heads were then washed three times for 5 minutes in PBS before being submerged in a 30% sucrose and PBS mixture at 4°C for 3 days. Afterwards, the heads were embedded in Optimal Cutting Temperature compound (Cat#:25608-930) and then sectioned with a cryostat to a thickness of 14 microns. Sections were transferred onto superfrost plus slides (Cat# 48311-703) and stored at -20C. The slides were washed in a solution of PBS +

0.1% Triton (0.1% PBST) before adding a primary antibody solution of anti-Tex15 mouse
monoclonal antibody (1H1A324-3) diluted at 1:10 in 4% goat serum and 0.1% PBST. The slides
were incubated with the primary antibody in a humidified chamber at 4°C overnight. The next
day, they were washed three times for 5 minutes with 0.1% PBST. The secondary antibody goat
anti mouse IgG1 Cy3 (Cat#: 115-165-205) and DAPI (Sigma-Aldrich: MBD0015-1ML) were
diluted at 1:1000 in 4% goat serum in 0.1% PBST and applied on the slides. They were
incubated in a humidified chamber at room temperature for 1 hour. The slides underwent three
more 5-minute washes in 0.1% PBST. They were then mounted with VectaShield Hardset
Mounting Medium and left to cure for 15 minutes in the dark. The images were taken with a
Leica TCS SP8 tauSted 3X confocal and imaged at 40x. Tilescan images were then stitched
together in FIJI with the Grid/Collection Stitch plugin (, et al, 2009).
354

Tex15 and Ngn-GFP Immunofluorescence
Tissue sections were prepared as described above, and placed into a blocking solution of
4% goat serum in 0.1% PBST for 1 hour at RT. The slides were then incubated with primary
antibodies for Tex15 (1H1A324-3) diluted at 1:10 and anti-GFP (Cat#: gfp-1020) at 1:1000 in
4% goat serum and 0.1% PBST. The slides were incubated in a humidified chamber at 4C
overnight. The next day, they were washed three times for 5 minutes with 0.1% PBST. The
slides were then incubated with secondary antibodies, goat anti-mouse IgG1 Cy3 (Cat#: 115-
165-205), IgY goat anti-chicken Alexa Fluor 488 (Cat#: A11039), and DAPI (Sigma-Aldrich:
MBD0015-1ML), diluted at 1:1000 in 4% goat serum 0.1% PBST in a humidified chamber at
room temperature for 1 hour. The slides were then washed three more times, each 5-minute with
0.1% PBST, and then were mounted with VectaShield Hardset Mounting Medium and left to
cure for 15 minutes in the dark. The images were taken with a Leica TCS SP8 tauSted 3X
confocal at 63x. The brightness and contrast of the Tex15 channel was adjusted to reduce non-
specific background reactivity.
369

RNA extraction and RNA-seq
The VNO was either dissected and used immediately or frozen at -80°C. RNA was
extracted by solubilizing the tissue in Trizol (Cat#: 15596026). The extracted RNA was isolated
from the Trizol solution using Zymo Research Direct-zol RNA Miniprep Kits (Cat# R2050).
Purified RNA was assessed on a Thermo Fisher Scientific Nanodrop One for concentration. It
was then quality controlled for RIN in an Agilent Tapestation 4150. Samples which had a RIN
greater than 7 were sent to Admera (South Plainfield, NJ, USA) for DNAse treatment, library
preparation based upon poly-A tail selection, and paired-end sequencing
378

379 **BaseScope Imaging and Counting**
380   BaseScope probes were generated by ACD. Hybridization and chromogenic staining
381   were performed using the ACD fixed/frozen protocol for the BaseScope duplex and BaseScope

382   single channel assay. Images were collected on a Zeiss Axio Imager.A1 with a lumineer camera.
383   Red, green, and blue images were taken and overlayed to create a full colour image.
384
C-Fos Imaging and Counting
Mice were placed in clean cages for 10 minutes to habituate. After that time, either clean
nesting material or soiled nesting material from male cages was added to the cage, and the mice
were incubated for 2 hours with the material. The mice were euthanised with CO_2 and their
olfactory bulbs were dissected out and fixed on ice for 20 minutes with 4% PFA in PBS. The
bulbs underwent two PBS washes on ice before being placed in 50% 0.5M EDTA in PBS at 4°C
overnight, followed by a solution of 30% sucrose in PBS at 4°C for 3 days or until the tissue
sank to the bottom. The bulbs were embedded sagittally in OCT and sectioned on a cryostat at 14
microns. For each animal, we collected three evenly-spaced 14-micron-thick sections of AOB
tissue.
The slides were blocked in a solution of 4% donkey serum and 4% goat serum in 0.1%
PBST for 1 hour. They were incubated in the primary antibody solution of c-Fos (Cat#:
ab190289) at 1:500 and anti GFP (Cat#: gfp-1020) at 1:1000 to enhance the fluorescence
overnight in a humidified chamber at 4°C. The next day, the slides were washed three times for 5
minutes with 0.1% PBST. The secondary antibody donkey anti rabbit Alexa 555, Alexa Fluor
488 (Cat#:A21206), and DAPI (Sigma-Aldrich: MBD0015-1ML) were diluted at 1:1000 in 4%
goat and donkey serum in 0.1% PBST and applied onto the slides. They were then incubated in a
humidified chamber at room temperature for 1 hour. The slides underwent three more 5-minute
washes in 0.1% PBST. The slides were mounted with VectaShield Hardset Mounting Medium
and left to cure for 15 minutes in the dark. The brightness and contrast of the c-Fos channel was
adjusted to reduce non-specific background reactivity. Images were collected on a Widefield
Leica DMI8 Thunder System at 40X. Each image included the glomerular and mitral layer of the
AOB.
408
Resident Intruder
Resident males were isolated for a period of 10 days prior to the trials. The resident males
were naive and had never been housed with a female. A wildtype, naive intruder of similar age
and size to the resident would be placed into the resident's cage and their interactions recorded
for 10 minutes and scored for social interactions, anogenital sniffing, and aggression based on
how many seconds they spent engaging in each behaviour. The trial was repeated 3 times over
the course of 3 days, each time with a new intruder who had not been previously used in a trial.
The recordings were scored by two independent and blinded scorers and an average of
the values and scores was generated by determining how many seconds the residents spent
engaging in certain behaviours. The scores of knockout residents and wildtype residents were
then compared.
420
421

Quantification and Statistical Analysis

Statistical analysis of histology and behavioral data was completed using mixed models to account for repeated measurements from the same subjects. Mixed models were generated, as described below, using the *afex* package in R. Post-hoc analyses were completed using the *emmeans* package in R.

Single Cell Analysis

Single cell RNA-seq analysis was performed in R using Seurat. Count matrices from GSE252365 were annotated with relevant metadata including age, sex, and replicate number. The replicates were combined into one object and then low quality cells and possible doublets were discarded by filtering on the fraction of mitochondrial reads, total counts, and total RNAs detected. The data was then log normalized, highly variable genes were identified, and these genes were used for principal components analysis. The different samples were then integrated using Seurat's "integrateLayers" function and Anchor-based Canonical Correlation Analysis. The integrated data was then clustered using 30 PCA dimensions and a clustering resolution of 4. Clusters from the neuronal lineage were identified by examining the expression of established marker genes (Hills et al., 2024), and these cells were selected for further analysis. As noted in Hills et al., 2024, there is a population of VSNs which do not follow the canonically established development trajectory and thus were excluded from the subset.

After subsetting, the cells from the neuronal lineage were reclustered, as described above, and then clusters were identified based on the expression of marker genes. Then a pseudotime analysis was carried out with Monocle3. For subsequent analyses of the V1R VSN and V2R VSN lineages, all cells prior to the lineage split (2 clusters) were retained and then we selectively retained either the V1R VSN or V2R VSN cells, as appropriate. These cells were once again reclustered and used for a lineage specific pseudotime analysis with Monocle3. We then plotted expression over pseudotime by combining pseudotime values for each cell with log normalised gene counts.

RNA-seq analysis

Adapters were removed from sequencing data using Trim Galore and cutadapt. STAR was used to align trimmed reads to the mouse genome (mm10). For alignment, the lab first used a custom transcriptome annotation file based upon ENSEMBL release GRCm38.102, but updated to include expert curated olfactory receptor and vomeronasal receptor gene annotation from Barnes et al 2020 and Dietschi et al 2022, respectively. Aligned reads were read into R using Rsubread, and then differential expression analysis was performed using DEseq2.

BaseScope Analysis

The Basescope images were analyzed to determine the number of nuclei that were red positive relative to negative nuclei. The specifications for a positive versus a negative cell were determined by comparing experimental BaseScope images to images that used a negative and

control bacterial probe provided by ACD. The BaseScope images were unaltered after
acquisition. The signal in the negative images served as a standard for background signal. A
positive cell required signal strength higher than the negative control in a concentrated area
surrounding or adjacent to a nucleus. FIJI counting plugin was used for counting expressing cells
in the epithelial layer of the VNO. Positive cells were divided by the total number of cells to
generate a percent positive in each image. Mixed models were used to analyze the number of
cells with positive BaseScope signal, with a fixed effect of genotype and a random effect of
subject.
470
C-Fos Analysis
Each condition included three Tex15KO and three wildtype controls, a total of n=12. The
brightness and contrast of the c-Fos and DAPI channels were adjusted to facilitate distinguishing
nuclear signal. The mitral layer was demarcated, and then a blinded investigator scored each
image by counting the total number of nuclei in the mitral layer and scoring them for the
presence of c-Fos. Nuclei in the mitral layer were considered positive for c-Fos expression if the
euchromatin of a nucleus was occluded by c-Fos expression on the same focal plane as the
nucleus. The FIJI counting plugin was used for counting c-Fos expressing nuclei in the mitral
layer of the accessory olfactory bulb. The percent of activated neurons is the fraction of mitral
cell nuclei that were identified as positive for c-Fos. Mixed models were used to analyze c-Fos
counts, with a fixed effect of the interaction of genotype and condition and a random effect of
subject. Post-hoc analysis compared the estimated marginal means of each condition within each
genotype.
484
Analysis of Behavioral Data
Attacks
Mice were labeled "attackers" if they attacked the intruder mouse during any of the three
trials. Fisher's exact test was used to analyze the proportion of attackers in each genotype.
Anogenital sniffing
Mixed models were used to analyze the duration of anogenital sniffing, with fixed effects
of genotype and trial and a random effect of subject.
492
Author contributions
Conceptualization: KM (Lead), NBG (Equal)
Data curation: NBG (lead), KM (equal)
Formal analysis: NBG (lead), KM (supporting), PK (Supporting), JD (Supporting) ZZ
(Supporting)
Funding acquisition: KM
Investigation: NBG (Lead), PK (supporting), RE (supporting), HP (supporting), NY
(supporting), ZZ (supporting)
Methodology: NBG (Lead), KM (Equal)

Project administration: KM (lead), NBG (supporting)

Resources: NBG (Lead), PK (Equal), KM (Supporting)

Supervision: KM (Lead), NBG (Supporting)

Validation: NBG (Lead), KM (Supporting)

Visualization: NBG (lead), KM (supporting)

Writing – original draft: NBG (lead), KM (Equal)

Writing – review & editing: NBG (lead), KM (Equal), JD (supporting)

Acknowledgements

The authors thank Irina Ruzina, Mackenzie Leung, Melina Updale, and Sanjana Jethani for their help with the resident intruder experiments. We would also like to thank Madison Freestone for helping refine and edit the manuscript.

The Tex15 monoclonal antibody was produced by Susan Brenner-Morton, Director of the Antibody Platform within the Scientific Platforms at the Zuckerman Institute, Columbia University. We gratefully acknowledge her expertise in the development of this antibody. We are grateful for the support of the Antibody Platform and the Zuckerman Institute Scientific Platforms.

We acknowledge the School of Arts and Science-Human Genetics Institute of NJ (SAS-HGINJ) Imaging Core Facility for use of their confocal and widefield microscope.

Funding

This work was supported by the National Institutes of Health [1R35GM146901 to K.M.] and a Rita Allen Scholar Award [K.M.].

Data availability

Raw and processed RNA-seq data reported in this paper are available from GEO under accession number GSE326692. VNO scRNA-seq data were previously described (Hills et al., 2024) and are available from GEO under accession number GSE252365.

Code availability

All code required to reproduce the figures in this manuscript is available at https://github.com/MonahanLab/Tex15_in_VNO/tree/main.

Conflict of Interest

The authors declare no competing financial interests.

 Figure 1

Figure 1. *Tex15* is transiently expressed by VSN progenitors before VR choice. (A) UMAP projection of cells from the neuronal VSN lineage, color coded by stage of differentiation. (B) *Tex15* transcript levels across the neuronal lineage. (C) Transcript levels and frequency of expression for developmental marker genes and *Tex15* in each cell population. (D) Pseudotime values calculated across the neuronal lineage. (E) Transcript levels of developmental markers and Tex15 over pseudotime in the isolated V1R lineage. (F) Transcript levels of developmental markers and Tex15 over pseudotime in the isolated V2R lineage. (G) Transcript levels of Tex15 and the transcript level of the most abundant V1R transcript level in each cell over pseudotime in the V1R lineage (H) Transcript levels of Tex15 and the transcript level of the most abundant V2R A/B/D

and V2R C in each cell in the V2R lineage. (I) Immunohistochemistry for TEX15 in VNO tissue sections from postnatal day eight (P8) mice. Images show the marginal zone of the VNO. Scalebar: 10μm. In all plots, transcript levels are expressed as log normalized counts.

554   **Figure 2**

555
556   Figure 2. *Tex15* knockout mice have altered VR expression and reduced diversity. (A) Coronal VNO section from a P2 Tex15KO mouse and
557   littermate control labeled with TEX15 antibodies (cyan) and DAPI, which labels DNA (gray). Tex15 immunoreactivity is detected in nuclei of

the control mice (arrows). Inset: Stitched tile scan of the full VNO. Scalebar: 10μm. (B) PCA analysis of VNO bulk-RNAseq libraries. (C) Abundance of *Vmn1r*, *Vmn2r*, and *Fpr* genes within the sets of significantly upregulated and significantly downregulated genes in the Tex15KO. (D) Scatter plot of V1R transcript levels in Tex15KO versus control VNOs. (E) Scatter plot of V2R A/B/D family transcript levels in Tex15KO versus control VNOs. (F) Scatter plot of V2R C-family transcript levels in Tex15KO versus control VNOs. (G) The cumulative expression of V1Rs ranked by expression within each genotype. (H) The cumulative expression of V2R A, B, and D family genes ranked by expression within each genotype. (I) Shannon's Diversity Index of all VR transcripts in control and Tex15KO libraries df = 3.9187, p-value = 6.491e-05 (t-test). For bulk RNA-seq, differentially expressed genes were determined based upon $p_{adj} < 0.05$ for a >50% change in expression (Wald test).

567   **Figure 3**

568
569   Figure 3. Tex15KO results in fewer cells expressing *Vmn1r202*. (A) The percentage of cells within the VSN layer expressing *Vmn1r202* in Tex15
570   knockout vs control (p=0.006). Each point-line represents the mean and standard error determined by counting multiple sections per mouse. (B)
571   Representative images of a knockout and control VNO. *Vmn1r202* BaseScope chromogenic signal is red. Scalebar: 100μm.
572

573   **Figure 4**

574
Figure 4. Loss of *Tex15* and VR diversity results in lower activation in the AOB. (A) Schematic showing the soiled bedding exposure assay. (B)
Representative immunohistochemistry images showing c-Fos and GFP signal in sagittal sections of the accessory olfactory bulb. The images
shown are tile scans that were stitched together. Scalebar: 250μm (C) The average percentage of activated neurons per mouse, grouped by
genotype and condition. Mean values for each mouse were calculated by averaging across 3 sections. Statistical significance was determined
using a mixed linear model, which included a fixed effect of genotype, condition, and the interaction of genotype and condition, and a random
effect of subject.
581

582   **Figure 5**

A

583
584   Figure 5. *Tex15* knockout males do not show aggressive behaviour towards intruders. (A) Diagram of the resident intruder assay conducted on the
585   mice. (B) Percent of trials in which the resident attacked the intruder. (C) Plot of the amount of time control and Tex15KO mice spent in each
586   trial performing anogenital sniffing behavior.
587

Figure S1

Figure S1. Identifying Tex15 expressing cells within the VNO. (A) UMAP projection of all cells present in VNO scRNA-seq from 14-day old mice. Abundant cell clusters are labeled based upon marker gene expression. (B) Dot plot showing transcript levels and frequency of expression for *Tex15* and VSN developmental markers across each identified cell cluster. (C) Transcript levels for developmental marker genes and Tex15 in all VNO cells. All gene expression UMAP and dot plots show transcript levels expressed as log-normalized counts. Atypical cells expressing both V1 and V2 VSN markers are called "Atypical VSNs".

 Figure S2

A

B

■DAPI ■TEX15

598
599   Figure S2 Detection of TEX15 expression. (A) Schematic showing the location and sequence used for generating a mouse monoclonal antibody
600   against TEX15. (B) The Tex15 specific immunoreactivity (arrowheads) observed in VSN nuclei in VNO tissue sections from control mice is lost
601   in Tex15KO VNO sections, while non-specific immunoreactivity typically observed with mouse monoclonal antibodies remains. TEX15 channel
602   brightness was adjusted identically in both images. Scalebar: 20μm

Figure S3

Figure S3. Transcriptional changes in the *Tex15* knockout. (A) Comparison of the total V1R and V2R transcript levels (sum of tpm) between the
control and experimental conditions. T-test was conducted to determine significance. (B) Scatter plot of FPR transcript levels in Tex15KO versus

control VNOs. Differentially expressed genes are in color. (C) Transcript levels of differentially expressed H2-MV genes. Significance was determined by a Wald Test on normalized counts. (D) Transcript levels of VR genes *Vmn2r81* and *Vmn2r82*, which are associated with the downregulated H2-MV genes. Significance was determined by a Wald Test on normalized counts. (E) Fold change of V1R transcript levels, with genes grouped by cluster. Differentially expressed genes are in color. (F) Fold change of A, B, and D V2Rs transcript levels with genes grouped by cluster. For bulk RNA-seq plots, differentially expressed genes were determined based upon $p_{adj} < 0.05$ for a >50% change in expression (Wald test). Genes that are significantly upregulated in the KO are blue, genes that are downregulated are red.

616   **Figure S4**

A

B

617
618   Figure S4 Tex15KO mice still express the downregulated VRs. (A) Representative images of *Vmn2r5* BaseScope assay on tissue
619   sections from control and Tex15KO mice. Scalebar: 100µm (B) Representative image showing a Tex15KO VNO with a cell
620   strongly expressing *Vmn1r202*. Scalebar: 100 µm.
621

References

Akiyoshi, S., Ishii, T., Bai, Z., & Mombaerts, P. (2018). Subpopulations of vomeronasal sensory neurons with coordinated coexpression of type 2 vomeronasal receptor genes are differentially dependent on Vmn2r1. *The European Journal of Neuroscience*, *47*, 887–900.

Bashkirova, E. V., Klimpert, N., Monahan, K., Campbell, C. E., Osinski, J. M., Tan, L., Schieren, I., Pourmorady, A., Stecky, B., Barnea, G., Xie, X. S., Abdus-Saboor, I., Shykind, B., Jones-Marlin, B., Gronostajski, R. M., Fleischmann, A., & Lomvardas, S. (2023). Opposing, spatially-determined epigenetic forces impose restrictions on stochastic olfactory receptor choice. In *bioRxiv*. https://doi.org/10.1101/2023.03.15.532726

Boillat, M., Carleton, A., & Rodriguez, I. (2021). From immune to olfactory expression: neofunctionalization of formyl peptide receptors. *Cell and Tissue Research*. https://doi.org/10.1007/s00441-020-03393-5

Brann, D. H., Tsukahara, T., Tau, C., Kalloor, D., Lubash, R., Thamarai Kannan, L., Klimpert, N., Kollo, M., Escamilla-Del-Arenal, M., Bintu, B., Bozza, T., & Datta, S. R. (2025). A spatial code governs olfactory receptor choice and aligns sensory maps in the nose and brain. In *bioRxivorg*. https://doi.org/10.1101/2025.05.02.651738

Devakinandan, G. V. S., Terasaki, M., & Dani, A. (2024). Single-cell transcriptomics of vomeronasal neuroepithelium reveals a differential endoplasmic reticulum environment amongst neuronal subtypes. *eLife*, *13*. https://doi.org/10.7554/eLife.98250

Dietschi, Q., Tuberosa, J., Fodoulian, L., Boillat, M., Kan, C., Codourey, J., Pauli, V., Feinstein, P., Carleton, A., & Rodriguez, I. (2022). Clustering of vomeronasal receptor genes is required for transcriptional stability but not for choice. *Science Advances*, *8*(46), eabn7450.

Dietz, A., Senf, K., & Neuhaus, E. M. (2025). Stem cell expression of CXCR4 regulates tissue composition in the vomeronasal organ. *Journal of Cell Science*, *138*(1). https://doi.org/10.1242/jcs.263451

Enomoto, T., Ohmoto, M., Iwata, T., Uno, A., Saitou, M., Yamaguchi, T., Kominami, R., Matsumoto, I., & Hirota, J. (2011). Bcl11b/Ctip2 controls the differentiation of vomeronasal sensory neurons in mice. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *31*(28), 10159–10173.

Fu, X., Yan, Y., Xu, P. S., Geerlof-Vidavsky, I., Chong, W., Gross, M. L., & Holy, T. E. (2015). A molecular code for identity in the vomeronasal system. *Cell*, *163*(2), 313–323.

Gong, S., Zheng, C., Doughty, M. L., Losos, K., Didkovsky, N., Schambra, U. B., ... & Heintz, N. (2003). A gene

expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature*, *425*(6961), 917-925.

Hills, M., Jr, Ma, L., Fang, A., Chiremba, T., Malloy, S., Scott, A., Perera, A., & Yu, C. R. (2024). Molecular, cellular, and developmental organization of the mouse vomeronasal organ at single cell resolution. In *eLife*. https://doi.org/10.7554/elife.97356

Ibarra-Soria, X., Levitin, M. O., & Logan, D. W. (2014). The genomic basis of vomeronasal-mediated behaviour. *Mammalian Genome: Official Journal of the International Mammalian Genome Society*, *25*(1-2), 75–86.

Imamura, F., Ito, A., & LaFever, B. J. (2020). Subpopulations of projection neurons in the olfactory bulb. *Frontiers in Neural Circuits*, *14*, 561822.

Ishii, T., & Mombaerts, P. (2011). Coordinated coexpression of two vomeronasal receptor V2R genes per neuron in the mouse. *Molecular and Cellular Neurosciences*, *46*(2), 397–408.

Itakura, T., Murata, K., Miyamichi, K., Ishii, K. K., Yoshihara, Y., & Touhara, K. (2022). A single vomeronasal receptor promotes intermale aggression through dedicated hypothalamic neurons. *Neuron*, *0*(0). https://doi.org/10.1016/j.neuron.2022.05.002

Katreddi, R. R., Taroc, E. Z. M., Hicks, S. M., Lin, J. M., Liu, S., Xiang, M., & Forni, P. E. (2022). Notch signaling determines cell-fate specification of the two main types of vomeronasal neurons of rodents. *Development* , *149*(13). https://doi.org/10.1242/dev.200448

Kondo, H., Iwata, T., Sato, K., Koshiishi, R., Suzuki, H., Murata, K., Spehr, M., Touhara, K., Nikaido, M., & Hirota, J. (2024). Impaired pheromone detection and abnormal sexual behavior in female mice deficient for ancV1R. *Current Biology: CB*. https://doi.org/10.1016/j.cub.2024.10.077

Koolhaas, J. M., Coppens, C. M., de Boer, S. F., Buwalda, B., Meerlo, P., & Timmermans, P. J. A. (2013). The resident-intruder paradigm: a standardized test for aggression, violence and social stress. *Journal of Visualized Experiments: JoVE*, *77*, e4367.

Lenth R, Piaskowski J (2026). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. R package version 2.0.2, https://rvlenth.github.io/emmeans/.

Leypold, B. G., Yu, C. R., Leinders-Zufall, T., Kim, M. M., Zufall, F., & Axel, R. (2002). Altered sexual and social behaviors in trp2 mutant mice. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(9), 6376–6381.

Lin, J. M., Taroc, E. Z. M., Frias, J. A., Prasad, A., Catizone, A. N., Sammons, M. A., & Forni, P. E. (2018). The

transcription factor Tfap2e/AP-2ε plays a pivotal role in maintaining the identity of basal vomeronasal sensory

neurons. *Developmental Biology*, *441*(1), 67–82.

Magklara, A., Yen, A., Colquitt, B. M., Clowney, E. J., Allen, W., Markenscoff-Papadimitriou, E., Evans, Z. A.,

Kheradpour, P., Mountoufaris, G., Carey, C., Barnea, G., Kellis, M., & Lomvardas, S. (2011). An epigenetic

signature for monoallelic olfactory receptor expression. *Cell*, *145*(4), 555–570.

Nakano, H., Iida, Y., Suzuki, M., Aoki, M., Umemura, M., Takahashi, S., & Takahashi, Y. (2015). Activating

transcription factor 5 (ATF5) is essential for the maturation and survival of mouse basal vomeronasal sensory

neurons. *Cell and Tissue Research*. https://doi.org/10.1007/s00441-015-2283-8

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M, Szoecs E,

Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Borman T, Carvalho G, Chirico M, De Caceres M,

Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, Martino C,

McGlinn D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter Braak C, Weedon J (2026). *vegan:*

Community Ecology Package. R package version 2.8-0, https://vegandevs.github.io/vegan/.

Pascarella, G., Lazarevic, D., Plessy, C., Bertin, N., Akalin, A., Vlachouli, C., Simone, R., Faulkner, G. J.,

Zucchelli, S., Kawai, J., Daub, C. O., Hayashizaki, Y., Lenhard, B., Carninci, P., & Gustincich, S. (2014).

NanoCAGE analysis of the mouse olfactory epithelium identifies the expression of vomeronasal receptors and

of proximal LINE elements. *Frontiers in Cellular Neuroscience*, *8*, 41.

Preibisch, S., Saalfeld, S., & Tomancak, P. (2009). Globally optimal stitching of tiled 3D microscopic image

acquisitions. *Bioinformatics*, *25*(11), 1463-1465.

Pourmorady, A. D., Bashkirova, E. V., Chiariello, A. M., Belagzhal, H., Kodra, A., Duffié, R., Kahiapo, J.,

Monahan, K., Pulupa, J., Schieren, I., Osterhoudt, A., Dekker, J., Nicodemi, M., & Lomvardas, S. (2024).

RNA-mediated symmetry breaking enables singular olfactory receptor choice. *Nature*, *625*(7993), 181–188.

Schöpp, T., Zoch, A., Berrens, R. V., Auchynnikava, T., Kabayama, Y., Vasiliauskaitė, L., Rappsilber, J., Allshire,

R. C., & O'Carroll, D. (2020). TEX15 is an essential executor of MIWI2-directed transposon DNA methylation

and silencing. *Nature Communications*, *11*(1), 3739.

Tierney N, Cook D (2023). "Expanding Tidy Data Principles to Facilitate Missing Data Exploration, Visualization

and Assessment of Imputations." *Journal of Statistical Software*, **105**(7), 1–31. doi:10.18637/jss.v105.i07.

706 Team, R. (2020). RStudio Team. RStudio: Integrated Development for R. *Boston, MA: RStudio Team*.

707 Wang, P. J., Page, D. C., & McCarrey, J. R. (2005). Differential expression of sex-linked and autosomal germ-cell-

708  specific genes during spermatogenesis in the mouse. *Human Molecular Genetics*, *14*(19), 2911–2918.

709 Wysocki, C. J. (1979). Neurobehavioral evidence for the involvement of the vomeronasal system in mammalian

710  reproduction. *Neuroscience and Biobehavioral Reviews*, *3*(4), 301–341.

711 Yang, F., Eckardt, S., Leu, N. A., McLaughlin, K. J., & Wang, P. J. (2008). Mouse TEX15 is essential for DNA

712  double-strand break repair and chromosomal synapsis during male meiosis. *The Journal of Cell Biology*,

713  *180*(4), 673–679.

714 Yang, F., Lan, Y., Pandey, R. R., Homolka, D., Berger, S. L., Pillai, R. S., Bartolomei, M. S., & Wang, P. J. (2020).

715  TEX15 associates with MILI and silences transposable elements in male germ cells. *Genes & Development*,

716  *34*(11-12), 745–750.

717 **Supplementary References**

719 Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D. M., Hill, A. J., ... & Shendure, J. (2019). The single-cell

720  transcriptional landscape of mammalian organogenesis. *Nature*, *566*(7745), 496-502.

721 Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., ... & Satija, R. (2021). Integrated

722  analysis of multimodal single-cell data. *Cell*, *184*(13), 3573-3587.

723 Kassambara A (2023). *rstatix: Pipe-Friendly Framework for Basic Statistical Tests*. R package version

724  0.7.2, https://rpkgs.datanovia.com/rstatix/.

725 Kassambara A (2026). *ggpubr: 'ggplot2' Based Publication Ready Plots*. R package version

726 0.6.3, https://rpkgs.datanovia.com/ggpubr/.

727 Lenth R, Piaskowski J (2026). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. R package version

728 2.0.3,

729 Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data

730  with DESeq2. *Genome Biology*, *15*(12), 550.

731 Mombaerts, P., Wang, F., Dulac, C., Chao, S. K., Nemes, A., Mendelsohn, M., ... & Axel, R. (1996). Visualizing an

732  olfactory sensory map. *Cell*, *87*(4), 675-686.

733 Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M, Szoecs E,

Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Borman T, Carvalho G, Chirico M, De Caceres M,

Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, Martino C,

McGlinn D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter Braak C, Weedon J (2026). *vegan:*

Community Ecology Package. R package version 2.8-0, https://vegandevs.github.io/vegan/.

R Core Team. (2016). R: A language and environment for statistical computing. R Foundation for Statistical

Computing, Vienna, Austria. *http://www. R-project. org/.*

Satija R, Butler A, Hoffman P, Stuart T (2023). *SeuratObject: Data Structures for Single Cell Data*. R package

version 4.1.4, https://github.com/mojaveazure/seurat-object.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., … Cardona, A. (2012). Fiji: an

open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. doi:10.1038/nmeth.2019

Singmann H, Bolker B, Westfall J, Aust F, Ben-Shachar M (2025). _afex: Analysis of Factorial Experiments_. R

package version 1.5-1, <https://CRAN.R-project.org/package=afex>.

Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-

24277-4, https://ggplot2.tidyverse.org.

Wickham H (2025). *forcats: Tools for Working with Categorical Variables (Factors)*. R package version

1.0.1, https://forcats.tidyverse.org/.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D. A., François, R., ... & Yutani, H. (2019).

Welcome to the Tidyverse. *Journal of open source software*, *4*(43), 1686.

Wickham H, François R, Henry L, Müller K, Vaughan D (2026). *dplyr: A Grammar of Data Manipulation*. R

package version 1.2.0, https://dplyr.tidyverse.org.

754