Dynamic alterations of DNA methylation and transcriptome in adaptation to and recovery from the space environment

Liang Lu,[1,4] Yangyang Hao,[2,4] Xue Lin,[3,4] Kai Li,[1,4] Tianzhang Zhai,[2] Fengji Liang,[1]
Linhuan Chen,[2] Liqing Wang,[2] Xueyin Mei,[2] Sidu Feng,[2] Ke Lv,[1] Yanhong Yuan,[1] Zhongquan Dai,[1]
Dong Liu,[1] Hongyu Zhang,[1] Chao Yang,[1] Anna Liu,[2] Linjie Wang,[1] Zhili Li,[1] Shujuan Liu,[1]
Xiaoqian Dai,[1] Chengjia Yang,[1] Chunyan Wang,[1] Peng Sun,[1] Liujia Shi,[1] Chi Zhang,[1]
Jianghui Xiong,[1] Ming Wei,[1] Chong Xu,[1] Zhaoxia Liu,[1] Lina Qu,[1,*] Jian Li,[1,2,*] Yinghui Li[1,*]
[1]State Key Laboratory of Space Medicine, China Astronaut Research and Training Center, Beijing 100094, China
[2]School of Life Science and Technology, Southeast University, Nanjing, Jiangsu Province 210096, China
[3]Department of Bioinformatics, School of Biomedical Engineering and Informatics, Nanjing Medical University,
Nanjing, Jiangsu Province 211166, China
[4]These authors contributed equally
*Correspondence: linaqu@263.net (L.-N.Q.), jianli2014@seu.edu.cn (J.L.), yinghuidd@vip.sina.com (Y.-H.L.)
12

SUMMARY

Space exploration presents tremendous health challenges. Here, we report time series of multi-omic and
phenotypic profiles of seventeen astronauts from six China Manned Space missions with continuous
spaceflight durations ranging from 13 to 180 days. We revealed a key role of DNA methylation regulation
in reshaping gene expression patterns to adapt to the space environment. Long-duration spaceflight
showed more alterations in epigenetic modifications correlated with alternative splicing and protein
acetylation. During recovery, an "overrange rebound" phenomenon was observed, furthermore, a
mathematical model was established to describe this implying important phenomenon. Moreover, we
revealed the correlations between molecular alterations and phenotypic changes such as coagulation
activation and bone intensity loss. Additionally, we performed ground-based simulation experiments to
estimate the impacts of individual stressors in the space environment on DNA methylation. In summary,
our study highlights the importance and complexity of epigenetic regulation in adaptation to and recovery
from the space environment.
26

KEYWORDS

28 Spaceflight, DNA methylation, epigenetics, adaptation to space environment, overrange rebound
29

INTRODUCTION

With the increasing frequency of space explorations[1-3], understanding the mechanisms of human
adaptation to and recovery from space environment has become increasingly important. In space,
stressors that affect human health include microgravity, radiation, confinement/isolation, and distance
from Earth[4], which collectively cause unique physiological changes in the human body[5-7]. Increasing
evidence shows that most of these physiological changes are mediated by molecular events[4]. The
International Space Station (ISS)[8], which serves as a space research laboratory, has already supported
many important molecular biological studies[1,9]. Over the last decade, China has launched a series of
manned spacecraft Shenzhou missions[10,11] and established the China Space Station (CSS), contributing
new knowledge regarding space adaptation[12]. Here, we present analyses of multi-omic and phenotypic
data from biological samples collected from seventeen astronauts who participated in the recent 10-years
China Manned Space missions (M13, a 13-day spaceflight mission; M15, a 15-day spaceflight mission;
M33, a 33-day spaceflight mission; M90, a 90-day spaceflight mission; M180-1, a 180-day spaceflight
mission; M180-2, another 180-day spaceflight mission) (Figure 1). Bulk RNA sequencing, DNA
methylation and phenotypic measurements were applied to analyze the biological samples of the
astronauts mainly at three key timepoints: preflight (T1, more than 1 month before launching), immediate
postflight (T2, around 1 day after returning to Earth) and recovery (T3, more than between 30 and 60

days after returning to Earth). In this study, an epigenetic characteristics of space environment adaptation
and recovery on the basis of the genome-wide DNA methylation landscape from a chronological
perspective was revealed. Furthermore, the results from the comparison between M90 and M180-1
missions were confirmed by the M180-2 mission, which presented the impact of long-duration spaceflight
on the human body. These experiments revealed an important role of DNA methylation as a mechanism
underlying human adaptation to and recovery from spaceflight on basis of data from multiple subjects and
missions. Corresponding ground-based simulation experiments were also conducted[13,14] (Figure 1 and
Table S1), and these findings were compared with those from spaceflights, enabling us to investigate the
impacts of individual space environmental stressors on the human body.

RESULTS

Increased DNA methylation alterations follow the accumulated space environment exposure time
We analyzed the peripheral blood DNA methylation profiles of seventeen astronauts from six China
Manned Space missions, including three short-duration spaceflights (M13, M15 and M33), one medium-
duration spaceflight (M90) and two long-duration spaceflights (M180-1 and M180-2). A comparison of the
DNA methylation profiles of astronauts from the spaceflights with different inflight durations revealed that
DNA methylation changes (T2 vs. T1) increased with increasing space environment exposure time
(Figure 2a). That is, for short periods, i.e., 13 days (M13) and 15 days (M15), little change in DNA
methylation was observed; a few DNA methylation alterations were identified in the 33-day spaceflight
(M33); and obvious DNA methylation changes were detected after 90 days of spaceflight (M90). More
significant global DNA methylation changes were present in the two 180-day spaceflights (M180-1 and
M180-2). Interestingly, decreased DNA methylation was more significant than increased DNA methylation
(T2 vs. T1) in almost all six spaceflights, suggesting that exposure to the space environment initiated an
epigenetic transition toward more open DNA (Figure 2a). Because the six spaceflight missions included in
this study span more than ten years, The DNA methylation analyses of the early spaceflight missions
(M13, M15 and M33) employed the Illumina 450K platform, the analysis of the latest three spaceflights
(M90, M180-1 and M180-2) employed the updated 850K platform. We used consistent probes between
the 450K and 850K platforms, the general tendency that more differentially methylated probes (DMPs)
were found in longer-duration spaceflights was maintained (data not shown).

Through the comparison of the changes in DNA methylation among the six spaceflight missions, several
consistent DNA methylation alteration regions were identified, which seemed to work as growth core
regions. A gradual extending pattern of DNA methylation change initiated from the growth core regions
was shown following increased exposure to the space environment (Figure 2b).

Long-duration spaceflight results in increased DNA demethylation and up-regulation of gene
expression
As significant changes in DNA methylation occurred during medium- and long-duration spaceflights, we
further investigated the similarities and differences in DNA methylation between these two cases and their
biological implications. A global change in the DNA methylation landscape occurred after spaceflight (T2
vs. T1), and most of the changes in DNA methylation were reversed during recovery, restoring the sites to
their preflight status (T3 vs. T2), showing an imperfect symmetrical curve (Figure 3a and Figure S1a).
Moreover, the number of hypomethylated DMPs was greater than the number of hypermethylated ones
during spaceflight, especially for the promoter and enhancer regions (T2 vs. T1) (Supplementary
Information Note S1, Figure 2a and Figure S1b), indicating genome-wide demethylation under exposure
to the space environment. The above phenomenon was consistently observed in M180-2 (Figures S2a
and S2b). Furthermore, longer-duration spaceflight resulted in more loci with DNA methylation changes
than did medium-duration spaceflight (11136 hypermethylated / 22033 hypomethylated DMPs in M90,
41476 hypermethylated / 47759 hypomethylated DMPs in M180-1 and 16468 hypermethylated /33884
hypomethylated DMPs in M180-2, *P* value < 0.05, |deltaBeta| > 0.05, ChAMP, with consistent changes
were observed among all astronauts in the same mission; see "Results of differential methylation analysis
of DNA methylation probes" in https://www.spacelifescience.cn/search/detail?id=24).

The results of the differentially methylated region (DMR) analysis also support the above conclusion that
M180-1 presents more DMRs of demethylation than M90 does in T2 vs. T1 (448 hypermethylated / 109
hypomethylated regions in M90, 78 hypermethylated / 976 hypomethylated regions in M180-1, adjusted P
value < 0.05, ChAMP) (Figure S3a and Table S2). Despite a global demethylation trend indicated by DMP
number in both missions and DMR number in M180-1, M90 exhibited more hypermethylated DMRs, yet
only the hypomethylated DMRs were functionally cohesive enough to yield significant GO enrichment
terms (Figure S3a and Table S2). Moreover, 24.1% (134) of the DMRs in M90 overlapped with the DMRs
in M180-1, including 45 completely overlapping DMRs (Figure S3a and Table S2). These DMRs present
in both missions were enriched for myeloid leukocyte activation, regulation of neutrophil activation and
leukocyte activation involved in the inflammatory response, etc. (Table S2). The demethylated genes
involved in DMRs in M90 were also predominantly immune and inflammation related genes, whereas the
DMRs in M180-1 involved a greater diversity of functional gene sets, such as "locomotory behavior",
"extracellular matrix organization" and "sensory organ morphogenesis" (Table S2). Both missions present
more hypermethylated DMRs than hypomethylated DMRs in T3 vs. T2 (Figure S3b). The DMRs in M180-
2 still exhibited considerable overlaps with those in M180-1 and M90 (Figure S3c).

As the mammalian DNA methylome is shaped by two antagonistic processes, methylation by DNA
methyltransferases (DNMTs) and demethylation by ten-eleven translocation (TET) dioxygenases[15], we
further investigated the expression of methylation-related enzymes. The upregulation of demethylases
(TET2 and TET3) and the downregulation of DNMT1 were observed during spaceflight (Figure 3b and
Table S3), which might be the underlying causes of the observed changes in global methylation patterns.
Similar alterations were observed in M180-2 and hind-limb unloading (HLU) model rats (Supplementary
Information Note S1 and Table S7).

Accordingly, longitudinal bulk RNA-seq data analysis of the astronauts' peripheral blood samples was
performed for M90 and M180-1, which revealed a dynamic global change in gene transcription with
numerous differentially expressed genes (DEGs) identified ($|\log_2 FC| > \log_2(1.5)$, and adjusted P value <
0.05, limma). Additionally, the total number of DEGs was greater in the samples from long-duration flights
than in those from medium-duration flights (Figure 3c). In the immediate postflight samples (T2 vs. T1),
there were more upregulated genes (274 in M90; 1204 in M180-1) than downregulated genes (196 in
M90; 194 in M180-1), revealing genome-wide significantly upregulated transcriptomic changes resulting
from exposure to the space environment (Figure 3c and Table S3). This phenomenon was highly similar
to that in M180-2 (Figure S4a-S4c). 202 upregulated and 74 downregulated genes overlapped in both
M90 and M180-1 (T2 vs. T1), while 180 upregulated and 748 downregulated genes overlapped after
recovery (T3 vs. T2) (Figure 3d).

A substantial negative correlation was observed between the changes in gene expression levels and the
changes in DNA methylation levels ($\log_2$FCs in T2 vs. T1) in the corresponding promoter and enhancer
regions for all expressed genes in both M90 and M180-1 (P value < 0.01, Pearson correlation test),
reflecting the impact of DNA methylation on the regulation of gene expression during adaptation to the
space environment (Figure 3e). This association between gene expression and DNA methylation was
more prominent in the DEGs (T2 vs. T1 or T3 vs. T2) and less pronounced in the unchanged genes.
Furthermore, longer spaceflight was associated with more pronounced negative correlations (Figure 3e).
The correlation between DNA methylation and gene expression observed in M90 and M180-1 was
confirmed in M180-2 (Figure S4d).

Association of differentially expressed alternative splicing isoforms with DNA methylation
Alternative splicing (AS) is a mechanism by which genes produce more than a single mature transcript,
which contributes to adaptation to a new environment[16]. In our study, we found no differential AS events
between the T1 and T3 transcriptomes in M90, but several differential AS events between the T1 and T2
transcriptomes (Table S4). In the comparison of AS between T2 and T1, M180-1 presented more
differential AS events than M90 did (Figure 4a). However, the genes exhibiting AS in the two spaceflight
durations had little overlap. The genes with differential AS events in M90 were enriched mainly in
immune-related pathways, whereas the genes in M180-1 were enriched primarily in epigenetics-related
pathways (Table S4), indicating that epigenetic regulation events may be enhanced during long-duration

spaceflight. In M180-1, histone acetylation-related genes were found to be differentially spliced. For
example, differentially expressed AS isoforms of the histone deacetylase complex subunit Sin3A-
associated protein 18 (*SAP18*)[17], as well as histone acetyltransferase-related genes *BRD1*[18] and
JADE1[19], were detected after long-duration spaceflight (M180-1). In M180-2, another 180-day spaceflight,
we observed differential splicing of the *MEN1* gene, which encodes the histone methyltransferase
complex Menin[20], in the comparison between T2 and T1 (Table S4), The above results suggest that
multiple epigenetic regulatory mechanisms are involved in prolonged adaptation to the space
environment.

The loci with altered DNA methylation levels in the differentially spliced genes, especially hypomethylated
loci, were significantly localized in gene body regions rather than randomly distributed (Figures 4b-4d and
Table S4). Previous studies have reported that DNA methylation plays a role in the identification of exons
involved in AS events, and that the loss of DNA methylation affects splicing patterns on a genome-wide
scale[21,22]. Accordingly, we speculate that DNA methylation may be involved in the regulation of AS events
during adaptation to the space environment.

Changes in proteomics and protein acetylation levels during spaceflight and recovery
Differential analysis of the proteomic profiles revealed 8 proteins that were differentially expressed
postflight in M90 and 14 proteins that were differentially expressed postflight in M180-1 (adjusted *P* value
< 0.05, limma). Reductions in hemoglobin levels were detected in astronauts from both flights (M90 and
M180-1) (Table S5). It has been reported that lower hemoglobin accelerates platelet aggregation, thereby
promoting coagulation[23]. The expression of Ig-like domain-containing, proteins, which are typically
involved in modulating immune responses, decreased postflight in M90[24,25]. The myeloblastin protein,
also known as PRTN3, showed increased expression postflight in M180-1. PRTN3 is a neutrophil-specific
protease primarily associated with the neutrophil immune response and inflammatory processes[26].

The number of proteins with acetylation modifications was generally reduced after both spaceflights (T2
vs. T1) (Table S5), especially M180-1 (*P* value = 0.077, Wilcoxon test, two-tailed) (Figure S5a).
Interestingly, all proteins that were differentially acetylated exhibited 'V' shaped profiles, presenting
decreases in acetylation during spaceflight (T2 vs. T1) and subsequent increases during recovery (T3 vs.
T2) (Figure S5b and Table S5). The functions of these differentially acetylated proteins are related
primarily to the cytoskeleton and metabolic functions. Among these, the deacetylation of ATP-citrate
synthase may affect cholesterol metabolism[27], which is notable as changes in blood cholesterol levels
postflight have been reported[9]. A change in the acetylation level of H1e was found only in M90, which
was consistent with the global acetylation profile, which presented a 'V' shape (Table S5). Changes in H4
histone acetylation at H4K12 and H4K16 were detected in both M90 and M180-1 but presented different
patterns. In M90, acetylation decreased at T2 and T3. However, in M180-1, H4 acetylation levels at
H4K12 and H4K16 increased (T2 vs. T1) and subsequently decreased (T3 vs. T2) (Table S5), suggesting
a difference in H4 histone modification between medium-duration spaceflight and long-duration
spaceflight.

Interestingly, transcriptomic analysis revealed that several histone deacetylase-encoding genes[28] *HDAC4*
($\log_2$FC = 0.58, adjusted *P* value = 0.001, limma), *HDAC5* ($\log_2$FC = 0.63, adjusted *P* value = 0.003,
limma) and *HDAC7* ($\log_2$FC = 0.51, adjusted *P* value = 0.005, limma) were significantly increased in
M180-1 (T2 vs. T1) (Table S3), with a decrease in DNA methylation in the *HDAC7* promoter region. None
of these transcriptomic changes were observed in M90.

Overrange rebound of gene expression during recovery and its underlying mechanisms
Most of the genes whose expression significantly changed during spaceflight tended to recover after
returning to Earth in both M90 and M180-1. A total of 89.8% of the upregulated genes and 90.8% of the
downregulated genes in M90 and 94.6% of upregulated genes and 74.2% of the downregulated genes in
M180-1 showed the reverse change in expression after landing, and the expression tended to return to
the preflight level (T1) (Figure 3c). However, the majority of these genes did not return to precisely the
same level of expression as T1 but rather exceeded the previous expression levels to a small extent,

showing "overshoot" (Figure 5a). During recovery, this overshoot was observed in 72.2% of the
upregulated genes and 74.7% of the downregulated genes in M90 and 73.2% of the upregulated genes
and 55.5% of the downregulated genes in M180-1 (Figure 5b). We name this phenomenon "overrange
rebound" and note that it presents an imperfect symmetrical curve (Figure 5c). This phenomenon was
also observed in M180-2 (Figures S4a and S4c). The functions of these overrange rebound DEGs
(DEG_over) vary, including roles in immunity, the inflammatory response, bone resorption, and
coagulation (Table S3). Furthermore, we established a mathematical model to describe the observed
overrange rebound overshoot (Supplementary Information Note S2).

Furthermore, we found that the expression values of the overrange rebound DEGs were significantly
negatively correlated with their promoter and enhancer regions DNA methylation levels (P value < 0.01,
Pearson correlation test), moreover this correlation was not significant for the unrecovered DEGs (Figure
5d). This finding suggests a role for DNA methylation in recovery. In addition, the overrange rebound
genes exhibited greater degree values in the constructed protein-protein interaction (PPI) network
("Construction of PPI networks related to environmental adaptation" in Methods) than non-rebound
overrange genes did (P value = 0.044 for M90 and P value = 2.5e-05 for M180-1, Wilcoxon test, two-
tailed) (Figure 5e), indicating that they interact with more genes and may be more susceptible to cascade
and feedback mechanisms. Overrange rebound overshoot may be a mechanism, by which the human
body can overcome the inertia established during adaptation to the space environment and accomplish
rapid recovery upon return to Earth.

Interactions of adaptation-related genes suggest that inflammation-like mechanisms coordinate
multisystem adaptation
Weighted gene coexpression network analysis (WGCNA) revealed genes whose expression exhibited a
characteristic chronological pattern according to the course of adaptation and recovery during spaceflight
(M90 and M180-1). These genes were categorized into 7 modules on the basis of their expression
correlation (Table S6), with the blue and brown modules significantly associated with spaceflight time (P
value < 0.05, ANOVA test) (Figure 6a and Table S6). Functional enrichment analysis revealed that the
genes in blue module were related mainly to immune system function, whereas the genes in brown
module were involved in multiple systems of interest (Figures 6b and 6c and Table S6), including
immune, bone resorption, coagulation, and oxidative stress-related gene sets. A predicted space-
adaptation-related PPI network was constructed on the basis of the genes in the brown module, which
reflects potential multisystemic changes induced by the space environment (Table S6). A gene set
interaction network was produced by calculating the interaction score for each pair of gene sets in the 476
biological processes enriched among the 1458 PPI nodes (Table S6). Inflammation-related pathways,
such as the chemokine, TNF signaling and Toll-like signaling pathways, interacted the most strongly with
multisystem-related pathways (Figure 6d and Table S6). Multisystem phenotypic changes induced by the
space environmental exposure, such as bone loss and blood coagulation activation, were identified in
M90 and M180-1 (Table S6).

The bone density of the six astronauts (M90 and M180-1) decreased to some extent after the
spaceflights, according to phenotypic measurements, including ultrasonic calcaneal bone density, lumbar
spine bone density, and left and right hip bone density. The 28-day HLU model experiment also revealed
a reduction in bone density (Supplementary Information Note S3, Figure S6, and Table S7). For
astronauts, the functions "osteoclast differentiation" and "NF-κB signaling pathway" were enriched in the
PPI network of the brown module and generally exhibited upregulated gene expression (Table S6).
Osteoclasts are specific cells responsible for bone matrix degradation[29]. Moreover, the expression of
these two gene sets was significantly negatively correlated with bone density (Table S6). Furthermore,
these genes displayed a general upregulation of expression, with some experiencing decreased DNA
methylation in the promoter region and overrange rebound.

The interaction of nuclear factor-kappa B receptor activator (RANK) with its ligand RANKL activates a
major signaling pathway for osteoclast formation, which regulates adequate osteoclast differentiation
through activation of NF-κB and its downstream targets, and genes related to these molecules were
generally upregulated after both spaceflights (Table S6). In addition, the formation of mature osteoclasts

may require costimulatory signals involving immune receptors associated with ITAM-carrying aptamers
such as FcRγ and DAP12, most of which were significantly upregulated after the two spaceflights (Table
S6). We also observed the upregulation of certain inflammatory factors, such as TLR-4[30,31] and IL1B[32]
(Table S3), that can stimulate the NF-kappa B signaling pathway[33]. The proinflammatory cytokine *IL1B*
was significantly coexpressed with the NF-κB encoding genes *NFKB1* (r = 0.533, P value = 0.020),
NFKB2 (r = 0.796, P value = 7.829e-05), and *RELB* (r = 0.873, P value = 2.241e-06) in the 18 samples
from the spaceflight durations. Moreover, the range of upregulation of the above genes was generally
greater in M180-1 than in M90 (Table S3). Additionally, a greater number of genes in M180-1 exhibited
decreased DNA methylation and overrange rebound (Table S6).

The functions "blood coagulation" and "platelet activation" were also enriched in the PPI network of the
brown module (Table S6). Two of the astronauts presented increased levels of thrombin antithrombin
complex (TAT) in M180-1, indicating activation of the coagulation system[34]. Moreover, all three astronauts
presented an increase in CD62P (P-selectin) after 180 days of spaceflight, indicating that platelet
activation may contribute to coagulation activation[35,36]. Moreover, the expression of "blood coagulation"
and "platelet activation" genes was positively correlated with CD62P and TAT. Similar phenotypic
changes were detected in the 28-day HLU model rats (Supplementary Information Note S3). Gene set
interaction analysis revealed that blood coagulation interacted significantly with platelet activation (Table
S6). Platelets carry a wide range of chemokines and cytokines and play important roles in a variety of
processes, such as hemostasis, wound repair, and pro- and anti-inflammatory activities[37]. This study
revealed a significant interaction between platelet activation and chemokines and cytokines signaling
associated pathways (Table S6). The expression levels of the chemokines *CXCL1* and *PF4*[38] increased
after both spaceflights (Table S3). *PF4* expression was also increased in 28-day HLU model rats.
Hemostasis-related genes encoding a vascular hemophilia factor, von Willegrand factor (vWF) were also
up-regulated after 180 days of spaceflight (Table S3). vWF interacts with collagen-GPVI, initiating a
series of signal transduction molecules and leading to platelet activation[39]. *TBXAS1*, αIIbβ3 integrin, and
ITGA2B, which encodes the coagulation-related protein TXA2 synthase[40], were also found to be up-
regulated (Table S3), which could serve as evidence of coagulation activation[41,42]. Inflammation and
coagulation are two major host defense systems that mutually interact[43,44]. The inflammatory cytokine IL-
1β may facilitate increased expression of COX-2[45,46] (*IL-1B* and *PTGS2*, r = 0.778, P value = 0.0001),
which in turn is linked to increased expression of *TBXAS1*. In addition, we detected elevated postflight
expression of the coagulation factor V encoding gene *F5* after both spaceflights (Table S3); coagulation
factor V is an important cofactor in the blood coagulation cascade. Overall, there was greater up-
regulation of coagulation-related genes in M180-1 than in M90, which may explain the coagulation
activation observed after 180 days of spaceflight. In addition, more genes in M180-1 showed decreased
DNA methylation and overrange rebound (Table S6).

In M180-2, a general postflight decrease in bone density and an increase in TAT were observed, which is
consistent with the findings from the M90 and M180-1 missions. Additionally, the plasma prothrombin
time (PT) and activated partial thromboplastin time (APTT) were reduced, further supporting the activation
of coagulation. However, the level of CD62P decreased. The above important functional gene set also
showed an increasing trend in expression in M180-2 and in HLU model rats, although individual variations
were observed.

We hypothesized that, during adaptation to and recovery from the space environment, many
inflammation-related genes and pathways are recruited to coordinate different organs and systems to
disrupt previous physiological homeostasis and rebuild novel fitness homeostasis as a fast response to
environmental change.

Comparison of ground-based simulation experiments and real spaceflight
Because of the key role of DNA methylation in human adaptation to and recovery from the space
environment revealed by the above findings from real spaceflights, we further investigated the DNA
methylation patterns of different components of space environmental stressors via ground-based
experiments. We compared the DNA methylation profiles of samples from M90 and M180-1 with those of
samples from 90-day head-down bed rest (HDBR) (simulating microgravity), 180-day Controlled

Ecological Life Support System (CELSS) (simulating 180-day isolation)[13], and Mars500 (simulating 520-
day isolation)[14] experiments. Overall, global DNA methylation differences were detected in the 90-day
HDBR and the Mars500 groups, whereas no significant global DNA methylation changes were detected
in the 180-day CELSS (Figures S7a-S7c) group. The group that experienced 90 days of HDBR (R90)
clearly presented significant differences compared with the prebed rest group (Pre15). A portion of
samples at day 60 of HDBR (R60) presented a change in DNA methylation levels, whereas samples from
day 30 exhibited little effect (Figure S7a). The DNA methylation data from Mars500 indicated notable
differences in samples taken before the 60th day compared with samples taken after the 120th day, and
especially samples from the 512th day (Figure S7c).

In all the aforementioned simulation studies, changes in bone phenotypes were observed. Among the
eight 90-day HDBR subjects, more than six presented decreased bone densities in the hip, tibia, and
calcaneus at R60 and R90 (Figure S7d). In the 180-day CELSS study, all four subjects showed a trend
toward reduced bone density in both weight-bearing bones (lumbar spine and femur) and non-weight-
bearing bones (radius and ulna)[47]. The Mars500 study indicated that prolonged confinement in closed
environments can lead to localized reductions in bone mineralization[48]. However, these scenarios, unlike
actual spaceflight and HLU models, induced localized decreases in bone density rather than a significant
overall reduction in mineral density. No direct evidence of coagulation activation was found in the
simulation studies. In the 90-day HDBR samples, the ADP-induced platelet aggregation rates decreased
and the PAC1 levels increased at R30 and R60, but the levels of CD62P and other markers did not
change significantly (Figure S7e). A previous 60-day HDBR study reported that 60 days of 6° HDBR did
not correlate with significant activation of the coagulation system as evaluated via intravascular thrombus
formation in healthy volunteers[49]. In the 180-day CELSS study, an increase in PAC1 was observed, but
CD62P did not exhibit consistent changes (Figure S7f). The 520-day isolation study (Mars500) reported a
slowdown in the intrinsic pathway and final-stage blood coagulation, with a relative increase in the activity
of anticoagulant proteins such as antithrombin III (ATIII) and protein C (PC) inhibitors[50].

Furthermore, our findings revealed that most genes related to osteoclast differentiation and the NF-κB
signaling pathway presented increased expression and decreased DNA methylation (T2 vs. T1) during
both M90 and M180-1 (Table S6 and Figures 7a and 7b). Among the ground-based experiments, the
DNA methylation decreased in the 90-day HDBR experiment but the decrease was minimal and reverted
quickly (Figure 7c). DNA methylation also decreased in the 180-day CELSS experiment on the 90th and
150th days but was not maintained (Figure 7d), and many of these genes were temporarily upregulated in
the 30- and 90-day samples (Table S6). Multiple fluctuating in DNA methylation also occurred during the
Mars500 mission (the 168th and 300th days) (Figure 7e). For the NF-κB signaling pathway, the
fluctuations in DNA methylation did not decrease after 90 days in either HDBR or 180-day CELSS,
suggesting that the effects of these experiments were not sufficient to induce stable modifications at the
DNA methylation level (Figures 7c-7e).

The DNA methylation levels of the coagulation gene set and platelet activation were not consistently
altered in the 90-day HDBR and 180-day CELSS studies, and only transient fluctuations were observed in
the Mars500 mission (Figures 7c-7e).

The range of DNA methylation fluctuations in the ground-based simulated experiments was generally
narrower than those in actual spaceflight, with the largest fluctuations occurring during the Mars500
mission. However, the effect of time on the range of fluctuations in the Mars500 mission experiment was
not found.

DISCUSSION

The NASA Twins Study was the first to systemically reveal the molecular profiles of long-duration
spaceflights on the basis of multi-omics data collected from an astronaut and his ground-based identical
twin[9]. In this study, we revealed a clear dynamic molecular process occurring during the course of
adaptation to and recovery from the space environment on the basis of molecular and phenotypic data
from seventeen astronauts from the six China Manned Space missions. Crucially, widespread promoter
DNA demethylation correlates with upregulation of gene expression, especially during long-duration

spaceflight, which may cause significant upregulation of many genes, suggesting a nonnegligible role for
DNA methylation in the adaptation of the human body to the space environment. With accumulated space
environment exposure time, DNA methylation alterations initiated at certain core seed loci gradually
extended to neighboring genomic regions to expand an open epigenetic state, reshaping the
transcriptome to adapt to the space environment. With increasing spaceflight time, in addition to the
gradual expansion of DNA methylation alterations from a few loci to the full genome, suggesting that
multiple steps and fine turning epigenetic regulations are involved in adaptation to the space environment.

Interestingly, numerous genes were found to use different transcript isoforms during spaceflight, which
indicates that AS may contribute to adaptation to the space environment. Notably, the genes with AS in
M180-1 and M180-2 are involved in histone acetylation. These findings indicate that long-duration
spaceflight results in more changes in epigenetic modifications than medium-duration spaceflight does
and suggest that cross-talk between multiple epigenetic modifications. These epigenetic modifications
such as DNA methylation and protein acetylation, orchestrate transcriptome reshaping adaptation to long-
term space environmental exposure. Our study revealed that epigenetic regulatory mechanisms may
determine the plasticity and elasticity of the human body in the context of rapid and long-term
environmental changes.

An overrange rebound during recovery was observed in multiple cohorts, which may drive the rapid
recovery upon return to Earth. Overrange rebound allows the system to respond more quickly to changes
or perturbations and can help the system overcome any resistance or inertia that might be present in the
system to achieve a more stable and well-regulated state in the long term. This phenomenon can be a
feature of adaptive systems, in which the system learns from its past experiences and adjusts its
response accordingly to refine its responses and adapt to changing environmental conditions. Our data
provide some insights into the regulatory mechanism of overrange rebound by revealing the relationship
between DNA methylation changes and gene expression. Further investigation is necessary to more fully
comprehend the mechanisms involved, which may be more complex.

Gene expression is correlated with phenotype, and a time effect dependent correlation was found
between gene sets associated with immunity and inflammation and those associated with space
environment-induced phenotypes such as bone loss and coagulation activation. Our findings highlight
that inflammation-like mechanisms at the physiological level connect role multiple systems in the
spaceflight adaptation and recovery. Silveira et al reported that mitochondrial stress, as a central
biological hub, drives the health risks of spaceflight[6]. As mitochondria are central hubs of the immune and
inflammatory system[51], our findings are consistent with those of Silveira's study.

Notably, renewed interest in inflammation biology has been driven by the realization that inflammation is
involved in a surprisingly broad range of biological processes, including tissue remodeling, metabolism,
thermogenesis, and nervous system function[52]. In the context of spaceflight, the human body recruits
molecular mediators that are enriched in canonical inflammation pathways, which are connected with the
stress response, metabolism and other biological processes and coordinate with major organs in the body
to quickly establish a physiological adaptation state and eliminate the source of perturbations from the
space environment.

Ground-based simulation experiments provide the opportunity to simulate different components of space
environmental stressors[53-55]. A series of systemic comparisons with samples from multiple spaceflights
revealed that the range of DNA methylation fluctuations in the simulation experiments was narrower than
that in actual spaceflight, suggesting that ground-based experiments may need to be carried out longer to
replicate the effects observed in space. The Mars500 mission and 180-day CELSS experiments revealed
little effect of time on the degree of methylation changes, which might be attributed to the ability of the
human body to recover to some extent during the isolation experiment. Conversely, the effect of time in
spaceflight suggests that DNA methylation changes in these gene sets may not be recoverable within the
space environment. Understanding the molecular mechanisms of dynamic space adaptation necessitates
the integration of multi-omics signals from multiple individuals and missions. Nevertheless, our study
reveals that DNA methylation plays an important role in space adaptation and recovery, providing new
insights in space life science and directions for measures to mitigate the health risks associated with
spaceflight.

Limitations of the study

Considering the limitations of actual spaceflight conditions, our study inevitably has several limitations.
First, proteomics and acetylation modification analyses were performed only for M90 and M180-1
because of the limited number of materials available. Further proteomics analyses are planned for future
spaceflights to investigate the roles of proteins and their modifications in human space adaptation.
Second, DNA methylation and the transcriptome were analyzed via bulk sequencing; thus, the potential
influence of changes in cell composition on these observations could not be eliminated. In the future, cell
sorting or single-cell sequencing could be applied to evaluate the cell composition at the single-cell level.
Finally, as a common limitation of spaceflight studies, few subjects are available. However, as
spaceflights and space exploration are becoming more frequent, and more accumulated data and
inspiring insights will allow space life science to thrive.

RESOURCE AVAILABILITY

Lead contact

- Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Yinghui Li (yinghuidd@vip.sina.com).

Materials availability

This study did not generate new unique reagents. All the reagents in this study were included in the key resources table.

Data and code availability

- All omics data have been deposited at GSA, GSA-Human and OMIX (The Genome Sequence Archive (GSA), https://ngdc.cncb.ac.cn/gsa/; GSA-Human, https://ngdc.cncb.ac.cn/gsa-human/; OMIX, https://ngdc.cncb.ac.cn/omix/). Identifiers of the datasets are documented in https://www.spacelifescience.cn/datasets. They are available upon request if access is granted. To request access, contact the corresponding author. In addition, processed datasets derived from these data have been deposited at LSOS (https://www.spacelifescience.cn) and are publicly available as of the date of publication.

- All original code has been deposited at https://github.com/lijian2014seu/LSOS and is publicly available as of the date of publication.

- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

This work was supported by the CSS Program (HYZHXMH01001, HYZHXM01004), the National Key R&D Program of China (2022YFA1104302, 2022YFA1604503), grants from the National Key Laboratory of Space Medicine, China Astronaut Research and Training Center (SMFA17B07, SMFA19A03, SMFA19B01, SMFA19C01, SMFA20C03), the National Natural Science Foundation of China (32270607, 31900473, 31800707), the Young Elite Scientists Sponsorship Program by CAST (2018QNRC001), and State Key Laboratory of Space Medicine, China Astronaut Research and Training Center (SKL2024K03, SKL2024Y04, SKL2024Y07). The authors would like to thank the astronauts for their support of this work.

AUTHOR CONTRIBUTIONS

Conceptualization, project administration and supervision: Y.H.L., J.L., and L.N.Q.; Investigation: L.L., Y.Y.H, X.L., K.L., T.Z.Z., F.J.L., L.H.C., L.Q.W., X.Y.M., S.D.F., K.L., Y.H.Y., Z.Q.D., D.L., H.Y.Z., C.Y., A.A.L., L.J.W., Z.L.L., S.J.L., X.Q.D., C.J.Y., C.Y.W., P.S., L.J.S., C.Z., J.H.X., M.W., C.X., Z.X.L., Q.L.N., J.L., and H.Y.L.; Writing – original draft: Y.Y.H.; Writing – review and editing: J.L., X.L., L.L., Y.H.L., and L.N.Q. Funding acquisition: Y.H.L., J.L., L.N.Q., L.L., K.L., Y.H.Y, and X.L.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

The authors did not use any generative AI and AI-assisted technologies in this study.

SUPPLEMENTAL INFORMATION

Document S1. Tables S1, FigureS1-S7, supplemental methods, supplemental notes, and supplemental references.

Table S2. Results of quantitative and functional enrichment analysis of DMRs.

Table S3. Gene differential expression analysis results.

Table S4. Differentially spliced genes.

Table S5. Differential expression analysis results of peptides and protein acetylation modification changes in mission M90 and M180-1.

Table S6. PPI network.

Table S7. HLU model rat.

480
481

Figure 1. Presentation of the experimental design and data in this study
We present DNA methylation and physiological (phenotypic) data from three cohorts of astronauts in the
China Manned Space Program (CMSP), and transcriptomic, DNA methylation, and physiological
(phenotypic) data from three cohorts of astronauts in the CSS. In addition, we performed ground-based
simulations, including microgravity-stimulated head-down bed rest (HDBR) and rat hindlimb unloading
(HLU), and included data from the 180-day CELSS isolation simulation and the Mars500 isolation
stimulation (520 days), to elucidate space environmental stressors.

Figure 2. DNA methylation changes in the six spaceflights
a, X-axis shows the six spaceflight times ranging from 13 to 180 days. The Y-axis shows the number of DMPs (T2 vs. T1), with increased methylation DMPs presented and decreased DMPs. **b,** Global changes in DNA methylation across six missions. The increased and decreased DMPs are shown above and below, respectively. The DMPs from M13, M15, M33, M90, M180-2 and M180-1 are plotted from outside to inside. Marks on the outermost circle represents the DMP enrichment region that DMPs keeps increasing following the mission duration increases, as consistent regions across all missions.

Figure 3. DNA methylation and transcriptome changes after spaceflight and recovery, and the correlation between DNA methylation and gene expression

a, The top panel shows the numbers of differentially methylated positions (DMPs) after spaceflight (T2 vs. T1) and after recovery (T3 vs. T2) in the 90-day spaceflight group (M90). The bottom panel shows the same data for the 180-day spaceflight group (M180-1). **b,** Comparison of the gene expression of demethylases (top) and methyltransferases (bottom) at three timepoints (T1, T2 and T3) in M90 and M180-1. The adjusted *P* values calculated by limma are listed. **c,** The upper panel shows the number of differentially expressed genes (DEGs) after spaceflight (T2 vs. T1) and after recovery (T3 vs. T2) from the M90 group. The bottom panel shows the same data from the M180-1 group. **d,** The number of overlaps of the DEGs between the two flights (left: T2 vs. T1, right: T3 vs. T2). The dots connected below the bars represent intersections, while the separate dots represent genes unique to each group. **e,** Correlation coefficients between the log$_2$FCs of the expression level and the log$_2$FCs of the DNA methylation level in the promoter and enhancer regions after spaceflight (T2 vs. T1) calculated for all expressed genes, all DEGs, and unchanged genes for the two spaceflight missions.

Figure 4. Alternative splicing events during adaptation to and recovery from spaceflight and their associations with gene body DNA methylation.
a, The number of differential AS events between three timepoints of the two spaceflights (M90 and M180-1). **b,** The number of DMPs occurring within genes with significant AS events, categorized and counted according to the regions of the gene, on which the probes are located. A hypergeometric test was performed to evaluate the significance of enrichment of the loci in the gene body regions. **c, d,** Similar to **b** but for the number of hypomethylated and hypermethylated probes, respectively. TSS, transcriptional start site; TSS200, 0–200 bases upstream of the TSS; TSS1500, 200–1500 bases upstream of the TSS; 5'UTR, within the 5' untranslated region, between the TSS and the ATG start site; Body, between the ATG and stop codon; irrespective of the presence of introns, exons, TSS, or promoters; 3'UTR, between the stop codon and poly A signal; ExonBnd, within 20 bases of an exon boundary (i.e., the start or end of an exon); IGR, intergenic regions.

Figure 5. The expression of the overrange rebound genes during recovery and the effects of DNA methylation

a, The expression profiles of all genes of all astronauts in M90 and M180-1 are shown, with genes grouped by the types of change in the T2 vs. T1 and T3 vs. T2 comparisons, e.g., "down_up" indicates downregulated in T2 vs. T1 and upregulated in T3 vs. T2, whereas "stable" indicates no significant change. **b,** Quantification of the overrange rebound of gene expression during recovery by calculating the log_2FCs comparing T3 and T1 for the DEGs in M90 (top) and M180-1 (bottom). **c,** Dynamic changes in the expression levels of DEGs from T2 vs. T1 postflight (T2) and recovery (T3) in M90 (left) and M180-1 (right). **d,** Correlation coefficients between the log_2FCs of the expression level and the log_2FCs of the DNA methylation level in the promoter and enhancer regions (T3 vs. T2) calculated for all DEGs, unrecovered DEGs, over-range rebound DEGs and other DEGs in M90 and M180-1. **e,** The node degrees of various gene groups within the PPI network (Wilcoxon test, two-tailed), including DEGs

549   (categorized as unrecovered DEGs, overrange rebound DEGs and other DEGs) and unchanged genes,
550   in M90 (left) and M180-1 (right).

Figure 6. Functional enrichment results for the genes included in various modules and the PPI network constructed from the brown module.
a, Significance of the correlation between the gene modules identified by WGCNA and time. Correlations were calculated separately in both spaceflights and jointly by ANOVA between module scores and sampling times. **b, c,** GO terms enriched among blue (**b**) and brown module (**c**) genes after clustering; keywords in different categories are displayed in descending order of occurrence. **d,** Interactions between gene sets enriched among the PPI network genes. Edges represent significant interactions, with darker to lighter colors corresponding to the intensity of interactions, and dots represent gene sets, which have been classified by function and marked with different colors.

Figure 7. Comparison of DNA methylation changes in the gene sets associated with space
environment adaptation in spaceflight and ground-based experiments.
a-e, DNA methylation alterations of the PPI network genes enriched in the "osteoclast differentiation",
"NF-κB signaling pathway", "blood coagulation", and "platelet activation" gene sets in M90 (**a**), M180-1
(**b**), the 90-day HDBR (**c**), the 180-day CELSS (**d**), and Mars500 (**e**).

Figure S1. Global landscape of DNA methylation alterations in M90 and M180-1
a, Dynamic changes in the methylation levels of DMPs in T2 vs. T1 postflight and recovery in M90 (left)
and M180-1 (right). **b,** Proportion of demethylated, hypermethylated DMPs distributed in promoter and
enhancer regions compared with total demethylated, hypermethylated DMPs in each experiment.

Figure S2. Alterations in DNA methylation during M180-2 spaceflight
a, Number of DMPs postflight (T2 vs. T1) and during recovery (T3 vs. T2) in M180-2. **b,** Dynamic changes
in the methylation levels of DMPs (T2 vs. T1) postflight and during recovery (T3 vs. T2) in M180-2.

Figure S3. Genomic distributions of DMRs in M90, M180-1, and M180-2

a, Distribution of DMRs in the genome of M90 and M180-1 in T2 vs. T1. The y-axis represents the DMR values, with the red line indicating the zero-value threshold. The blue points correspond to hypomethylated DMRs, and the orange points represent hypermethylated DMRs. Distribution of DMRs present in both M90 and M180-1 (T2 vs. T1). This includes DMRs from M180-1 that overlap with those from M90 and DMRs from M90 that overlap with those from M180-1. Distribution of DMRs that are uniquely present in either the M90 or M180-1 T2 vs. T1. Number of overlapping and unique DMRs in M90 and M180-1 (T2 vs. T1). **b,** Distribution of DMRs in the genome of M90 and M180-1 in T3 vs. T2. **c,** Distribution of DMRs in the genome of M180-2 in T2 vs. T1 and T3 vs. T2, and overlap with DMRs in M90 and M180-1. The number of overlapping and unique DMRs between M180-2 and the other groups (M90, M180-1) is shown for both time comparisons.

Figure S4. Alterations in gene expression and the correlation between DNA methylation and gene expression in an independent 180-day spaceflight (M180-2)

a, The expression trends of all genes in spaceflight M180-2. Genes were grouped by the type of change in the T2 vs. T1 and T3 vs. T2 comparisons, e.g., "down_up" indicates downregulated in T2 vs. T1 and upregulated in T3 vs. T2, whereas "stable" indicates no significant change. **b,** Number of DEGs in T2 vs. T1 and the changes in these DEGs after recovery (T3 vs. T2) in M180-2. **c,** Dynamic changes in the expression levels of DEGs from T2 vs. T1 postflight (T2) and recovery (T3) in M180-2. **d,** The correlation coefficient of log$_2$FCs of gene expression and log$_2$FCs of promoter and enhancer area methylation in T2 vs. T1 was calculated for all expressed genes, all DEGs, and other unchanged genes.

614 **Figure S5. Changes in protein acetylation modifications in M90 and M180-1**
615 **a,** The number of proteins with acetylation modifications in samples from the M90 and M180-1
616 spaceflights. **b,** Differentially acetylated proteins in M90 and M180-1. Each line represents a protein and
617 the vertical axis corresponds to the number of acetylation sites of that protein. The source data for Figure
618 S5 are presented in Table S5.

28-day HLU model

Control

622 **Figure S6. Reconstructed microCT images of bone morphology in 28-day HLU model rats and**
623 **control rats**
624 Representative reconstructed microCT images of the longitudinal section of distal femurs collected from
625 the groups of 28-day HLU model rats (**a**) and control rats (**c**). Images depicted cross sections of the same
626 femurs from the HLU rats (**b**) and control rats (**d**), respectively.

Figure S7. Global alterations in methylation and phenotype alterations in ground-based simulation experiments

a-c, Demonstration of dimensionality reduction by tSNE before and after removal of batch effects for the 90-day HDBR (**a**), 180-day CELSS (**b**) and Mars500 (**c**). Annotations show the sample's source (subject) and the time of sampling (time).

d-e, Bone and coagulation related phenotype alterations in the 90-day HDBR. (**d**)**,** Bone related phenotypes includes Bone Mineral Density (BMD) and Z-score of hip bone, tibia, and calcaneus. The prefix "Pre" of the measurement timepoint represents how many days before HDBR, "R" represents the number of days the experiment is conducted, and "Post" represents how many days after HDBR. (**e**), Coagulation related phenotypes includes ADP-induced platelet aggregation rate, AA-induced platelet aggregation rate, PAC-1 expression rate, CD62P expression rate, TAT Complex, MPV, and PLCR.

(**f**), Coagulation related phenotype alterations in the 180-day CELSS. Coagulation related phenotypes includes PAC-1 expression rate and CD62P expression rate. The prefix "Pre" of the measurement time point represents how many days before the experiment, "R" represents the number of days the experiment is conducted, and "Post" represents how many days after the experiment.

REFERENCES

1. Mason, C.E., Green, J., Adamopoulos, K.I., Afshin, E.E., Baechle, J.J., Basner, M., Bailey, S.M., Bielski, L., Borg, J., Borg, J., et al. (2024). A second space age spanning omics, platforms and medicine across orbits. Nature *632*, 995-1008. 10.1038/s41586-024-07586-8.

2. Jones, C.W., Overbey, E.G., Lacombe, J., Ecker, A.J., Meydan, C., Ryon, K., Tierney, B., Damle, N., MacKay, M., Afshin, E.E., et al. (2024). Molecular and physiological changes in the SpaceX Inspiration4 civilian crew. Nature *632*, 1155-1164. 10.1038/s41586-024-07648-x.

3. Overbey, E.G., Kim, J., Tierney, B.T., Park, J., Houerbi, N., Lucaci, A.G., Garcia Medina, S., Damle, N., Najjar, D., Grigorev, K., et al. (2024). The Space Omics and Medical Atlas (SOMA) and international astronaut biobank. Nature *632*, 1145-1154. 10.1038/s41586-024-07639-y.

4. Afshinnekoo, E., Scott, R.T., MacKay, M.J., Pariset, E., Cekanaviciute, E., Barker, R., Gilroy, S., Hassane, D., Smith, S.M., Zwart, S.R., et al. (2020). Fundamental Biological Features of Spaceflight: Advancing the Field to Enable Deep-Space Exploration. Cell *183*, 1162-1184. 10.1016/j.cell.2020.10.050.

5. Gertz, M.L., Chin, C.R., Tomoiaga, D., MacKay, M., Chang, C., Butler, D., Afshinnekoo, E., Bezdan, D., Schmidt, M.A., Mozsary, C., et al. (2020). Multi-omic, Single-Cell, and Biochemical Profiles of Astronauts Guide Pharmacological Strategies for Returning to Gravity. Cell Rep *33*, 108429. 10.1016/j.celrep.2020.108429.

6. da Silveira, W.A., Fazelinia, H., Rosenthal, S.B., Laiakis, E.C., Kim, M.S., Meydan, C., Kidane, Y., Rathi, K.S., Smith, S.M., Stear, B., et al. (2020). Comprehensive Multi-omics Analysis Reveals Mitochondrial Stress as a Central Biological Hub for Spaceflight Impact. Cell *183*, 1185-1201.e1120. 10.1016/j.cell.2020.11.002.

7. Jirak, P., Mirna, M., Rezar, R., Motloch, L.J., Lichtenauer, M., Jordan, J., Binneboessel, S., Tank, J., Limper, U., and Jung, C. (2022). How spaceflight challenges human cardiovascular health. Eur J Prev Cardiol *29*, 1399-1411. 10.1093/eurjpc/zwac029.

8. Thirsk, R., Kuipers, A., Mukai, C., and Williams, D. (2009). The space-flight environment: the International Space Station and beyond. Cmaj *180*, 1216-1220. 10.1503/cmaj.081125.

9. Garrett-Bakelman, F.E., Darshi, M., Green, S.J., Gur, R.C., Lin, L., Macias, B.R., McKenna, M.J., Meydan, C., Mishra, T., Nasrini, J., et al. (2019). The NASA Twins Study: A multidimensional analysis of a year-long human spaceflight. Science *364*. 10.1126/science.aau8650.

10. Ball, P. (2003). China's first astronaut launched. Nature. 10.1038/news031013-5.

11. Gu, Y. (2022). The China Space Station: a new opportunity for space science. Natl Sci Rev *9*, nwab219. 10.1093/nsr/nwab219.

12. Human performance in space: Advancing astronautics research in China. (2014). Science *345*, 1522-1522. doi:10.1126/science.345.6203.1522-d.

13. Yuan, M., Custaud, M.A., Xu, Z., Wang, J., Yuan, M., Tafforin, C., Treffel, L., Arbeille, P., Nicolas, M., Gharib, C., et al. (2019). Multi-System Adaptation to Confinement During the 180-Day Controlled Ecological Life Support System (CELSS) Experiment. Front Physiol *10*, 575. 10.3389/fphys.2019.00575.

14. Liang, F., Lv, K., Wang, Y., Yuan, Y., Lu, L., Feng, Q., Jing, X., Wang, H., Liu, C., Rayner, S., et al. (2019). Personalized Epigenome Remodeling Under Biochemical and Psychological Changes During Long-Term Isolation Environment. Front Physiol *10*, 932. 10.3389/fphys.2019.00932.

15. Shi, J., Xu, J., Chen, Y.E., Li, J.S., Cui, Y., Shen, L., Li, J.J., and Li, W. (2021). The concurrence of DNA methylation and demethylation is associated with transcription regulation. Nat Commun *12*, 5285. 10.1038/s41467-021-25521-7.

16. Wright, C.J., Smith, C.W.J., and Jiggins, C.D. (2022). Alternative splicing as a source of phenotypic diversity. Nat Rev Genet *23*, 697-710. 10.1038/s41576-022-00514-4.

17. Matyash, A., Singh, N., Hanes, S.D., Urlaub, H., and Jäckle, H. (2009). SAP18 promotes Krüppel-dependent transcriptional repression by enhancer-specific histone deacetylation. J Biol Chem *284*, 3012-3020. 10.1074/jbc.M806163200.

18. Mishima, Y., Wang, C., Miyagi, S., Saraya, A., Hosokawa, H., Mochizuki-Kashio, M., Nakajima-Takagi, Y., Koide, S., Negishi, M., Sashida, G., et al. (2014). Histone acetylation mediated by Brd1 is crucial for Cd8 gene activation during early thymocyte development. Nat Commun *5*, 5872. 10.1038/ncomms6872.

19. Han, J., Lachance, C., Ricketts, M.D., McCullough, C.E., Gerace, M., Black, B.E., Côté, J., and

Marmorstein, R. (2018). The scaffolding protein JADE1 physically links the acetyltransferase
subunit HBO1 with its histone H3-H4 substrate. J Biol Chem *293*, 4498-4509.
10.1074/jbc.RA117.000677.
20. Jiao, Y., Shi, C., Edil, B.H., de Wilde, R.F., Klimstra, D.S., Maitra, A., Schulick, R.D., Tang, L.H.,
Wolfgang, C.L., Choti, M.A., et al. (2011). DAXX/ATRX, MEN1, and mTOR pathway genes are
frequently altered in pancreatic neuroendocrine tumors. Science *331*, 1199-1203.
10.1126/science.1200609.
21. Shayevitch, R., Askayo, D., Keydar, I., and Ast, G. (2018). The importance of DNA methylation of
exons on alternative splicing. Rna *24*, 1351-1362. 10.1261/rna.064865.117.
22. Lev Maor, G., Yearim, A., and Ast, G. (2015). The alternative role of DNA methylation in splicing
regulation. Trends Genet *31*, 274-280. 10.1016/j.tig.2015.03.002.
23. Williams, A., Bissinger, R., Shamaa, H., Patel, S., Bourne, L., Artunc, F., and Qadri, S.M. (2023).
Pathophysiology of Red Blood Cell Dysfunction in Diabetes and Its Complications.
Pathophysiology *30*, 327-345. 10.3390/pathophysiology30030026.
24. Hatherley, D., Lea, S.M., Johnson, S., and Barclay, A.N. (2013). Structures of CD200/CD200
receptor family and implications for topology, regulation, and evolution. Structure *21*, 820-832.
10.1016/j.str.2013.03.008.
25. Chen, X., Zhou, Z., Huang, C., Zhou, Z., Kang, S., Huang, Z., Jiang, G., Hong, Z., Chen, Q., Yang,
M., et al. (2021). Crystal Structures of Bat and Human Coronavirus ORF8 Protein Ig-Like Domain
Provide Insights Into the Diversity of Immune Responses. Front Immunol *12*, 807134.
10.3389/fimmu.2021.807134.
26. Crisford, H., Sapey, E., and Stockley, R.A. (2018). Proteinase 3; a potential target in chronic
obstructive pulmonary disease and other chronic inflammatory diseases. Respir Res *19*, 180.
10.1186/s12931-018-0883-z.
27. Huynh, K. (2019). Targeting ATP citrate lyase to reduce LDL cholesterol. Nat Rev Cardiol *16*, 385.
10.1038/s41569-019-0201-6.
28. Milazzo, G., Mercatelli, D., Di Muzio, G., Triboli, L., De Rosa, P., Perini, G., and Giorgi, F.M.
(2020). Histone Deacetylases (HDACs): Evolution, Specificity, Role in Transcriptional Complexes,
and Pharmacological Actionability. Genes (Basel) *11*. 10.3390/genes11050556.
29. de Vries, T.J., El Bakkali, I., Kamradt, T., Schett, G., Jansen, I.D.C., and D'Amelio, P. (2019).
What Are the Peripheral Blood Determinants for Increased Osteoclast Formation in the Various
Inflammatory Diseases Associated With Bone Loss? Front Immunol *10*, 505.
10.3389/fimmu.2019.00505.
30. Inada, M., Matsumoto, C., Uematsu, S., Akira, S., and Miyaura, C. (2006). Membrane-bound
prostaglandin E synthase-1-mediated prostaglandin E2 production by osteoblast plays a critical
role in lipopolysaccharide-induced bone loss associated with inflammation. J Immunol *177*, 1879-
1885. 10.4049/jimmunol.177.3.1879.
31. Andreev, D., Liu, M., Weidner, D., Kachler, K., Faas, M., Grüneboom, A., Schlötzer-Schrehardt, U.,
Muñoz, L.E., Steffen, U., Grötsch, B., et al. (2020). Osteocyte necrosis triggers osteoclast-
mediated bone loss through macrophage-inducible C-type lectin. J Clin Invest *130*, 4811-4830.
10.1172/jci134214.
32. Liao, R., Feng, Z., Li, W., Liu, R., Xu, X., Yao, S., and Tian, J. (2021). Interleukin-1 induces
receptor activator of nuclear factor-κB ligand-independent osteoclast differentiation in RAW264.7
cells. Exp Ther Med *21*, 640. 10.3892/etm.2021.10072.
33. Abu-Amer, Y. (2013). NF-κB signaling and bone resorption. Osteoporos Int *24*, 2377-2386.
10.1007/s00198-013-2313-x.
34. Hundalani, S.G., Nguyen, K.T., Soundar, E., Kostousov, V., Bomgaars, L., Moise, A., Hui, S.K.,
and Teruya, J. (2014). Age-based difference in activation markers of coagulation and fibrinolysis
in extracorporeal membrane oxygenation. Pediatr Crit Care Med *15*, e198-205.
10.1097/pcc.0000000000000107.
35. Blair, T.A., and Frelinger, A.L., 3rd (2020). Platelet surface marker analysis by mass cytometry.
Platelets *31*, 633-640. 10.1080/09537104.2019.1668549.
36. George, R., Bhatt, A., Narayani, J., Thulaseedharan, J.V., Sivadasanpillai, H., and Tharakan, J.A.
(2016). Enhanced P-selectin expression on platelet-a marker of platelet activation, in young
patients with angiographically proven coronary artery disease. Mol Cell Biochem *419*, 125-133.
10.1007/s11010-016-2756-4.

37.   Sang, Y., Roest, M., de Laat, B., de Groot, P.G., and Huskens, D. (2021). Interplay between
758   platelets and coagulation. Blood Rev *46*, 100733. 10.1016/j.blre.2020.100733.
759   38.   Greinacher, A., and Warkentin, T.E. (2023). Platelet factor 4 triggers thrombo-inflammation by
760   bridging innate and adaptive immunity. Int J Lab Hematol *45 Suppl 2*, 11-22. 10.1111/ijlh.14075.
761   39.   Peyvandi, F., Garagiola, I., and Baronciani, L. (2011). Role of von Willebrand factor in the
762   haemostasis. Blood Transfus *9 Suppl 2*, s3-8. 10.2450/2011.002s.
763   40.   Li, L., He, Z.Y., Wang, Y.Z., Liu, X., and Yuan, L.Y. (2018). Associations between thromboxane A
764   synthase 1 gene polymorphisms and the risk of ischemic stroke in a Chinese Han population.
765   Neural Regen Res *13*, 463-469. 10.4103/1673-5374.228729.
766   41.   Nieswandt, B., Varga-Szabo, D., and Elvers, M. (2009). Integrins in platelet activation. J Thromb
767   Haemost *7 Suppl 1*, 206-209. 10.1111/j.1538-7836.2009.03370.x.
768   42.   Nakahata, N. (2008). Thromboxane A2: physiology/pathophysiology, cellular signal transduction
769   and pharmacology. Pharmacol Ther *118*, 18-35. 10.1016/j.pharmthera.2008.01.001.
770   43.   Levi, M., and van der Poll, T. (2010). Inflammation and coagulation. Crit Care Med *38*, S26-34.
771   10.1097/CCM.0b013e3181c98d21.
772   44.   Huilcaman, R., Venturini, W., Fuenzalida, L., Cayo, A., Segovia, R., Valenzuela, C., Brown, N.,
773   and Moore-Carrasco, R. (2022). Platelets, a Key Cell in Inflammation and Atherosclerosis
774   Progression. Cells *11*. 10.3390/cells11061014.
775   45.   Gross, S., Tilly, P., Hentsch, D., Vonesch, J.L., and Fabre, J.E. (2007). Vascular wall-produced
776   prostaglandin E2 exacerbates arterial thrombosis and atherothrombosis through platelet EP3
777   receptors. J Exp Med *204*, 311-320. 10.1084/jem.20061617.
778   46.   Dixon, D.A., Tolley, N.D., Bemis-Standoli, K., Martinez, M.L., Weyrich, A.S., Morrow, J.D.,
779   Prescott, S.M., and Zimmerman, G.A. (2006). Expression of COX-2 in platelet-monocyte
780   interactions occurs via combinatorial regulation involving adhesion and cytokine signaling. J Clin
781   Invest *116*, 2727-2738. 10.1172/jci27209.
782   47.   Chao, -.Y., Zihan, -.X., Kai, -.L., Hongyu, -.Z., Hailong, -.W., Feng, -.W., Zi, -.X., Zhongquan, -.D.,
783   and Yinghui, -.L. (2021). - Effects of 180-day's Isolation on Bone, Glycolipid Metabolism and Their
784   Correlation Analysis. - Chinese Journal of Space Science - *41*, - 293. - 10.11728/cjss2021.02.293.
785   48.   Novikov, V., Oganov, V., Kabitskaya, O., Murashko, L., Naidina, V., and Chernikhova, E. (2018).
786   Mineral Bone Density and Body Composition of Participants in Experiment Mars-500. Human
787   Physiology *44*, 815-818. 10.1134/S0362119718070101.
788   49.   Haider, T., Gunga, H.C., Matteucci-Gothe, R., Sottara, E., Griesmacher, A., Belavý, D.L.,
789   Felsenberg, D., Werner, A., and Schobersberger, W. (2013). Effects of long-term head-down-tilt
790   bed rest and different training regimes on the coagulation system of healthy men. Physiol Rep *1*,
791   e00135. 10.1002/phy2.135.
792   50.   Kuzichkin, D.S., Markin, A.A., Vorontsov, A.L., Morukov, B.V., Zhuravleva, O.A., Zabolotskaya, I.V.,
793   and Vostrikova, L.V. (2016). Comprehensive assessment of the hemostasis system in the
794   participants of the Mars-500 experiment. Human Physiology *42*, 782-787.
795   10.1134/S0362119716070082.
796   51.   Andrieux, P., Chevillard, C., Cunha-Neto, E., and Nunes, J.P.S. (2021). Mitochondria as a Cellular
797   Hub in Infection and Inflammation. Int J Mol Sci *22*. 10.3390/ijms222111338.
798   52.   Medzhitov, R. (2021). The spectrum of inflammatory responses. Science *374*, 1070-1075.
799   10.1126/science.abi5200.
800   53.   Feuerecker, M., Crucian, B.E., Quintens, R., Buchheim, J.I., Salam, A.P., Rybka, A., Moreels, M.,
801   Strewe, C., Stowe, R., Mehta, S., et al. (2019). Immune sensitization during 1 year in the Antarctic
802   high-altitude Concordia Environment. Allergy *74*, 64-77. 10.1111/all.13545.
803   54.   Buchheim, J.I., Billaud, J.N., Feuerecker, M., Strewe, C., Dangoisse, C., Osterman, A., Mehta, S.,
804   Crucian, B., Schelling, G., and Choukér, A. (2020). Exploratory RNA-seq analysis in healthy
805   subjects reveals vulnerability to viral infections during a 12- month period of isolation and
806   confinement. Brain Behav Immun Health *9*, 100145. 10.1016/j.bbih.2020.100145.
807   55.   Ponomarev, S., Kalinin, S., Sadova, A., Rykova, M., Orlova, K., and Crucian, B. (2021).
808   Immunological Aspects of Isolation and Confinement. Front Immunol *12*, 697435.
809   10.3389/fimmu.2021.697435.
810   56.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
811   and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.
812   10.1093/bioinformatics/bts635.

813   57.   Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with
814   or without a reference genome. BMC Bioinformatics *12*, 323. 10.1186/1471-2105-12-323.
815   58.   Zhang, Y., Parmigiani, G., and Johnson, W.E. (2020). ComBat-seq: batch effect adjustment for
816   RNA-seq count data. NAR Genom Bioinform *2*, lqaa078. 10.1093/nargab/lqaa078.
817   59.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
818   powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
819   Acids Res *43*, e47. 10.1093/nar/gkv007.
820   60.   Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., Pimentel, H., Salzberg, S.L.,
821   Rinn, J.L., and Pachter, L. (2012). Differential gene and transcript expression analysis of RNA-
822   seq experiments with TopHat and Cufflinks. Nat Protoc *7*, 562-578. 10.1038/nprot.2012.016.
823   61.   Morris, T.J., Butcher, L.M., Feber, A., Teschendorff, A.E., Chakravarthy, A.R., Wojdacz, T.K., and
824   Beck, S. (2014). ChAMP: 450k Chip Analysis Methylation Pipeline. Bioinformatics *30*, 428-430.
825   10.1093/bioinformatics/btt684.
826   62.   Tian, Y., Morris, T.J., Webster, A.P., Yang, Z., Beck, S., Feber, A., and Teschendorff, A.E. (2017).
827   ChAMP: updated methylation analysis pipeline for Illumina BeadChips. Bioinformatics *33*, 3982-
828   3984. 10.1093/bioinformatics/btx513.
829   63.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al.
830   (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innovation
831   (Camb) *2*, 100141. 10.1016/j.xinn.2021.100141.
832   64.   Lizio, M., Harshbarger, J., Shimoji, H., Severin, J., Kasukawa, T., Sahin, S., Abugessaisa, I.,
833   Fukuda, S., Hori, F., Ishikawa-Kato, S., et al. (2015). Gateways to the FANTOM5 promoter level
834   mammalian expression atlas. Genome Biology *16*, 22. 10.1186/s13059-014-0560-6.
835   65.   Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network
836   analysis. BMC Bioinformatics *9*, 559. 10.1186/1471-2105-9-559.
837   66.   Gu, Z., and Hübschmann, D. (2023). simplifyEnrichment: A Bioconductor Package for Clustering
838   and Visualizing Functional Enrichment Results. Genomics Proteomics Bioinformatics *21*, 190-202.
839   10.1016/j.gpb.2022.04.008.
840
841

842 **STAR★METHODS**

843 **KEY RESOURCES TABLE**

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Bacterial and virus strains		
Biological samples		
Chemicals, peptides, and recombinant proteins		
Critical commercial assays		
Deposited data		
Transcriptomics data	This paper	HRA: HRA008486, HRA008487, HRA008488 (https://ngdc.cncb.ac.cn/gsa-human); OMIX: OMIX007304 (https://ngdc.cncb.ac.cn/omix); CRA: CRA018849, CRA018831 (https://ngdc.cncb.ac.cn/gsa)

DNA methylation data	This paper	OMIX: OMIX007289, OMIX007291, OMIX007300, OMIX007301, OMIX007290, OMIX008613, OMIX008614, OMIX008615, OMIX007302 (https://ngdc.cncb.ac.cn/omix)
Proteomics data	This paper	OMIX: OMIX007327, OMIX007328 (https://ngdc.cncb.ac.cn/omix)
Protein acetylation data	This paper	OMIX: OMIX007319, OMIX007320 (https://ngdc.cncb.ac.cn/omix)
Phenotypic data	This paper	OMIX: OMIX007309, OMIX007310, OMIX007312, OMIX007306, OMIX007307, OMIX007359, OMIX010025, OMIX010026, OMIX010027, OMIX007311 (https://ngdc.cncb.ac.cn/omix)
Experimental models: Cell lines		
Experimental models: Organisms/strains		
Oligonucleotides		
Recombinant DNA		
Software and algorithms		

R 4.2.1	http://www.r-project.org/	RRID:SCR_001905
edgeR	http://bioconductor.org/packages/edgeR/	RRID:SCR_012802
UpSetR	https://github.com/hms-dbmi/UpSetR	RRID:SCR_026112
ChAMP	http://www.bioconductor.org/packages/2.13/bioc/html/ChAMP.html	RRID:SCR_012891
SVA	http://www.omicsexpress.com/sva.php	RRID:SCR_002155
Nextflow	https://github.com/nextflow-io/nextflow	RRID:SCR_024135
FastQC	http://www.bioinformatics.babraham.ac.uk/projects/fastqc/	RRID:SCR_014583
MultiQC	http://multiqc.info/	RRID:SCR_014982
Trim Galore	http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/	RRID:SCR_011847
STAR	http://code.google.com/p/rna-star/	RRID:SCR_004463
RSEM	http://deweylab.biostat.wisc.edu/rsem/	RRID:SCR_000262
clusterProfiler	http://bioconductor.org/packages/release/bioc/html/clusterProfiler.html	RRID:SCR_016884
WGCNA	http://www.genetics.ucla.edu/labs/horvath/CoexpressionNetwork/	RRID:SCR_003302
Cytoscape	http://cytoscape.org	RRID:SCR_003032
Other		

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

METHOD DETAILS

Overview of the experiments and datasets

Here, we provide independent multi-omics data from multiple Chinese astronauts in one 13-day (M13), one 15-day (M15), one 33-day (M33), one 90-day (M90) and two 180-day spaceflights (M180-1 and M180-2) (Figure S1). For all the six spaceflights, DNA methylation and phenotypes were analyzed. For M90, M180-1 and M180-2, both transcriptome and DNA methylation were analyzed using peripheral blood cells, and the corresponding phenotypes were measured as well. For M180-1, proteome and protein acetylation were analyzed. Concurrently, we introduced and performed a variety of ground-based simulation experiments to estimate the effects of different space environment factors independently (Figure 1). The Mars500 (520-day isolation simulation[14]) and 180-day Controlled Ecological Life Support System (CELSS) [13] experiments were introduced to stimulate isolation in space. A 90-day head-down bed

rest (HDBR) experiment and 28-day (rat) and 35-day (rat) hind-limb unloading (HLU) model experiments
were conducted to simulate microgravity on Earth. Details of the ground-based simulation experiments
are described in the Supplementary Methods.

All animal experimental protocols in this study were performed in accordance with standard ethical
guidelines and approved by the China Astronaut Research and Training Center Animal Ethics Review
Committee (NO: ACC-IACUC-2021-015).

Participants
A total of seventeen Chinese astronauts participating in one 13-day (M13, three astronauts), one 15-day
(M15, three astronauts), one 33-day (M33, two astronauts), one 90-day (M90, three astronauts)
spaceflight and two 180-day spaceflights (M180-1, three astronauts, and M180-2, three astronauts) were
included. We calculated the statistics of the ages of all the seventeen astronauts from the six spaceflights.
The minimum value is 33.42 years, the 1st quarter value is 41.75 years, the median value is 45.67 years,
the mean value is 45.17 years, the 3rd quarter value is 48.75 years, the maximum value is 56.75 years,
the standard deviation is 6.73 years. There exists no outlier value among all the seventeen age values,
indicating consistent age range. The study subjects signed informed consent according to the declaration
of Helsinki and Article 1008 Civil Code of the People's Republic of China for the collection and use of
sample materials and data in research protocols at China Astronaut Research and Training Center and
the collaborating institutions. The study protocols were approved by the China Astronaut Research and
Training Center Animal Medical Ethics Review Committee.

Sample collection
For each of the six spaceflights, blood samples from each astronaut at three or four timepoints (Table S1)
were collected into 2 mL BD Vacutainer® EDTA Tubes (Cat # 367841) per the manufacturer's
recommendations. For the latest three spaceflights (M90, M180-1 and M180-2), additional blood samples
were collected into 2.5 mL BD PAXgene® Blood RNA Tubes (Cat # 762165) for transcriptome
sequencing. Sample collection was performed at 6:30 o'clock a.m. each time to avoid circadian rhythmic
effects. To minimize batch effects within each spaceflight, the blood samples were stored at -80 ℃ until all
the samples from the same spaceflight were fully prepared.

Multi-omics measurement
For multi-omics data acquisition, DNA methylation was profiled via the Illumina Infinium MethylationEPIC
BeadChip. Samples from the earlier spaceflight missions (M13, M15 and M33) were analyzed using 450K
BeadChip, whereas, later spaceflight missions (M90, M180-1 and M180-2) were analyzed using the 850K
BeadChip, reflecting the updates of the DNA methylation assay by Illumina. RNA sequencing was
conducted on an Illumina NovaSeq 6000 PE150 platform using the standard method. Proteomics and
protein acetylation were quantified on a Bruker timsTOF Pro mass spectrometer. The full details are
available in the Supplementary Information.

QUANTIFICATION AND STATISTICAL ANALYSIS

Expression quantification and differential expression analysis
To facilitate transcriptomic analysis, the following systematic pipeline was employed for the preprocessing
of raw RNA sequencing data, utilizing the human genome from Ensembl version 107 as the reference.
FastQC software (v0.12.1) (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) was used to
evaluate the quality of the original FASTQ files. Using parameters including --length 36, a quality
threshold of --quality 25, and a trimming stringency of --stringency 3, TrimGalore software (v0.6.10)
(https://github.com/FelixKrueger/TrimGalore/releases/tag/0.6.10) was used to trim and filter the initial
FASTQ files, efficiently eliminating low-quality nucleotides and adapter sequences. STAR (v2.7.10a)
(https://github.com/alexdobin/STAR/)[56] was modified to align the trimmed FASTQ files to the Ensembl
version 107 reference genome (hg38). The RSEM tool (v1.3.1) (https://github.com/deweylab/RSEM)[57]
was used to compute gene expression abundance according to matched BAM files to generate an
expression profile. The above analysis was performed on CentOS Linux version 7.2.1511. The Rtsne
package (v0.16) was used for dimensionality reduction and visualization. To mitigate the potential
influence of interindividual variability and batch effects, the combat method was employed. We used
ComBat_seq from the sva package (v3.46.0)[58] to remove the individual-specific characteristics from each
dataset by treating individual sources as batches to remove these environmentally irrelevant effects. After
batch effect removal, the two flight datasets were merged for further analysis. Differential expression

analysis was conducted using the limma package (v3.54.2)[59], and DEGs ($|\log_2FC| > \log2(1.5)$, and
adjusted P value < 0.05, DEGs) were identified across all groups. All of the R packages used in this study
were applied in R version 4.2.1 (https://cran.r-project.org/bin/windows/base/old/4.2.1/). The DEG
classification method is available in the Supplementary Methods.

Alternative splicing analysis
The clean RNA-seq data obtained via the above TrimGalore processing were subjected to alternative
splicing analysis using the TopHat Cufflinks pipeline[60]. The analysis was carried out with Bowtie2-2.5.1-
linux-x86_64 (https://sourceforge.net/projects/bowtie-bio/files/bowtie2/2.5.1/), TopHat v2.1.1
(https://ccb.jhu.edu/software/tophat/downloads/), and Cufflinks v2.2.1 (https://cole-trapnell-
lab.github.io/cufflinks/install/) on CentOS Linux release 7.2.1511. After quality control, the clean FASTQ
data were first aligned using TopHat to the reference genome
Homo_sapiens.GRCh38.dna_sm.primary_assembly.fa, which was downloaded from ENSEMBL, with
guidance from Homo_sapiens.GRCh38.107.gtf, which contains transcript information. The Cufflinks
package was subsequently used to assemble transcripts, compare the assemblies to annotations, merge
the assemblies and detect differential splicing and promoter use. The differentially expressed alternatively
spliced transcripts were counted and then enriched with STRING (https://cn.string-db.org/).

DNA methylation raw data processing
DNA methylation data from space- and human ground-based experiments were processed with ChAMP
and then visualized with the Rtsne package (0.16) in R version 4.2.1[61,62] for dimensionality reduction. As
in the transcriptomic data, there was a batch effect between the two flight datasets, with samples in each
flight grouped primarily according to individual sources. To remove the batch effects, we carried out a
procedure similar to that used for the transcriptome data, applying the champ.runCombat method in the
ChAMP package (2.28.0) (R version 4.2.1).

Differential methylation analysis
Differential methylation analysis was performed using ChAMP. Positions with P value < 0.05, |deltaBeta|
> 0.05, and a consistent change across all three subjects (either all demethylated or all hypermethylated)
were considered differentially methylated between the two groups and defined as differentially methylated
positions (DMPs). Differentially methylated regions (DMRs) were identified using the champ.DMR
function, employing the Bumphunter method with an adjusted P value threshold set to 0.05. DMRs with
containment relationships (including complete overlap) in the two flight experiments were identified by
comparing their positions on the chromosome. The functional enrichment of DMRs was performed on the
genes hosting the DMPs distributed in the region via the clusterProfiler package (v4.6.2)[63].

Methylation change trend analysis
We compared the trends in genomic methylation changes in spaceflight (M90 and M180-1) and ground-
based simulation experiments (90-day HDBR, 180-day CELSS, and Mars500) for a set of genes enriched
by nodes of the adaptation-related PPI network, that is, the functional genes of interest. To select
appropriate loci exhibiting consistent changes, the following criteria were employed: loci on the promoter
region (including "TSS1500", "TSS200", "5'UTR", and "1stExon") with a corrected P value < 0.05 as
determined by an ANOVA test and grouped by time, as well as loci exhibiting a range of fluctuating beta
values greater than 0.05. The DNA methylation levels of these selected loci were subsequently averaged
across a group of experimental subjects to facilitate visualization of the data. The visualization method
began with data normalization of the matrix of average methylation values for the loci, followed by the
construction of a graph depicting the trends in the methylation levels of the locus over time. The graphs
used filled regions from light to dark to represent the range of intervals of loci methylation levels,
illustrating changes in gene set methylation levels over time.

Correlation between DNA methylation and expression
The significance of the connection between promoter region methylation and expression was evaluated
by performing a Pearson correlation test on the $\log_2FC$ values of gene expression and the $\log_2FC$ values
(deltaBeta) of promoter/enhancer region DNA methylation (the median deltaBeta of loci on the
promoter/enhancer region). DMPs located in the promoter regions were automatically annotated by the
champ.DMP function. We identified DMPs in the enhancer regions through annotating against the
human-specific FANTOM5 enhancer database[64].

Proteomics and protein acetylation data processing

The data from the machine were identified using MaxQuant's (v.1.5.3.30) integrated Andromeda engine;
then, quantitative analysis was performed using MaxQuant on the basis of the peak intensity, peak area
and retention time of peptide segments related to primary mass spectrometry, and the targeted modified
peptide segments were extracted. We used the UniProt protein database and genome annotation-based
protein database for protein identification. The data obtained were filtered using the Andromeda engine
integrated in MaxQuant, filtering at the PSM-level with an FDR <= 1% at the spectrum level and further
filtering at the protein-level FDR <= 1% at the protein level. Additionally, for modification sites, filtering
was performed at the modification site level with an FDR of 1% for identification (site decoy fraction <=
1%) to obtain significant modification results. The modified peptides were counted for further differential
analysis.

Differential expression analysis of proteomics data
The differential expression analysis of expressed peptides was performed with limma (v3.54.2) in R
version 4.2.1. Quantile normalization was first applied to the data before linear regression. The
significance threshold for differentially expressed peptides was an adjusted P value <= 0.05.

Differential expression analysis of protein acetylation data
The differential fold changes in the modified peptide segments in each comparison group were
subsequently calculated on the basis of the defined comparison groups and statistically tested using
edgeR 3.40.2. The data were first quantile normalized with the voom function, and then a linear model
was fitted to the expression data for each gene. An adjusted P value of less than 0.05 was used as the
threshold for differentially modified peptides.

Construction of PPI networks related to environmental adaptation
Initially, the WGCNA package (v1.72-5)[65] was employed to partition genes into modules consisting of
closely associated genes. Subsequently, the correlation P values between all modules and time (T1, T2,
T3) were calculated. The brown and blue modules were identified on the basis of a P value threshold for
significance of 0.05. The ClusterProfiler (v4.6.2)[63] package was used to perform functional enrichment
analysis of the module genes. The simplifyGO function from the simplifyEnrichment[66] (v1.8.0) package
was employed to cluster and visualize the GO terms. The biological functions within each cluster were
summarized and presented as word clouds, which were then linked to a similarity heatmap displaying the
shared biological functions across the clusters. The importance of each keyword was determined using
Fisher's exact test, and their significance was represented by the size of the font used.

To investigate the interplay between different functional genes within the brown module, gene-gene
interactions were assessed by integrating the gene coexpression coefficients, which were calculated
using WGCNA, with the gene interaction scores from the STRING database. A threshold of WGCNA
weights > 0.125 (equivalent to the absolute value of the coexpression coefficient > 0.5 where power = 3)
and a STRING score > 0.4 were applied to identify interacting genes, among which 1727 genes were
identified. This step ultimately led to the construction of a predicted protein-protein interaction (PPI)
network, with a total of 12490 interactions and 1458 nodes.

Gene set interaction analysis
To calculate the interaction scores between geneset1 and geneset2, the following steps were taken. First,
sub_PPI_net, a PPI network subset, was extracted containing genes from both geneset1 and geneset2.
The calculation formula for the interaction score is as follows:

$$InteractionScore_{(GeneSet1,GeneSet2)} = \frac{\sum_{j}^{GeneSet2} \sum_{i}^{GeneSet1} (Weights_{WGCNA\ (i,j)} + Score_{STRING(i,j)})}{Length_{GeneSet1} \times Length_{GeneSet2}}$$

1005

To assess the significance of the interaction score, 100 random gene sets with the same size as
geneset1 and geneset2 were randomly generated, and the interaction scores were calculated for these
sets. The frequency of actual interaction scores greater than the scores obtained from the 100 random
sets was calculated as the P value for interaction significance. Gene set pairs with an interaction score P
less than 0.05 were considered to have significant interactions. The importance of each gene set was
determined by the sum of the interaction scores between the gene set and all other gene sets.