In form for a swarm: programmable neutrophil swarming impacts infection outcome

Salik M. Borbora[1][§], Ivanna Williantarra[1][§], Giulia Rinaldi[1], Can Cui[1], Echo Yongqi Luo[1], Hazel A. Walker[1], Holly M. Craven[1,2], Auxence Desrentes[1], Nir S. Gov[1,3], Milka Sarris[1][*]

Affiliations:

[1]University of Cambridge, Department of Physiology, Development and Neuroscience, Downing Site, Cambridge, CB2 3DY, UK.

[2]Roving Researcher Programme, School of Biological Sciences, University of Cambridge, United Kingdom

[3]Department of Chemical and Biological Physics, Weizmann Institute of Science, Rehovot, Israel

§ These authors contributed equally

*Correspondence to: ms543@cam.ac.uk

Abstract

Migrating cells often pattern the distribution of chemoattractants to support their accumulation in target tissues. Neutrophil swarming represents a prominent example of such self-organisation. During this process, neutrophils engage in coordinated autocrine/paracrine chemoattractant signalling that drives rapid clustering at sites of injury or infection. Whether neutrophil swarming can be regeared to influence immune outcomes is unclear. Here, we show that neutrophil swarming is subject to reprogramming after microbial experience and amenable to targeted genetic engineering with measurable effects on infection outcome. We established a model of bacterial wound infection in zebrafish, focusing on *P. aeruginosa*, an important opportunistic pathogen. We demonstrate that zebrafish larvae exposed to prior microbial experience (with non-pathogenic *E. coli*) are more resistant to subsequent wound infection by *P. aeruginosa,* in comparison to naive animals. Through live imaging and chemical perturbations, we show that these training effects are in part attributed to changes in neutrophil swarming and associated gene expression. Finally, we demonstrate that genetic enhancement of 5-lipoxygenase in neutrophils is sufficient to maximise neutrophil swarming and improve infection outcome. Together, these data suggest new routes for reprogramming neutrophil accumulation in tissues, via manipulating their ability to shape chemoattractant landscapes.

Introduction

Directed cell migration is essential for tissue development and surveillance. Guidance of migrating cells relies on interpretation of gradients of chemoattractants within tissues. Traditional mechanistic models portray migrating cells as receivers of information, responding to chemoattractant landscapes patterned by extrinsic tissue cells. However, several lines of evidence highlight an important role for self-organisation in the process, through cell-autonomous generation of chemical gradients by migratory cells[1–5]. For example, during development, migrating tissue collectives sculpt chemical gradients through asymmetric distribution of dedicated chemoattractant scavenging receptors[3,5]. In cancer metastasis, melanoma cells disperse via catabolism of ligand and local micro-gradient generation[6]. In chemotactic mazes, amoeboid cells detect dead ends using breakdown of chemoattractant[7]. During immune responses, leukocytes pattern autologous chemical gradients through chemoattractant scavenging or through autocrine/paracrine secretion[8–12]. A fundamental question raised by these observations is whether the migratory patterns of cells are intrinsically programmable, in a manner sufficient to impact disease at the tissue and organism scale.

To explore this question, we focus on the prominent paradigm of neutrophil swarming, as a self-organised migration behaviour. Neutrophils are first responders to tissue injury and infection and directly fight microbes through a variety of effector mechanisms, such as phagocytosis, secretion of antimicrobial molecules, and release of neutrophil extracellular traps (NETs)[13]. Neutrophils also produce chemokines and cytokines that influence the accumulation and function of other cells and ultimately shape the course of inflammation and disease. The recruitment of neutrophils is initially prompted by primary signals produced by tissue cells afflicted by injury or infection[13]. Thereafter, neutrophils begin to secrete further chemoattractants, generating tissue-scale gradients and/or waves of chemoattractants, which amplify the range of recruitment and determine the final magnitude of the response[8–11,14–16]. The process of neutrophil swarming is highly conserved and thought to impart functional benefits in fighting infection, particularly in the face of large and focal targets, such as wound infection or fungal invasions[9,17,18]. Specifically, swarms are characterized by rapid kinetics of recruitment and dense accumulation of neutrophils[8–11,14], which in principle would support prompt containment of pathogens, physical insulation of the pathogen-afflicted area and local concentration of antimicrobial molecules. On the other hand, neutrophil accumulation is also

known to prolong pathological inflammation when not properly regulated and resolved[13,19].

Therefore, the mechanisms and rules of organisation underlying neutrophil swarm

development have important functional implications in disease.

Several conserved mechanisms have been determined that drive the initiation and evolution of

neutrophil swarming. This includes the key neutrophil-derived chemoattractants driving swarm

development (notably, the eicosanoid leukotriene B4 (LTB4) and the chemokine Cxcl8[8,9,14,20]),

signal inducers of chemoattractant secretion (such as coordinated calcium signals[9,10,21]) as well

as factors that self-limit the evolution of swarms (such as receptor desensitization and reactive

oxygen species[10,22,23]). Additional contributing factors have also been described, such as

damage associated molecular patterns (for example ATP and fMLP[9,11]) and cellular behaviours,

such as neutrophil NETosis and target cloaking by other immune cells[20,24]. Despite the overall

conservation of neutrophil swarming and associated mechanisms, the dynamics of neutrophil

swarm behaviours within a given experimental system show significant heterogeneity, ranging

from small, short-lived clusters, to persistent and larger clusters[9,25]. Although some of this

heterogeneity can be attributed to variation in the intensity and composition of tissue-derived

signals[22,24], it remains unclear whether cell-intrinsic properties account for variation in

neutrophil propensity to swarm and whether these are subject to re-programming.

Immune cell gene expression and function can be re-programmed in at least two ways. The

first is through experience and 'training', whereby immune cells exposed to a primary insult

acquire epigenetic marks that poise their gene expression profile and responsiveness to a

subsequent insult[26–28]. Neutrophils have been shown to undergo such durable training after

exposure to primary infection, through epigenetic re-programming in relevant long-lived

progenitors[29,29–32]. Innate immune training increases resistance to infection in a variety of

tumour and infection settings and involves a range of epigenome and transcriptome

changes[29,30,33,34]. Whilst it is well-established that neutrophils enhance their killing capacity

and associated gene signatures after immune training[29,35–37], it remains unclear to what extent

their ability to swarm can also undergo re-programming under these conditions. A second route

to re-programming immune function is through genetic engineering, as exemplified in tumor

immunotherapies. Specifically, chimeric antigen receptors (CAR) can be engineered in

immune cells to programme their ability to recognise and kill certain tumours[38]. Whilst this has

been developed to the largest extent for T lymphocytes[38], new approaches for CAR therapies

are geared towards the manipulation of myeloid cells, including macrophages and

neutrophils[39–42]. However, the ability to genetically reprogramme the infiltration of immune

cells to target diseased tissues remains a challenging frontier. In this context, the manipulation

of self-generated gradients is a relatively unexplored but interesting route of manipulation, as

it offers the potential to modify the composition of inflammatory niches through the

reprogramming of ligands delivered by immune cells to the site.

Here we exploit the zebrafish as a tractable model to explore the link between cell behaviours

at the microscopic with disease outcome at the whole-organism scale. We developed and

characterised a model of wound infection in which neutrophil swarm heterogeneities can be

visualized in real time in individual animals, concomitant with monitoring of disease

progression and infection outcome. We first establish factors that determine susceptibility of

wound infection, notably the role of osmotic surveillance programmes in the epithelial barrier.

We then show that training challenge of zebrafish with non-pathogenic microbes enhances

resistance to subsequent wound infections and that, under these conditions, neutrophils exhibit

enhanced swarming prowess with reduced swarm heterogeneity across individuals. We found

that this correlates with upregulation of swarm-related gene expression signatures and that

inhibition of swarm machinery (LTB4 biosynthesis pathway) limits the protective effects of

prior microbial insult. Finally, we show that targeted genetic enhancement of swarm machinery

enhances neutrophil swarming and resistance to wound infection, indicating that neutrophil

swarm reprogramming is sufficient to influence infection outcome.

Results

Tissue damage detection through osmotic surveillance shapes susceptibility of zebrafish larvae to pathogenic wound infections

To investigate the dynamics and functional consequences of neutrophil swarms, we employed a wound infection model. *Pseudomonas aeruginosa* is an opportunistic bacterial pathogen, present in water and soil, able to colonise wounds in multiple species. In humans, wound infections can occur through contact of injured tissue with contaminated water or surfaces[43,44]. These infections can be severe and lead to sepsis and mortality, particularly in immunocompromised individuals or in the case of large surgery-associated infections[43,44]. We previously introduced a model whereby zebrafish larvae are subjected to wounding in the presence of aqueous medium contaminated with *P. aeruginosa* PAO1 strain[9]. This experimental model is advantageous in that it recapitulates a natural route of infection, in comparison to more common microinjection routes[45], and provides a relevant setting to study the role of neutrophil swarming in defence. However, these experimental infections are inherently more variable, since heterogeneities in the injured tissue can influence the titres of microbial colonisation and the ensuing immune response.

To understand and control sources of variation in this model, we examined known aspects of the tissue response to damage. Previous studies have shown that damage detection through osmotic surveillance is important for epithelial wound closure[46], inflammatory signal production and recruitment of neutrophils[47]. In addition, these tissue damage detection pathways were shown to promote optimal defence from microbes, when the latter are microinjected into hosts[48]. We therefore aimed to establish whether wound infections are influenced by osmotic surveillance pathways, with a view to minimising variation in infection.

To test the role of osmotic surveillance, we subjected zebrafish larvae to two types of wounding, mechanical and laser wounding **(Fig. 1A, G)**, in two common zebrafish buffers E3 medium (hypotonic), Ringer medium (isotonic), as well as isotonic E3 medium (with the same sodium levels and tonicity as Ringer). The use of two wounding assays offers complementary advantages. Whilst mechanical wounding can be performed on a large scale, to screen for disease phenotypes, laser wounding permits real-time visualisation of neutrophil responses

immediately upon wounding[9,49]. In both types of wound assays, we found that larvae in

hypotonic medium were largely resistant to wound infection, whereas larvae in isotonic

medium (iso E3, Ringer) were susceptible **(Fig. 1E, H)**. In all media conditions, non-wounded

larvae were resistant to infection, confirming the wound-dependent nature of infection **(Fig.**

1E, H). To characterise the disease progression in a more granular fashion, we created a disease

scoring matrix, with phenotypes ranging from alterations in swimming behaviour, to more

extensive morphological abnormalities **(Fig. 1B)**. Larvae infected in hypotonic media showed

minimal disease symptoms and bacterial burdens in comparison to larvae infected in isotonic

media (**Fig 1C, J**).

Wound closure can be delayed in isotonic media due to lack of tissue damage detection by the

ruptured epithelium[46]. We hypothesised that this may render wounds more permissive to

bacterial colonisation, contributing to the increased burden of infection in these conditions

(Fig. 1D, I). To test whether early wound colonisation might differ in the different media, we

measured bacterial burden shortly after wound infection (10 min and 30 min), at time points

shorter than the duplication time of *P. aeruginosa*[50]. We found that at this early stage the

bacterial titres were already significantly higher in isotonic conditions, suggesting either a

difference in bacterial colonisation or a difference in the early containment by immune cells

(Fig. 1F). To test direct effects of the medium composition on the growth of bacteria, we

examined the growth of PAO1 bacteria in hypotonic E3, isotonic E3, Ringer and PBS solutions.

We found comparable bacterial growth after 30 min exposure in the different media, suggesting

that the disease phenotypes of the larvae depended on the host response **(Fig. S1)**.

Together these results suggested a role for osmotic surveillance in wound defence and allowed

us to establish the optimal conditions for recapitulating wound infection and associated disease

in larvae.

Training through prior non-pathogenic microbial challenge alters neutrophil swarm-

related gene expression and enhances resistance to wound infections

We thereafter used the susceptible model of wound infection (isotonic conditions for

infection/Ringer solution) to explore whether we can reprogramme the innate immune response

to infection through training. To mimic vaccination challenge, we used superficial

microinjection as a route. We injected non-pathogenic, CFP-expressing *Escherichia coli*

MG1655[51] in the otic vesicle at 2 days post-fertilisation **(Fig. 2A)**. We found this pre-challenge increased the resistance to wound infection by *P. aeruginosa*, two days after the first challenge, as compared with controlled PBS-pre-challenged larvae **(Fig. 2B)**. Notably, in the absence of wounding, larvae were not susceptible to infection in either PBS *or E. coli*-pre-challenge, confirming again the opportunistic nature of infection **(Fig. S2A)**. Varying the dose of *E. coli* injection showed that a minimum dose was required to trigger protective effects **(Fig. S2B)**. One way of distinguishing innate immune 'training' from immune 'priming' is by assessing whether the protective effects persist after the immune response to the first challenge has passed its peak or resolved. To explore the nature of the protection observed, we determined the resolution kinetics of the first challenge, by measuring titres of *E. coli* and levels of inflammatory cytokines **(Fig 2E)**. Both markers indicated that the response to *E. coli* had largely subsided by the time the second challenge was given **(Fig 2F-J)**.

Neutrophil training is characterised by changes in transcriptomic and epigenetic signatures in differentiated neutrophils and progenitors[32]. Since the training challenge led to resistance of infection in a heterologous tissue site (tail versus ear), we hypothesised neutrophils from hematopoietic sites might be centrally altered. To explore this, we isolated neutrophils from the caudal hematopoietic tissue (CHT) **(Fig. 2K)** and found that CHT-derived neutrophils showed increased expression of genes associated with inflammation and cell migration, in particular *cxcl8a*, *5-lo* and cytoskeletal proteins, such as *tubg1* **(Fig 2L-R)**. This suggested that neutrophils may be centrally altered by training in a manner that changes their gene expression profile.

Together, these analyses revealed that innate immune responses to wound infections can be trained by prior microbial challenge and suggested that, among the many likely effects of training, neutrophil genes associated with cell migration (cytoskeletal proteins) or swarming (autologous chemoattractant ligands) can be subject to reprogramming.

Neutrophil swarming is enhanced through prior training challenge, and these phenotypic changes contribute to disease resistance

The changes in gene expression patterns in neutrophils upon training suggested that their migratory behaviour is altered. To relate neutrophil swarm patterns with disease outcome, we combined early imaging of wound infections (0-2 hours post infection-hpi) with monitoring of

disease outcome over a longer timescale (up to 24 hpi) in the same larvae **(Fig. 3A)**. We confirmed that in this laser-wound assay, PBS pre-challenged, control fish were more susceptible to PAO1 infection than *E. coli* pre-challenged fish **(Fig. S3A)**. We then compared the behaviour of neutrophils upon wounding and found that in PBS-pre-challenged, control animals, neutrophils exhibited both small-scale clustering and large-scale clustering **(Fig. 3B, Video S1)**. By contrast, in *E. coli*-pre-challenged animals, we observed a more consistent scale of clustering **(Fig. 3B, Video S2)**. To relate swarm variation to disease outcome, we plotted the scale of neutrophil recruitment (number of cells at 2 hpi) versus the bacterial burden (number of CFU at 24 hpi) in individual larvae. We confirmed that the initial number of neutrophils in the field of view in PBS versus *E. coli*-pre-challenged fish was not significantly different **(Fig. S3B)**. However, PBS-pre-challenged fish were characterised by greater variation in neutrophil recruitment, higher disease scores and bacterial counts in comparison to *E. coli*-challenged fish **(Fig. 3C)**. To understand these variations further, we stratified fish into four categories: i) weak neutrophil recruiters with high bacterial burden (Group 1, G1, in **Fig. 3C**), ii) strong recruiters with high bacterial burden (Group 2, G2, in **Fig. 3C**), iii) strong recruiters with weak bacterial burden (Group 3. G3, in **Fig. 3C**) and iv) weak recruiters with weak bacterial burden (Group 4, G4, in **Fig. 3C**). We considered as 'strong recruiters' those larvae in which the cell number exceeded 10 neutrophils. This was based on the distribution of the numbers seen in our experiments and on previous papers indicating a minimal number of clustering neutrophils to correlate with swarming kinetics of recruitment[15,16,52]. We considered as 'high burden', the lowest level of CFU count at which disease symptoms were observed across all experiments with these two groups of larvae. The number of larvae within G4 did not differ between PBS and *E. coli*-pre-challenged fish, and we reason this group may represent larvae in which bacterial colonisation was weak, as there was neither evidence for neutrophil recruitment nor bacterial burden. Most notably, we found that upon *E. coli* training there was a significant increase of larvae in G3 and a decrease in G1 **(Fig. 3D)**. This suggested that in a proportion of control larvae (G1), the swarm potential of neutrophils is sub-maximal and limits resistance to infection, whereas in *E. coli*-trained larvae, the swarm potential is maximised across different animals.

As an additional analysis, we measured the maximal rate of neutrophil recruitment across the 2 h imaging window. Here, we plotted the number of new neutrophils arriving at the wound over time and determined the slope of the curve during the phase of maximal recruitment **(Fig. S3C, D)**. We then plotted the maximal rate of recruitment against the bacterial burden for the

same embryo. Here, we observed a trend for higher rates of recruitment in *E. coli*-pre-
challenged larvae and lower burdens compared with PBS controls. We performed similar
stratification of larvae, considering 'fast recruiters' and 'slow recruiters' as those that had a
maximal rate of recruitment of either above or below the mean maximal rate, respectively,
across all larvae in both PBS and *E. coli*-pre-challenge groups. This analysis indicated a trend
for a higher probability of larvae to show fast maximal rate of recruitment and low bacterial
burden **(Fig. S3D)**, although the difference was not found to be significant with this sample
size **(Fig. S3E)**.
276
To investigate whether a change in neutrophil swarm potential contributes to wound infection
resistance, we used a specific inhibitor for 5-lipoxygenese (5-LO), zileuton, to target the
production of the swarm-driving chemoattractant, LTB4. When fish were wounded and
infected in the presence of zileuton, the *E. coli*-pre-challenged fish did not show as marked an
increase of infection resistance (over PBS-pre-challenged fish), in comparison to when control
vehicle (DMSO) was used **(Fig. 3E-H)**.
283
284 Together, these results indicated that innate immune training leads to changes in neutrophil
285 swarm patterns upon secondary pathogenic challenge, which in turn influence the probability
286 to resolve infection successfully.
287
288 **5-lipoxygenase enhancement in neutrophils is sufficient to enhance neutrophil swarming**
289 **and resistance from wound infection**
290
To determine whether reprogramming of neutrophil swarming is sufficient to influence wound
infections, we used genetic engineering of 5-LO in neutrophils. We hypothesised that
neutrophils expressing maximal levels of 5-LO would have enhanced swarming capacity and
that this may contribute to containment of infection. We first compared the resistance to
mechanical wound infection in transgenic larvae in which neutrophils overexpress 5-LO ('5-
LO-Neu'; for 5-lipoxygenase overexpressing neutrophils) versus control wild type larvae. We
found that 5-LO-Neu larvae showed enhanced resistance to infection, as measured by an
improvement in survival rates, disease score and bacterial burden **(Fig. 4A-C)**.
299
To relate neutrophil swarm patterns to disease susceptibility, we combined again live imaging
during the early stage of wound infection, with subsequent monitoring of disease progression

in individual larvae, using 5-LO-Neu transgenic larvae crossed with GCaMP-expressing
neutrophils. As with our mechanical wound model, we found that the 5-LO-Neu larvae showed
greater resistance to infection in our laser wound infection model **(Fig. S4B)**. By observation
of the videos, we noted heterogeneity in swarm magnitude across control larvae, which
appeared to be less pronounced in 5-LO-Neu larvae **(Fig. 4D** and **Videos S3** and **S4)**. This was
not explained by developmental differences in the number of the cells, as the number of
neutrophils in the field of view at the start of the videos was comparable **(Fig. S4C)**. To quantify
these differences, we again used the number of recruited neutrophils or the maximal rate of
recruitment as a measure of recruitment magnitude and speed respectively. Accordingly, we
stratified fish into four groups: i) weak or slow neutrophil recruiters with high bacterial burden
(Group 1, G1, in **Fig S4D** and **Fig. 4E**, respectively), ii) strong or fast recruiters with high
bacterial burden (Group 2, G2, in **Fig. S4D** and **Fig. 4E**, respectively), iii) strong or fast
recruiters with weak bacterial burden (Group 3. G3, in **Fig. S4D** and **Fig. 4E**, respectively) and
iv) weak or slow recruiters with weak bacterial burden (Group 4, G4, in **Fig. S4D** and **Fig. 4E**,
respectively). We found that in control larvae, the main outcomes were in groups G1 and G2
whereas in 5-LO-Neu larvae there was increased proportion of outcomes in group G3 and a
decrease in group G1. Similar trends were observed whether we used the number of cells
recruited or the maximal rate of recruitment **(Fig. 4E and Fig. S4D)**, but the results reached
statistical significance only in the latter case **(Fig. 4F)**.
321
322 Together these results suggested that 5-LO enhancement in neutrophils is sufficient to enhance
323 neutrophil swarming rates and that this can provide a beneficial outcome in a subset of larvae.
324

Discussion

Here, we examine the influence of neutrophil swarming on infection outcome and the extent to
which this migratory behaviour can be reprogrammed to alter disease outcome. We show that
neutrophil swarming and the expression of genes for swarm machinery are altered during
natural reprogramming ('training') by microbial experience. Through chemical inhibition, we
found that these alterations contribute to the enhanced resistance of trained animals to
pathogenic infection. Finally, we show that genetic engineering of 5-LO in neutrophils is
sufficient to accelerate neutrophil swarming and improve bacterial clearance and disease
progression.

Whilst exploring tissue variables influencing bacterial wound infection, we found that osmotic surveillance during tissue damage plays a major role in resistance to opportunistic wound infection. This extends findings reported previously, using *P. aeruginosa* microinjection in the context of wound infection[48]. Previous studies conversely reported that damage detection through osmotic surveillance compromised *Candida albicans* wound infection outcome[53]. Mechanistically, this was attributed to keratinocyte movement in response to hypotonic medium detection, which, rather than limiting microbial colonisation, appeared to favour tissue spreading of *C. albicans*. These discrepancies suggest that the role of osmotic surveillance in wound infection outcome depends on the type of invading microbe. For example, *P. aeruginosa* bacteria are highly mobile[54] and thus may rely less on host tissue movements for spreading than *C. albicans*.

We observed improved resistance to bacterial wound infection after a pre-challenge with non-pathogenic heterologous bacteria. These protective effects were present at a stage when the primary challenge had largely resolved, consistent with innate immune training. A limitation here is that we have not profiled the epigenetic signatures of neutrophils or sampled the resistance effects at longer time scales. However, other studies have shown that zebrafish neutrophils undergo epigenetic modifications during training with *Shigella flexneri* (which are phylogenetically close to *E. coli)* in a comparable timeframe with our study[37]. In addition, other studies have demonstrated that neutrophil training in zebrafish larvae arises through changes in hematopoietic stem cells within similar timeframes post-infection, as has been reported in adult mammals[35]. Our observations are consistent with such central reprogramming of neutrophils, as the protective effects were seen across heterologous tissue sites and the gene expression changes were detectable in neutrophils isolated from hematopoietic tissue. Though the detailed mechanisms remain to be investigated, our study highlights novel routes for training in wound infection settings. Probiotic *E. coli* strains have been reported to have beneficial effects on viral resistance in mouse models[55], but to our knowledge such training has not been reported for wound infection.

We found that neutrophil swarm capacity is amenable to reprogramming after microbial experience. Specifically, neutrophils upregulated expression of genes associated with swarming and larvae were less likely to show poor swarming and high susceptibility to infection. Whether these larvae represent a biologically distinct group of 'less trained' larvae

or part of a normal distribution is difficult to ascertain with this experimental size. However, it
is interesting to note that individual variations in training have been reported in humans
exposed to Bacillus Calmette-Guérin (BCG) vaccine, wherein training improved outcome in a
subpopulation of individuals with sub-maximal immunological profiles at the time of
vaccination[56]. Another noteworthy observation in our data was that training reduced the
higher-end values for swarm kinetics (with migration kinetics more closely clustered around a
medium range). One potential explanation is that the effect of increasing neutrophil numbers
has non-linear effects on bacterial clearance, whereby at low neutrophil numbers the changes
in neutrophil kinetics can make a significant difference in bacterial clearance whereas beyond
a certain number of recruited neutrophils, the benefit decreases. A question raised is why
neutrophil swarming has not evolved to be maximal regardless of immune experience and
whether it is detrimental to have high levels of swarm machinery. We speculate that adaptive
reprogramming of neutrophil swarming may reflect a strategy to economise energy or a means
to balance the risks of overt inflammation.
383
Our discovery that engineering 5-LO is sufficient to improve bacterial clearance has relevance
to cellular immunotherapies. Strategies for genetic modification of immune cells are growing
and extending to cell types beyond lymphocytes, due to advances in genetic engineering and
stem cell biology[39,40,42,57]. In this backdrop, the engineering of chemoattractant receptors has
been conceptualised to improve the efficacy of CAR-engineered immune cell deployment to
tumors[58]. Our study suggests that harnessing the ability of immune cells to seed
chemoattractant gradients in target tissues may offer new routes for reprogramming immune
cell migratory patterns.
392
393
394

Acknowledgements

The *P. aeruginosa* strains were kindly provided by Prof. Martin Welch, Department of Biochemistry, University of Cambridge and the *E. coli* strains were kindly provided by Dr. Somenath Bakshi, Department of Engineering, University of Cambridge. S.M.B., I.W., G.R., C.C., Y.L. were supported by the European Research Council (ERC) under the Horizon 2020 program and UKRI, Grant agreement No. EP/Y02799X/1. Y.L. is also supported by a BBSRC DTP programme BB/X010899/1. N.S.G acknowledges support from the Royal Society Wolfson Visiting Fellowship RSWVF\R2\2320010. H.A.W. was supported by an MRC DTP studentship (MR/N013433/1) and a Leverhulme Trust grant (RPG-2021-226). H.M.C. was supported by the Roving Researcher scheme in the School of Biological Sciences, University of Cambridge. A.D. was supported by a MIRES scholarship from IdEx Université Paris Cité ANR-18-IDEX-0001, funded by the French Government through its 'Investments for the Future' program. The authors gratefully acknowledge the Cambridge Imaging Centre for their support and assistance in this work.

Competing Interests

The authors declare no competing interests

Methods

Experimental model and subject details

All zebrafish were maintained in accordance with UK Home Office regulations under the UK Animals (Scientific Procedures) Act 1986 (PPL number PP4458156), with approval from the University Biomedical Service Committee. Adult fish were maintained and bred following established protocols[59]. Briefly, zebrafish were bred and maintained under standard conditions at $28.5 \pm 0.5\,°C$ on a 14 h light:10 h dark cycle. Embryos were collected from natural spawnings at 3 hours post-fertilization (hpf), bleached for 5 minutes in 0.003% sodium hypochlorite (NaOCl; Cleanline, CL3013), and rinsed three times with E3 medium. They were subsequently incubated at 28 °C in E3 medium supplemented with 0.1 μg /ml methylene blue (Sigma-Aldrich, M9140-25G). For imaging experiments, E3 medium was additionally supplemented with 0.003% 1-phenyl-2-thiourea (PTU; Sigma Aldrich, P7629) to inhibit pigmentation and preserve optical transparency.

For experiments that required *E. coli* training, *E. coli* was administered to larvae at 2 days post-fertilization (dpf) and subsequent re-challenge experiments were done at the 4 dpf stage. In experiments without training, larvae at 3-4 dpf were used. For live imaging, larvae were anesthetized in appropriate buffer containing 160-200 mg/L MS-222 (Sigma-Aldrich, E10521-50G). Where indicated, larvae were treated with Zileuton (20 μM, Tocris, 3308) or appropriate volume of vehicle (DMSO (Sigma-Aldrich, 276855-100ML)) solution by immersion.

The following transgenic zebrafish lines were used: Tg(*lyz*:GCamp6F)*cu104* and Tg(*lyz*:tRFP-5LO) *cu106* from previously described work[9] ,Tg(*mpx*:GFP)i114[60], Tg(*runx1+23*:mCherry).

Bacterial culture and preparation

For *E. coli* injection, cyan fluorescent protein (CFP)-expressing *E. coli* MG1655 7740 strain was developed as described previously[61]. A single colony was obtained from a four-way streak on antibiotic-free LB Agar (Melford, A20020-500.0) plate. The colony was inoculated into 50 ml of antibiotic-free LB broth (Formedium, LBX0102) and cultured at 37 °C with shaking until the OD_{600} reading reached 0.8-0.9. The culture was then aliquoted into 1 ml stocks in antibiotic-free LB broth with 20% v/v glycerol (Fisher Scientific, G/0650/17), snap-frozen with liquid nitrogen, and stored at -80 °C until usage. On the day of the experiment, aliquots of *E. coli* were thawed at 37 °C, pelleted by centrifugation at 5,000 × *g* for seven minutes at 4 °C, and

washed twice with PBS (Sigma Aldrich, D8537). The bacteria were finally suspended in PBS

with 10% v/v phenol red (Sigma, P0290).

For *P. aeruginosa* infection, PAO1 strain was used[9]. One day prior to infection, a single colony

of *P. aeruginosa* was obtained from a four-way streak on Pseudomonas Isolation Agar (PIA;

BD Difco, 292710) supplemented with cetrimide and nalidixic acid (E&O Laboratories Ltd,

LS0006) and glycerol. The colony was inoculated into 5 ml of antibiotic-free LB broth and

cultured at 37 °C with shaking for 24 h. On the day of the experiment, the overnight culture

was diluted in fresh LB and incubated to mid-logarithmic phase (OD_{600} = 0.6–0.8). Bacterial

density was adjusted to 3×10^5 CFU/mL using spectrophotometric estimation. Bacteria were

pelleted at $1700 \times g$ for five minutes and washed twice with PBS to remove residual growth

medium, then resuspended in PBS.

PBS and *E. coli* injections

On the day of injections, *E. coli* injection mix was prepared as described above. PBS injection

mix was prepared as PBS with 10% v/v phenol red for control. Larvae were anaesthetised in

E3 medium supplemented with 0.1 µg/ml methylene blue and 160-200mg/L MS-222. The left

otic vesicle of each zebrafish larva was then injected with 2 drops of 0.125 nL of PBS or *E.*

coli injection mix with a pressure of 20 psi using Picospritzer III microinjector (Intracell), as

previously described[62]. To confirm the dose of *E. coli* injected, 2 drops of 0.125 nL of injection

mix was directly injected into a microcentrifuge tube containing 100 µl PBS after the injections

and were then plated onto antibiotic-free LB Agar plates overnight at 37 °C at 10^0 and 10^{-1}

dilution. The bacterial colony forming units (CFU) were then determined by counting the

number of CFP^+ colonies after overnight incubation. Only when the *E. coli* injection dose

exceeded 400 CFU/ml were the larvae included for subsequent experiments. After injections,

larvae were transferred to fresh E3 medium supplemented with 0.1 µg/mL methylene blue and

kept at 28 °C until further use.

Tail transection, wound infection and survival assay

Zebrafish larvae were transferred to a Petri dish containing Ringer's solution and 160-200 mg/L

MS-222. Tail transection was done using a surgical blade (Swann Morton, 0510) at the distal

boundary of the notochord, avoiding injury to the notochord. Non-transected larvae served as

controls. Immediately after tail transection, larvae were transferred to a six-well plate

containing Ringer's solution with 3×10^5 CFU/mL *P. aeruginosa* (for the role of specific

buffers, E3/isotonic E3/Ringer were used as indicated) and incubated at 32 °C for 3 h. After
incubation, larvae were washed five times in E3 medium (without methylene blue). Individual
larvae were then placed in wells of a 24-well plate containing 1 mL E3 (without methylene
blue) and monitored for survival and disease progression. For 5-LO inhibition assays, 20 μM
zileuton treatment (diluted in DMSO) was initiated 1 h prior to tail transection and maintained
until the end of the experiment.

Survival was assessed immediately after washing, at 3 hpi, and 24 hpi. Assessment was also
done at intermediate time points (7 hpi and 10 hpi) when experiment was done for 24 h. For
48h time point experiments, analysis was done at 3 h, 24 h and 48 h. Phenotypic disease scoring
was done on a 0–4 scale **(Fig. 1B)** based on behavioural responses, transparency, tissue
integrity, circulation, and survival. Between observation timepoints, larvae were returned to 32
°C. A score of 4 was defined as death of the organism that was determined by the absence of
heartbeat as observed under the microscope.

Bacterial burden quantification

At indicated timepoints, zebrafish larvae from each condition were pooled into groups of three
larvae and homogenised in 300 μL PBS containing 0.1% v/v Triton X-100 (VWR, 28817.295)
using sterile pestles. Serial dilutions (up to 10^{-3}) were prepared in PBS and plated on PIA
supplemented with cetrimide and nalidixic acid (as per manufacturer's instructions) for *P.*
aeruginosa or antibiotic-free LB agar for *E. coli.* CFU were then counted after 24 hours
incubation.

Fluorescence-Activated Cell Sorting (FACS)

Approximately 400 embryos from a cross between Tg(*mpx*:GFP)[i114] and
Tg(*runx1+23*:mCherry) transgenic lines were injected with either *E. coli* or PBS (200 per
group) in the otic vesicles at 2 dpf as described above. At 4 dpf, larvae were collected into
microcentrifuge tubes, rinsed twice with 1x HBSS (prepared by diluting 10x HBSS (Gibco™,
14175095) in UltraPure™ DNase/RNase-Free Distilled Water (Invitrogen™, 10977035)), and
placed on ice for 10 minutes to immobilise the larvae. The caudal haematopoietic tissue (CHT)
regions from the larvae were dissected on sterile petri dishes using ethanol-sterilised scalpels
and centrifuged to remove excess HBSS.

For tissue dissociation, 500 μL of digestion solution (2% w/v collagenase (Gibco™, 17104019)
and 40% v/v trypsin (Gibco™, 15400054) in 1x HBSS) was added to each tube. Samples were

incubated at 30 °C for 30 minutes with periodic trituration by pipetting. Digestion was halted with 1x HBSS containing 10% v/v heat-inactivated fetal bovine serum (FBS; Gibco™, A5670701). All subsequent steps were performed on ice. Cells were centrifuged at 500 × g for 5 minutes at 4 °C and then resuspended in sorting solution (1x HBSS supplemented with 10 mM HEPES 15630056) and 2% FBS) and filtered through a pre-wetted 40 µm cell strainer (Falcon, 352340). Filtered cells were centrifuged and resuspended in 1 ml sorting solution in a microcentrifuge tube. Cell viability and concentration were assessed using trypan blue (Gibco, 15250) staining and a hemocytometer, respectively.

GFP-positive neutrophils were FAC-sorted using a BD FACS Aria III (Becton Dickinson) directly into 100 µL of lysis solution supplied with the RNAqueous® Micro Kit (ThermoFisher, AM1931).

RNA extraction and cDNA synthesis

For transcriptomic analysis, RNA isolation was done from the sorted cells using the RNAqueous® Micro Kit (Thermo Fisher, AM1931) according to the manufacturer's instructions, including DNase I digestion step to eliminate genomic DNA contamination.

For cytokine expression analysis following PBS/ *E. coli* injections, at 2 dpf, RNA was extracted from groups of 10 wild-type zebrafish per condition (non-injected, PBS-injected, and *E. coli*-injected) after euthanasia with MS-222 at 4.5 hours post-injection using the RNeasy® Mini Kit (Qiagen, 74104). Subsequently, from 3 to 10 dpf, groups of 8-10 larvae per condition were euthanised with MS-222 prior to RNA extraction.

The obtained RNA concentrations were measured using the Nanodrop. Purified RNA was then reverse transcribed into cDNA using the Maxima H Minus First Strand cDNA Synthesis Kit (ThermoFisher, K1651) following the manufacturer's protocol. Finally, cDNA was stored at -20 °C until further use for quantitative PCR (qPCR) analysis.

Gene expression analysis by qRT-PCR

The qRT-PCR was performed using the PowerTrackTM SYBR™ Green Master Mix (ThermoFisher, A46012) and carried out in the LightCycler® 480 Instrument II (Roche). For transcriptomic analysis, each reaction contained 0.5 ng of cDNA, 10 nM of primer pairs, and 1X master mix. The zebrafish gene *rpl32* served as the reference for normalisation.

For cytokine expression analysis, each reaction contained 5 ng cDNA, 500 nM primer pairs,

and 1X master mix and the zebrafish gene *ef1a* served as the reference for normalisation.

The mRNA levels of target genes relative to a housekeeping gene (**Table 1**) were calculated

using the Livak ($2^{-\Delta\Delta Cqt}$) method. All qRT-PCR reactions for transcriptomic analysis were

performed in duplicate with five biological repeats for transcriptomic analysis while the

cytokine expression experiments were performed in triplicate with three biological repeats.

Table 1

Gene	Primer Sequences (5′ → 3′)
5-lo	Forward: GAGATGGCAGGGCTCGATTG
	Reverse: TACTCATCGGGAAGCCAGGA
cx43	Forward: GCTCTCCACTCTTTACTTCTTTCCAG
	Reverse: GTATTGCACTTGAAAGCTGACTGC
cxcl8a	Forward: GTCGCTGCATTGAAACAGAA
	Reverse: CTTAACCCATGGAGCAGAGG
cxcl8b	Forward: CTACCGAGACGTGGGTGATT
	Reverse: GCTCGGTGAATGGTCATTTT
cxcr2	Forward: TGACCTGCTTTTTTCCCTCACT
	Reverse: TGACCGGCGTGGAGGTA
cxcr1	Forward: GTGATCGTACGCGCTATGGA
	Reverse: ATTCGGGTTGCTAATCGCCA
tubg1	Forward: GGATACACACCGCTGACCAC
	Reverse: AGCCTCCTCATCACATCCAGA
rpl32	Forward: CCCTCACCAAACCTAAGATCGT
	Reverse: CTCCAGTTTGCCCTGATCTTG
ef1a *(housekeeping)*	Forward: AAGGAGAAGATCGACCGTCG
	Reverse: TGATTGAGGGAAATTCACTTGG
cxcl18b	Forward: TGCCAGTGTGAAGAGTCATC
	Reverse: TCGCAGTTTGGTCTTTTAGGG
il6	Forward: AGACGAGCAGTTTGAGAGAGATT
	Reverse: GTTTGAGGAGAGGAGTGCTGAT
il1b	Forward: ATCAAACCCCAATCCACAGAGT

	Reverse: GGCACTGAATCCACCACGTT

Two-photon imaging and laser wounding

Two photon imaging was performed in Tg(*lyz*:GCaMP6f)[cu104] larvae or larvae from crosses between Tg(*lyz*:GCaMP6f) x Tg(*lyz*:tRFP-5LO). Transgenic larvae expressing the calcium indicator GCaMP6f were screened to ensure high reporter expression. Larvae exhibiting strong fluorescence and normal morphology were selected for imaging and infection. Selected larvae were anaesthetised and mounted in a mixture of 1.2% low-melting-point agarose (Invitrogen, 16520) and the medium (E3, isotonic E3, Ringer) containing 160-200 mg/L MS-222. Imaging under sterile conditions was performed as as previously described[49]. For sterile imaging, larvae were mounted on a 35 mm glass-bottom dish with a 14 mm microwell #1.5 coverglass. Excess agarose was removed before solidification to embed the larvae in a thin layer. For infection imaging, larvae were mounted in a custom-built coverslip chamber consisting of glass coverslips sealing both the top and bottom of the sample onto a metallic ring. Larvae were oriented laterally, and the agarose was allowed to solidify. After agarose solidification, the urogenital region was exposed by carefully removing surrounding agarose under a dissecting microscope using a pair of forceps, allowing bacterial access post-wounding. The chamber (for infection conditions) or dishes (for sterile conditions) were filled with 1 mL or 4 mL of Ringer's solution containing 160-200 mg/L MS-222 and 3×10^5 CFU/mL PAO1. For sterile conditions, appropriate buffer (Ringer/ E3 or isotonic E3) was used as indicated.

Laser ablation and time-lapse imaging were performed on a LaVision TriM Scope multiphoton microscope equipped with an electro-optic modulator for rapid power modulation. A Spectra-Physics Insight DeepSee dual-line laser was tuned to 900 nm for imaging and 1,040 nm for ablation, with imaging power set to ~500 mW at the specimen plane. Image acquisition was controlled using ImSpector Pro software (version 5.0.284.0; LaVision Biotec, ©1998–2016). Two-photon imaging was conducted with a 25×/NA 1.05 water-dipping objective. A 40 μm-diameter region of interest was defined on a single superficial focal plane (240 nm/pixel, 15 μs dwell time). Z-stacks (~20 planes, 2 μm step size) were acquired every 20 s. Laser wounding was initiated after a 2 min pre-wound baseline, followed by 2 h of post-wound imaging. Focus was maintained throughout acquisition. Image stacks were processed in Fiji using maximum-intensity z-projections to generate neutrophil-swarming time-lapse videos for subsequent analysis.

Image analyses

Defining maximum rate of recruitment

After image acquisition, each video was manually analysed using Fiji, to determine the number of cells recruited to the wound overtime. Using the number of cells recruited at set time points (at 0 min, 5 min and then every 15 min), the maximum rate of recruitment's function was calculated. In this, only the active recruitment timepoints were considered (i.e. if the recruitment started at 15 min, the timepoints 0 and 5 min were not considered. Similarly, if the recruitment of cells plateaued at later timepoints, those timepoints were not considered for the calculation).

Defining the quadrants used for stratification of the imaging data

For each video, a graph was generated using the logarithmic value of the absolute number of PAO1 CFUs on the y-axis, and either the number of cells recruited or the rate of recruitment function on the x-axis. Thresholds were set in PBS or control groups and then applied to *E. coli* or 5-LO-Neu groups. The threshold for good recruitment was set to more than or equal to 10 cells recruited at the wound overtime. To determine fast recruitment rate, the mean of the maximum recruitment rate of all experiments (PBS and *E. coli* for training and CTR and 5-LO-Neu) was calculated and set as threshold. To evaluate the implication of CFU in disease outcomes, a threshold CFU was set for experiments involving both trained larvae and 5-LO-Neu larvae. The lowest CFU value that resulted in disease manifestations in the observed larvae (disease score ≥ 1) was set as the threshold for the specific set of experiment. Employing the mentioned parameters, we stratified the imaging data into four categories of larvae (G1, G2, G3, G4).

Cell trajectories and Statistics

Neutrophils were semi-automatically tracked using Imaris v8.2 (Bitplane AG, Zürich, Switzerland) on 2D maximum intensity projections of the 4D time-lapse videos. All statistical tests were performed in Prism10 (GraphPad Software, La Jolla, CA). The statistical test and the n number are indicated in the figure legends. The error bars show standard error of the mean except for bacterial burden data where error bars represent 95% confidence intervals of the

623   median. Where the distribution was verified as normal, outliers were removed by applying

624   Rout test. Live imaging experiments were acquired in minimum five independent experiments.

625   A normal distribution test was performed prior to any *t*-test statistical analyses. Normal

626   distribution was assessed using *Shapiro-Wilk test*.

References

1. Schrope, J. H., Robertson, T. F., Sarris, M. & Huttenlocher, A. Chemical and Mechanical Regulation of Leukocyte Migration. *Cold Spring Harb. Perspect. Biol.* a041752 (2025) doi:10.1101/cshperspect.a041752.

2. Insall, R. H., Paschke, P. & Tweedy, L. Steering yourself by the bootstraps: how cells create their own gradients for chemotaxis. *Trends Cell Biol.* **32**, 585–596 (2022).

3. Stock, J., Kazmar, T., Schlumm, F., Hannezo, E. & Pauli, A. A self-generated Toddler gradient guides mesodermal cell migration. *Sci. Adv.* **8**, eadd2488 (2022).

4. Uçar, M. C., Alsberga, Z., Alanko, J., Sixt, M. & Hannezo, E. Self-generated chemotaxis of mixed cell populations. *Proc. Natl. Acad. Sci.* **122**, e2504064122 (2025).

5. Dalle Nogare, D. & Chitnis, A. B. A framework for understanding morphogenesis and migration of the zebrafish posterior Lateral Line primordium. *Mech. Dev.* **148**, 69–78 (2017).

6. Muinonen-Martin, A. J. *et al.* Melanoma Cells Break Down LPA to Establish Local Gradients That Drive Chemotactic Dispersal. *PLoS Biol.* **12**, e1001966 (2014).

7. Tweedy, L. *et al.* Seeing around corners: Cells solve mazes and respond at a distance using attractant breakdown. *Science* **369**, eaay9792 (2020).

8. Lämmermann, T. *et al.* Neutrophil swarms require LTB4 and integrins at sites of cell death in vivo. *Nature* **498**, 371–375 (2013).

9. Poplimont, H. *et al.* Neutrophil Swarming in Damaged Tissue Is Orchestrated by Connexins and Cooperative Calcium Alarm Signals. *Curr. Biol.* **30**, 2761-2776.e7 (2020).

10. Strickland, E. *et al*. Self-extinguishing relay waves enable homeostatic control of

human neutrophil swarming. *Dev. Cell* **59**, 2659-2671.e4 (2024).

11. Afonso, P. V. *et al*. LTB4 Is a Signal-Relay Molecule during Neutrophil Chemotaxis. *Dev.*

Cell **22**, 1079–1091 (2012).

12. Alanko, J. *et al*. CCR7 acts as both a sensor and a sink for CCL19 to coordinate

collective leukocyte migration. *Sci. Immunol*. **8**, eadc9584 (2023).

13. Burn, G. L., Foti, A., Marsman, G., Patel, D. F. & Zychlinsky, A. The Neutrophil.

Immunity **54**, 1377–1391 (2021).

14. Reátegui, E. *et al*. Microscale arrays for the profiling of start and stop signals

coordinating human-neutrophil swarming. *Nat. Biomed. Eng*. **1**, 0094 (2017).

15. Ng, L. G. *et al*. Visualizing the Neutrophil Response to Sterile Tissue Injury in Mouse

Dermis Reveals a Three-Phase Cascade of Events. *J. Invest. Dermatol*. **131**, 2058–

2068 (2011).

16. Chtanova, T. *et al*. Dynamics of Neutrophil Migration in Lymph Nodes during Infection.

Immunity **29**, 487–496 (2008).

17. Hopke, A. *et al*. Neutrophil swarming delays the growth of clusters of pathogenic

fungi. *Nat. Commun*. **11**, 2031 (2020).

18. Saraswat, D. *et al*. Neutrophil swarming is crucial for limiting oral mucosal infection

by *Candida albicans*. *J. Leukoc. Biol*. **117**, qiae239 (2025).

19. Herrero-Cervera, A., Soehnlein, O. & Kenne, E. Neutrophils in chronic inflammatory

diseases. *Cell. Mol. Immunol*. **19**, 177–191 (2022).

20. Isles, H. M. *et al*. Pioneer neutrophils release chromatin within in vivo swarms. *eLife*

10, e68755 (2021).

21. Khazen, R. *et al.* Spatiotemporal dynamics of calcium signals during neutrophil
cluster formation. *Proc. Natl. Acad. Sci.* **119**, e2203855119 (2022).

22. Kienle, K. *et al.* Neutrophils self-limit swarming to contain bacterial growth in vivo.
Science **372**, eabe7729 (2021).

23. Song, Z., Bhattacharya, S., Clemens, R. A. & Dinauer, M. C. Molecular regulation of
neutrophil swarming in health and disease: Lessons from the phagocyte oxidase.
iScience **26**, 108034 (2023).

24. Uderhardt, S., Martins, A. J., Tsang, J. S., Lämmermann, T. & Germain, R. N. Resident
Macrophages Cloak Tissue Microlesions to Prevent Neutrophil-Driven Inflammatory
Damage. *Cell* **177**, 541-555.e17 (2019).

25. Lämmermann, T. In the eye of the neutrophil swarm—navigation signals that bring
neutrophils together in inflamed and infected tissues. *J. Leukoc. Biol.* **100**, 55–63
(2016).

26. Netea, M. G. *et al.* Defining trained immunity and its role in health and disease. *Nat.*
Rev. Immunol. **20**, 375–388 (2020).

27. Kleinnijenhuis, J. *et al.* Bacille Calmette-Guérin induces NOD2-dependent
nonspecific protection from reinfection via epigenetic reprogramming of monocytes.
Proc. Natl. Acad. Sci. **109**, 17537–17542 (2012).

28. Quintin, J. *et al.* Candida albicans Infection Affords Protection against Reinfection via
Functional Reprogramming of Monocytes. *Cell Host Microbe* **12**, 223–232 (2012).

29. Kalafati, L. *et al.* Innate Immune Training of Granulopoiesis Promotes Anti-tumor
Activity. *Cell* **183**, 771-785.e12 (2020).

30. Kato, Y. & Kumanogoh, A. The immune memory of innate immune systems. *Int.*
Immunol. **37**, 195–202 (2025).

31. Mitroulis, I. *et al.* Modulation of Myelopoiesis Progenitors Is an Integral Component of

Trained Immunity. *Cell* **172**, 147-161.e12 (2018).

32. Kaufmann, E. *et al.* BCG Educates Hematopoietic Stem Cells to Generate Protective

Innate Immunity against Tuberculosis. *Cell* **172**, 176-190.e19 (2018).

33. Schlüter, T., Van Elsas, Y., Priem, B., Ziogas, A. & Netea, M. G. Trained immunity:

induction of an inflammatory memory in disease. *Cell Res.*

https://doi.org/10.1038/s41422-025-01171-y (2025) doi:10.1038/s41422-025-01171-

y.

34. Fanucchi, S., Domínguez-Andrés, J., Joosten, L. A. B., Netea, M. G. & Mhlanga, M. M.

The Intersection of Epigenetics and Metabolism in Trained Immunity. *Immunity* **54**,

32–43 (2021).

35. Darroch, H. *et al.* Infection-experienced HSPCs protect against infections by

generating neutrophils with enhanced mitochondrial bactericidal activity. *Sci. Adv.* **9**,

eadf9904 (2023).

36. Mariotti, B. *et al.* Innate immune reprogramming in circulating neutrophils of COPD

patients. *J. Allergy Clin. Immunol.* **156**, 373–384 (2025).

37. Gomes, M. C., Brokatzky, D., Bielecka, M. K., Wardle, F. C. & Mostowy, S. *Shigella*

induces epigenetic reprogramming of zebrafish neutrophils. *Sci. Adv.* **9**, eadf9706

(2023).

38. Patel, K. K., Tariveranmoshabad, M., Kadu, S., Shobaki, N. & June, C. From concept to

cure: The evolution of CAR-T cell therapy. *Mol. Ther.* **33**, 2123–2140 (2025).

39. Abdin, S. M., Paasch, D. & Lachmann, N. CAR macrophages on a fast track to solid

tumor therapy. *Nat. Immunol.* **25**, 11–12 (2024).

40. Chang, Y. *et al*. CAR-neutrophil mediated delivery of tumor-microenvironment

responsive nanodrugs for glioblastoma chemo-immunotherapy. *Nat. Commun.* **14**,

2266 (2023).

41. Chang, Y. *et al*. Engineering chimeric antigen receptor neutrophils from human

pluripotent stem cells for targeted cancer immunotherapy. *Cell Rep.* **40**, 111128

(2022).

42. Majumder, A. *et al*. iPSC-Derived CAR Neutrophils Possess Potent Activity Against

Solid Tumors In Vivo. *Blood* **144**, 86–86 (2024).

43. Sathe, N. *et al*. Pseudomonas aeruginosa: Infections and novel approaches to

treatment "Knowing the enemy" the threat of Pseudomonas aeruginosa and exploring

novel approaches to treatment. *Infect. Med.* **2**, 178–194 (2023).

44. Ruffin, M. & Brochiero, E. Repair Process Impairment by Pseudomonas aeruginosa in

Epithelial Tissues: Major Features and Potential Therapeutic Avenues. *Front. Cell.*

Infect. Microbiol. **9**, 182 (2019).

45. Clatworthy, A. E. *et al*. *Pseudomonas aeruginosa* Infection of Zebrafish Involves both

Host and Pathogen Determinants. *Infect. Immun.* **77**, 1293–1303 (2009).

46. Gault, W. J., Enyedi, B. & Niethammer, P. Osmotic surveillance mediates rapid wound

closure through nucleotide release. *J. Cell Biol.* **207**, 767–782 (2014).

47. Enyedi, B., Kala, S., Nikolich-Zugich, T. & Niethammer, P. Tissue damage detection by

osmotic surveillance. *Nat. Cell Biol.* **15**, 1123–1130 (2013).

48. Huang, C. & Niethammer, P. Tissue Damage Signaling Is a Prerequisite for Protective

Neutrophil Recruitment to Microbial Infection in Zebrafish. *Immunity* **48**, 1006-

1013.e6 (2018).

49. Williantarra, I., Georgantzoglou, A. & Sarris, M. Visualising Neutrophil Actin Dynamics

in Zebrafish in Response to Laser Wounding Using Two-Photon Microscopy. *BIO-*

Protoc. **14**, (2024).

50. LaBauve, A. E. & Wargo, M. J. Growth and Laboratory Maintenance of *Pseudomonas*

aeruginosa. *Curr. Protoc. Microbiol.* **25**, (2012).

51. Potvin-Trottier, L., Lord, N. D., Vinnicombe, G. & Paulsson, J. Synchronous long-term

oscillations in a synthetic gene circuit. *Nature* **538**, 514–517 (2016).

52. Park, S. A., Choe, Y. H., Park, E. & Hyun, Y.-M. Real-time dynamics of neutrophil

clustering in response to phototoxicity-induced cell death and tissue damage in

mouse ear dermis. *Cell Adhes. Migr.* 1–20 (2018)

doi:10.1080/19336918.2018.1471322.

53. Horn, A. *et al.* Isotonic medium treatment limits burn wound microbial colonisation

and improves tissue repair. *Wound Repair Regen.* **33**, e13242 (2025).

54. Kazmierczak, B. I., Schniederberend, M. & Jain, R. Cross-regulation of Pseudomonas

motility systems: the intimate relationship between flagella, pili and virulence. *Curr.*

Opin. Microbiol. **28**, 78–82 (2015).

55. Huang, L. *et al.* Engineered probiotic Escherichia coli elicits immediate and long-term

protection against influenza A virus in mice. *Nat. Commun.* **15**, 6802 (2024).

56. Moorlag, S. J. C. F. M. *et al.* Multi-omics analysis of innate and adaptive responses to

BCG vaccination reveals epigenetic cell states that predict trained immunity.

Immunity **57**, 171-187.e14 (2024).

57. Wang, W. *et al.* Breakthrough of solid tumor treatment: CAR-NK immunotherapy. *Cell*

Death Discov. **10**, 40 (2024).

58. Sterner, R. C. & Sterner, R. M. CAR-T cell therapy: current limitations and potential

strategies. *Blood Cancer J.* **11**, 69 (2021).

59. Bradford, Y. M. *et al.* Zebrafish information network, the knowledgebase for *Danio*

rerio research. *Genetics* **220**, iyac016 (2022).

60. Renshaw, S. A. *et al.* A transgenic zebrafish model of neutrophilic inflammation.

Blood **108**, 3976–3978 (2006).

61. Bakshi, S. *et al.* Tracking bacterial lineages in complex and dynamic environments

with applications for growth control and persistence. *Nat. Microbiol.* **6**, 783–791

(2021).

62. Sarris, M. *et al.* Inflammatory Chemokines Direct and Restrict Leukocyte Migration

within Live Tissues as Glycan-Bound Gradients. *Curr. Biol.* **22**, 2375–2382 (2012).

Figure Legends

FIGURE_1

A. Schematic detailing the mechanical wound (MW) wound-infection model by tail fin (TF) transection method in the presence of *P. aeruginosa* PAO1.

B. Parameters assessed in the disease scoring matrix to accord a disease progression stage to each zebrafish larva.

C. Disease progression in the specified group of larvae at the indicated time points. Representative of three independent experiments, with n ≥ 24 larvae per group.

D. Bacterial burden of *P. aeruginosa* in the specified groups of zebrafish larvae at 24 hpi (hours post infection). CFU, colony forming units. (Representative of three independent experiments with n ≥9 larvae per group used for CFU analysis in each experiment). (mean ± SD). *One-way ANOVA with Tukey's multiple comparisons test, ***P < 0.001*

E. Survival curves from zebrafish larvae when exposed to indicated buffers and *P. aeruginosa* infection upon MW. (Representative of three independent experiments with n ≥ 24 larvae per group for the wounded-infected; n ≥ 9 larvae per group for the non-wounded infected, in each experiment). *Log-rank (Mantel-Cox) test; ***P < 0.001*

F. Bacterial burden of *P. aeruginosa* in the specified groups of zebrafish larvae at early time points (10 mins and 30 mins) of infection. (Representative of three independent experiments with n ≥ 12 larvae per group per time point in each experiment. (mean ± SD). CFU, colony forming units. *One-way ANOVA with, Šídák's multiple comparisons test, **P < 0.01*

G. Schematic detailing the laser wound (LW) infection model. CHT: caudal hematopoietic tissue; VF: ventral fin.

H. Survival curves from zebrafish larvae when exposed to indicated buffers and *P. aeruginosa* infection upon LW. (Representative of three independent experiments with n ≥ 24 larvae per group for the wounded-infected; n≥12 larvae per group for the non-wounded infected, in each experiment). *Log-rank (Mantel-Cox) test, *P < 0.05*

I. Bacterial burden of *P. aeruginosa* in the specified groups of zebrafish larvae at 24 hpi (hours post infection). Representative of three independent experiments, with n ≥12 larvae per group in each experiment. (mean ± SD). CFU, colony forming units. *Unpaired t test, *P < 0.05*

J. Disease progression in the specified group of larvae at the indicated time points.
Representative of three independent experiments with n ≥ 24 larvae per group.

FIGURE _2

A. Schematic of the training experiment. Zebrafish larvae pre-challenged by PBS or a dose
of *E. coli* in the otic vesicle (OV) at 2 days post fertilization (dpf) stage and
subsequently subjected to mechanical wounding in the tail fin (TF) and *P. aeruginosa*
PAO1 infection at 4 dpf.

B. Survival curves from control (PBS) and trained (EC) zebrafish larvae upon *P.*
aeruginosa infection following MW. (Representative of three independent experiments
with n ≥ 36 larvae per group in each experiment). *Log-rank (Mantel-Cox) test; ****P*
< 0.0001.

C. Bacterial burden of *P. aeruginosa* in the control and trained zebrafish larvae at 24 hpi
(hours post infection). (Representative of three independent experiments, with n ≥ 9
larvae per group used for CFU analysis in each experiment. (mean ± SD). CFU, colony
forming units. *One-way ANOVA with Tukey's multiple comparisons test, *P < 0.05*

D. Disease progression in the control and trained zebrafish larvae at the indicated time
points. Representative of three independent experiments with n ≥ 36 larvae per group
in each experiment.

E. Schematic detailing the assessment of *E. coli* burden and cytokine profile in whole
zebrafish larvae after primary challenge (PBS/ *E. coli*) for trained immunity induction.

F. Bacterial counts at the indicated time points after injection of *E. coli* in the otic vesicle
(OV) of the zebrafish larvae. Representative of three independent experiments, with n
≥ 12 larvae for each time point.

G-J. Change in the expression profile of the indicated cytokines in naive and trained
zebrafish larvae at the specific time points after primary challenge. Representative of
three independent experiments, with n ≥ 8 larvae at each time point.

K. Schematic describing the isolation of neutrophils by Fluorescence Activated Cell
Sorting (FACS) from the hematopoietic site (CHT: Caudal hematopoietic tissue) of
zebrafish larvae, followed by transcript analysis using RT-PCR.

L-R. Change in the expression profile of the indicated genes in naive and trained zebrafish
larvae at 4dpf (2 days after primary challenge). Quantitative PCR data are presented as
mean ± S.E.M., with expression levels normalised to PBS-treated control groups (n=5

independent experiments, with each replicate representing a pool of 150-200 larvae at

4 dpf). Statistical significance was assessed using the Mann-Whitney test/ t-test (*p <

0.05, **p < 0.01).

FIGURE_3

A. Schematic of experiment with control, PBS pre-challenged or trained, *E. Coli* pre-

challenged (EC) zebrafish larvae subjected to laser wounding (LW), followed by

infection with *P. aeruginosa* (PAO1) and simultaneous confocal imaging.

B. Time lapse images of representative videos of control and trained larvae at 1 min prior

LW (-1 min), at the time of LW (0 min), at 30 min and 90 min post LW. White dashed

lines correspond to the CHT: caudal hematopoietic tissue and VF: ventral fin. Red

dashed lines correspond to the site of the LW. On the far-right, neutrophil trajectories

of same video are depicted wherein each colour corresponds to a different neutrophil

trajectory. Scale bar 50 μm.

C. Summary of individual larva-based analyses of control (PBS) *(left)* and trained (EC)

animals *(right)*. The y-axis on the graph depicts bacterial burden in larva at 24 hours

post infection (hpi) (after LW and infection), and the x-axis indicates cumulative

number of neutrophils recruited at 120 minutes post LW. Thresholds are set at the

minimum PAO1 CFU value in which disease symptoms were observed (stage 1 or

above, light blue square), and 10 neutrophils recruited over time. Black arrows point to

videos used for representative images in B (PBS example 1= PBS G1; PBS example 2=

PBS G2; EC example 1= EC G2; EC example 2= EC G3). Different shapes and colour

show the individual disease stages of each larva.

D. Contingency analysis of groups in C. n=16 (PBS) and n=17 (EC). *Fisher's exact test **

p < 0.05.

E-F. Survival curves from control (PBS) and trained (EC) zebrafish larvae upon PAO1

infection following mechanical wound. Larvae were pre-treated with DMSO or 20μM

Zileuton (Zil), 1h prior to wounding and kept in this medium until the end of the

experiment Representative of three independent experiments with n ≥28 larvae per

group for each experiment in E, and n ≥ 20 larvae in F. *Log-rank (Mantel-Cox) test; **

p < 0.05 , ns p > 0.05.

G-H. Bacterial burden of *P. aeruginosa* in larvae from groups in E and F respectively.

Representative of three independent experiments with n ≥ 9 larvae per group used for

CFU analysis in each experiment. (mean ± SD). CFU, colony forming units. *Unpaired*
t test with Welch's correction. * p < 0.05, ns p > 0.05.

FIGURE_4

A. Survival curves from control (CTR) zebrafish larvae and larvae overexpressing 5-LO
in neutrophils (5-LO-Neu) upon *P. aeruginosa* infection following MW.
Representative of three independent experiments with $n \geq 50$ larvae per group in each
experiment. *Log-rank (Mantel-Cox) test; * p < 0.05.*
B. Bacterial burden of fish in survival experiments (A). Representative of three
independent experiments with $n \geq 9$ larvae per group used for CFU analysis in each
experiment. *Unpaired t test with Welch's correction; * p < 0.05, ** p < 0.01.*
C. Disease progression in the specified group of larvae at the indicated time points.
Representative of three independent experiments with $n \geq 50$ larvae per group in each
experiment.
D. Time lapse images of representative videos of CTR and 5-LO-Neu at 1 min prior to LW
(-1 min), at the time of LW (0 min), at 30 min and 90 min after LW. White dashed lines
indicated the CHT: caudal hematopoietic tissue and VF: ventral fin. Red dashed line
corresponds to the site of the laser wound (LW). On the right column neutrophil
trajectories of same video are represented wherein each colour indicates a different
neutrophil trajectory. Scale bar 50 μm.
E. Summary of individual larva-based analyses of control (CTR) larvae (*left*) and 5-LO-
Neu larvae (*right*). The y-axis on the graph depicts bacterial burden PAO1 in larva at
24 hours post infection (hpi) (after LW and infection), and the x-axis indicates the
maximum rate of neutrophil recruitment. Thresholds are set at the minimum CFU value
in which disease symptoms were observed (stage 1, light blue square) and the mean of
maximum rate of recruitment for both CTR and 5-LO-Neu experiments. Black arrows
point to videos used for representative images in (D) (arrow in CTR example 1= CTR
G1, CTR example 2=CTR G2; 5-LO- example 1=5-LO-Neu G2, 5-LO-Neu example
2=5-LO-Neu G3). Different shapes and colour show disease stages of individual larva.
Videos of larva wherein recruiting cells were observed at only one time point were
excluded.
F. Contingency analysis of groups in E. n=15 (CTR) and n=18 (5LO). *Fisher's exact test*
* p < 0.05.

916
917