Striatal ensembles specify and control granular forelimb actions

Ines Rodrigues-Vaz[1-4]†, Vivek R. Athalye[1,3,4]†, Darcy S. Peterka[1,4,5], and Rui M. Costa[1,3,4,5]*

[1] Zuckerman Mind Brain Behavior Institute, Departments of Neuroscience and Neurology; Columbia University; New York, NY, 10027; USA

[2] Champalimaud Neuroscience Programme, Champalimaud Centre for the Unknown, Lisbon 1400-038, Portugal

[3] Allen Institute, Allen Institute for Neural Dynamics, Seattle, WA, 98109; USA

[4] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD20815

[5] Kavli Institute for Brain Science, Columbia University; New York, NY 10027; USA

† These authors contributed equally to this work.

*Corresponding author. rui.costa@alleninstitute.org (RMC)

The ability of the brain to control specific fine actions is crucial for survival. Although striatal dysfunction underlies numerous movement disorders[1–9], its activity has classically been studied in the context of movement reinforcement and invigoration [5,10–13]. Recent studies have revealed that striatal activity encodes specific whole-body [14,15] and forelimb movements[16,17]. However, it is unclear if this activity actually controls specific movements. We developed a closed-loop system to identify and manipulate action-specific neuronal ensembles in real time using holographic optogenetics through a GRIN lens. We designed an isometric force task in which mice performed push or pull actions on an immobile joystick without overt kinematics—actions that engage the same forelimb muscles but with different activation patterns. Using 2-photon calcium imaging, we simultaneously recorded D1- and D2-medium spiny neurons (MSNs) in the dorsolateral striatum - classically viewed as promoting versus inhibiting movement[2,3,18] – and found that both populations equally encoded isometric action identity. Stimulation of action-specific D1- and D2-MSN ensembles increased ongoing force, but only when the stimulated ensemble was congruent with the ongoing action. These results reveal that specific ensembles of D1- and D2-MSNs control specific actions, as granular as distinct muscle activation patterns of the same forelimb. These findings provide a novel framework for understanding how striatal dysfunction can produce highly specific movement impairments in Huntington's disease[2,9] and dystonia[2,6].

The brain must generate signals that enable the execution of appropriate actions, many of which demand granular motor control. Writing, for example, requires precisely coordinated muscle co-contractions in the arm to apply specific forces to a pen. Whereas motor cortices, brainstem, and spinal circuits are classically linked[19] to the online specification and execution of movement kinematics and force, the striatum has been primarily framed as a center for invigoration[5,10–13] and reinforcement[12,13,20–26] rather than as a controller of ongoing action per se. This view has been supported by the fact that Parkinson's disease, where dopamine inputs to striatum are depleted, leads to deficits in action initiation[4], vigor[5], and reinforcement[7,8]. Alternatively, accumulating evidence suggests that the striatum specifies actions[14–17] rather than merely broadcasting a more generic "go/no-go"[27] or vigor signal. This is consistent with

45 observations that in disorders where striatal neurons are affected, like in Huntington's disease[9]
46 or dystonia[2,6], the execution of particular specific movements is disrupted.
47
Recent work shows that the activity of specific striatal ensembles encodes particular whole-body
[14,15] and forelimb movements[16,17]. However, it remains untested if this specific activity does
indeed control the ongoing actions. To address this, we developed a real-time system to
identify neuronal ensembles with a statistical model and manipulate them with cellular precision
in closed-loop with behavior. Our all-optical system measured neuronal activity with two-photon
calcium imaging and stimulated targeted neurons with holographic optogenetics[28–31] deep in the
brain through a GRIN lens[30,32].
55
Using this system, we asked whether the striatum controls ongoing, granular actions. We
designed a self-paced isometric force task, in which head-fixed mice executed push or pull
forelimb actions on an immobile joystick, activating the same forelimb muscles but with different
patterns. This task allows us to test orthogonal aspects to recent studies using cued movements
with subtly different kinematics[17], and enables us to test whether the striatum can encode and
control different self-paced forelimb actions with similar force and no overt kinematics, just
different patterns of activation of the same muscles.
63
Striatal projection neurons are two distinct cell types, D1- and D2-MSNs, with different
dopamine receptors and projection patterns, that have been canonically assigned opponent
roles in movement promotion and suppression, respectively [2,3,18,33]. However, they both receive
cortical inputs that carry specific movement plans[34–38], both are activated at movement
initiation[39,40], and both seem necessary for execution[40,41]. Therefore, we first tested whether D1-
and D2-MSNs would encode the identity of specific isometric actions differently or similarly. We
found that both the activity of specific D1- and D2-MSN ensembles differentially encoded the
identity of the different isometric actions.
72
Then we tested if activity of these specific ensembles could control particular ongoing
movements. We modeled the dimensions of population activity that predict specific actions,
identified small, action-specific ensembles within spatially intermingled D1- and D2-MSNs, and
stimulated those ensembles precisely during self-paced actions with millisecond latency. We
thus tested whether stimulating action-specific ensembles increases the force of the ongoing
action, and whether this specificity holds for both D1- and D2-MSNs.
79
80 Together, this strategy allowed us to determine that the striatum specifies ongoing actions, with
81 specific ensembles of both D1- and D2-MSNs controlling specific actions, as granular as distinct
82 patterns of forelimb muscles without overt movement.
83
An isometric force task to study granular forelimb actions
To investigate whether the striatum encodes and controls granular actions of the same body
part, we developed a task in which mice performed two forelimb actions (Figures 1A and S4A),
consisting of a push or pull force on an immobile joystick in the absence of overt movement (i.e.
isometric co-contractions, STAR Methods). Mice initiated actions in a self-paced manner, in the

absence of external cues. The task enabled analysis of push and pull actions with matched
continuous parameters including force, duration, and licking (Figure S3A-H, STAR Methods).
91
Mice performed isometric actions in order to receive sucrose reinforcement. We defined two
targets as push and pull that exceed thresholds of force and duration determined for each
session (Figure 1B,C, Table 2,3, STAR Methods), and these actions triggered delayed sucrose
reward (5 μl per reward). Mice selectively increased performance of whichever action was
reinforced, confirming that they performed actions to obtain sucrose (Figures 1D, S2A-C, STAR
Methods).
98
We characterized the simultaneous activity of four forelimb muscles - biceps, triceps, palmaris
longus (PL), and the extensor digitorium communis (EDC) - during the performance of these
isometric actions (Figure 1E). All four muscles were activated for both isometric push and pull
(Figure 1F,G), but the pattern across muscles was action-specific (Fig. 1H). Thus, both
isometric push and pull were executed through activation of the same forelimb muscles -- co-
contraction of extensor and flexor pairs -- but with action-specific activation patterns.

105
Figure 1: An isometric force task engages the same forelimb muscles with action-specific
activation patterns.
(A) Task schematic.
(B) Task events for reinforcement of self-paced actions. Trace of force applied to the joystick on
an example trial.
(C) Trial-averaged force trace of 6g cross actions time locked to force peak. By our convention,
force in the push direction is positive and pull direction is negative. Data are mean+s.e.m.
across 109 sessions from 8 mice.
(D) Rate of 6g cross actions. Mice performed both actions and increased the action which led to
sucrose reinforcement. Data are mean+s.e.m. from 8 mice, averaging over sessions in each
block (STAR Methods). Mice performed push more than pull when both actions were reinforced
(Table S1, ref. 1.1). Mice performed pull more than push when pull was reinforced (Table S1,
ref. 1.2). Mice performed push more than pull when push was reinforced (Table S1, ref. 1.3).
(E) Schematic of forelimb muscles simultaneously recorded with EMG.
(F) Trial-averaged z-scored EMG of 6g cross push actions locked to 6g cross, force-matched
with pull actions (STAR Methods). Each muscle significantly activated at 6g cross relative to 1
second before (Table S1, ref. 1.4-1.7).

(G) Same as (F) but for pull action, force-matched with push actions. Each muscle significantly
activated at 6g cross relative to 1 second before (Table S1, ref. 1.8-1.11).
(H) Vector magnitude of EMG difference at 6g cross. The difference across actions is greater
than within each action (Table S1, ref. 1.12,1.13).
(F-H) Data are mean+s.e.m. over n=41 sessions from 4 mice.
128

Striatal ensembles encode the identity of isometric actions
To test whether the striatum encodes the identity of isometric forelimb actions, we measured the
simultaneous activity of D1- and D2-MSNs in the forelimb region of dorsolateral striatum[42,43]
(DLS) (Figure 2A,B). To identify D1- and D2-MSNs within the same subject[44], we used
transgenic mice that expressed the red label tdTomato either in D1-MSNs (T6 and Ai9xEY217)
or D2-MSNs (Ai9xAdora2cre) (Table 1), and we developed an automated method to classify red
versus non-red neurons (Figure S1). tdTomato-positive neurons were thus identified as D1-
MSNs (or D2-MSNs, depending on the mouse line), and tdTomato-negative neurons were
classified as the other subtype; together, D1- and D2-MSNs account for approximately 95% of
all striatal neurons, so non-red neurons are overwhelmingly the other MSN subtype[44–46]. We
virally expressed GCaMP6f in both MSN populations in DLS using the *Camk2a* promoter[44] and
used 2-photon microscopy through a 1mm diameter GRIN lens to image neuronal calcium
activity as a proxy for action potentials. We were able to image up to ~200 neurons
simultaneously, and the fluorescence signal in both color channels was stable across months of
animal training (Figure S3K). We identified ongoing isometric actions that exceeded 3g of force
for ~66 ms duration (two imaging frames) and that were temporally isolated (i.e. had no
preceding action in the 0.5 seconds before onset). The activity of D1- and D2-MSNs was
modulated for both actions (Figure 2B,C), with slightly greater activity in D1-MSNs (Figure 2C).
147

Many individual neurons modulated their activity only during a specific action (Figure 2D). We
therefore asked whether the spatiotemporal activity of MSNs encoded the identity of ongoing
actions. We extracted the vector of MSN population activity averaged in 150 ms bins (5 imaging
frames) centered at time lags relative to force peak for each trial, and we decoded single-trial
population activity into action identity (push or pull) using a linear support vector machine
(training in 90% of data, testing in 10% held out data). For time lags before (after) force peak,
we analyzed trials with no other action in the 0.5 seconds before (after) force peak. Population
activity predicted action identity well above chance with peak accuracy at the time of peak force
(Figure 2E,F). In addition, population activity predicted action identity in advance of force peak
(Figure 2F). Interestingly, both D1- and D2- MSNs encoded action identity with equal accuracy
and timing (Figure 2G).

Figure 2: D1- and D2-MSNs both encode identity of isometric forelimb actions.

(A) Schematic of experimental procedures. Top: GCaMP6f virus injection and lens implant in

transgenic mice to image D1- and D2-MSNs simultaneously. Bottom: Schematic of automated

method for identifying D1- and D2-MSNs based on transgenic expression of tdTomato (STAR

Methods).

(B) Example trace of force, individual neuronal activity for all ROIs in this field of view,

population activity averaged across neurons, averaged activity for D1-MSNs (n=21 neurons)

and averaged activity for D2-MSNs (n=21 neurons). Neural activity is $\Delta F/F_0$.

(C) Population activity locked to peak force of push and pull actions, averaged across D1-, D2-,

and all MSNs. Data are mean+s.e.m. across 86 sessions from 8 mice. Activity increased at

push force peak ("0 s") relative to "-0.5 s" for D1-MSNs (Table S1, ref. 2.1) and D2-MSNs (Table

S1, ref. 2.2). Activity increased at pull force peak ("0 s") relative to "-0.5 s" for D1-MSNs (Table

S1, ref. 2.3) and D2-MSNs (Table S1, ref. 2.4). D1-MSNs were more active than D2-MSNs at

force peak (Table S1, ref. 2.5,2.6). Bar plots were $\Delta F/F_0$ averaged across 3 event-centered

imaging frames.

(D) Example traces of individual D1- or D2-MSN activity locked to force peak for push and pull

actions.

(E) Decoding all MSN activity to predict action identity on single trials with a model fit at each

time lag relative to force peak. Accuracy at force peak was greater than a model ("shuffle")

trained with shuffled action identity (Table S1, ref. 2.7). Accuracy was greater than shuffle when

averaged over time points in the 4 seconds before force peak (Table S1, ref. 2.8) and in the 0.5

seconds before force peak during which there was no other action (Table S1, ref. 2.9). Data are
mean+s.e.m. across n=87 sessions from 8 mice.
(F) Decoding all MSN activity overlayed on the average force of actions (normalized to 0 at -4s
and 1 at peak of 0s).
(G) Decoding D1- or D2- MSN activity separately. Accuracy at force peak was greater than
shuffle for D1-MSNs (Table S1, ref. 2.10) and D2-MSNs (Table S1, ref. 2.11), with no significant
difference between D1- and D2-MSNs (Table S1, ref. 2.12). Accuracy was greater than shuffle
when averaged over time points in the 4 seconds before force peak for D1-MSNs (Table S1, ref.
2.13) and D2-MSNs (Table S1, ref. 2.14) and in the 0.5 seconds before force peak for D1-MSNs
(Table S1, ref. 2.15) and D2-MSNs (Table S1, ref. 2.16). Data are mean+s.e.m. across n=87
sessions from 8 mice.
(H-L) Data are mean+s.e.m. across 86 sessions from 8 mice.
(H) Decoding MSN activity at force peak for push and pull, predicting action identity between
pairs of actions including push, pull, touch, and licking. Accuracy was greater than shuffle for all
action pairs (Table S1, ref. 2.17-2.22).
(I) Same as (G), decoding D1- and D2-MSNs activity. Decoding push versus pull was not
significantly different between D1- and D2-MSNs (Table S1, ref. 2.23). Accuracy was greater
than shuffle for all action pairs for D1-MSNs (Table S1, ref. 2.24-2.29) and D2-MSNs (Table S1,
ref. 2.30-2.35).
(J) Projection of population activity (z-scored $\Delta F/F_0$) on the SVM dimension locked to force
peak. Inset: average SVM projection at 500 ms before force peak ("-0.5 s") and at force peak ("0
s"). For the push action, the SVM projection at force peak ("0 s") increased relative to "-0.5 s"
(Table S1, ref. 2.36). For the pull action, the SVM projection at force peak ("0 s") decreased
relative to "-0.5 s" (Table S1, ref. 2.37). The SVM projection at force peak was greater for push
than for pull (Table S1, ref. 2.38).
(K) Ensemble activity (z-scored $\Delta F/F_0$) averaged across push weighted neurons, locked to force
peak for push and pull actions. Inset: average activity at -0.5 s before force peak and at force
peak for each action. The activity of push weighted neurons increased at push force peak ("0 s")
relative to "-0.5 s" (Table S1, ref. 2.39). Activity at push force peak was greater than at pull force
peak (Table S1, ref. 2.40).
(L) Ensemble activity (z-scored $\Delta F/F_0$) averaged across pull weighted neurons, locked to force
peak for push and pull actions. Inset: average activity at -0.5 s before force peak and at force
peak for each action. The activity of pull weighted neurons increased at pull force peak ("0 s")
relative to "-0.5 s" (Table S1, ref. 2.41). Activity at pull force peak was greater than at push force
peak (Table S1, ref. 2.42).
(M) Same as (K) for D1- and D2-MSNs. Activity of push weighted D1- and D2-MSNs around
push force peak (-0.5 s to 0.5 s) is not significantly different (Table S1, ref. 2.43). Data are
mean+s.e.m. across 63 sessions from 8 mice.
(N) Same as (L) for D1- and D2-MSNs. Activity of pull weighted D1- and D2-MSNs around pull
force peak (-0.5 s to 0.5 s) is not significantly different (Table S1, ref. 2.44). Data are
mean+s.e.m. across 73 sessions from 8 mice.

Population activity at force peak predicted action identity even for trials when other actions were
allowed in the 0.5s temporal window before and after (Figure 2H). D1- and D2-MSNs encoded
these actions with equal accuracy (Figure 2I). Given that the striatum encodes continuous
kinematics[16,17], we verified action identity was encoded for trials with matched continuous
parameters including force, duration, and licking (Figure S3I,J).

To understand the structure of this encoding, we examined how action identity was embedded
within the high dimensional space of population activity where each neuron's instantaneous
activity is one axis. The SVM identifies a single dimension that best predicts the animal's action
at the time of the force peak. It follows that the projection of population activity on that SVM
dimension (a weighted sum over neurons) showed differential modulation at force peak for push
and pull actions (Figure 2J). The SVM's weights partition neurons into action-specific groups
whose activation biases the decoder to predict push or pull. The push-weighted neurons
increased activity at push force peak, but not for pull force peak (Figure 2K). Similarly, the pull
weighted neurons increased activity at pull force peak but not for push force peak (Figure 2L).
D1- and D2-MSNs showed no difference in SVM weights and action-specific modulation of
push- and pull-weighted neurons (Figure 2M,N).

The striatum has a classical role in the reinforcement of actions, with D1- and D2-MSNs thought
to have opponent[13,25] or different[26] roles. Therefore, we examined if encoding of action identity
would change whether the action is reinforced or not. We found that both types of MSNs encode
ongoing action identity regardless of whether both or one of the particular actions were
reinforced (Figure S2D-G). Further, we tested whether striatal plasticity is necessary for action
encoding. Notably, we found that striatal activity encoded the identity of specific actions even in
mice with impaired striatal plasticity[47] who could not learn which actions were reinforced[48–50]
(Figure S2H-J). These results reveal that although the striatum has a clear role in reinforcement
and learning, it can encode ongoing actions independently of these roles.

Together, these results demonstrate that striatal population activity robustly encodes the identity
of ongoing forelimb actions, with equal contributions from D1- and D2-MSNs.

Specific stimulation of striatal neurons encoding specific actions
We next addressed our central question: whether these ensembles causally control the specific
actions they encode. One critical technical challenge was that we needed to target perturbations
to specific and spatially-intermingled D1- and D2- MSNs that encode action identity. For this, we
developed an approach to stimulate individual MSNs with 2-photon optogenetics through a
GRIN lens[30] in mice co-expressing GCaMP6f and red-shifted activating opsin ChRmine[29]
(Figure 3A-C). We used a spatial light modulator to holographically target a group of user-
defined MSNs for simultaneous stimulation[28]. Another technical challenge was that action
duration was very short, and we needed to trigger stimulation upon self-initiated actions with low
latency, as force onsets rapidly over milliseconds (Figure 1C). For this, we developed an
experimental platform that allowed us to flexibly select the trigger action and stimulation pattern
and begin the stimulation with ~2ms total latency (Figures 3D and S4B).

Figure 3: Closed-loop modeling and manipulation of neuronal ensembles that encode action

identity.

(A) Schematic of GCaMP6f and ChRmine viral injection and lens implant in transgenic mice to

image and photostimulate D1- and D2-MSNs.

(B) Example field of view with a single neuron targeted for stimulation. Inset: zoom in onto

targeted neuron in black.

(C) Example activity traces of neurons in (B) inset. Targeted neuron (in black) and neighboring

non-targeted neurons' activity during 2-photon optogenetic stimulation.

(D) Schematic of closed-loop system to select the pattern of holographic stimulation and the

action that triggers stimulation (Figure S4B for more details).

(E) Cartoon depicting how activating specific neurons can manipulate a dimension of population

activity that encodes and predicts specific actions.

(F) Schematic for conducting the modeling and manipulation experiment.

(G) Left: example field of view with regions of interest labeled based on their decoder weights

and identification as D1-MSNs. Right: Decoder weights for identified D1-MSNs, colored by

assignment into action-specific ensembles.

(H) Z-scored fluorescence of each ROI member of each action-specific ensemble, and the

projection of population activity on the SVM decoder dimension. Four consecutive stimulation

trials concatenated from the beginning, middle, and end of the example session.

(I) Trial-averaged traces of targeted neurons and projection of population activity on the action-

predictive SVM dimension.

(J-N) Data are mean+s.e.m. across n=21 sessions and 10 mice.
(J) Average decoder weights of action-specific ensembles. The top four neurons with the largest
weights are shown.
(K) Average activity (z-scored fluorescence) of action-specific ensembles in 100 ms before force
peak during trials in the calibration block. Push ensemble activity was larger for push than pull
(Table S1, ref. 3.1). Pull ensemble activity was larger for pull than push (Table S1, ref. 3.2).
(L-M) Neural activity averaged across neurons (z-scored fluorescence) within each ensemble,
locked to stimulation of each ensemble. Inset: average ensemble activity in 100 ms after
stimulation.
(L) After push ensemble stimulation, activity of the push ensemble was greater than the pull
ensemble (Table S1, ref. 3.3) and non-targeted neurons (Table S1, ref. 3.4). There was no
significant difference between the pull ensemble and non-targeted neurons (Table S1, ref. 3.5).
(M) After pull ensemble stimulation, activity of the pull ensemble was greater than the push
ensemble (Table S1, ref. 3.6) and non-targeted neurons (Table S1, ref. 3.7). There was no
significant difference between the push ensemble and non-targeted neurons (Table S1, ref.
3.8).
(N) Projection of population activity on the SVM dimension locked to ensemble stimulation.
Inset: Average SVM projection in the 100 ms before stimulation ("pre") and after stimulation
("post"). Stimulating the push ensemble increased the "post" SVM projection relative to "pre"
(Table S1, ref. 3.9). Stimulating the pull ensemble decreased the "post" SVM projection relative
to "pre" (Table S1, ref. 3.10). The "post" SVM projection was greater after push ensemble
stimulation than pull ensemble stimulation (Table S1, ref. 3.11).
310
Within a single session, we identified action-specific MSN ensembles and stimulated them in
closed-loop during specific, self-paced actions (Figure 3E,F). In a calibration block, we collected
behavior trials and fit a decoder to predict action identity from MSN activity. As before (Figure
2K-N), we identified action-specific ensembles (Figure 3G) using the weights of the force peak
SVM. We defined the action-specific ensembles as equally sized groups of 4-11 neurons with
the largest decoder weights (mean ensemble size was 6.7 neurons over 21 sessions). Then in
the stimulation block, specific self-paced actions triggered stimulation of action-specific
ensembles (Figure 3H,I). We interleaved 6 conditions: two actions that triggered stimulation (3g
push, 3g pull) and three stimulation patterns for each action (push ensemble, pull ensemble, no
stimulation) (Figure 3F).
321
The identified ensembles had similar magnitude decoder weights (Figure 3J) and modulated
specifically for their corresponding actions (Figure 3K). Stimulation specifically activated the
targeted ensemble (Figure 3L,M). We verified that this stimulation of a small number of action-
specific neurons bi-directionally drove the action-specific SVM dimension of population activity
(Figure 3N). Thus, selective stimulation of small, action-specific striatal ensembles was
sufficient to bias population activity toward encoding the corresponding action (Figure 3E).
328
Specific striatal ensembles control specific actions
We used this approach to test whether action-specific striatal ensembles control specific actions
or simply invigorate any ongoing action. If ensembles control specific actions, stimulating a pull

ensemble should selectively increase pull force, with no effect or an opposing effect during
push. If ensembles merely invigorate movement, any ensemble stimulation should increase
force regardless of action identity. We refer to the pairing of an ongoing action with ensemble
stimulation specific to that action as "congruent" (e.g. pull ensemble stimulation during pull
action), and the pairing of an ongoing action with ensemble stimulation specific to the other
action as "non-congruent" (e.g. push ensemble stimulation during pull action) (Figure 3F). This
also allowed us to test whether D1-MSNs promote and D2-MSNs suppress action, or both
promote action.

340
Figure 4: Action-specific ensembles of D1- and D2- MSNs control congruent actions.
(A-B) Trial-averaged force traces locked to 3g cross that triggers holographic stimulation. In this
plotting convention, force is positive for the action executed at time 0 (push or pull) and negative
for the other action.
(A) Stimulation of D1-MSN ensembles. Data are mean+s.e.m. across n=12 sessions and 9
mice.
(B) Stimulation of D2-MSN ensembles. Data are mean+s.e.m. across n=9 sessions and 5 mice.
(C-D) Force perturbation, i.e. the trial-averaged force trace on stimulation trials minus non-
stimulation trials. Right panel is the force perturbation at the end of stimulation, averaged from
100ms to 110ms.
(C) Force perturbation from stimulation of D1-MSN ensembles. The perturbation was greater
than 0 with congruent stimulation (Table S1, ref. 4.12) and was greater than non-congruent
stimulation (Table S1, ref. 4.11) at the end of stimulation. The perturbation was not significantly
different than 0 with non-congruent stimulation (Table S1, ref. 4.13). The perturbation first
became significantly different between congruent and non-congruent stimulation at 40ms after
stimulation onset (Table S1, ref. 4.5).
(D) Force perturbation from stimulation of D2-MSN ensembles. The perturbation was greater
than 0 with congruent stimulation (Table S1, ref. 4.25) and was greater than non-congruent
stimulation (Table S1, ref. 4.24) at the end of stimulation. The perturbation was not significantly
different than 0 with non-congruent stimulation (Table S1, ref. 4.26). The perturbation first
became significantly different between congruent and non-congruent stimulation at 100ms after
stimulation onset (Table S1, ref. 4.24).
363

Stimulating specific D1-MSN ensembles congruent with the ongoing action increased ongoing
force (Figure 4A left). In contrast, no change in ongoing force was observed after stimulating the
non-congruent ensemble (Figure 4A right). Notably, stimulating specific D2-MSN ensembles
congruent with the ongoing action also increased ongoing force (Figure 4B left), whereas
stimulation of the non-congruent ensemble did not (Figure 4B right). In other words, stimulation
of the same ensemble during different ongoing actions had different effects (e.g. stimulating the
pull ensemble increased force during pull, but not during push).
371
We quantified the force perturbation as the difference between force traces in stimulation trials
and non-stimulation trials. Stimulation of action-specific ensembles increased congruent but not
non-congruent force for both D1- and D2-MSNs (Figure 4C,D). Congruent stimulation of D1-
MSN ensembles increased perturbation relative to non-congruent stimulation as early as 40 ms
after start of stimulation, whereas D2-MSNs ensemble stimulation increased perturbation later,
around 100 ms after start of stimulation. Thus, the dynamics of the stimulation of D1- vs D2-
MSNs were slightly different, suggesting that D1- and D2 populations do not have a redundant
role. Together, these results demonstrate that specific ensembles of both D1- and D2-MSNs
causally control specific ongoing actions, increasing force selectively for the action they encode.
381

Discussion
Here we show that striatal ensembles that encode specific ongoing actions can indeed control
them. Further, this control operates at the granularity of distinct muscle activation patterns of the
same forelimb. We identified and stimulated small, action-specific ensembles of D1- and D2-
MSNs during self-paced isometric actions and found that stimulation selectively increased force
for the congruent action. Despite the striatum's indirect access to spinal circuits, these results
establish it as an online controller of specific actions and forces — a role beyond its classical
contributions to invigoration and reinforcement. To achieve this, we developed a model-guided,
all-optical[28–30,32] closed-loop system to identify and manipulate action-specific ensembles in real
time, even deep in the brain — addressing a critical challenge in motor neuroscience: linking the
encoding properties of specific neural ensembles to their role in the execution of movement.
393
These results offer new views on the canonical model that D1- and D2-MSNs respectively
promote and suppress movement[3,9,18,33] via the striatonigral and striatopallidal pathways. While
bulk manipulations have supported opposing effects[13,18], subsequent manipulations[40,41]
suggested that their coordinated activity are necessary for movement execution. Here we
identified and stimulated action-specific ensembles of D1- and D2-MSNs and found that both
increase the force of congruent actions. Thus, D2-MSNs can not only promote movement, but
also control specific actions in real time, online. This finding suggests that D2-MSN projections
to GPe can have effects that are independent of canonical inhibition of GPi/SNr, possibly
through different anatomical circuits linking GPe directly to thalamus, cortex and brainstem[51]. It
is also important to note that we focused our recordings and manipulations on forelimb region of
dorsolateral striatum, and understanding the function of D1- and D2-MSNs would likely depend
on the striatal subregion[16,33,52], each receiving distinct topographic inputs from cortex[42,43] and
thalamus, and also projecting to specific outputs. Finally, the faster onset of perturbations

evoked by D1-MSN congruent stimulation suggests different rather than opposing roles for D1-
and D2-MSNs in the control of movement.
409
This study positions the striatum as a center for controlling granular ongoing actions, despite its
indirect connections to the spinal cord. Although basal ganglia outputs tonically suppress
downstream motor centers[27,53–55], recent work reveals highly specific output dynamics during
forelimb behavior[56]. Our work provides a foundation for future work on a fundamental problem:
to understand how action-specific ensembles in the center of the brain can have such
remarkably specific control over downstream motor centers. Conceptually, we hypothesize that
the striatum guides motor centers into specific subspaces of neural population dynamics which
implement action plans and execution[57], consistent with evidence that striatal activity[58] and
corticostriatal plasticity[48] are required for the brain to learn and control low-dimensional cortical
dynamics[59–61] that directly operate brain-machine interfaces. More generally, our experimental
approach can be applied to motor circuits across the brain to test not only what neurons encode
but also what their activity causally produces[62] — both in terms of downstream population
dynamics and movement.
423
424 The discovery that striatal ensembles control granular actions offers a framework for
425 understanding movement disorders with striatal dysfunction characterized by impaired, highly
426 specific motor patterns—such as task-specific dystonias[63] (e.g. writer's cramp, musician's
427 dystonia) and the choreic disruptions of selective movements observed in Huntington's
428 disease[2,9].
429

Resource Availability
430
Lead contact
431
432 Further information and requests should be directed to the lead contact, Rui M. Costa.
433

Materials availability
434
435 No unique reagents were generated in this study.
436

Acknowledgements
437
We thank A. Vaz, M. Correia and G. Martins for mouse colony management. We thank the
members of the Scientific Hardware Platform of the Champalimaud Institute for help with
hardware and software. We thank H. Rodrigues for help with experimental setups. We thank T.
Tabachnik and R. Hormigo for technical advice and hardware support. We thank L. Hammond
and the Zukerman Institute's Cellular Imaging Platform for guidance in imaging and analysis.
We thank A.M. Vicente for blinding the experimenter and discussions about mouse training. We
thank Bruker and M.J. Fox for help developing the Bruker Software API. We thank K.
Deisseroth, Y.S. Kim, and C. Ramakrishnan for providing the ChRmine virus for holographic
stimulation. This work was supported by Fundacao para Ciencia e Tecnologia (FCT
PD/BD/105950/2014) to I.R.-V. This work was supported by BRAIN Initiative National Institute of
Mental Health postdoctoral fellowship (1F32MH118714-01) and NIH NINDS Pathway to
Independence Award (1K99NS128250-01) to V.R.A. This work was supported by NS
(NS104649), the Aligning Science Across Parkinson's (ASAP-020551) through the Michael J.

Fox Foundation for Parkinson's Research (MJFF) and Simons-Emory International Consortium
(Simons 717104) to R.M.C, and by the National Institutes of Health (NIH, U19: 5U19NS104649)
to D.S.P and R.M.C.
454

455 **Author contributions**
456 I.R.-V., V.R.A., D.S.P. and R.M.C. designed the study, interpreted results and wrote the
457 manuscript. I.R.-V. performed experiments. I.R.-V. and V.R.A. analyzed data and developed the
458 model-guided system for closed-loop stimulation.
459

Declaration of interests
D.S.P. is listed as an inventor of the following patent: "Devices, apparatus and method for
providing photostimulation and imaging of structures" (United States Patent 9846313). He also
receives income as a consultant to the Bruker Corporation related to the above patent.

I.R.-V., V.R.A., and R.M.C. declare no competing interests.

STAR Methods
Experimental model and study participant details
Animals
All experiments and procedures were performed according to National Institutes of Health (NIH)
guidelines and approved by the Institutional Animal Care and Use Committee (IACUC) of Columbia
University. Adult mice, aged 2–6 months, were used for these experiments. For labeling D1-MSNs and
D2-MSNs we used the following transgenic lines (Table 1): Tg(Drd1-cre)EY217Gsat (RRID:MGI:4366805),
Tg(Drd1-cre)EY217Gsat crossed with Ai9(RCL-tdT) (RRID:IMSR_JAX:007905), Tg(Adora2a-
cre)KG139Gsat (RRID:MMRRC_036158-UCD), Tg(Adora2a-cre)KG139Gsat crossed with Ai9(RCL-tdT)
and Drd1a-tdTomato line 6 (RRID:IMSR_JAX:007905). For experiments testing the necessity of plasticity in
the striatum we used the following lines (Table 1): RGS9-cre (RRID:IMSR_JAX:020550) crossed with
NMDAR1-loxP (RRID:IMSR_JAX:036352) [47]. For comparison, we used the negative Cre littermates as
controls. Mice used for experiments were individually housed and kept under a 12-h light-dark cycle.

Line name	Extended name	Source	Available at	Code	Ref.
Tg(Drd1-cre)EY217Gsat	STOCK Tg(Drd1-cre)EY217Gsat/Mmucd	Gerfen lab*	MMRRC	030778	Gong et al, 2007
Ai9(RCL-tdT)	B6;129S6-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J	Jackson Laboratories	Jackson Laboratories	007905	Madisen et al, 2010
Drd1a-tdTomato line 6	B6.Cg-Tg(Drd1a-tdTomato)6Calak/J	Jackson Laboratories	Jackson Laboratories	016204	Ade et al, 2011
Tg(Adora2a-cre)KG139Gsat	B6.FVB(Cg)-Tg(Adora2a-cre)KG139Gsat/Mmucd	MMRRC*	MMRRC	036158	Gong et al, 2007
RGS9-cre	B6;129S-Rgs9$^{tm1.1(cre)Yql}$/J	NIH from Yuqing Li*	Jackson Laboratories	020550	Dang et al, 2006
NMDAR1-loxP	B6.129S2(Cg)-Grin1^{tm1Yql}/NkzaJ	NIH from Yuqing Li*	Jackson Laboratories	036352	Dang et al, 2006

478 * These lines were kept in house and backcrossed for more than 10 generations.
479 Table 1 – Mouse lines used in experiments.
480

481 **Randomization and blinding**
482 The experimenter was blinded to the experimental groups for plasticity experiments.
483
484 All animals were run on each day in a fixed random order.
485
486 Holographic manipulations were performed according to a pre-defined protocol and controlled by Matlab
487 in closed-loop with the mouse's behavior - not manually by the experimenter. Sessions for stimulating D1-
488 and D2-MSNs were interleaved as much as possible.
489
490 **Method Details**
491 **Stereotaxic Surgery**
492 Surgical procedures for imaging of neurons in the dorsolateral striatum (DLS) were performed in two
493 steps consisting of viral injection at least 5 days prior to lens implant, following protocols described in
494 protocols.io ("Viral injection" and "Lens implant").
495
496 Before each surgery, analgesia was administered to ensure 72 hours of pain relief. The mouse was
497 placed in a stereotaxic frame under isoflurane anesthesia on a heated pad.
498
For viral injection, capillary tubes (Drummond 3.5" #3-000-203-G/X) were pulled using a Pipette puller
(Sutter P-2000; Heat 360, FIL 5, Vel 25, Del 10, Pul 50) and backfilled with mineral oil. A craniotomy was
made using a dental drill, and the capillary tube was lowered to the target coordinates. For imaging, 300
nl of AAV5.CaMKII.GCaMP6f (titer: 1×10^{12}–1×10^{13} GC/ml) was injected into the left DLS (AP: +0.75 mm,
ML: 2.5 mm, DV: 2.4 mm) at a rate of ~5 nl/pulse. For holographic optogenetics experiments, an
additional 300 nl of AAV8.CaMKII.ChRmine (titer: ~1×10^{11} GC/ml) was co-injected with
AAV5.CaMKII.GCaMP6f at a 1:1 ratio into the left DLS at the same coordinates. Following injection, the
capillary tube was left in place for at least 5 minutes before slow withdrawal, and the skin was sealed with
sutures.
508
509 At least 5 days after viral injection, a 1 mm GRIN lens implant surgery was performed. Following
510 craniotomy and cortex removal – using vacuum aspiration with 1 mm blunt needle – the GRIN lens was
511 implanted at the same coordinates as the viral injection at depth of DV: 2.0 mm. A head bar was secured
512 and a protective cap was built around the lens, which was optionally covered with silicone sealant (Kwick-
513 Sil or Kwick-Cast) for protection.
514
For experiments requiring electromyographic (EMG) recordings of forelimb muscle activity, we fabricated
EMG electrodes in-house as previously described[64] and in protocols.io ("EMG fabrication"). Briefly, two
pieces of insulated braided stainless-steel wire were cut, knotted together, and secured to a working
surface. Under a microscope, 0.5 mm portions of insulation were stripped from each strand just below the
knot, leaving exposed contact sites separated by 0.5 mm. The wire ends were inserted into hypodermic
needles and crimped in place to facilitate muscle insertion, and the opposing ends were soldered into
corresponding sites on a miniature Omnetics connector.
522
EMG electrodes were then implanted into forelimb muscles (biceps, triceps, palmaris longus, and
extensor digitorum communis) following lens implant[36], as described in protocols.io ("EMG implant"). Prior
to surgery, mice were habituated to an Elizabethan collar and gel food for 4 days to minimize post-
surgical disruption. On the day of surgery, analgesia was administered as described above. Electrodes
were inserted into each target muscle using the hypodermic needle as a guide and secured in place by
knotting the wire at the exit point to fix the recording sites within the muscle. Each incision was closed
with sutures and covered with antibiotic ointment. The EMG connector was attached to the back of the

skull or existing head cap using dental cement (Metabond), and any neck incision was similarly closed
with sutures and antibiotic ointment. After surgery, the Elizabethan collar was placed on the mouse for at
least 5 days. Food gel and soften pellets were provided for at least 5 days post-surgery.
533
534 **Histology**
535 Histology was performed as described in protocols.io (Histology) to verify lens coordinates and viral
536 expression. Cardiac perfusion used 4% PFA solution to ensure fixation of the brain tissue. At least 24
537 hours after the cardiac perfusion, the brain was moved into 0.05% Thimerosal-PBS and stored in the
538 fridge until histological processing.
539
Brains were sliced coronally at 50 µm using a vibratome (Leica VT1000) and collected into 1X PBS.
Immunohistochemistry was performed as follows:
• To localize the imaging site, sections were incubated with an Alexa Fluor 488-conjugated GFP
antibody at 1:1000 in 0.4% Triton-PBS overnight at room temperature.
• To identify tdTomato somatic expression, sections were incubated with a primary anti-RFP
antibody at 1:2000 in 0.4% Triton-PBS overnight at 4°C, followed by a secondary anti-Rabbit
Alexa Fluor 647 antibody at 1:1000 in 0.4% Triton-PBS for 2 hours at room temperature.
547
548 All sections were counterstained with DAPI at 1:1000 in PBS for 15 minutes and then mounted on glass
549 slides with Mowiol solution, allowed to dry overnight, and sealed with nail polish.
550
Sections were imaged using a custom-built automated slide scanner equipped with a 4x 0.4-NA Plan Apo
objective (Nikon Instruments Inc.) and P200 slide loader (Prior Scientific), controlled by NIS-Elements
using custom acquisition scripts, from the Zuckerman Institute's Cellular Imaging Platform. For detection
of DAPI, Alexa Fluor 488, and Alexa Fluor 647, the following excitation wavelengths and filters were used,
respectively: 405 nm (filter 435/26 nm), 488 nm (filter 525/30 nm), and 647 nm (filter 705/72 nm). Image
processing and analysis were performed using Brain J (RRID:SCR_027061).
557
Two-action isometric task
Setup
Mice were head-fixed and placed in a 3D-printed cup[65] with a sucrose spout placed within tongue reach.
The right forelimb interacted with an immobile joystick (3 mm screw) mounted on 3D-printed frames
attached to two load cell sensors (Phidgets 3139_0), one measuring push/pull and the other measuring
left/right forces. The left forelimb rested on an auxiliary pole equipped with two additional load sensors
measuring push/pull and vertical forces. The cup holding the mouse was mounted on a fifth load sensor
to capture overall body movement.
566
567 Signals from all load sensors were recorded at 1 kHz using the Champalimaud Foundation Scientific
568 (CFS) "Scientific Board v1.3" with "Quad Load Cell v1.2" module, which also detected threshold crossings
569 based on experimenter-defined parameters. Data were visualized and stored using the CFS "Load Cells
570 Visualizer" software.
571
A pyControl "Lickometer v1.0" board detected licking and contact with the joystick and auxiliary pole. Task
control and sucrose delivery were managed by a pyControl[66] "Breakout 1.2" board, which logged events
at 1 kHz and opened solenoid to dispense the reinforcer - 5 µl of 10% sucrose solution. Sucrose flowed
via gravity through tubing and was delivered through a blunt 16G needle spout positioned near the
animal's mouth. Solenoid timing was calibrated before each session, and the sucrose solution was
prepared fresh weekly. Hardware schematics are shown in Figure S4a.
578

Training
Mice were habituated to head fixation for 15 min on a running wheel over four days and then food-
restricted to 80–85% of initial body weight. Training for the two-action isometric task was performed as
described in protocols.io ("Behavior training"). Mice pushed or pulled an immobile, pressure-sensitive
joystick in a self-paced manner to receive reinforcer. A task action was defined as a force exceeding a
task-defined threshold for longer than the task-defined duration. Thresholds, durations, and reinforcement
delays were progressively adjusted during training – these are referred to as "Task Parameters" (Table 2).
Reinforcement followed the end of a valid action after a delay. A 3 g quiescence threshold was required
for trial initiation.
588

Reinforcement block	Parameters label	Task action threshold (g)	Task action duration (ms)	Reinforcement delay (ms)
Both	1	4	10	100
Both	2	4	20	200
Both	3	6	20	300
A or B	1	4	20	300
A or B	2	6	20	300
A or B	3	6	50	300

589 Table 2 – Changes in task parameters for different reinforcement blocks.
590
591 Each trial consisted of: (1) a 1 s inter-trial interval (ITI); if forces were below 3 g at 1 s, a new trial began,
592 otherwise the ITI reset; (2) a trial period, during which a task action (push or pull exceeding the threshold
593 for the task duration) triggered delayed sucrose delivery. Trials terminated upon reinforcement or when
594 force fell below task-defined threshold after an action that did not meet task-defined duration.
595
Training comprised three sequential blocks with distinct reinforcement contingencies. In Block Both, push
and pull actions were both reinforced, up to 50 reinforcements per action per session. On the final
session of Block Both, the less frequent action was designated action A and the more frequent as action
B. In Block A, only action A was reinforced – starting after last session of Block Both; in Block B, only
action B was reinforced – starting after last session of Block A. Block Both sessions ended after 50
reinforcements of each action or 30 min. Blocks A and B ended after 200 reinforced actions or 30 min.
Task parameters were adjusted during training, advancing according to Table 2 when mice executed >50
reinforced task actions in a session. Block Both concluded when mice exceeded 50 reinforced actions in
two consecutive sessions at Parameters 3. Blocks A and B each concluded when the reinforced action
comprised ≥80% of total actions - defined as reinforced task actions divided by total action A and action B
executions - over three consecutive sessions at Parameters 3. Conclusion of Block B marked the end of
training, after which animals were returned to ad libitum feeding.
608
609 For Rgs9-LCre::Grin1tm1Yql homozygous mice (Striatal NR1-KO) and Cre-negative control littermates,
610 the same training protocol was used, with two modifications. Each block was capped at 1.2× the number
611 of sessions required by the third fastest mouse in the cohort to reach Parameters 3 (Table 2). Animals
612 were also advanced to the next block if they met give-up criteria, defined as fewer than ten 3g threshold
613 crossings (≥50 ms duration) averaged over three consecutive sessions.
614
615 **Selected sessions for analysis of behavioral training**
616 Because animals were trained to criterion, the number of sessions per block varied across individuals. To
617 enable cross-animal comparisons, specific sessions were selected for analysis. In Block Both, four

618   sessions were chosen, and in Blocks A and B, five sessions were chosen, following the selection rules
619   detailed in Table 3. These sessions were used in Figures S2 and S3.
620
621

Reinforcement block	Selected session	Selection Rules
Both	1	1st day at Parameters 1 for Block Both
Both	2	1st day at Parameters 2 for Block Both
Both	3	1st day at Parameters 3 from Block Both
Both	4	Last day before change in reinforcement schedule
A or B	1	1st day at Parameters 1 for Block A or B
A or B	2	1st day at Parameters 2 or Block A or B
A or B	3	1st day at Parameters 3 or Block A or B
A or B	4	Middle day between selected session 3 and 5
A or B	5	Last day of current reinforcement Block

622   Table 3 – Selected sessions for comparison across animals for all reinforcement blocks.
623

Adaptive Isometric task
Setup
Mice were trained on this task to test causal striatal control of specific ongoing actions. To do that, we
developed a modified version of the two-action forelimb task that promoted balanced performance of both
actions. Reinforcement probability was adjusted online based on recent performance: the proportion of
push and pull actions was tracked over the last 10 trials. When both actions occurred in fewer than 70%
of recent trials, both were reinforced; if one action exceeded 70%, that action was unrewarded.
631

The setup was otherwise identical to the two-action isometric task (Figure S4A), with additional hardware
for closed-loop holographic stimulation (Figure S4B). This system required flexibility to: (1) select, on a
single-trial basis, the threshold crossing to trigger stimulation; (2) trigger laser with millisecond latency;
and (3) specify the holographic 2-photon pattern, on a single-trial basis. Features 1 and 2 required an
improved hardware system for processing force sensors, while feature 3 required a software update to
the two-photon microscope's stimulation interface.
638

Features 1 and 2 were accomplished using an updated load-cell system ("HARP Load Cells Acquisition
v1.1" and "HARP Load Cell Interface v1.1," CFS Hardware Platform), which evaluated in parallel when
the force exerted on the joystick crossed six experimenter-defined thresholds. In the holographic
experiments, six thresholds were defined: 3 g, 4 g, and 6 g for push and pull actions. To select which
threshold would trigger stimulation on a given trial, and to restrict the trigger signal to specific time
windows within each trial, a custom circuit board consisting of a MUX and an AND gate was built to route
the appropriate trigger signal to the microscope while minimizing the delay between threshold crossing
and stimulation onset. The MUX and AND gate was controlled by custom MATLAB scripts running on the
microscope control computer, which determined which force threshold should serve as the trigger on
each trial and when to enable it. Digital communication between the task boards, the computer, and the
MUX and AND gate was implemented using a NIDAQ DIO board (USB-6001).
650

651   Feature 3 was accomplished through a newly implemented function in the PrairieLink API of the Bruker
652   microscope, which allowed two-photon stimulation patterns to be specified dynamically during an ongoing
653   imaging experiment. Schematics of the full setup, including both the behavioral and stimulation hardware,
654   are presented in Figures S4.
655

656   *Training*
657   This task was used for holographic stimulation experiments and followed the two-action forelimb task with
658   modified reinforcement to promote balanced performance of both actions. Mice were trained for 3 weeks
659   with adaptive reinforcement and no stimulation. Training was conducted partly in behavior boxes in the
660   absence of imaging, prior to head-fixed sessions under the microscope.
661

2-photon microscope
The optical setup comprised two femtosecond pulse lasers coupled to a custom-modified two-photon
laser scanning microscope (Ultima In Vivo, Bruker Corporation, Billerica, Massachusetts) equipped with
an 8 kHz resonant scanner. The laser source for imaging was a pulsed Ti:sapphire laser (Chameleon
Vision-S, Coherent, Inc., Saxonburg, Pennsylvania) tuned to 920 nm for GCaMP6f imaging, with power
controlled by a Pockels cell (350–105-BK, 302RM controller, Conoptics, Inc., Danbury, Connecticut). The
laser beam was expanded by a 1:2 telescope (Thorlabs GBE02-B) and further scaled in the microscope
by a 1:1.33 telescope before being coupled into a scan lens (f = 75 mm), tube lens (f = 180 mm), and
objective lens (16X 0.8NA N16XLWD-PF, Nikon Corporation, Tokyo, Japan). The laser could also be
directed to a non-resonant scanning path where both X and Y scanning were controlled by galvanometric
mirrors. The fluorescence signal was collected through the objective lens, split from the IR beam with a
670 nm long-pass dichroic (ZT670rdc, Chroma Technology Corp., Bellows Falls, Vermont), and coupled
into a liquid light guide assembly. The light was split by a 565 nm long-pass dichroic mirror (HQ565dcxr,
Chroma Technology Corp.), with the green and red channels each directed to GaAsP PMTs through
510/20 and 607/45 bandpass filters, respectively (Chroma Technology Corp.).
677

The photostimulation path was largely independent from the imaging path. The laser source for
photostimulation was a low repetition rate (1 MHz) amplified laser (Monaco 1035–40, Coherent, Inc.,
Saxonburg, Pennsylvania) operating at 1035 nm, with power controlled by an integrated acousto-optic
modulator. The beam was expanded by a 1:2 telescope (Thorlabs GBE02-B) and directed into the
microscope, where it could be steered either by a dedicated pair of galvanometers or coupled to a
customized light path including a spatial light modulator (Bruker Neurolight 3D, Billerica, Massachusetts)
for holographic stimulation. The photostimulation beam was combined and made coaxial with the imaging
beam just before the scan lens via a 1030 nm short-pass dichroic (T1030SP, Chroma Technology Corp.).
686

687   Imaging and photostimulation were controlled by PrairieView (Bruker Corporation, Billerica,
688   Massachusetts). Custom software defined stimulation patterns and triggered closed-loop photostimulation
689   based on behavioral events - MATLAB (The Mathworks, Inc., Natick, Massachusetts) and Python,
690   interfaced with PrairieView via the PrairieLink API.
691

692   **2-photon imaging**
693   Before each experimental cohort, the imaging and stimulation laser powers were measured at the output
694   of 16X Nikon objective and confirmed stable across months. A custom computerized goniometer was
695   used to precisely and reproducibly angle the head of the mouse such that the imaging lens was
696   orthogonal to the beam path.
697

698   2-photon imaging was performed as described in protocols.io ("2-photon imaging"). Prior to each session,
699   laser alignment was confirmed at the objective output. The mouse was head-fixed in the behavioral setup
700   under the microscope, and the top of the GRIN lens was located and cleaned if necessary. Imaging power
701   was continuously monitored using a beam pickoff (Thorlabs BSF10-B) and power meter (Thorlabs
702   PDA100A2), with the Pockels cell bias adjusted accordingly throughout the session.
703

704   The typical imaging power was <50 mW, and could be up to 80 mW for imaging deeper than ~250 µm
705   below the terminus of the GRIN lens. Images were acquired at 30 Hz over a 512 × 512 pixel field of view.
706   For Figures 1 and 2, a 1.5x zoom was used over 496.1 µm × 496.1 µm; for Figures 3 and 4, a 1.5x zoom
707   over 550.4 µm × 550.4 µm and a 2x zoom over 412.7 µm × 412.7 µm were used.

708
709   For cell type identification, 1000 frames of functional and structural images were acquired using green
710   and red PMTs respectively. These data were then processed as described below (STAR methods –
711   Distinguishing D1- and D2-MSNs). Structural images and z-stacks were acquired at the end of each
712   session using both green and red channels to confirm ROI identity (Figure S1).

713
714   Most mice were imaged for the entire duration of training, with the same field of view tracked across
715   sessions. ROIs identified in previous sessions were used to help locate the same field of view, and
716   custom code was written to track the same ROIs across days.

717
718   **Protocol to image the same field of view and neurons across sessions**
719   For holographic stimulation experiments, the same field of view and neurons were imaged across paired
720   sessions. After the first session, imaging data were processed with Suite2p[67] to extract active ROIs. In the
721   paired session, PrairieView's "Brightness over Time" (BOT) feature was used to annotate these neurons
722   in the GUI. XYZ coordinates were adjusted to best match the previous session's field of view.

723
Calibration of 2-photon stimulation spatial targeting and power
When doing holographic stimulation, the spatial targeting of 2-photon stimulation was calibrated before
each session (protocols.io, "2-photon stimulation calibration"). Using PrairieView's "burn spots" routine,
we iteratively aligned burned spots on a fluorescent sample to target locations, allowing the software to
calibrate galvos' control for accurate stimulation. Calibration was stable across sessions, requiring only
minor adjustments. This calibration was done using the same objective used for the experimental
recordings.

731
732   Before experiments, we calibrated stimulation power and duration for each ROI. Using custom Python
733   scripts with PrairieView's "BOT" feature, we visualized ROI responses while sweeping power (3–7 mW
734   per target) and duration (50-100 ms) to identify the minimum parameters that reliably activated the
735   targeted neurons.

736
737   **Holographic optogenetics experiments**
738   We performed holographic optogenetics experiments to test the effect of stimulating specific neuronal
739   ensembles on self-paced forelimb actions during the Adaptive Isometric task.

740
Experiments relied on pairs of sessions in which we imaged the same field of view and neurons. On the
first session, we performed two-photon imaging as mice performed the task, used Suite2p to extract
functional ROIs, and automatically labelled ROIs as D1- or D2-MSNs. On the second session, we
performed the stimulation experiment, consisting of a calibration block followed by a stimulation block
(Figure 3F). For each stimulation session, cell type choice (D1- or D2-MSNs) was determined by
balancing the number of sessions per cell type and the number of neurons available for stimulation.

747
Within the calibration block, we performed two-photon imaging as mice performed the adaptive isometric
task for up to 30 minutes. Using PrairieView's "BOT" feature, we pre-defined ROIs identified from the
previous session and rapidly extracted their fluorescence, averaged over the pixels within a circle defined
by each Suite2p mask's center and radius. We extracted behavioral data and identified force threshold
crossings of 3 g (hereafter "3g cross actions"). For either D1- or D2-MSNs, we z-scored each ROI's

fluorescence across the recording and fit an SVM to predict push versus pull using single trials of neural
activity averaged over the 5 imaging frames centered at force peak. If SVM accuracy on test trials
exceeded 55%, we used the SVM weights to identify push- and pull-weighted ensembles with equal
numbers of neurons, and plotted their activity time-locked to the force peak of 3g cross actions. If the
push (pull) ensemble showed greater activity for push (pull) actions, we proceeded to holographic
stimulation.
759
We then defined a software command via the PrairieLink API to specify holographic stimulation for each
ensemble. Stimulation was targeted to the center of each ROI in the ensemble, with total power
calculated as the product of the number of neurons per ensemble and the desired average power per
neuron. Stimulation duration was typically 100 ms - sufficient to reliably activate neurons expressing
ChRmine. To cover each ROI's circular extent, dedicated galvos traced 5 spirals per stimulation with a
diameter matching the average ROI diameter. Stimulation was configured to be delivered upon a trigger
signal corresponding to a force threshold crossing.
767
Within the stimulation block, we performed two-photon imaging and closed-loop holographic stimulation
as mice performed the adaptive isometric task for 30 minutes. We pre-defined a schedule interleaving
conditions based on which action triggered stimulation (3g cross push or pull) and which stimulation
pattern was delivered (push ensemble, pull ensemble, or no stimulation). Trials were organized into
alternating blocks of 6, triggered by one action (either 3g cross push or 3g cross pull), with the following
stimulation sequence per block: (1) no stimulation, (2) push ensemble, (3) no stimulation, (4) no
stimulation, (5) pull ensemble, (6) no stimulation. A 2-second inter-trial interval was imposed between
trials.
776
Our custom setup (Figures 3 and S4B) enabled selection of which force threshold crossing served as the
trigger on interleaved trials and stimulation delivery within milliseconds of threshold crossing — critical
given the rapid timescale of force changes. As described in the setup section above, the microscope
computer ran custom MATLAB code to step through the scheduled trials, using the PrairieLink API to
send stimulation commands and a NIDAQ DIO board (USB-6001) to select and enable force thresholds
via the MUX and AND gate. The trigger signal output from the MUX was also received by the computer to
track stimulation delivery and advance to the next inter-trial interval and trial.
784
EMG recordings
EMG recordings were performed as described in protocols.io ("EMG recordings"). Custom-made
electrodes were implanted in forelimb muscles and connected to an Omnetics connector. Prior to each
session, mice were head-fixed in the behavioral setup under the microscope and the 16-channel ZC16
headstage (Tucker-Davis Technologies – TDT) was connected to the implanted connector. EMG
recordings were set to true differential mode in Synapse recording software, and the signal quality was
confirmed when the mouse moved. Signal quality was optimized by grounding the mouse to the air table.
Signals were acquired via the ZC16 headstage connected to a TDT RZ5D Bioamplifier and PZ5-32
preamplifier, and recorded at 24 kHz. For analysis, EMG signals were downsampled to 1 kHz, high-pass
filtered at 40 Hz, rectified, and smoothed with a Gaussian kernel (SD = 25 ms)[68]. Data included 41
recording sessions from 4 mice performing the two-action isometric task (Figure 1).
796
797 **Quantification and Statistical Analysis**
798 Analysis was performed using custom scripts in Python.
799
800 **Data extraction and synchronization**

801 All data streams—task events (1 kHz), joystick forces (1 kHz), EMG (24 kHz), 2-photon imaging/voltage
802 (30 Hz/1 kHz) - were synchronized via the ITI signal, providing a unified timestamp across modalities for
803 precise alignment of behavior, forces, EMG and imaging.
804
Calibrated force measurement
The joystick voltage signal was converted to force using a linear model:
Force (g) = (Voltage – baseline) × conversion factor
Baseline voltage was measured before each session and defined as the average voltage with no weight
on the sensors. The conversion factor was determined by applying known weights (0, 2, 5, 20 g) and
fitting a linear model. Conversion was stable and re-calibrated weekly to correct drift.
811
Definition of actions
For analysis, we defined several actions from force and touch signals. Task actions were push and pull
events identified by the hardware from 1 kHz force traces, with intervals from threshold rise to fall. 3 g
cross actions were pushes or pulls exceeding 3 g for ≥66 ms with joystick touch ≥100 ms before crossing
to ensure isometric behavior; force traces were downsampled to 30 Hz. 6 g cross actions were a subset of
3 g crosses peaking above 6 g, the final training threshold. Touch was defined as joystick contact <2 g for
>1 s, and licks as ≥5 licks with <0.5 s inter-lick interval, excluding overlap with 3 g crosses. We also
analyzed special subsets: pre-isolated 3 g crosses had no other 3 g crosses in the 0.5 s before, post-
isolated 3 g crosses had none in the 0.5 s after, and matched actions were push/pull trials matched for
continuous-valued features (STAR Methods - Matching trials across actions).
822
823 **Action events**
824 For push and pull actions, analysis was aligned to two key events within the action interval: the moment
825 the force threshold was crossed (rising edge) and the moment of peak force. For touch and lick actions,
826 analysis was aligned to the midpoint of the action interval.
827
828 **Matching trials across actions**
829 We tested whether muscle and neural activity encoded push and pull actions using trials matched for
830 average features (Figures 1F-H and S3). Tolerances were: average force ≤0.1 g (Figure 1F-H); peak force
831 ≤0.1 g, two-axis force magnitude at peak ≤0.2 g, action duration ≤30 ms, and lick rate/probability during
832 actions and at peak ≤0.5 Hz and ≤0.1, respectively (Figure S3).
833
Trials were iteratively removed to achieve matching while minimizing loss of trials. Features were z-
scored, then for each iteration: (1) define action 1 as the one with fewer trials, action 2 as the other; (2)
compute the target vector as the difference in mean features across actions; (3) zero in-tolerance features
(i.e. set the difference to zero for each feature that is within-tolerance); (4) compute dot products of action
2 trials with the target; (5) remove the trial with the smallest dot product; (6) repeat until all features met
tolerance.
840
841 For Figure S3, trials were pre-selected such that the time between solenoid opening and force threshold
842 crossing was at least 4 seconds, **ensuring that analyzed actions did not occur during sucrose**
843 **consumption**.
844
845 **Time course of action force**
846 Trial-averaged joystick force was computed as peri-event time histograms (PETHs) locked to action
847 events: 6 g cross push/pull actions were aligned to force peak (Figure 1C). In Figure 2F, -4–0 s shows
848 pre-isolated and 0–4 s shows post-isolated 3 g cross force, illustrating trials used to predict action identity
849 from neural activity.

Quantification of actions across reinforcement

Action performance was quantified across the reinforcement schedule. For task actions, we quantified action rate for each session (Figure S2B,C). For 6 g cross actions, we quantified action rate pooled over sessions in each reinforcement block (Figures 1D). For trials matched across 3g cross actions, we quantified number of actions, peak and average force, two-dimensional force magnitude, average lick probability, action duration, action rate, and action proportion (Figure S3A-H).

For Striatal NR1-KO mice, we quantified 6g cross actions with the overall action rate for each session (Figure S2H), and the action rate pooled over sessions in each reinforcement block (Figure S2I).

EMG analysis

We analyzed pre-processed EMG (STAR Methods – EMG Recordings) from four forelimb muscles for 6 g cross actions with force matched within 0.1 g. EMG channels were z-scored across each session and aligned to the 6 g cross event. We compared the vector magnitude of EMG difference across actions to the difference within each individual action. Concretely, for each session, we split trials for each action into two halves and calculated the EMG PETH for each half. At each time point locked to 6g cross, we calculated the vector magnitude of difference across halves within an action, as well as the difference across actions using the PETH from one half. We compared across-action and within-action difference in the 100ms centered at 6g cross (Figure 1H).

Neuron detection and activity extraction with Suite2p

2-photon imaging data were processed with Suite2p[67] for motion correction, neuron detection, and fluorescence extraction; ROIs were classified using a classifier trained on sparse striatal MSN activity, neuropil-corrected ($F_{corrected} = F - 0.7 \times F_{neu}$), and neural activity was reported as z-scored $\Delta F/F_0$ with F_0 as the 1-minute running median. Figure 3H,I,K-M shows z-scored, raw fluorescence averaged over ROI pixels from PrairieView's "BOT".

Distinguishing D1- and D2-MSNs

We developed an automated thresholding method to identify D1- and D2-MSNs based on fluorophore expression (Figure S1). Functional imaging (green channel) was used to detect ROIs with Suite2p, while structural imaging (red channel) determined co-localization of cell-type–specific fluorophores. The overall workflow is outlined in Figure S1A. The method involved three main steps: (1) aligning and processing structural and functional data, (2) labeling ROIs manually (ground truth) and automatically using red-channel thresholds, and (3) comparing manual and automatic labeling.

1) **Structural data processing:**
   Functional data were motion-corrected and ROIs extracted using Suite2p (Figure S1B). As fluorophore expression is soma-restricted, only pixels within a 12-pixel diameter circle of each ROI center were used. Structural images (red channel, 1000 frames) were motion-corrected against the functional template, then background-subtracted using a 25-pixel rolling ball filter in ImageJ (Figure S1C,D).

2) **ROI labeling:**
   ROIs near the edge of the GRIN lens were excluded to avoid aberrations; the center of the lens was estimated from the red channel by computing the median coordinates of pixels above the 20th percentile intensity. Valid ROIs fell within a circle around this center.

   a. *Manual labeling:*
      Manual labels were assigned on average, motion-corrected, background-subtracted structural images overlaid with functional ROIs (Figure S1E). Labels were "red," "non-red," or

"no-ID" if uncertain. Each mouse had at least two sessions labeled three times (folds); only
ROIs consistently labeled across all folds were retained (Figure S1F,G). Thresholds of 0.7
(red) and 0.3 (non-red) were used for comparisons in Figure S1.
b. *Automatic labeling:*
Structural images were processed with Ilastik[69] (v1.3.3) to estimate per-pixel probability of
being red (Figure S1H). ROIs with >0.7 red pixels were labeled "red," <0.3 as "non-red," and
0.3–0.7 as "no-ID" (Figure S1I).
3) **Manual vs. automatic comparison:**
Label accuracy, swapping errors, and single-method errors were quantified across threshold pairs
to optimize performance (Figure S1J-M).
Thresholds of 0.3/0.7 (non-red/red) were used for general characterization (Figure 2), and 0.45/0.55
thresholds were used for holographic stimulation experiments (Figures 3 and 4).
910
Time course of neuronal activity
Neuronal activity was analyzed time-locked to action events. An example trace of activity averaged over
D1-, D2-, and all MSNs is shown alongside force (Figure 2B). We analyzed session-averaged PETHs
aligned to the force peak of pre-isolated 3 g cross actions (Figure 2C). We specifically analyzed activity at
force peak and 0.5 seconds before force peak, computed as z-scored F/F0 averaged over 3 frames
centered at each time point. Averaging over neurons of each cell type, we compared activity at force peak
versus 0.5 seconds before force peak (paired t-test). We also compared population-averaged activity
across actions at force peak (paired t-test). We plotted PETHs of example individual D1- and D2-MSNs
(Figure 2D). Only sessions with ≥20 trials per action and ≥10 neurons were analyzed (same as in STAR
Methods – Support Vector Machine (SVM) decoder model).
921
Support Vector Machine (SVM) decoder model
To test whether MSNs encoded specific actions, we trained a linear SVM (sklearn.svm.LinearSVC) to
predict action identity (e.g. push versus pull) from single-trial population activity. Each trial vector
comprised the $\Delta F/F_0$ of each neuron averaged over 5 frames (~167 ms) centered on the action event (i.e.
the frame at the event and 2 frames before and after) (Figures 2H,I and S2). Only sessions with ≥20 trials
per action and ≥10 neurons were analyzed, and trials were weighted by 1/(number of trials per action
class) to prevent bias. Trials were randomly split 90/10 for training/testing, repeated 100 times. The
regularization parameter (C) was optimized over a range of 1×10^{-5} to 100 via 10-fold cross-validation on
the training set (with 80% of training data used for sub-training and 20% for sub-testing in each sub-fold).
Accuracy was computed for each action separately and overall, and compared to a shuffle control **in**
which the model was trained on randomized action labels but tested on non-randomized labels.
933
934   **Matching number of D1- and D2-MSNs for SVM decoding**
935   To compare D1- and D2-MSNs, we equalized their numbers for SVM decoding (Figure S3K). For the cell
936   type with more neurons, a random subset was selected in each training/testing fold to match the number
937   of neurons in the other cell type.
938
SVM decoding across sessions
We reported SVM decoding accuracy for individual sessions and pooled across sessions, under different
reinforcement conditions.
SVM accuracy across each pair of actions, pooling the selected sessions across all reinforcement blocks:
• Figure 2H decoded the population activity vector including all MSNs. For push and pull, 3g cross
actions were used, and neural activity was centered at force peak.
• Figure 2I is the same as Figure 2H, but decoding population activity of a cell type (D1- or D2-).

• Figure S3I is the same as Figure 2H, and Figure S3J is the same as Figure 2I, except for
matched 3g cross actions and separating by action A and B rather than push and pull.
SVM accuracy for each selected session across reinforcement:
• Figure S2D decoded all MSNs at force peak for 3g cross actions, predicting push versus pull.
• Figure S2E (left) is the same as Figure S2D, but decoding population activity of a cell type.
• Figure S2F (left) is the same as Figure S2D, except with one SVM predicting action A versus
touch and another SVM predicting action B versus touch.
SVM accuracy pooling selected sessions within each reinforcement block:
• Figure S2E (right) decoded population activity of a cell type at force peak for 3g cross actions,
predicting push versus pull.
• Figure S2F (right) decoded all MSNs at force for 3g cross actions, with one SVM predicting action
A versus touch and another SVM predicting action B versus touch.
• Figure S2G is the same as Figure S2F (right), but decoding population activity of a cell type.
• Figure S2J is the same as Figure S2E (right), except decoding all MSNs for NR1-KO and
littermate control mice.
961
Time course of action decoding
We used the SVM to analyze the time course of action decoding. We extracted and decoded neural
activity centered at specific time points relative to action events. In Figure 2E-G, we analyzed non-
overlapping 5 frame bins locked to force peak. The bin at time 0 was centered at force peak. From -4
seconds to 0 seconds, the SVM was applied on trials from pre-isolated 3g cross actions. From after 0
seconds to 4 seconds, the SVM was applied on trials from post-isolated 3g cross actions. The SVM used
all MSNs in Figure 2EF and matched numbers of D1- or D2-MSNs in Figure 2G.
969
970 **Overlay of decoder accuracy and force**
971 Decoder accuracy and force were visualized in normalized units (Figure 2F) by centering data at -4
972 seconds from force peak (i.e. subtracting the force at -4 seconds from all time points) and scaling by the
973 window's (-4 s to +4 s) maximum trial-averaged value (i.e. dividing by the maximum).
974
SVM weights and dimension
The SVM assigned positive (negative) weights to neurons whose activation biased the SVM to predict
push (pull). For analysis of the weights directly, we z-scored the weights for each decoder (Figure 3G,J).
The weights define a dimension of population activity that best discriminates action identity. We extract
this dimension by using the decoder's weight vector and dividing by its L2 norm. This is a vector $w \in R^N$,
where N is the number of neurons and the vector magnitude is 1 ($\|w\|_2 = 1$).
981
Time course of SVM projection
We visualized the time course of action-specific population activity by projecting neural activity onto the
SVM dimension. At each time point, the vector of population activity $x \in R^N$ (where each entry is one
neuron's activity) was projected onto the SVM dimension: $w^T x \in R^1$, resulting in a scalar that represents
the weighted sum of neuronal activity. In Figure 2J, we used the SVM trained at force peak and projected
activity from pre-isolated 3g cross actions time locked to force peak. In Figure 3H,I and N, we used the
SVM trained at force peak and projected activity locked to holographic stimulation.
989
SVM-weighted neural ensembles
We defined neuronal ensembles that biased the SVM toward push or pull. For each action, we selected
the top $k *$ ensemble_frac neurons with the largest positive (push) or negative (pull) weights, where k is
the smaller of the number of push- or pull-weighted neurons. We used ensemble_frac = 0.25 to visualize

the most strongly weighted neurons (Figure 2K-N) and ensemble_frac = 1 to select as many neurons as
possible for holographic optogenetic stimulation (Figure 3G-N).
996
Time course of SVM-weighted neural ensembles
We analyzed the activity of push-weighted and pull-weighted neural ensembles that biased the SVM to
predict push or pull.
• We averaged the activity over the neurons in the push-weighted and pull-weighted ensembles
time-locked to force peak of 3g cross actions (Figure 2K,L).
• We separately analyzed the average activity of the identified D1- and D2-MSNs within these
ensembles (Figure 2M,N).
• In Figure 3K, we averaged ensemble activity in the 100ms before force peak of 3g cross actions.
• In Figure 3L,M, we time-locked ensemble activity to holographic stimulation.
1006
1007   **Effect of holographic stimulation on actions**
1008   We compared force in stimulation trials versus no-stimulation trials that immediately preceded stimulation
1009   trials. No-stimulation trials that occurred closer in time to the previous stimulation trial than the next
1010   stimulation trial were excluded. We analyzed force sampled at 1 kHz and low-pass filtered at 30 Hz (10th-
1011   order Butterworth filter), providing both high resolution visualization and de-noising.
1012
Given that stimulation occurred with a random (but small) latency from 3g cross, we performed an
alignment procedure to match the force of no-stimulation trials to stimulation trials in the time leading up
to stimulation. We identified the time in the no-stimulation trial that best matched the following stimulation
trial on 1) the force at time of stimulation ("target instantaneous force") and 2) the average force in the
100 ms window preceding time of stimulation ("target preceding force"). Concretely, we identified the time
in each no-stimulation trial that minimized a cost, which was the sum of two terms: 1) the absolute
difference between force at the time in the no-stimulation trial and the "target instantaneous force", and 2)
the absolute difference between the average force in the 100 ms preceding the time in the no-stimulation
trial and the "target preceding force." This alignment was performed on the raw 1 kHz force trace before
filtering at 30 Hz.
1023
We analyzed the difference between stimulation trials locked to stimulation time and no-stimulation trials
locked to this aligned time. Trials were pooled into "congruent stimulation" and "non-congruent
stimulation" conditions (Figure 4). Congruent stimulation consisted of 1) stimulation of the push ensemble
triggered on push 3g cross, and 2) stimulation of the pull ensemble triggered on pull 3g cross. Non-
congruent stimulation consisted of 1) stimulation of the push ensemble triggered on pull 3g cross, and 2)
stimulation of the pull ensemble triggered on push 3g cross. Accordingly, the number of data points for
statistics (Figure 4C,D) was 2×num_sessions.
1031
1032   **Statistical analysis**
1033   Statistical analysis was performed using custom scripts in Python. We used the 'scipy.stats' package to
1034   perform paired t-tests ('ttest_rel' method), unpaired t-tests ('ttest_ind' method), and linear regression
1035   ('linregress' method). Linear mixed effects models were performed using the 'statsmodels' package and
1036   'mixedlm' method. Significance was set at P=0.05. Statistical information in Tables S1 and S2.
1037
References

1. Donaldson, I., Marsden, C.D., Schneider, S., and Bhatia, K. (2012). Marsden's Book of
Movement Disorders (Oxford University Press)
https://doi.org/10.1093/med/9780192619112.001.0001.

2. Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia
disorders. Trends Neurosci. *12*, 366–375. https://doi.org/10.1016/0166-2236(89)90074-
X.
3. DeLong, M.R. (1990). Primate models of movement disorders of basal ganglia origin.
Trends Neurosci. *13*, 281–285. https://doi.org/10.1016/0166-2236(90)90110-V.
4. Giladi, N., McMahon, D., Przedborski, S., Flaster, E., Guillory, S., Kostic, V., and Fahn, S.
(1992). Motor blocks in Parkinson's disease. Neurology *42*, 333–333.
https://doi.org/10.1212/WNL.42.2.333.
5. Mazzoni, P., Hristova, A., and Krakauer, J.W. (2007). Why Don't We Move Faster?
Parkinson's Disease, Movement Vigor, and Implicit Motivation. J. Neurosci. *27*, 7105.
https://doi.org/10.1523/JNEUROSCI.0264-07.2007.
6. Simonyan, K., Cho, H., Hamzehei Sichani, A., Rubien-Thomas, E., and Hallett, M. (2017).
The direct basal ganglia pathway is hyperfunctional in focal dystonia. Brain *140*, 3179–
3190. https://doi.org/10.1093/brain/awx263.
7. Shohamy, D., Myers, C.E., Grossman, S., Sage, J., Gluck, M.A., and Poldrack, R.A. (2004).
Cortico-striatal contributions to feedback-based learning: converging data from
neuroimaging and neuropsychology. Brain J. Neurol. *127*, 851–859.
https://doi.org/10.1093/brain/awh100.
8. Knowlton, B.J., Mangels, J.A., and Squire, L.R. (1996). A Neostriatal Habit Learning
System in Humans. Science *273*, 1399–1402.
https://doi.org/10.1126/science.273.5280.1399.
9. Albin, R.L., Reiner, A., Anderson, K.D., Penney, J.B., and Young, A.B. (1990). Striatal and
nigral neuron subpopulations in rigid Huntington's disease: Implications for the
functional anatomy of chorea and rigidity-akinesia. Ann. Neurol. *27*, 357–365.
https://doi.org/10.1002/ana.410270403.
10. Panigrahi, B., Martin, K.A., Li, Y., Graves, A.R., Vollmer, A., Olson, L., Mensh, B.D.,
Karpova, A.Y., and Dudman, J.T. (2015). Dopamine Is Required for the Neural
Representation and Control of Movement Vigor. Cell *162*, 1418–1430.
https://doi.org/10.1016/j.cell.2015.08.014.
11. Dudman, J.T., and Krakauer, J.W. (2016). The basal ganglia: from motor commands to
the control of vigor. Neurobiol. Cogn. Behav. *37*, 158–166.
https://doi.org/10.1016/j.conb.2016.02.005.
12. Turner, R.S., and Desmurget, M. (2010). Basal ganglia contributions to motor control: a
vigorous tutor. Mot. Syst. – Neurobiol. Behav. *20*, 704–716.
https://doi.org/10.1016/j.conb.2010.08.022.
13. Yttri, E.A., and Dudman, J.T. (2016). Opponent and bidirectional control of movement
velocity in the basal ganglia. Nature *533*, 402–406.
https://doi.org/10.1038/nature17639.
14. Klaus, A., Martins, G.J., Paixao, V.B., Zhou, P., Paninski, L., and Costa, R.M. (2017). The
Spatiotemporal Organization of the Striatum Encodes Action Space. Neuron *95*, 1171-
1180.e7. https://doi.org/10.1016/j.neuron.2017.08.015.
15. Markowitz, J.E., Gillis, W.F., Beron, C.C., Neufeld, S.Q., Robertson, K., Bhagat, N.D.,
Peterson, R.E., Peterson, E., Hyun, M., Linderman, S.W., et al. (2018). The Striatum

Organizes 3D Behavior via Moment-to-Moment Action Selection. Cell *174*, 44-58.e17.
https://doi.org/10.1016/j.cell.2018.04.019.

16. Dhawale, A.K., Wolff, S.B.E., Ko, R., and Ölveczky, B.P. (2021). The basal ganglia control the detailed kinematics of learned motor skills. Nat. Neurosci. *24*, 1256–1269. https://doi.org/10.1038/s41593-021-00889-3.

17. Park, J., Polidoro, P., Fortunato, C., Arnold, J., Mensh, B., Gallego, J.A., and Dudman, J.T. (2025). Conjoint specification of action by neocortex and striatum. Neuron *113*, 620-636.e6. https://doi.org/10.1016/j.neuron.2024.12.024.

18. Kravitz, A.V., Freeze, B.S., Parker, P.R.L., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer, A.C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. Nature *466*, 622–626. https://doi.org/10.1038/nature09159.

19. Arber, S., and Costa, R.M. (2018). Connecting neuronal circuits for movement. Science *360*, 1403–1404. https://doi.org/10.1126/science.aat5994.

20. Joel, D., Niv, Y., and Ruppin, E. (2002). Actor–critic models of the basal ganglia: new anatomical and computational perspectives. Neural Netw. *15*, 535–547. https://doi.org/10.1016/S0893-6080(02)00047-3.

21. Niv, Y. (2009). Reinforcement learning in the brain. J. Math. Psychol. *53*, 139–154. https://doi.org/10.1016/j.jmp.2008.12.005.

22. Ito, M., and Doya, K. (2011). Multiple representations and algorithms for reinforcement learning in the cortico-basal ganglia circuit. Curr. Opin. Neurobiol. *21*, 368–373. https://doi.org/10.1016/j.conb.2011.04.001.

23. Samejima, K., Ueda, Y., Doya, K., and Kimura, M. (2005). Representation of Action-Specific Reward Values in the Striatum. Science *310*, 1337–1340. https://doi.org/10.1126/science.1115270.

24. Reynolds, J.N.J., Hyland, B.I., and Wickens, J.R. (2001). A cellular mechanism of reward-related learning. Nature *413*, 67–70. https://doi.org/10.1038/35092560.

25. Kravitz, A.V., Tye, L.D., and Kreitzer, A.C. (2012). Distinct roles for direct and indirect pathway striatal neurons in reinforcement. Nat. Neurosci. *15*, 816–818. https://doi.org/10.1038/nn.3100.

26. Vicente, A.M., Galvão-Ferreira, P., Tecuapetla, F., and Costa, R.M. (2016). Direct and indirect dorsolateral striatum pathways reinforce different action strategies. Curr. Biol. *26*, R267–R269. https://doi.org/10.1016/j.cub.2016.02.036.

27. Hikosaka, O., Takikawa, Y., and Kawagoe, R. (2000). Role of the Basal Ganglia in the Control of Purposive Saccadic Eye Movements. Physiol. Rev. *80*, 953–978. https://doi.org/10.1152/physrev.2000.80.3.953.

28. Yang, W., Carrillo-Reid, L., Bando, Y., Peterka, D.S., and Yuste, R. (2018). Simultaneous two-photon imaging and two-photon optogenetics of cortical circuits in three dimensions. eLife *7*, 1–21. https://doi.org/10.7554/eLife.32671.

29. Marshel, J.H., Kim, Y.S., Machado, T.A., Quirin, S., Benson, B., Kadmon, J., Raja, C., Chibukhchyan, A., Ramakrishnan, C., Inoue, M., et al. (2019). Cortical layer–specific critical dynamics triggering perception. Science *365*, eaaw5202–eaaw5202. https://doi.org/10.1126/science.aaw5202.

30. Piantadosi, S.C., Zhou, Z.C., Pizzano, C., Pedersen, C.E., Nguyen, T.K., Thai, S., Stuber, G.D., and Bruchas, M.R. (2024). Holographic stimulation of opposing amygdala

ensembles bidirectionally modulates valence-specific behavior via mutual inhibition.
Neuron *112*, 593-610.e5. https://doi.org/10.1016/j.neuron.2023.11.007.

31. Gill, J.V., Lerman, G.M., Zhao, H., Stetler, B.J., Rinberg, D., and Shoham, S. (2020).
Precise Holographic Manipulation of Olfactory Circuits Reveals Coding Features
Determining Perceptual Detection. Neuron *108*, 382-393.e5.
https://doi.org/10.1016/j.neuron.2020.07.034.

32. Vinograd, A., Nair, A., Kim, J.H., Linderman, S.W., and Anderson, D.J. (2024). Causal
evidence of a line attractor encoding an affective state. Nature *634*, 910–918.
https://doi.org/10.1038/s41586-024-07915-x.

33. Cruz, B.F., Guiomar, G., Soares, S., Motiwala, A., Machens, C.K., and Paton, J.J. (2022).
Action suppression reveals opponent parallel control via striatal circuits. Nature *607*,
521–526. https://doi.org/10.1038/s41586-022-04894-9.

34. Kaufman, M.T., Churchland, M.M., Ryu, S.I., and Shenoy, K.V. (2014). Cortical activity in
the null space: permitting preparation without movement. Nat. Neurosci. *17*, 440–448.
https://doi.org/10.1038/nn.3643.

35. Inagaki, H.K., Chen, S., Daie, K., Finkelstein, A., Fontolan, L., Romani, S., and Svoboda,
K. (2022). Neural Algorithms and Circuits for Motor Planning. Annu. Rev. Neurosci. *45*,
249–271. https://doi.org/10.1146/annurev-neuro-092021-121730.

36. Nelson, A., Abdelmesih, B., and Costa, R.M. (2021). Corticospinal populations
broadcast complex motor signals to coordinated spinal and striatal circuits. Nat.
Neurosci. *24*, 1721–1732. https://doi.org/10.1038/s41593-021-00939-w.

37. Economo, M.N., Viswanathan, S., Tasic, B., Bas, E., Winnubst, J., Menon, V., Graybuck,
L.T., Nguyen, T.N., Smith, K.A., Yao, Z., et al. (2018). Distinct descending motor cortex
pathways and their roles in movement. Nature *563*, 79–84.
https://doi.org/10.1038/s41586-018-0642-9.

38. Churchland, M.M., and Shenoy, K.V. (2024). Preparatory activity and the expansive null-
space. Nat. Rev. Neurosci. *25*, 213–236. https://doi.org/10.1038/s41583-024-00796-z.

39. Cui, G., Jun, S.B., Jin, X., Pham, M.D., Vogel, S.S., Lovinger, D.M., and Costa, R.M.
(2013). Concurrent activation of striatal direct and indirect pathways during action
initiation. Nature *494*, 238–242. https://doi.org/10.1038/nature11846.

40. Sheng, M., Lu, D., Shen, Z., and Poo, M. (2019). Emergence of stable striatal D1R and
D2R neuronal ensembles with distinct firing sequence during motor learning. Proc.
Natl. Acad. Sci. *116*, 11038–11047. https://doi.org/10.1073/pnas.1901712116.

41. Tecuapetla, F., Jin, X., Lima, S.Q., and Costa, R.M. (2016). Complementary
Contributions of Striatal Projection Pathways to Action Initiation and Execution. Cell
166, 703–715. https://doi.org/10.1016/j.cell.2016.06.032.

42. Foster, N.N., Barry, J., Korobkova, L., Garcia, L., Gao, L., Becerra, M., Sherafat, Y., Peng,
B., Li, X., Choi, J.-H., et al. (2021). The mouse cortico–basal ganglia–thalamic network.
Nature *598*, 188–194. https://doi.org/10.1038/s41586-021-03993-3.

43. Hintiryan, H., Foster, N.N., Bowman, I., Bay, M., Song, M.Y., Gou, L., Yamashita, S.,
Bienkowski, M.S., Zingg, B., Zhu, M., et al. (2016). The mouse cortico-striatal
projectome. Nat. Neurosci. *19*, 1100–1114. https://doi.org/10.1038/nn.4332.

44. Parker, J.G., Marshall, J.D., Ahanonu, B., Wu, Y.-W., Kim, T.H., Grewe, B.F., Zhang, Y., Li,
J.Z., Ding, J.B., Ehlers, M.D., et al. (2018). Diametric neural ensemble dynamics in

parkinsonian and dyskinetic states. Nature *557*, 177–182.
https://doi.org/10.1038/s41586-018-0090-6.

45. Gerfen, C.R., and Surmeier, D.J. (2011). Modulation of Striatal Projection Systems by
Dopamine. Annu. Rev. Neurosci. *34*, 441–466.
https://doi.org/https://doi.org/10.1146/annurev-neuro-061010-113641.

46. Maltese, M., March, J.R., Bashaw, A.G., and Tritsch, N.X. (2021). Dopamine differentially
modulates the size of projection neuron ensembles in the intact and dopamine-
depleted striatum. eLife *10*, e68041. https://doi.org/10.7554/eLife.68041.

47. Dang, M.T., Yokoi, F., Yin, H.H., Lovinger, D.M., Wang, Y., and Li, Y. (2006). Disrupted
motor learning and long-term synaptic plasticity in mice lacking NMDAR1 in the
striatum. Proc. Natl. Acad. Sci. *103*, 15254–15259.
https://doi.org/10.1073/pnas.0601758103.

48. Koralek, A.C., Jin, X., Long II, J.D., Costa, R.M., and Carmena, J.M. (2012). Corticostriatal
plasticity is necessary for learning intentional neuroprosthetic skills. Nature *483*, 331–
335. https://doi.org/10.1038/nature10845.

49. Jin, X., and Costa, R.M. (2010). Start/stop signals emerge in nigrostriatal circuits during
sequence learning. Nature *466*, 457–462. https://doi.org/10.1038/nature09263.

50. Santos, F.J., Oliveira, R.F., Jin, X., and Costa, R.M. (2015). Corticostriatal dynamics
encode the refinement of specific behavioral variability during skill learning. eLife *4*,
e09423. https://doi.org/10.7554/eLife.09423.

51. Dong, J., Hawes, S., Wu, J., Le, W., and Cai, H. (2021). Connectivity and Functionality of
the Globus Pallidus Externa Under Normal Conditions and Parkinson's Disease. Front.
Neural Circuits *Volume 15-2021*. https://doi.org/10.3389/fncir.2021.645287.

52. Yin, H.H., Mulcare, S.P., Hilário, M.R.F., Clouse, E., Holloway, T., Davis, M.I., Hansson,
A.C., Lovinger, D.M., and Costa, R.M. (2009). Dynamic reorganization of striatal circuits
during the acquisition and consolidation of a skill. Nat. Neurosci. *12*, 333–341.
https://doi.org/10.1038/nn.2261.

53. Arber, S., and Costa, R.M. (2022). Networking brainstem and basal ganglia circuits for
movement. Nat. Rev. Neurosci. *23*, 342–360. https://doi.org/10.1038/s41583-022-
00581-w.

54. Redgrave, P., Prescott, T.J., and Gurney, K. (1999). The basal ganglia: a vertebrate
solution to the selection problem? Neuroscience *89*, 1009–1023.
https://doi.org/10.1016/S0306-4522(98)00319-4.

55. Grillner, S., Robertson, B., and Stephenson-Jones, M. (2013). The evolutionary origin of
the vertebrate basal ganglia and its role in action selection. J. Physiol. *591*, 5425–5431.
https://doi.org/10.1113/jphysiol.2012.246660.

56. Falasconi, A., Kanodia, H., and Arber, S. (2025). Dynamic basal ganglia output signals
license and suppress forelimb movements. Nature *644*, 749–758.
https://doi.org/10.1038/s41586-025-09066-z.

57. Athalye, V.R., Carmena, J.M., and Costa, R.M. (2020). Neural reinforcement: re-entering
and refining neural dynamics leading to desirable outcomes. Neurobiol. Behav. *60*,
145–154. https://doi.org/10.1016/j.conb.2019.11.023.

58. Neely, R.M., Koralek, A.C., Athalye, V.R., Costa, R.M., and Carmena, J.M. (2018).
Volitional Modulation of Primary Visual Cortex Activity Requires the Basal Ganglia.
Neuron *97*, 1356-1368.e4. https://doi.org/10.1016/j.neuron.2018.01.051.

59. Athalye, V.R., Ganguly, K., Costa, R.M., and Carmena, J.M. (2017). Emergence of
Coordinated Neural Dynamics Underlies Neuroprosthetic Learning and Skillful Control.
Neuron *93*, 955-970.e5. https://doi.org/10.1016/j.neuron.2017.01.016.

60. Athalye, V.R., Khanna, P., Gowda, S., Orsborn, A.L., Costa, R.M., and Carmena, J.M.
(2023). Invariant neural dynamics drive commands to control different movements.
Curr. Biol. *33*, 2962-2976.e15. https://doi.org/10.1016/j.cub.2023.06.027.

61. Athalye, V.R., Santos, F.J., Carmena, J.M., and Costa, R.M. (2018). Evidence for a neural
law of effect. Science *359*, 1024–1029. https://doi.org/10.1126/science.aao6058.

62. Kao, T.-C., and Hennequin, G. (2019). Neuroscience out of control: control-theoretic
perspectives on neural circuit dynamics. Comput. Neurosci. *58*, 122–129.
https://doi.org/10.1016/j.conb.2019.09.001.

63. Stahl, C.M., and Frucht, S.J. (2017). Focal task specific dystonia: a review and update. J.
Neurol. *264*, 1536–1541. https://doi.org/10.1007/s00415-016-8373-z.

64. Akay, T., Acharya, H.J., Fouad, K., and Pearson, K.G. (2006). Behavioral and
Electromyographic Characterization of Mice Lacking EphA4 Receptors. J. Neurophysiol.
96, 642–651. https://doi.org/10.1152/jn.00174.2006.

65. Mosberger, A.C., Sibener, L.J., Chen, T.X., Rodrigues, H.F.M., Hormigo, R., Ingram, J.N.,
Athalye, V.R., Tabachnik, T., Wolpert, D.M., Murray, J.M., et al. (2024). Exploration biases
forelimb reaching strategies. Cell Rep. *43*.
https://doi.org/10.1016/j.celrep.2024.113958.

66. Akam, T., Lustig, A., Rowland, J.M., Kapanaiah, S.K., Esteve-Agraz, J., Panniello, M.,
Márquez, C., Kohl, M.M., Kätzel, D., Costa, R.M., et al. (2022). Open-source, Python-
based, hardware and software for controlling behavioural neuroscience experiments.
eLife *11*, e67846. https://doi.org/10.7554/eLife.67846.

67. Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Rossi, L.F., Dalgleish, H.,
Carandini, M., and Harris, K.D. (2017). Suite2p: beyond 10,000 neurons with standard
two-photon microscopy. bioRxiv, 061507. https://doi.org/10.1101/061507.

68. Miri, A., Warriner, C.L., Seely, J.S., Elsayed, G.F., Cunningham, J.P., Churchland, M.M.,
and Jessell, T.M. (2017). Behaviorally Selective Engagement of Short-Latency Effector
Pathways by Motor Cortex. Neuron *95*, 683-696.e11.
https://doi.org/10.1016/j.neuron.2017.06.042.

69. Sommer, C., Straehle, C., Köthe, U., and Hamprecht, F.A. (2011). Ilastik: Interactive
learning and segmentation toolkit. In 2011 IEEE International Symposium on
Biomedical Imaging: From Nano to Macro, pp. 230–233.
https://doi.org/10.1109/ISBI.2011.5872394.

1253