1   Mothbox: inexpensive, lightweight, automated light trap for scalable insect biodiversity

2   monitoring

3   Hubert A. Szczygieł[1,2]*, Daisy H. Dent[1,4,5], Andrew Quitmeyer[3,6]
4
[1] Smithsonian Tropical Research Institute; Apartado 0843-03092, Balboa, Ancón, República
de Panamá.
[2] Department of Biological and Environmental Sciences, Gothenburg Global Biodiversity
Centre, University of Gothenburg, Medicinaregatan 7B, 413 90 Göteborg, Sweden
[3] Digital Naturalism Laboratories, Sibert Ave 123B, Gamboa, Colón Province, Panama
[4] Department of Environmental Systems Science, ETH, Zürich, Switzerland.
[5] Max Planck Institute of Animal Behavior, Konstanz, Germany
[6] Department of Human Centered Design & Engineering, University of Washington, 3960
Benton Lane NE 428, Seattle, WA 98195
14

15 *Corresponding author. Email: hubert.70@gmail.com
16

Abstract

1. Insects represent the most diverse group of organisms on Earth, and comprise the

majority of known species, yet they are seldom accounted for in large-scale

biodiversity monitoring systems and conservation planning.

2. We have developed the *Mothbox*: an automated light trap that uses open science

hardware to make insect monitoring accessible to non-specialists and scalable for

scientific and conservation purposes. We have also developed *Mothbot*: a machine

learning tool and user interface for processing data from automated light traps. The

Mothbox is portable, durable to harsh environments, and low cost to produce, while

Mothbot prioritizes human-in-the-loop data validation of computer vision outputs

to ensure data quality and aid in developing custom image reference libraries.

3. Mothbox has now been extensively tested in field conditions; with > 185

deployments, spanning > 450 nights and > 100 sampling locations. The two case

studies presented here demonstrate the potential of Mothbox data for behavioural

studies and conservation projects.

4. As the young field of automated insect monitoring continues to develop, Mothbox

provides an affordable entry point for a wide range of stakeholders. It enables

entomologists to explore new research questions by tracking insect diversity in real

time and across entire landscapes. Mothbox will help build global insect-monitoring

capacity, improving our ability to detect biodiversity loss and guide effective

conservation action.

Keywords: biodiversity; monitoring; computer vision; insects; light trap; open source;

artificial intelligence

1 Introduction

The majority of the world's species are insects and most insects are active at night

(Wong & Didham, 2024). Insects are incredibly valuable ecologically and economically, and

rapidly declining in many parts of the world (Basset & Lamarre, 2019; Scudder, 2017;

Wagner et al., 2021). However, there are limited options for measuring insect diversity at

scale. Traditional approaches require specialised knowledge and considerable time

investment (Montgomery et al., 2021), while newer technological approaches have their

own limitations. For example, genetic approaches (e.g. DNA metabarcoding) remain

expensive, rely on reference libraries with limited coverage for the tropics (Alvarado-

Robledo et al., 2024), and are poorly suited to assessing abundance (Sickel et al., 2023),

while satellite-based remote-sensing lacks the resolution for direct monitoring of insects

(Rhodes et al., 2022).

Insects present a high-resolution solution for quantifying biodiversity as their high

abundance, exceptional taxonomic richness, and relatively limited dispersal mean they

often show clear and measurable responses to environmental change. For example, a week

of conventional camera trapping at one site for mammals can often yield no data (Kays et

al., 2020), in contrast, a week of insect monitoring at the same location can produce

hundreds of records for dozens of species (Brehm & Axmacher, 2006). In addition, great

diversification of insects is partially the result of specialisation, which makes them

particularly relevant biodiversity indicators. For instance, Lepidoptera larvae often feed on

only one species or genus of plant (Dyer et al., 2007), and adults can similarly be adapted to

pollinating particular host plants (Hahn & Brühl, 2016). Consequently, monitoring

Lepidoptera can provide inferences about the plant community (Duran et al., 2022a; Tyler,

2020). Ultimately, insects are rarely included in conservation planning, despite their great

diversity, ecological significance, and significant evidence of their global decline, largely

because of the difficulty of standardized large-scale insect monitoring (Chowdhury et al.,

2023; Høye et al., 2021).

Traditional approaches to monitoring nocturnal insects are time-consuming and

require high degrees of specialised knowledge. Moth-sheeting, i.e. attracting moths and

other nocturnal insects to lights and setting a sheet for them to land on, is an effective

method for visually assessing insect populations or targeting particular taxa. However,

documenting every insect that lands on the sheet is often impractical, especially in areas

with abundant and diverse insect communities, and requires researchers to attend sheets

throughout the night (Duran et al., 2022a). Bucket traps designed to attract and capture

insects do not require ongoing maintenance throughout a night, but do require extensive

manual sample processing once collected (Brehm & Axmacher, 2006). To bypass the

difficulties with traditional methods, an automated light trap that uses a programmed light

source and computer vision for identification was developed in 2021 (Bjerge et al., 2021).

This proof of concept inspired the creation of a number of automated light trap devices,

including the Automatic Moth Trap (Möglich et al., 2023), Automated Monitoring of Insects

(*UKCEH AMI System | UK Centre for Ecology & Hydrology*, n.d.), Insect Detect (Sittinger et al.,

2024), DIOPSIS (*Camera | Diopsis.Eu*, n.d.), and others (Beuchert, 2024; Hobern, 2021;

Korsch et al., 2023). The field of automated non-lethal insect monitoring is in its infancy,

and the devices developed thus far have advantages and disadvantages depending on the

monitoring context. There are trade-offs in total cost, portability, image resolution, power

consumption, insect attraction lights, 4G capabilities, ease of repair, ease of modification,

weather resistance and durability, to name a few.

Here we describe the development and proof-of-concept of the Mothbox, an open-

source, automated light trap designed for scalable insect monitoring in challenging field

conditions. The Mothbox device integrates a low-cost Raspberry Pi computer, a high-

resolution camera, and specialized attractant lighting to capture time-lapse images of

nocturnal insects landing on a target sheet. Unlike currently available insect traps, the

Mothbox is optimized for portability, adaptability, high-resolution images, and weather

resistance. Furthermore, the low materials cost (~$300) makes it possible to deploy many

Mothboxes as a part of a sensor network. There are currently two Mothbox versions:

Mothbox 4.5, which is a modular design oriented towards Do-It-Yourself construction and

modification, and Mothbox 5.0, which is oriented towards mass production.

We also describe Mothbot, an open-source (MIT license) data processing and

organization platform for automated light trap photos. The Mothbot platform detects

insects in images, identifies them to a user-defined taxonomic threshold, facilitates rapid

human validation and refinement of the automatic identifications, then produces a Darwin

Core (Wieczorek et al., 2012) formatted CSV file with the results.

We use two case studies conducted in Panama using the Mothbox and Mothbot to

demonstrate potential applications in behavioural research and conservation. The first

study investigates patterns in insect activity and community turnover over the night,

capturing temporal dynamics often missed by traditional sampling methods. The second

study examines the impacts of land-use (teak plantation, secondary and mature forest) on

nocturnal insect diversity and composition, providing insights into how land-use change

influences community composition.

2 Materials and methods

2.1 Design

2.1.1 Mothbox

The Mothbox is an automated light trap for scalable insect monitoring (Fig. 1).

Inspired by initial prototypes developed in 2021 (Bjerge et al., 2021), we set out to design a

device that is robust and lightweight enough for use in remote fieldwork. Our vision was to

create a tool that 1. Requires no specialised knowledge to run, 2. Produces real ground-

based biodiversity data describing insect communities, 3. Is affordable, durable and

sufficiently portable for use in large-scale field-based monitoring and conservation

projects. Development has focused on deployment in off-grid, remote tropical forests, as no

other automated insect monitoring tools are currently suited to this use case. In addition

to housing the most biodiversity, the tropics also present the most hostile conditions for

electronics. Thus, if the Mothbox can operate and collect usable data in tropical forest

environments, it will be capable of monitoring insects in more forgiving environments. The

Mothbox design prioritises low cost, simplicity, and repairability, and all documentation is

open-source to maximise potential for scaling and broad use.

The Mothbox is composed of a Raspberry Pi miniature computer that controls a 64

megapixel camera (9248 × 6944 resolution), lights for attracting nocturnal insects, and

lights to illuminate the insects for taking photographs (akin to a camera flash). The system

is housed in a watertight plastic box and powered by a 156Wh lithium battery

(commercially available and legal on most flights). All electronic components are within the

watertight housing in the default configuration. Ports in the base of the housing allow for

connection to a solar panel, additional batteries, or external light sources as required. The

camera faces a 'target' - a piece of 3mm acrylic with felted acrylic fabric on one side that

provides a surface diffusely illuminated by the attractant light and with enough texture for

insects to land. The default sheet size is 29 cm x 20.5 cm, and is attached to the Mothbox via

clear acrylic arms positioned 30cm from the camera. This gives a functional resolution of

25 px/mm per photograph, at this resolution most insects > 3mm in length can be readily

identified to family level (Fig. 2). At this distance, the camera field of view can

accommodate a sheet of up to 29 cm x 58 cm, however this leaves less margin for error in

positioning the arms and sheet and so we suggest use of the default sheet size. The

Mothbox can easily be adapted to using larger target sheets at greater distances to the

cameras, although this will result in functionally lower resolution photographs (fewer

pixels per insect). Mothbox can also be positioned to focus on any flat surface, without the

need for a dedicated target sheet, as the camera auto-focuses. A complete Mothbox weighs

2 - 3 kg , depending on the model and configuration, and is 28 x 18 x 12 cm.

We have prioritised ease of Do-It-Yourself construction and repair in the Mothbox

4.5 design. Minimal tools are required for construction, and most repairs can be

accomplished with a screwdriver, knife, and hot glue stick. We opted for using lever nuts

for electrical connections to avoid soldering and enable ease of modification and repair.

The internal framework as well as the arms and sheet backing are made with laser-cut

3mm acrylic, and all components can be cut from a single 50 x 60 cm sheet of acrylic. While

still 3D-printable, this flat pack design choice enables easy field servicing, as replacement

components can be cut and drilled by hand from acrylic or wood of the appropriate

thickness. In terms of programming, setting up a Mothbox in default configuration requires

simply loading a ready to go operating system image onto an SD card. No coding expertise

is needed. The main customization required is changing the sampling schedule, which can

be achieved by editing a scheduling CSV file placed in the data storage drive. Mothbox 5.0

combines many of the electronic components of Mothbox 4.5 into one printed circuit board

(PCB), decreasing construction time, number of components, and cost per unit. Mothbox

5.0 includes physical switches for setting the sampling schedule, which increases ease of

use in field conditions. Mothbox 5.0 electronics are housed in a custom 3D-printed box,

while 4.5 are in a commonly available clear acrylic box.

Motion-activated photo saving was the default on the original automated light trap

prototype (Bjerge et al., 2021). We use timelapse (photos taken at regular intervals)

instead to conserve battery life and to reduce costs. Motion-activated functions are

computationally heavy. Keeping on-board computational requirements to a minimum

decreases energy consumption and enables use of less expensive Raspberry Pi models.

Furthermore, we have observed that the increase in richness of insect assemblages

attracted to the sheet tends to be cumulative if the attractant lights remain on. If an insect

moves off of the sheet while the attractant lights are on, it still typically remains in the area

of the Mothbox until the lights turn off, flying around it or landing on different surfaces.

Mothboxes can be programmed for any sampling schedule. In Panama, where daylength is

consistently between 11.35 and 12.40 hours with sunset at 18:20 ($\pm$ 20 mins annually) and

sunrise at 6:20 ($\pm$ 20 mins), we programmed the following default schedule: Five one-hour

sampling intervals spaced throughout the night (at 19:00, 21:00, 23:00, 2:00 and 4:00) with

photographs taken every minute. Each Mothbox produces 300 photos (approximately

5.2GB of data) per night on this schedule. We chose this as the default schedule over

continuous recording to save battery life while still sampling insect assemblages at

different times of night. This approach has the added advantage of clearing the sheet of

insects between sampling sessions, which anecdotally decreases the abundance of

predators (geckoes, mantids, and ants) on the sheet and allows for insect turnover

throughout the night. The Mothbox lasts for approximately 12 hours of run time per

battery, or two nights with the default schedule.

The single integrated battery can be augmented by attaching an additional box

containing up to three more batteries connected in parallel, increasing run time to 48

hours. Portable solar panels ranging from 10-60 watts can be added to extend run time.

Larger solar panels will allow more hours of sampling per night or fewer required hours of

direct sunlight. We caution that solar panels are of limited usefulness in closed-canopy

tropical forests, especially in the wet season. The Mothbox system can run on any schedule

with no hardware modification, the only limitation is battery life and so for continuous

operation the Mothbox can be connected to mains power.

The hardware for both Mothbox versions is fully open source with a CERN-OHL-W

license (Digital Naturalism Laboratories, 2025a), the operating software has an MIT

licence, and the documentation is CC BY 4.0. Full specifications, including instructions for

construction, programming, use, and adaptation are available on Github (*Mothbox Github*,

2024).

2.1.2 Mothbeam

To complement the Mothbox, we used an open-source printed circuit board (PCB)

with light emitting diodes (LEDs) that emit a range of light frequencies designed to

specifically attract insects (the Mothbeam). The Mothbeam, originally created and open-

sourced by Moritz Buttlar of LabLab (Buttlar, 2024), was designed in collaboration with the

Mothbox team as an inexpensive, open-source equivalent of a LepiLED (Brehm, 2017).

There are two Mothbeam PCB configurations, each with three LEDs. One configuration

targets violet light (365, 395, 405 nm) and the other targets visible light (one full spectrum

white LED, 450, 520 nm). By default, each Mothbox 4.5 comes with one of each PCB and

both PCBs are activated when collecting data. Mothbox 5 has three PCB slots that are each

individually controllable. The Mothbox can be programmed to only use one PCB, and the

PCBs can be modified to deactivate individual LEDs or dim light intensity, to target

particular taxonomic groups or reduce battery consumption. We have also designed stand-

alone Mothbeams for use with traditional moth sheeting. Either two or four PCBs are

mounted to an aluminium heat sink, housed in water-resistant acrylic housing. Material

costs for one standalone mothbeam is approximately $30. Just like the Mothbox, the

Mothbeam runs on 12V power (it is operable between 9 and 36V). The addition of an

inexpensive ($10) step-up converter allows use of the Mothbeam with any standard USB

power bank (5V). Like the Mothbox, Mothbeams are fully open source with a CERN-OHL-W

licence (Digital Naturalism Laboratories, 2025b).

2.1.3 Mothbot

We developed 'Mothbot', an open-source machine learning tool and user interface

for processing data from automated light traps. There are two parts of Mothbot: 'Mothbot

Process' and 'Mothbot Classify'. The first detects insects from sourceImages, identifies

them, and spatially clusters them, the second is designed for human validation of computer

vision identification and manual finer-taxonomy classification, as well as data export.

Mothbot Process uses a combination of two open-sourced AI programs: YOLO (Redmon et

al., 2016) and BioCLIP (Stevens et al., 2024). A custom YOLO model trained on Mothbox

data from 4500 hand-labeled images automatically detects and crops individual insects

(down to ~1 mm total length) from novel imagery. These cropped images (called

'detections') are then fed to a modified BioCLIP model with filters for taxonomic and

geographic limitations to assist detection. The detections are then processed with a custom

clustering algorithm. The detections' visual features are extracted with DINOv2

(https://github.com/facebookresearch/dinov2) and then grouped by similarity with

HDBSCAN (Hierarchical Density-Based Spatial Clustering of Applications with Noise -

https://pypi.org/project/hdbscan/). These visual clusters are further sub-grouped

temporally (i.e. by the time the photo they appear on was taken) to create hierarchical

groupings of detected organisms that are not only visually similar but exist in temporal

'tracks' over a set of sequentially-taken images. The metadata associated with each

detection are then added into each detection image via EXIF.

The resulting output is then loaded into the 'Mothbot Classify', a user interface that

allows for human validation of the BioCLIP identifications. Mothbot Classify includes an

easily searchable, hierarchical taxonomic classification system that can be rapidly applied

to detections in a dataset. Mothbot Classify facilitates designating morphospecies that can

be linked to iNaturalist observations and exports data to a Darwin Core formatted CSV file

(Wieczorek et al., 2012). The human-in-the-loop workflow for refining taxonomy is not

required for generating an output CSV file, but we believe it is currently an essential step in

generating high-quality data from automated insect monitoring devices, as it is for mammal

camera trap data (Huebner et al., 2024; Leorna & Brinkman, 2022).

2.2 Field assembly and deployment

When the arms and target sheet are in place and the Mothbox has been switched on,

it can be mounted to a tree using a strap through the handle (Fig 1). Straps used for

conventional camera traps work well for this. In our case studies we deployed the Mothbox

at chest height on trees, which minimises additional installation equipment while still

elevating the Mothbox above the ground to increase attractant light range. Mothboxes can

be deployed in the canopy with a throw line, on the ground, in open areas, and mounted to

tripods or frames with minor modifications. In some areas, for example the tropical forests

we used for the case studies, deploying a Mothbox on the ground not only decreases the

range of the attractant lights, but also invites predatory ants and spiders to the target sheet,

and so elevating the device is advised. An early version of the Mothbox was developed for

drone deployment into rainforest canopies during the 2023 Rainforest XPRIZE competition

in Singapore. After collection, photos can be retrieved via a USB drive or SD card, and

processed with the Mothbot program.

2.3 Data processing

Processing data with the Mothbot user interface (UI) starts with selecting file paths

for a folder containing Mothbox photos, a CSV file that contains deployment metadata, and

a CSV file containing a species list. The formatting for the metadata CSV file is based on the

Camera Trap Metadata Standard (Forrester et al., 2016) and Darwin Core (Wieczorek et al.,

2012) (Table S1). The species list CSV file is in the format of Global Biodiversity

Information Facility (GBIF) downloads (Table S2). Species lists from GBIF require cleaning

to remove synonyms and other errors (SI Script 1). For our case studies, we generated a

CSV file from GBIF for all insects recorded in Panama and Costa Rica. We also tried running

Mothbot with a species list that included arachnids as well as insects, because arachnids

are also occasionally attracted to Mothboxes. However, we found that model accuracy

improved when arachnids are excluded, and the analysis process was more time efficient

when manually correcting the identifications of spiders.

After selection of the required filepaths, the photos can be processed using the tabs

on the Mothbot Process user interface. The first tab finds detections on the source image,

the second identifies them, the third adds metadata to JSON files associated with each

sourceImage, the fourth clusters detections perceptually, and the fifth adds EXIF data,

including GPS coordinates and identification, to cropped detections. Complete instructions

for data analysis using the Mothbot scripts can be found on the Mothbox website (*Mothbox*

Github, 2024). The final output of Mothbot is a CSV file with occurrences and identifications

of each organism photographed in a given deployment, following Darwin Core formatting

(Wieczorek et al., 2012).

2.3.2 Terminology

To standardize terminology in automated insect monitoring research, and align with

standards used for other automated biodiversity data collection (Bubnicki et al., 2024;

Wieczorek et al., 2012), we propose a set of terms used to describe different parts of the

data collection and analysis workflows. For example, *deployment* describes a discrete

sampling interval associated with one location and one device; *sourceImage* describes the

whole un-cropped image taken by an automated insect monitoring device; and *detection*

specific subset of a sourceImage that covers only a portion of the sourceImage that contains

one item / organism (generally an insect). See Table S3 for a more extensive list of

terminology. We understand that terminology will continue to evolve alongside

developments in the field of automated insect monitoring, and do not intend for this to be a

final, definitive list, rather an entry point for cross-platform collaboration and

standardization.

2.4 Case Studies

For both case studies we used tree-mounted Mothbox 4.5 devices. For each night

sampled, we recorded data for five hour periods starting at 19:00, 21:00, 23:00, 2:00, and

4:00), with timelapse set to capture one photo every minute (defined as sampling schedule

'scheduleA' in our metadata). After collecting the Mothboxes from the field we processed

the data using the Mothbot system set to order level automatic identification, we then

manually identified each insect detected to the finest taxonomic level possible and assigned

morphospecies to all insects that were not identified to species level.

2.4.1 Case study 1: Insect activity and time of night

The Mothbox is well suited to answer questions about insect behaviour as it is

unsupervised and collects standardised data. Most flying insects use light to maintain flight

altitude and control, via the dorsal-light-response (Fabian et al., 2024). Because of this, it

has been suggested that insects are only trapped by lights in passing, rather than being

attracted to them from farther away. Based on the assumption that insects attracted to light

traps, like the Mothbox, are active only in the immediate sampling area, then the Mothbox

can assess activity patterns of nocturnal insects and shifts in composition at different times

of night. Understanding how the activity patterns of different insects vary throughout the

night can help describe ecological relationships that are otherwise difficult to study, such as

overlapping activity patterns of different insect groups and species of specialist

insectivorous bats (Rydell et al., 1996).

This case study aimed to assess how insect activity patterns changed over the night.

A single Mothbox was deployed with an extra battery for four continuous nights 10 - 13

April 2025. The sampling site (7.49388, -80.31782) was located on a ridgetop in a remote

part of the Azuero Peninsula of Panama accessed by offroad driving and hiking.

2.4.3 Case study 2: Land use in fragmented landscape

This study demonstrates that the Mothbox has significant potential as a tool for

conservation and restoration monitoring. As tropical ecosystem restoration accelerates in

response to the twin crises of climate change and biodiversity loss, the need for effective,

scalable biodiversity monitoring has never been greater (Collen et al., 2008; Schmeller et

al., 2017). While carbon sequestration is relatively straightforward to quantify, assessing

biodiversity - particularly insect diversity - remains a major challenge (Wauchope et al.,

2024). Traditional survey methods are often labor-intensive, taxonomically limited, and

spatially constrained, making it difficult to capture the full complexity of ecological

recovery. As a result, large-scale reforestation projects tend to be evaluated and funded

based on carbon outcomes, while biodiversity gains remain poorly documented and

undervalued (Cole et al., 2024; Ermgassen & Löfqvist, 2024). Panama currently recognizes

monoculture plantations of introduced timber species as a type of reforestation (*DECRETO*

LEY No. 1, 1994), while ecological restoration success tends to be higher for natural

regeneration than for active restoration in tropical forests (Crouzeilles et al., 2017). To

assess the effect of different reforestation systems on insect diversity, we deployed four

Mothboxes simultaneously on the Azuero Peninsula of Panama at four sampling points for

two consecutive nights 11 - 13 April 2025. The Mothboxes were deployed in two stands of

teak plantation, a young secondary forest planted with native species, and a mature

secondary forest. All sampling points were separated by at least 250m.

3 Results and Discussion

3.1 Deployments

During the insect activity and time of night case study Mothbot generated a total of

6,271 detections, with nine orders and 43 families represented across all nights of the

deployment. A total of 219 unique species were recognised but only 36 of the species could

be readily identified to species level by a general entomologist (not an expert taxonomist),

the remaining 183 (84%) were assigned morphospecies names. A total of 1046 images

were collected during this deployment, less than the expected 1200 due to early battery

discharge on the last night of the deployment (consequently, there is no data for the final

session of night 4). The target sheet was clean of insects at the beginning of each hour-long

session (Fig. 3 c and d), indicating that when the Mothbeam attractant lights were off all

insects left the sheet. During each session, both number of detections per photo (Fig. 3 c)

and morphospecies richness (Fig. 3 d) increased with time since attractant lights turned on.

The first session (19:00 - 20:00) had the greatest number of cumulative detections and the

greatest morphospecies richness (Fig. 3 a and b) (P = .0002 and .0021, respectively,

ANOVA).

During the land use case study Mothbot generated a total of 2,755 detections across

all four sampling sites and both nights of the deployments. Eight orders and 40 families

were represented, and a total of 158 species were recorded (though 94% were designated

morphospecies). Both mature and native restoration forests had greater species richness

and number of detections than the teak plantations. The mature forest site had greater

richness than the native restoration forest, while the converse was true for the total

number of detections (Fig. 4).

Some limitations encountered during the deployments included the limited battery

life restricting the monitoring time, and the potential for insects to land outside the camera

frame (e.g. on the bark of the deployment tree). Battery life could be extended with use of

additional batteries or a solar panel positioned in the canopy or forest clearings. However

such logistical difficulties may not be required for obtaining ecologically significant results,

given that it is possible to differentiate forest types from collecting moth data for just one

night (Duran et al., 2022b). Both case studies occurred towards the end of the dry season,

when the moon was close to full and above the horizon during all sampling intervals. Both

of these variables suggest that total richness and activity were underestimated for these

sample sites (Janzen, 1987; McGeachie, 1989). It did not rain during our chosen case

studies as it was the dry season in Panama, however Mothboxes withstood daily

downpours during wet season testing in Panama for several weeks at a time. Due to the

remoteness and rugged terrain of the field sites used for our case studies, we deployed the

Mothboxes on trees using camera trap straps. This is the fastest, lightest, and easiest

system, short of putting them on the ground, however it does mean that some moths

attracted to the Mothboxes likely landed on the trunk of the tree instead of on the target

sheet.

3.2 Data processing

The Mothbot system allows users to choose the taxonomic level at which the AI

identifies detections. Model accuracy depends largely on the quality and quantity of

reference data, as well as the presence of cryptic species. Finer taxonomic resolution

introduces accuracy trade-offs and increases the time required for manual validation. We

found the model sufficiently accurate at the Order level for rapid human validation. In

highly diverse regions with limited reference data, such as the tropical forests used in our

case studies, the model performs poorly at species level. With more manual identification

of automated insect detections, we expect that future AI models will achieve finer

resolution, with reliable family-level identification for most groups and genus- or species-

level identification for selected taxa.

Our data analysis system is based on the assumption that automated monitoring

systems for insects empower experts and do not replace them. Expert taxonomists will be

invaluable in the training and refinement of AI models. Consequently, we prioritized

accuracy and ease of human verification over fine taxonomic identification at this stage.

When expert taxonomists are not available to aid in data validation and refinement, we

recommend uploading one example of each morphospecies (unique-looking insect that is

distinct from others but cannot be identified to species level) from each deployment to

iNaturalist for verification (Campbell et al., 2023). Because BioCLIP uses GBIF data, these

iNaturalist uploads will eventually feed back into improving the accuracy of Mothbot

identifications.

4 Future Directions

4.1 Hardware

Mothbox can be adapted for specific use cases beyond those showcased in the case

studies. For example, Mothbox hardware includes extra relays that can be used for adding

additional functions such as triggering a release of pheromones at programmed times to

attract targeted species, or to control nectar release from a synthetic flower. The main

limiting factor for deployment length is battery life. If deployed in a location with

continuous power, Mothboxes could operate for entire field seasons rather than just short

deployments in remote locations. While we have focused on nocturnal insects and have

only conducted tests at night, the additions described above could expand the range of

operation into daytime. The Mothbox is designed as a standalone sensor, but it is ideally

suited for use in multi-sensor monitoring arrangements, for example with simultaneous

acoustic monitoring with Audiomoths (Hill et al., 2017), and conventional camera traps. In

addition, synchronizing Mothbox data collection with environmental variables, including

ambient light, wind speed, humidity, precipitation, and temperature, would help identify

the drivers of changes in insect activity throughout the night and shifts in insect richness

across seasons.

While Mothbox development has focused on deployment in remote tropical

locations, the system can be applied in any regions or ecosystem where there is an interest

in monitoring nocturnal insects. Inclusion of Mothboxes in large-scale ecological

monitoring networks like the Long Term Ecological Research (LTER) network and the

National Ecological Observatory Network (NEON) would yield an additional valuable

dataset for insect population trends (*LTER*, n.d.; *NSF NEON*, n.d.). Mothbox would also be

well suited to monitor pollinator activity and diversity in agriculture, or the spread of

invasive species.

4.2 Data processing

The accuracy of computer vision models improves when the training data closely

matches the data being classified, and Mothbox raw images provide ideal conditions:

insects are captured against a flat white background, with consistent lighting and a fixed

distance from the camera. The same cannot be said for the massive and visually diverse

datasets like those on GBIF, which include many dead, pinned insects, and insect *in-situ* in

their environments. In time, with larger Mothbox datasets, we will augment the BioCLIP

model with human-validated data identified to finer taxonomic resolution data, producing

a region-specific improved version of the Mothbot that can accurately identify taxa below

family level. We believe that this tiered regional approach will result in the highest quality

and efficient data collection, whereby an initial round of monitoring and identification to a

coarse taxonomic resolution using the default Mothbot is followed by human validation to

create a regional reference library. Future collections in the region can then be run through

an augmented Mothbot system at finer-scale resolution. We believe that broad use of this

approach will rapidly change our ability to classify species so that all insects photographed

with automated insect monitors can be confidently identified to Genus or Family level by

Mothbot. In the meantime, higher taxonomic levels can be used as surrogates of species-

level data (Báldi, 2003; Zou et al., 2020), an approach that is key to current large-scale

insect monitoring in the tropics while taxonomic expertise, genetic reference libraries, and

photographic reference libraries catch up with monitoring needs.

Future success of automated insect monitoring will require greater investment in

traditional taxonomy and ecological research. The accuracy of the computer vision models

employed by automated insect monitors like the Mothbox are limited by training data.

Improving training datasets will require manual validation and refinement of existing

images and ongoing data collection. Automated insect monitors are likely to document

undescribed species, presenting exciting opportunities for prioritising sampling sites and

taxa for identification and description of novel species.

Perhaps the most important need for automated insect monitoring is a centralised

repository for human-validated image storage and model development. Such repositories

exist for mammal camera trapping (Vélez et al., 2023). A comparable platform for insect

light-trap data would allow projects worldwide to upload their observations, with human-

validated records feeding into regionally refined machine-learning models. In return, users

would receive AI-based identifications and organised datasets. As participation grows, the

expanding volume of training data would further improve identification accuracy. To

reduce cloud storage requirements for such a repository only cropped images that have

undergone an initial on-board processing step could be accepted, rather than bulk raw data

which is often largely composed of blank sheet space.

5 Conclusion

Mothbox will offer for insect monitoring what low-cost camera trapping offers for

mammals. By decreasing the time, cost, and specialised knowledge required for monitoring

insects, Mothbox enables standardised insect monitoring at the landscape scale. This

technology increases scientific capacity for monitoring global trends in insect populations,

and acts as a tool for exploring new ecological relationships and behavioural patterns.

Importantly, Mothbox will make biodiversity monitoring more accessible to community

conservation groups, empowering restoration and conservation.

Figures

498
499 **Figure 1** Mothbox 4.5 deployed in a tropical forest in Panama.
500

501
502   **Figure 2** Unedited photo taken by a Mothbox in Panama, with an inset to demonstrate the
503   capacity for documenting small insects like this 1.5 mm long minute pirate bug
504   (Anthocoridae). The resolution of the full photo is 9248 x 6944 pixels. The scale provided
505   in the inset photo is in millimeters.

(c)

Figure 3 data from insect activity case study: (a) total number of detections (individual organisms in each photo, for all photos) for five discontinuous one-hour sampling sessions; (b) total species richness within five discontinuous one-hour sampling sessions; (c) total number of insect detections in each photo by minute from start of sampling, for the five sampling sessions; (d) species richness per photo across the five sampling sessions. All four panels represent data averaged across the four nights of the deployment. The effect of session on cumulative detections and richness was significant (P = .0002 and .0021, respectively). Error bars in (a) and (b) represent standard error.

515
Figure 4 Insect detections and richness for different land uses. Four Mothboxes were
deployed simultaneously for two nights at points with different land use histories. Two
points were in non-native teak (*Tectona grandis*) plantations, one point was in an area
planted with native tree species and undergoing natural regeneration, and one point in
mature native forest. Data from both nights was pooled for each point, and teak plantation
data was averaged. (a) shows the total number of detections for each land use type, and (b)
shows the total species richness for each land use type.
523

References

Alvarado-Robledo, E. J., Hernández-Velázquez, I. M., Guillén-Navarro, K., Diego-García, E.,
Zarza, E., & Zamora-Briseño, J. A. (2024). Metabarcoding: Opportunities for accelerating
monitoring and understanding insect tropical biodiversity. *Journal of Insect Conservation*,
28(4), 589–604. https://doi.org/10.1007/s10841-024-00584-1

Báldi, A. (2003). Using higher taxa as surrogates of species richness: A study based on 3700
Coleoptera, Diptera, and Acari species in Central-Hungarian reserves. *Basic Appl. Ecol*,
4, 589–593. https://doi.org/10.1078/1439-1791-00193

Basset, Y., & Lamarre, G. P. A. (2019). Toward a world that values insects. *Science*, *364*(6447),
1230–1231. https://doi.org/10.1126/science.aaw7071

Beuchert, J. (2024). *Prototyping a Low-Cost, Low-Power System for Automated Monitoring of
Nocturnal Insects*.

Bjerge, K., Nielsen, J. B., Sepstrup, M. V., Helsing-Nielsen, F., & Høye, T. T. (2021). An
Automated Light Trap to Monitor Moths (Lepidoptera) Using Computer Vision-Based
Tracking and Deep Learning. *Sensors (Basel, Switzerland)*, *21*(2), 343.
https://doi.org/10.3390/s21020343

Brehm, G. (2017). A new LED lamp for the collection of nocturnal Lepidoptera and a spectral
comparison of light-trapping lamps. *Nota Lepidopterologica*, *40*, 87–108.
https://doi.org/10.3897/nl.40.11887

Brehm, G., & Axmacher, J. C. (2006). A Comparison of Manual and Automatic Moth Sampling
Methods (Lepidoptera: Arctiidae, Geometridae) in a Rain Forest in Costa Rica.
Environmental Entomology, *35*(3), 757–764. https://doi.org/10.1603/0046-225X-35.3.757

Bubnicki, J. W., Norton, B., Baskauf, S. J., Bruce, T., Cagnacci, F., Casaer, J., Churski, M.,
Cromsigt, J. P. G. M., Farra, S. D., Fiderer, C., Forrester, T. D., Hendry, H., Heurich, M.,
Hofmeester, T. R., Jansen, P. A., Kays, R., Kuijper, D. P. J., Liefting, Y., Linnell, J. D. C.,

… Desmet, P. (2024). Camtrap DP: An open standard for the FAIR exchange and

archiving of camera trap data. *Remote Sensing in Ecology and Conservation*, *10*(3),

283–295. https://doi.org/10.1002/rse2.374

Buttlar, M. (2024). *Mothbeam · CircuitHub*.

https://circuithub.com/projects/Moritz/Mothbeam/revisions/57895/parts

Camera | diopsis.eu. (n.d.). Retrieved September 18, 2024, from https://diopsis.eu/en/camera/

Campbell, C. J., Barve, V., Belitz, M. W., Doby, J. R., White, E., Seltzer, C., Di Cecco, G.,

Hurlbert, A. H., & Guralnick, R. (2023). Identifying the identifiers: How iNaturalist

facilitates collaborative, research-relevant data generation and why it matters for

biodiversity science. *BioScience*, *73*(7), 533–541. https://doi.org/10.1093/biosci/biad051

Chowdhury, S., Jennions, M. D., Zalucki, M. P., Maron, M., Watson, J. E. M., & Fuller, R. A.

(2023). Protected areas and the future of insect conservation. *Trends in Ecology &*

Evolution, *38*(1), 85–95. https://doi.org/10.1016/j.tree.2022.09.004

Cole, R. J., Werden, L. K., Arroyo, F. C., Quirós, K. M., Cedeño, G. Q., & Crowther, T. W.

(2024). Forest restoration in practice across Latin America. *Biological Conservation*, *294*,

110608. https://doi.org/10.1016/j.biocon.2024.110608

Collen, B., Ram, M., Zamin, T., & McRae, L. (2008). The Tropical Biodiversity Data Gap:

Addressing Disparity in Global Monitoring. *Tropical Conservation Science*, *1*(2), 75–88.

https://doi.org/10.1177/194008290800100202

Crouzeilles, R., Ferreira, M. S., Chazdon, R. L., Lindenmayer, D. B., Sansevero, J. B. B.,

Monteiro, L., Iribarrem, A., Latawiec, A. E., & Strassburg, B. B. N. (2017). Ecological

restoration success is higher for natural regeneration than for active restoration in

tropical forests. *Science Advances*, *3*(11), e1701345.

https://doi.org/10.1126/sciadv.1701345

DECRETO LEY No. 1. (1994). Autoridad Nacional del Ambiente.

https://difor.miambiente.gob.pa/wp-content/uploads/2025/03/Ley-1-del-3-de-febrero-de-

1994.pdf

Digital Naturalism Laboratories. (2025a, May 15). *Mothbox OSHWA Certification*. Open Source

Hardware Association Certification. https://certification.oshwa.org/pa000003.html

Digital Naturalism Laboratories. (2025b, October 31). *Mothbeam OSHWA Certification*. Open

Source Hardware Association Certification. https://certification.oshwa.org/pa000004.html

Duran, D. P., Timar, M., & Rothauser, B. (2022a). Single Night Surveys of Moth Communities

Can Serve as Ultra-Rapid Biodiversity Assessments. *Insects*, *13*(12), 1135.

https://doi.org/10.3390/insects13121135

Duran, D. P., Timar, M., & Rothauser, B. (2022b). Single Night Surveys of Moth Communities

Can Serve as Ultra-Rapid Biodiversity Assessments. *Insects*, *13*(12), 1135.

https://doi.org/10.3390/insects13121135

Dyer, L. A., Singer, M. S., Lill, J. T., Stireman, J. O., Gentry, G. L., Marquis, R. J., Ricklefs, R.

E., Greeney, H. F., Wagner, D. L., Morais, H. C., Diniz, I. R., Kursar, T. A., & Coley, P.

D. (2007). Host specificity of Lepidoptera in tropical and temperate forests. *Nature*,

448(7154), 696–699. https://doi.org/10.1038/nature05884

Ermgassen, S. O. S. E. zu, & Löfqvist, S. (2024). Financing ecosystem restoration. *Current*

Biology, *34*(9), R412–R417. https://doi.org/10.1016/j.cub.2024.02.031

Fabian, S. T., Sondhi, Y., Allen, P. E., Theobald, J. C., & Lin, H.-T. (2024). Why flying insects

gather at artificial light. *Nature Communications*, *15*(1), 689.

https://doi.org/10.1038/s41467-024-44785-3

Forrester, T., O'Brien, T., Fegraus, E., Jansen, P., Palmer, J., Kays, R., Ahumada, J., Stern, B.,

& McShea, W. (2016). An Open Standard for Camera Trap Data. *Biodiversity Data*

Journal, *4*, e10197. https://doi.org/10.3897/BDJ.4.e10197

Hahn, M., & Brühl, C. A. (2016). The secret pollinators: An overview of moth pollination with a

focus on Europe and North America. *Arthropod-Plant Interactions*, *10*(1), 21–28.

https://doi.org/10.1007/s11829-016-9414-3

Hill, A. P., Prince, P., Piña Covarrubias, E., Doncaster, C. P., Snaddon, J. L., & Rogers, A.

(2017). AudioMoth: Evaluation of a smart open acoustic device for monitoring

biodiversity and the environment. *Methods in Ecology and Evolution*, *9*(5), 1199–1211.

https://doi.org/10.1111/2041-210X.12955

Hobern, D. (2021, October 16). *Autonomous Moth Trap Project – Stangeia*.

https://stangeia.hobern.net/autonomous-moth-trap-project/

Høye, T. T., Ärje, J., Bjerge, K., Hansen, O. L. P., Iosifidis, A., Leese, F., Mann, H. M. R.,

Meissner, K., Melvad, C., & Raitoharju, J. (2021). Deep learning and computer vision will

transform entomology. *Proceedings of the National Academy of Sciences*, *118*(2),

e2002545117. https://doi.org/10.1073/pnas.2002545117

Huebner, S., Palmer, M., & Packer, C. (2024). Human Supervision is Key to Achieving Accurate

AI-assisted Wildlife Identifications in Camera Trap Images. *Citizen Science: Theory and

Practice*, *9.* https://doi.org/10.5334/cstp.752

Janzen, D. H. (1987). How Moths Pass the Dry Season in a Costa Rican Dry Forest.

International Journal of Tropical Insect Science, *8*(4), 489–500.

https://doi.org/10.1017/S1742758400022530

Kays, R., Arbogast, B. S., Baker-Whatton, M., Beirne, C., Boone, H. M., Bowler, M., Burneo, S.

F., Cove, M. V., Ding, P., Espinosa, S., Gonçalves, A. L. S., Hansen, C. P., Jansen, P.

A., Kolowski, J. M., Knowles, T. W., Lima, M. G. M., Millspaugh, J., McShea, W. J.,

Pacifici, K., … Spironello, W. R. (2020). An empirical evaluation of camera trap study

design: How many, how long and when? *Methods in Ecology and Evolution*, *11*(6), 700–

713. https://doi.org/10.1111/2041-210X.13370

Korsch, D., Bodesheim, P., Brehm, G., & Denzler, J. (2023). *Automated Visual Monitoring of

Nocturnal Insects with Light-based Camera Traps* (No. arXiv:2307.15433). arXiv.

https://doi.org/10.48550/arXiv.2307.15433

Leorna, S., & Brinkman, T. (2022). Human vs. machine: Detecting wildlife in camera trap

628 images. *Ecological Informatics*, *72*, 101876. https://doi.org/10.1016/j.ecoinf.2022.101876

629 *LTER*. (n.d.). LTER. Retrieved November 24, 2025, from https://lternet.edu/

630 McGeachie, W. J. (1989). The effects of moonlight illuminance, temperature and wind speed on

631 light-trap catches of moths. *Bulletin of Entomological Research*, *79*(2), 185–192.

632 https://doi.org/10.1017/S0007485300018162

633 Möglich, J. M., Lampe, P., Fickus, M., Younis, S., Gottwald, J., Nauss, T., Brandl, R., Brändle,

634 M., Friess, N., Freisleben, B., & Heidrich, L. (2023). Towards reliable estimates of

635 abundance trends using automated non-lethal moth traps. *Insect Conservation and*

636 *Diversity*, *16*(5), 539–549. https://doi.org/10.1111/icad.12662

637 Montgomery, G. A., Belitz, M. W., Guralnick, R. P., & Tingley, M. W. (2021). Standards and Best

638 Practices for Monitoring and Benchmarking Insects. *Frontiers in Ecology and Evolution*,

639 *8*. https://doi.org/10.3389/fevo.2020.579193

640 *Mothbox Github*. (2024). Mothbox. https://digital-naturalism-laboratories.github.io/Mothbox/

641 *NSF NEON*. (n.d.). Retrieved November 24, 2025, from https://www.neonscience.org/

642 Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You Only Look Once: Unified, Real-

643 Time Object Detection. *Conference on Computer Vision and Pattern Recognition*.

644 https://doi.org/10.48550/arXiv.1506.02640

645 Rhodes, M. W., Bennie, J. J., Spalding, A., ffrench-Constant, R. H., & Maclean, I. M. D. (2022).

646 Recent advances in the remote sensing of insects. *Biological Reviews*, *97*(1), 343–360.

647 https://doi.org/10.1111/brv.12802

648 Rydell, J., Entwistle, A., & Racey, P. A. (1996). Timing of Foraging Flights of Three Species of

649 Bats in Relation to Insect Activity and Predation Risk. *Oikos*, *76*(2), 243–252.

650 https://doi.org/10.2307/3546196

651 Schmeller, D. S., Böhm, M., Arvanitidis, C., Barber-Meyer, S., Brummitt, N., Chandler, M.,

652 Chatzinikolaou, E., Costello, M. J., Ding, H., García-Moreno, J., Gill, M., Haase, P.,

653 Jones, M., Juillard, R., Magnusson, W. E., Martin, C. S., McGeoch, M., Mihoub, J.-B.,

Pettorelli, N., … Belnap, J. (2017). Building capacity in biodiversity monitoring at the

global scale. *Biodiversity and Conservation*, *26*(12), 2765–2790.

https://doi.org/10.1007/s10531-017-1388-7

Scudder, G. G. E. (2017). The Importance of Insects. In *Insect Biodiversity* (pp. 9–43). John

Wiley & Sons, Ltd. https://doi.org/10.1002/9781118945568.ch2

Sickel, W., Zizka, V., Scherges, A., Bourlat, S. J., & Dieker, P. (2023). Abundance estimation

with DNA metabarcoding – recent advancements for terrestrial arthropods.

Metabarcoding and Metagenomics, *7*, e112290. https://doi.org/10.3897/mbmg.7.112290

Sittinger, M., Uhler, J., Pink, M., & Herz, A. (2024). Insect detect: An open-source DIY camera

trap for automated insect monitoring. *PLOS ONE*, *19*(4), e0295474.

https://doi.org/10.1371/journal.pone.0295474

Stevens, S., Wu, J., Thompson, M. J., Campolongo, E. G., Song, C. H., Carlyn, D. E., Dong, L.,

Dahdul, W. M., Stewart, C., Berger-Wolf, T., Chao, W.-L., & Su, Y. (2024). *BioCLIP: A*

Vision Foundation Model for the Tree of Life. 19412–19424.

https://openaccess.thecvf.com/content/CVPR2024/html/Stevens_BioCLIP_A_Vision_Fo

undation_Model_for_the_Tree_of_Life_CVPR_2024_paper.html

Tyler, T. (2020). Relationship between moth (night active Lepidoptera) diversity and vegetation

characteristics in southern Sweden. *Journal of Insect Conservation*, *24*(6), 1005–1015.

https://doi.org/10.1007/s10841-020-00270-y

UKCEH AMI system | UK Centre for Ecology & Hydrology. (n.d.). Retrieved September 18,

2024, from https://www.ceh.ac.uk/solutions/equipment/automated-monitoring-insects-

trap

Vélez, J., McShea, W., Shamon, H., Castiblanco-Camacho, P. J., Tabak, M. A., Chalmers, C.,

Fergus, P., & Fieberg, J. (2023). An evaluation of platforms for processing camera-trap

data using artificial intelligence. *Methods in Ecology and Evolution*, *14*(2), 459–477.

https://doi.org/10.1111/2041-210X.14044

Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R., & Stopak, D. (2021). Insect

decline in the Anthropocene: Death by a thousand cuts. *Proceedings of the National*

Academy of Sciences, *118*(2), e2023989118. https://doi.org/10.1073/pnas.2023989118

Wauchope, H. S., zu Ermgassen, S. O. S. E., Jones, J. P. G., Carter, H., Schulte to Bühne, H.,

& Milner-Gulland, E. J. (2024). What is a unit of nature? Measurement challenges in the

emerging biodiversity credit market. *Proceedings of the Royal Society B: Biological*

Sciences, *291*(2036), 20242353. https://doi.org/10.1098/rspb.2024.2353

Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., Robertson, T., &

Vieglais, D. (2012). Darwin Core: An Evolving Community-Developed Biodiversity Data

Standard. *PLOS ONE*, *7*(1), e29715. https://doi.org/10.1371/journal.pone.0029715

Wong, M. K. L., & Didham, R. K. (2024). Global meta-analysis reveals overall higher nocturnal

than diurnal activity in insect communities. *Nature Communications*, *15*(1), 3236.

https://doi.org/10.1038/s41467-024-47645-2

Zou, Y., Van der Werf, W., Liu, Y., & Axmacher, J. (2020). Predictability of species diversity by

family diversity across global terrestrial animal taxa. *Global Ecology and Biogeography*.

https://doi.org/10.1111/geb.13043