1   Mothbox and Mothbot: automated light trap and data processing system for scalable insect

2   monitoring

3   Hubert A. Szczygieł[1,2]*, Brianna Johns[3,4], Bernat Fortet[5], Daisy H. Dent[2,6,7], Kit Quitmeyer[4],
4   Andrew Quitmeyer[4,8]
5

[1] Department of Biological and Environmental Sciences, Gothenburg Global Biodiversity
Centre, University of Gothenburg, Medicinaregatan 7B, 413 90 Göteborg, Sweden
[2] Smithsonian Tropical Research Institute; Apartado 0843-03092, Balboa, Ancón, República
de Panamá.
[3] School of Biological Sciences, Monash University, Clayton, Victoria, Australia, 3800
[4] Digital Naturalism Laboratories, Sibert Ave 123B, Gamboa, Colón Province, Panama
[5] Restoration Scope, 411 Walnut Street #24052, Green Cove Springs, FL 32043-3443, USA
[6] Department of Environmental Systems Science, ETH, Zürich, Switzerland.
[7] Max Planck Institute of Animal Behavior, Konstanz, Germany
[8] Department of Human Centered Design & Engineering, University of Washington, 3960
Benton Lane NE 428, Seattle, WA 98195, USA

*Corresponding author. Email: hubert.70@gmail.com

Abstract

1. Insects represent the most diverse group of organisms on Earth, and comprise the

majority of known species, yet they are seldom accounted for in large-scale

biodiversity monitoring systems and conservation planning.

2. We have developed the *Mothbox* - an open source automated light trap that makes

insect monitoring accessible to non-specialists and scalable for scientific and

conservation purposes - and the *Mothbot* - a machine learning tool and user

interface for processing data from automated light traps. The Mothbox is portable,

durable, and low-cost, while Mothbot prioritizes human-in-the-loop data validation

of computer vision outputs to ensure data quality and aid in developing custom

image reference libraries.

3. Mothbox has been extensively tested in field conditions, with > 185 deployments,

spanning > 450 nights and > 100 sampling locations. We present proof-of-concept

studies that examine shifts in insect activity and richness across i. time intervals

throughout the night at a single location, and ii. different habitat types, to assess

whether these patterns can be captured through automated sampling with Mothbox

hardware and Mothbot processing.

4. As the field of automated insect monitoring continues to develop, Mothbox provides

an affordable entry point for a range of stakeholders. It enables entomologists to

explore new research questions by tracking high temporal resolution changes in

insect activity, as well as systematically monitoring insects across landscapes.

Mothbox will help build global insect-monitoring capacity via autonomous sampling,

improving our ability to detect biodiversity loss and guide effective conservation

action.

Keywords: biodiversity; monitoring; computer vision; insects; light trap; open source;

artificial intelligence

1 Introduction

The majority of the world's species are insects, and most insects are active at night

(Wong & Didham, 2024). Insects are incredibly valuable ecologically and economically, and

rapidly declining in many parts of the world (Basset & Lamarre, 2019). Specialisation

makes insects particularly relevant biodiversity indicators; for example, lepidoptera larvae

often feed on only one species or genus of plant (Dyer et al., 2007), and adults can similarly

be adapted to pollinate single host plant species (Hahn & Brühl, 2016), thus, monitoring

Lepidoptera can provide inferences about the plant community (Duran et al., 2022; Tyler,

2020). However, insects are rarely included in conservation planning, despite their great

diversity, ecological significance, and evidence of global decline, because of the difficulty of

conducting standardised large-scale insect monitoring (Chowdhury et al., 2023; Høye et al.,

2021).

Traditional approaches to monitoring nocturnal insects require specialised

knowledge and considerable time investment (Montgomery et al., 2021). Light-sheeting, i.e.

attracting nocturnal insects to an illuminated sheet, is an effective method for visually

assessing insect populations or targeting particular taxa. However, documenting every

insect is often impractical and requires researchers to work through the night (Duran et al.,

2022). Bucket traps, which attract and capture insects, do not require ongoing supervision,

but are typically deployed for the purpose of lethal collection and require extensive manual

sample processing once collected (Brehm & Axmacher, 2006). New technological

approaches to biodiversity monitoring also face unique challenges when applied to insects.

For example, genetic approaches (e.g. DNA metabarcoding) remain expensive, rely on

reference libraries with limited coverage for many regions and taxa (Alvarado-Robledo et

al., 2024), and are poorly suited to assessing abundance (Sickel et al., 2023), while satellite-

based remote-sensing lacks the resolution for direct monitoring of insects (Rhodes et al.,

2022). To bypass difficulties with these monitoring methods, the first automated light trap

that used a programmed light source and computer vision for identification was developed

in 2021 (Bjerge et al., 2021). This proof of concept inspired the creation of a range of

automated insect monitoring systems, including the Automatic Moth Trap (Möglich et al.,

2023), Automated Monitoring of Insects (Lord et al., 2026), DIOPSIS (Huijbers et al., 2025),

and others (Beuchert, 2024; Hobern, 2021; Korsch et al., 2023). Automated non-lethal

insect monitoring remains an emerging field, and existing devices, including the one

presented in this manuscript, represent complementary approaches that balance cost,

portability, image resolution, insect-attraction lighting, 4G connectivity, ease of

modification, and durability as the research community collectively works toward further

technological refinement.

2. Mothbox and Mothbot: An open-source automated light trap and image-

processing platform

Here we describe the Mothbox, an automated light trap designed for scalable

insect monitoring, and Mothbot, a platform for automated light trap data processing

and validation. The Mothbox device integrates a low-cost Raspberry Pi computer, a

high-resolution camera, and specialized attractant lights to capture images of

nocturnal insects. The Mothbox has been developed specifically to optimize

portability, adaptability, high-resolution images, and weather resistance.

Furthermore, the low materials cost of the latest model (Mothbox Pro; approximately

USD 375) enables the simultaneous deployment of multiple units. Mothbot is an

open-source tool designed to extract insect occurrence data from images captured by

Mothbox and similar devices. We present two proof-of-concept studies from Panama

that employ the Mothbox and Mothbot to detect and identify night-flying insects,

demonstrating the potential for biodiversity monitoring and ecological research.

2.1 Terminology

The emerging field of automated insect monitoring is generating new concepts and components that require clear and consistent terminology. Inconsistent naming across research groups has created challenges in comparing monitoring systems and protocols. To promote standardization, we propose a set of terms to describe key aspects of hardware design, data collection, and data analysis. Where possible, our terminology is aligned with existing standards for automated biodiversity data collection (Bubnicki et al., 2024; Wieczorek et al., 2012). We use *deployment* to describe a discrete sampling interval associated with one location and one device; *sourceImage* to describe the whole un-cropped image taken by an automated insect monitoring device; and *detection* to refer to a specific subset of a sourceImage that covers only a portion of the sourceImage that contains one item or organism (see Table S3 for a list of terminology).

2.2 Mothbox

The Mothbox is an automated light trap for insect monitoring designed to produce accurate insect censuses, while maintaining affordability, durability and portability (Fig. 1). Two Mothbox versions are currently available: Mothbox DIY, which emphasizes device adaptability and supports do-it-yourself (DIY) construction, and Mothbox Pro, which is designed for streamlined manufacturing and large-scale deployment. Full descriptions of components and construction for both Mothbox versions is fully open source with a CERN-OHL-W license (Digital Naturalism Laboratories, 2025a), the operating firmware has an MIT licence, and the documentation is CC BY 4.0. Full specifications, including instructions for construction, programming, use, and adaptation of the Mothbox are on Github (*Mothbox Github*, 2024), and on the Mothbox website (https://mothbox.org/).

121   The Mothbox consists of a Raspberry Pi 5 computer that controls a 64 megapixel

122 Arducam Owlsight camera (9248 × 6944 resolution), custom 'Mothbeam' lights to attract

123 nocturnal insects, and lights to illuminate the insects for photography (like a camera flash)

124 (Fig. 2). The system is housed in a watertight plastic box and powered by a 156Wh lithium

125 battery. All electronic components are within the watertight housing, however ports in the

126 base of the housing allow for connection to a solar panel, additional batteries, or external

127 light sources. Mothbox Pro models also include an ambient light sensor, internal

128 temperature sensor, slot for attachment of an optional external temperature sensor, and

129 slots for adding an optional GPS unit. Both Mothbox DIY and Mothbox Pro can be

130 programmed via editing a CSV file on the SD card (by plugging the SD card into a different

131 computer or accessing it remotely via a Virtual Network Computing (VNC) system. The Pro

132 model also comes with the addition of manual switches for deployment schedule setting as

133 a backup for easily adjusting settings in the field (Fig. 2). The camera faces a target panel; a

134 29 cm x 20.5 cm x 0.3 cm acrylic sheet with felted acrylic fabric on one side to create a

135 textured surface diffusely illuminated by the attractant light. The target panel is attached to

136 the Mothbox using clear acrylic arms positioned 28.5cm from the camera. This gives a

137 functional resolution of approximately 625 px / mm^2 per photograph. At this resolution

138 most insects > 3mm in length can be identified to family level (Fig. 3). The Mothbox can be

139 adapted to use larger target sheets at greater distances from the cameras, which will result

140 in functionally lower resolution photographs (fewer pixels per insect). Mothbox can also be

141 positioned to focus on any flat surface as the camera auto-focuses. Mothbox Pro dimensions

142 are 25.5 x 18 x 13 cm, the arms are 39 x 7 cm, and a full kit with spare arms and a strap

143 weighs 2.7 kg

144   We have prioritised ease of Do-It-Yourself construction and repair in the Mothbox

145 DIY design. Minimal tools are required for construction (drill, screw driver, wire strippers,

146 epoxy, and a 53mm hole saw drill bit), while most repairs can be managed with a

147 screwdriver, knife, and a hot glue stick. We use lever nuts for electrical connections to

148 avoid soldering to enable modification and repair. The internal framework, arms and target

149 panel can be cut from a single sheet of 50 x 60 x 0.3 cm acrylic. While still 3D-printable, this

150 flat-pack design choice enables easy field servicing, as replacement components can be cut

151 and drilled by hand from acrylic or wood. To set up a Mothbox requires loading a ready-to-

152 go operating system image onto an SD card; no coding expertise is needed. Users can

153 change the sampling schedule by editing a scheduling CSV file in the data storage drive.

154 Mothbox Pro combines the electronic components of Mothbox DIY into a single printed

155 circuit board (PCB), which decreases construction time, number of components, and cost.

156 Mothbox Pro electronics are housed in a custom 3D-printed box, while DIY are in a Plano

157 1460-00 clear acrylic box. Motion-activated photo saving was the default on the original

158 automated light trap prototype (Bjerge et al., 2021) however for the Mothbox we use time-

159 lapse (photos taken at regular intervals). Time-lapse imaging reduces computational load

160 (and thus extending battery life and eliminating the need for higher-cost Raspberry Pi

161 models), however it can also miss rapid insect activity between frames, potentially

162 underestimating abundance relative to high-sensitivity motion-detection systems.

163 Mothboxes can be programmed for any sampling schedule with no hardware modification.

164 The current default sampling schedule which we use in the tropics is five one-hour

165 intervals at 19:00, 21:00, 23:00, 02:00, and 04:00, with photographs captured every minute

166 (300 images; ~5.2 GB per night). This intermittent design samples insect assemblages

throughout the night, reduces battery consumption and allows the target panel to clear

between sessions, promoting insect turnover and reducing repeated observations of the

same individuals. Mothbox DIY lasts for 10-12 hours of runtime per battery, or two nights

with the default schedule, while Mothbox Pro can last for up to 18 hours, depending on

light intensity settings. The single integrated battery can be augmented with an additional

box containing up to three more batteries connected in parallel, increasing total run time. A

second Plano box with a charging port added to the bottom works for this purpose. The

Mothbox can also be connected to portable solar panels (up to 100 watt, 12-20 volt output)

and mains power via charging ports in the bottom.

The relatively small size and durability enables the Mothbox to be used in a range of

habitats and locations. Once the arms and target panel are in place, and the Mothbox is

switched on, it can be mounted to a tree using a strap through the handle (Fig 1), mounted

to tripods with the addition of a tripod mount, positioned in the canopy using a throw line,

or placed on the ground. Deploying a Mothbox on the ground reduces the range of the

attractant lights and gives access to predatory ants and spiders, so elevating the device is

advised. After collection, photos are retrieved via a USB drive or SD card and processed

with Mothbot.

2.3 Mothbeam

The Mothbeam is an open-source printed circuit board (PCB) with light-emitting

diodes (LEDs) that emit a range of light frequencies to attract nocturnal insects. The

Mothbeam, originally created by Moritz Buttlar of LabLab (Buttlar, 2024), was designed in

collaboration with the Mothbox team as an inexpensive, open-source equivalent of a

LepiLED (Brehm, 2017). There are two Mothbeam configurations, one which targets violet

light (365, 395, 405 nm) and another that targets visible light (full-spectrum white, 450,

520 nm). Mothbeams have four switches that control light intensity. No switches flipped

produces the least powerful output (10 mA), two switches add 70mA each, and two

switches add 150 mA each. We currently do not advise having all switches engaged at the

same time as this might result in overheating the Mothbeam. By default, each Mothbox 4.5

comes with one of each PCB and both PCBs are activated when collecting data. Mothbox 5

has three LED PCB slots. Mothboxes can be programmed to only use one PCB, and the PCBs

can be modified to deactivate individual LEDs or dim light intensity, to target particular

taxonomic groups or reduce battery consumption. We have also designed stand-alone

Mothbeams for use with traditional moth sheeting. Like the Mothbox, Mothbeams are fully

open source with a CERN-OHL-W licence (Digital Naturalism Laboratories, 2025b).

2.4 Mothbot

Mothbot is an open-source machine learning tool and user interface for processing

data from automated light traps. Mothbot consists of two user interfaces: 'Mothbot Process'

and 'Mothbot Classify'. The first automatically detects insects from source images, identifies

them to a user-defined taxonomic threshold, then spatially clusters them. The second

facilitates rapid human validation of computer vision identification and manual finer-

taxonomy classification, as well as data export. Mothbot Process uses a combination of two

open-sourced AI programs: YOLO (Redmon et al., 2016) and BioCLIP (Stevens et al., 2024).

A custom YOLO11 model trained on Mothbox data from 4500 hand-labelled images

automatically detects and crops individual insects (down to ~1 mm total length) from

novel imagery. These cropped detections are then fed to a modified BioCLIP model. The

model outputs are restricted to taxa on a user-defined species list, generated with

taxonomic and geographic filters on GBIF. A custom algorithm then clusters the detections,

extracting the detections' visual features with DINOv2

(https://github.com/facebookresearch/dinov2) and then grouping the detections by

similarity with HDBSCAN (Hierarchical Density-Based Spatial Clustering of Applications

with Noise - https://pypi.org/project/hdbscan/). These visual clusters are sub-grouped

temporally (i.e. by the time their source image was taken) to create hierarchical groupings

of detected organisms that exist in temporal 'tracks' over a set of sequentially-taken

images.

Processing data with Mothbot Process starts with selecting file paths to a folder

containing Mothbox JPG photos, a CSV file that contains deployment metadata, and a CSV

file containing a species list. The formatting for the metadata CSV file is based on Camtrap

DP (Bubnicki et al., 2024) and Darwin Core (Wieczorek et al., 2012) (Table S1). The species

list CSV file is in the Global Biodiversity Information Facility (GBIF) downloads format

(Table S2), and requires cleaning to remove synonyms and other errors (SI Script 1). For

our proof-of-concept studies, our species list included all insect species recorded on GBIF

in Panama and Costa Rica. We tried running Mothbot with a species list that also included

all arachnids, however, we found that model accuracy decreased with this addition, and

data validation was more time-efficient when manually correcting arachnid identifications.

After selecting the required file paths, the photos can be processed using the tabs on in the

Mothbot Process user interface. The first tab finds detections on the source image, the

second identifies them, the third adds metadata to JSON files associated with each

sourceImage, the fourth clusters detections perceptually, and the fifth adds EXIF data,

including GPS coordinates and identification, to cropped detection photos.

Mothbot Classify is a React-based user interface is designed for human validation of

the BioCLIP identifications. Mothbot Classify includes an easily searchable, hierarchical

taxonomic classification system, and allows designation of morphospecies that can be

linked to iNaturalist observations. The interface exports occurrence data for each organism

photographed to a Darwin Core formatted CSV file (Wieczorek et al., 2012). The human-in-

the-loop workflow for refining taxonomy is not required for generating an output CSV file,

but we believe it is currently an essential step in generating high-quality data from

automated insect monitoring devices, as it is for mammal camera trap data (Huebner et al.,

2024; Leorna & Brinkman, 2022).

3. Proof of concept: Field deployment in Panama

To demonstrate the capabilities of Mothbox devices in field conditions, we

conducted proof-of-concept studies examining shifts in insect activity and richness across

1. intra-night temporal dynamics at a single location, and 2. different habitat types assessed

simultaneously, to assess whether patterns could be captured through automated sampling

with Mothbox hardware and Mothbot processing.

Mothbox development has focused on creating devices that can be deployed in off-

grid, remote tropical locations, as no other currently available automated light traps are

suited for this use case. The tropics pose some of the most challenging environmental

conditions for electronic equipment, including high humidity, elevated temperatures, and

biological agents (e.g. insects, rodents and fungi) capable of infiltrating or degrading

devices, making tropical forests optimal testing environments for Mothbox robustness and

durability. In April 2025, we tested five Mothbox DIY units in two study locations:

1. We deployed one Mothbox with an additional battery for four continuous nights (10

- 13 April 2025) in a tropical forest in the southern Azuero Peninsula, Panama to

assess intra-night temporal shifts in insect assemblages.

2. We deployed three Mothboxes simultaneously in a teak plantation, a regenerating

forest stand and a neighbouring mature forest near Venao in the southern Azuero

Peninsula, Panama for two consecutive nights (11 - 13 April 2025) to assess shifts in

insect assemblages with habitat type. Mothbox sampling points were separated by

at least 250m.

For both studies we used Mothbox DIY devices mounted on trees at chest height,

programmed with a sampling schedule of one-hour sampling sessions starting at 19:00,

21:00, 23:00, 02:00, and 04:00. The Mothboxes were programmed to take a photo every

minute, and thus generate 300 images per night. After collecting the Mothboxes from the

field, we processed the data using the Mothbot interface set to order-level automatic

identification, then manually identified each insect detected to the finest taxonomic level

possible and assigned morphospecies where necessary.

4 Performance

4.1 Detection of patterns in insect number and diversity

In the first study, we recorded 6,271 detections from nine orders, 43 families, and 219

unique morphospecies to assess intra-night temporal shifts in insect assemblages. Only 37

of the 219 morphospecies could be identified to species level by a general entomologist

(not an expert taxonomist), the remaining 183 (84%) were assigned morphospecies

names. A total of 1046 images were collected over 3 nights (i.e. 19 hours; no data collected

04:00-05:00 on night 4 due to early battery discharge. The target panel had no insects at

the beginning of each hour-long session (Fig. 4 c and d), confirming that when the

Mothbeam lights were off, all insects left the target panel encouraging insect turnover and

reducing repeat recordings. Most insects attracted to the Mothbeams fly around the

Mothbox for several minutes before they settle, as they do on conventional moth sheets.

During each 1-hr session, the number of detections per photo (Fig. 4 c) and morphospecies

richness (Fig. 4 d) increased with time since attractant lights were turned on. The earliest

sampling session (19:00-20:00) had the most cumulative detections and the highest

morphospecies richness compared to later sampling sessions (Fig. 4 a and b; P =0.0002 and

0.0021, respectively, ANOVA).

In the second study, we recorded 2,755 detections in total with 8 orders and 40

families represented, and a total of 158 unique morphospecies (with 94% designated

morphospecies) across four sampling sites to quantify shifts in insect numbers and species

richness with habitat type. Mature forest and native restoration had higher numbers of

detections and morphospecies richness (1058 detections / 81 morphospecies and 1140

detections / 64 morphospecies, respectively) than teak plantation (138 detections and 19

morphospecies). Mature forests had higher species richness than the native restoration

forest, while native restoration had higher numbers of detections (Fig. 5). Overall, our

results illustrate that the Mothbox and Mothbot can effectively detect differences in insect

activity and richness across temporal and land-use gradients.

4.2 Identification rates among taxonomic groups

Large, distinctively coloured insects which do not need dissection or microscopic

examination were more likely to be identifiable at finer taxonomic levels. For example, of

the species which were identified to species level, 87.5% were Lepidoptera, mostly in

families Crambidae, Erebidae, and Noctuidae. Conversely, none of the 297 Diptera

detections were identified to species level, as most of them were quite small. Some insect

detections had poor image resolution due to their small size, and consequently could not be

confidently identified to even Order level. For example, it can be difficult to differentiate

between the tiniest Diptera and Hymenoptera, or Coleoptera and Hemiptera. While not

employed for this analysis, Mothbot Classify has the ability to set size filters, which could be

used to narrow the scope of a study to monitoring only larger taxa.

5 Implications and Future Directions

5.1 Mothbox configuration and considerations

Mothbox can be used to monitor nocturnal insect activity, species richness and

composition and can be adapted for other scientific uses. For example, Mothboxes can be

modified to release pheromones or nectar from an artificial flower using the spare plug

already built-in (Fig. 2 j). If continuous power is available, a Mothbox unit can run on

continuous schedules and operate for entire field seasons rather than just short

deployments. The Mothbox functions as a standalone sensor, but integrates well with other

sensors. For example, acoustic monitors, conventional camera traps, and environmental

sensors (light, wind, humidity, precipitation) can all synchronize with Mothbox data

collection. This would help identify the drivers of changes in insect activity throughout the

night and or across seasons.

While Mothbox development has focused on deployment in remote tropical

locations, the system can be applied in any regions or ecosystem where there is an interest

in monitoring nocturnal insects. Inclusion of Mothboxes in large-scale ecological

monitoring networks like the Long Term Ecological Research (LTER) network and the

National Ecological Observatory Network (NEON) would yield an additional valuable

dataset for insect population trends (Crossley et al., 2020; Dantzer et al., 2023). Mothbox

would also be well suited to monitor pollinator activity and diversity in agriculture, or the

spread of invasive species.

5.2 Data processing and Mothbot development

The Mothbot Process interface allows users to select the taxonomic level for AI-

based identification. Model accuracy depends on the quality and quantity of reference data

(in this case training image datasets) and the presence of cryptic species, with finer

taxonomic resolution reducing accuracy and increasing manual validation time. In the

context of insect identification models, accuracy refers to the correctness of the automated

identification relative to a human validation. In our proof-of-concept studies, the model

was sufficiently accurate at the order level for rapid human verification, while species-level

identification remained unreliable in highly diverse tropical regions with limited reference

data. As more detections are manually identified and incorporated into training datasets,

future models are expected to achieve reliable family-level identification for most groups

and genus- or species-level identification for selected taxa. Our data analysis framework

assumes that automated insect monitoring systems support, rather than replace,

taxonomic experts, who remain essential for model training and systematic results

validation. We therefore prioritized taxonomic accuracy and ease of human verification

over fine-scale taxonomic resolution. Where expert taxonomists are unavailable, we

recommend uploading representative images of each morphospecies from each

deployment to iNaturalist for community-based verification (Campbell et al., 2023).

The accuracy of computer vision models improves when the training data closely

matches the data being classified, and Mothbox raw images provide ideal conditions:

subjects against a well-lit flat white background, at a fixed distance from the camera. The

same cannot be said for the massive and visually diverse datasets that underpin BioCLIP

and other current insect classifiers, which include many dead pinned and living but *in-situ*

insects. In time, with larger Mothbox datasets, we will augment the Mothbot system with a

new model based on human-validated data identified to a finer taxonomic resolution,

producing region-specific versions that can accurately identify taxa below family level. We

believe that this approach will rapidly change our ability to classify species so that all

insects photographed with automated insect monitors can be confidently identified to

Genus or Family level by Mothbot. A centralised repository for human-validated image

storage and model development in the realm of automated insect monitoring would greatly

aid the process of refining classification models, as it has for mammal camera trapping

(Vélez et al., 2023). In the meantime, higher taxonomic levels can be used as surrogates of

species-level data (Zou et al., 2020), an approach that is key to current large-scale insect

monitoring in the tropics while taxonomic expertise, genetic reference libraries, and

photographic reference libraries catch up with monitoring needs.

5 Conclusion

Mothbox aims to offer for insect monitoring what low-cost camera trapping offers

for mammals. By decreasing the time, cost, and specialised knowledge required for

monitoring insects, Mothbox enables standardised insect monitoring at the landscape scale.

Keeping taxonomists in the loop is critical for ensuring data quality given constraints of

insect classification models, and Mothbot offers a platform for rapid validation of computer

vision outputs. This technology increases capacity for monitoring global trends in insect

populations, and acts as a tool for exploring new ecological relationships and behavioural

patterns. Pairing Mothbox and Mothbot has the potential to drive development of human-

validated insect image repositories to support improved classification models. Importantly,

Mothbox will make biodiversity monitoring more accessible to community conservation

groups, empowering restoration and conservation.

Figures

Figure 1 Mothbox DIY (a) and Mothbox Pro (b) deployed in a tropical forest in Panama.

Figure 2 Front (left) and back (right) of Mothbox Pro printed circuit board (PCB) with
selected components in blue boxes. The front view is from a complete Mothbox looking in
through the clear acrylic front, the back view is a computer-aided design. **a** three
Mothbeam PCBs in the attractant light slots; **b** 64 MP Arducam camera; **c** example of three
LEDs for illuminating photos (54 total across the front of the PCB); **d** e-paper low-power
display with device name, battery status, photo status, and deployment settings; **e** ambient
light sensor; **f** Raspberry Pi 5 slot (real time clock battery module attaches to Pi); **g** battery
location (voltage regulator attaches to battery, which in turn is fastened to the PCB with zip
ties); **h** power raw input (9-36V); **i** 12 V regulator control; **j** external attractant slot; **k**
physical schedule programming switches (as a backup option for computer schedule
programming); **l** mode programming switches; **m** integrated power monitor; **n** optional 4
or 5 pin GPS slot; **o** temperature sensor with optional external temperature sensor slot.
399

Figure 3 Unedited photo taken by a Mothbox in Panama, with an inset to demonstrate the capacity for documenting small insects like this 1.5 mm long minute pirate bug (Anthocoridae). The resolution of the full photo is 9248 x 6944 pixels. The scale provided is in millimetres.

Figure 4 Insect activity proof-of-concept study: (a) total number of detections (individual organisms in each photo, for all photos) for five one-hour sampling sessions spaced throughout the night; (b) total species richness within the five sessions; (c) total number of insect detections in each photo by minute from start of sampling, for the five sampling sessions; (d) species richness per photo across the five sampling sessions. All four panels represent data averaged across the four nights of the deployment. The effect of session on cumulative detections and richness was significant (p = 0.0002 and 0.0021, respectively). Error bars in (a) and (b) represent standard error. This study demonstrates that insect activity and richness exhibit intra-night temporal dynamics detectable with automated monitoring

Figure 5 Land use proof-of-concept study. Three Mothboxes were deployed simultaneously for two nights at points with different land use histories. One point was in non-native teak (Tectona grandis) plantation, one point was in an area planted with native tree species, and one point in mature native forest. (a) shows mean detections per hour for each land use type, and (b) shows mean species richness per hour for each land use type.

442 **Author contributions:** H.A.S and AQ conceived the ideas and designed methodology; H.A.S,
443 B.J., K.Q. and AQ collected the data; H.A.S analyzed the data; H.A.S led the writing of the
444 manuscript; All authors contributed critically to the drafts and gave final approval for
445 publication.
446
Acknowledgments: The Mothbox project would not have been possible without many
contributors over the years. Special thank you to Wildlabs, Earthshot, Ponterra, Michigan
State University, Experiment.com and a Google Carbon Removal Research Award for
funding the development of the Mothbox. Thank you to Yash Sondhi for support on early
Mothbox development. Thank you to Pro Eco Azuero, Totumas, Daniel Murcia, Josue
Santos, Mulget Amaru, and Evidelio Garcia for helping us deploy Mothboxes across Panama.
Thank you to the Automated Insect Monitoring community for helping us define
terminology.
455

References

Alvarado-Robledo, E. J., Hernández-Velázquez, I. M., Guillén-Navarro, K., Diego-García, E.,

Zarza, E., & Zamora-Briseño, J. A. (2024). Metabarcoding: Opportunities for

accelerating monitoring and understanding insect tropical biodiversity. Journal of

Insect Conservation, 28(4), 589–604. https://doi.org/10.1007/s10841-024-00584-

1

Basset, Y., & Lamarre, G. P. A. (2019). Toward a world that values insects. Science,

364(6447), 1230–1231. https://doi.org/10.1126/science.aaw7071

Beuchert, J. (2024). Prototyping a Low-Cost, Low-Power System for Automated Monitoring

of Nocturnal Insects.

Bjerge, K., Nielsen, J. B., Sepstrup, M. V., Helsing-Nielsen, F., & Høye, T. T. (2021). An

Automated Light Trap to Monitor Moths (Lepidoptera) Using Computer Vision-

Based Tracking and Deep Learning. Sensors (Basel, Switzerland), 21(2), 343.

https://doi.org/10.3390/s21020343

Brehm, G. (2017). A new LED lamp for the collection of nocturnal Lepidoptera and a

spectral comparison of light-trapping lamps. Nota Lepidopterologica, 40, 87–108.

473  https://doi.org/10.3897/nl.40.11887

474 Brehm, G., & Axmacher, J. C. (2006). A Comparison of Manual and Automatic Moth Sampling

475  Methods (Lepidoptera: Arctiidae, Geometridae) in a Rain Forest in Costa Rica.

476  Environmental Entomology, 35(3), 757–764. https://doi.org/10.1603/0046-225X-

477  35.3.757

478 Bubnicki, J. W., Norton, B., Baskauf, S. J., Bruce, T., Cagnacci, F., Casaer, J., Churski, M.,

479  Cromsigt, J. P. G. M., Farra, S. D., Fiderer, C., Forrester, T. D., Hendry, H., Heurich, M.,

480  Hofmeester, T. R., Jansen, P. A., Kays, R., Kuijper, D. P. J., Liefting, Y., Linnell, J. D. C., …

481  Desmet, P. (2024). Camtrap DP: An open standard for the FAIR exchange and

482  archiving of camera trap data. Remote Sensing in Ecology and Conservation, 10(3),

483  283–295. https://doi.org/10.1002/rse2.374

484 Buttlar, M. (2024). Mothbeam · CircuitHub.

485  https://circuithub.com/projects/Moritz/Mothbeam/revisions/57895/parts

486 Campbell, C. J., Barve, V., Belitz, M. W., Doby, J. R., White, E., Seltzer, C., Di Cecco, G., Hurlbert,

487  A. H., & Guralnick, R. (2023). Identifying the identifiers: How iNaturalist facilitates

488  collaborative, research-relevant data generation and why it matters for biodiversity

489  science. BioScience, 73(7), 533–541. https://doi.org/10.1093/biosci/biad051

490 Chowdhury, S., Jennions, M. D., Zalucki, M. P., Maron, M., Watson, J. E. M., & Fuller, R. A.

491  (2023). Protected areas and the future of insect conservation. Trends in Ecology &

492  Evolution, 38(1), 85–95. https://doi.org/10.1016/j.tree.2022.09.004

493 Crossley, M. S., Meier, A. R., Baldwin, E. M., Berry, L. L., Crenshaw, L. C., Hartman, G. L.,

494  Lagos-Kutz, D., Nichols, D. H., Patel, K., Varriano, S., Snyder, W. E., & Moran, M. D.

495  (2020). No net insect abundance and diversity declines across US Long Term

Ecological Research sites. Nature Ecology & Evolution, 4(10), 1368–1376.

https://doi.org/10.1038/s41559-020-1269-4

Dantzer, B., Mabry, K. E., Bernhardt, J. R., Cox, R. M., Francis, C. D., Ghalambor, C. K., Hoke, K.

L., Jha, S., Ketterson, E., Levis, N. A., McCain, K. M., Patricelli, G. L., Paull, S. H., Pinter-

Wollman, N., Safran, R. J., Schwartz, T. S., Throop, H. L., Zaman, L., & Martin, L. B.

(2023). Understanding Organisms Using Ecological Observatory Networks.

Integrative Organismal Biology, 5(1), obad036.

https://doi.org/10.1093/iob/obad036

Digital Naturalism Laboratories. (2025a, May 15). Mothbox OSHWA Certification. Open

Source Hardware Association Certification.

https://certification.oshwa.org/pa000003.html

Digital Naturalism Laboratories. (2025b, October 31). Mothbeam OSHWA Certification.

Open Source Hardware Association Certification.

https://certification.oshwa.org/pa000004.html

Duran, D. P., Timar, M., & Rothauser, B. (2022). Single Night Surveys of Moth Communities

Can Serve as Ultra-Rapid Biodiversity Assessments. Insects, 13(12), 1135.

https://doi.org/10.3390/insects13121135

Dyer, L. A., Singer, M. S., Lill, J. T., Stireman, J. O., Gentry, G. L., Marquis, R. J., Ricklefs, R. E.,

Greeney, H. F., Wagner, D. L., Morais, H. C., Diniz, I. R., Kursar, T. A., & Coley, P. D.

(2007). Host specificity of Lepidoptera in tropical and temperate forests. Nature,

448(7154), 696–699. https://doi.org/10.1038/nature05884

Hahn, M., & Brühl, C. A. (2016). The secret pollinators: An overview of moth pollination

with a focus on Europe and North America. Arthropod-Plant Interactions, 10(1), 21–

28. https://doi.org/10.1007/s11829-016-9414-3

Hobern, D. (2021, October 16). Autonomous Moth Trap Project – Stangeia.

https://stangeia.hobern.net/autonomous-moth-trap-project/

Høye, T. T., Ärje, J., Bjerge, K., Hansen, O. L. P., Iosifidis, A., Leese, F., Mann, H. M. R.,

Meissner, K., Melvad, C., & Raitoharju, J. (2021). Deep learning and computer vision

will transform entomology. Proceedings of the National Academy of Sciences,

118(2), e2002545117. https://doi.org/10.1073/pnas.2002545117

Huebner, S., Palmer, M., & Packer, C. (2024). Human Supervision is Key to Achieving

Accurate AI-assisted Wildlife Identifications in Camera Trap Images. Citizen Science:

Theory and Practice, 9. https://doi.org/10.5334/cstp.752

Huijbers, C., Hogeweg, L., Jongejans, E., & Peterse, S. (2025). INSECTEN automatisch

vastgelegd. In Statusrapport Nederlandse biodiversiteit 2025 (pp. 102–104).

Naturalis Biodiversity Center. https://doi.org/10.5281/ZENODO.15350844

Kays, R., Arbogast, B. S., Baker-Whatton, M., Beirne, C., Boone, H. M., Bowler, M., Burneo, S.

F., Cove, M. V., Ding, P., Espinosa, S., Gonçalves, A. L. S., Hansen, C. P., Jansen, P. A.,

Kolowski, J. M., Knowles, T. W., Lima, M. G. M., Millspaugh, J., McShea, W. J., Pacifici,

K., … Spironello, W. R. (2020). An empirical evaluation of camera trap study design:

How many, how long and when? Methods in Ecology and Evolution, 11(6), 700–713.

https://doi.org/10.1111/2041-210X.13370

Korsch, D., Bodesheim, P., Brehm, G., & Denzler, J. (2023). Automated Visual Monitoring of

Nocturnal Insects with Light-based Camera Traps (arXiv:2307.15433). arXiv.

https://doi.org/10.48550/arXiv.2307.15433

Leorna, S., & Brinkman, T. (2022). Human vs. machine: Detecting wildlife in camera trap

images. Ecological Informatics, 72, 101876.

https://doi.org/10.1016/j.ecoinf.2022.101876

Lord, W., Teagle, S., Alton, J., Bannister, G., Beuchert, J., Bjerge, K., Carbone, D., Segura, A. G.,

Howson, T., Høye, T. T., Lawson, J., Ravivarma, A., Roy, D. B., Rylett, D., Skinner, G.,

Warwick, A., & August, T. (2026). Automated Monitoring of Insects system: Building

a non-lethal and scalable solution for monitoring of nocturnal insects (p.

2026.02.06.704388). bioRxiv. https://doi.org/10.64898/2026.02.06.704388

Möglich, J. M., Lampe, P., Fickus, M., Younis, S., Gottwald, J., Nauss, T., Brandl, R., Brändle, M.,

Friess, N., Freisleben, B., & Heidrich, L. (2023). Towards reliable estimates of

abundance trends using automated non-lethal moth traps. Insect Conservation and

Diversity, 16(5), 539–549. https://doi.org/10.1111/icad.12662

Montgomery, G. A., Belitz, M. W., Guralnick, R. P., & Tingley, M. W. (2021). Standards and

Best Practices for Monitoring and Benchmarking Insects. Frontiers in Ecology and

Evolution, 8. https://doi.org/10.3389/fevo.2020.579193

Mothbox Github. (2024). Mothbox. https://digital-naturalism-

laboratories.github.io/Mothbox/

Redmon, J., Divvala, S., Girshick, R., & Farhadi, A. (2016). You Only Look Once: Unified, Real-

Time Object Detection. Conference on Computer Vision and Pattern Recognition.

https://doi.org/10.48550/arXiv.1506.02640

Rhodes, M. W., Bennie, J. J., Spalding, A., ffrench-Constant, R. H., & Maclean, I. M. D. (2022).

Recent advances in the remote sensing of insects. Biological Reviews, 97(1), 343–

360. https://doi.org/10.1111/brv.12802

Rydell, J., Entwistle, A., & Racey, P. A. (1996). Timing of Foraging Flights of Three Species of

Bats in Relation to Insect Activity and Predation Risk. Oikos, 76(2), 243–252.

https://doi.org/10.2307/3546196

Schmeller, D. S., Böhm, M., Arvanitidis, C., Barber-Meyer, S., Brummitt, N., Chandler, M.,

Chatzinikolaou, E., Costello, M. J., Ding, H., García-Moreno, J., Gill, M., Haase, P., Jones,

M., Juillard, R., Magnusson, W. E., Martin, C. S., McGeoch, M., Mihoub, J.-B., Pettorelli,

N., … Belnap, J. (2017). Building capacity in biodiversity monitoring at the global

scale. Biodiversity and Conservation, 26(12), 2765–2790.

https://doi.org/10.1007/s10531-017-1388-7

Sickel, W., Zizka, V., Scherges, A., Bourlat, S. J., & Dieker, P. (2023). Abundance estimation

with DNA metabarcoding – recent advancements for terrestrial arthropods.

Metabarcoding and Metagenomics, 7, e112290.

https://doi.org/10.3897/mbmg.7.112290

Sittinger, M., Uhler, J., Pink, M., & Herz, A. (2024). Insect detect: An open-source DIY camera

trap for automated insect monitoring. PLOS ONE, 19(4), e0295474.

https://doi.org/10.1371/journal.pone.0295474

Stevens, S., Wu, J., Thompson, M. J., Campolongo, E. G., Song, C. H., Carlyn, D. E., Dong, L.,

Dahdul, W. M., Stewart, C., Berger-Wolf, T., Chao, W.-L., & Su, Y. (2024). BioCLIP: A

Vision Foundation Model for the Tree of Life. 19412–19424.

https://openaccess.thecvf.com/content/CVPR2024/html/Stevens_BioCLIP_A_Visio

n_Foundation_Model_for_the_Tree_of_Life_CVPR_2024_paper.html

Tyler, T. (2020). Relationship between moth (night active Lepidoptera) diversity and

vegetation characteristics in southern Sweden. Journal of Insect Conservation,

24(6), 1005–1015. https://doi.org/10.1007/s10841-020-00270-y

UKCEH AMI system | UK Centre for Ecology & Hydrology. (n.d.). Retrieved September 17,

2024, from https://www.ceh.ac.uk/solutions/equipment/automated-monitoring-

insects-trap

Vélez, J., McShea, W., Shamon, H., Castiblanco-Camacho, P. J., Tabak, M. A., Chalmers, C.,

Fergus, P., & Fieberg, J. (2023). An evaluation of platforms for processing camera-

trap data using artificial intelligence. Methods in Ecology and Evolution, 14(2), 459–

477. https://doi.org/10.1111/2041-210X.14044

Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., Robertson, T., &

Vieglais, D. (2012). Darwin Core: An Evolving Community-Developed Biodiversity

Data Standard. PLOS ONE, 7(1), e29715.

https://doi.org/10.1371/journal.pone.0029715

Wong, M. K. L., & Didham, R. K. (2024). Global meta-analysis reveals overall higher

nocturnal than diurnal activity in insect communities. Nature Communications,

15(1), 3236. https://doi.org/10.1038/s41467-024-47645-2

Zou, Y., Van der Werf, W., Liu, Y., & Axmacher, J. (2020). Predictability of species diversity

by family diversity across global terrestrial animal taxa. Global Ecology and

Biogeography. https://doi.org/10.1111/geb.13043