**The Gibbs free energy landscape based on liver metabolome revealed thermodynamic**

**robustness against fasting and obesity**

Takumi Abekawa[1], Satoshi Ohno[1,2,3,*], Akiyoshi Hirayama[4], Tomoyoshi Soga[4,5],

Shinya Kuroda[1,2,*]

[1]Department of Biological Sciences, Graduate School of Science, The University of Tokyo,

7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[2]Molecular Genetics Research Laboratory, Graduate School of Science, The University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, Japan.

[3]Department of AI Systems Medicine, M&D Data Science Center, Institute of Integrated

Research, Institute of Science Tokyo, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.

[4]Institute for Advanced Biosciences, Keio University, 246-2 Mizukami, Kakuganji,

Tsuruoka, Yamagata 997-0052, Japan

[5]Human Biology-Microbiome-Quantum Research Center (WPI-Bio2Q), Keio University,

Tsuruoka, Yamagata 997-0052, Japan

*Correspondence to: s-ohno.dsc@tmd.ac.jp (S.O.)

*Correspondence to: skuroda@bs.s.u-tokyo.ac.jp (S.K.)

**Abstract**

Mammalian liver metabolism undergoes a substantial shift during fasting. Thermodynamic principles impose fundamental constraints on metabolism, and the Gibbs free energy change of reaction ($\Delta_r G'$) indicates the reaction's direction and distance from equilibrium. However, $\Delta_r G'$ landscapes in intact mammalian organs remain largely uncharacterized. Here, we mapped $\Delta_r G'$ profile of glucose metabolism in mouse liver during fasting, using experimentally measured absolute metabolite concentrations and a newly developed computational method, GLEAM. We found that despite large metabolite fluctuations during fasting, the corresponding $\Delta_r G'$s remained robust, even for reactions reversing the direction between glycolysis and gluconeogenesis. A remarkable thermodynamic robustness is found in maintaining potential candidates of rate-limiting steps during fasting. This robustness is achieved by expending substantial costs for enzyme expressions, contributing to efficient switching from glycolysis to gluconeogenesis. Furthermore, obese mouse liver also showed thermodynamic robustness despite obesity-induced metabolic disruption. Our framework provided a novel thermodynamic perspective on intact organ metabolism, demonstrating that the liver robustly maintains thermodynamic characteristics favorable for metabolic control by buffering metabolite concentration differences.

37 **Glossary**

| Metabolite abbreviations | Metabolite names |
| --- | --- |
| G6P | Glucose 6-phosphate |
| F6P | Fructose 6-phosphate |
| F1,6BP | Fructose 1,6-bisphosphate |
| ADP | ADP |
| ATP | ATP |
| DHAP | Glycerone phosphate |
| GAP | Glyceraldehyde 3-phosphate |
| NAD | $NAD^+$ |
| NADH | NADH |
| Pi | Orthophosphate |
| 3PG | 3-Phospho-D-glycerate |
| 2PG | 2-Phospho-D-glycerate |
| PEP | Phosphoenolpyruvate |
| H2O | H2O |
| G1P | Glucose 1-phosphate |
| G3P | sn-Glycerol 3-phosphate |
| Pyr | Pyruvate |
| CoA | Coenzyme A |
| CO2 | CO2 |
| AcCoA | Acetyl-CoA |

| | |
|---|---|
| OAA | Oxaloacetate |
| Cit | Citrate |
| AKG | alpha-Ketoglutaric acid |
| SucCoA | Sucinyl-CoA |
| Suc | Succinate |
| GDP | GDP |
| GTP | GTP |
| Fum | Fumarate |
| FAD | FAD |
| FADH2 | FADH2 |
| Mal | Malate |
| Glu | Glutamate |
| NH3 | Ammonia |
| Ala | L-Alanine |
| Lac | Lactate |
| HCO3 | $HCO_3^-$ |
| NADP | $NADP^+$ |
| NADPH | NADPH |

| Reaction abbreviations | Reaction names |
|---|---|
| GLUT | Glucose transporter |
| GCK | Glucokinase |

| HK | Hexokinase |
|---|---|
| GPI | Glucose-6-phosphate isomerase |
| PFK | Phosphofructokinase |
| ALDO | Aldolase |
| TPI | Triosephosphate isomerase |
| GAPDH:PGK | Glyceraldehyde-3-phosphate dehydrogenase and Phosphoglycerate kinase |
| PGAM | Phosphoglycerate mutase |
| ENO | Enolase |
| PGM | Phosphoglucomutase |
| FBP | Fructose-1,6-bisphosphatase |
| G6PC | Glucose-6-phosphatase |
| GPD | Glycerol-3-phosphate dehydrogenase |
| PKLR | Pyruvate kinase |
| PDH | Pyruvate dehydrogenase |
| CS | Citrate synthase |
| ACO:IDH | Aconitase and Isocitrate dehydrogenase |
| OGDH | 2-Oxoglutarate dehydrogenase |
| LSC | Succinate-CoA ligase |
| SDH | Succinate dehydrogenase |
| FH | Fumarate hydratase |
| MDH | Malate dehydrogenase |
| GDH | Glutamate dehydrogenase |

| ALT | Alanine aminotransferase |
| LDH | Lactate dehydrogenase |
| MCT | Monocarboxylate transporter |
| PC | Pyruvate carboxylase |
| PCK | Phosphoenolpyruvate carboxykinase |
| ME | Malic enzyme |

**Introduction**

Blood glucose homeostasis is crucial in mammals for providing stable energy sources to

organs and maintaining health. During fasting, the liver plays a central role in maintaining

blood glucose homeostasis by shifting its metabolism from glycolysis to gluconeogenesis (*1–*

*5*). This metabolic shift is a highly regulated process that involves the directional reversal of

metabolic reactions and fluctuations in the concentrations of numerous metabolites (*6–8*).

Obesity induces metabolic dysregulation in the liver (*7–12*), leading to impaired glucose

homeostasis (*13–17*) and the development of further diseases, including type 2 diabetes (*9,*

*18–20*).

Metabolic processes are governed by the laws of thermodynamics. According to the second

law of thermodynamics, in the cellular environment, metabolic reactions proceed in the

direction where the Gibbs free energy change of reaction ($\Delta_r G'$) is negative, and $\Delta_r G'$ is

zero when the reaction reaches equilibrium. Therefore, $\Delta_r G'$ indicates the reaction distance

from equilibrium. $\Delta_r G'$ can be calculated as the difference between the Gibbs free energy

of formation ($\Delta_f G'$) of the product and substrate metabolites:

$$\Delta_r G' = \sum_i S_i \Delta_f G'_i \tag{1}$$

55  where $S_i$ is the stoichiometric coefficient of metabolite *i* in the reaction. $\Delta_f G'$ depends on

56  the metabolite activity, typically substituted by the concentration:

$$\Delta_f G' = \Delta_f G'^\circ + RT \ln c \tag{2}$$

where $\Delta_f G'^\circ$ is standard Gibbs free energy of formation representing $\Delta_f G'$ at 1M of

metabolite concentration, $R$ is the gas constant, $T$ is temperature, and $c$ is the metabolite

concentration. Although calculation of $\Delta_r G'$ requires absolute metabolite concentrations,

mass spectrometry-based metabolomics analyses often suffer from ion suppression, a

phenomenon where co-eluting compounds compromise quantitative accuracy by reducing

analyte ionization efficiency (*21*). Nevertheless, previous studies have successfully achieved

absolute quantification of intracellular metabolites, primarily by employing isotope-labeled

internal standards (*22, 23*). These techniques have enabled the investigation of absolute

metabolite concentrations and subsequent thermodynamic analyses, such as $\Delta_r G'$

landscapes, in microorganisms and cultured cells. (*24–27*). Despite these advancements,

thermodynamic profiles have rarely been investigated at *in vivo* mammalian organ levels.

Furthermore, all prior thermodynamic studies have focused on glycolysis-dominant

conditions where glucose is the primary nutrient, leaving a large gap in our understanding of

metabolism during fasting, a condition dominated by gluconeogenesis.

Mathematical metabolic modeling is an essential framework for a quantitative system-level

understanding of metabolism. The thermodynamic principle regarding reaction direction has

been widely applied as a constraint to increase the reliability of metabolic models (*28–35*).

Furthermore, the significance of $\Delta_r G'$ extends beyond physical aspects such as distance

from equilibrium and reaction directionality to several physiological meanings, including

pathway rate-limiting steps quantified by the flux control coefficient (FCC) (*36–38*), and

enzyme demands for carrying metabolic flux (*39–42*). However, these thermodynamic

frameworks have hardly been applied to experimental data, especially those from

multicellular organisms.

In this study, we explored $\Delta_r G'$ landscape in the liver metabolism of mice at different

fasting levels using time-series absolute quantitative metabolomic data measured by CE-MS,

which can reduce the effect of ion suppression (*43*, *44*), without using isotopically labeled

standards. Our results are the first *in vivo* identification of large-scale $\Delta_r G'$ landscapes in

mammalian organisms, and also the first elucidation of $\Delta_r G'$ behavior during fasting in any

cell types. Since metabolomic data contains uncertainties, we developed GLEAM to estimate

$\Delta_r G'$ with thermodynamically consistent metabolite concentrations from experimental

metabolomic data. The estimated $\Delta_r G'$ showed that the global $\Delta_r G'$ landscape in mouse

liver is robust during fasting, despite large fluctuations in metabolite concentrations. We

further integrated the $\Delta_r G'$ and enzymatic kinetics to investigate the physiological

significance of the $\Delta_r G'$ landscape in terms of rate-limiting steps and efficiency of

metabolic switching. In addition to the thermodynamic robustness of mouse liver against

fasting, we also demonstrated the robustness against obesity using leptin-deficient obese

(*ob*/*ob*) mice (*45*, *46*). Thus, our study offers novel insights into glucose metabolism in *in*

*vivo* mammalian liver from a thermodynamic standpoint. These findings extend our

understanding of mammalian metabolisms beyond individual metabolite concentrations.

**Results**

**$\Delta_r G'$ landscape is robust against fasting despite metabolite concentration fluctuations**

In this study, we developed GLEAM to estimate $\Delta_r G'$ from time-series metabolome and

$\Delta_f G'^\circ$. One of the challenges in estimating $\Delta_r G'$ in an entire reaction pathway is inherent

uncertainties and missing values in metabolome and $\Delta_f G'^\circ$ data. To address this issue,

GLEAM utilizes thermodynamic constraints and variances and covariances of metabolome

and $\Delta_f G'^\circ$ data, and estimates a complete set of thermodynamically consistent metabolite

concentrations and $\Delta_f G'^\circ$, and subsequent $\Delta_r G'$ (see Methods). Briefly, GLEAM fits

metabolite concentrations and $\Delta_f G'^\circ$ to data from the experiment or database within a range

that every reaction holds $\Delta_r G' < 0$ by the weighted least mean squares method. The weights

of the residuals are determined by variances and covariances of metabolite concentrations

and $\Delta_f G'^\circ$ data.

By applying GLEAM to time-series metabolomic data from *ad libitum* feeding to fasting in

mice, and $\Delta_f G'^\circ$ data from eQuilibrator (data file S1), we estimated the $\Delta_r G'$ landscape

(Fig. 1A, B, data file S2) in glycolysis, gluconeogenesis, and TCA cycle (data file S3) in the

liver. At *ad libitum* feeding (0-hour fasting), when glycolysis is dominant, most of the

irreversible reactions were markedly far from equilibrium, such as $\Delta_r G' = -21.56$ kJ/mol in

PFK and $\Delta_r G' = -34.34$ kJ/mol in PDH. Among the 11 irreversible reactions, only three

reactions (ACO:IDH, PC, and PCK) exhibited $\Delta_r G' > -10$ kJ/mol, and PC and PCK were

exceptionally near equilibrium, with $\Delta_r G' = -3.48$ kJ/mol and $-3.70$ kJ/mol, respectively.

In contrast, most of the reversible reactions exhibited $\Delta_r G'$ close to zero at *ad libitum*

feeding, such as $\Delta_r G' = -0.24$ kJ/mol in GPI and $\Delta_r G' = -0.03$ kJ/mol in SDH, suggesting

that these reactions were near equilibrium. Four reversible reactions (ALDO, LSC, MDH,

and ME) were exceptions and exhibited $\Delta_r G' < -5$ kJ/mol. MDH was largely far from

equilibrium in the opposite direction of the oxidative TCA cycle, with $\Delta_r G' = -18.13$ kJ/mol.

The fasting-induced metabolic shifts in the liver from glycolysis to gluconeogenesis lead to

changes in thermodynamic characteristics of metabolic reactions, including reversal of the

sign of $\Delta_r G'$ of some reactions *i.e.* reversal of their direction. After 16-hour fasting, when

gluconeogenesis is dominant in the liver, all irreversible reactions exhibited $\Delta_r G' < -5$

kJ/mol, indicating that the overall trend of irreversible reactions being far from equilibrium

was maintained across *ad libitum* feeding and after fasting (Fig. 1C) despite large differences

in metabolite concentrations (Fig. 1D). Although PC and PCK had been near equilibrium at

*ad libitum* feeding, these reactions moved away from equilibrium to exhibit $\Delta_r G' = -9.07$

kJ/mol and $-5.25$ kJ/mol after fasting, respectively. As for reversible reactions, the directions

of some reactions are reversed to switch the liver's metabolism from glycolysis to

gluconeogenesis. However, all the reversible reactions with $\Delta_r G' > -5$ kJ/mol at *ad libitum*

feeding remained $\Delta_r G' > -5$ kJ/mol, indicating that the overall trend of reversible reactions

being close to equilibrium was maintained across *ad libitum* feeding and after fasting.

To examine whether the thermodynamic characteristics of metabolic reactions observed in

mouse liver are conserved across species, we compared the estimated $\Delta_r G'$ in glycolytic

reactions in mouse liver with those in other biological systems. The previous study (*26*)

determined $\Delta_r G'$ in glycolysis of immortalized baby mouse kidney epithelial (iBMK) cells,

yeast, and *Escherichia coli*. Similar to the mouse liver, all of these cell types showed

reversible reactions close to equilibrium, and the irreversible reaction PFK far from

equilibrium (Fig. 1E). This result suggests that the thermodynamic characteristics in

glycolysis are widely conserved across species. An exception is PKLR, whose $\Delta_r G'$ in

mouse liver was more than 20 kJ/mol smaller than those in the other cell types. Among the

metabolite concentrations (or ratios) involved in PKLR, pyruvate concentration and

ATP/ADP were smaller in mouse liver than in mammalian iBMK cells, yeast, and *E. coli*

(Table S2). In particular, pyruvate concentrations were two orders of magnitude smaller in

mouse liver ($3.77 \times 10^{-2}$ mM) compared to the other three cell types (5.88 mM in iBMK cells,

9.40 mM in yeast, and 3.66 mM in *E. coli*).

**$\Delta_r G'$ maintenance during fasting is associated with metabolite concentration**

**fluctuations in different ways among pathways**

To characterize thermodynamic changes in each metabolic reaction accompanying the

fasting-induced metabolic shift, we calculated the $\Delta_r G'_{16h} - \Delta_r G'_{0h}$ during fasting, defined

as $\Delta_r G'$ variations (Fig. 2A). Among the seven irreversible reactions of glycolysis and

gluconeogenesis, four reactions (GCK, PFK, PKLR, and PC) exhibited an absolute $\Delta_r G'$

variation during fasting greater than 5 kJ/mol. In contrast, only one reversible reaction

(ALDO) out of 11 showed an absolute $\Delta_r G'$ variation during fasting exceeding 5 kJ/mol,

despite the reversal of reversible reaction directions between glycolysis and gluconeogenesis

under these conditions. This result suggests that reversible reactions maintained $\Delta_r G'$ more

strictly than irreversible reactions even though the sign of $\Delta_r G'$ is reversed. In the TCA

cycle, the absolute $\Delta_r G'$ variations during fasting were less than 3 kJ/mol for all reactions,

indicating that the TCA cycle is thermodynamically stable in response to fasting.

As $\Delta_r G'$ variations during fasting differed between irreversible and reversible reactions in

glycolysis and gluconeogenesis, we further examined the relationship between the distance

from equilibrium ($|\Delta_r G'|$) at *ad libitum* feeding and the absolute $\Delta_r G'$ variation (Fig. 2B).

In glycolysis and gluconeogenesis, the absolute $\Delta_r G'$ variations during fasting were less

than 3.5 kJ/mol for near-equilibrium reactions with $|\Delta_r G'| < 5$ kJ/mol at *ad libitum* feeding

except for PC. PC moved away from equilibrium only when gluconeogenesis is dominant in

response to fasting. In contrast, among the reactions with $|\Delta_r G'| \geq 5$ kJ/mol at *ad libitum*

feeding, the absolute $\Delta_r G'$ variations ranged approximately between 0 and 10 kJ/mol. This

result suggests that $\Delta_r G'$ of near-equilibrium reactions in glycolysis and gluconeogenesis

tended to be significantly strictly maintained, while this tendency was lost among reactions

displaced from equilibrium (p value = 0.037 < 0.05 calculated by two-tailed Welch's *t*-test, n

= 12 for near-equilibrium reactions with $|\Delta_r G'| < 5$ and n = 6 for the other reactions with

$|\Delta_r G'| \geq 5$ kJ/mol). In the TCA cycle, $\Delta_r G'$ was strictly maintained regardless of the distance

from equilibrium. The statistical test was not performed for the TCA cycle due to a lack of

near-equilibrium reactions.

To investigate the reason maintaining $\Delta_r G'$ in near-equilibrium reactions in glycolysis and

gluconeogenesis, as well as in the TCA cycle reactions, we compared the absolute $\Delta_r G'$

variation with the total absolute $\Delta_f G'$ variation of metabolites involved in the reaction (Fig.

2C). The total absolute $\Delta_f G'$ variation of metabolites is defined as $\sum \left| S\left(\Delta_f G'_{16h} - \Delta_f G'_{0h}\right)\right|$,

where $S$ is the stoichiometric coefficient of a metabolite involved in the reaction. The $\Delta_r G'$

variation is associated in various ways with fluctuations in the $\Delta_f G'$ of substrates and

products, which depend on the metabolite concentrations. For example, when substrate and

product concentrations remain relatively constant and changes in their $\Delta_f G'$ remain small,

the $\Delta_r G'$ variation is small (*e.g.*, GPD in Fig. 2D). Even when substrate and product

concentrations largely fluctuate and changes in their $\Delta_f G'$ are large, the $\Delta_r G'$ variation can

remain small when the changes in $\Delta_f G'$ of substrate and products are cancelled out. This

can happen, for instance, when both the substrate and product $\Delta_f G'$ values increase

simultaneously (*e.g.*, GAPDH:PGK in Fig. 2D). In contrast, when the changes in $\Delta_f G'$ of

substrate and products are not cancelled out, the $\Delta_r G'$ variation becomes large (*e.g.*, GCK

in Fig. 2D). In glycolysis and gluconeogenesis, reactions with small absolute $\Delta_r G'$ variation

due to small fluctuation in metabolite concentrations, such as GPD, and due to the

cancellation of $\Delta_f G'$ variations such as GAPDH:PGK, were both observed (Fig. 2C). The

existence of the latter type of reaction suggests that while metabolite concentrations fluctuate

in response to fasting in glycolysis and gluconeogenesis, the fluctuations in metabolites

involved in near-equilibrium reactions were coordinated, thereby maintaining $\Delta_r G'$ of these

reactions. In the TCA cycle, the reason for maintaining the absolute $\Delta_r G'$ variations were

small fluctuations in metabolite concentrations for all the reactions.

**Thermodynamic Rate-limiting candidates (TRCs) were maintained during fasting**

One of the physiological consequences of reaction distances from equilibrium is the control of steady-state pathway fluxes. The extent to which perturbation of each reaction can control pathway steady-state flux is quantified by the flux control coefficient (FCC) in metabolic control analysis (MCA) (*36, 47*). In this study, we analytically calculated FCCs under both arbitrary $\Delta_r G'$ landscapes and those observed in mouse liver within glycolysis (Fig. 3A), gluconeogenesis (Fig. 3B), and the TCA cycle (Fig. 3C) from an enzymatic rate law using the MCA's theorems. To incorporate reaction distance from equilibrium, we used reversible kinetics called the direct binding modular rate law (*48*) as the enzymatic rate law. Under this framework, a flux $v$ can be decomposed into the effect of non-equilibrium $\eta_{\text{rev}}$ (Fig. 3D) and substrate saturation $\eta_{\text{sat,s}}$ (*49*):

$$v = E k_{\text{cat}}^{+} \eta_{\text{rev}} \eta_{\text{sat,s}}, \tag{3}$$

$$\eta_{\text{rev}} = 1 - \exp\left(\frac{\Delta_r G'}{RT}\right), \tag{4}$$

$$\eta_{\text{sat,s}} = \frac{\prod_{i \in M^s}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}}{1 + \prod_{i \in M^s}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|} + \prod_{i \in M^p}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}} \tag{5}$$

where $E$ is an enzyme concentration, $k_{\text{cat}}^{+}$ is the forward catalytic constant, $K_{\text{m}}$ is the Michaelis-Menten constant, $M^s$ is substrates, and $M^p$ is products. Using this reaction rate raw, we calculated FCC based on the connectivity theorem and summation theorem in MCA (*50*) (see Methods). This FCC calculation depends on metabolite concentrations, $\Delta_f G'^{\circ}$, and $K_{\text{m}}$. For $\Delta_f G'^{\circ}$, we used sampling values for the confidence interval calculation in GLEAM.

$K_m$ values were sampled from distributions based on BRENDA database data (*51*).

Metabolite concentrations were obtained by the following two approaches.

We investigated FCC for metabolite concentrations in mouse liver using sampled values for

confidence interval calculations in GLEAM, and FCC for arbitrary $\Delta_r G'$ landscapes to

identify thermodynamic conditions for large FCC. FCC for arbitrary $\Delta_r G'$ landscapes

showed that moderate or large FCC required $\eta_{rev} \approx 1$ in reactions except for ones located

upstream in a pathway (Fig. S1 A to D). In glycolysis, reactions with $\eta_{rev} \approx 1$ at *ad libitum*

feeding included irreversible reactions GCK, PFK, PKLR, and the reversible reaction ALDO

in mouse liver. Among these, reactions that can achieve FCC ≈ 1 were GCK and PFK, while

others can achieve at most moderate FCC values (Fig. 3E). We also found that GPI, which is

located upstream in the pathway, can achieve moderate FCC at moderate $\eta_{rev}$. In

gluconeogenesis, reactions with $\eta_{rev} \approx 1$ after 16-hour fasting were the irreversible

reactions PC, PCK, FBP, and G6PC, all of which can achieve FCC ≈ 1 (Fig. 3F). All the other

reactions exhibited only small FCC values. In the TCA cycle, similar FCC distributions were

observed at *ad libitum* feeding and after fasting (Fig. 3G, S3). Reactions with $\eta_{rev} \approx 1$ were

irreversible reactions CS, ACO:IDH, OGDH, and a reversible reaction LSC. Among these,

CS and OGDH can achieve FCC ≈ 1, while ACO:IDH and LSC can achieve at most moderate

FCC values. Generally, reversible reactions are considered to exhibit only small FCC values

and thus are not recognized as rate-limiting reactions. However, our results demonstrated that

the reversible reactions GPI, ALDO, and LSC can achieve moderate FCC values comparable

to several irreversible reactions. Considering the results of arbitrary $\Delta_r G'$ landscapes, these

FCC values of the reversible reactions can be attributed to GPI being located relatively

upstream in glycolysis, and ALDO and LSC being sufficiently displaced from equilibrium to

achieve $\eta_{\text{rev}} \approx 1$ despite being closer to equilibrium than irreversible reactions.

As described above, a reaction with a large FCC seems to be either far from equilibrium ($\eta_{\text{rev}}$

$\approx 1$), or located upstream in a pathway. Because the location of a reaction in a pathway

depends on the definition of the pathway, a reaction being far from equilibrium is a key

determinant for the reaction to have a large FCC, sometimes being referred to as the rate-

limiting step. Therefore, we introduced Thermodynamic Rate-limiting Candidates (TRC) as

reactions with $\eta_{\text{rev}} \geq 0.9$, and examined which reaction is thermodynamically capable of

being a rate-limiting step for any pathway definitions at *ad libitum* feeding (Fig. 3H, 0h) and

after fasting (Fig. 3H, 16h). TRC reactions both at *ad libitum* feeding and after fasting

included GCK, PFK, CS, and seven other reactions, indicating these reactions can be a rate-

limiting step regardless of fasting levels. The parameter $\eta_{\text{rev}}$ and TRC status (either TRC

or not) are sensitive to $\Delta_r G'$ variations at a near equilibrium because $\eta_{\text{rev}}$ saturates around

$\Delta_r G' = -10$ kJ/mol (Fig. 3D). Due to this property, 15 out of 18 reactions remained either

TRC (red line in Fig. 3H) or non-TRC (blue line in Fig. 3H) across at *ad libitum* feeding and

after fasting in glycolysis and gluconeogenesis, where only $\Delta_r G'$ of near-equilibrium

reactions were strictly maintained. This result suggests that fasting-induced fluctuations in

metabolite concentrations occur within a range that maintains the TRC status for most

reactions. The three reactions of ALDO, PC, and PCK were the exceptions that did not remain

TRC or non-TRC (gray line in Fig. 3H), and whether these reactions are rate-limiting or not

would depend on fasting levels. TRC status described above remains stable across the

threshold from $\eta_{\text{rev}} = 0.88$ to $0.95$.

**271**  **Glycolysis and Gluconeogenesis Achieve Efficient Directional Switching by Expending**

**272**  **Large Enzyme Costs**

A fundamental question in biology is to understand what biological systems are optimized

for. While objective functions optimized in microbial metabolism—such as maximizing

growth or ATP production (*52*) and minimizing enzyme demands (*42*)—have been reported,

the objective function for *in vivo* mammalian metabolism remains completely unknown.

Therefore, we examined the optimality of metabolism in mouse liver from the aspect of

metabolite concentrations and $\Delta_r G'$ landscape obtained from GLEAM.

By solving the equation of the direct binding modular rate law (1) with respect to the enzyme

amount, we obtain

$$E \ = \frac{v}{k_{cat}^+} \eta_{rev}^{-1} \, \eta_{sat,s}^{-1}. \tag{6}$$

This equation shows that the nearer a reaction is to equilibrium (the smaller $\eta_{rev}$ is), the

greater the enzyme amount required to achieve the same flux $v$. Here we defined Enzyme

Cost (EC) as the product of enzyme amount and enzyme molecular weight, and Enzyme Cost

Minimum (ECM) as the theoretical state that minimizes the total EC within the entire

pathway (see Method). By comparing ECs and metabolite concentrations in mouse liver to

those in ECM, we investigated whether mouse liver metabolism is optimized for enzyme

demands reduction. As with FCC calculation, we calculated EC using sampling values of

metabolite concentrations and $\Delta_f G'^\circ$ used in the confidence interval calculation in GLEAM.

For $K_m$ values, we obtained them as geometric means of data in the BRENDA database (*51*).

We calculated EC and ECM for the reactions in glycolysis at *ad libitum* feeding,

gluconeogenesis after 16-hour fasting, and the TCA cycle under both these conditions (data

file S4). The liver showed larger total EC compared to ECM, with a difference between

pathways. The liver/ECM ratio of total EC was on the order of $10^4$ for glycolysis and

gluconeogenesis, but only $10^1$ for the TCA cycle (Fig. 4A). Additionally, we compared ECM-

predicted metabolite concentrations with liver metabolite concentrations in each pathway,

and found typical fold errors of $10^{\mathrm{RMSE}} = 29.79$ (RMSE: root mean square error on a $\log_{10}$-

scale) for glycolysis and gluconeogenesis and 14.40 for the TCA cycle (Fig. 4B). These

findings show that enzyme and metabolite concentrations in the liver are not optimized for

EC, and this is more evident in glycolysis and gluconeogenesis than in the TCA cycle.

To identify a key reaction contributing to the large total EC of glycolysis and gluconeogenesis

in mouse liver, we broke down the total EC of pathways into ECs of individual reactions

(data file S4). The GAPDH:PGK reaction in mouse liver exhibited the highest EC of $10^6$ both

in glycolysis and gluconeogenesis (Fig. 4C), while EC of reactions in the TCA cycle (*e.g.*,

SDH, ACO:IDH, and FH) are at most on the order of $10^2$. The EC values of GAPDH:PGK

are 99% and 89% of the total EC of glycolysis and gluconeogenesis, respectively, suggesting

that GAPDH:PGK outstandingly elevates the total EC of glycolysis and gluconeogenesis.

The high EC for GAPDH:PGK both in glycolysis and gluconeogenesis is attributed to two

factors. One factor is low substrate saturation (Fig. 4C, blue), which is given by the lower

substrate metabolite concentrations ($3.4 \times 10^{-3}$ mM of GAP *ad libitum* feeding and $1.3 \times 10^{-2}$

mM of 3PG after fasting) than their respective $K_{\mathrm{m}}$ values ($1.9 \times 10^{-1}$ mM for GAP and

$3.5 \times 10^{-1}$ mM for 3PG), resulting in large $\eta_{\mathrm{sat,s}}^{-1}$. From equation (4), a large $\eta_{\mathrm{sat,s}}^{-1}$, indicating

low substrate saturation, leads to an increase in EC to carry a certain flux. Under this

condition, an increase in flux from higher enzyme levels is mitigated by fluctuations in

metabolite concentrations. The other factor of the high EC for GAPDH:PGK is liver

thermodynamics (Fig. 4C, red), where the reaction was close to equilibrium ($\Delta_r G' = -0.01$

kJ/mol both at *ad libitum* feeding and after fasting), resulting in large $\eta_{\mathrm{rev}}^{-1}$. When the reaction

is close to equilibrium, increasing the enzyme abundance enhances both the forward and

reverse fluxes, thus limiting the net flux increase. Thus, the elevated EC for this reaction is a

result of both low substrate saturation and high reaction reversibility.

The liver glycolysis and gluconeogenesis required much larger EC than ECM, especially in

the midstream reversible reactions due to large $\eta_{\mathrm{rev}}^{-1}$ values. We hypothesized that the reason

for this may be that the enzyme costs are required for enabling switching from glycolysis to

gluconeogenesis during fasting. To assess the metabolic hurdle for directional switching from

glycolysis to gluconeogenesis, we introduced the Metabolite Equilibrium Gap (MEG) as the

typical fold change in metabolite concentration required for the directional switching (data

file S5) (see Methods). In glycolysis and gluconeogenesis, mouse liver exhibited MEG =

1.35 while ECM exhibited MEG = 16.01 (Fig. 4D), suggesting that the liver achieves

directional switching from glycolysis to gluconeogenesis with much smaller metabolite

concentration changes than ECM. Unlike glycolysis and gluconeogenesis, the TCA cycle,

which does not undergo directional switching during fasting, exhibited a larger MEG (6.85)

in the liver than that of glycolysis and gluconeogenesis. The mouse liver expends a much

larger enzyme cost in glycolysis and gluconeogenesis than the theoretical minimum, allowing

it to achieve directional switching with smaller changes in metabolite concentration.

Conversely, a smaller enzyme cost is expended for the TCA cycle—which does not undergo

directional switching—than for glycolysis and gluconeogenesis.

**Robustness of $\Delta_r G'$ in The Liver Against Obesity**

Obesity disrupts liver metabolism, leading to impaired homeostasis in response to fasting (*7*,

*8*, *10*) and subsequent development of various diseases (*9*, *18–20*). Recent studies have

reported that the obesity-associated dysregulation of liver metabolism occurs even at *ad*

*libitum* feeding (*11*, *12*). To reveal the impact of obesity-induced metabolic disruption on the

thermodynamic characteristics of reactions, we calculated $\Delta_r G'$ in the *ob*/*ob* mouse liver

during fasting.

The $\Delta_r G'$ landscape in *ob*/*ob* mouse liver was similar to that in WT mouse liver both at ad

libitum feeding and after 16-hour fasting (Pearson's correlation coefficient $r = 0.99$ at ad

libitum feeding and 0.98 after 16-hour fasting) (Fig. 5A, B, data file S2). Consequently, the

overall trend of irreversible reactions being far from equilibrium and reversible reactions

being near equilibrium was preserved even under obesity. More than 80% of all 29 reactions

showed markedly small $\Delta_r G'$ differences (less than 3 kJ/mol) between WT and *ob*/*ob* mice

both at *ad libitum* feeding and after 16-hour fasting, with only three reactions (TPI after 16-

hour fasting, CS at the both timepoints) exceeding a 5 kJ/mol differences (Fig. 5C). The

similar $\Delta_r G'$ values between WT and *ob*/*ob* mouse indicate that the concentration ratios of

substrates and products for the reactions are comparable between the genotypes. Individual

metabolite concentrations (data file S2), however, showed discrepancies between WT and

*ob*/*ob* mice, with Pearson's $r = 0.015$ at ad libitum feeding and $-0.46$ after 16-hour fasting

(Fig. 5D). This result suggests that the thermodynamic characteristics of reactions remain

robust against obesity despite the changes in individual concentrations. Particularly, in most

reversible reactions of glycolysis and gluconeogenesis, the differences in $\Delta_f G'$ between WT

and *ob/ob* mouse were not reflected in the differences in $\Delta_r G'$ (Fig. 5E). This result

prominently demonstrates that the reaction thermodynamic characteristics are robust to

obesity while the individual metabolite concentrations are sensitive.

**Discussion**

We revealed the $\Delta_r G'$ landscape of glucose metabolism in *in vivo* mouse liver for the first time from fasting time-series absolute quantitative metabolome data using the newly developed GLEAM. In mouse liver, overall trend of $\Delta_r G'$ landscape was maintained regardless of fasting level (Fig. 1B, C). The $\Delta_r G'$ maintenance was markedly strict in near-equilibrium reactions in glycolysis and gluconeogenesis such as GAPDH:PGK (Fig. 2B), despite large fluctuations in metabolite concentrations (Fig. 1D, 2C), while this strictness was not true for reactions far from equilibrium such as GCK. In the TCA cycle, overall $\Delta_r G'$ variations were small due to minimal metabolite concentration fluctuations during fasting (Fig. 2B, C). This thermodynamic robustness against fasting led to maintenance of the reactions that could have large FCC (*i.e.*, rate-limiting step candidates) during fasting (Fig. 3H) because FCC is sensitive only in the near-equilibrium region. The thermodynamic landscape in the liver could also contribute to an efficient switching from glycolysis to gluconeogenesis by expending a larger enzyme expression than that theoretically minimized (Fig. 4A to D). We also found similar $\Delta_r G'$ landscapes between WT and *ob/ob* mice, while metabolite concentrations were quite different (Fig. 5A to E). Our results revealed that the liver robustly maintains $\Delta_r G'$ landscapes favorable for metabolic control in contrast to fluctuations in individual metabolite concentrations caused by fasting and obesity.

Methods for estimating thermodynamically consistent metabolite concentrations, $\Delta_f G'^\circ$, and $\Delta_r G'$ from absolute quantitative metabolome data have been proposed in previous studies (*26*, *27*). In these methods, metabolite concentrations and $\Delta_f G'^\circ$ were determined by

weighted least squares, where uncertainties of individual concentration or $\Delta_f G'^\circ$ were

independently accounted with the weights. However, metabolites sharing common chemical

group, such as ADP and ATP, show high correlation in their $\Delta_f G'^\circ$, because $\Delta_f G'^\circ$ depends

on structures of metabolites. GLEAM incorporates the covariance of $\Delta_f G'^\circ$ from

eQuilibrator (*53*) and enabled more precise $\Delta_r G'$ estimation compared to the previous

methods.

Our study narrowed down the reactions that could achieve large FCC in the pathways using

the obtained $\Delta_r G'$ landscape in mouse liver (Fig. 3E to G, S2). A previous study

experimentally measured FCC in glycolysis in iBMK cells, and reactions exhibiting large

FCCs are hexokinase (HK), which catalyzes the same reaction as GCK, and PFK (*54*).

Another previous study exhibited PC having dominant FCC in gluconeogenesis in rat liver

(*55, 56*). Although no studies have measured FCC in the TCA cycle to our knowledge, CS,

IDH, and OGDH have conventionally been regarded as rate-limiting steps. All these reactions

are included in the rate-limiting candidates in this study, suggesting that rate-limiting

reactions are maintained through cell types. However, the advantage of controlling flux via

these reactions remains unclear and is an important future topic for understanding system-

level metabolic regulation.

When calculating EC, we made two assumptions: having common turnover number, $k_{cat}$

values for all reactions and setting saturation ($c/K_m$) to the geometric mean of the pathway

when $K_m$ could not be obtained from BRENDA. These assumptions were required because

of the difficulty of obtaining kinetic parameters in mammals. The former assumption may

cause inaccurate weighting of individual enzymes in the total EC minimization, and the latter

may lead to inaccurate $\eta_{\mathrm{sat},s}^{-1}$. However, the metabolite concentrations close to ECM were

reported with a modeling where $k_{\mathrm{cat}}$ and $K_{\mathrm{m}}$ information was lacking and only the

influence of $\eta_{\mathrm{rev}}^{-1}$ was considered (*42*). This means that the $\Delta_r G'$ landscape in ECM would

be similar under any $k_{\mathrm{cat}}$ and $K_{\mathrm{m}}$ values. Therefore, our conclusion of mouse liver

metabolism not being optimized for enzyme cost reduction would be reasonable regardless

of $k_{\mathrm{cat}}$ and $K_{\mathrm{m}}$ assumptions.

According to equations (23), (24), (30) to (35), and (39), the design of $\Delta_r G'$ landscape faces

a trade-off between FCC concentration and enzyme cost minimization. If only a few reactions

in a metabolic pathway are far from equilibrium and exhibit $\eta_{\mathrm{rev}}$ close to 1, large FCCs are

likely to be concentrated in these reactions. In this case, $\eta_{\mathrm{rev}}$ of other reactions are close to

zero, leading to a large EC. This trade-off can be rephrased as a trade-off between flexibility

responding to multiple conditions and survival cost reduction under a single condition

because the pathway flux can be efficiently regulated through a minimal number of enzyme

activity changes when FCCs are concentrated in limited reactions. A similar trade-off also

occurs between MEG reduction and enzyme cost minimization. In mouse liver, particularly

in glycolysis and gluconeogenesis, reactions that could achieve high FCC were concentrated

in a few reactions (Fig. 3E to G, S2), and MEG was remarkably small (Fig. 4D), while there

were large deviations between mouse liver and ECM in total EC and metabolite

concentrations (Fig. 4A, B). These results indicate that metabolic regulation in mouse liver

adopts a strategy of flexibly responding to changes in environmental and physiological

conditions rather than optimizing for a certain condition. In contrast, the previous study in *E.*

*coli* showed that both enzyme concentrations and metabolite concentrations are well

predicted by ECM within a typical fold error range of $10^0$ (*42*) (Fig. S3). Taken together,

these facts suggest differences in metabolic regulation strategies between cell types; mouse

liver expends costs to perform flexible metabolic regulation, while *E. coli* optimized survival

cost in a certain condition. Furthermore, our study also showed that this difference in

metabolic regulation strategy may also exist between pathways by demonstrating the large

differences in TRC proportions and total EC optimality between glycolysis and

gluconeogenesis, and the TCA cycle. Note that the different total EC optimality among cell

types and pathways is consistent with their common tendency of reversible reactions being

near equilibrium (Fig. 1A, B) because $\eta_{\mathrm{rev}}^{-1}$ reaches $10^2$ at $\Delta_r G' = -5$ kJ/mol, meaning EC

takes greatly different values even in the range of near equilibrium.

One of the limitations of this study is that we assumed hepatocytes as a single compartment

and calculated $\Delta_r G'$ from metabolite concentrations averaged within the cell. As glycolysis

operates in the cytosol and the TCA cycle operates in mitochondria, more accurate $\Delta_r G'$

calculation requires concentrations in each of these compartments. Furthermore, localized

concentrations can be affected by substrate channeling (*57*). Substrate channeling is a

phenomenon where enzymes of consecutive reactions form complexes, allowing the product

of the former enzyme to be directly transferred to the active site of the latter enzyme without

being released into the solvent. For reactions involved in substrate channeling, $\Delta_r G'$ is

dependent on the channeled metabolite concentration in a limited space among the enzymes.

Although substrate channeling has not yet been confirmed in living organisms, the

channeling of oxaloacetate between MDH and CS has been indirectly supported by

experiments (*58*, *59*), and ALDO, TPI, and GAP have also been suggested to form complexes

(*60*, *61*). Another limitation is that the FCC calculations were based solely on stoichiometric

relationships and did not account for allosteric regulations because of the severe lack of $K_i$

in mammals and simplification of the calculation. Inclusion of allosteric regulations would

alter FCC values, allowing for further narrowing down the reactions that can have a large

FCC.

In conclusion, we developed GLEAM to estimate $\Delta_r G'$ in the glucose metabolism in mouse

liver and revealed the thermodynamic robustness against fasting and obesity. Furthermore,

we suggested that these thermodynamic characteristics in the liver contribute to efficient

metabolic control in two ways: the maintenance of rate-limiting step candidates and the

switching between glycolysis and gluconeogenesis with minimal metabolite fluctuations by

expending large enzyme expressions. This study thus provides crucial insights into the

principles underlying the highly flexible and robust metabolic regulation observed in

mammalian liver, paving the way for future investigations into therapeutic targets for

metabolic diseases.

**Materials and Methods**

**Metabolic network for glucose metabolism in mice**

A metabolic network for glucose metabolism in mice was constructed to calculate $\Delta_r G'$. The

network consists of total 55 reactions and 80 metabolites in the liver, muscle, and blood (the

same reactions and metabolites in different compartments were counted as distinct ones), and

involves glycolysis, gluconeogenesis, TCA cycle, glycogen metabolism, alanine metabolism,

and transporters (Fig. S4, data file S3). Among the 55 reactions, we defined 23 reversible

reactions and 32 irreversible reactions based on the tissue-specific genome-scale metabolic

network (*62*). All the transporters were defined as reversible reactions.

Reversible reactions GAPDH and PGK were combined into a single reversible reaction

GAPDH:PGK because 1,3-bisphosphoglycerate was not measured in the metabolomic data

(*8*). According to the experimentally measured metabolite concentrations, a reversible

reaction ACO proceeded in the reverse direction of the oxidative TCA cycle with a little

thermodynamic driving force at all the time points in WT and *ob*/*ob* mice, breaking a

sequential pathway from CS to FH. Since this long sequential pathway is necessary for the

downstream analysis, such as FCC and MEC calculation, we summed ACO and irreversible

reaction, IDH, which have a large thermodynamic driving force in the direction of the

oxidative TCA cycle, as a single irreversible reaction, ACO:IDH.

In the skeletal muscle, FBP, G6PC, and PCK in gluconeogenesis were not included in the

metabolic network. Although PC is also involved in gluconeogenesis, PC is included in the

metabolic network of skeletal muscle because we could not exclude the possibility of its

expression in the skeletal muscle (*63*, *64*).

In this study, we assumed the liver and skeletal muscle as a single compartment of

hepatocytes and myocytes, respectively, and cytosolic and mitochondrial compartments in

these cells were not considered for simplification of the modeling. Therefore, the $\Delta_r G'$

calculated in this study represents the average values within the cells.

**Algorithm of GLEAM**

We developed GLEAM as a method to estimate complete set of thermodynamically

consistent metabolite concentrations and $\Delta_f G'^\circ$ values, from metabolomic data on a

logarithmic scale, $\ln \boldsymbol{c} \in \mathbb{R}^{m \times h \times g}$ , and $\Delta_f G'^\circ$ data calculated by the component

contribution method (*65*), $\boldsymbol{\Delta_f G'^\circ} \in \mathbb{R}^m$, where *m*, *h*, *g* are the number of metabolites,

timepoints, and genotypes, respectively. Since metabolomic and $\Delta_f G'^\circ$ data contain

uncertainties and missing values, directly substituting them into the definition equation of

$\Delta_r G'$ often violates the second law of thermodynamics. To address this issue, GLEAM

corrects for the uncertainties and missing values by calculate the most plausible values based

on measured metabolomic data within a thermodynamically feasible space.

We first formulate the quadratic programming problem to minimize the departures of

estimated metabolite concentrations and $\Delta_f G'^\circ$ from the experimental and database values

under thermodynamic and physiological constraints:

$$\min_{y\in\{\ln c,\Delta_f G'^\circ\},a} Z_{\text{QP}}$$

$$= \sum_y \frac{1}{\#y^{\text{exp}}} \left\{ (y + a - y^{\text{exp}})^{\text{T}} \Sigma_{(y)}^{-1} (y + a - y^{\text{exp}}) + \sum_i \lambda \left| \frac{(P_i)^{\text{T}} a}{\sqrt{D_{ii}}} \right| \right\}$$

$$\text{s. t.} \qquad \Delta_r G'_j < 0 \quad \forall i \in M, \forall j \in N$$

$$\Delta_r G'_j = \sum_{i \in M} S_{i,j} \left( \Delta_f G'^\circ_i + RT \ln c_i \right) \quad \forall j \in N^{\text{cell}}$$

(7)

$$\Delta_r G'_{\text{GLUT}} = RT \ln \frac{[\text{Glu[h]}]}{[\text{Glu[b]}]}$$

$$\Delta_r G'_{\text{LDH}} = RT \ln \frac{[\text{H}^+[\text{b}]][\text{Lac[b]}]}{[\text{H}^+[\text{h}]][\text{Lac[h]}]}$$

$$b_i \leq \ln c_i \leq ub_i \quad \forall i \in M$$

$$b_i \leq \ln \frac{c_{i_1}}{c_{i_2}} \leq ub_i \quad \forall (i_1, i_2) \in M^{\text{cofactor}}$$

where $c_i$ is concentration of metabolite $i$, $S_{ij}$ is stoichiometric coefficient of metabolite $i$

in reaction $j$, $R$ is gas constant, and $T$ is temperature. $M$, $N$, and $N^{\text{cell}}$ are sets of

metabolites, reactions, and intracellular reactions, respectively. $M^{\text{cofactor}}$ is the cofactor

pairs of which the ratio was constrained. $\Sigma$ is variance-covariance matrix of which the eigen

decomposition is $\Sigma = \text{PDP}^{\text{T}}$. $a$ is systematic error, and $\lambda$ is regularization parameter. The

superscript exp means measured or database value. The first term in curly brackets of the

objective function represents the variance and covariance-weighted residuals of estimated

metabolite concentrations and $\Delta_f G'^\circ$, which considers systematic errors from experimental

operations, while the second term represents a regularization term for systematic errors to

avoid overfitting. By using the covariances in $\Delta_f G'^\circ$ for the weight of residuals, the

estimation reflects the correlations in $\Delta_f G'^\circ$ due to shared chemical groups. In contrast, all

covariances between metabolite concentrations were set to zero, making $\Sigma$ a diagonal

matrix, since no prior correlation in metabolite concentrations is expected.

The solution of the optimization problem (7) may not provide unique concentrations,

especially for unmeasured metabolites, which can lead to undetermined $\Delta_r G'$ of the

reactions involving the unmeasured metabolites. To estimate unique $\Delta_r G'$, two terms with

variance of unmeasured metabolite concentrations were added to the objective function:

$$
Z_{\text{QP\_unique}}(y, a) = \sum_y \frac{1}{\#y^{\exp}} \left\{ (y + a - y^{\exp})^{\mathrm{T}} \Sigma_{(y)}^{-1} (y + a - y^{\exp}) + \sum_i \lambda \left| \frac{(P_i)^{\mathrm{T}} a}{\sqrt{D_{ii}}} \right| \right\}
$$

$$
+ \epsilon_1 \sum_{(h_1, h_2) \in H^{\text{adj}}} \frac{1}{(h_2 - h_1)} \sum_{i \in M^{\text{unmeasured}}} \left( \ln c_{i, h_2} - \ln c_{i, h_1} \right)^2 \tag{8}
$$

$$
+ \epsilon_2 \sum_{i \in M^{\text{unmeasured}}} \left( \ln c_i - \widehat{\ln c_i} \right)^2
$$

where $H^{\text{adj}}$ is pairs of adjacent timepoints, $M^{\text{unmeasured}}$ is unmeasured metabolites, $\widehat{\ln c}$

is median of upper and lower bounds of metabolite concentration, and $\epsilon_1$ is small values

enough to prioritize the first term over the second term. Similarly, $\epsilon_2$ $(\epsilon_1 \gg \epsilon_2)$ allows the

second term to take precedence over the third term. The second term of the objective function

(the first additional term) reduces the sum of concentration changes of unmeasured

metabolites across time points, making smooth concentration changes more likely to be

estimated than unnatural repetition of increases and decreases. The third term of the objective

function (the second additional term) departures of the estimated unmeasured metabolite

concentrations from the median of their upper and lower bounds. This term can be justified

by the trade-off in increasing concentrations between elevation of enzyme efficiency and

cellular burden from osmotic pressure and toxicity (*40*, *66*). Since the objective function

includes terms with greatly different orders of magnitude due to $\epsilon \in \{\epsilon_1, \epsilon_2\}$, it has difficulty

in optimized at once. Therefore, in implementation, we first solve the optimization problem

only for the first term, then, with the measured metabolite concentrations and $\Delta_f G'^{\circ}$ fixed

to the obtained solutions, and then solve the optimization problem for the second and third

terms.

Metabolic networks can include reversible reactions and their directions are unknown under

a condition of interest. By introducing binary variables representing reaction, GLEAM model

is finally formalized by the following mixed integer quadratic programming problem:

$$\min_{y\in\{\ln c,\Delta_f G'^\circ\},a,v,d} Z_{\text{MIQP}}$$

$$= \sum_y \frac{1}{\#y^{\text{exp}}} \left\{ (y + a - y^{\text{exp}})^{\text{T}} \Sigma_{(y)}^{-1} (y + a - y^{\text{exp}}) + \sum_i \lambda \left| \frac{(P_i)^{\text{T}} a}{\sqrt{D_{ii}}} \right| \right\}$$

$$+ \epsilon_1 \sum_{(h_1,h_2)\in H^{\text{adj}}} \frac{1}{(h_2 - h_1)} \sum_{i\in M^{\text{unmeasured}}} \left( \ln c_{i,h_2} - \ln c_{i,h_1} \right)^2$$

$$+ \epsilon_2 \sum_{i\in M^{\text{unmeasured}}} \left( \ln c_i - \widehat{\ln c_i} \right)^2$$

s.t.
$$d_j \cdot \Delta_r G_j' < 0 \quad \forall i \in M, \forall j \in N$$

$$\Delta_r G_j' = \sum_{i\in M} S_{i,j} \left( \Delta_f G_i'^\circ + RT \ln c_i \right) < 0 \quad \forall i \in M, \forall j \in N^{\text{cell}}$$

$$\Delta_r G_{\text{GLUT}}' = RT \ln \frac{[\text{Glu}[h]]}{[\text{Glu}[b]]}$$

(9)

$$\Delta_r G_{\text{LDH}}' = RT \ln \frac{[\text{H}^+[b]][\text{Lac}[b]]}{[\text{H}^+[h]][\text{Lac}[h]]}$$

$$d_j \in \{-1,1\} \quad \forall j \in N^{\text{rev}}$$

$$d_j \in \{1\} \quad \forall j \in N^{\text{irrev\_f}}$$

$$d_j \in \{-1\} \quad \forall j \in N^{\text{irrev\_b}}$$

$$d_j \cdot v_j > 0 \quad j \in N$$

$$\sum_{j\in N} S_{i,j} v_j = 0 \quad i \in M^{\text{steady}}$$

$$lb_i \leq \ln c_i \leq ub_i \quad \forall i \in M$$

$$lb_i \leq \ln \frac{c_{i_1}}{c_{i_2}} \leq ub_i \quad \forall (i_1, i_2) \in M^{\text{cofactor}}$$

where $v_j$ is metabolic flux through reaction $j$, $d_j = \{1, -1\}$ is a binary variable

representing direction of reaction $j$, and $N^{\mathrm{rev}}$ is reversible reactions. $N^{\mathrm{irrev\_f}}$ and $N^{\mathrm{irrev\_b}}$

are irreversible reactions of which the direction is fixed to forward and backward with respect

to the definition of metabolic network, respectively. $M^{\mathrm{steady}}$ is metabolites assumed to be

in steady state. To solve this mixed integer quadratic programming problem efficiently,

GLEAM treats it as a bilevel optimization problem, where the optimization of binary

variables is the upper-level problem and the optimization of continuous variables is the

lower-level problem:

$$\min_{d} \quad Z_{\mathrm{MIQP}}\left(y^*, a^*, v^*, d\right)$$

$$\mathrm{s.\,t.} \quad y^*, a^*, v^* \in \arg\min_{y,a,v} Z_{\mathrm{MIQP}}(y, a, v, d) \qquad (10)$$

$$y, a, v, d \in \Omega_{\mathrm{GLEAM}}$$

where $\Omega_{\mathrm{GLEAM}}$ is the solution space defined in the problem (9). In the bilevel optimization

problem, binary variables are optimized by a genetic algorithm, while continuous variables

are optimized by the interior-point method under the individuals of binary variables (Fig. S5).

The genetic algorithm was performed by the MATLAB function ga with termination

condition of FunctionTolerance = $10^{-9}$ and population size of PopulationSize = 500. The

lower-level quadratic programming problem was solved using Gurobi (*67*) MATLAB API

with dual feasibility tolerance of OptimalityTol = $10^{-9}$ and primal feasibility tolerance of

FeasibilityTol = $10^{-9}$. This bilevel optimization was performed 10 times from random initial

values, and the solution with the minimum objective function value was selected as the

optimal solution.

**Calculation of confidence intervals**

The confidence intervals of the estimation by GLEAM were calculated using parametric

bootstrap sampling of metabolite concentrations and $\Delta_f G'^\circ$. These data were first resampled

by adding perturbation based on the variance and residual of each metabolite concentration

and $\Delta_f G'^\circ$ in the $\Delta_r G'$ estimation by equation (9) using the experimental and database

values (27, 68). To generate $\Delta_f G'^\circ$ samples reflecting their covariance, the perturbations

were applied after the basis of $\Delta_f G'^\circ$ vector was transformed by transposed matrix of the

eigenvectors of the variance-covariance matrix:

$$\ln c_i^{b,\exp} = \ln c_i^{\exp} + \sigma_i \delta_i^b, \quad \forall b \in B, \tag{11}$$

$$\left(\boldsymbol{P}_{(\Delta_f G'^\circ)i}\right)^{\mathrm{T}} \boldsymbol{\Delta_f G'^{o,b,\exp}} = \left(\boldsymbol{P}_{(\Delta_f G'^\circ)i}\right)^{\mathrm{T}} \boldsymbol{\Delta_f G'^{o,\exp}} + \sqrt{D_{(\Delta_f G'^o)ii}}\,\delta_i^b, \tag{12}$$

$$\boldsymbol{\Sigma}^b = \boldsymbol{\Sigma}, \tag{13}$$

$$\delta_i^b \sim \mathcal{N}(0, (sd^*)^2), \tag{14}$$

$$sd^* = \sqrt{\frac{1}{\#y^{\exp} - 1} \sum_{i \in \text{columns}} \left(\frac{\left(\boldsymbol{P}_{(y)i}\right)^{\mathrm{T}} \boldsymbol{y}^* + \left(\boldsymbol{P}_{(y)i}\right)^{\mathrm{T}} \boldsymbol{a}^* - \left(\boldsymbol{P}_{(y)i}\right)^{\mathrm{T}} \boldsymbol{y}^{\exp}}{\sqrt{D_{(y)ii}}} - err^*\right)^2}, \tag{15}$$

$$err^* = \frac{1}{\#y^{\exp}} (\boldsymbol{y}^* + \boldsymbol{a}^* - \boldsymbol{y}^{\exp})^{\mathrm{T}} \boldsymbol{\Sigma}_{(y)}^{-1} (\boldsymbol{y}^* + \boldsymbol{a}^* - \boldsymbol{y}^{\exp}) \tag{16}$$

where $b$ is the indices of bootstrap samples $B$, $*$ represents the optimized values by

equation (9) using the experimental and database values. $\delta$ is gaussian noise generated from

the normal distribution with mean 0 and standard deviation $sd^*$, which represents the

standard deviation of metabolite concentration and $\Delta_f G'^\circ$ residuals in the $\Delta_r G'$ estimation

from the experimental and database values. Using the resampled metabolite concentrations

and $\Delta_f G'^\circ$ values, we conducted the estimation of thermodynamically consistent metabolite

concentrations and $\Delta_f G'^\circ$ in the same manner as the estimation from the experimental and

database values. In this optimization, the directions of reversible reactions were fixed to those

estimated from the experimental and database values. This resampling and optimization were

performed 2000 times. By applying the percentile method to the resulting 2000 sets of the

estimated values, we obtained the 95% confidence intervals.

**Selection of regularization parameter**

The regularization parameter $\lambda$ for systematic errors in GLEAM was selected by the

reduced chi-square ( $\chi^2_{\text{red}}$ ). $\chi^2_{\text{red}}$ was calculated using the estimated metabolite

concentrations and $\Delta_f G'^\circ$ from resampled values (*27, 68*):

$$\chi^2_{\text{red}} = \frac{\#y^{\text{exp}} EPE}{df}, \tag{17}$$

$$EPE = \frac{1}{\#B} \sum_{b \in B} \frac{1}{\#y^{\text{exp}}} \left(\boldsymbol{y}^{b*} + \boldsymbol{a}^{b*} - \boldsymbol{y}^{\text{exp}}\right)^{\text{T}} \boldsymbol{\Sigma}^{-1}_{(y)} \left(\boldsymbol{y}^{b*} + \boldsymbol{a}^{b*} - \boldsymbol{y}^{\text{exp}}\right), \tag{18}$$

$$df = \frac{\#y^{\text{exp}}}{2sd^*} (EPE - err^*) \tag{19}$$

where $b*$ represents the optimized values calculated from the resampled values. The model

is interpreted as fitting the data ideally when $\chi^2_{\text{red}} = 1$, as overfitting when $\chi^2_{\text{red}} < 1$, and

as poorly fitting when $\chi^2_{\text{red}} > 1$. In this study, $\chi^2_{\text{red}} = 1.03$ when $\lambda = 0.18$, and therefore it

was selected as the best regularization parameter.

**Application of GLEAM to mice data**

We applied GLEAM to absolute quantitative metabolome data in the liver, skeletal muscle,

and plasma from WT and *ob/ob* mice (n = 5) after 0, 2, 4, 6, 8, 12, and 16 hours of fasting

(data file S1), which were measured in our previous studies (*8*, *69*). We used the logarithmic

mean and variance of the metabolites, which were successfully measured in three or more

samples per condition, and metabolites with two or fewer measured samples were treated as

unmeasured metabolites. The cofactors ATP, ADP, GTP, GDP, NAD$^+$, NADH, FAD, FADH2,

NADP$^+$, and NADPH were treated as unmeasured metabolites regardless of the number of

measured samples due to sample degradation, and instead, their ratios were constrained by

previous reports (Table S1). Intracellular lactate was also treated as an unmeasured

metabolite in the same manner as the cofactors because it might leach out of the container

during measurement. The upper and lower bounds for many metabolites in GLEAM were set

to a physiologically typical range of intracellular metabolite concentration (lower bound: $10^-$

$^7$ M, upper bound: $10^{-2}$ M) (*26*), while the bounds for some unmeasured metabolites were

individually defined based on biological knowledge (Table S1).

$\Delta_f G'^\circ$ data calculated by the component contribution method (*65*) were retrieved from

eQuilibrator database (data file S1) (*53*). We used the identical $\Delta_f G'^\circ$ data for both liver and

skeletal muscle under the condition in eQulibrator of pH = 7.2, pMg = 3.0, and ionic strength

= 0.15 M (*26*). We also set the condition in blood to pH = 7.4 (*70*) because $\Delta_r G'$ for MCT

depends on pH in the cells and blood. Temperature was input to $T$ = 310.15 K for all the

compartments.

In the liver, glycolysis was assumed to be dominant at the 0h-fasting state (*ad libitum* feeding)

while gluconeogenesis was assumed to take over after no later than 16h-fasting, both in WT

(*71–75*) and *ob*/*ob* mouse (*76*). At these two timepoints, the directions of reversible reactions

involved in glycolysis and gluconeogenesis (GPI, ALDO, TPI, GAPDH:PGK, PGAM, ENO,

PGM, GPD, LDH) were fixed and treated as those of the irreversible reactions in the

calculation of GLEAM.

Since the reaction rates are sufficiently faster than the temporal changes in metabolite and

enzyme levels during starvation, our metabolic model was assumed to be in a pseudo-steady

state at each measured time point. This assumption leads to mass balance equilibrium for the

intermediate metabolites (G6P, F6P, F1,6BP, DHAP, GAP, 3PG, 2PG, PEP, Pyruvate, AcCoA,

Cit, Isocit, AKG, SucCoA, Suc, Fum, Mal, OAA) within the metabolic network.

**Derivation of FCC**

We derived flux control coefficients (FCCs) within glycolysis (Fig. 3A), gluconeogenesis

(Fig. 3B), and the TCA cycle (Fig. 3C) from the direct binding modular rate law (*48*), which

is equivalent to the reversible Michaelis-Menten kinetics (*77*) for simple reversible uni-uni

reactions, using theorems of Metabolic Control Analysis (MCA) (*36*, *47*). In the direct

binding modular rate law, flux can be described by the effect of the maximal rate, the

thermodynamics, and the saturation of substrate (*49*):

$$v = E k_{\text{cat}}^{+} \eta_{\text{rev}} \eta_{\text{sat,s}}, \tag{20}$$

$$\eta_{\text{rev}} = 1 - \exp\left(\frac{\Delta_r G'}{RT}\right), \tag{21}$$

$$\eta_{\text{sat,s}} = \frac{\prod_{i \in M^s}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}}{1 + \prod_{i \in M^s}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|} + \prod_{i \in M^p}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}} \tag{22}$$

650    where $M^s$ is substrates, $M^p$ is products. From equation (20), we calculated the elasticity of

651    substrate $\epsilon_{s,i}^v$ and product $\epsilon_{p,i}^v$:

$$\epsilon_{s,i}^v = \frac{\partial v}{\partial c_i}\frac{c_i}{v} = |S_i|(\eta_{\text{rev}}^{-1} - \eta_{\text{sat,s}}), \tag{23}$$

$$\epsilon_{p,i}^v = \frac{\partial v}{\partial c_i}\frac{c_i}{v} = |S_i|(1 - \eta_{\text{rev}}^{-1} - \eta_{\text{sat,p}}), \tag{24}$$

$$\eta_{\text{sat,p}} = \frac{\prod_{i \in M^p}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}}{1 + \prod_{i \in M^s}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|} + \prod_{i \in M^p}\left(\frac{c_i}{K_{\text{m},i}}\right)^{|S_i|}}. \tag{25}$$

652    The stoichiometric matrices of the pathways we analyzed here are full rank, and each of their

653    null spaces is one-dimensional, which means the degree of freedom of steady-state flux is

654    one. For these pathways, the following equations (26) and (27) hold from the connectivity

655    theorem and the summation theorem in MCA, respectively (*50*):

$$\mathbf{C}_v^J \boldsymbol{\epsilon}_c^v = \mathbf{0}, \tag{26}$$

$$\sum_j C_{v_j}^J = 1, \tag{27}$$

$$\mathbf{C_v^J} = \begin{pmatrix} C_{v_1}^J & \cdots & C_{v_n}^J \\ \vdots & \ddots & \vdots \\ C_{v_1}^J & \cdots & C_{v_n}^J \end{pmatrix}, \tag{28}$$

$$\boldsymbol{\epsilon_c^v} = \begin{pmatrix} \epsilon_{c_1}^{v_1} & \cdots & \epsilon_{c_k}^{v_1} \\ \vdots & \ddots & \vdots \\ \epsilon_{c_1}^{v_n} & \cdots & \epsilon_{c_k}^{v_n} \end{pmatrix} \tag{29}$$

656    where $\epsilon_c^v = \{\epsilon_s^v, \epsilon_p^v\}$. Integrating equations (23), (24), (26) and (27), we obtained FCC as:

$$C_{v_l}^J = \frac{\alpha_l}{\sum_j \alpha_j}, \tag{30}$$

657    where $\alpha$ for GAPDH:PGK in glycolysis is described as

$$\alpha = \prod_{j=0}^{\text{ALDO}} \left(-\epsilon_p^{v_j}\right) \left(\epsilon_s^{v_{\text{TPI}}} - \epsilon_p^{v_{\text{TPI}}}\right) \prod_{j=l+1}^{n+1} \epsilon_s^{v_j}, \tag{31}$$

658    $\alpha$ for downstream reactions of GAPDH:PGK in glycolysis is described as

$$\alpha = \prod_{j=0}^{\text{ALDO}} \left(-\epsilon_p^{v_j}\right) \left(\epsilon_s^{v_{\text{TPI}}} - \epsilon_p^{v_{\text{TPI}}}\right) \prod_{j=\text{GAPDH:PGK}}^{l-1} \left(-\epsilon_p^{v_l}\right) \prod_{j=l+1}^{n+1} \epsilon_s^{v_j}, \tag{32}$$

659    $\alpha$ for ALDO in gluconeogenesis is described as

$$\alpha = \prod_{j=0}^{\text{GAPDH:PGK}} \left(-\epsilon_p^{v_j}\right) \left(\epsilon_s^{v_{\text{TPI}}} - \epsilon_p^{v_{\text{TPI}}}\right) \prod_{j=l+1}^{n+1} \epsilon_s^{v_j}, \tag{33}$$

660    $\alpha$ for downstream reactions of ALDO in gluconeogenesis is described as

$$\alpha = \prod_{j=0}^{\text{GAPDH:PGK}} \left(-\epsilon_p^{v_j}\right) \left(\epsilon_s^{v_{\text{TPI}}} - \epsilon_p^{v_{\text{TPI}}}\right) \prod_{j=\text{ALDO}}^{l-1} \left(-\epsilon_p^{v_j}\right) \prod_{j=l+1}^{n+1} \epsilon_s^{v_j}, \tag{34}$$

661    $\alpha$ for other reactions is described as

$$\alpha_l = \prod_{j=0}^{l-1}(-\epsilon_\mathrm{p}^{v_j}) \prod_{j=l+1}^{n+1} \epsilon_\mathrm{s}^{v_j}. \tag{35}$$

Note that reaction indices $j \in \{1, \ldots, N\}$ are denoted such that 1 represents the most

upstream reaction and N represents the most downstream reaction in the pathway, and $\epsilon_\mathrm{s}^{v_0} =$

$\epsilon_\mathrm{p}^{v_0} = \epsilon_\mathrm{s}^{v_{n+1}} = \epsilon_\mathrm{p}^{v_{n+1}} = 1$.

**Metabolite concentrations and $\Delta_f G'^\circ$ values used in FCC calculation**

We calculated FCC under arbitrary $\Delta_r G'$ landscapes as well as under the hepatic one. For

both conditions, we used $\Delta_f G'^\circ$ values sampled for confidence interval calculation in

GLEAM. To generate the hepatic $\Delta_r G'$ landscapes, we used metabolite concentrations

sampled in the confidence interval calculation. To generate arbitrary $\Delta_r G'$ landscapes,

logarithmic metabolite concentrations were uniformly sampled from the physiologically

reasonable solution space $\Omega_{\ln c}$:

$$\Omega_{\ln c} = \left\{ \ln c \,\middle|\, \begin{array}{c} \Delta_r G_j' < 0 \\[6pt] lb_i \le \ln c_i \le ub_i \\[6pt] lb_i = ub_i = \ln c_i^{\mathrm{blood}} \quad i \in M^{\mathrm{blood}} \end{array} \right\} \tag{36}$$

673    where $M^{\mathrm{blood}}$ represents blood metabolites. The upper and lower bounds of metabolite

674    concentrations were the same as those used in GLEAM (Table S1). This sampling was

675    performed 20 times for each $\Delta_f G'^\circ$ set using MATLAB function polySampler in the

676    Volume-and-Sampling package (*78*). Thus, we obtained 2000 × 20 = 40000 arbitrary $\Delta_r G'$

677    landscapes in total.

### $K_m$ values used in FCC calculation

In FCC calculation, we sampled Michaelis-Menten constants $K_m$ for each $\Delta_r G'$ landscape

to calculate reaction saturation level. To sample $K_m$, we first retrieved $K_m$ and the turnover

number, $k_{cat} \in \{k_{cat}^+, k_{cat}^-\}$ values for enzymes of *Mus musculus*, *Rattus norvegicus*, *Homo*

*sapiens*, *Bos taurus*, *Sus scrofa*, *Canis lupus familiaris*, and *Mammalia* (*79*) from BRENDA

database (*79*). For $K_m$ and $k_{cat}$ values from three or more data without reports of mutant

enzymes, we calculated their logarithmic means and standard deviations. For $K_m$ and $k_{cat}$

from fewer than three data, we set the logarithmic mean of $k_{cat}$ (/s) to 1 with a standard

deviation of 1.5, and the logarithmic mean of $K_m$ (mM) to −1 with a standard deviation of

1 (*80*). With the means and standard deviations of $K_m$ and $k_{cat}$, we uniformly sampled

logarithmic $K_m$, $k_{cat}$ values for each enzyme from the solution space $\Omega_{KP}$:

$\Omega_{KP,j}$

$$= \left\{ \begin{array}{c} (\log K_m, \\ \log k_{cat}) \end{array} \left| \begin{array}{c} \log \mu^{K_{m,i,j}} - 2\sigma^{K_{m,i,j}} \leq \log K_{m,i,j} \leq \log \mu^{K_{m,i,j}} + 2\sigma^{K_{m,i,j}} \\ \log \mu^{k_{cat,j}} - 2\sigma^{k_{cat,j}} \leq \log k_{cat,j} \leq \log \mu^{k_{cat,j}} + 2\sigma^{k_{cat,j}} \\ \log k_{cat,j}^+ - \log k_{cat,j}^- - \sum_{i \in M^s} \log K_{m,i,j}^{|S_{i,j}|} + \sum_{i \in M^p} \log K_{m,i,j}^{|S_{i,j}|} = \log K_{eq,j} \\ K_{eq,j} = \exp\left( -\frac{1}{RT} \sum_{i \in M} S_{i,j} \Delta_f G_i'^\circ \right) \end{array} \right\}, \quad (37)$$

$\forall i \in M, \forall j \in N$

where $\mu$ is mean, $\sigma$ is standard deviation, and $K_{eq}$ is equilibrium constant. The third

constraint equation reflects the Haldane relationship (*80*). These samplings were performed

20 times for each $\Delta_r G'$ landscape using MATLAB function polySampler in the Volume-

and-Sampling package (*78*). Thus, we finally calculated FCC under the arbitrary and hepatic

$\Delta_r G'$ landscapes with $20 \times 20 \times 2000 = 800000$ and $20 \times 2000 = 40000$ sets of metabolite

concentrations, $\Delta_f G'^{\circ}$ and $K_{\mathrm{m}}$ values, respectively.

**Definitions of EC and total EC**

By solving the equation of the direct binding modular rate law with respect to enzyme amount,

we can obtain enzyme demand needed to carry a certain flux:

$$E \; = \frac{v}{k_{\mathrm{cat}}^{+}} \eta_{\mathrm{rev}}^{-1} \, \eta_{\mathrm{sat,s}}^{-1}. \tag{39}$$

700    With this equation, we defined Enzyme Cost (EC) as the product of the enzyme demand and

701    the enzyme molecular weight *w*:

$$\mathrm{EC} = wE. \tag{40}$$

702    The enzyme molecular weights were retrieved from the UniProt database (*8*, *69*). We also

703    defined pathway's total EC as the sum of ECs in the pathway:

$$\mathrm{total\ EC} = \sum_{j \in N^{\mathrm{pathway}}} \mathrm{EC}_j \tag{41}$$

where $N^{\mathrm{pathway}}$ is reactions included in the pathway. We calculated ECs in glycolysis (Fig.

3A), gluconeogenesis (Fig. 3B), and the TCA cycle (Fig. 3C), which were the same as those

used in FCC calculations, under the hepatic $\Delta_r G'$ landscape using metabolite concentrations

and $\Delta_f G'^{\circ}$ values sampled for confidence interval calculation in GLEAM, and geometric

mean of $K_{\mathrm{m}}$ values from the BRENDA database (*51*) obtained in FCC calculation (data file

S4). For $K_{\mathrm{m}}$ with fewer than three data found in the BRENDA, we set their $c/K_{\mathrm{m}}$ values

to the geometric mean of the pathway. $k_{cat}^+$ values were assumed to be the same within a

pathway, and their effect can be ignored in this study because we only focused on the relative

values of EC.

**Minimization of total EC**

715    Since EC is dependent on metabolite concentrations, we defined Enzyme Cost Minimum

716    (ECM) as a theoretical condition where total EC is minimized with metabolite concentrations.

717    We obtained ECM by solving the following non-linear optimization problem (*26*):

$$
\begin{aligned}
&\min_{\ln c} \quad && \text{total EC} = \sum_{j \in N^{\text{pathway}}} \text{EC}_j \\
&\text{s.t.} \quad && \text{EC}_j = w v \eta_{\text{rev},j}^{-1} \eta_{\text{sat,s},j}^{-1}, \quad j \in N \\
& && \eta_{\text{rev},j} = 1 - \exp\left(\frac{\Delta_r G_j'}{RT}\right) \\
& && \eta_{\text{sat,s},j} = \frac{\prod_{i \in M^{\text{s}}} \left(\frac{c_i}{K_{\text{m},i,j}}\right)^{|S_{i,j}|}}{1 + \prod_{i \in M^{\text{s}}} \left(\frac{c_i}{K_{\text{m},i,j}}\right)^{|S_{i,j}|} + \prod_{i \in M^{\text{p}}} \left(\frac{c_i}{K_{\text{m},i,j}}\right)^{|S_{i,j}|}} \\
& && \Delta_r G_j' = \sum_{i \in M} S_{i,j}\left(\Delta_f G_i'^{\text{o,liver}} + RT \ln c_i\right) < 0, \quad j \in N \\
& && lb_i \leq \ln c_i \leq ub_i, \quad i \in M \\
& && lb_i = ub_i = \ln c_i^{\text{blood}}, \quad i \in M^{\text{blood}} \\
& && lb_i = ub_i = \ln c_i^{\text{liver}}, \quad i \in M^{\text{terminal}} \\
& && \ln \frac{c_{i_1}}{c_{i_2}} = \left(\ln \frac{c_{i_1}}{c_{i_2}}\right)^{\text{liver}}, \quad \forall(i_1, i_2) \in M^{\text{cofactor}}.
\end{aligned}
\tag{42}
$$

*w* and *v* are normalized by the minimum molecular weight and flux, respectively. The relative

flux values for a pathway we analyzed were unique because the degree of freedom of these

pathways' steady flux is 1. $M^{\text{blood}}$ represents blood metabolites, and $M^{\text{terminal}}$ represents

intracellular metabolites at the termini of each pathway: Glucose and Pyr for glycolysis and

gluconeogenesis, and AcCoA, OAA, and Mal for the TCA cycle. By fixing these

concentrations and cofactor ratios to those in mouse liver, the total Gibbs free energy change

across the entire pathway in ECM is consistent with that in the mouse liver. ECM was

calculated by MATLAB function fmincon with termination tolerance on the first-order

optimality of OptimalityTolerance = $10^{-7}$, tolerance on the constraint violation of

ConstraintTolerance = $10^{-9}$, maximum number of function evaluations of

MaxFunctionEvaluations = 1000000, and maximum number of iterations of MaxIterations =

100000. The optimization was performed 100 times for a single problem, and the solution

minimizing the objective function was selected as ECM.

**Calculation of MEG**

To quantify the efficiency of metabolic switching in terms of metabolite concentration

changes, we defined the Metabolite Equilibrium Gap (MEG) as the minimum concentration

changes required to reverse the direction of metabolism:

$$\text{MEG} = \exp A$$

$$A = \min_{\ln c} \sqrt{\frac{1}{\#M} \sum_{i \in M} \left( \ln c_i - \ln c_i^{\text{sys}} \right)^2}$$

$$\text{s.t.} \quad \Delta_r G_j' = \sum_{i \in M} S_{i,j} \left( \Delta_f G_i'^\circ + RT \ln c_i \right) = 0, \quad j \in N^{\text{shift}} \tag{43}$$

$$\Delta_r G_j' = \sum_{i \in M} S_{i,j} \left( \Delta_f G_i'^\circ + RT \ln c_i \right) < 0, \quad j \in N \notin N^{\text{shift}}$$

$$b_i \leq \ln c_i \leq ub_i, \quad j \in M$$

where sys represents a value in a target system corresponding to the liver or ECM in this

study. $N^{\text{shift}}$ is the reactions that must proceed in the reverse direction when switching

metabolisms; it includes reversible reactions in glycolysis and gluconeogenesis, and all

reactions in the TCA cycle. $\Delta_f G'^\circ$ values used in MEG calculation were the same as those

in the liver (and also the same as those in ECM). MEG was calculated using Gurobi (*67*)

MATLAB API with dual feasibility tolerance of OptimalityTol = $10^{-9}$ and primal feasibility

tolerance of FeasibilityTol = $10^{-9}$.

**Implementation and statistical analysis**

Statistical tests, optimizations, computational samplings, and calculations of $\Delta_r G'$, FCC, and

EC were performed using MATLAB 2021b (The MathWorks Inc.) and Gurobi (version

10.0.3) (*67*). Database handlings were performed using Python (version 3.10. All details of

statistical tests are described in the Results section. The MATLAB package Plot Groups of

Stacked Bars (*82*) was used to create the figures.

**Supplementary Materials**

Figs. S1 to S5

Tables S1 to S2

Data files S1 to S5

References (*81*) to (*91*)

## References

1.   M. C. Scrutton, M. F. Utter, The Regulation of Glycolysis and Gluconeogenesis in
Animal Tissues. *Annu. Rev. Biochem.* **37**, 249–302 (1968).

2.   D. L. Rothman, I. Magnusson, L. D. Katz, R. G. Shulman, G. I. Shulman, Quantitation
of Hepatic Glycogenolysis And Gluconeogenesis in Fasting Humans With $^{13}$ C NMR.
*Science* **254**, 573–576 (1991).

3.   R. C. Nordlie, J. D. Foster, A. J. Lange, Regulation of glucose production by the liver.
*Annu. Rev. Nutr.* **19**, 379–406 (1999).

4.   M. C. Moore, K. C. Coate, J. J. Winnick, Z. An, A. D. Cherrington, Regulation of hepatic
glucose uptake and storage in vivo. *Adv. Nutr.* **3**, 286–294 (2012).

5.   H.-S. Han, G. Kang, J. S. Kim, B. H. Choi, S.-H. Koo, Regulation of glucose metabolism
from a liver-centric perspective. *Exp. Mol. Med.* **48**, e218 (2016).

6.   J. Katz, J. A. Tayek, Gluconeogenesis and the Cori cycle in 12-, 20-, and 40-h-fasted
humans. *American Journal of Physiology-Endocrinology and Metabolism* **275**, E537–
E542 (1998).

7.   R. Egami, T. Kokaji, A. Hatano, K. Yugi, M. Eto, K. Morita, S. Ohno, M. Fujii, K.-I.
Hironaka, S. Uematsu, A. Terakawa, Y. Bai, Y. Pan, T. Tsuchiya, H. Ozaki, H. Inoue, S.
Uda, H. Kubota, Y. Suzuki, M. Matsumoto, K. I. Nakayama, A. Hirayama, T. Soga, S.
Kuroda, Trans-omic analysis reveals obesity-associated dysregulation of inter-organ
metabolic cycles between the liver and skeletal muscle. *iScience* **24**, 102217 (2021).

8.   K. Morita, A. Hatano, T. Kokaji, H. Sugimoto, T. Tsuchiya, H. Ozaki, R. Egami, D. Li,
A. Terakawa, S. Ohno, H. Inoue, Y. Inaba, Y. Suzuki, M. Matsumoto, M. Takahashi, Y.
Izumi, T. Bamba, A. Hirayama, T. Soga, S. Kuroda, Structural robustness and temporal
vulnerability of the starvation-responsive metabolic network in healthy and obese mouse
liver. *Sci. Signal.* **18**, eads2547 (2025).

9.   V. T. Samuel, G. I. Shulman, The pathogenesis of insulin resistance: integrating signaling
pathways and substrate flux. *J. Clin. Invest.* **126**, 12–22 (2016).

10.  J. H. Stern, G. I. Smith, S. Chen, R. H. Unger, S. Klein, P. E. Scherer, Obesity
dysregulates fasting-induced changes in glucagon secretion. *J. Endocrinol.* **243**, 149–
160 (2019).

11.  B. Stocks, A. Gonzalez-Franquesa, M. L. Borg, M. Björnholm, L. Niu, J. R. Zierath, A.
S. Deshmukh, Integrated Liver and Plasma Proteomics in Obese Mice Reveals Complex
Metabolic Regulation. *Mol. Cell. Proteomics* **21**, 100207 (2022).

12. Y. Bai, K. Morita, T. Kokaji, A. Hatano, S. Ohno, R. Egami, Y. Pan, D. Li, K. Yugi, S.
Uematsu, H. Inoue, Y. Inaba, Y. Suzuki, M. Matsumoto, M. Takahashi, Y. Izumi, T.
Bamba, A. Hirayama, T. Soga, S. Kuroda, Trans-omic analysis reveals opposite
metabolic dysregulation between feeding and fasting in liver associated with obesity.
*iScience* **27**, 109121 (2024).

13. I. Magnusson, D. L. Rothman, L. D. Katz, R. G. Shulman, G. I. Shulman, Increased rate
of gluconeogenesis in type II diabetes mellitus. A 13C nuclear magnetic resonance study.
*J. Clin. Invest.* **90**, 1323–1327 (1992).

14. R. S. Hundal, M. Krssak, S. Dufour, D. Laurent, V. Lebon, V. Chandramouli, S. E.
Inzucchi, W. C. Schumann, K. F. Petersen, B. R. Landau, G. I. Shulman, Mechanism by
which metformin reduces glucose production in type 2 diabetes. *Diabetes* **49**, 2063–2069
(2000).

15. A. Wajngot, V. Chandramouli, W. C. Schumann, K. Ekberg, P. K. Jones, S. Efendic, B.
R. Landau, Quantitative contributions of gluconeogenesis to glucose production during
fasting in type 2 diabetes mellitus. *Metabolism* **50**, 47–52 (2001).

16. K. F. Petersen, T. B. Price, R. Bergeron, Regulation of Net Hepatic Glycogenolysis and
Gluconeogenesis during Exercise: Impact of Type 1 Diabetes. *J. Clin. Endocrinol. Metab.*
**89**, 4656–4664 (2004).

17. M. C. Petersen, D. F. Vatner, G. I. Shulman, Regulation of hepatic glucose metabolism
in health and disease. *Nat. Rev. Endocrinol.* **13**, 572–587 (2017).

18. T. K. Schramm, G. H. Gislason, L. Køber, S. Rasmussen, J. N. Rasmussen, S. Z.
Abildstrøm, M. L. Hansen, F. Folke, P. Buch, M. Madsen, A. Vaag, C. Torp-Pedersen,
Diabetes Patients Requiring Glucose-Lowering Therapy and Nondiabetics With a Prior
Myocardial Infarction Carry the Same Cardiovascular Risk. *Circulation* **117**, 1945–1954
(2008).

19. E. A. Spencer, K. L. Pirie, R. J. Stevens, V. Beral, A. Brown, B. Liu, J. Green, G. K.
Reeves, Diabetes and modifiable risk factors for cardiovascular disease: the prospective
Million Women Study. *Eur. J. Epidemiol.* **23**, 793–799 (2008).

20. S. Gancheva, T. Jelenik, E. Álvarez-Hernández, M. Roden, Interorgan metabolic
crosstalk in human insulin resistance. *Physiol. Rev.* **98**, 1371–1415 (2018).

21. T. M. Annesley, Ion suppression in mass spectrometry. *Clin. Chem.* **49**, 1041–1044
(2003).

22. M. R. Mashego, L. Wu, J. C. Van Dam, C. Ras, J. L. Vinke, W. A. Van Winden, W. M.
Van Gulik, J. J. Heijnen, MIRACLE: mass isotopomer ratio analysis of U-13C-labeled
extracts. A new method for accurate quantification of changes in concentrations of
intracellular metabolites. *Biotechnol. Bioeng.* **85**, 620–628 (2004).

23. M. Sugimoto, A. Hirayama, M. Robert, S. Abe, T. Soga, M. Tomita, Prediction of
metabolite identity from accurate mass, migration time prediction and isotopic pattern
information in CE-TOFMS data. *Electrophoresis* **31**, 2311–2318 (2010).

24. B. D. Bennett, E. H. Kimball, M. Gao, R. Osterhout, S. J. Van Dien, J. D. Rabinowitz,
Absolute metabolite concentrations and implied enzyme active site occupancy in
Escherichia coli. *Nat. Chem. Biol.* **5**, 593–599 (2009).

25. A. B. Canelas, C. Ras, A. ten Pierick, W. M. van Gulik, J. J. Heijnen, An in vivo data-
driven framework for classification and quantification of enzyme kinetics and
determination of apparent thermodynamic data. *Metab. Eng.* **13**, 294–306 (2011).

26. J. O. Park, S. A. Rubin, Y. F. Xu, D. Amador-Noguez, J. Fan, T. Shlomi, J. D. Rabinowitz,
Metabolite concentrations, fluxes and free energies imply efficient enzyme usage. *Nat.*
*Chem. Biol.* **12**, 482–489 (2016).

27. B. Niebel, S. Leupold, M. Heinemann, An upper limit on Gibbs energy dissipation
governs cellular metabolism. *Nature Metabolism* **1**, 125–132 (2019).

28. M. L. Mavrovouniotis, Identification of localized and distributed bottlenecks in
metabolic pathways. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* **1**, 275–283 (1993).

29. C. S. Henry, M. D. Jankowski, L. J. Broadbelt, V. Hatzimanikatis, Genome-Scale
Thermodynamic Analysis of Escherichia coli Metabolism. *Biophys. J.* **90**, 1453–1461
(2006).

30. A. M. Feist, C. S. Henry, J. L. Reed, M. Krummenacker, A. R. Joyce, P. D. Karp, L. J.
Broadbelt, V. Hatzimanikatis, B. Ø. Palsson, A genome-scale metabolic reconstruction
for Escherichia coli K-12 MG1655 that accounts for 1260 ORFs and thermodynamic
information. *Mol. Syst. Biol.* **3**, 121 (2007).

31. C. S. Henry, L. J. Broadbelt, V. Hatzimanikatis, Thermodynamics-based metabolic flux
analysis. *Biophys. J.* **92**, 1792–1805 (2007).

32. K. C. Soh, V. Hatzimanikatis, Network thermodynamics in the post-genomic era. *Curr.*
*Opin. Microbiol.* **13**, 350–357 (2010).

33. J. J. Hamilton, V. Dwivedi, J. L. Reed, Quantitative assessment of thermodynamic
constraints on the solution space of genome-scale metabolic models. *Biophys. J.* **105**,
512–522 (2013).

34. L. Seep, Z. Razaghi-Moghadam, Z. Nikoloski, Reaction lumping in metabolic networks
for application with thermodynamic metabolic flux analysis. *Sci. Rep.* **11**, 8544 (2021).

35. M. Janasch, N. Crang, J. Asplund-Samuelsson, E. Sporre, M. Bruch, A. Gynnå, M. Jahn, E. P. Hudson, Thermodynamic limitations of PHB production from formate and fructose in Cupriavidus necator. *Metab. Eng.* **73**, 256–269 (2022).

36. H. Kacser, J. A. Burns, The control of flux. *Symp. Soc. Exp. Biol.* **27**, 65–104 (1973).

37. Z. Dai, J. W. Locasale, Thermodynamic constraints on the regulation of metabolic fluxes. *J. Biol. Chem.* **293**, 19725–19739 (2018).

38. D. D. van Niekerk, E. Rust, F. Bruggeman, H. V. Westerhoff, J. L. Snoep, Is distance from equilibrium a good indicator for a reaction's flux control? *Biosystems.* **232**, 104988 (2023).

39. A. Flamholz, E. Noor, A. Bar-Even, W. Liebermeister, R. Milo, Glycolytic strategy as a tradeoff between energy yield and protein cost. *Proceedings of the National Academy of Sciences* **110**, 10039–10044 (2013).

40. N. Tepper, E. Noor, D. Amador-Noguez, H. S. Haraldsdóttir, R. Milo, J. Rabinowitz, W. Liebermeister, T. Shlomi, Steady-State Metabolite Concentrations Reflect a Balance between Maximizing Enzyme Efficiency and Minimizing Total Metabolite Load. *PLoS One* **8**, e75370 (2013).

41. E. Noor, A. Bar-Even, A. Flamholz, E. Reznik, W. Liebermeister, R. Milo, Pathway Thermodynamics Highlights Kinetic Obstacles in Central Metabolism. *PLoS Comput. Biol.* **10** (2014).

42. E. Noor, A. Flamholz, A. Bar-Even, D. Davidi, R. Milo, W. Liebermeister, The Protein Cost of Metabolic Fluxes: Prediction from Enzymatic Rate Laws and Cost Minimization. *PLoS Comput. Biol.* **12**, e1005167 (2016).

43. T. Soga, Y. Ohashi, Y. Ueno, H. Naraoka, M. Tomita, T. Nishioka, Quantitative metabolome analysis using capillary electrophoresis mass spectrometry. *J. Proteome Res.* **2**, 488–494 (2003).

44. T. Soga, T. Ishikawa, S. Igarashi, K. Sugawara, Y. Kakazu, M. Tomita, Analysis of nucleotides by pressure-assisted capillary electrophoresis-mass spectrometry using silanol mask technique. *J. Chromatogr. A* **1159**, 125–133 (2007).

45. A. M. Ingalls, M. M. Dickie, G. D. Snell, Obese, a new mutation in the house mouse. *J. Hered.* **41**, 317–318 (1950).

46. P. Lindström, The physiology of obese-hyperglycemic mice [ob/ob mice]. *ScientificWorldJournal* **7**, 666–685 (2007).

47. R. Heinrich, T. A. Rapoport, A linear steady-state treatment of enzymatic chains. General properties, control and effector strength. *Eur. J. Biochem.* **42**, 89–95 (1974).

48.  W. Liebermeister, J. Uhlendorf, E. Klipp, Modular rate laws for enzymatic reactions:
thermodynamics, elasticities and implementation. *Bioinformatics* **26**, 1528–1534 (2010).

49.  E. Noor, A. Flamholz, W. Liebermeister, A. Bar-Even, R. Milo, A note on the kinetics of
enzyme action: A decomposition that highlights thermodynamic effects. *FEBS Lett.* **587**,
2772–2777 (2013).

50.  J. H. Hofmeyr, A. Cornish-Bowden, Co-response analysis: a new experimental strategy
for metabolic control analysis. *J. Theor. Biol.* **182**, 371–380 (1996).

51.  A. Chang, L. Jeske, S. Ulbrich, J. Hofmann, J. Koblitz, I. Schomburg, M. Neumann-
Schaal, D. Jahn, D. Schomburg, BRENDA, the ELIXIR core data resource in 2021: new
developments and updates. *Nucleic Acids Res.* **49**, D498–D508 (2021).

52.  R. Schuetz, N. Zamboni, M. Zampieri, M. Heinemann, U. Sauer, Multidimensional
Optimality of Microbial Metabolism. *Science* **336**, 601–604 (2012).

53.  M. E. Beber, M. G. Gollub, D. Mozaffari, K. M. Shebek, A. I. Flamholz, R. Milo, E.
Noor, eQuilibrator 3.0: a database solution for thermodynamic constant estimation.
*Nucleic Acids Res.* **50**, D603–D609 (2022).

54.  L. B. Tanner, A. G. Goglia, M. H. Wei, T. Sehgal, L. R. Parsons, J. O. Park, E. White, J.
E. Toettcher, J. D. Rabinowitz, Four key steps control glycolytic flux in mammalian cells.
*Cell Syst.* **7**, 49-62.e8 (2018).

55.  A. K. Groen, C. W. van Roermund, R. C. Vervoorn, J. M. Tager, Control of
gluconeogenesis in rat liver cells. Flux control coefficients of the enzymes in the
gluconeogenic pathway in the absence and presence of glucagon. *Biochem. J* **237**, 379–
389 (1986).

56.  M. Rigoulet, X. M. Leverve, P. J. Plomp, A. J. Meijer, Stimulation by glucose of
gluconeogenesis in hepatocytes isolated from starved rats. *Biochem. J.* **245**, 661–668
(1987).

57.  M. Milani, A. Pesce, M. Bolognesi, A. Bocedi, P. Ascenzi, Substrate channeling:
Molecular bases. *Biochem. Mol. Biol. Educ.* **31**, 228–233 (2003).

58.  A. Datta, J. M. Merz, H. O. Spivey, Substrate channeling of oxalacetate in solid-state
complexes of malate dehydrogenase and citrate synthase. *J. Biol. Chem.* **260**, 15008–
15012 (1985).

59.  F. Wu, S. Minteer, Krebs cycle metabolon: structural evidence of substrate channeling
revealed by cross-linking and mass spectrometry. *Angew. Chem. Int. Ed Engl.* **54**, 1851–
1854 (2015).

60. L. E. Anderson, I. M. Goldhaber-Gordon, D. Li, X. Y. Tang, M. Xiang, N. Prakash,
Enzyme-enzyme interaction in the chloroplast: glyceraldehyde-3-phosphate
dehydrogenase, triose phosphate isomerase and aldolase. *Planta* **196**, 245–255 (1995).

61. D. Araiza-Olivera, N. Chiquete-Felix, M. Rosas-Lemus, J. G. Sampedro, A. Peña, A.
Mujica, S. Uribe-Carvajal, A glycolytic metabolon in Saccharomyces cerevisiae is
stabilized by F-actin. *FEBS J.* **280**, 3887–3905 (2013).

62. A. Bordbar, A. M. Feist, R. Usaite-Black, J. Woodcock, B. O. Palsson, I. Famili, A multi-
tissue type genome-scale metabolic network for analysis of whole-body systems
physiology. *BMC Syst. Biol.* **5** (2011).

63. A. H. Lancha Jr, M. B. Recco, R. Curi, Pyruvate carboxylase activity in the heart and
skeletal muscles of the rat. Evidence for a stimulating effect of exercise. *Biochem. Mol.*
*Biol. Int.* **32**, 483–489 (1994).

64. A. D. Minet, M. Gaster, Pyruvate carboxylase is expressed in human skeletal muscle.
*Biochem. Biophys. Res. Commun.* **402**, 196–197 (2010).

65. E. Noor, H. S. Haraldsdóttir, R. Milo, R. M. T. Fleming, Consistent Estimation of Gibbs
Energy Using Component Contributions. *PLoS Comput. Biol.* **9** (2013).

66. S. Schuster, R. Heinrich, Minimization of intermediate concentrations as a suggested
optimality principle for biochemical networks: I. Theoretical analysis. *J. Math. Biol.* **29**,
425–442 (1991).

67. L. L. C. Gurobi Optimization, Gurobi Optimizer Reference Manual (2024).
https://www.gurobi.com/.

68. T. Hastie, R. Tibshirani, J. Friedman, *The Elements of Statistical Learning: Data Mining,*
*Inference, and Prediction, Second Edition* (Springer, New York, NY, ed. 2,
2009)*Springer series in statistics*.

69. D. Li, K. Morita, T. Kokaji, A. Hatano, A. Hirayama, T. Soga, Y. Suzuki, M. Matsumoto,
T. Tsuchiya, H. Ozaki, S. Ohno, H. Inoue, Y. Inaba, H. Sugimoto, Y. Pan, S. Kuroda,
Global loss of responsiveness in key regulator metabolites and elevated enzyme proteins
as metabolic dysregulation in skeletal muscle and liver of obese mice during starvation,
*bioRxiv* (2025)p. 2025.01.22.634184.

70. H. S. Haraldsdóttir, I. Thiele, R. M. T. Fleming, Quantitative assignment of reaction
directionality in a multicompartmental human metabolic reconstruction. *Biophys. J.* **102**,
1703–1711 (2012).

71. S. C. Burgess, F. M. H. Jeffrey, C. Storey, A. Milde, N. Hausler, M. E. Merritt, H. Mulder,
C. Holm, A. D. Sherry, C. R. Malloy, Effect of murine strain on metabolic pathways of
glucose production after brief or prolonged fasting. *Am. J. Physiol. Endocrinol. Metab.*
**289**, E53-61 (2005).

72. S. Satapati, N. E. Sunny, B. Kucejova, X. Fu, T. T. He, A. Méndez-Lucas, J. M. Shelton,
J. C. Perales, J. D. Browning, S. C. Burgess, Elevated TCA cycle function in the
pathology of diet-induced hepatic insulin resistance and fatty liver. *J. Lipid Res.* **53**,
1080–1092 (2012).

73. C. M. Hasenour, M. L. Wall, D. E. Ridley, C. C. Hughey, F. D. James, D. H. Wasserman,
J. D. Young, Mass spectrometry-based microassay of (2)H and (13)C plasma glucose
labeling to quantify liver metabolic fluxes in vivo. *Am. J. Physiol. Endocrinol. Metab.*
**309**, E191-203 (2015).

74. C. M. Hasenour, M. Rahim, J. D. Young, In vivo estimates of liver metabolic flux
assessed by 13C-propionate and 13C-lactate are impacted by tracer recycling and
equilibrium assumptions. *Cell Rep.* **32**, 107986 (2020).

75. Y. Wang, H. Kwon, X. Su, F. E. Wondisford, Glycerol not lactate is the major net carbon
source for gluconeogenesis in mice during both short and prolonged fasting. *Molecular*
*Metabolism* **31**, 36–44 (2020).

76. S. M. Turner, P. A. Linfoot, R. A. Neese, M. K. Hellerstein, Sources of plasma glucose
and liver glycogen in fasted ob/ob mice. *Acta Diabetol.* **42**, 187–193 (2005).

77. J. B. S. Haldane, *Enzymes* (Longmans, Green and Co., London, England, 1930).

78. B. Cousins, Volume-and-Sampling - File Exchange - MATLAB CentralFile Exchange -
MATLAB                   Central                  (2022).
https://www.mathworks.com/matlabcentral/fileexchange/43596-volume-and-sampling.

79. S. Ohno, L.-E. Quek, J. R. Krycer, K. Yugi, A. Hirayama, S. Ikeda, F. Shoji, K. Suzuki,
T. Soga, D. E. James, S. Kuroda, Kinetic trans-omic analysis reveals key regulatory
mechanisms for insulin-regulated glucose metabolism in adipocytes. *iScience* **23**,
101479 (2020).

80. T. Lubitz, M. Schulz, E. Klipp, W. Liebermeister, Parameter balancing in kinetic models
of cell metabolism. *J. Phys. Chem. B* **114**, 16298–16303 (2010).

81. UniProt Consortium, UniProt: The universal protein knowledgebase in 2023. *Nucleic*
*Acids Res.* **51**, D523–D531 (2023).

82. Evan, Plot Groups of Stacked Bars - File Exchange - MATLAB CentralFile Exchange -
MATLAB Central (2025). https://jp.mathworks.com/matlabcentral/fileexchange/32884-
plot-groups-of-stacked-bars.

**Acknowledgments**

We thank our laboratory members for critically reading this manuscript. The computational

analysis of this work was performed in part with support of the supercomputer system of the

National Institute of Genetics (NIG), Research Organization of Information and Systems

(ROIS). We used ChatGPT, DeepL, and Claude to assist in manuscript editing and coding.

**Funding**

This study was supported by the Japan Society for the Promotion of Science (JSPS)

KAKENHI (Grant Numbers JP17H06300, JP17H06299, JP18H03979, JP21H04759,

JP22K17992, JP23H04946, JP23H04939) (S.K.); the Japan Science and Technology Agency

(JST) as part of CREST (JPMJCR2123, JPMJCR25T3) and as part of ASPIRE

(JPMJAP24B1) (S.K.); The Uehara Memorial Foundation (S.K.); AMED Grant Number

JP21zf0127001 (T.S.); JST, CREST Grant Number JPMJCR2123 (T.S.); MEXT KAKENHI

Grant Number JP23H04946 (T.S.); World Premier International Research Center Initiative

(WPI), Human Biology-Microbiome-Quantum Research Center (Bio2Q), MEXT, Japan

(T.S.); JSPS KAKENHI JP24K11417 (A.H.); and the FOREST Program of the Japan Science

and Technology Agency (JPMJFR2052) (A.H.).

1010

**Author contributions**

Conceptualization: T.A., S.O., and S.K.

Methodology: T.A., S.O., A.H., and T.S.

Investigation: T.A., A.H., and T.S.

Visualization: T.A.

Funding acquisition: A.H., T.S., and S.K.

Project administration: S.K.

Supervision: S.K.

Writing – original draft: T.A., S.O., and S.K.

Writing – review & editing: T.A., S.O., A.H., T.S., and S.K.

1021

1022     **Competing interests**

1023     The authors declare no competing interests.

1024

**Data and materials availability**

The information on kinetic parameter (Michaelis-Menten constant and turnover number)

distributions used for FCC calculation, the FCC values generated in this study, and the

MATLAB code for GLEAM and the analysis of FCC and EC are deposited in GitHub

(https://github.com/tabekawa/Thermodynamics_MouseLiver). All other data needed to

evaluate the conclusions in the paper are present in the paper or the supplementary materials.

1031

1032

**Fig. 1: $\Delta_r G'$ landscape is robust against fasting despite metabolite concentration fluctuations.**

(**A**) Overview of the application of GLEAM to estimate $\Delta_r G'$ landscape in the mouse liver. Metabolomic data were acquired from the liver, skeletal muscle, and plasma of WT and *ob/ob* mice during fasting (data file S1) (*8, 68*). *n* = 5 biological replicates were used per group. The estimated metabolite concentrations, $\Delta_f G'^\circ$, and $\Delta_r G'$ values by GLEAM are shown in data file S2. (**B**) $\Delta_r G'$ landscape of glucose metabolism in mouse liver at *ad libitum* feeding and after 16-hour fasting. $\Delta_r G'$ is depicted by arrow colors. Shaded texts represent

reaction names; boxed reactions are irreversible, while unboxed reactions are reversible. Blue

shadings indicate near-equilibrium reactions with $\Delta_r G' > -5$ kJ/mol, and gray shadings

indicate the other reactions with $\Delta_r G' \leq -5$ kJ/mol. Unshaded texts represent metabolites.

[h], [m], and [b] indicate hepatic, myocytic, and blood metabolites, respectively.

Abbreviations of reactions and metabolites are defined in the glossary and data file S3. (**C**)

Comparison of $\Delta_r G'$ in WT mouse liver at *ad libitum* feeding and after 16-hour fasting. The

forward direction ($\Delta_r G' < 0$) is defined as the reaction direction at *ad libitum* feeding.

Pearson's correlation coefficient (r) is shown. Error bars show 95% confidence intervals. (**D**)

Comparison of metabolite concentrations in WT mouse liver at *ad libitum* feeding and after

16-hour fasting. Plotted metabolite concentrations represent those normalized by the mean

value at all seven time points and two genotypes (14 conditions in total) to eliminate scale

differences in concentrations among metabolites. Pearson's correlation coefficient (r) is

shown. Error bars show 95% confidence intervals. Error bars show 95% confidence intervals.

(**E**) Comparison of $\Delta_r G'$ in glycolysis across cell types and organisms. The data in iBMK

cell, yeast, and *E. coli* were measured using LC-MS and isotope tracers (*26*). Standard Gibbs

free energy was calculated by treating all metabolite concentrations as 1 M. The vertical axis

represents cumulative $\Delta_r G'$ from GPI. Blue and orange bars depict negative and positive

$\Delta_r G'$, respectively. Error bars show 95% confidence intervals.

1059

1060

1061

**Fig. 2: $\Delta_r G'$ maintenance during fasting is associated with metabolite concentration fluctuations in different ways among pathways.**

(**A**) Absolute $\Delta_r G'$ variation = $|\Delta_r G'_{16h} - \Delta_r G'_{0h}|$. Bars depict irreversible reactions in glycolysis and gluconeogenesis (orange), reversible reactions in glycolysis and gluconeogenesis (blue), irreversible reactions in the TCA cycle (yellow), reversible reactions in the TCA cycle (purple), and the other reactions in the metabolic network (grey), respectively. Error bars show 95% confidence intervals. (**B**) Relationship between absolute

$\Delta_r G'$ variation and reaction distance from equilibrium represented by absolute $\Delta_r G'$ at *ad*

*libitum* feeding. Error bars show 95% confidence intervals. The reaction PC is highlighted

and discussed in the Results section. (**C**) Comparison of absolute $\Delta_r G'$ variation of reactions

and total absolute $\Delta_f G'$ variation of metabolites. Absolute $\Delta_f G'$ variation is dependent on

concentrations of metabolites involved in the reaction, and defined by $\sum \left| S \left( \Delta_f G'_{16h} - \right.\right.$

$\left.\left. \Delta_f G'_{0h} \right) \right|$. When $\Delta_f G'$ variation $= S \left( \Delta_f G'_{16h} - \Delta_f G'_{0h} \right)$ of the metabolites involved in the

reaction are cancelled out, absolute $\Delta_r G'$ variation is smaller than total absolute $\Delta_f G'$

variation, resulting in the reaction being plotted in the lower right. Plotted reactions for

glycolysis and gluconeogenesis, and the TCA cycles are the same as Fig. 2B. Error bars show

95% confidence intervals. The reactions GPD, GAPDH:PGK and GCK are highlighted and

discussed in the Results section. (**D**) Time courses of $\Delta_r G'$ variation and $\Delta_f G'$ variation

for representative reactions: GPD (small $\Delta_f G'$ variations), GAPDH:PGK (large $\Delta_f G'$

variations which are canceled out), and GCK (large $\Delta_f G'$ variations which are not canceled

out). Red and grey lines depict $\Delta_r G'$ variation and $\Delta_f G'$ variation, respectively. Error bars

show 95% confidence intervals.

1084

**Fig. 3: Thermodynamic Rate-limiting candidates (TRCs) were maintained during**

**fasting.**

(**A**) Glycolysis pathway where FCCs were calculated. (**B**) Gluconeogenesis pathway where

FCCs were calculated. (**C**) The TCA cycle where FCCs were calculated. MDH was excluded

from this TCA cycle because it proceeded in the opposite direction to the oxidative TCA

cycle. (**D**) Correspondence between $\Delta_r G'$ and $\eta_{rev}$. (**E-G**) Relationship between $\eta_{rev}$ and

FCC in glycolysis (**E**), gluconeogenesis (**F**), and the TCA cycle (**G**) in the mouse liver. $\eta_{rev}$

and FCC values of each point were calculated from each sampled set of metabolite

concentrations and $\Delta_f G'^\circ$ values for the calculation of confidence intervals and sampled

$K_m$ values based on BRENDA database (see Methods). Metabolite concentrations at *ad*

*libitum* feeding were used for glycolysis (**E**) and the TCA cycle (**G**), and those after 16-hour

fasting were used for gluconeogenesis (**F**). Vertical variation is attributed to differences in

$K_m$ values and condition of the other reactions, while horizontal variation reflects differences

in metabolite concentrations and $\Delta_f G'^\circ$. (**H**) Alterations of TRC status based on transitions

in $\eta_{rev}$ for reactions in glycolysis and gluconeogenesis before and after starvation. Red, blue,

and grey lines depict reactions remaining TRC (5 reactions), reactions remaining non-TRC

(10 reactions), and reactions altering their TRC status (either TRC or not) during fasting (3

reactions), respectively. Error bars show 95% confidence intervals.

1104

1105

1106

**Fig. 4: Glycolysis and gluconeogenesis achieve efficient switching by expending large**

**enzyme costs.**

(**A**) Ratio of total enzyme cost under metabolite concentrations in the mouse liver and ECM

(Liver/ECM). The total EC was calculated for glycolysis, gluconeogenesis, and the TCA

cycle, which are the same as the pathways used for FCC calculation (Fig. 3A-C). Metabolite

concentrations at *ad libitum* feeding and after 16-hour fasting were used for glycolysis and

gluconeogenesis, respectively. $K_m$ values for EC calculation were the geometric mean of

data in the BRENDA database (data file S4). Error bars show 95% confidence intervals. EC

in mouse liver, and metabolite concentrations and EC at ECM are shown in data file S4. (**B**)

Comparison of metabolite concentrations in the mouse liver and ECM. Plotted data are for

the measured metabolites, and the scatter plot for the TCA cycle includes the data both at *ad*

*libitum* feeding and after 16-hour fasting. RMSE is the root mean square error on a $\log_{10}$-

scale of ECM's prediction compared to metabolite concentrations in the mouse liver, and

typical fold error (typical FE) = $10^{\text{RMSE}}$. Error bars show 95% confidence intervals. Plotted

data are for the measured metabolites. (**C**) EC of each reaction in glycolysis, gluconeogenesis,

and the TCA cycle. The left and right bars of each reaction are EC in the mouse liver and

ECM, respectively. Since the vertical axis represents a logarithmic scale, EC can be depicted

by the stacked bar of normalized enzyme molecular weight (green), the effect of

thermodynamics, $\eta_{\text{rev}}^{-1}$ (red), and saturation, $\eta_{\text{sat},s}^{-1}$ (blue). Note that although the total

Gibbs free energy change across the entire pathway in ECM is fixed to that in the mouse liver

(see Methods), the sum of $\eta_{\text{rev}}^{-1}$ differs due to the non-linear increase of $\eta_{\text{rev}}$ over $\Delta_r G'$

(Fig. 3D). Error bars show 95% confidence intervals. (**D**) Comparison of metabolite

concentrations in the liver or ECM, and those for MEG calculation, where the reactions

required to reverse for metabolic switching are at equilibrium. Plotted data are for the

measured metabolites. Error bars show 95% confidence intervals. Metabolite concentrations

where the reactions required to reverse for metabolic switching are at equilibrium are shown

in data file S5.

1134

1135

**Fig. 5: Robustness of $\Delta_r G'$ in the liver against obesity.**

(**A**) $\Delta_r G'$ landscape of glucose metabolism in *ob/ob* mouse liver at *ad libitum* feeding and after 16-hour fasting. $\Delta_r G'$ is depicted by arrow colors. Shaded texts represent reaction names; boxed reactions are irreversible, while unboxed reactions are reversible. Blue shadings indicate near-equilibrium reactions with $\Delta_r G' > -5$ kJ/mol, and gray shadings indicate the other reactions with $\Delta_r G' \leq -5$ kJ/mol. Unshaded texts represent metabolites.

The estimated metabolite concentrations, $\Delta_f G'^\circ$, and $\Delta_r G'$ values by GLEAM are shown

in data file S2. (**B**) Comparison of $\Delta_r G'$ in WT and *ob*/*ob* mouse liver. The forward direction

($\Delta_r G' < 0$) is defined as the reaction direction in WT at *ad libitum* feeding. Pearson's

correlation coefficients (r) are shown. Error bars show 95% confidence intervals. (**C**)

Difference in $\Delta_r G'$ between WT and *ob*/*ob* mouse liver ($\Delta_r G'_{ob/ob} - \Delta_r G'_{WT}$). Reactions are

arranged from left to right in order of decreasing absolute difference in $\Delta_r G'$. Error bars show

95% confidence intervals. (**D**) Comparison of metabolite concentrations in WT and *ob*/*ob*

mouse liver. Plotted data represent the concentrations normalized by the mean value at all

seven time points and two genotypes *(i.e.*, 14 conditions) to eliminate scale differences in

concentrations among metabolites. Pearson's correlation coefficients (r) are shown. Error

bars show 95% confidence intervals. (**E**) Comparison of absolute difference in $\Delta_r G'$

variation and total absolute deference in $\Delta_f G' = \Sigma |S(\Delta_f G'_{ob/ob} - \Delta_f G'_{WT})|$, between WT

and *ob*/*ob* mouse liver. Plotted data includes those at *ad libitum* feeding and after 16-hour

fasting. The reactions for glycolysis and gluconeogenesis, and the TCA cycles are the same

as Fig. 2B. Error bars show 95% confidence intervals.

1157

1158

**Fig. S1: Relationship between $\eta_{\mathrm{rev}}$ and FCC for arbitrary $\Delta_r G'$ landscapes.**

(**A-D**) Arbitrary $\Delta_r G'$ landscapes were generated by the uniform sampling of intrasellar

metabolite concentrations under the blood metabolite concentrations at *ad libitum* feeding

and after 16-hour fasting (see Methods). The arbitrary $\Delta_r G'$ landscapes at *ad libitum* feeding

were used for FCC calculation in glycolysis (**A**) and the TCA cycle (**C**), while those after 16-

hour fasting were used for gluconeogenesis (**B**), and the TCA cycle (**D**). Plotted data

represents each sampled set of metabolite concentrations and $\Delta_f G'^\circ$, and $K_m$ values.

1166

Fig. S2: Relationship between $\eta_{\mathrm{rev}}$ and FCC in the TCA cycle in the mouse liver after

16-hour fasting.

Plotted data represents each set of metabolite concentrations and $\Delta_f G'^{\circ}$ values sampled for

the calculation of confidence intervals, and $K_{\mathrm{m}}$ values sampled based on BRENDA database

(see Methods).

1174

**Fig. S3: Typical fold error of ECM's prediction compared to metabolite concentrations in the mouse liver and *E. coli*.**

Glyc. and Gluc. means glycolysis and gluconeogenesis, respectively. Minimizations of total EC in *E. coli* were conducted for the metabolic network, which consists of glycolysis, the pentose phosphate pathway, and the TCA cycle, assuming four different enzymatic rate equations (see Methods).

1181

1182

**Fig. S4: A whole metabolic network used to estimate $\Delta_r G'$ landscape.**

Shaded texts represent reaction names; boxed reactions are irreversible, while unboxed

reactions are reversible. Unshaded texts represent metabolites. Although the network consists

of liver, blood, and skeletal muscle to improve the accuracy of $\Delta_f G'^\circ$ estimation, only the

$\Delta_r G'$ landscape of reactions and transporters in the liver were discussed in this study because

the concentration of unmeasured major metabolites, such as glucose in skeletal muscle,

cannot be estimated with sufficient accuracy due to a lack of reactions.

1190

1191 **Fig. S5: Flowchart of the bilevel optimization problem in GLEAM.**

1192 The optimization problems for the continuous and binary variables are treated as the lower-

1193 level and upper-level problems, respectively. The optimal solution of GLEAM was selected

1194 after solving this bilevel optimization 10 times.

1195

1196 **Supplementary tables**

1197 **Table S1: Upper and lower bounds of metabolite concentrations and cofactor ratios.**

| Metabolites, cofacer ratios | Lower bounds | Upper bounds | References |
|---|---|---|---|
| Intracellular glucose[*1] | $1.0 \times 10^{-3}$ M | $1.5 \times 10^{-2}$ M | (*81*) |
| Hepatic Pi | $3.0 \times 10^{-3}$ M | $4.3 \times 10^{-3}$ M | (*82*) |
| Myocytic Pi | $3.0 \times 10^{-3}$ M | $9.0 \times 10^{-3}$ M | (*82, 83*) |
| $CO_2$, $HCO_3^-$ | $1.5 \times 10^{-3}$ M | $2.5 \times 10^{-3}$ M | (*84*) |
| $NH_3$ | $2.0 \times 10^{-4}$ M | $1.0 \times 10^{-3}$ M | (*85*) |
| Other intracellular metabolites[*1] | $1.0 \times 10^{-7}$ M | $1.0 \times 10^{-2}$ M | (*26*) |
| Blood metabolites | $1.0 \times 10^{-7}$ M | $1.0 \times 10^{-1}$ M | (*86*) |
| ATP/ADP | 1 | 10 | (*83, 87, 88*) |
| $NAD^+$/NADH | 3 | 10 | (*89*) |
| GTP/GDP | 1 | 10 | (*90*) |
| $NADP^+$/NADPH | 0.05 | 0.2 | (*91*) |

1198 *1: The upper bounds of intracellular glucose, G6P, F6P, F1,6BP, DHAP, GAP, 3PG, 2PG,

1199 PEP, G1P, G3P, and Lac in *ob*/*ob* mouse are $1.0 \times 10^{-1}$ M.

1200

1201 **Table S2: Metabolite concentrations and a cofactor ratio involved in PKLR between**

1202 **mouse liver and other cell types.**

| Metabolites, cofacer ratios | Mouse liver | iBMK cell | Yeast | *E. coli* |
|---|---|---|---|---|
| PEP | $2.02 \times 10^{-5}$ M | $1.16 \times 10^{-5}$ M | $2.91 \times 10^{-5}$ M | $1.84 \times 10^{-4}$ M |
| Pyr | $3.77 \times 10^{-5}$ M | $5.88 \times 10^{-3}$ M | $9.40 \times 10^{-3}$ M | $3.66 \times 10^{-3}$ M |
| ATP/ADP | 1.00 | 8.21 | 3.95 | 17.35 |

1203 Metabolite concentrations in iBMK cell, Yeast, and E. coli were measured by LC-MS (*86*).

1204

**Supplementary data file legends**

**Data File S1: Metabolomic and $\Delta_f G'^\circ$ data used as input for GLEAM.**

This file includes the input metabolomic and $\Delta_f G'^\circ$ data for GLEAM (logarithmic mean of

metabolome concentrations, $\Delta_f G'^\circ$ values, and variance-covariance matrix of metabolite

concentrations and $\Delta_f G'^\circ$s). The metabolite concentrations were measured form from WT

and *ob*/*ob* mice in previous studies (*8, 69*), and $\Delta_f G'^\circ$ data were retrieved from eQuilibrator

database (*53*). NaN represents unmeasured metabolite. [h], [m], and [b] indicate hepatic,

myocytic, and blood compartment, respectively. The logarithmic mean of metabolite

concentrations shown in this file is converted from a logarithmic scale to a linear scale.

1214

1215 **Data File S2: Thermodynamically consistent metabolite concentrations, $\Delta_f G'^\circ$, and**

1216 **$\Delta_r G'$ estimated by GLEAM.**

1217 This file includes the thermodynamically consistent metabolite concentrations, $\Delta_f G'^\circ$, and

1218 $\Delta_r G'$ values in mouse liver and blood estimated by GLEAM. For glucose and lactate, hepatic

1219 and blood metabolites are distinguished by the [h] and [b], respectively.

1220

**Data File S3: Metabolites and reactions in the metabolic network for glucose**

**metabolism in mice.**

This file includes definitions of metabolite abbreviations, reaction abbreviations, and the

metabolic network *i.e.* the stoichiometric matrix for glucose metabolism used in this study.

In the Reactions and Stoichiometric matrix sheet, [h], [m], and [b] indicate hepatic, myocytic,

and blood compartments, respectively.

1227

**Data File S4: EC calculation in mouse liver and ECM.**

This file includes EC values within glycolysis, gluconeogenesis, and the TCA cycle in mouse

liver and EC, metabolite concentrations in ECM, and geometric mean of Michaelis-Menten

constant retrieved from BRENDA database (*51*) used for EC calculation. In EC calculation,

metabolite concentrations at *ad libitum* feeding were used for glycolysis and the TCA cycle,

and those after 16-hour fasting were used for gluconeogenesis and the TCA cycle. EC is

product of normalized molecular weight, $1/\eta_{rev}$, and $1/\eta_{sat,s}$.

1235

**Data File S5: Metabolite concentrations where the reactions required to reverse for**

**metabolic switching are at equilibrium, as determined by MEG calculation.**

This file includes the metabolite concentrations that are the optimal solution to the

optimization problem in MEG calculation for glycolysis, gluconeogenesis, and the TCA

cycle. At these concentrations, the reactions required to reverse for directional switching of

the pathways are at equilibrium with the minimum concentration changes from mouse liver

or ECM.

1243

**Supplementary references**

81. P. A. Wals, J. Katz, A concentration gradient of glucose from liver to plasma. *Metabolism* **42**, 1492–1496 (1993).

82. U. Till, D. Brox, H. Frunder, "Orthophosphate Turnover in the Extracellular and Intracellular Space of Mouse Liver" (1969); https://febs.onlinelibrary.wiley.com/doi/full/10.1111/j.1432-1033.1969.tb00806.x.

83. R. L. Veech, J. W. Lawson, N. W. Cornell, H. A. Krebs, Cytosolic phosphorylation potential. *J. Biol. Chem.* **254**, 6538–6547 (1979).

84. D. A. Beard, H. Qian, Thermodynamic-based computational profiling of cellular regulatory control in hepatocyte metabolism. *American Journal of Physiology-Endocrinology and Metabolism* **288**, E633–E644 (2005).

85. A. J. Meijer, W. H. Lamers, R. A. Chamuleau, Nitrogen metabolism and ornithine cycle function. *Physiol. Rev.* **70**, 701–748 (1990).

86. H. Ando, K. Ushijima, S. Shimba, A. Fujimura, Daily Fasting Blood Glucose Rhythm in Male Mice: A Role of the Circadian Clock in the Liver. *Endocrinology* **157**, 463–469 (2016).

87. S. Mörikofer-Zwez, P. Walter, Binding of ADP to rat liver cytosolic proteins and its influence on the ratio of free ATP/free ADP. *Biochem. J* **259**, 117–124 (1989).

88. T. W. Traut, Physiological concentrations of purines and pyrimidines. *Mol. Cell. Biochem.* **140**, 1–22 (1994).

89. S.-J. Lin, L. Guarente, Nicotinamide adenine dinucleotide, a metabolic regulator of transcription, longevity and disease. *Curr. Opin. Cell Biol.* **15**, 241–246 (2003).

90. J. L. Bos, H. Rehmann, A. Wittinghofer, GEFs and GAPs: Critical Elements in the Control of Small G Proteins. *Cell* **129**, 865–877 (2007).

91. R. L. Veech, L. V. Eggleston, H. A. Krebs, The redox state of free nicotinamide–adenine dinucleotide phosphate in the cytoplasm of rat liver. *Biochem. J* **115**, 609–619 (1969).