High-throughput screening reveals mechanisms of environmental control of germination in a fungal thermophile

Olusola A. Ogunyewo[1], Kristopher Fleming[1], Madeleine Morris[1], Kayleigh Fort[1], Lori B. Huberman*[2], Rachel B. Brem*[1]

[1]Department of Plant and Microbial Biology, University of California, Berkeley, United States
[2]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, NY, United States

*Corresponding authors: rbrem@berkeley.edu, huberman@cornell.edu

Abstract

Thermothelomyces thermophilus is a filamentous fungus isolated from self-heating compost. Unlike most of the fungal kingdom, this species exhibits a growth optimum at 45°C and is intolerant of temperatures below 30°C. To dissect the mechanisms of thermophilicity in this system, we implemented a large-scale insertional mutagenesis approach. We first made thousands of *T. thermophilus* mutants and cultured them at different temperature extremes in standard medium. Phenotyping-by-sequencing identified dozens of disrupted loci that represented candidate determinants of thermophilicity, including several annotated in metal transport. We then validated a subset of screen hits with a directed, single-gene knockout paradigm. The results revealed a temperature-dependent regulatory logic for germination, the developmental decision by which a fungal spore initiates growth under replete conditions. Surprisingly, most of our *T. thermophilus* mutants germinated far better at 50°C than the wild-type in standard medium and showed markedly slower germination at lower temperatures. We hypothesized that *T. thermophilus* has evolved sophisticated regulatory machinery to block germination at high temperature unless environmental conditions are favorable. As a proof of concept, we surveyed media conditions and established that elevated zinc dampened germination of wild-type *T. thermophilus* at 50°C but promoted it at lower temperature; mutation experiments made clear that such sensitivity was mediated in part by the zinc transporter *zip*. We interpret these results under a model in which *T. thermophilus* integrates temperature and nutrient availability to control the transition from spore dormancy to vegetative growth.

Significance

Fungal thermophiles thrive at temperatures that represent the upper limits of eukaryotic life. The molecular mechanisms that underlie their unique capacities remain poorly understood. In this work, we elucidate how *Thermothelomyces thermophilus* integrates temperature cues with other environmental inputs during germination, a key life-cycle stage for dispersal. These insights are of basic-biological interest and provide a foundation for rational strategies to boost thermotolerance in industrial strains, with applications for high-temperature bioprocessing.

Key Words

BarSeq / TnSeq, high-throughput screening, temperature response, *Thermothelomyces thermophilus*, germination, insertional mutagenesis, zinc signaling

Introduction

Thermophilic fungi occupy the upper temperature range of eukaryotic life and operate under a physiological ceiling approaching ~60°C (1). Microbial ecology surveys have outlined the narrow diversity of truly thermophilic fungal species adapted to composts and geothermal soils (2, 3). In thermophilic bacteria, elevated growth temperatures have been linked to genome streamlining, higher GC content, thermostable protein folds, membrane lipid saturation, and enhanced chaperone and DNA-repair capacities (4, 5). Whether similar mechanisms govern thermophily in fungi remains unclear, particularly given the added complexity of organelles, chromatin regulation, and developmental programs in these systems. Thus, against the backdrop of years

62 of work on mesophile and thermotolerant fungi (6–9), the mechanisms of high-temperature
63 growth preference in fungal thermophiles are still incompletely understood.
64
Beyond their fundamental biological interest, thermophilic fungi also have practical relevance as
potential hosts for industrial bioprocessing. Microbial cell factories are widely engineered to
produce fuels, pharmaceuticals, and other industrially valuable chemicals; however, the cost of
cultivation substrates remains a major constraint to sustainable biomanufacturing (10, 11).
Fungi represent a potential solution toward this end, when they have the capacity to digest
abundant, low-cost crop wastes like rice straw and corn stover. For such substrates, industrial
processes commonly apply high-temperature pretreatments to accelerate biomass
deconstruction. In this pipeline, subsequent cooling prior to microbial inoculation adds cost and
energy demand (12). Thermophilic fungi therefore represent a promising chassis for
bioproduction, as they could bridge pretreatment and fermentation steps without the need for
cooling, directly converting pretreated biomass into target products.
76
Among thermophilic fungi, *Thermothelomyces thermophilus* (formerly *Myceliophthora
thermophila*) has emerged as a compelling model with robust growth near 50°C and a
carbohydrate active enzyme repertoire capable of plant cell-wall deconstruction, sparking
interest in its use as a thermophilic cell factory (13, 14). Genetic tools in this system now enable
precise engineering (15), including the potential to optimize high-temperature performance(16–
18). We set out to use *T. thermophilus* as a testbed for the dissection of mechanisms of fungal
thermophily at a genomic scale. Our approach was to pioneer the application of a recently
developed method for randomly barcoded insertional mutagenesis with screening via
sequencing in filamentous fungi (19) for the *T. thermophilus* system. We sought to use this
strategy for an unbiased genetic survey of temperature-responsive growth behaviors in *T.
thermophilus*. We used the results as a jumping-off point for exploration of the temperature
dependence of the asexual stages of the filamentous fungal life cycle and the underlying
genetics. The results reveal how a thermophilic eukaryote coordinates stress signaling and
development at high temperature.
91

Results

Temperature-dependent growth and development of *T. thermophilus*

With the ultimate goal of understanding mechanisms of thermophily in *T. thermophilus*, we first
surveyed growth behaviors of *T. thermophilus* across temperatures and conditions. On solid
agar, we documented the asexual growth cycle, from hyphal expansion through formation of
aerial hyphae to production of spores (conidia) with visible pigmentation (Figure 1A). We
observed a temperature-responsive lag in development as temperature increased, even with
high-percentage agar and high-humidity incubation to maintain plate integrity (Figure 1A-B). As
a complement, we also assayed growth in liquid culture, in which we inoculated *T. thermophilus*
conidia into standard liquid medium and quantified the resulting mycelial biomass accumulation
after incubation with shaking. The results revealed evidence for *bona fide* thermophilicity by *T.
thermophilus*, with an optimum at 45°C during liquid growth and a gradual falloff of performance
under warmer and cooler conditions (Figure 1C). We thus considered liquid culture as a growth
format suitable for genetic analyses to reveal mechanisms of high-temperature adaptation in *T.
thermophilus*.

Construction and mapping of a *T. thermophilus* insertional mutant library

To develop a genomic resource for mapping of genotype to phenotype in *T. thermophilus*, we constructed a pool of barcoded insertional mutants. Our process used random insertion from *Agrobacterium tumefaciens* harboring hygromycin-marked, barcoded transfer DNA (T-DNA) (19), with which we carried out ectopic transformation of mononucleate *T. thermophilus* conidia via established procedures (20) with minor modifications to conidial density, co-cultivation duration, and selection conditions, followed by expansion and aliquoting of replicate sub-pools (Figure 2A). To catalog the diversity and chromosomal distribution of T-DNA insertions across the mutants, we isolated DNA from replicate cultures of the pool grown in liquid medium at 37°C and prepared sequencing libraries enriched for T-DNA–genome junctions. *In silico* analysis of reads mapped to the *T. thermophilus* reference genome identified the T-DNA right-arm junction, extracted adjacent genomic sequence, and assigned uniquely mapped barcodes to genomic coordinates. From these data, we detected 17,155 distinct insertion sites distributed uniformly across all seven chromosomes (Figure 2B). Of these, 3,426 insertions were located within annotated genes (Supplementary File 1). These data established our pool as a partial-coverage mutant collection sufficient to pursue genetic principles via quantitative fitness profiling in *T. thermophilus*.

Genome-wide quantification of temperature-dependent mutant fitness

We expected that screens with our insertional mutant pool could enable dissection of temperature response in *T. thermophilus*. To this end, we sought to use the resource to identify genes whose mutation effect on fitness was temperature-dependent. We designed an experimental scheme using growth assays of our mutant pool at 37°C and 50°C, representing temperatures that approached the lower and upper extremes, respectively, of wild-type temperature tolerance in liquid culture (Figure 1C). For each biological replicate, we cultured independent aliquots of the *T. thermophilus* mutant pool in parallel at 37°C and 50°C, and we quantified mutant abundance by sequencing the barcode regions of the inserted T-DNA from the resulting genomic DNA (Figure 3A). In the resulting abundance data, reproducibility between biological replicates was high across both temperatures (Pearson r > 0.994; Figure 3B and Supplementary Figure S1). We used the complete data set as input into a quality control and analysis pipeline (see Methods) that assessed differences between temperatures in mutant abundance on a per-gene basis, finding 783 mutants at adjusted *p*-value < 0.05 and absolute $\log_2$(fold-change) ≥ 1 (Supplementary File 2). Though these genes spanned a range of annotated functions (Supplementary File 2), their behavior in the screen was remarkably consistent: for all but six gene hits, mutant abundance was higher at 50°C than at 37°C (Figure 3C and Supplementary File 2). These data attested to a highly complex genetic architecture of temperature response in *T. thermophilus,* whose contributors, when mutated, almost invariably conferred defects in cooler conditions and/or benefits at high temperature.

Single-gene validation of temperature-dependent screen hits

To validate the inferences from our genome-wide random mutagenesis screen, we took a single-gene approach. For this purpose we chose eight screen hits (Figure 3C and Table 1) with high coverage in the mutant pool and robust mutant effects favoring either 50°C or 37°C. These included putative regulators (*zincTF, syg1*, *pty-5*, and *nrc-1*), transporters (*zip* and *mfs*), and housekeeping genes (*ech* and *MYCTH_2313252*) (Table 1). We generated targeted deletion mutants for each by homologous recombination of a hygromycin‑resistance cassette at the endogenous *T. thermophilus* locus in a strain harboring a deletion in *ku80*, to ensure efficient deletion by eliminating non-homologous end joining. Correct integration events were confirmed by PCR and sequencing (Supplementary Figure S2).

In designing experiments to characterize these single-gene mutant strains, we reasoned that temperature could influence multiple stages of the *T. thermophilus* life cycle, and that distinguishing between such effects would maximize our insights into the mechanisms of temperature response. We elected to focus on the phases relevant for the growth scheme of the high-throughput screen (Figure 3), namely germination of conidia and the extension of hyphae after vegetative growth had established. To this end, we established an assay to quantify temperature impacts on conidial germination of a given strain by microscopy, at a relatively early timepoint where germlings could easily be resolved from one another in an imaging field (Figure 4A). Separately, we designed an assay of temperature-dependent vegetative hyphal extension for a given strain, avoiding confounding effects from germination by using a standardized pre-germinated inoculum (Figure 4B). We applied both assays to each mutant and to their isogenic parent, the otherwise wild-type Δ*ku*80 strain.

From germination measurements across this strain set, we observed a modest drop in performance at elevated temperature by the *Δku80* parent strain, which we used as a baseline for comparison of the remainder of the single-gene mutants (Figure 4A). Among these strains, 6 of 8 genotypes exhibited a marked advantage at 50°C, germinating up to 2-fold more avidly than the parent; by contrast, at 37°C, 6 of 8 mutants exhibited reduced germination relative to the parent (Figure 4A). In these validated cases, mutations at a given locus (*ech, mfs, nrc-1, pty-5, syg1, zip*) most often exerted effects in opposite directions at the two temperatures (Figure 4A), attesting to dual roles of a given gene with respect to germination at the two extremes. Their trends mirrored results from our large-scale screen, in which fitness phenotypes of mutants made by random integration almost universally favored growth at 50°C (Figure 3B). Interestingly, in three exceptions to the latter (*mfs*, *ech,* and *zip*), for which mutant abundance in the screen was highest at 37°C (Figure 3B and Table 1), our single-gene validation experiments nonetheless established that the respective mutants germinated preferentially at 50°C. This observation, and the fact that a single-gene deletion of *MYCTH_2313252* conferred a germination defect at the lower temperature (Figure 4A) but random insertional mutants at this locus in the screen were highly abundant at 37°C relative to 50°C (Figure 3B and Table 1), suggested cases in which performance in the growth scheme of the screen reflected contributions from behaviors other than those interrogated in our germination assay. Apart from Δ*MYCTH_2313252*, however, the most salient and pervasive pattern from our observations of germination in single-gene mutants was that of improvements at high temperature (Figure 4A). We thus formulated a model in which the wild-type alleles of many loci function as repressors of germination at high temperature, such that their deletion would release this brake and allow development to proceed despite heat stress; the same loci would act as activators of germination under cooler conditions.

We next turned to assays of biomass accumulation in the hyphal growth phase of *T. thermophilus*, inoculating with pre-germinated mycelium at an equal starting concentration for all strains, including the Δ*ku80* parent (see Methods). For wild-type *T. thermophilus*, high temperature retarded growth in this scenario, with 1.5-fold lower biomass accumulation at 50°C than at 37°C (Figure 4B). Most single-gene mutants phenocopied this behavior almost exactly, such that we identified few cases in which a mutant grew differently from the wild-type parent at either temperature (Figure 4B). Among these exceptions, we noted an advantage of the *nrc-1* deletion mutant relative to wild-type at 37°C; defects at 50°C in the Δ*pty-5* and Δ*ech* strains (representing a plausible mechanism for increased abundance of mutants at these loci at 37°C in our original large-scale screen [Figure 3B and Table 1]); and increased growth by Δ*syg-1* at 50°C (Figure 4B). But as the latter represented outliers across the gene set, we concluded that growth was not a primary life cycle stage for the roles of most of our focal genes in temperature response, particularly in light of the striking impact of their deletions during germination (Figure

214 4A). Given the origin of our focal genes as representative of the hundreds of hits from our large-
215 scale screen (Figure 3B), we infer that the overall trends in our screen results were also likely
216 dominated by temperature-dependent effects of gene disruption on germination.
217
218
Zinc availability modulates germination efficiency

Is there a regulatory logic that governs repression of germination in wild-type *T. thermophilus* at
high temperature? We hypothesized that the fungus integrates multiple environmental inputs,
including temperature and nutrient availability, in the decision to germinate. As a test case, we
considered environmental metal exposure, in light of the validated role in temperature-
dependent germination of the putative zinc-iron permease *zip* (Figure 4A). We formulated a
model in which zinc availability acts to gate *T. thermophilus* germination under heat stress. To
explore this, we performed a zinc titration series for wild-type *T. thermophilus*, assaying
germination as a function of zinc concentration and temperature. The results revealed a dose-
responsive benefit in germination from increasing zinc at 37°C and the inverse relationship at
50°C, with germination falling off markedly as zinc concentrations increased (Figure 5A). At the
lowest zinc concentrations, germination at high temperature far outstripped that observed at
37°C (Figure 5A). Such a pattern dovetailed with the benefit to germination at 50°C that we had
observed in standard medium (5 mg/L zinc) upon deletion of *zip* (Figure 4A), as expected if zinc
uptake were limited in the Δ*zip* mutant. To test the role of *zip* directly, we repeated germination
assays in the Δ*zip* strain and found no impact of zinc concentration on germination efficiency in
this background (Figure 5B). Rather, germination remained higher at 50°C than at 37°C across
zinc conditions in the absence of *zip* (Figure 5B). Together, these data demonstrate that zinc is
a key environmental input into developmental decision-making in *T. thermophilus*, and that zinc-
dependent modulation of germination requires *zip*.

Discussion

240
Thermophilic fungi are a powerful model for the basic biology of how eukaryotes adapt to
extreme environments, and they also are of interest in industrial applications. In this work, we
have established insertional mutagenesis as a tool for mechanistic dissection of temperature
response in *T. thermophilus.* High-throughput forward genetics in filamentous fungi is in its
infancy (19), and our method is well-positioned to accelerate advances in the *T. thermophilus*
system, as has now become standard in bacteria and yeasts (21–25). The principles that have
emerged from our study break new ground in the understanding of fungal thermophilicity.
248
We have shown that hundreds of genes, when disrupted in our high-throughput screening
approach, affect high-temperature performance by *T. thermophilus.* Among these, our validation
of focal gene hits has found that most act as negative regulators of conidial germination at high
temperature, in that they improve germination when deleted. The capacity to make spores
represents a mechanism for stress resistance common to essentially all fungi, and exit from this
dormant state is tightly regulated even in mesophiles (26–28), to ensure that a new colony only
commits to growth under favorable conditions. It is tempting to speculate that for thermophiles,
evolution has optimized the dormancy-breaking decision in unique ways, taking into account the
cost or toxicity of growth programs at extreme temperatures. If so, the logic in *T. thermophilus*
would be to sense environmental signals associated with the worst risk at high temperature and
use them to trigger a block on germination.
260

Our screen and validation data raise the possibility that zinc processing represents one such
environmental risk factor. If cellular functions involving zinc were particularly deleterious during
T. thermophilus germination under high-heat conditions, it would mirror previous studies in
mesophilic fungi in which metal exposure compromises spore germination more than it does
hyphal growth (29). Plausibly, such effects could be mediated by reactive oxygen species, which
have been associated with both zinc (30, 31) and elevated temperature (32, 33) in other
systems. Formally, however, our data leave open the question of whether and how zinc exerts
damaging effects *per se* in *T. thermophilus*. What our experiments have shown is that levels of
zinc that shut down germination at high temperature fostered the best germination at 37°C. This
pattern is consistent with a regulated delay of germination under combined zinc and heat stress,
rather than solely a passive consequence of zinc-induced damage. Furthermore, although our
validation experiments have focused on zinc, we expect it to be only one of many environmental
inputs that *T. thermophilus* integrates with temperature when regulating life-cycle transitions.
Under this model, the annotated regulator genes whose deletion boosts germination at high
temperature represent candidate nodes of a complex signal-integration network controlling the
T. thermophilus life cycle. Such a function could reflect repurposing from the known roles, in
mesophiles, of our focal genes *pty-5* (a tyrosine phosphatase) and *nrc-1* (a MAP3 kinase) in
conidiation and germling fusion respectively (34, 35); the function of another of our focal factors,
syg-1, likely in phosphate-based signaling (36), is unknown even in model fungi. For the
annotated effector genes we have highlighted here—*ech* (encoding enoyl-CoA hydratase, a β-
oxidation enzyme whose disruption alters growth and development in other fungi) and the
zinc/iron permease gene *zip* (37, 38)—we interpret their mutant effects as eliminating repressive
environmental inputs farther upstream, allowing germination to proceed even under heat stress.
284
Meanwhile, although we have emphasized relief of germination repression in *T. thermophilus*
mutants, our work has also uncovered genes for which we infer positive effects on mycelial
growth (focal genes *ech* and *pty-5*): that is, their deletion compromises this phase of the life
cycle at high temperature. We anticipate that many more such pro-growth factors remain to be
identified in *T. thermophilus*. Indeed, it is reasonable to assume that principles from thermophilic
bacteria and archaea that lack multiple life stages, in which evolution has adjusted
housekeeping genes all over the genome to promote growth at high temperature (39), are likely
to bear out in fungi. Our screening approach for *T. thermophilus* may have been particularly
likely to miss pro-growth loci of this kind if they are essential, in that their disruption mutants
would fail to propagate in our stocks at any temperature and would be lost from our analysis
pipeline. As such, and in light of the partial coverage of our mutant pool, our screen results
represent only a first window onto the complete suite of genes that mediate thermophilicity in
this system. As they stand, however, our findings underscore the importance of germination as a
nexus of regulatory decisions for *T. thermophilus*, integrating temperature with other
environmental inputs—which could well include nutrient availability, pH, or osmotic or oxidative
stress, as well as metal homeostasis—to optimize fitness.
301
302 ## Conclusion
303
We have used an insertional mutagenesis screen and validation follow-up in *Thermothelomyces*
thermophilus to uncover principles of thermophilicity in fungi. Our results shed light for the first
time on the regulatory landscape of decisions by these organisms about whether to germinate in
the heat. And in an industrial setting, the deletion mutants we study here represent compelling
candidate cases in which genetic manipulations could improve performance by *T. thermophilus*
even at the extreme of its heat tolerance.
310

Methods

Strains, media, and growth conditions

Thermothelomyces thermophilus ATCC 42464 was obtained from the American Type Culture Collection. The Δ*ku80* strain MJK20.2 was generously provided by Vera Meyer (40, 41). The strains were maintained on Vogel's minimal medium (VMM) (42) supplemented with 2% sucrose. For solid cultures, VMM with the indicated carbon source was solidified with 3% (w/v) agar. Liquid cultures were grown in 250 mL Erlenmeyer flasks at 200 rpm with 50 mL culture volume. Unless otherwise stated, cultures were incubated at 37°C, 45°C, or 50°C in a closed, dark incubator using the New Brunswick Innova 40 incubator. Conidia were harvested from 7-day-old plates by gentle flooding with sterile 0.01% (v/v) Tween-80, filtered through sterile Miracloth to remove hyphal fragments, counted using a hemocytometer, and used immediately or stored at -80°C until use. All other strains and plasmids generated from the study are listed in Table 2.

Temperature-dependent growth and development assays of wild-type *T. thermophilus*

To characterize thermal performance, conidia were inoculated onto solid VMM + 2% sucrose plates at a final density of $1.2{\times}10^6$ conidia/spot and incubated at 37°C, 45°C, or 50°C for 168 h in a closed incubator under standard dark conditions. Growth was observed daily using a flatbed scanner. Colony pigmentation and aerial hyphae formation were documented under standardized lighting conditions. Conidiation output was quantified from cultures grown on solid medium for 168 h. Conidia were harvested as described above and counted using a hemocytometer in triplicate biological replicates per temperature. For liquid growth assays, conidia were inoculated at $1{\times}10^\square$ conidia/mL into 50 mL VMM + 2% sucrose in 250 mL Erlenmeyer flasks and incubated at 30°C, 37°C, 45°C, 50°C, or 55°C for 24 h at 200 rpm under standard dark conditions. Biomass accumulation was quantified by dry weight after filtration onto pre-weighed Whatman GF/A filters and dried at 80°C until a constant mass was achieved.

Construction and Tnseq of the insertional mutant pool

We generated genome-wide random insertional mutant libraries in *Thermothelomyces thermophilus* ATCC 42464 using *Agrobacterium tumefaciens*–mediated transformation. To obtain fresh starting material, we prepared conidia on VMM containing 2% sucrose and 2% agar and incubated the plates at 37°C for 5–7 days in the dark. Conidia were harvested in sterile water, filtered through Miracloth to remove hyphal debris, counted with a hemocytometer, and adjusted to ~$4{\times}10^7$ conidia/mL. For *Agrobacteriu*m transformation, we used *the A. tumefaciens* strain EHA105, which carries a binary vector encoding a barcoded T-DNA cassette and a hygromycin resistance marker (19). The strain was cultured in low-salt LB with kanamycin in an exponential phase (OD_{600} ~0.5–0.8), then transferred to induction medium (IM) containing 19.5 g/L 2-(N-morpholino)ethane sulfonic acid (MES) at pH 5.5 supplemented with 500 μM acetosyringone for 18 h to induce virulence following the modified method of Michielse *et. al.* (43). Grown *A. tumefaciens* (OD_{600} ~0.3–0.7) was mixed with *T. thermophilus* conidia at a 1:1.5 ratio and co-incubated for 1 h at room temperature with gentle mixing to prevent settling.

We spread 1 mL of the co-culture mixture onto the surface of a sterile ultra clear cellophane (#RPI-1090) placed on a 150 × 15 mm IM agar plate supplemented with 500 μM acetosyringone. Plates were incubated at 28°C for 96 h before transferring the cellophane overlays to VMM + 2% sucrose selection plates containing 200 μg/mL hygromycin and 100 μg/mL carbenicillin.

Plates were incubated at 37°C until hygromycin-resistant colonies emerged (4–7 days), and
conidiation was allowed to proceed for an additional ~14 days. This full workflow was performed
twice, generating two independent mutant pools: Batch 1 (~4,000 transformants) and Batch 2
(~13,000 transformants). For each batch, conidia from all selection plates were harvested,
pooled, filtered through Miracloth, and stored as 10% glycerol stocks at –80°C. Together, these
two batches comprised ~20,000 unique insertion mutants.

For mapping T-DNA insertion sites (Tnseq), for each batch we inoculated pooled mutant conidia
(1×10□ conidia/mL) into 100 mL VMM + 2% sucrose and grew the culture for 24 h at 37^{o}C and
200 rpm in the dark before extracting genomic DNA using the Quick-DNA Fungal/Bacterial
Miniprep Kit (Zymo Research). No dropout mix was used at any stage, as *T. thermophilus* is not
maintained as an auxotrophic system; transformants were selected solely on hygromycin-
containing medium, and only disruptions in essential genes would be expected to drop out of
the pooled library. We verified DNA integrity by agarose gel electrophoresis and quantified with
a Qubit dsDNA HS assay. We prepared T-DNA genome junction sequencing libraries following
the general TnSeq workflow of Wetmore et al. (2), but using an enzymatic-fragmentation
workflow based on the NEBNext Ultra II FS DNA Library Prep Kit with Sample Purification
Beads for Illumina (New England Biolabs #E6177S). Library preparation followed the
manufacturer's instructions with modifications appropriate for TnSeq-style T-DNA junction
enrichment as described previously (44). Briefly, FS-based fragmentation and end-repair,
adaptor ligation, and dual-SPRI size selection were performed using components of the
NEBNext kit, after which nested PCR enrichment of junction fragments was carried out using
the Nspacer_barseq_universal and P7_MOD_TS_index primers (sequences in Supplementary
File 4). Final libraries were sequenced as paired-end 150 bp reads on an Illumina MiSeq at the
UC Berkeley Vincent J. Coates Genomics Sequencing Laboratory.

Tnseq analysis/mapping of the insertional mutant pool

To determine the genomic coordinates of each barcoded insertion, sequencing reads from
Tnseq from Batch 1 and Batch 2 (see above) were combined and then processed using a
modified version of the pipeline described by Kim et al. (45). All custom modifications and
scripts used in this study are available in the accompanying GitHub repository
(https://github.com/oaogunyewo-git/Thermothelomyces-Tnseq-
project/tree/main/TnSeq_BarSeq_sequencings_notebooks_and_scripts) and permanently
archived on Zenodo (46). Briefly, the algorithm first identified reads containing the T-DNA right-
arm junction sequence, allowing up to two mismatches to accommodate sequencing error, and
extracted the adjacent genomic flanking sequence. These flanking sequences were aligned to
the *T. thermophilus* reference genome (assembly MycTherm_2019, available at the DOE JGI
MycoCosm portal) (13) using BLASTn with a minimum alignment identity of 95% and an E-value
threshold of 0.1. A barcode was considered to map uniquely only if the highest-scoring genome
alignment exceeded the second-best alignment by at least 10 bitscore units. Barcodes differing
by single-base sequencing errors were collapsed, and barcodes mapping to nearly identical
positions (<10 bp apart) were resolved by retaining the most abundantly supported locus.
Barcodes classified as multi-locus, ambiguous, or insert-derived were excluded. Only single-
locus barcodes supported by at least one uniquely mapped read were retained. Each insertion
was assigned a classification (single, multi-locus, concatameric, or ambiguous) based on
mapping structure, and only single-locus insertions were retained for downstream mutant fitness
calculations. The final poolfile reported for each barcode, the chromosomal coordinate, strand
orientation, gene annotation, and corresponding read abundance. The resulting poolfile was
then annotated by intersecting insertion coordinates with gene boundaries defined in the *T.*

thermophilus genome annotation (GFF) file, thereby assigning each barcode to a specific gene
(or intergenic region) and producing a final gene-level insertion index used for subsequent
fitness scoring.

High-throughput phenotyping and quantification of temperature-dependent fitness in the
insertional mutant pool

For a given replicate of quantification of abundance of mutant across the pool of insertional
mutants (Barseq), we proceeded as follows. A 1 mL frozen aliquot of a given mutant pool (Batch
1 or Batch 2; see above) was spread onto two 150 x 15 mm VMM +2% sucrose agar plates
(500 uL per plate) and incubated at 37^{o}C for 6 days in the dark to allow full conidiation. Conidia
harvested from these expanded plate cultures were pooled and split into two inocula for growth:
50 mL liquid VMM + 2% sucrose was inoculated with 1×10□ conidia/mL in each case, after
which one was incubated at 37°C and the other at 50°C for 24 h at 200 rpm under dark
conditions. For each, mycelial biomass was harvested for genomic DNA extraction using Quick-
DNA Fungal/Bacterial Miniprep Kit (Zymo Research D6005). Unique barcode regions were
amplified using Q5 High-Fidelity DNA Polymerase with GC Enhancer (NEB M0491S) and
primers targeting constant priming sites flanking each barcode, yielding a ~185 bp Illumina-
ready amplicon (47). Amplicons were purified using Zymo DNA Clean & Concentrator (D4014),
quantified using a Qubit 3.0 fluorometer, pooled in equimolar ratios within each batch.
Replicates of Batch 1 (the ~4,000-mutant library) were sequenced on an Illumina MiSeq
platform, and replicates of Batch 2 (the ~13,000-mutant library) were sequenced on an Illumina
NextSeq P2 platform (100-cycle single-end). When necessary, libraries were size-selected
using a Pippin Prep (Sage Biosciences) and spiked with 15% PhiX to increase base diversity.

Sequencing reads were processed using a modified RBseq_Count_BarCodes pipeline (46).
Barcode counts were normalized by total-count scaling and linked to genomic insertion sites via
the annotated poolfile; gene-level abundance estimates were derived from uniquely mapped,
single-locus barcodes only. As an internal quality control, genome-wide insertion density was
summarized in 50-kb chromosomal bins, and replicate concordance was assessed by pairwise
scatter plotting of normalized barcode abundances with coefficients of determination (R^2) from
ordinary least-squares regression. Barcodes whose insertion position from Tnseq mapped to
multiple promoters or genes were eliminated from downstream analysis, as was any gene for
which all insertion mutants in all replicates had fewer than 5 observed reads.

To test for differential abundance between temperatures of the mutants of a given gene, we
proceeded as follows. We calculated the mean normalized abundance across all insertional
mutants from all 37°C replicates and divided the value of the normalized abundance from each
insertional mutant in each 50°C replicate by this value. $\log_2$ transformation of these ratios
yielded per-gene fold-change estimates. For this baseline-normalized analysis, statistical testing
was restricted to genes represented by ≤15 uniquely mapped barcodes, to prevent genes with
very high insertion density from disproportionately influencing variance structure. For each such
gene, a one-sample Wilcoxon signed-rank test was used to determine whether the median
$\log_2$(50°C/37°C) differed from 0 (i.e., ratio = 1). P-values were corrected using the Benjamini–
Hochberg procedure, and genes were designated as temperature-responsive when the absolute
value of the mean $\log_2$(50°C/37°C) ≥ 1 and FDR < 0.05. We also computed the standard
deviation and coefficient of variation (CV%) across all available replicate $\log_2$(50°C/37°C)
measurements.

Single-gene validation of screen hits

 Construction of single-gene deletion mutants

Single-gene knockout mutants were generated in the Δ*ku80* background of *T. thermophilus* to
promote homologous recombination (48). For each target gene, we PCR-amplified the upstream
and downstream genomic flanking regions, typically 500 bp to 1 kb each, depending on the
locus. We assembled these flanking regions on either side of the hygromycin resistance marker
(hph), which was amplified from the pLH159RB plasmid. The flanking regions and hph cassette
were inserted into the pCAMBIA1302 binary vector, which had been linearized by *Xho*I and
*BssH*II digestion to remove the native GFP cassette and create compatible ends for assembly.
Constructs were assembled using NEBuilder HiFi DNA Assembly Master Mix (New England
Biolabs, #E5520S). The assembled plasmids were transformed into *Escherichia coli* DH5α, and
transformants were selected on LB agar supplemented with 50 µg/mL kanamycin. We screened
positive colonies by colony PCR, and plasmid DNA was purified using a miniprep kit (QIAprep
Spin Miniprep Kit #27106). Correct assembly was verified by diagnostic restriction enzyme
digest and Sanger sequencing across the junction regions to confirm integrity of the flanking
regions and hph cassette. The complete list of plasmids generated in this study is provided in
Table 2, and all primers used for cassette construction are listed in Supplementary File 4.

The verified pCAMBIA1302-based deletion vectors were introduced into *Agrobacterium*
tumefaciens strain LBA4404 by electroporation. Transformants were recovered on low-salt LB
agar supplemented with 100 µg/mL kanamycin and 50 µg/mL rifampicin to select for plasmid
maintenance and the bacterial background, respectively. Gene deletion in *T. thermophilus* was
then carried out by Agrobacterium-mediated transformation following the protocol of Michielse *et*
al., 2008 (43). Briefly, freshly harvested conidia (~4×10☐ per plate) were co-cultivated with *A.*
tumefaciens on cellophane membrane placed directly on IM agar plates containing 500 µM
acetosyringone and incubated at 28°C for 96 h as above. After a 96 h incubation period, the
cellophane membrane containing the grown co-cultivation mixture was subsequently transferred
to the selection plates (VMM+ 2% sucrose supplemented with 200 µg/mL hygromycin and 100
µg/mL carbenicillin) and incubated at 37°C for 4–5 days to recover primary transformants.
Hygromycin-resistant colonies were then re-streaked to secondary selection plates containing
the same antibiotic and incubated for an additional 10 days to promote full conidial development
and ensure stable genomic integration prior to strain purification and downstream verification.

Molecular verification of single-gene validation mutants

To verify each gene deletion, we first grew conidiospores from the hygromycin-resistant
transformants in 50 mL VMM + 2% sucrose for 72 h at 37°C and 200 rpm in the dark and
extracted genomic DNA using the Quick-DNA Fungal/Bacterial Miniprep Kit (Zymo Research,
D6005). We confirmed the gene deletions using a two-step PCR strategy. First, we performed 5′
and 3′ verification PCRs using primer pairs located outside the upstream and downstream
flanking regions of the targeted locus. This allowed us to distinguish between the wild-type allele
and the deletion allele based on product size. For each mutant, we observed a single band
corresponding to the expected size of the deletion allele, which differed from the native allele
band amplified from the Δ*ku80* parental strain. Second, we used internal ORF-specific primers
to test for the presence or absence of the native gene. The deletion strains showed no internal
ORF product, while the Δku80 parental strain produced the expected amplicon, confirming loss
of the target coding region. PCRs were performed using Q5 High-Fidelity DNA Polymerase
(NEB, M0491S) with GC enhancer, and products were resolved on 1% agarose gels. Strains
that met both criteria—correct external PCR size and absence of the internal ORF—were
purified further by single-conidium isolation to ensure clonal homogeneity. For a subset of

mutants, we also sequenced the cassette–genome junctions to confirm precise integration and
rule out partial or ectopic events. Primer sequences used for verification are provided in
Supplementary File 4.

<u>Phenotypic validation of single-gene validation mutants</u>

For the liquid growth assay of single-gene validation mutants, because strains differed in
germination efficiency at 37°C and 50°C, we designed a scheme in which all strains began the
hyphal-growth phase with a similar biomass input, allowing us to assess post-germination
hyphal growth independently of initial germination rate. We achieved this by pre-germinating
each strain and replicate in turn at 45°C and ensuring consistent growth-ready inocula in liquid
culture, as follows. For a given replicate, conidia were first harvested from fully sporulated
cultures, washed once in sterile water, and counted using a hemocytometer. We next used this
to establish a pre-culture by inoculating 10 mL of the conidial suspension (~2 x 10^7
condiospores/mL) into 90 mL of VMM + 2% sucrose in a 500 mL Erlenmeyer flask, sealing with
foil, and pre-germinating for 24 h at 45°C with shaking at 200 rpm under standard dark
conditions. We then dispensed 5 mL pre-germinated culture (containing mycelium suspended in
medium) into each of three 250 mL Erlenmeyer flasks containing 45 mL fresh VMM + 2%
sucrose to establish technical replicate cultures. Replicate flasks were incubated in VMM + 2%
sucrose at 37°C or 50°C for 18 h with shaking at 200 rpm under dark conditions. At the end of
the incubation, mycelia were harvested by filtration through Miracloth, dried overnight at 50°C,
and weighed to obtain final dry biomass (mg). Following incubation, mycelia were collected by
filtration through Miracloth, washed with sterile water, and dried overnight at 50°C to constant
weight. The dry biomass (mg) was recorded for each culture. In a separate checking step for
each replicate, we also took an additional 5 mL of the pre-germinated culture (apart from the
inocula used for growth assays), dried it overnight at 50°C, and weighed it. The results from
these checks revealed little variation between replicates or genotypes (7.0 – 11.0 mg),
confirming that 5-mL aliquots used as inocula for growth would be consistent with respect to
growth-ready biomass. Biological replicates were performed with $n \geq 3$ per strain per condition.
For visualization in heatmaps, replicate measurements were aggregated as mean values. Full
replicate-level data are provided in Supplementary File 3.

Germination efficiency was assessed as follows. For a given replicate, 150 µL of a conidial
suspension (~2 x 10^7 condiospores/mL) was inoculated into 1.5 mL VMM + 2% sucrose in each
well of a 24-well plate (Cellvis #P24-1.5H-N). Plates were incubated at 37°C or 50°C with
shaking at 200 rpm for 10 h in the dark. To ensure uniform sampling, wells were briefly mixed by
pipetting with a P1000 immediately before microscopy. This mixing step resuspended settled
conidia and did not alter germination conditions. For microscopy, we withdrew 10 µL of the
culture from each well and mounted it on a microscope slide for imaging. Germination was
scored using a compound light microscope (National Ecoline D-ELDB Binocular Digital
Microscope) at 400× magnification under brightfield illumination. A conidium was defined as
germinated when the emerging germ tube exceeded the diameter of the spore body. For each
replicate, ≥200 conidia were counted manually to calculate percent germination. Each strain and
temperature condition was assayed in ≥ 3 biological replicates.

Assays of zinc effects on germination

Zinc-dependent effects on germination were assessed in liquid VMM supplemented with 2%
sucrose using *T. thermophilus* Δ*ku*80 or Δ*zip* strains, as indicated. Vogel's minimal medium was
prepared with graded $ZnSO_4$ concentrations by adjusting the trace element stock to yield final

zinc levels of 0.5, 1, 3, 5, or 10 mg/L ZnSO$_4$ in the working 1× medium. (Standard VMM
contains 5 mg/L ZnSO$_4$.) For each zinc condition, 150 µL of a conidial suspension (4 x 10^7
condiospores/mL) was inoculated into 1.5 mL zinc-adjusted Vogel's medium in each well of a
24-well plate (Cellvis #P24-1.5H-N). Plates were incubated at 37°C or 50°C with shaking at 200
rpm for 10 h in the dark. Wells were briefly mixed by pipetting with a P1000 immediately prior to
sampling to ensure uniform suspension. For microscopy, 10 µL of culture was mounted on a
glass slide and imaged using a compound light microscope at 400× magnification under
brightfield illumination. A conidium was considered germinated when the germ tube length
exceeded the diameter of the spore body. For each replicate, at least 200 conidia were
manually counted to calculate the percent germination. Each zinc condition and temperature
combination was assayed in ≥3 biological replicates. For Δ*zip* assays shown in Figure 5B,
germination was quantified at low (0.5 mg/L) and high (10 mg/L) zinc concentrations using the
same conditions, incubation times, and scoring criteria as for wild-type assays.

Statistical analysis

All data processing and statistical analyses were performed in Python (v3.10) using the pandas,
numpy, scipy, statsmodels, matplotlib, and seaborn packages available in the scientific Python
ecosystem (49). Barcode abundance tables from the pooled mutant screens were normalized
by total-count scaling. Gene-level fitness effects were quantified using two complementary
statistical frameworks: (i) a Mann–Whitney U test comparing barcode abundance distributions
between 37°C and 50°C, and (ii) a baseline-normalized log$_2$ fold-change approach in which the
mean normalized abundance across 37°C replicates was used as a reference, and a one-
sample Wilcoxon signed-rank test assessed whether the median log$_2$(50°C/37°C) differed from
zero. Resulting p-values were corrected for multiple testing using Benjamini–Hochberg false
discovery rate (FDR) control, and genes were classified as temperature-responsive when mean
log$_2$FC ≥ 1 and FDR < 0.05. Biological replicate reproducibility was evaluated by pairwise
scatter analysis of normalized barcode abundances, with coefficients of determination (R²)
computed from ordinary least-squares regression. Volcano plots and coefficient-of-variation
(CV) summaries were generated directly from the baseline-normalized log$_2$ fold-change tables.

For germination and growth phenotyping assays, temperature and genotype effects were
assessed using two-way ANOVA (Strain × Temperature) implemented in the statsmodels
package within the Python scientific ecosystem (49). Where a significant strain × temperature
interaction was detected, post-hoc comparisons were conducted. For growth, Tukey's HSD was
applied to the strain factor to identify genotypes that differed from Δ*ku80* across temperatures,
with significance determined by the Tukey-adjusted p-value. Heatmaps were generated with
seaborn, and significance markers were placed directly on the row margins. All figures were
generated in Python and finalized in Inkscape software version 1.4.2.

Data availability

574
Raw sequencing data are available from the NCBI Sequence Read Archive (SRA) under
accession number PRJNA1355313. Germination assay micrographs are available on Figshare
at https://doi.org/10.6084/m9.figshare.30593345. T-DNA sequencing data are in Supplementary
File 1, and barcode sequencing data are in Supplementary File 2. Growth, conidiation and
germination data for wild-type and single-gene validation mutants are in Supplementary File 3.
Plasmids generated in this study are listed in Table 2, and all primers used for cassette
construction and strain verification are provided in Supplementary File 4.

Acknowledgements

We thank Prof. Louise Glass, Dr. Florian Drescher, Dr. José Manuel Villalobos-Escobedo, and Dr. Adam Deutschbauer for invaluable guidance and support of this work. We also thank the members of the Brem Lab for insightful suggestions. O.A.O. acknowledges financial support from the Energy & Biosciences Institute (EBI) at UC Berkeley through the EBI-Shell Postdoctoral Fellow program. This work was supported in part by NIH R01GM120430 to R.B.B.

References

1. Tansey MR, Brock TD. 1972. The Upper Temperature Limit for Eukaryotic Organisms. Proc Natl Acad Sci USA 69:2426.

2. Maheshwari R, Bharadwaj G, Bhat MK. 2000. Thermophilic Fungi: Their Physiology and Enzymes. Microbiol Mol Biol Rev 64:461–488.

3. Wiegant WM. 1992. Growth Characteristics of the Thermophilic Fungus *Scytalidium thermophilum* in Relation to Production of Mushroom Compost. Appl Environ Microbiol 58:1301.

4. Siliakus MF, van der Oost J, Kengen SWM. 2017. Adaptations of archaeal and bacterial membranes to variations in temperature, pH and pressure. Extremophiles 21:651–670.

5. Wang Q, Cen Z, Zhao J. 2015. The survival mechanisms of thermophiles at high temperatures: An angle of omics. Physiology 30: 97–106

6. Xiao W, Zhang J, Huang J, Xin C, Li MJ, Song Z. 2022. Response and regulatory mechanisms of heat resistance in pathogenic fungi. Appl Microbiol Biotechnol 106:5415.

7. Fabri JHTM, Rocha MC, Fernandes CM, Persinoti GF, Ries LNA, Cunha AF da, Goldman GH, Del Poeta M, Malavazi I. 2021. The Heat Shock Transcription Factor HsfA Is Essential for Thermotolerance and Regulates Cell Wall Integrity in *Aspergillus fumigatus*. Front Microbiol 12:656548.

8. Francisco CS, McDonald BA, Palma-Guerrero J. 2023. A transcription factor and a phosphatase regulate temperature-dependent morphogenesis in the fungal plant pathogen *Zymoseptoria tritici*. Fungal Genet Biol 167:103811.

9. Yaakoub H, Mina S, Calenda A, Bouchara JP, Papon N. 2022. Oxidative stress response pathways in fungi. Cell Mol Life Sci 79:333.

10. Carruthers DN, Lee TS. 2022. Translating advances in microbial bioproduction to sustainable biotechnology. Front Bioeng Biotechnol 10:968437.

11.   Adegboye MF, Ojuederie OB, Talia PM, Babalola OO. 2021. Bioprospecting of microbial
strains for biofuel production: metabolic engineering, applications, and challenges.
Biotechnol Biofuels 14:1–21.

12.   Yuan SF, Guo GL, Hwang WS. 2017. Ethanol production from dilute-acid steam exploded
lignocellulosic feedstocks using an isolated multistress-tolerant Pichia kudriavzevii strain.
Microb Biotechnol 10:1581.

13.   Berka RM, Grigoriev I V., Otillar R, Salamov A, Grimwood J, Reid I, Ishmael N, John T,
Darmond C, Moisan MC, Henrissat B, Coutinho PM, Lombard V, Natvig DO, Lindquist E,
Schmutz J, Lucas S, Harris P, Powlowski J, Bellemare A, Taylor D, Butler G, De Vries RP,
Allijn IE, Van Den Brink J, Ushinsky S, Storms R, Powell AJ, Paulsen IT, Elbourne LDH,
Baker SE, Magnuson J, Laboissiere S, Clutterbuck AJ, Martinez D, Wogulis M, De Leon
AL, Rey MW, Tsang A. 2011. Comparative genomic analysis of the thermophilic biomass-
degrading fungi *Myceliophthora thermophila* and *Thielavia terrestris*. Nat Biotechnol
29:922–927.

14.   Contato AG, Borelli TC, Buckeridge MS, Rogers J, Hartson S, Prade RA, Polizeli M de LT
de M. 2024. Secretome Analysis of *Thermothelomyces thermophilus* LMBC 162
Cultivated with Tamarindus indica Seeds Reveals CAZymes for Degradation of
Lignocellulosic Biomass. J Fungi 10(121): 1-14.

15.   Liu Q, Gao R, Li J, Lin L, Zhao J, Sun W, Tian C. 2017. Development of a genome-editing
CRISPR/Cas9 system in thermophilic fungal *Myceliophthora* species and its application
to hyper-cellulase production strain engineering. Biotechnol Biofuels 10:1–14.

16.   Liu Q, Zhang Y, Li F, Li J, Sun W, Tian C. 2019. Upgrading of efficient and scalable
CRISPR-Cas-mediated technology for genetic engineering in thermophilic fungus
Myceliophthora thermophila. Biotechnol Biofuels 12:1–19.

17.   Zhang C, Li N, Rao L, Li J, Liu Q, Tian C. 2022. Development of an Efficient C-to-T Base-
Editing System and Its Application to Cellulase Transcription Factor Precise Engineering
in Thermophilic Fungus *Myceliophthora thermophila*. Microbiol Spectr 10:e02321-21.

18.   Kwon MJ, Schütze T, Spohner S, Haefner S, Meyer V. 2019. Practical guidance for the
implementation of the CRISPR genome editing tool in filamentous fungi. Fungal Biol
Biotechnol 6:1–11.

19.   Huberman LB, Villalobos-Escobedo JM, Skerker JM, Shi R, Rico-Ramirez AM, Adams C,
Arkin AP, Deutschbauer AM, Glass L. 2025. Construction of a randomly barcoded

insertional mutant library in the filamentous fungus *Trichoderma atroviride*. bioRxiv
2025.11.30.691285.

20.   Xu J, Li J, Lin L, Liu Q, Sun W, Huang B, Tian C. 2015. Development of genetic tools for
671   *Myceliophthora thermophila*. BMC Biotechnol 15:35.

673   21.   Billmyre RB, Craig CJ, Lyon JW, Reichardt C, Kuhn AM, Eickbush MT, Zanders SE. 2025.
674   Landscape of essential growth and fluconazole-resistance genes in the human fungal
675   pathogen *Cryptococcus neoformans.* PLoS Biol 23:e3003184.

677   22.   Segal ES, Gritsenko V, Levitan A, Yadav B, Dror N, Steenwyk JL, Silberberg Y, Mielich K,
678   Rokas A, Gow NAR, Kunze R, Sharan R, Berman J. 2018. Gene essentiality analyzed by
679   in vivo transposon mutagenesis and machine learning in a stable haploid isolate of
680   *candida albicans.* mBio 9:1–21.

682   23.   Michel AH, Hatakeyama R, Kimmig P, Arter M, Peter M, Matos J, De Virgilio C, Kornmann
683   BT. 2017. Functional mapping of yeast genomes by saturated transposition. Elife
684   6:e23570

686   24.   Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl J V., Melnyk RA,
687   Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM,
688   Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP, Deutschbauer AM. 2018.
689   Mutant phenotypes for thousands of bacterial genes of unknown function. Nature
690   557:503–509.

692   25.   Blake Billmyre R, Eickbush MT, Craig CJ, Lange JJ, Wood C, Helston RM, Zanders SE.
693   2022. Genome-wide quantification of contributions to sexual fitness identifies genes
694   required for spore viability and health in fission yeast. PLoS Genet 18:e1010462.

696   26.   Ijadpanahsaravi M, Wösten HAB. 2024. Germination strategies of stress-resistant
697   Aspergillus conidia. Curr Opin Food Sci 57:101169.

699   27.   Wyatt TT, Wösten HAB, Dijksterhuis J. 2013. Fungal spores for dispersion in space and
700   time. Adv Appl Microbiol 85:42–91

702   28.   Osherov N, May GS. 2001. The molecular mechanisms of conidial germination. FEMS
703   Microbiol Lett 199:153–160.

29.   Amir H, Pineau R. 1998. Effects of metals on the germination and growth of fungal
isolates from New Caledonian ultramafic soils. Soil Biol Biochem 30:2043–2054.

30.   Feigl G, Lehotai N, Molnár Á, Ördög A, Rodríguez-Ruiz M, Palma JM, Corpas FJ, Erdei L,
Kolbert Z. 2014. Zinc induces distinct changes in the metabolism of reactive oxygen and
nitrogen species (ROS and RNS) in the roots of two Brassica species with different
sensitivity to zinc stress. Ann Bot 116:613.

31.   Santos TA de O, Soares LW, Oliveira LN, Moraes D, Mendes MS, Soares CM de A,
Bailão AM, Bailão MGS. 2024. Zinc Starvation Induces Cell Wall Remodeling and
Activates the Antioxidant Defense System in *Fonsecaea pedrosoi*. J Fungi 10:118

32.   Egan MJ, Wang ZY, Jones MA, Smirnoff N, Talbot NJ. 2007. Generation of reactive
oxygen species by fungal NADPH oxidases is required for rice blast disease. Proc Natl
Acad Sci USA 104:11772–11777.

33.   Zhang Z, Chen Y, Li B, Chen T, Tian S. 2020. Reactive oxygen species: A generalist in
regulating development and pathogenicity of phytopathogenic fungi. Comput Struct
Biotechnol J 18:3344.

34.   Fischer MS, Glass NL. 2019. Communicate and Fuse: How Filamentous Fungi Establish
and Maintain an Interconnected Mycelial Network. Front Microbiol 10:619.

35.   Ghosh A, Servin JA, Park G, Borkovich KA. 2014. Global analysis of serine/threonine and
tyrosine protein phosphatase catalytic subunit genes in *Neurospora crassa* reveals
interplay between phosphatases and the p38 mitogen-activated protein kinase. G3
(Bethesda) 4:349–365.

36.   Desmarini D, Lev S, Furkert D, Crossett B, Saiardi A, Kaufman-Francis K, Li C, Sorrell
TC, Wilkinson-White L, Matthews J, Fiedler D, Teresa Djordjevic J. 2020. Ip7-spx domain
interaction controls fungal virulence by stabilizing phosphate signaling machinery. mBio
11:1–20.

37.   Tamayo E, Gómez-Gallego T, Azcón-Aguilar C, Ferrol N. 2014. Genome-wide analysis of
copper, iron and zinc transporters in the arbuscular mycorrhizal fungus *Rhizophagus*
irregularis. Front Plant Sci 5:113084.

38. Patkar RN, Ramos-Pamplona M, Gupta AP, Fan Y, Naqvi NI. 2012. Mitochondrial β-
oxidation regulates organellar integrity and is necessary for conidial germination and
invasive growth in *Magnaporthe oryzae*. Mol Microbiol 86:1345–1363.

39. Leuenberger P, Ganscha S, Kahraman A, Cappelletti V, Boersema PJ, Von Mering C,
Claassen M, Picotti P. 2017. Cell-wide analysis of protein thermal unfolding reveals
determinants of thermostability. Science (1979) 355.

40. Drescher F, Li Y, Villalobos-Escobedo JM, Haefner S, Huberman LB, Glass NL. 2024.
Transcriptomic and genetic analysis reveals a Zn2Cys6 transcription factor specifically
required for conidiation in submerged cultures of *Thermothelomyces thermophilus*. mBio
https://doi.org/10.1128/mbio.03111-24.

41. Kwon MJ, Schütze T, Spohner S, Haefner S, Meyer V. 2019. Practical guidance for the
implementation of the CRISPR genome editing tool in filamentous fungi. Fungal Biol
Biotechnol 6:15.

42. Vogel HJ. 1956. A Convenient Growth Medium for Neurospora crassa. Microbiol Genet
Bull 13:42.

43. Michielse CB, Hooykaas PJJ, van den Hondel CAMJJ, Ram AFJ. 2008. Agrobacterium-
mediated transformation of the filamentous fungus *Aspergillus awamori*. Nat Protoc
3:1671–1678.

44. Helmann TC, Filiatrault MJ, Stodghill P V. 2022. Genome-Wide Identification of Genes
Important for Growth of *Dickeya dadantii* and *Dickeya dianthicola* in Potato (Solanum
tuberosum) Tubers. Front Microbiol 13:778927.

45. Kim J, Coradetti ST, Kim YM, Gao Y, Yaegashi J, Zucker JD, Munoz N, Zink EM, Burnum-
Johnson KE, Baker SE, Simmons BA, Skerker JM, Gladden JM, Magnuson JK. 2021.
Multi-Omics Driven Metabolic Network Reconstruction and Analysis of Lignocellulosic
Carbon Utilization in *Rhodosporidium toruloides*. Front Bioeng Biotechnol 8.

46. oaogunyewo-git. oaogunyewo-git/Thermothelomyces-Tnseq-project: Initial code release
for manuscript reproducibility https://doi.org/10.5281/ZENODO.17586924.

47. Coradetti ST, Pinel D, Geiselman GM, Ito M, Mondo SJ, Reilly MC, Cheng Y-F, Bauer S,
Grigoriev I V, Gladden JM, Simmons BA, Brem RB, Arkin AP, Skerker JM. 2018. .

Functional genomics of lipid metabolism in the oleaginous yeast *Rhodosporidium*
toruloides 7:e32110.

48. Siebecker B, Schütze T, Spohner S, Haefner S, Meyer V. 2023. Transcriptomic insights
into the roles of the transcription factors Clr1, Clr2 and Clr4 in lignocellulose degradation
of the thermophilic fungal platform *Thermothelomyces thermophilus*. Front Bioeng
Biotechnol 11:1279146.

49. Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser
E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk MH, Brett M,
Haldane A, del Río JF, Wiebe M, Peterson P, Gérard-Marchant P, Sheppard K, Reddy T,
Weckesser W, Abbasi H, Gohlke C, Oliphant TE. 2020. Array programming with NumPy.
Nature 585:357–362.

50. Randhawa A, Ogunyewo OA, Eqbal D, Gupta M, Yazdani SS. 2018. Disruption of zinc
finger DNA binding domain in catabolite repressor Mig1 increases growth rate, hyphal
branching, and cellulase expression in hypercellulolytic fungus *Penicillium funiculosum*
NCIM1228. Biotechnol Biofuels 11:15.

51. Rizwan HM, Yang Q, Yousef AF, Zhang X, Sharif Y, Kaijie J, Shi M, Li H, Munir N, Yang X,
Wei X, Oelmüller R, Cheng C, Chen F. 2021. Establishment of a novel and efficient
agrobacterium-mediated in planta transformation system for passion fruit (Passiflora
edulis). Plants 10:2459.

Figure captions

Figure 1. Growth and developmental measurements of *T. thermophilus* across temperatures. (A) Colony morphology of the wild-type strain grown on solid Vogel's medium at 37°C, 45°C, and 50°C and imaged at 72 h, 120 h, and 168 h post-inoculation. (B) Conidial density measured from cultures grown at 37°C, 45°C, and 50°C after 168 h. The *x*-axis indicates growth temperature (°C), and the y-axis indicates conidia per milliliter (log scale). Bars represent mean ± SD from three biological replicates. (C) Dry mycelial biomass measured from 50 ml liquid cultures incubated for 24 h at the indicated temperatures. The *x*-axis indicates incubation temperature (°C), and the *y*-axis indicates dry mycelial weight (mg per 50 ml culture). Bars represent mean ± SD from three biological replicates.

Figure 2. Construction and genomic distribution of the *T. thermophilus* barcoded insertional mutant library. (A) Overview of the Agrobacterium-mediated transformation workflow used to generate the barcoded insertion mutant library. *T. thermophilus* conidia were co-cultivated with *A. tumefaciens* containing plasmids with barcoded T-DNA. Transformants were selected on hygromycin-containing medium. Transformant colonies were bulked by allowing them to conidiate on selection plates, and conidia from all plates were harvested and pooled to generate the barcoded insertion mutant library (B) Genome-wide mapping of T-DNA insertions across the seven chromosomes of *T. thermophilus*. The *x*-axis indicates chromosomal position (grouped by chromosome), and the *y*-axis indicates the number of mapped T-DNA insertions per 50 kb genomic window. Colors correspond to individual chromosomes.

Figure 3. Genome-wide BarSeq fitness profiling identifies temperature-dependent mutant defects. (A) Experimental workflow for barcode sequencing (BarSeq) of the *T. thermophilus* mutant pool. Conidia were inoculated from the pool stock onto solid medium, harvested, and used to start liquid cultures. Cultures were incubated for 24 h at 37°C or 50°C. Genomic DNA was extracted from each culture, and barcode regions were amplified and sequenced. Sequencing reads were used to quantify barcode abundance for downstream fitness analysis. (B) Example comparison of mutant abundances between two biological replicates grown at 37°C. Each point represents one mutant, and the *x*- and *y*-axes report normalized abundance in the indicated replicate. (C) Each point reports the results of a test for differential abundance of insertional mutants in one gene between temperatures. The *x*-axis reports the log of the ratio of two quantities: the average abundance, across genotypes and biological replicates, of mutants at the respective gene at 50°C, and the analogous quantity at 37°C. The *y*-axis indicates – $\log_{10}$(adjusted *p*-value) from statistical testing. Points in red indicate loci meeting quality control thresholds for temperature-dependent abundance change in terms of significance, effect size, and variability (see Methods). Genes chosen for single-gene follow-up are highlighted and labeled with arrows.

Figure 4. Germination and biomass measurements for temperature-tested single-gene deletion strains. (A) Top, germination assay. Conidia were inoculated into microtiter wells and incubated at a temperature of interest, followed by scoring via microscopy. Bottom, each row reports results from germination assays of the indicated *T. thermophilus* strain at the indicated temperatures. Each cell reports the average, across replicates, of the percent of conidia scored as germinating. (B) Top, biomass assay. Conidia were inoculated into liquid medium and incubated to allow germination and pre-growth; the resulting mycelium was used as inoculum into a second culture, followed by incubation at a temperature of interest, and biomass harvest and quantification. Bottom, each row reports results from growth assays of the indicated *T. thermophilus* strain at the indicated temperatures. Each cell reports the average, across

replicates, of dry biomass (mg) for each strain at each temperature. Statistical significance was
evaluated using a two-way ANOVA (strain × temperature) with post-hoc comparisons relative to
the Δku80 reference strain at each temperature. Asterisks denote significance (*$p < 0.05$; **$p <$
0.01; ***$p < 0.001$).

Figure 5: Zinc-dependent modulation of germination in *T. thermophilus*. (A) Germination of
wild-type *T. thermophilus* across zinc concentrations and temperatures**.** The *x*-axis indicates
zinc concentration in the medium (mg/L), and the *y*-axis indicates the percentage of germinated
conidia. Each point represents the mean value, and error bars denote standard deviation across
biological replicates. (B) Germination of *T. thermophilus* Δ*zip* across zinc concentrations and
temperatures. The *x*-axis indicates temperature and the *y*-axis is as in (A). Each bar represents
the mean value, and error bars denote standard deviation across biological replicates. ns, not
significant in a *t*-test comparing germination between zinc concentrations for the indicated
temperatures.

Supplementary figure captions

Supplementary Figure 1. Reproducibility of barcode abundance measurements across biological replicates. Data are as in Figure 3B of the main text, except that panels A–C correspond to replicate comparisons at 37°C for the Batch 1 mutant pool (sets A1–A2, B1–B2, and C1–C2); D–E show replicate comparisons at 50°C for the Batch 1 mutant pool (sets A1–A2 and B1–B2); F-G correspond to replicate comparisons at 37°C for the Batch 2 mutant pool; H-I correspond to replicate comparisons at 50°C for the Batch 2 mutant pool. The coefficient of determination (R^2) for each comparison is shown in the top left of each plot.

Supplementary Figure 2. PCR confirmation of targeted gene deletions in *T. thermophilus*. Diagnostic PCR was used to verify homologous replacement of each target locus in the deletion strains (Δ*mfs*, Δ*syg1*, Δ*nrc-1*, Δ*pty-5*, Δ*zincTF*, Δ*zip*, Δ*MYCTH_2313252* and Δ*ech*). For each gene, lanes labeled C correspond to the parental Δ*ku80* control, and adjacent lanes show amplicons from independent deletion-mutant isolates. Loss of the wild-type band and/or the presence of the expected deletion-specific product confirms successful gene replacement at the target locus. M refers to the DNA size marker. The expected size bands for each of the wild type and transformant PCRs are listed in Supplementary File 4.

Supplementary file captions

Supplementary File S1. T-DNA junction sequencing of *T. thermophilus* insertional mutants. Listed are all mapped insertion mutants in the *T. thermophilus* genome. Columns report barcode and reverse-complement sequences (barcode, rcbarcode), total read support (nTot) and primary-locus support (n), genomic location (scaffold, strand, pos), mapping classification (type), read support for the main mapped site (nMainLocation) and for T-DNA border–derived reads (nInsert), all alternate genomic alignments (All genomic mappings), and annotated overlapping or nearest gene (gene).

Supplementary File S2. Genome-wide BarSeq fitness profiles of *T. thermophilus* at 37°C and 50°C. This file contains three worksheets reporting results of gene-level tests for differential abundance between temperatures from BarSeq data. In each sheet, GeneID is the annotated gene identifier; Mean_log2FC is the mean $\log_2$ fold change in mutant abundance (50°C/37°C); P-Value is the nominal *p*-value from the Wilcoxon rank-sum test for differential abundance between temperatures; Adjusted_P-Value is the multiple-testing–corrected *p*-value (Benjamini–Hochberg); protein_names is the predicted or annotated protein name; Std_log2FC is the standard deviation of mean $\log_2$ fold change in mutant abundance (50°C/37°C) across mutants and replicates; and CV_log2FC_percent is the coefficient of variation of mean $\log_2$ fold change in mutant abundance (50°C/37°C) across mutants and replicates, expressed as a percentage. Worksheet 1 includes results from all genes with BarSeq data that met our quality control threshold (≤15 detected insertions; see Methods). Worksheet 2 reports genes whose mutants exhibited temperature-responsive fitness that met significance and effect-size thresholds for which mean $\log_2$ fold change in mutant abundance (50°C/37°C) ≥ 1. Worksheet 3 reports genes that met significance and effect-size thresholds for which mean $\log_2$ fold change in mutant abundance (50°C/37°C) ≤ −1.

Supplementary File S3. Raw measurements of conidiation, biomass accumulation, and germination of wild-type and single-gene validation mutant *T. thermophilus*. This file contains five worksheets reporting unprocessed measurements underlying Figures 1, 4, and 5. Worksheet 1 reports raw counts of conidia harvested from wild-type cultures grown on solid VMM + 2% sucrose at 37°C, 45°C, and 50°C. Columns report temperature and conidia count. Worksheet 2 reports dry-weight biomass measurements for wild-type cultures grown in liquid VMM containing 2% sucrose at 30°C, 37°C, 45°C, 50°C, and 55°C. Columns report temperature and growth (mg). Worksheet 3 reports Biomass accumulation dry-weight biomass measurements for all deletion mutants and the Δ*ku*80 parent grown in liquid VMM + 2% sucrose at 37°C and 50°C. Columns report strain, temperature, and growth (mg). Worksheet 4 reports microscopy-based germination measurements for all deletion mutants and the Δku80 parent at 37°C and 50°C. Columns report strain, temperature, total number of conidia, number of germinated conidia, and percentage germination. Worksheet 5 reports the raw germination data for Δku80 conidia exposed to varying Zn concentrations at 37°C and 50°C. Columns report zinc concentration, temperature, total number of conidia, number of germinated conidia, and percentage germination.

Supplementary File S4. Primer sequences used for construction of deletion cassettes and validation of gene deletions by PCR. This file contains three worksheets detailing the oligonucleotides used in this study for TnSeq/BarSeq library preparation, deletion-cassette construction, and PCR-based mutant verification. Worksheet 1 reports the primers used for Tnseq and Barseq amplifications, indexing, and nested enrichment of T-DNA junction fragments. The columns report primer_name, primer_sequence, index_sequence (for multiplexing), and index_reverse (custom P1 index primer). Worksheet 2 reports the primers used to assemble deletion cassettes and to verify correct integration in transformants by PCR.

938   Columns report GeneID, primer_name, primer_sequence, and purpose (e.g., 5′ flank
939   amplification, junction PCR, confirmation PCR). Worksheet 3 reports the expected amplicon
940   sizes for wild-type (Δku80 background) and deletion mutants after cassette integration. Columns
941   report gene, primer_set, expected_size_in_Δku80 (kb), and expected_size_in_mutant (kb).

GeneID	Gene symbol	Protein name	Mutant effect on abundance, $\log_2$(50°C/37°C)	Adjusted P-Value
MYCTH_2139072	zincTF	Zn(2)-C6 fungal-type domain-containing protein	1.98	1.12 E-05
MYCTH_2300403	pty-5	Tyrosine-specific protein phosphatase-domain-containing protein	1.95	5.84 E-02
MYCTH_100809	syg1	SPX domain-containing protein	1.90	2.01E-08
MYCTH_2305107	nrc-1	MAPKK kinase	1.15	8.6 E-04
MYCTH_116881	mfs	MFS general substrate transporter	-0.98	0.0184
MYCTH_2303755	ech	Enoyl-CoA hydratase	-1.40	0.0078
MYCTH_2307966	zip	Zinc/iron permease	-1.45	0.0184
MYCTH_2313252	MYCTH_2313252	Putative UPF0658 Golgi apparatus membrane protein	-1.83	0.0171

943
Table 1. Hits from high-throughput screen subjected to single-gene validation. Each row
reports effects of mutation of one gene on temperature-dependent fitness as measured by
large-scale pooled T-DNA mutagenesis, from Figure 3C, for a locus chosen for follow-up single-
gene validation analysis. For each gene, the table reports the *T. thermophilus* gene identifier
(GeneID); the corresponding gene symbol; the annotated protein name; the log of the ratio of
two quantities: the average abundance, across genotypes and biological replicates, of mutants
at the respective gene at 50°C, and the analogous quantity at 37°C; and the adjusted *p*-value
from a Wilcoxon rank-sum test for differential abundance of the respective mutants between
temperatures.
953

Strains	Description	References / sources
Escherichia coli DH5α	F- Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17	Invitrogen
Agrobacterium tumefaciens LBA4404	Ach5 with pAch5, Δ*tra* Δ*occ* ΔT$_L$ ΔT$_R$ Rif^r	(50)
Thermothelomyces thermophilus ATCC 42464	Wild-type reference strain used as the parental background in this study	(40)
MJK20.2 (Δ*ku*80)	Mutant strain of *T. thermophilus* having deletion of *ku*80 gene (MYCTH_2118116)	(40)
Δ*ech*	Mutant strain of *T. thermophilus* having deletion of *ech* (MYCTH_2303755) and *ku*80 (MYCTH_2118116) genes	This study
Δ*mfs*	Mutant strain of *T. thermophilus* having deletion of *mfs* (MYCTH_116881) and *ku*80 (MYCTH_2118116) genes	This study
Δ*nrc-1*	Mutant strain of *T. thermophilus* having deletion of *nrc-1* (MYCTH_2305107) and *ku*80 (MYCTH_2118116) genes	This study
Δ*pty-5*	Mutant strain of *T. thermophilus* having deletion of *pty-5* (MYCTH_2300403) and *ku*80 (MYCTH_2118116) genes	This study
Δ*syg1*	Mutant strain of *T. thermophilus* having deletion of *syg1* (MYCTH_100809) and *ku*80 (MYCTH_2118116) genes	This study
Δ*MYCTH_2313252*	Mutant strain of *T. thermophilus* having deletion of *MYCTH_2313252* and *ku*80 (MYCTH_2118116) genes	This study
Δ*zincTF*	Mutant strain of *T. thermophilus* having deletion of *zincTF* (MYCTH_2139072) and *ku*80 (MYCTH_2118116) genes	This study
Δ*zip*	Mutant strain of *T. thermophilus* having deletion of *zip* (ΔMYCTH_2307966) and *ku*80 (MYCTH_2118116) genes	This study
Plasmids	**Description**	**References / sources**
pCambia 1302	Kanamycin and hygromycin resistant vector containing EGFP	(51)
pLH159RB	Plasmids for *A. tumefaciens*-mediated	(19)

	transformation with barcoded T-DNA	
pOORB007	pCAMBIA1302-based plasmid bearing the *nrc-1* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *nrc-1* ORF	This study
pOORB008	pCAMBIA1302-based plasmid bearing the *syg1* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *syg1* ORF	This study
pOORB009	pCAMBIA1302-based plasmid bearing the *ech* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *ech* ORF	This study
pOORB012	pCAMBIA1302-based plasmid bearing the *zincTF* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *zincTF* ORF	This study
pOORB014	pCAMBIA1302-based plasmid bearing the *pty-5* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *pty-5* ORF	This study
pOORB018	pCAMBIA1302-based plasmid bearing the *mfs* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *mfs* ORF	This study
pOORB020	pCAMBIA1302-based plasmid bearing the MYCTH_2313252 deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the MYCTH_2313252 ORF	This study
pOORB021	pCAMBIA1302-based plasmid bearing the *zip* deletion cassette, consisting of ~1 kb upstream and downstream homology arms flanking the hph selectable marker in place of the *zip* ORF	This study

957 **Table 2. List of strains and plasmids used for this study.** For each row, the first column lists
958 the strain or plasmid, the second column provides its description, and the third column indicates
959 the corresponding reference or source for strain construction and plasmid generation.
960

	37°C	45°C	50°C
72 h			
120 h			
168 h			

A

B

Conidia per ml (millions, log scale)

Temperature (°C)

C

Dry mycelia weight (mg/50ml)

Temperature (°C)

A

Aliquot of mutant pool → *T. thermophilus* mutant expansion → → 37 °C / 24 hr / 50 °C → Grown mutant pool → PCR amplify barcodes → → Barcode abundance readout

B

Mutant abundance at 37°C, replicate 2 vs Mutant abundance at 37°C, replicate 1

C

-log10(adjusted p-value) vs Mean log₂ (abundance, 50°C/abundance, 37°C)

Not significant / Significant

syg1, zincTF, nrc-1, pty-5, mfs, ech, zlp, MYCTH_2313252