Convergent evolution on *RBA1* regulates *Candidozyma auris* host adaptation

Laura A. Doorley[1], Alyssa A. La Bella[2], Sarah J. Jones[1], Guolei Zhao[3], Mohammad Asadzadeh[4], Quanita J. Choudhury[1], Suhail Ahmad[4], Eiman Mokaddas[4,5], Inaam Al-Obaid[6], Wadha Alfouzan[4,7], Khalifa Al-Banwan[4,8], Teresa R. O'Meara[3] Felipe H. Santiago Tirado[2], Ana L. Flores Mireles[2], Jeffrey M. Rybak[1*]

[1] Department of Pharmacy and Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN 38105, USA.
[2] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556, USA
[3] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI 48109, USA.
[4] Department of Microbiology, Faculty of Medicine, Kuwait University, Jabriya 46300, Kuwait.
[5] Department of Microbiology, Ibne-Sina Hospital, Shuwaikh, Kuwait
[6] Department of Microbiology, Al-Sabah Hospital, Shuwaikh, Kuwait
[7] Department of Microbiology, Farwaniya Hospital, Farwaniya, Kuwait
[8] Department of Microbiology, Amiri Hospital, Kuwait City, Kuwait

[*] Corresponding author: Jeffrey.Rybak@stjude.org

Jeffrey M. Rybak, PharmD, PhD
Assistant Member
Department of Pharmacy and Pharmaceutical Sciences
St. Jude Children's Research Hospital
262 Danny Thomas Place
Chili's Care Center, I5304, Mail Stop #313
Memphis TN, 38105

Office: 901.595.2368
Fax: 901.595.3125

ABSTRACT

Candidozyma auris is an emerging healthcare-associated fungal pathogen, increasingly isolated from clinical outbreaks, with a high propensity to colonize patients and the medical environment. Here, we leveraged *C. auris* isolates from multi-facility clinical outbreaks to identify genomic patho-adaptations promoting persistence and dissemination in the healthcare environment. Genomic and phylogenetic analyses revealed loss-of-function mutations in the uncharacterized *C. auris* transcription factor gene *RBA1* to have independently emerged multiple times within these outbreaks. We demonstrate loss of *RBA1* increases *C. auris* adhesion to plastic and human keratinocytes, enhances biofilm formation, and exacerbates fungal burden in a mouse model of catheter-associated urinary tract infection. Finally, we uncover mutations in *RBA1* have emerged during multiple *C. auris* outbreaks across the globe, and that *RBA1* mutant lineages are present among large ongoing clinical outbreaks. These results reveal loss-of-function mutations in *C. auris* *RBA1* as a novel and clinically relevant genetic determinant of enhanced outbreak characteristics.

INTRODUCTION

Candidozyma auris (formerly *Candida auris*) is an emerging pathogenic yeast first identified in 2009 and later recognized by the World Health Organization as a critical threat to global public health in 2022[1, 2]. Initially isolated from a limited number of individual patient infections, *C. auris* was soon revealed as the cause of multiple nosocomial outbreaks of invasive fungal infections across the globe[3]. Now, *C. auris* outbreaks have been reported from more than 50 countries, representing thousands of documented infections annually, including 4,514 cases in 2023 in the United States alone[4, 5, 6]. *C. auris* is distinguished from other fungal pathogens as it is not known to be ubiquitous within common natural environmental reservoirs nor is it a member of the normal human microbiota[7]. Instead, *C. auris* infections are healthcare-associated and typically occur following a lateral transmission to vulnerable patients within hospital settings[8]. Multiple studies have demonstrated that once introduced into the medical environment, *C. auris* rapidly spreads among both patients and medical surfaces, acting as an invasive species in these healthcare settings. As a result, clinical *C. auris* infections routinely occur in series or as clustered clinical outbreaks, sometimes spanning multiple facilities or institutions within an affected healthcare network[9, 10].

As with other healthcare-associated and multi-drug-resistant pathogens, the characterization and tracking of emergent microbial lineages with traits impacting clinical outcomes is paramount to infection prevention and treatment. Prior study of *C. auris* evolution during infection and exposure to clinically relevant stressors has largely focused on the emergence of antifungal resistance[11, 12, 13, 14]. We and others have shown, through both *in vitro* evolution experiments and by studying sets of isogenic clinical isolates, that *C. auris* is capable of rapidly adapting to antifungal-induced stress and that multiple independent drug-resistant lineages can emerge after only a single antifungal exposure[11, 12, 13, 14]. However, investigations of patho-adaptation in the clinical setting or during outbreaks of infections remains unexplored.

In this work, we leveraged a collection of *C. auris* clinical isolates from multi-institutional

outbreaks of infection to identify novel genetic markers and link them to traits driving host

adaptation and outbreak potential in clinical settings. We identified multiple independent lineages

of *C. auris* harboring distinct mutations in a previously uncharacterized fungal transcription factor,

here named, Repressor of Biofilm and Adhesion 1 (*RBA1*). Genetic and transcriptomic

interrogation revealed loss of *RBA1* to result in enhanced biofilm formation, increased adhesion

to multiple surfaces including human keratinocytes, and significantly elevated expression of

C. auris adhesins and virulence factors. Moreover, loss of *RBA1* was found to significantly

increase fungal burden in a mouse model of catheter-associated urinary tract infection (CAUTI),

an infection characteristically involving biofilms in urine where *C. auris* is often isolated[15]. Finally,

analysis of publicly available whole genome sequencing data revealed numerous distinct *RBA1*

mutations present in hundreds of clinical isolates, demonstrating consistent and convergent loss-

of-function mutations from isolates from well-documented and ongoing clinical outbreaks of

C. auris infections across the globe. Collectively these findings demonstrate *RBA1* encodes a

repressor of *C. auris* biofilm and adhesion and implicate mutations in *RBA1* as clinically relevant

adaptations promoting *C. auris* survival during healthcare-associated outbreaks of infection.

RESULTS

Clinical emergence of multiple *C. auris* outbreak lineages carrying *RBA1* mutations.

Whole genome sequencing (WGS) analysis revealed all *C. auris* isolates belonged to

Clade I, one of three major genetic clades previously linked to healthcare-associated outbreaks[16].

This included 34 and 36 isolates from the previously described subclades Ib and Ic, respectively[17].

A total of 155,135 SNPs were identified among the 70 included isolates as compared to the Clade

I reference genome (B8441v3), with 46,231 predicted to impact coding regions. As anticipated

from characteristically fluconazole-resistant Clade I isolates, all isolates were found to harbor one

or more mutations in the genes *ERG11*, *TAC1B*, and/or *MRR1*, which we and others have

previously shown to directly contribute to fluconazole resistance[14, 18, 19, 20]. Additionally, 16 isolates

were found to possess mutations known to contribute to echinocandin resistance (altering *FKS1*)

and 4 isolates possessed mutations in genes associated with amphotericin B (AmB) resistance

(*ERG3*, 1 isolate; *ERG6*, 3 isolates)[12, 21, 22].

Phylogenetic analysis revealed the population structure of this collection to consist of 69

isolates grouping into three major clusters, with 35, 24, and 10 isolates grouping into clusters A,

B, and C, respectively (**Fig. 1**). A single isolate (SKU043) did not group into a larger cluster and

was more distantly related. Within clusters, isolates were found to be largely clonal, with an

average of 5.3 to 12.6 coding region SNPs separating each isolate (**Fig. S1**), while SKU043

differed from all other isolates by 43 to 113 coding region SNPs. Clusters A and C each contained

isolates originating from multiple healthcare facilities within Kuwait, while cluster B originated

entirely from a single tertiary care hospital. Notably, multiple instances were identified where

isolates from different patients and even different hospitals were found to be more genetically

similar than serial isolates originating from the same patient case, as might be expected in the

setting of a clinical outbreak.

To investigate potential patho-adaptations occurring during outbreaks of *C. auris*

infections, we further evaluated the WGS data to identify genes where mutations were acquired

on multiple occasions and/or mutations emerged and persisted among the outbreak population.

Intriguingly, this revealed 19 isolates harboring mutations in a previously uncharacterized *C. auris*

gene B9J08_03209 (identified as B9J08_002136 in B8441v2 and hereafter named *RBA1*), which

is predicted to encode a fungal-specific zinc-cluster transcription factor (ZCF). Although initially

identified as an ortholog of *WOR2* in *C. albicans,* the proteins are not syntenic, with only 22%

sequence identity, primarily located in the DNA-binding domain. The 19 isolates with mutations in

RBA1 included representatives from all three major phylogenetic clusters, including all 10 isolates

in cluster C, as well as the non-clustered isolate SKU043. Four different mutations in *RBA1* were

identified, all considered likely loss-of-function mutations, with three encoding an early stop codon

(upstream of the predicted DNA-binding domain) and one altering a highly conserved cysteine

residue in the $Zn(II)_2Cys_6$ DNA-binding motif. The early stop codon mutation, $RBA1^{E47*}$, was found

to have emerged and persisted among two intensive care unit (ICU) patients, these cases and

the associated isolates were further evaluated to identify the potential clinical impact of *RBA1*

mutations.

Loss of *RBA1* enhances *C. auris* biofilm formation.

A subset of isolates from patient cases 51 and 53 have been previously characterized with

consideration to antifungal susceptibility while investigating the emergence of echinocandin and

amphotericin B resistance-conferring mutations, respectively[12, 23, 24]. In brief, both were adult

patients with complex medical histories including hematologic malignancies admitted to the ICU

of a tertiary care center and received multiple courses of the antifungals liposomal amphotericin

B (AmB) and caspofungin (CSF). Over the course of approximately eight months, a total of 14 *C.*

auris isolates were cultured from the respiratory tract, urinary tract, and blood of patients 51 and

53 (**Fig. 2a**). Notably, while the earliest isolate in these cases, SKU013, harbored a wildtype *RBA1*

allele ($RBA1^{WT}$), an early stop codon encoding mutation in *RBA1* ($RBA1^{E47*}$) emerged in an isolate

obtained from culturing the urinary catheter (SKU018). This mutation was subsequently found in

all but one *C. auris* isolate cultured from patient 53. Further details of these cases can be found

in **Supplemental Data**

Of the 14 isolates cultured from patient 51 and patient 53, 13 were found to be closely

related (separated by 2 to 8 SNPs within coding regions, **Fig. S1**) and grouped into cluster B.

While *RBA1* mutant isolates were observed to exhibit increased biofilm formation under standard

laboratory conditions, there was no association between antifungal susceptibility and *RBA1*

genotype, (**Fig. 2b, c**). As the earliest isolate where the $RBA1^{E47*}$ mutation was present (SKU018)

was cultured from the urinary tract of patient 51, and multiple subsequent isolates with this

mutation were isolated from the urinary tract of patient 53, we used our fibrinogen-urine biofilm

model [25] that utilizes serum-supplemented urine and fibrinogen-coated surfaces to evaluate

biofilm formation in conditions mimicking the bladder during CAUTI. In this model, isolates

carrying the $RBA1^{E47*}$ mutation exhibited enhanced biofilm formation as compared to the earliest

isolate SKU013 (approximately 2 to 6-fold higher), while biofilm formation was relatively

unchanged among closely related $RBA1^{WT}$ isolates (**Fig. 2d**). To evaluate if other clinically

emergent predicted loss-of-function mutations in $RBA1$ had a similar effect on biofilm formation,

we also compared the biofilm formation of isolates harboring the $RBA1^{Q174*}$ (SKU055), $RBA1^{Q225*}$

(SKU043), and $RBA1^{C303S}$ (SKU056) mutations to their closest available genetic relatives

(SKU021, SKU034, and SKU037, respectively) using the same model. In each case, biofilm

formation was observed to be significantly higher in the isolates harboring mutations in $RBA1$

(**Fig. 2e**), further bolstering the connection between loss of $RBA1$ and biofilm formation.

From these observations, we hypothesized that loss-of-function mutations in $RBA1$ are a

genetic determinant of enhanced *C. auris* biofilm formation. To test this hypothesis, we next

generated the $RBA1$-knockout (*Δrba1*) and $RBA1$-revertant (*Δrba1*::$RBA1^+$) control strains in the

SKU013 clinical isolate background and evaluated biofilm formation. In the fibrinogen-urine

model, *Δrba1* strains were found to exhibit significantly increased biofilm formation (3.9 to 4.8-fold

greater than SKU013; p<0.0001). This enhancement of biofilm formation resembled the isogenic

clinical isolate SKU018 which carries the $RBA1^{E47*}$ mutation (3.3-fold greater than SKU013;

p<0.0001) (**Fig. 3a**). Importantly, restoration of the $RBA1$ gene in *Δrba1*::$RBA1^+$ control strains

returned biofilm formation to levels comparable to SKU013.

We further investigated the impact of $RBA1$ using a plasma-RPMI biofilm model that

utilized serum-supplemented RPMI culture media and human plasma-coated surfaces to evaluate

biofilm production that could apply more generally to clinical settings. Once again, *Δrba1* strains

were found to produce more robust biofilms (6.2 to 6.8-fold greater than SKU013; p<0.0001),

while *Δrba1*::$RBA1^+$ control strains closely matched the biofilm formation of SKU013 (**Fig. 3b**).

Examination of the plasma-RPMI biofilms using scanning electron microscopy, showed scattered

patches of biofilm predominantly consisting of one to two layers of cells for both SKU013 and

Δrba1::*RBA1*⁺ (**Fig. 3c**). By comparison, *Δrba1* strains formed more confluent biofilms of variable

but generally greater thickness with greater production of extracellular matrix. (**Fig. 3d**).

Loss of *RBA1* increases expression of *C. auris* adhesins and virulence factors.

As *RBA1* is predicted to encode a transcription factor and our earlier experiments had

shown loss of *RBA1* significantly enhances biofilm formation, we next evaluated the impact of

RBA1 on the *C. auris* transcriptome during both planktonic and biofilm growth conditions using

our *Δrba1* strain and the parental *RBA1*^WT isolate SKU013. When cells were grown in RPMI tissue

culture media under planktonic conditions, transcriptional profiling revealed loss of *RBA1* to have

a modest impact, and only 55 of 5,594 *C. auris* genes (0.98%) were differentially expressed (≥|1.5|

fold-change and false discovery rate [FDR] ≤0.05) (**Fig. 4a**). Of these, 42 genes were upregulated

in the absence of *RBA1*, and these genes were enriched for those involved in biofilm formation

(p= 1.60e-07) and cell aggregation (p= 2.18e-07), and no enrichment for biological processes was

found among the 13 down-regulated genes. By comparison, when cells were harvested from

actively growing biofilms, 701 of 5,594 genes (12.53%) were observed to be differentially

expressed, including 250 and 451 genes which were up- and down-regulated in the *Δrba1* strain,

respectively (**Fig. 4b**). Up-regulated genes were enriched for those involved in transmembrane

transport (p= 0.01351), regulation of RNA biosynthetic process (p=0.02583), cell aggregation

(p=0.05187), and regulation of primary metabolic process (p=0.07722), while down regulated

genes were enriched for 18 different biological processes including protein biosynthetic process

(p= 2.35e-28), macromolecule biosynthetic process (p= 4.05e-10), premeiotic DNA replication (p=

0.00273), and steroid biosynthetic process (p= 0.02579). Thirty-two *C. auris* genes were up-

regulated in the absence of *RBA1* under both growth conditions, and these genes were enriched

for those involved in biofilm formation and cell aggregation (**Fig. 4c**). Notably, this included four

C. auris adhesins (*ALS4112/ALS5, IFF4*, *IFF4109*, and *SCF1*), three of which we have previously

shown to impact virulence in a mouse model of invasive candidiasis[26, 27].

As many genes in the *C. auris* genome remain uncharacterized and/or currently have

incomplete annotations, we further evaluated differential expression among *C. auris* genes with

homology to those associated with adhesion, biofilm formation, and cell wall architecture in the

model pathogenic yeast *Candida albicans.* This revealed additional *C. auris*-specific adhesins,

cell wall-associated proteins, GPI-anchored proteins, and aspartic-type endopeptidases

exhibiting differential expression under biofilm conditions in the absence of *RBA1* (**Fig. 4d**)*.

Notably, four of the differentially expressed genes associated with the fungal cell wall,

B9J08_002109, B9J08_003350, *PHR1*, and *XOG1*, are the predicted orthologs of *C. albicans* cell

wall proteins known to be involved in cell wall remodeling, cell wall masking from immune

detection, and response to host immune cells[28, 29, 30].

As *RBA1* appears to repress biofilm formation in *C. auris,* we also examined the

expression of *C. auris* orthologs of genes known to regulate biofilm formation in the model

pathogenic yeast, *C. albicans* (**Fig. 4d**)*. UME6* and B9J08_002529 (ortholog of *C. albicans*

BRG1), exhibited significantly increased expression in the *Δrba1* strain under planktonic

conditions. In *C. albicans,* these genes are known to be important positive regulators of hyphal

growth and biofilm formation*. Further, UME6* has previously been shown to impact *C. auris* biofilm

formation when overexpressed and artificially hyper-activated through HA-tagging of the putative

transcription factor activation domain[31]. Under biofilm conditions, 14 genes orthologous to those

known to regulate biofilm formation in *C. albicans* were up-regulated, notably again including the

C. auris orthologs of *BRG1.* These findings demonstrate that *RBA1* significantly impacts

regulation of *C. auris* gene expression during biofilm formation and suggest that *RBA1* may play

a central regulatory role in this clinically impactful process.

Loss of *RBA1* enhances *C. auris* adhesion to plastics and human keratinocytes

A key clinical characteristic of *C. auris* is its ability to adhere to and persist on a variety of

surfaces in the nosocomial environment, including medical equipment and patient skin. As our

transcriptional profiling revealed loss of *RBA1* enhanced the expression of an assortment of

C. auris adhesin-encoding genes, we hypothesized that loss of *RBA1* would enhance *C. auris*

adhesion to multiple substrates. To test this, we first evaluated the adherence of our *Δrba1* and

Δrba1::*RBA1*⁺ strains, as well as the isogenic clinical isolates SKU013 and SKU018, to

polystyrene plastic under a variety of conditions. In RPMI media, SKU018 and the *Δrba1* strains

exhibited greater adhesion than SKU013 to both polystyrene and tissue culture-treated

polystyrene (**Fig. 5a, b**). This difference was magnified when the polystyrene well plates were

pre-treated with human plasma, with or without the addition of 10% human serum to the media

(**Fig. 5c, d**). By contrast, *Δrba1*::*RBA1*⁺ control strains phenocopied the parental SKU013 under

all conditions.

Next, we evaluated whether loss of *RBA1* promotes adhesion of *C. auris* to human

keratinocytes. *Δrba1* strains were observed to exhibit enhanced adherence to a confluent culture

of human N/TERT keratinocytes, while neither *Δrba1*::*RBA1*⁺ controls strains nor SKU018 differed

from the SKU013 control (**Fig. 5e** (top)**, f**). Outbreak isolates harboring the *RBA1*^Q174* (SKU055),

RBA1^Q225* (SKU043), and *RBA1*^C303S (SKU056) mutations were similarly found to exhibit

enhanced adhesion to keratinocytes as compared to their closest available genetic relatives (**Fig.**

5e (bot)**, g**) further supporting the connection between loss of *RBA1* and enhanced adherence to

human keratinocytes. Taken together, these results confirm that loss of *RBA1* significantly

enhances *C. auris* adhesion characteristics under a variety of conditions.

Loss of *RBA1* increases fungal burden during catheter-associated urinary tract infections.

As loss of *RBA1* increased *C. auris* adhesion, biofilm formation, and the expression of the

previously characterized virulence factors *SCF1*, *IFF4109*, and *ALS4112/ALS5*, we next sought

to evaluate the impact of *RBA1* on *C. auris* biofilm-related infection *in vivo.* To accomplish this,

we utilized our mouse model of CAUTI [25, 32], where immunocompetent mice were catheterized

and the bladder of each mouse was challenged with *C. auris* transurethrally 24 hours prior to

sacrifice for evaluation of fungal burden and systemic dissemination of infection. Importantly, this

mimics a clinical scenario where *C. auris* is often cultured from patients with urinary catheters and

is relevant to multiple patient populations at increased risk of *C. auris* colonization and infection.

In this model, the *Δrba1* strain was found to exhibit a 6.8-fold increase in bladder tissue fungal

burden (p= 0.0435) by 24 hours post-infection, as compared to SKU013 (**Fig. 5h**). Moreover, a

trend toward increased dissemination of the *Δrba1* strain to kidney and spleen tissues was also

observed, and one mouse infected with the *Δrba1* strain was found to have detectable fungal

burden in cardiac tissue. These findings suggest that loss of *RBA1* may also enhance virulence

of *C. auris* during clinical infections such as CAUTI.

Mutations in *RBA1* have emerged during numerous *C. auris* clinical outbreaks

271   We next hypothesized that *RBA1* mutations have also likely occurred during other *C. auris*

outbreaks as well. To test this, we leveraged publicly available *C. auris* WGS data collected for

infection control and outbreak tracking to identify mutations in the *RBA1* gene. Ninety-four

mutations in *RBA1* were identified, and 310 of 2,181 evaluable isolates were found to carry one

or more of 58 identified non-synonymous or nonsense *RBA1* mutations, including *C. auris* isolates

representing genetic clades I, II, and IV. The majority of the identified non-synonymous/nonsense

mutations are considered as likely loss-of-function mutations (40; 61.5%) (**Fig. 6a**), similar to the

mutations we initially identified from the Kuwait outbreak collection. Evaluating meta data and

publications associated with identified isolates revealed 10 non-synonymous *RBA1* mutations

originating from documented outbreaks of *C. auris* infection on four continents and were caused

by isolates representing both genetic clades I and IV (**Fig. 6b**) [10, 16, 33, 34, 35, 36]. Notably, this included

two instances where *RBA1* mutations arose during Indian outbreaks[37] initially reported in 2013,

as well as two instances of *RBA1* mutations occurring during ongoing outbreaks in the United

States[34, 38]. These findings clearly demonstrate convergent mutations in *RBA1* across multiple of

C. auris outbreaks and further suggest that these mutations confer a competitive advantage to *C.*

auris in the healthcare environment.

DISCUSSION

289   *C. auris* is frequently associated with clinical outbreaks of infection and multi-drug

resistance thus posing a challenge to clinicians reminiscent of the healthcare-associated bacterial

pathogens, *Acinetobacter baumannii* and *Enterococcus faecium*, [3, 8]. Unlike other agents of

invasive candidiasis, *C. auris* is not considered as part of the human normal flora and is not known

to be widely distributed in environmental reservoirs. Instead, *C. auris* infections are currently

thought to be acquired predominantly through dissemination of recently emergent drug-resistant,

and possibly infection-adapted, lineages within medical environments, leading to the infection of

vulnerable patient populations[3, 16]. Thus, as with other healthcare-associated pathogens, the

identification of *C. auris* evolution during clinical outbreaks and characterization of traits promoting

the persistence among human patients is essential to both preventing infections and mitigating

prolonged outbreaks of infection.

300   Here, we analyzed samples from outbreaks in Kuwait to identify genomic and

microbiological elements which may promote transmission and persistence of *C. auris* in clinical

settings. Rather than distinct clustering of serial isolates from individual patient cases,

phylogenetic analysis identified isolate clusters originating from multiple patients indicating

multiple instances of pathogen lateral transmission consistent with clinical dissemination.

Interestingly, while numerous cases of isolates acquiring mutations associated with echinocandin

and AmB resistance were identified, persistence and spread of these multi-drug-resistant lineages

was not observed in this study. Conversely, among both phylogenetic clusters B and C, closely

related isolates harboring acquired *RBA1* mutations were cultured from multiple patients over

periods of time spanning many months. While there was no association between clinical

antifungal susceptibility and *RBA1* genotype, *C. auris* isolates carrying clinically acquired loss-of-

function mutations in *RBA1* were observed to exhibit enhanced biofilm formation as compared to

those with wildtype *RBA1* sequences.

Biofilms are critically important to human public health as they promote microbial survival

in a variety of environments, demonstrating increased resilience to disinfectants and

antimicrobials, and contribute to pathogen dissemination and the establishment of new infections.

In fact, the majority of human infections involve biofilms either within human tissues or on the

surface of in-dwelling medical devices such as catheters or prosthetic implants [39]. As numerous

clinical isolates from the studied clinical outbreaks in Kuwait were isolated from the urine of

catheterized patients or the respiratory secretions of patients with endotracheal tubes, we

hypothesized that the identified loss-of-function mutations in *C. auris RBA1* represented a

clinically emergent adaptation promoting pathogen success in the medical environment through

enhanced biofilm formation. Genetic interrogation of the impact of *C. auris RBA1* definitively

demonstrated loss of this gene enhanced biofilm formation when evaluated using *in vitro* models

mimicking clinically relevant conditions including those specific to CAUTI. This enhancement of

biofilm formation may represent a selective pressure which promoted the emergence and

persistence of *RBA1* mutant lineages during the studied clinical outbreaks.

Subsequent transcriptional profiling revealed a narrow role in *C. auris* cells growing under

planktonic conditions, and a much wider influence on gene expression when cells were grown in

biofilms. Under both conditions, however, *Δrba1* strains exhibited increased expression of a

multitude of cell surface adhesins, including multiple genes we have previously shown to

significantly impact *C. auris* virulence during infection. The two previously characterized *C. auris*-

specific adhesins, *SCF1* and *ALS4112/ALS5,* were substantially elevated in the *Δrba1* strain

grown in planktonic conditions, with both genes exceeding a 7-$\log_2$-fold increase over the SKU013

parental control. As these adhesins are known to be critical to the ability of *C. auris* to adhere to

a variety of substrates including human skin[26, 27], this suggested that *RBA1* may impact the

propensity of *C. auris* to both colonize patients and establish new biofilms. Further investigation

of gene expression in biofilms demonstrated a wide variety of genes predicted to encode *C. auris*

cell surface proteins, cell wall constituents, secreted enzymes, and transcriptional regulators were

differentially expressed in the *Δrba1* strain. Intriguingly, this also included the *C. auris* orthologs

of *C. albicans ENG1* and *YWP1*, two secreted beta-glucanases which are known to play critical

roles in cell wall beta-glucan masking[28, 29, 30]. This finding, combined with the altered expression

of a multitude of GPI-anchored cell surface proteins and the differences in extracellular matrix

production observed by scanning electron microscopy, suggests that the loss of *RBA1* may also

facilitate evasion of *C. auris* detection from host defenses during infection.

We next examined whether loss of *RBA1* and subsequent increased expression of *SCF1*

and *ALS4112/ALS5* would impact the ability of *C. auris* to adhere to surfaces where *SCF1* and

ALS4112/ALS5 play a significant role[26, 27]. Strains lacking *RBA1* were found to exhibit enhanced

adhesion to human keratinocytes as well as polystyrene surfaces both with and without human

plasma coatings, which mimics the deposition of host factors such as fibrinogen on medical device

surfaces. Moreover, evaluation of three *C. auris* clinical isolates from the Kuwait outbreaks

harboring three additional loss-of-function mutations, as compared to their respective closest

available genetic relative with a *RBA1*[WT] allele, again demonstrated loss of *RBA1* to increase

adhesion of *C. auris* to human keratinocyte. This finding is of substantial clinical relevance, as it

supports that loss of *RBA1* can promote the dissemination of *C. auris* and colonization of patients

during a clinical outbreak. Further, patients colonized with *C. auris* strains harboring *RBA1* loss-

of-function mutations may pose a greater risk of transmission within the hospital environment.

We also examined the impact of *RBA1 in vivo* using a murine CAUTI model which

replicates a biofilm-related infection of a site where *C. auris* is commonly isolated. Contrary to

other commonly utilized models of pathogenic yeast infection, this model uses mice that have not

been immunosuppressed. Instead, immunocompetent mice are made vulnerable to *C. auris*

infection through the implantation of a silicone catheter, similar to urinary catheters commonly

used among human patients. In this model, the *Δrba1* strain was found to develop a significantly

greater fungal burden in the mouse bladder, and a trend towards greater dissemination of the

Δrba1 strain from the urinary tract to other tissues including the heart was observed. These

findings further suggest loss of *RBA1* is advantageous during *C. auris* clinical infection. This is

clinically significant as CAUTIs often lead to urosepsis with a ~30% 7-day mortality rate [15, 40].

Since the urinary tract is a source of ~25% of the all the sepsis cases, patients infected with

C. auris strains harboring *RBA1* mutation may be predisposed to severe clinical complication.

Finally, leveraging the growing body of publicly available WGS data generated through

ongoing *C. auris* infection control and prevention efforts, we were able to investigate whether

mutations in *RBA1* have emerged during other clinical outbreaks. This approach uncovered

dozens of distinct *RBA1* mutations, impacting hundreds of clinical isolates and more than 10% of

evaluable samples. *RBA1* mutations were identified among isolates representing three of the

major genetic subgroups of *C. auris,* but were notably absent among isolates from clade III*,*

suggesting either clade-specific impacts of *RBA1* mutations or a potential bias within samples

currently available in the public data set. Like the four *RBA1* mutations identified from the Kuwait

outbreaks, the mutations identified from the global collection were predominantly predicted loss-

of-function mutations such as those encoding early stop codons, frameshift mutations, or altering

highly conserved residues in the ZCF DNA-binding domain. Identified isolates with *RBA1*

mutations were found to date back to some of the earliest documented clinical outbreaks of

C. auris, including those in India, England, and Columbia. Additionally, isolates from outbreak

lineages actively disseminating in the United States were also found to harbor *RBA1* mutations.

Taken together, these findings clearly demonstrate that mutations in *RBA1* have emerged

repeatedly during clinical outbreaks of *C. auris* infection and strongly support the notion that the

enhancement in adhesion and biofilm formation conferred by these mutations provided a

competitive advantage within the medical environment in numerous previous outbreaks.

In conclusion, these results call attention to the capacity of *C. auris* to evolve and enhance

traits beyond drug resistance during outbreaks of infection by highlighting the unique nature of

C. auris among other fungal pathogens and identifying mutations in *RBA1* as a potential indicator

of outbreak-adapted lineages. As traits conferred by *RBA1* mutations are advantageous to

C. auris spread and persistence, and as we have identified numerous instances where these

mutations have arisen during global outbreaks of infection, the identification and tracking of

lineages carrying *RBA1* mutations may be merited as part of infection prevention efforts.

MATERIALS AND METHODS

Ethics Statement

C. auris isolates were cultured from various clinical specimens of hospitalized patients in Kuwait. The need for informed consent was waived by the Health Sciences Center Ethics Committee (Approval letter Ref: VDR/EC/3724 dated January 31, 2021) as the clinical specimens were collected as part of routine diagnostic work-up for the isolation of bacterial/fungal pathogens for suspected infections. The results are reported here on deidentified samples without revealing patient identity. All animal care was consistent with the Guide for the Care and Use of Laboratory Animals from the National Research Council. The University of Notre Dame Institutional Animal Care and Use Committee approved all mouse infections and procedures as part of protocol number 25-01-9009. For urine collection, all donors signed an informed consent form, and protocols were approved by the Institutional Review Board of the University of Notre Dame under study #19-04-5273.

C. auris Isolates

Seventy *C. auris* isolates from previously described multi-institution outbreaks of *C. auris* infections in Kuwait, were obtained by the Mycology Reference Laboratory Department of Microbiology, Kuwait University, Kuwait [41]. All isolates (including names used in publication of their original description) and strains used in this study are listed in **Table S1**. All *C. auris* strains were stored in 40% glycerol stocks at -80°C and cultured at 35°C in liquid YPD (1% yeast extract, 2% peptone, 2% glucose) media before plating on Sabouraud dextrose (SD) agar (BD Difco) media to ensure standardized growth for phenotypic testing.

Plasmid and strain construction

EPIC-mediated gene editing was used to generate *RBA1* (B8441v3: B9J08_03209) modifications in *C. auris* clinical isolate SKU013 (also known as Kw1432/17). Duplexed guide

oligos (Integrated DNA Technologies) (**Table S2**) targeting disruption or reversion of

B9J08_03209 were added to the pJMR19 vector using restriction enzyme cloning[22]. Plasmids

were maintained in *E. coli* DH5α cells cultured in Luria-Bertani broth supplemented with 5mg/L

ampicillin and isolated using QIAprep® Spin Minipreps (Qiagen). Repair templates for *RBA1*-

disruption and reversion were amplified from a gBlock sequence (Integrated DNA Technologies)

and SKU013 gDNA, respectively, via PCR and purified with QIAquick PCR Purification Kit

(Qiagen). All oligos, plasmids, and gBlocks used in this study are listed in **Table S2.** *C. auris*

Lithium acetate transformations were then performed as previously described [22] with positive

selection on 200μg/mL nourseothricin-supplemented YPD agar. *RBA1* editing was confirmed by

both Sanger sequencing at the Hartwell Center at St. Jude Children's Research Hospital (SJCRH)

and with long read nanopore sequencing by Plasmidsaurus.

Whole genome sequencing

Genomic DNA (gDNA) was extracted from 1mL of overnight liquid YPD cultures using

MasterPure™ Yeast DNA Purification kits (Biosearch Technologies) according to manufacturer

instructions. Sample libraries were prepared from gDNA by the Hartwell Center (SJCRH) using

KAPA HyperPrep (Roche) and sequencing targeting 100x coverage for paired 150bp reads was

performed via Illumina NovaSeq. Paired-reads FASTQ files were analyzed using CLC genomics

workbench v24.0 (QIAGEN) software. Low quality reads were trimmed with Trim Reads v2.8 and

subsequently mapped to the B8441v3 genome (GCA002759435.3) according to CLC-workbench

recommendations for match scores, mismatch, insertion, and deletion costs. Local Realignment

v2.6 was used to refine unaligned ends and mutations were identified within annotated coding

sequences via Fixed Ploidy Variant Detection v2.6 with ploidy=1.

Phylogenetic analysis was initiated with Create SNP Tree v1.9 using the Maximum

Likelihood Phylogeny v1.4 algorithm. The resulting SNP phylogenetic tree was visualized from

Neighbor Joining construction methods specifying Jukes Cantor nucleotide substitution modeling

with estimated topology and bootstrapping for 100 replicates. Sequences were deposited in

National Center for Biotechnology Information Sequence Read Archive.

Biofilm Production Assays

To assess biofilm production in the presence of human plasma, 96-well flat-bottomed

plates (VWR, 10861-562) were coated with 100 μL pooled human plasma (Innovative Research

Inc, IPLAWBNAC50ML) diluted 1:5 in carbonate-bicarbonate buffer (Sigma C3041-50CAP) and

incubated overnight at 4°C. Colonies from overnight SD agar plates were used to generate

standardized OD_{600} = 0.5 cell suspensions[42] in RPMI-Biofilm media, consisting of RPMI medium

1640 (Gibco) + 3-(4-Morpholino)propane sulfonic acid (MOPS, Fisher Bioreagent) + 2% dextrose,

buffered to pH 7 and supplemented with 10% heat-inactivated/male AB human serum (HuS,

Sigma-Aldrich). Plasma coatings were gently removed the plates and 200 μL of the cell

suspensions were added to the wells and plates were sealed with a breathable (EASYSEAL)

membrane. After 90-minutes at 37°C, 250 RPM, plates were gently washed sterile PBS to remove

non-adhered cells. Fresh 200 μL of RPMI-Biofilm media was then added back to each well and

plates were incubated for 24 hrs at 37°C, 250 RPM. For quantification of biofilm production,

supernatants were removed and wells were gently washed twice with PBS. Biofilms were stained

with 0.4% crystal violet as previously described[42]. Crystal violet absorbance measured at 595nm

on an Agilent BioTek Synergy H1 Microplate Reader. Average media only negative controls were

subtracted from the sample reads and values were normalized to SKU013.

To replicate biofilm growth replicating the urinary tract environment, 96-well flat-bottomed

plates (VWR, 10861-562) were coated with 100 µl of Fg (450 µg/ml) and incubated overnight at

4°C.[25] The various strains were grown overnight, shaking in YPD, and the inoculum was

normalized to ~1 × 10^6 CFUs/ml. Cultures were then diluted (1:100) into human urine

supplemented with 10% heat inactivated human serum and incubated in the wells of the 96-well

plate at 37°C for 48 hours. Human urine from at least two healthy female donors between the

ages of 20 to 35 were collected and pooled. Donors did not have a history of kidney disease,

diabetes, or recent antibiotic treatment. Urine was sterilized with a 0.22-µm filter (VWR 29186-

212), pH was normalized to 6.0 to 6.5, and urine was used immediately for the assays. Following

biofilm formation, supernatants were removed, and plates were incubated in 200 µl of 0.5% crystal

violet for 15 min. Crystal violet stain was removed, and the plate was washed with water to remove

the remaining stain. Plates were dried and then incubated with 200 µl of 33% acetic acid for 15

min. In another plate, 100 µL of the acetic acid solution was transferred, and absorbance values

were measured via a plate spectrophotometer at 595 nm (Molecular Devices SpectraMax ABS).

Values were then normalized to SKU013.

RNA Extraction

Normalized cell suspensions (OD_{600} = 0.1) from SD replicate platings were diluted ~1:50

in 10 mL RPMI-Growth media, prepared from RPMI medium 1640 (Gibco) + MOPS (Fisher

Bioreagent) buffered to pH 7 and grown overnight at 35°C, 220 RPM to an OD_{600} = 2.0 $\pm$ 0.3.

Concentrated cultures were diluted in >20mL RPMI-Growth media to an OD_{600} = 0.2 then split

into two 10mL aliquots for direct comparison of planktonic growth and biofilm development.

Planktonic cells were grown over 4 hours in 50mL conical at 35°C, 220RPM and pelleted at 4000

RPM. For biofilm development cells were transferred to T75 flask pre-coated with 20% human

plasma in carbonate-bicarbonate buffer prepared as described above. Following 2 hr incubation

at 37°C, non-adhered cells were gently removed and fresh 10mL RPMI-Growth media was added

for an additional 4 hr incubation at 37°C. Biofilms were gently washed once with PBS and biofilms

were then removed by scraping in the presence of 10mL sterile PBS and pipette transfer to a

50mL conical tube for centrifuge pelleting. All pellets were flash frozen in liquid nitrogen and stored

at -80ºC. RNA was isolated using the Invitrogen RiboPure™ Yeast kit (ThermoFisher Scientific™,

USA) per manufacturer's instructions.

RNA Sequencing

The Hartwell Center (SJCRH) prepared the RNA libraries using Illumina chemistries per

manufacturer's instructions and sequenced the RNA on an Illumina NextSeq (Illumina Inc., USA)

using the total stranded RNA-Seq application for paired end runs targeting 150bp reads.

Sequenced reads were analyzed with CLC Genomics Workbench version 24.0.1 (QIAGEN).

Imported reads were trimmed using default settings for failed reads and adaptor sequences and

then subsequently mapped to the *C. auris* B8441 reference genome (GenBank accession:

GCA_002759435.3) with paired reads counted as one and expression values set to RPKM.

Principal component analysis was utilized for the initial clustering assessment of biological

replicates and mismatch, insertion, and deletion costs were set to default parameters for RNA-

Seq Analysis 2.7. Differential expression was calculated using negative binomial GLM modeling

with whole transcriptome RNA-Seq defaults for normalization with Trimmed Mean of M values,

using Differential gene expression v1.4 (CLC genomics workbench, Qiagen). Comparisons were

made to the wildtype *SKU013* clinical isolate for each condition. Differential expression was

defined as fold changes ≥ 2 or ≤ -2 (FDR p-value ≤0.05)

Scanning Electron Microscopy

Sterile silicon chips were loaded into 24-well plate and coated overnight with 0.5mL 20%

human plasma in carbonate-bicarbonate buffer. Suspensions of SKU013, *Δrba1*::*RBA1*⁺_a , and

Δrba1_a were cultured in RPMI-Biofilm media as described above for biofilm production assays

with 0.5mL of cells loaded into the wells. Following 24 hour incubation, wells were gently washed

with 1mL PBS and biofilms were fixed with 0.1M sodium cacodylate buffer at pH 7.4 with 0.15%

alcian blue, 1% tannic acid, 2.5% glutaraldehyde, 2% paraformaldehyde for 2 hours and stored

at 4°C. Preparation, staining, and imaging the fixed biofilms was performed by the Electron

Microscopy Division of the Cell and Tissue Imaging Center (CTIC-EM) at SJCRH for SEM Fixed

biofilms were washed with 0.1M sodium cacodylate buffer + 0.15% alcian blue prior to osmium

tetroxide staining in the presence of 0.15% alcian blue. Imaging was performed on a Zeiss

GeminiSEM 460. Scale bars were embedded using ImageJ[43].

Polystyrene adhesion assay

Adhesion to flat-bottom 96-well non-tissue culture treated polystyrene plates (CS100-

701311, NEST SCIENTIFIC) was assessed in RPMI-Growth+ media, defined as RPMI medium

1640 + MOPS + 2% dextrose, buffered to pH 7. Adhesion to tissue culture treated plates

(TP92696, TPP Midwest Scientific), was assessed for RPMI-Growth+ media and RPMI-Biofilm

media in the presence and absence of human plasma, coated as described above. cells cultured

and plates were loaded as described for biofilm production assays. Following a 90-minute

incubation at 37°C, 250RPM, wells were gently washed once with PBS to remove non-adhered

cells. Adhered cells were stained with 100μL of a 10:1 solution of XTT sodium salt and phenazine

methosulfate (VWR International LLC). Absorbances at OD_{492} and OD_{630} were read after 60

minutes and values were normalized to SKU013.

Keratinocyte adhesion assay

Adhesion of *C. auris* to human keratinocytes was determined as previously described[26].

Human N/TERT keratinocytes[44] were grown in 96-well plates to complete confluence in

keratinocyte-SFM medium (Gibco, 10724-011) supplemented with 30 μg/ml bovine pituitary

extract, 0.2 ng/ml epidermal growth factor, and 0.31 mM $CaCl_2$. Overnight *C. auris* cultures were

diluted to an OD_{600} of 0.03, and 100 μl of this suspension was added to each well containing

keratinocytes. In parallel, an equal volume of culture was added to empty wells, centrifuged (210

g, 1 min), and imaged to determine input cell counts, while plates containing keratinocytes were

centrifuged under the same conditions and incubated at 37 °C with 5% CO_2 for 1 h. Non-adherent

yeast cells were then removed by washing three times with PBS. The remaining cells were fixed

in 4% formaldehyde and stained with FITC-conjugated anti-*Candida* antibody (Meridian

Bioscience, B65411F; 1:1000). Keratinocyte nuclei and cell bodies were labeled with Hoechst

(Thermo Fisher, 62249; 1:1000) and CellMask Deep Red (Thermo Fisher, H32721; 1:5000),

respectively.

Fluorescent images were acquired at 10X magnification using a BioTek Lionheart FX

microscope. The number of adherent *C. auris* cells was quantified using a customized CellProfiler

analysis pipeline[26, 45]. Numbers of input cells were determined from brightfield images pre-

processed in Fiji/ImageJ (v2.14.0)[43] using a custom macro for edge-based segmentation and

binary mask generation. The proportion of adherent to input cells was calculated to assess the

adhesion efficiency of each *C. auris* strain.

Catheter-Associated Urinary Tract Infection (CAUTI) Modeling

Female C57BL/6 mice (Jackson Laboratory) were subjected to transurethral implantation

of a silicone catheter followed by transurethral inoculation of the bladder lumen with ~5 × 10^5

colony forming units (CFU) in PBS, as previously described [32]. Mice were euthanized at 24 hours

post-infection by cervical dislocation after anesthesia inhalation and the catheter, bladder,

kidneys, spleen, and heart were aseptically harvested. Bladders were weighed and then

homogenized for CFUs. The other organs were homogenized, and catheters were cut into small

pieces before sonication for fungal CFU enumeration.

Statistics and Reproducibility

Statistical parameters and applied analysis methods including exact sample numbers (n)

and definitions for representative central value and spread (mean ± SEM, CFU quantifications;

representative modal value (n=3), minimum inhibitory concentrations) are annotated in

corresponding figure legends. Data from at least three experiments were pooled for each

adhesion and biofilm assay and twice for each *in vivo* mouse assay. One-way ANOVA tests, two-

tailed Welch's t-tests, and two-tailed Mann-Whitney *U* tests were performed with GraphPad Prism

10 software (GraphPad Software) for comparisons of described in-vitro assays and in vivo CAUTI

experiments, respectively.

For RNA sequencing, statistical analysis was performed using CLC Genomics Workbench

version 20.0 (QIAGEN) with paired reads counted as one and expression values set to RPKM.

Whole transcriptome differential gene expression analysis was performed with mismatch,

insertion, and deletion costs set to default parameters, and a Wald test was used to compare

SKU013 against Δ*rba1_a*. Differential expression determined by a fold change of ≥ 2 or ≤ -2 with

a False discovery rate (FDR) corrected p value ≤0.05.

ACKNOWLEDGMENTS

This work was supported in part by the St. Jude Children's Research Hospital Children's Infection Defense Center and Center for Infectious Diseases Research awards granted to JMR, by the National Institutes of Health (NIH)/ National Institute of Allergy and Infectious Diseases (NIAID) grant U19-AI181767 awarded to TRP, and by the NIH/ NIAID F32 AI181164 awarded to GZ. This work was supported by institutional funds from the University of Notre Dame (to FHST and ALFM), and by grants from the Good Venture Foundation (Open Philanthropy) (to ALFM), from the National Institutes of Health R01AI177875 and R21AI171742 (to FHST), and R01DK128805 (to ALFM). This research included experiments conducted by the Hartwell Center for Bioinformatics & Biotechnology, the St. Jude Center for Advanced Genome Engineering and the Cell and Tissue Imaging Center - Electron Microscopy (CTIC-EM) which are supported in part by ALSAC and the National Cancer Institute at the National Institutes of Health [P30 CA021765]. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTIBUTIONS

Conceptualization, J.M.R., T.R.O., F.H.S.T., and A.L.F.M.; methodology, J.M.R., L.A.D., A.A.L., S.J.J., Q.Z., T.R.O., and A.L.F.M.; Investigation, L.A.D., A.A.L., S.J.J., Q.Z., Q.J.C., and J.M.R.; writing—original draft, L.A.D. and J.M.R.; writing—review & editing, L.A.D., A.A.L., S.J.J., G.Z., M.A., Q.J.C., S.A., E.M., I.A., W.A., K.A., T.R.O., F.H.S.T., A.L.F.M., and J.M.R.; funding acquisition J.M.R., T.R.O., F.H.S.T., A.L.F.M., and Q.Z.; supervision, T.R.O., F.H.S.T., A.L.F.M., and J.M.R..

REFERENCES

1. World Health Organization. *WHO fungal priority pathogens list to guide research, development and public health action*. (World Health Organization, Geneva, 2022).

2. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol Immunol* **53**, 41-44 (2009).

3. Lockhart SR*, et al.* Simultaneous Emergence of Multidrug-Resistant *Candida auris* on 3 Continents Confirmed by Whole-Genome Sequencing and Epidemiological Analyses. *Clin Infect Dis* **64**, 134-140 (2017).

4. European Centre for Disease Prevention and Control. *Survey on the epidemiological situation, laboratory capacity and preparedness for Candidozyma (Candida) auris, 2024*. (Publications Office of the European Union, 2025).

5. Sabino R, Veríssimo C, Pereira Á A, Antunes F. *Candida auris*, an Agent of Hospital-Associated Outbreaks: Which Challenging Issues Do We Need to Have in Mind? *Microorganisms* **8**, (2020).

6. Baker AD, Gold JAW, Forsberg K, Jones S, Lyman MM. Progression from *Candida auris* Colonization Screening to Clinical Case Status, United States, 2016-2023. *Emerg Infect Dis* **31**, 1613-1617 (2025).

7. Akinbobola AB, Kean R, Hanifi SMA, Quilliam RS. Environmental reservoirs of the drug-resistant pathogenic yeast *Candida auris*. *PLoS Pathog* **19**, e1011268 (2023).

8. Proctor DM*, et al.* Clonal *Candida auris* and ESKAPE pathogens on the skin of residents of nursing homes. *Nature* **639**, 1016-1023 (2025).

9. Pacilli M*, et al.* Regional Emergence of *Candida auris* in Chicago and Lessons Learned From Intensive Follow-up at 1 Ventilator-Capable Skilled Nursing Facility. *Clin Infect Dis* **71**, e718-e725 (2020).

10. Chow NA*, et al.* Multiple introductions and subsequent transmission of multidrug-resistant *Candida auris* in the USA: a molecular epidemiological survey. *The Lancet Infectious Diseases* **18**, 1377-1384 (2018).

11. Codda G*, et al.* In vivo evolution to echinocandin resistance and increasing clonal heterogeneity in *Candida auris* during a difficult-to-control hospital outbreak, Italy, 2019 to 2022. *Euro Surveill* **28**, (2023).

652 12. Rybak JM*, et al.* In vivo emergence of high-level resistance during treatment reveals the
653   first identified mechanism of amphotericin B resistance in *Candida auris*. *Clin Microbiol*
654   *Infect* **28**, 838-843 (2022).

655
656 13. Carolus H*, et al.* Genome-Wide Analysis of Experimentally Evolved *Candida auris* Reveals
657   Multiple Novel Mechanisms of Multidrug Resistance. *mBio* **12**, (2021).

658
659 14. Rybak JM*, et al.* Mutations in *TAC1B*: a Novel Genetic Determinant of Clinical Fluconazole
660   Resistance in *Candida auris*. *mBio* **11**, (2020).

661
662 15. La Bella AA, Santiago-Tirado FH, Flores-Mireles AL. *Candida auris* is emerging as a
663   prevalent urinary pathogen. *PLoS Pathog* **21**, e1013138 (2025).

664
665 16. Chow NA*, et al.* Tracing the Evolutionary History and Global Expansion of *Candida auris*
666   Using Population Genomic Analyses. *mBio* **11**, (2020).

667
668 17. Muñoz JF*, et al.* Clade-specific chromosomal rearrangements and loss of subtelomeric
669   adhesins in *Candida auris*. *Genetics* **218**, (2021).

670
671 18. Barker KS*, et al.* Relative contributions of the *ERG11*[(VF125AL)] and *MRR1A*[(N647T)] mutations
672   to fluconazole resistance in Clade III *Candidozyma (Candida) auris* clinical isolates. *Clin*
673   *Microbiol Infect*, (2025).

674
675 19. Rybak JM, Sharma C, Doorley LA, Barker KS, Palmer GE, Rogers PD. Delineation of the
676   Direct Contribution of *Candida auris ERG11* Mutations to Clinical Triazole Resistance.
677   *Microbiol Spectr* **9**, e0158521 (2021).

678
679 20. Mayr EM, Ramírez-Zavala B, Krüger I, Morschhäuser J. A Zinc Cluster Transcription
680   Factor Contributes to the Intrinsic Fluconazole Resistance of *Candida auris*. *mSphere* **5**,
681   (2020).

682
683 21. Massic L*, et al.* Acquired amphotericin B resistance attributed to a mutated *ERG3* in
684   *Candidozyma auris*. *Antimicrob Agents Chemother* **69**, e0060125 (2025).

685
686 22. Doorley LA, Meza-Perez V, Jones SJ, Rybak JM. A *Candidozyma (Candida) auris*-
687   Optimized Episomal Plasmid-Induced Cas9-Editing System Reveals the Direct Impact of
688   the S639F-Encoding *FKS1* Mutation. *J Infect Dis* **232**, e529-e536 (2025).

689
690 23. Spruijtenburg B*, et al.* Whole genome sequencing analysis demonstrates therapy-induced
691   echinocandin resistance in *Candida auris* isolates. *Mycoses* **66**, 1079-1086 (2023).

693 24. Asadzadeh M*, et al.* Molecular characterisation of *Candida auris* isolates from
694 immunocompromised patients in a tertiary-care hospital in Kuwait reveals a novel mutation
695 in *FKS1* conferring reduced susceptibility to echinocandins. *Mycoses* **65**, 331-343 (2022).

696
697 25. La Bella AA*, et al.* The catheterized bladder environment promotes Efg1- and Als1-
698 dependent *Candida albicans* infection. *Science Advances* **9**, eade7689 (2023).

699
700 26. Zhao G*, et al.* Adhesin Als4112 promotes *Candida auris* skin colonization through
701 interactions with keratinocytes and extracellular matrix proteins. *Nat Commun* **16**, 5673
702 (2025).

703
704 27. Santana DJ*, et al.* A *Candida auris*-specific adhesin, Scf1, governs surface association,
705 colonization, and virulence. *Science* **381**, 1461-1467 (2023).

706
707 28. de Assis LJ*, et al.* Nature of β-1,3-Glucan-Exposing Features on *Candida albicans* Cell
708 Wall and Their Modulation. *mBio* **13**, e0260522 (2022).

709
710 29. Hopke A, Nicke N, Hidu EE, Degani G, Popolo L, Wheeler RT. Neutrophil Attack Triggers
711 Extracellular Trap-Dependent *Candida* Cell Wall Remodeling and Altered Immune
712 Recognition. *PLoS Pathog* **12**, e1005644 (2016).

713
714 30. Yang M*, et al.* Control of β-glucan exposure by the endo-1,3-glucanase Eng1 in *Candida*
715 *albicans* modulates virulence. *PLoS Pathog* **18**, e1010192 (2022).

716
717 31. Louvet M*, et al.* Ume6-dependent pathways of morphogenesis and biofilm formation in
718 Candida auris. *Microbiol Spectr* **12**, e0153124 (2024).

719
720 32. Conover MS, Flores-Mireles AL, Hibbing ME, Dodson K, Hultgren SJ. Establishment and
721 Characterization of UTI and CAUTI in a Mouse Model. *J Vis Exp*, e52892 (2015).

722
723 33. Escandón P*, et al.* Notes from the Field: Surveillance for *Candida auris* - Colombia,
724 September 2016-May 2017. *MMWR Morb Mortal Wkly Rep* **67**, 459-460 (2018).

725
726 34. Barbian HJ*, et al. Candida auris* Outbreak and Epidemiologic Response in Burn Intensive
727 Care Unit, Illinois, USA, 2021-2023. *Emerg Infect Dis* **31**, 438-447 (2025).

728
729 35. Rhodes J*, et al.* Genomic epidemiology of the UK outbreak of the emerging human fungal
730 pathogen *Candida auris*. *Emerg Microbes Infect* **7**, 43 (2018).

731
732 36. Araúz AB*, et al.* Isolation of *Candida auris* from 9 patients in Central America: Importance
733 of accurate diagnosis and susceptibility testing. *Mycoses* **61**, 44-47 (2018).

735   37.   Chowdhary A, *et al.* New clonal strain of Candida auris, Delhi, India. *Emerg Infect Dis* **19**,
736   1670-1673 (2013).

737
738   38.   Gorzalski A, *et al.* The use of whole-genome sequencing and development of
739   bioinformatics to monitor overlapping outbreaks of Candida auris in southern Nevada.
740   *Front Public Health* **11**, 1198189 (2023).

741
742   39.   Mishra A, Aggarwal A, Khan F. Medical Device-Associated Infections Caused by Biofilm-
743   Forming Microbial Pathogens and Controlling Strategies. *Antibiotics (Basel)* **13**, (2024).

744
745   40.   Molina JJ, Flores-Mireles AL. CAUTIon - not all UTIs are the same. *Nat Rev Urol* **22**, 799-
746   814 (2025).

747
748   41.   Ahmad S, Khan Z, Al-Sweih N, Alfouzan W, Joseph L. *Candida auris* in various hospitals
749   across Kuwait and their susceptibility and molecular basis of resistance to antifungal
750   drugs. *Mycoses* **63**, 104-112 (2020).

751
752   42.   Gulati M, *et al.* In Vitro Culturing and Screening of *Candida albicans* Biofilms. *Curr Protoc*
753   *Microbiol* **50**, e60 (2018).

754
755   43.   Schindelin J, *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
756   *Methods* **9**, 676-682 (2012).

757
758   44.   Dickson MA, *et al.* Human keratinocytes that express hTERT and also bypass a
759   p16(INK4a)-enforced mechanism that limits life span become immortal yet retain normal
760   growth and differentiation characteristics. *Mol Cell Biol* **20**, 1436-1447 (2000).

761
762   45.   Lamprecht MR, Sabatini DM, Carpenter AE. CellProfiler: free, versatile software for
763   automated biological image analysis. *Biotechniques* **42**, 71-75 (2007).

764
765   46.   Khan Z, Ahmad S, Al-Sweih N, Joseph L, Alfouzan W, Asadzadeh M. Increasing
766   prevalence, molecular characterization and antifungal drug susceptibility of serial *Candida*
767   *auris* isolates in Kuwait. *PLoS One* **13**, e0195743 (2018).

768
769   47.   Al-Obaid I, *et al.* Fatal Breakthrough Candidemia in an Immunocompromised Patient in
770   Kuwait Due to *Candida auris* Exhibiting Reduced Susceptibility to Echinocandins and
771   Carrying a Novel Mutation in Hotspot-1 of FKS1. *J Fungi (Basel)* **8**, (2022).

772
773   48.   Alfouzan W, *et al.* Molecular Epidemiology of *Candida auris* Outbreak in a Major
774   Secondary-Care Hospital in Kuwait. *J Fungi (Basel)* **6**, (2020).

   FIGURES

778
779
780   **Fig 1. Clinical emergence and spread of *C. auris* isolates harboring mutations in *RBA1* during an**
781   **outbreak of infections.**
782   Phylogenetic analysis of 70 isolates from documented outbreaks of *C. auris* infections in Kuwait. Nodes
783   indicate branch length termination. Patient cases, culture sources, and *RBA1* mutations are indicated on
784   the right.
785

	Culture Source	SNPs vs SKU013	*RBA1*	Susceptibility		
				FLU	**AmB**	**Echino**
Patient 51						
SKU013	Endotracheal asp.	-	WT	R	S	S
SKU014	Endotracheal asp.	2	WT	R	S	S
SKU015	Endotracheal asp.	4	WT	R	S	S
SKU016	Endotracheal asp.	4	WT	R	S	S
SKU017	Endotracheal asp.	4	WT	R	S	S
SKU018	Urine	8	E47*	R	S	R
SKU020	Blood	5	WT	R	S	S
Patient 53						
SKU022	Endotracheal asp.	6	E47*	R	S	S
SKU023	Endotracheal asp.	6	E47*	R	S	S
SKU026	Tracheal sec.	7	E47*	R	S	S
SKU031	Urine	7	E47*	R	R	S
SKU032	Urine	89	WT	R	S	S
SKU033	Urine	8	E47*	R	R	S
SKU035	Urine	8	E47*	R	S	R

786
787
Fig 2. *RBA1* mutations emerge during patient infection and are associated with increased biofilm
production. a, Timeline of patient cases 51 and 53 with *C. auris* isolations and antifungal treatment courses
indicated, AMB: liposomal amphotericin B, CSF: caspofungin. Isolate numbers (SKU) with *RBA1*[E47*]
mutations indicated by green background. **b,** Biofilm formation on sidewall of polystyrene (PS) tubes
following overnight growth in RPMI culture media. Arrow indicates visible biofilm ring. **c,** Isolate details
including name, culture source, number of coding region single nucleotide polymorphisms (SNPs) relative
to the earliest isolate SKU013, *RBA1* genotype, and interpretation of susceptibility to fluconazole (FLU),
amphotericin B (AmB), and echinocandins (Echino) based on current CDC tentative breakpoints. **d, e,**
Biofilm production of isolates formed over 48 hours on fibrinogen-treated PS in urine media supplemented
with 10% serum. Biofilm assessment via crystal violet staining with absorbance read at 595 nm. Reads
were normalized to the initial isolate SKU013 with means (bars) and standard error of the means plotted in
GraphPad Prism. (**d**) gray-blue bars indicate wildtype *RBA1* containing isolates and green indicates
RBA1[E47*] mutation. Clinical isolates harboring (**e**) *RBA1*[Q174*], *RBA1*[Q225*], and *RBA1*[C303S] mutations are
indicated with light blue, red and orange bars, respectively, and plotted alongside a corresponding closest
genetic relative harboring a wildtype *RBA1* (gray-blue bars). Kruskal-Wallis statistical one-way ANOVA
testing with Dunn's correction for multiple comparisons showed significant differences between SKU013
and *RBA1*[E47*] containing isolates (**d**; ****P = <0.0001) and *RBA1*[Q174*], *RBA1*[Q225*], and *RBA1*[C303S] isolates
and corresponding closest genetic relative (**e**; ***P= <0.001).

806
Fig 3. Loss of *RBA1* enhances biofilm production
a,b, Biofilm production of *C. auris RBA1* strains formed on (**a**) fibrinogen coated microtiter plates in human
urine media supplemented with 10% human serum after 48hrs and (**b**) on human plasma coated microtiter
plates in RPMI tissue culture media supplemented with 10% human serum after 24hrs. Biofilms stained
with crystal violet and absorbance measured at 595 nm. Absorbance reads were normalized to SKU013
(*RBA1*^{WT}) and graphed (GraphPad Prism) showing mean (bars) and standard error of the mean. Kruskal-
Wallis statistical testing with Dunn's correction for multiple comparisons showed significant differences
between *Δrba1_a, Δrba1_b*, and SKU018 (*RBA1*^{E47*}) when compared to SKU013 (*RBA1*^{WT}) (****P=
<0.0001). **c, d,** Representative SEM images for (**c**) SKU013 (*RBA1*^{WT}) and (**d**) *Δrba1* biofilms formed over
24 hours in RPMI media supplemented with 10% human serum on silicone disks treated with human
plasma. Dried samples coated with 5nm iridium and imaged in the Zeiss Gemini 460 (CITC-SJCRH) using
Secondary Electrons detector at 2kV, 77pA, and 5mm working distance.
819

Fig 4. Loss of *RBA1* increases expression of *C. auris* adhesins and virulence factors under both planktonic and biofilm conditions.

a,b Volcano plot of differentially expressed genes between clinical isolate SKU013 and *Δrba1* strain during exponential growth phase when cultured under (a) planktonic growth conditions (10mL RPMI media in 50mL conical tube incubated at 35°C with 220RPM) and (**b**) biofilm-forming conditions (10mL RPMI media in plasma-treated T75 flasks for static incubation at 37°C). Blue, purple, and red highlighted dots represent cell wall associated, ZCF, and cell aggregation or biofilm formation encoding genes, respectively. Dotted lines represent lower limit equivalents for fold changes of ≥ |1.5| on the x-axis, and a false discovery rate (FDR) p-value ≤0.05 on the y-axis. **c**, Venn diagram comparing differentially expressed genes (≥ |1.5| on the x-axis, and a false discovery rate p-value ≤0.05) exhibiting up-regulation or down-regulation in the *Δrba1* strain under planktonic or biofilm growth conditions. **d**, Differential gene expression encoding predicted GPI-anchored proteins, cell surface adhesions, fungal cell wall associated proteins, aspartic-type endopeptidases, and genes orthologous to transcriptional regulators of biofilm formation in *Candida albicans* with planktonic expression on the left and biofilm expression on the right for each column. *C. albicans* genes are listed in gray with true orthologs bolded. Significant differential expression (fold change ≥ |1.5|; FDR p-value ≤ 0.05) is indicated with a black dot.

838
Fig 5. Loss of *RBA1* increases adhesion and fungal burden in murine CAUTI model.
a, b, c, d, Adhesion of *C. auris* isolates and *RBA1* strains grown in RPMI culture media after 90 minutes to
(**a**) untreated polystyrene (PS), (**b**) tissue culture (TC) -treated PS, (**c**) human plasma coated, TC-treated
PS, and (**d**) human plasma coated, TC-treated PS grown in RPMI supplemented with 10% human serum.
Adhesion assessed by absorbance at 490 nm following XTT treatment. Statistical significance was
determined using Kruskal-Wallis one-way ANOVA tests followed by a Dunn's test for multiple comparisons
to the *RBA1*^WT control SKU013 (*$P \le 0.05$; ****$P < 0.0001$). **e,** Representative immunofluorescence images
showing adherence of *C. auris* cells to human N/TERT keratinocytes. *C. auris* was stained with FITC-
conjugated anti-*Candida* antibody (yellow) and keratinocytes were stained using CellMask Deep Red
(magenta) with a Hoechst counterstain for nuclei identification (blue). **f, g,** Quantification of *C. auris* isolates
strains adhered to human N/TERT keratinocytes as measured by the proportion of attached cells remaining
after washing. Bars represent mean ± standard deviation for three biological replicates. Statistically
significant adhesion increases were determined using unpaired two-tailed Welch's t-tests for (**e**) *Δrba1*
compared to SKU013 (*$P \le 0.05$) and (**g**) SKU055 (*RBA1*^Q174*), SKU043 (*RBA1*^Q225*), and SKU056
(*RBA1*^C303S) compared to *RBA1*^WT isolates, SKU021, SKU034, and SKU037, respectively (*$P \le 0.05$;
***$P \le 0.001$). **i,** Fungal burden of SKU013 and the derivative *Δrba1* strain obtained from the catheter material
or bladder, kidney, spleen, and heart tissues after infection in a murine model of catheter associated urinary
tract infection (CAUTI). Two-tailed Mann-Whitney tests showed significant differences in fungal burden
recovered from bladder tissue (*$P \le 0.05$, n=8 mice per group). CFU, colony forming units; LOD, lower limit
of detection.

860
Fig 6. Identification of *RBA1* mutations emerging during numerous outbreaks of *C. auris* infections
worldwide.
a, Diagram depicting non-synonymous mutations in *RBA1* identified by analyzing publicly available *C. auris*
whole genome sequencing data. Green lines on top of figure indicate early stop codons and frameshift
mutations. Blue lines on bottom of figure indicate missense mutations. N241H and *354Q were identified
among all clinical isolates classified as clade IV. DBD: DNA-binding domain. **b,** World map indicating
locations of documented outbreaks of *C. auris* infections where mutations in *RBA1* were identified from
publicly available data. Green fills indicate isolates representing genetic Clade I and blue fills indicate
isolates representing genetic clade IV.

 SUPPLEMENTAL DATA

 Patient 51 and 53 Case Descriptions:

Patient 51 was a 54-year-old male with a past medical history significant for multiple

myeloma admitted to the intensive care unit (ICU) of Hospital F, a tertiary care center associated

with Kuwait Cancer Control Center in Kuwait which cares for immunocompromised patients

including cancer patients, burn patients and transplant recipients diagnosed and requiring

treatment for infectious diseases. Patient 51 was receiving a 14-day course of empiric

caspofungin treatment when *C. auris* isolate SKU013 ($RBA1^{WT}$; wildtype-*RBA1*) was first

identified from a culture of their endotracheal tube and considered to represent colonization

(**Fig. 2a**). Four successive isolates (**Fig. 2c**) were identified from endotracheal tube cultures, and

the presence of respiratory symptoms prompted a 21-day course of empiric caspofungin

treatment. During this treatment, isolate SKU018 ($RBA1^{E47*}$) was obtained from the patient's

urinary catheter. SKU018 notably exhibited *in vitro* resistance to echinocandins ($FKS1^{S639F}$),

although it was still thought to represent likely colonization. SKU020 was obtained from a blood

culture the day before patient 51 expired. SKU020 also exhibited elevated *in vitro* echinocandin

resistance and differed from the earlier respiratory tract isolates SKU016 and SKU017 by only

one coding region SNP ($FKS1^{R1354H}$), indicating likely dissemination from the respiratory tract.

Patient 53 was a 33-year-old female in the ICU of the same tertiary care hospital (Hospital

F) with a significant past medical history for Hodgkin's lymphoma and advanced stage nodular

sclerosis. Approximately one month after receiving a 21-day empiric course of liposomal

amphotericin B (AmB) for suspected respiratory infection, the patient's respiratory condition

worsened and *C. auris* (SKU022) was isolated from the patient's endotracheal tube. Notably,

isolate SKU022 harbored the same $RBA1^{E47*}$ mutation present in the isolate previously cultured

from the urine of patient 51 (SKU018), and genetically these two isolates differed by four coding

region SNPs, suggestive of pathogen lateral transmission. Following completion of a 14-day

course of caspofungin treatment, two subsequent *C. auris* isolates (both $RBA1^{E47*}$) were cultured

from endotracheal tube and tracheal aspirate cultures, prompting a second 14-day course of

liposomal AmB. After this second course of AmB, all cultures from respiratory sources remained

negative. However, one month later the patient was suspected to have developed a CAUTI and

urine cultures yielded isolate SKU031 ($RBA1^{E47*}$). A second 14-day course of caspofungin was

initiated after SKU031 was found to be AmB-resistant. Subsequently, AmB-susceptible *C. auris*

(SKU032) was cultured from the patient's urine and a third course of liposomal AmB was given.

During this third course of AmB, two additional isolates were obtained from urine cultures,

SKU033 ($RBA1^{E47*}$) and SKU035 ($RBA1^{E47*}$), and exhibited decreased susceptibility to AmB and

echinocandins, respectively. Patient 53 was then transferred to another care facility, precluding

further follow-up.

Supplemental Table 1. *Candidozyma auris* isolates and strains

CLINICAL ISOLATES

SKU_ID	Mycology Reference Laboratory ID	Patient ID	Subclade	Isolation source	Previous citations
SKU001	Kw1983-16	47	Ic	Tissue	[46]
SKU002	Kw2031-16	47	Ic	Ab. drain fluid	[46]
SKU003	Kw2236-16	47	Ic	Pus	[46]
SKU004	Kw2254-16	47	Ic	Tissue	[46, 24]
SKU005	Kw2442-16	30	Ic	Urine	[23, 46]
SKU006	Kw2515-16	30	Ic	Tracheal sec.	[23, 46]
SKU007	Kw2555-16	30	Ic	Urine	[23, 46]
SKU008	Kw369-17	30	Ic	Tracheal sec.	[23, 46]
SKU009	Kw1138-17	30	Ic	Urine	[23, 46]
SKU010	Kw1239-17	30	Ic	Urine	[23, 46]
SKU011	Kw1308-17	30	Ic	Urine	[23, 46]
SKU012	Kw1388-17	30	Ic	Tracheal sec.	[23, 46]
SKU013	Kw1432-17	51	Ib	ETT	[46, 24, 23]
SKU014	Kw1695-17	51	Ib	Tracheal aspirate	[46, 24]
SKU015	Kw1825-17	51	Ib	Tracheal aspirate	[46, 24]
SKU016	Kw1856-17	51	Ib	Tracheal aspirate	[46, 24]
SKU017	Kw1893-17	51	Ib	ETT	[46, 24, 23]
SKU018	Kw1994-17	51	Ib	Urine	[46, 24, 23]
SKU019	Kw2040-17	49	Ib	Vaginal swab	[46, 24]
SKU020	Kw2074-17	51	Ib	Blood	[46, 24, 23]
SKU021	Kw2260-17	52	Ib	Blood	[46, 24, 23]
SKU022	Kw2501-17	53	Ib	ET aspirate	[46, 24, 23]
SKU023	Kw2647-17	53	Ib	ET aspirate	[46, 24]
SKU024	Kw2713-17	52	Ib	Urine	[46, 24, 23]
SKU025	Kw2742-17	52	Ib	Urine	[24, 23]
SKU026	Kw2871-17	53	Ib	Tracheal sec.	[24]
SKU027	Kw2968-17	57	Ib	Urine	[24]
SKU028	Kw2989-17	57	Ib	ETT	[24]
SKU029	Kw3092-17	57	Ib	Catheter tip	[24, 23]
SKU030	Kw3134-17	57	Ib	Urine	[24, 23]
SKU031	Kw3135-17	53	Ib	Urine	[24, 23]
SKU032	Kw3367-17	53	Ib	Urine	[24]
SKU033	Kw3424-17	53	Ib	Urine	[24, 23]
SKU034	Kw3506-17	59	Ic	Urine	[47]
SKU035	Kw3521-17	53	Ib	Urine	[24, 23]
SKU036	Kw3525-17	59	Ic	Tracheal sec.	[47]

Strain ID	Background				Citations
SKU037	Kw3584-17	59	Ic	Urine	[47]
SKU038	Kw55-18	65	Ib	Sputum	[24, 23]
SKU039	Kw60-18	59	Ic	Tracheal sec.	[23, 47]
SKU040	Kw87-18	59	Ic	Blood	[23, 47]
SKU041	Kw93-18	59	Ic	Tracheal sec.	[23, 47]
SKU042	Kw108-18	59	Ic	Tracheal sec.	[23, 47]
SKU043	Kw137-18	64	Ic	Blood	[41]
SKU044	Kw147-18	65	Ib	Blood	[24, 23]
SKU045	Kw257-18	68	Ic	Urine	[41]
SKU046	Kw334-18	65	Ib	Oral swab	[24, 23]
SKU047	Kw532-18	71	Ic	Urine	[41]
SKU048	Kw1014-18	79	Ic	Blood	[41]
SKU049	Kw1148-18	80	Ic	Urine	[41]
SKU050	Kw1201-18	77	Ic	Bronchial washing	[41]
SKU051	Kw1252-18	81	Ic	Blood	[41]
SKU052	Kw1517-18	68	Ic	Urine	[41]
SKU053	Kw1610-18	84	Ic	Urine	[41]
SKU054	Kw1674-18	89	Ib	Urine	[24]
SKU055	Kw1753-18	89	Ib	Blood	[24]
SKU056	Kw2103-18	92	Ic	Urine	[41]
SKU057	Kw2201-18	100	Ib	Urine	[41]
SKU057	Kw2201-18	100	Ib	Urine	[41]
SKU058	Kw2272-18	102	Ib	Blood	[41]
SKU059	Kw2427-18	103	Ib	Blood	[48]
SKU060	Kw2466-18	105	Ic	Blood	[41]
SKU061	Kw2518-18	101	Ib	Urine	[41]
SKU062	Kw2570-18	100	Ib	Urine	[41]
SKU063	Kw2717-18	104	Ic	Tracheal sec.	[41]
SKU064	Kw2821-18	100	Ib	Urine	[41]
SKU065	Kw2885-18	100	Ib	Blood	[41]
SKU066	Kw2898-18	113	Ic	Urine	[41]
SKU067	Kw2999-18	113	Ic	Urine	[41]
SKU068	Kw3014-18	118	Ib	Urine	[41]
SKU069	Kw3144-18	125	Ic	Blood	[41]
SKU070	Kw3204-18	123	Ic	Groin swab	[41]

Strain ID	Background	RBA1 Genotype			Citations
Δrba1_a	SKU013	B9J08_003209:c.950CAGT>CTTAT			This study
Δrba1_b	SKU013	B9J08_003209:c.950CAGT>CTTAT			This study
Δrba1::RBA1+_a	*Δrba1_b*	*RBA1*WT			This study
Δrba1::RBA1+_b	*Δrba1_a*	*RBA1*WT			This study

909 **Supplemental Table 2. Oligos and Primers**

Name	Sequence	Notes
RBA1_RT_AMP-F	CGCCAATCCATGTAACCCTC	Amplify Repair Template sequence: *Δrba1_RT-gBlock* for *Δrba1;* SKU013 genomic DNA for *RBA1+*
RBA1_RT_AMP-R	GTTGTTGTTGGATATGAGGTGG	
Δrba1_gRNA_TOP	CCACAGCACGGGGTGAGCCAGTC	Guide sequences inserted into pJMR19 for *Δrba1* in SKU013
Δrba1_gRNA_BOT	AACGACTGGCTCACCCCGTGCTG	
RBA1+_gRNA_TOP	CCACTGGACTCGAGCCCCAGATA	Guide sequences inserted into pJMR19 for *RBA1+*in *Δrba1*
RBA1+_gRNA_BOT	AACTATCTGGGGCTCGAGTCCAG	
*RBA1*_scn_F	CCGACCCTTGTTCCCTCCTC	PCR Amplification of *RBA1* for Screening
*RBA1*_scn_R	GTGGGAGACCGGCGTTG	
RBA1_SeqF	CCTACCTCCTCCATGCCC	Sanger Sequencing Primer
Δrba1_RT-gBlock	TTCGGTGTCTGAGTTGATTGTGGGCGCC CAGCTCTCGTCCCACCTCGACCGCTCTC CTAACCACATGACAGTGAAGCACCTGGA CTCGAGCCCCAGATAAGGCTCACCCCGT GCTGCCTCACCTGGAAAAAAGCACTTTT GCCTCCTCGGTGCCGCCGCCTCTGGCG CCCTCGTTACATCCACCCGTGGCGTTTC C	Repair template gBlock for *Δrba1* in SKU013

910
911

912
Supplementary Figure 1. Matrix of identified single nucleotide polymorphisms (SNP) in
coding regions of genome. *C. auris* isolate Genomes aligned to reference genome
GCA002759435.3 (clade I, B8441_v3) for SNP identifications. SNP calling performed with fixed
ploidy variant detection in CLC genomics workbench (Qiagen). 46,231 SNPs identified among
the annotated open reading frame with identification of 46,055 SNPs leading to amino acid
changes across all isolates.
919

920
921 **SUPPLEMENTAL REFERENCES**
922

923 1.   Khan Z, Ahmad S, Al-Sweih N, Joseph L, Alfouzan W, Asadzadeh M. Increasing
924   prevalence, molecular characterization and antifungal drug susceptibility of serial *Candida*
925   *auris* isolates in Kuwait. *PLoS One* **13**, e0195743 (2018).

926
927 2.   Asadzadeh M*, et al.* Molecular characterisation of *Candida auris* isolates from
928   immunocompromised patients in a tertiary-care hospital in Kuwait reveals a novel mutation
929   in *FKS1* conferring reduced susceptibility to echinocandins. *Mycoses* **65**, 331-343 (2022).

930
931 3.   Spruijtenburg B*, et al.* Whole genome sequencing analysis demonstrates therapy-induced
932   echinocandin resistance in *Candida auris* isolates. *Mycoses* **66**, 1079-1086 (2023).

933
934 4.   Al-Obaid I*, et al.* Fatal Breakthrough Candidemia in an Immunocompromised Patient in
935   Kuwait Due to *Candida auris* Exhibiting Reduced Susceptibility to Echinocandins and
936   Carrying a Novel Mutation in Hotspot-1 of *FKS1*. *J Fungi (Basel)* **8**, (2022).

937
938 5.   Ahmad S, Khan Z, Al-Sweih N, Alfouzan W, Joseph L. *Candida auris* in various hospitals
939   across Kuwait and their susceptibility and molecular basis of resistance to antifungal
940   drugs. *Mycoses* **63**, 104-112 (2020).

941
942 6.   Alfouzan W*, et al.* Molecular Epidemiology of *Candida auris* Outbreak in a Major
943   Secondary-Care Hospital in Kuwait. *J Fungi (Basel)* **6**, (2020).