Emergence and spread of outbreak-adapted *Candidozyma auris* harboring mutations in *RBA1*

Laura A. Doorley[1], Alyssa A. La Bella[2], Sarah J. Jones[1], Guolei Zhao[3], Mohammad Asadzadeh[4], Quanita J. Choudhury[1], Suhail Ahmad[4], Eiman Mokaddas[4,5], Inaam Al-Obaid[6], Wadha Alfouzan[4,7], Khalifa Al-Banwan[4,8], Teresa O'Meara[3] Felipe H. Santiago Tirado[2], Ana L. Flores Mireles[2], Jeffrey M. Rybak[1*]

[1] Department of Pharmacy and Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN 38105, USA.
[2] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556, USA
[3] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI 48109, USA.
[4] Department of Microbiology, Faculty of Medicine, Kuwait University, Jabriya 46300, Kuwait.
[5] Department of Microbiology, Ibne-Sina Hospital, Shuwaikh, Kuwait
[6] Department of Microbiology, Al-Sabah Hospital, Shuwaikh, Kuwait
[7] Department of Microbiology, Farwaniya Hospital, Farwaniya, Kuwait
[8] Department of Microbiology, Amiri Hospital, Kuwait City, Kuwait

[*] Corresponding author: Jeffrey.Rybak@stjude.org

Jeffrey M. Rybak, PharmD, PhD
Assistant Member
Department of Pharmacy and Pharmaceutical Sciences
St. Jude Children's Research Hospital
262 Danny Thomas Place
Chili's Care Center, I5304, Mail Stop #313
Memphis TN, 38105

Office: 901.595.2368
Fax: 901.595.3125

ABSTRACT

Candidozyma auris (formerly *Candida auris*) is an emerging healthcare-associated fungal

pathogen, increasingly isolated from clinical outbreaks of infection, and with a high propensity to

colonize and persist on patients and within the medical environment. Here, we leveraged a large

collection of *C. auris* isolates from multi-facility clinical outbreaks within the same urban area to

identify genomic patho-adaptations that promote persistence and dissemination in the healthcare

environment. Genomic and phylogenetic analyses revealed loss-of-function mutations in the

uncharacterized *C. auris* transcription factor gene *RBA1* to have independently emerged multiple

times within these outbreaks. We provide evidence that loss of *RBA1* results in increased *C. auris*

adhesion to plastic and human keratinocytes, enhances biofilm formation in multiple conditions,

and exacerbates fungal burden in a mouse model of catheter-associated urinary tract infection.

Finally, we uncover that loss-of-function mutations in *RBA1* have independently emerged during

multiple outbreaks of *C. auris* infections across the globe, and that *RBA1* mutant lineages are

present among isolates collected from large ongoing clinical outbreaks. Taken together, the

results of this study reveal loss-of-function mutations in the *C. auris RBA1* as a novel and clinically

relevant genetic determinant of enhanced outbreak characteristics and virulence factor-gene

expression.

INTRODUCTION

Candidozyma auris (formerly *Candida auris*) is an emerging pathogenic yeast first identified in 2009 and later recognized by the World Health Organization as a critical threat to global public health in 2022[1, 2]. Initially isolated from a limited number of individual patient infections, *C. auris* was soon revealed as the cause of multiple nosocomial outbreaks of invasive fungal infections across the globe[3]. Now, *C. auris* outbreaks have been reported from more than 50 countries, representing thousands of documented infections annually, including 4,514 cases in 2023 in the United States alone[4, 5, 6]. When appropriately treated, outcomes of *C. auris* infections are similar to those previously associated with invasive candidiasis, with crude mortality rates ranging from 30 to 75%, depending on affected patient population and severity of disease[7, 8]. However, timely and effective treatment of *C. auris* infections is often complicated both by challenges in the clinical identification of *C. auris* using conventional techniques, and by high rates of resistance to commonly employed antifungal drugs[3].

Further distinguishing *C. auris* from other fungal pathogens, *C. auris* is not known to be ubiquitous within common natural environmental reservoirs nor is it a member of the normal human microbiota[9]. *C. auris* infections are, instead, healthcare-associated and typically occur following a horizontal transfer to vulnerable patients within hospital settings[10]. Multiple studies have demonstrated that once introduced into the medical environment, *C. auris* rapidly spreads among both patients and medical surfaces, acting as an invasive species in these healthcare settings. As a result, clinical *C. auris* infections routinely occur in series or as clustered clinical outbreaks, sometimes spanning multiple facilities or institutions within an affected healthcare network[11, 12]. Further, while *C. auris* isolates exhibiting decreased susceptibility to the polyene and echinocandin classes of antifungals were initially associated with acquisition of resistance-conferring mutations during treatment of individual patients, dissemination of antifungal-resistant and treatment-adapted lineages has now been repeatedly documented[13, 14, 15].

As with other healthcare-associated and multi-drug-resistant pathogens, the

characterization and tracking of emergent microbial lineages with traits impacting clinical

outcomes is paramount to infection prevention and treatment. Prior study of *C. auris* evolution

during infection and exposure to clinically relevant stressors has largely focused on the

emergence of antifungal resistance[15, 16, 17, 18]. We and others have shown, through both *in vitro*

evolution experiments and by studying sets of isogenic clinical isolates, that *C. auris* is capable

of rapidly adapting to antifungal-induced stress and that multiple independent drug-resistant

lineages can emerge after only a single antifungal exposure[15, 16, 17, 18]. However, investigations of

patho-adaptation in the clinical setting or during outbreaks of infections remains unexplored.

In this work, we leveraged a collection of 70 *C. auris* clinical isolates from multi-institutional

outbreaks of infection to identify novel *C. auris* genetic markers and link them to traits driving

persistence and outbreak potential in clinical settings. We identified multiple independent lineages

of *C. auris* harboring distinct mutations in a previously uncharacterized fungal transcription factor,

here named, Repressor of Biofilm and Adhesion 1 (*RBA1*). Genetic and transcriptomic

interrogation revealed loss of *RBA1* to result in enhanced biofilm formation, increased adhesion

to multiple surfaces including human keratinocytes, and significantly elevated expression of

C. auris adhesins and virulence factors. Moreover, loss of *RBA1* was found to significantly

increase fungal burden in a mouse model of catheter-associated urinary tract infection (CAUTI),

an infection characteristically involving biofilms in urine where *C. auris* is often isolated[19]. Finally,

analysis of publicly available whole genome sequencing data revealed numerous distinct *RBA1*

mutations present in hundreds of clinical isolates, demonstrating consistent and convergent loss-

of-function mutations from isolates from well-documented and ongoing clinical outbreaks of

C. auris infections across the globe. Collectively these findings demonstrate *RBA1* encodes a

repressor of *C. auris* biofilm and adhesion and implicate mutations in *RBA1* as clinically relevant

adaptations promoting *C. auris* survival during healthcare-associated outbreaks of infection.

RESULTS

Clinical emergence of multiple *C. auris* outbreak lineages carrying *RBA1* mutations.

WGS analysis revealed all *C. auris* isolates belonged to Clade I, one of three major genetic clades previously linked to healthcare-associated outbreaks[20]. This included 34 and 36 isolates from the previously described subclades Ib and Ic, respectively[21]. A total of 155,135 SNPs were identified among the 70 included isolates as compared to the Clade I reference genome (B8441v3), with 46,231 predicted to impact coding regions. As anticipated from characteristically fluconazole-resistant Clade I isolates, all isolates were found to harbor one or more mutations in the genes *ERG11*, *TAC1B*, and/or *MRR1*, which we and others have previously shown to directly contribute to fluconazole resistance[18, 22, 23, 24]. Additionally, 16 isolates were found to possess mutations known to contribute to echinocandin resistance (altering *FKS1*) and 4 isolates possessed mutations in genes associated with amphotericin B resistance (*ERG3*, 1 isolate; *ERG6*, 3 isolates)[15, 25, 26].

Phylogenetic analysis revealed the population structure of this collection to consist of 69 isolates grouping into three major clusters, with 35, 24, and 10 isolates grouping into clusters A, B, and C, respectively. The remaining single isolate (SKU043) did not group into a larger cluster and was more distantly related (**Fig. 1**). Within clusters, isolates were found to be largely clonal, with an average of 5.3 to 12.6 coding region SNPs separating each isolate (**Fig. S1**), while SKU043 differed from all other isolates by 43 to 113 coding region SNPs. Clusters A and C each contained isolates originating from multiple healthcare facilities within Kuwait, while cluster B originated entirely from a single tertiary care hospital. Notably, multiple instances were identified where isolates from different patients and even different hospitals were found to be more genetically similar than serial isolates originating from the same patient case, as might be expected in the setting of a clinical outbreak.

To investigate potential patho-adaptations occurring during outbreaks of *C. auris* infections, we further evaluated the WGS data to identify genes where mutations were acquired

on multiple occasions and/or mutations emerged and persisted among the outbreak population.

Intriguingly, this revealed 19 isolates harboring mutations in a previously uncharacterized *C. auris*

gene B9J08_03209 (identified as B9J08_002136 in B8441v2 and hereafter named *RBA1*), which

is predicted to encode a fungal-specific zinc-cluster transcription factor. Although initially identified

as an ortholog of *WOR2* in *C. albicans,* the proteins are not syntenic, with only 22% sequence

identity, primarily located in the DNA-binding domain. The 19 isolates with mutations in *RBA1*

included representatives from all three major phylogenetic clusters, including all 10 isolates in

cluster C, as well as the non-clustered isolate SKU043. Four different mutations in *RBA1* were

identified, all considered likely loss-of-function mutations, with three encoding an early stop codon

(upstream of the predicted DNA-binding domain) and one altering a highly conserved cysteine

residue in the $Zn(II)_2Cys_6$ DNA-binding motif. The early stop codon mutation, $RBA1^{E47*}$, was found

to have emerged and persisted among two intensive care unit patients over the course of eight

months, these cases and the associated 13 cluster B isolates were further evaluated to identify

the potential clinical impact of *RBA1* mutations.

Loss of *RBA1* enhances *C. auris* biofilm formation.

A subset of isolates from patient cases 51 and 53 have been previously characterized with

consideration to antifungal susceptibility while investigating the emergence of echinocandin and

amphotericin B resistance-conferring mutations, respectively[15, 27, 28]. In brief, patient 51 was a 54-

year-old male with a past medical history significant for multiple myeloma admitted to the intensive

care unit (ICU) of Hospital F, a tertiary care center associated with Kuwait Cancer Control Center

in Kuwait which cares for immunocompromised patients including cancer patients, burn patients

and transplant recipients diagnosed and requiring treatment for infectious diseases. Patient 51

was receiving a 14-day course of empiric caspofungin treatment when *C. auris* isolate SKU013

($RBA1^{WT}$; wildtype-*RBA1*) was first identified from a culture of their endotracheal tube and

considered to represent colonization (**Fig. 2a**). Four successive isolates (**Table 1**) were identified

from endotracheal tube cultures, and the presence of respiratory symptoms prompted a 21-day

course of empiric caspofungin treatment. During this treatment, isolate SKU018 ($RBA1^{E47*}$) was

obtained from the patient's urinary catheter. SKU018 notably exhibited *in vitro* resistance to

echinocandins ($FKS1^{S639F}$), although it was still thought to represent likely colonization. SKU020

was obtained from a blood culture the day before patient 51 expired. SKU020 also exhibited

elevated *in vitro* echinocandin resistance and differed from the earlier respiratory tract isolates

SKU016 and SKU017 by only one coding region SNP ($FKS1^{R1354H}$), indicating likely dissemination

from the respiratory tract.

Patient 53 was a 33-year-old female in the ICU of the same tertiary care hospital (Hospital

F) with a significant past medical history for Hodgkin's lymphoma and advanced stage nodular

sclerosis. Approximately one month after receiving a 21-day empiric course of liposomal

amphotericin B (AmB) for suspected respiratory infection, the patient's respiratory condition

worsened and *C. auris* (SKU022) was isolated from the patient's endotracheal tube. Notably,

isolate SKU022 harbored the same $RBA1^{E47*}$ mutation present in the isolate previously cultured

from the urine of patient 51 (SKU018), and genetically these two isolates differed by four coding

region SNPs, suggestive of pathogen lateral transmission. Following completion of a 14-day

course of caspofungin treatment, two subsequent *C. auris* isolates (both $RBA1^{E47*}$) were cultured

from endotracheal tube and tracheal aspirate cultures, prompting a second 14-day course of

liposomal AmB. After this second course of AmB, all cultures from respiratory sources remained

negative. However, one month later the patient was suspected to have developed a CAUTI and

urine cultures yielded isolate SKU031 ($RBA1^{E47*}$). A second 14-day course of caspofungin was

initiated after SKU031 was found to be AmB-resistant. Subsequently, AmB-susceptible *C. auris*

(SKU032) was cultured from the patient's urine and a third course of liposomal AmB was given.

During this third course of AmB, two additional isolates were obtained from urine cultures,

SKU033 ($RBA1^{E47*}$) and SKU035 ($RBA1^{E47*}$), and exhibited decreased susceptibility to AmB and

echinocandins, respectively. Patient 53 was then transferred to another care facility, precluding

further follow-up.

Of the 14 isolates cultured from patient 51 and patient 53, 13 were found to be closely

related (separated by 2 to 8 SNPs within coding regions, **Fig. S1**) and grouped into cluster B.

Seven of these isolates, starting with SKU018 and including all but one isolate later cultured from

patient 53, encoded the early stop codon mutation in *RBA1.* While there was no association

between antifungal susceptibility and *RBA1* genotype, all seven *RBA1* mutant isolates were

observed to exhibit increased biofilm formation under standard laboratory conditions (**Fig. 2b**). As

the earliest isolate where the $RBA1^{E47*}$ mutation was identified (SKU018) originated from culturing

the urinary tract of patient 51, and multiple subsequent isolates with this mutation were isolated

from the urinary tract of patient 53, we next used our fibrinogen-urine biofilm model [29] that utilizes

serum-supplemented urine and fibrinogen-coated surfaces to evaluate biofilm formation in

conditions mimicking the bladder during CAUTI. In this model, isolates carrying the $RBA1^{E47*}$

mutation exhibited enhanced biofilm formation as compared to the earliest isolate SKU013

(approximately 2 to 6-fold higher), while biofilm formation was relatively unchanged among closely

related $RBA1^{WT}$ isolates (**Fig. 2c**). To evaluate if other clinically emergent predicted loss-of-

function mutations in *RBA1* had a similar effect on biofilm formation, we also compared the biofilm

formation of isolates harboring the $RBA1^{Q174*}$ (SKU055), $RBA1^{Q225*}$ (SKU043), and $RBA1^{C303S}$

(SKU056) mutations to their closest available genetic relatives (SKU021, SKU034, and SKU037,

respectively) using the same model. In each case, biofilm formation was observed to be

significantly higher in the isolates harboring mutations in *RBA1* (**Fig. 2d**), further bolstering the

connection between loss of *RBA1* and biofilm formation.

As we observed a strong association between *RBA1* genotype and biofilm formation, we

hypothesized that loss-of-function mutations in *RBA1* are a genetic determinant of enhanced

C. auris biofilm formation. To test this hypothesis, we next generated the *RBA1*-knockout (*Δrba1*)

and *RBA1*-revertant (*Δrba1*::$RBA1^+$) control strains in the SKU013 clinical isolate background and

evaluated biofilm formation. In the fibrinogen-urine model, *Δrba1* strains were found to exhibit

significantly increased biofilm formation (3.9 to 4.8-fold greater than SKU013; p<0.0001). This

enhancement of biofilm formation resembled the isogenic clinical isolate SKU018 which carries

the *RBA1*^E47* mutation (3.3-fold greater than SKU013; p<0.0001) (**Fig. 3a**). Importantly,

restoration of the *RBA1* gene in *Δrba1*::*RBA1*⁺ control strains returned biofilm formation to levels

comparable to SKU013.

We further investigated the impact of *RBA1* using a plasma-RPMI biofilm model that

utilized serum-supplemented RPMI culture media and human plasma-coated surfaces to evaluate

biofilm production that could apply more generally to clinical settings. Once again, *Δrba1* strains

were found to produce more robust biofilms (6.2 to 6.8-fold greater than SKU013; p<0.0001),

while *Δrba1*::*RBA1*⁺ control strains closely matched the biofilm formation of SKU013 (**Fig. 3b**).

Examination of the plasma-RPMI biofilms using scanning electron microscopy, showed scattered

patches of biofilm predominantly consisting of one to two layers of cells for both SKU013 and

Δrba1::*RBA1*⁺ (**Fig. 4a**). By comparison, *Δrba1* strains formed more confluent biofilms of variable

but generally greater thickness with greater production of extracellular matrix. (**Fig. 4b**).

Loss of *RBA1* increases expression of *C. auris* adhesins and virulence factors.

As *RBA1* is predicted to encode a transcription factor and our earlier experiments had

shown loss of *RBA1* significantly enhances biofilm formation, we next evaluated the impact of

RBA1 on the *C. auris* transcriptome during both planktonic and biofilm growth conditions using

our *Δrba1* strain and the parental *RBA1*^WT isolate SKU013. When cells were grown to exponential

growth phase in RPMI tissue culture media under planktonic conditions, transcriptional profiling

by RNAseq revealed loss of *RBA1* to have a modest impact, and only 55 of 5,594 *C. auris* genes

(0.98%) were differentially expressed (≥|1.5| fold-change and false discovery rate [FDR] ≤0.05)

(**Fig. 5a**). Of these, 42 genes were upregulated in the absence of *RBA1*, and these genes were

enriched for those involved in biofilm formation (p= 1.60e-07) and cell aggregation (p= 2.18e-07),

and no enrichment for biological processes was found among the 13 down-regulated genes. By

comparison, when cells were harvested from actively growing biofilms, 701 of 5,594 genes

(12.53%) were observed to be differentially expressed, including 250 and 451 genes which were

up- and down-regulated in the *Δrba1* strain, respectively (**Fig.5b**). Up-regulated genes were

enriched for those involved in transmembrane transport (p= 0.01351), regulation of RNA

biosynthetic process (p=0.02583), cell aggregation (p=0.05187), and regulation of primary

metabolic process (p=0.07722), while down regulated genes were enriched for 18 different

biological processes including protein biosynthetic process (p= 2.35e-28), macromolecule

biosynthetic process (p= 4.05e-10), premeiotic DNA replication (p= 0.00273), and steroid

biosynthetic process (p= 0.02579). Thirty-two *C. auris* genes were up-regulated in the absence

of *RBA1* under both growth conditions, and these genes were enriched for those involved in

biofilm formation and cell aggregation (**Fig. 5c**). Notably, this included four *C. auris* adhesins

(*ALS4112/ALS5, IFF4*, *IFF4109*, and *SCF1*), three of which we have previously shown to impact

virulence in a mouse model of invasive candidiasis[30, 31].

As many genes in the *C. auris* genome remain uncharacterized and/or currently have

incomplete annotations, we further evaluated differential expression among *C. auris* genes with

homology to those associated with adhesion, biofilm formation, and cell wall architecture in the

model pathogenic yeast *Candida albicans.* This revealed four additional *C. auris*-specific adhesins

(B9J08_004110, B9J08_004451, B9J08_004195, and *HYR3*), six additional cell wall-associated

proteins (B9J08_002109, B9J08_005545, B9J08_003550, *HGC1*, *PHR1*, and *XOG1*), seven

GPI-anchored proteins (B9J08_000892, B9J08_001409, B9J08_00116, B9J08_000438,

B9J08_003591, B9J08_00560, and *PGA52*), and three aspartic-type endopeptidases (*SAP1*,

SAP6, and *SAP7*) exhibiting differential expression under biofilm conditions in the absence of

RBA1 (**Fig. 5d**). Notably, four of the differentially expressed genes associated with the fungal cell

wall, B9J08_002109, B9J08_003350, *PHR1*, and *XOG1*, are the predicted orthologs of

C. albicans cell wall proteins known to be involved in cell wall remodeling, cell wall masking from

immune detection, and response to host immune cells[32, 33, 34]. Finally, as *RBA1* appears to repress

biofilm formation in *C. auris,* we also examined the expression of *C. auris* orthologs of genes

known to regulate biofilm formation in the model pathogenic yeast, *Candida albicans.* Two and

17 of these genes were found to be differentially expressed under planktonic and biofilm

conditions, respectively. *UME6* and B9J08_002529 (ortholog of *C. albicans BRG1*), exhibited

significantly increased expression in the *Δrba1* strain under planktonic conditions, and in

C. albicans both of these genes are known to be important positive regulators of hyphal growth

and biofilm formation*.* Further, *UME6* has previously been shown to impact *C. auris* biofilm

formation when overexpressed and artificially hyper-activated through HA-tagging of the putative

transcription factor activation domain[35]. Under biofilm conditions, 14 genes orthologous to those

known to regulate biofilm formation in *C. albicans* were up-regulated, notably including the

C. auris orthologs of *AHR1*, *EFG1*, *RFG1*, *BCR1*, *BRG1*, and *WOR1*, and 3 genes were down-

regulated (orthologs of *C. albicans CRZ2*, *CZF1*, and *ZCF4*). These findings demonstrate that

RBA1 significantly impacts regulation of *C. auris* gene expression during biofilm formation and

suggest that *RBA1* may play a central regulatory role in this clinically impactful process.

Loss of *RBA1* enhances *C. auris* adhesion to plastics and human keratinocytes

A key clinical characteristic of *C. auris* is its ability to adhere to and persist on a variety of

surfaces in the nosocomial environment, including medical equipment and patient skin. As our

transcriptional profiling revealed loss of *RBA1* enhanced the expression of an assortment of

C. auris adhesin-encoding genes, we hypothesized that loss of *RBA1* would enhance *C. auris*

adhesion to multiple substrates. To test this, we first evaluated the adherence of our *Δrba1* and

Δrba1::*RBA1*⁺ strains, as well as the isogenic clinical isolates SKU013 and SKU018, to

polystyrene plastic under a variety of conditions. After 90 minutes of exposure in RPMI tissue

culture media, SKU018 and the *Δrba1* strains exhibited greater adhesion than SKU013 to both

polystyrene and tissue culture-treated polystyrene (**Fig. 6a, b**). This difference was magnified

when the polystyrene well plates were pre-treated with human plasma, with or without the addition

of 10% human serum to the RPMI culture media (**Fig. 6c, d**). By contrast, $\Delta rba1$::$RBA1^+$ control

strains phenocopied the parental SKU013 under all conditions.

Next, we evaluated whether loss of *RBA1* promotes adhesion of *C. auris* to human

keratinocytes. After 60 minutes of exposure, $\Delta rba1$ strains were observed to exhibit enhanced

adherence to a confluent culture of human N/TERT keratinocytes, while neither $\Delta rba1$::$RBA1^+$

controls strains nor SKU018 differed from the SKU013 control (**Fig. 6e, f**). Outbreak isolates

harboring the $RBA1^{Q174*}$ (SKU055), $RBA1^{Q225*}$ (SKU043), and $RBA1^{C303S}$ (SKU056) mutations

were similarly found to exhibit enhanced adhesion to keratinocytes as compared to their closest

available genetic relatives (**Fig. 6g, h**) further supporting the connection between loss of *RBA1*

and enhanced adherence to human keratinocytes. Taken together, these results confirm that loss

of *RBA1* significantly enhances *C. auris* adhesion characteristics under a variety of conditions.

Loss of *RBA1* increases fungal burden during catheter-associated urinary tract infections.

As we had observed that loss of *RBA1* increased *C. auris* adhesion, biofilm formation, and

the expression of the previously characterized virulence factors *SCF1*, *IFF4109*, and

ALS4112/ALS5, we next sought to evaluate the impact of *RBA1* on *C. auris* virulence *in vivo*

during a model of biofilm-related infection. To accomplish this, we utilized our mouse model of

CAUTI [29, 36], where immunocompetent mice were catheterized and the bladder of each mouse

was challenged with *C. auris* transurethrally 24 hours prior to sacrifice for evaluation of fungal

burden and systemic dissemination of infection. Importantly, this mimics a clinical scenario where

C. auris is often cultured from patients with urinary catheters and is relevant to multiple patient

populations at increased risk of *C. auris* colonization and infection. In this model, the $\Delta rba1$ strain

was found to exhibit a 6.8-fold increase in bladder tissue fungal burden (*p*= 0.0435) by 24 hours

post-infection, as compared to SKU013 (**Fig. 7**). Moreover, a trend toward increased

dissemination of the *Δrba1* strain to kidney and spleen tissues was also observed, and one mouse

infected with the *Δrba1* strain was found to have detectable fungal burden in cardiac tissue. These

findings suggest that loss of *RBA1* may also enhance virulence of *C. auris* during clinical infections

such as CAUTI.

Mutations in *RBA1* have emerged during numerous *C. auris* clinical outbreaks

Having identified a consistent signature for *RBA1* mutations during documented clinical

outbreaks of *C. auris* infections in Kuwait and after demonstrating that these mutations confer

phenotypes advantageous to *C. auris* persistence, we hypothesized that *RBA1* mutations have

also likely occurred during *C. auris* outbreaks in other countries as well. To test this, we leveraged

publicly available *C. auris* WGS data collected for infection control and outbreak tracking to

identify mutations in the *RBA1* gene. WGS data are freely available for 2181 *C. auris* clinical

isolates. A total of 94 mutations in *RBA1* were identified, and a total of 310 isolates were found to

carry one or more of the 58 identified non-synonymous or nonsense *RBA1* mutations, including

representatives of the *C. auris* major genetic clades I, II, and IV. The majority of the identified non-

synonymous/nonsense mutations are considered as likely loss-of-function mutations (40; 61.5%),

resulting in early stop codons, frameshift mutations, or alterations within the highly conserved

DNA-binding domain (**Fig. 8a**), similar to the mutations we initially identified from the Kuwait

outbreak collection. We next evaluated meta data and associated publications characterizing

identified isolates harboring non-synonymous *RBA1* mutations to determine those which could be

definitively tied to documented outbreaks of infection. This revealed 10 *RBA1* mutations from

documented outbreaks of *C. auris* infection which occurred on four continents and were caused

by isolates representing both genetic clades I and IV (**Fig. 8b**) [12, 20, 37, 38, 39, 40]. Notably, this included

two instances where *RBA1* mutations arose during Indian outbreaks[41] initially reported in 2013,

one of the earliest reported outbreaks in South America[37] where three distinct *RBA1* mutant

linages emerged, as well as two instances of *RBA1* mutations occurring during ongoing outbreaks

in the United States[38, 42]. These findings clearly demonstrate mutations in *RBA1* have previously

occurred during a multitude of *C. auris* outbreaks, and further suggest that these mutations confer

a competitive advantage to *C. auris* in the healthcare environment.

DISCUSSION

Unique among fungal pathogens, *C. auris* poses challenges to clinicians reminiscent of

healthcare-associated bacterial pathogens such as *Acinetobacter baumannii* and *Enterococcus*

faecium, frequently being associated with clinical outbreaks of infection and multi-drug

resistance[3, 10]. Unlike other agents of invasive candidiasis, *C. auris* is not considered as part of

the human normal flora and is not known to be widely distributed in environmental reservoirs.

Instead, *C. auris* infections are currently thought to be acquired predominantly through

dissemination of recently emergent drug-resistant, and possibly infection-adapted, lineages within

medical environments, leading to the infection of vulnerable patient populations[3, 20]. Thus, as with

other healthcare-associated pathogens, the identification of *C. auris* evolution during clinical

outbreaks and characterization of traits promoting the persistence among human patients is

essential to both preventing infections and mitigating prolonged outbreaks of infection.

Previous studies of isolates from infection outbreaks have largely focused on poor clinical

or microbiologic outcomes driven by mutations leading to antifungal resistance[15, 17, 18, 25, 27, 43, 44].

Here, we analyzed samples from outbreaks in Kuwait to identify genomic and microbiological

elements which may promote transmission and persistence of *C. auris* in clinical settings. Rather

than distinct clustering of serial isolates from individual patient cases, phylogenetic analysis

identified isolate clusters originating from multiple patients indicating multiple instances of

pathogen lateral transmission consistent with clinical dissemination. Interestingly, while numerous

cases of isolates acquiring mutations associated with echinocandin and AmB resistance were

identified, persistence and spread of these multi-drug-resistant lineages was not observed in this

study. Conversely, among both phylogenetic clusters B and C, closely related isolates harboring

acquired *RBA1* mutations were cultured from multiple patients over periods of time spanning

many months. While there was no association between antifungal susceptibility and *RBA1*

genotype, *C. auris* isolates carrying clinically acquired loss-of-function mutations in *RBA1* were

observed to exhibit enhanced biofilm formation as compared to those with wildtype *RBA1*

sequences.

Biofilms are critically important to human public health as they promote microbial survival

in a variety of environments, demonstrating increased resilience to disinfectants and

antimicrobials, and contribute to pathogen dissemination and the establishment of new infections.

In fact, the majority of human infections involve biofilms either within human tissues or on the

surface of in-dwelling medical devices such as catheters or prosthetic implants [45]. As numerous

clinical isolates from the studied clinical outbreaks in Kuwait were isolated from the urine of

catheterized patients or the respiratory secretions of patients with endotracheal tubes, we

hypothesized that the identified loss-of-function mutations in *C. auris RBA1* represented a

clinically emergent adaptation promoting pathogen success in the medical environment through

enhanced biofilm formation. Genetic interrogation of the impact of *C. auris RBA1* definitively

demonstrated loss of this gene enhanced biofilm formation when evaluated using *in vitro* models

mimicking clinically relevant conditions including those specific to CAUTI, an infection from where

C. auris is frequently isolated. Microscopic evaluation revealed biofilms formed by *Δrba1* strain to

be thicker, more confluent, and to include more extracellular matrix on the biofilm surface. This

enhancement of biofilm formation may represent a selective pressure which promoted the

emergence and persistence of *RBA1* mutant lineages during the studied clinical outbreaks.

Subsequent characterization of the impact of *RBA1* on gene expression by transcriptional

profiling revealed a narrow role in *C. auris* cells growing under planktonic conditions, and a much

wider influence on gene expression when cells were grown in biofilms. Under both conditions,

however, *Δrba1* strains exhibited increased expression of a multitude of cell surface adhesins,

including multiple genes we have previously shown to significantly impact *C. auris* virulence

during infection. The two previously characterized *C. auris*-specific adhesins, *SCF1* and

ALS4112/ALS5, were substantially elevated in the *Δrba1* strain grown in planktonic conditions,

with both genes exceeding a 7-log$_2$-fold increase over the SKU013 parental control. As these

adhesins are known to be critical to the ability of *C. auris* to adhere to a variety of substrates

including human skin[30, 31], this suggested that *RBA1* may impact the propensity of *C. auris* to both

colonize patients and establish new biofilms. Further investigation of gene expression in biofilms

demonstrated that a wide variety of genes predicted to encode *C. auris* cell surface proteins, cell

wall constituents, secreted enzymes, and transcriptional regulators were differentially expressed

in the *Δrba1* strain. Intriguingly, this also included the *C. auris* orthologs of *C. albicans ENG1* and

YWP1, two secreted beta-glucanases which are known to play critical roles in cell wall beta-

glucan masking[32, 33, 34]. This finding, combined with the altered expression of a multitude of GPI-

anchored cell surface proteins and the differences in extracellular matrix production observed by

scanning electron microscopy, suggests that the loss of *RBA1* may also facilitate evasion of

C. auris detection from host defenses during infection.

Following the identification of dramatically increased expression of *C. auris* adhesins

SCF1 and *ALS4112/ALS5* in the *Δrba1* strain under planktonic growth conditions, we next

examined whether loss of *RBA1* would impact the ability of *C. auris* to adhere to surfaces where

we have previously shown *SCF1* and *ALS4112/ALS5* to play a significant role[30, 31]. Strains lacking

RBA1 were found to exhibit enhanced adhesion to human keratinocytes as well as polystyrene

surfaces both with and without human plasma treatment, which mimics the deposition of host

factors such as fibrinogen on medical device surfaces. Moreover, evaluation of three *C. auris*

clinical isolates from the Kuwait outbreaks harboring three additional loss-of-function mutations,

as compared to their respective closest available genetic relative with a *RBA1*^{WT} allele, again

demonstrated loss of *RBA1* to increase adhesion of *C. auris* to human keratinocyte. This finding

is of substantial clinical relevance, as it supports that loss of *RBA1* can promote the dissemination

of *C. auris* and colonization of patients during a clinical outbreak. Further, patients colonized with

C. auris strains harboring *RBA1* loss-of-function mutations may pose a greater risk of

transmission within the hospital environment.

We also examined the impact of *RBA1 in vivo* using a murine CAUTI model which

replicates a biofilm-related infection of a site where *C. auris* is commonly isolated. Contrary to

other commonly utilized models of pathogenic yeast infection, this model uses mice that have not

been immunosuppressed. Instead, immunocompetent mice are made vulnerable to *C. auris*

infection through the implantation of a silicone catheter, similar to urinary catheters commonly

used among human patients. In this model, the *Δrba1* strain was found to develop a significantly

greater fungal burden in the mouse bladder, and a trend towards greater dissemination of the

Δrba1 strain from the urinary tract to other tissues including the heart was observed. These

findings further suggest loss of *RBA1* is advantageous during *C. auris* clinical infection. This is

clinically significant as CAUTIs often lead to urosepsis with a ~30% 7-day mortality rate [19, 46].

Since the urinary tract is a source of ~25% of the all the sepsis cases, patients infected with

C. auris strains harboring *RBA1* mutation may be predisposed to severe clinical complication.

Finally, leveraging the growing body of publicly available WGS data generated through

ongoing *C. auris* infection control and prevention efforts, we were able to investigate whether

mutations in *RBA1* have emerged during clinical outbreaks in other countries. This approach

uncovered dozens of distinct *RBA1* mutations, impacting hundreds of clinical isolates and more

than 10% of evaluable samples. *RBA1* mutations were identified among isolates representing

three of the major genetic subgroups of *C. auris,* but were notably absent among isolates from

clade III*,* suggesting either clade-specific impacts of *RBA1* mutations or a potential bias within

samples currently available in the public data set. Like the four *RBA1* mutations identified from

the Kuwait outbreaks, the mutations identified from the global collection were predominantly

predicted loss-of-function mutations such as those encoding early stop codons, frameshift

mutations, or altering highly conserved residues in the zinc-cluster transcription factor DNA-

binding domain. Identified isolates with *RBA1* mutations were found to date back to some of the

earliest documented clinical outbreaks of *C. auris*, including those in India, England, and

Columbia. Additionally, isolates from outbreak lineages actively disseminating in the United States

were also found to harbor *RBA1* mutations. Taken together, these findings clearly demonstrate

that mutations in *RBA1* have emerged repeatedly during clinical outbreaks of *C. auris* infection

and strongly support the notion that the enhancement in adhesion and biofilm formation conferred

by these mutations provided a competitive advantage within the medical environment in

numerous previous outbreaks.

In conclusion, these results call attention to the capacity of *C. auris* to evolve and enhance

traits beyond drug resistance during outbreaks of infection by highlighting the unique nature of

C. auris among other fungal pathogens and identifying mutations in *RBA1* as a potential indicator

of outbreak-adapted lineages. As traits conferred by *RBA1* mutations are advantageous to

C. auris spread and persistence, and as we have identified numerous instances where these

mutations have arisen during global outbreaks of infection, the identification and tracking of

lineages carrying *RBA1* mutations may be merited as part of infection prevention efforts.

MATERIALS AND METHODS

Ethics Statement

C. auris isolates were cultured from various clinical specimens of hospitalized patients in Kuwait. The need for informed consent was waived by the Health Sciences Center Ethics Committee (Approval letter Ref: VDR/EC/3724 dated January 31, 2021) as the clinical specimens were collected as part of routine diagnostic work-up for the isolation of bacterial/fungal pathogens for suspected infections. The results are reported here on deidentified samples without revealing patient identity. All animal care was consistent with the Guide for the Care and Use of Laboratory Animals from the National Research Council. The University of Notre Dame Institutional Animal Care and Use Committee approved all mouse infections and procedures as part of protocol number 25-01-9009. For urine collection, all donors signed an informed consent form, and protocols were approved by the Institutional Review Board of the University of Notre Dame under study #19-04-5273.

C. auris Isolates

Seventy *C. auris* isolates from previously described multi-institution outbreaks of *C. auris* infections in Kuwait, were obtained by the Mycology Reference Laboratory Department of Microbiology, Kuwait University, Kuwait [47]. All isolates (including names used in publication of their original description) and strains used in this study are listed in **Table S1**. All *C. auris* strains were stored in 40% glycerol stocks at -80°C and cultured at 35°C in liquid YPD (1% yeast extract, 2% peptone, 2% glucose) media before plating on Sabouraud dextrose (SD) agar (BD Difco) media to ensure standardized growth for phenotypic testing.

Plasmid and strain construction

EPIC-mediated gene editing was used to generate *RBA1* (B8441v3: B9J08_03209) modifications in *C. auris* clinical isolate SKU013 (also known as Kw1432/17). Duplexed guide

oligos (Integrated DNA Technologies) (**Table S2**) targeting disruption or reversion of

B9J08_03209 were added to the pJMR19 vector using restriction enzyme cloning[26]. Plasmids

were maintained in *E. coli* DH5α cells cultured in Luria-Bertani broth supplemented with 5mg/L

ampicillin and isolated using QIAprep® Spin Minipreps (Qiagen). Repair templates for *RBA1-*

disruption and reversion were amplified from a gBlock sequence (Integrated DNA Technologies)

and SKU013 gDNA, respectively, via PCR and purified with QIAquick PCR Purification Kit

(Qiagen). All oligos, plasmids, and gBlocks used in this study are listed in **Table S2.** *C. auris*

Lithium acetate transformations were then performed as previously described [26] with positive

selection on 200µg/mL nourseothricin-supplemented YPD agar. *RBA1* editing was confirmed by

both Sanger sequencing at the Hartwell Center at St. Jude Children's Research Hospital (SJCRH)

and with long read nanopore sequencing by Plasmidsaurus.

Whole genome sequencing

Genomic DNA (gDNA) was extracted from 1mL of overnight liquid YPD cultures using

MasterPure™ Yeast DNA Purification kits (Biosearch Technologies) according to manufacturer

instructions. Sample libraries were prepared from gDNA by the Hartwell Center (SJCRH) using

KAPA HyperPrep (Roche) and sequencing targeting 100x coverage for paired 150bp reads was

performed via Illumina NovaSeq. Paired-reads FASTQ files were analyzed using CLC genomics

workbench v24.0 (QIAGEN) software. Low quality reads were trimmed with Trim Reads v2.8 and

subsequently mapped to the B8441v3 genome (GCA002759435.3) according to CLC-workbench

recommendations for match scores, mismatch, insertion, and deletion costs. Local Realignment

v2.6 was used to refine unaligned ends and mutations were identified within annotated coding

sequences via Fixed Ploidy Variant Detection v2.6 with ploidy=1.

Phylogenetic analysis was initiated with Create SNP Tree v1.9 using the Maximum

Likelihood Phylogeny v1.4 algorithm. The resulting SNP phylogenetic tree was visualized from

Neighbor Joining construction methods specifying Jukes Cantor nucleotide substitution modeling

with estimated topology and bootstrapping for 100 replicates. Sequences were deposited in

National Center for Biotechnology Information Sequence Read Archive.

Biofilm Production Assays

To assess biofilm production in the presence of human plasma, 96-well flat-bottomed

plates (VWR, 10861-562) were coated with 100 μL pooled human plasma (Innovative Research

Inc, IPLAWBNAC50ML) diluted 1:5 in carbonate-bicarbonate buffer (Sigma C3041-50CAP) and

incubated overnight at 4°C. Colonies from overnight SD agar plates were used to generate

standardized OD_{600} = 0.5 cell suspensions[48] in RPMI-Biofilm media, consisting of RPMI medium

1640 (Gibco) + 3-(4-Morpholino)propane sulfonic acid (MOPS, Fisher Bioreagent) + 2% dextrose,

buffered to pH 7 and supplemented with 10% heat-inactivated/male AB human serum (HuS,

Sigma-Aldrich). Plasma coatings were gently removed the plates and 200 μL of the cell

suspensions were added to the wells and plates were sealed with a breathable (EASYSEAL)

membrane. After 90-minutes at 37°C, 250 RPM, plates were gently washed sterile PBS to remove

non-adhered cells. Fresh 200 μL of RPMI-Biofilm media was then added back to each well and

plates were incubated for 24 hrs at 37°C, 250 RPM. For quantification of biofilm production,

supernatants were removed and wells were gently washed twice with PBS. Biofilms were stained

with 0.4% crystal violet as previously described[48]. Crystal violet absorbance measured at 595nm

on an Agilent BioTek Synergy H1 Microplate Reader. Average media only negative controls were

subtracted from the sample reads and values were normalized to SKU013.

To replicate biofilm growth replicating the urinary tract environment, 96-well flat-bottomed

plates (VWR, 10861-562) were coated with 100 µl of Fg (450 µg/ml) and incubated overnight at

4°C.[29] The various strains were grown overnight, shaking in YPD, and the inoculum was

normalized to ~1 × 10^6 CFUs/ml. Cultures were then diluted (1:100) into human urine

supplemented with 10% heat inactivated human serum and incubated in the wells of the 96-well

plate at 37°C for 48 hours. Human urine from at least two healthy female donors between the

ages of 20 to 35 were collected and pooled. Donors did not have a history of kidney disease,

diabetes, or recent antibiotic treatment. Urine was sterilized with a 0.22-µm filter (VWR 29186-

212), pH was normalized to 6.0 to 6.5, and urine was used immediately for the assays. Following

biofilm formation, supernatants were removed, and plates were incubated in 200 µl of 0.5% crystal

violet for 15 min. Crystal violet stain was removed, and the plate was washed with water to remove

the remaining stain. Plates were dried and then incubated with 200 µl of 33% acetic acid for 15

min. In another plate, 100 µL of the acetic acid solution was transferred, and absorbance values

were measured via a plate spectrophotometer at 595 nm (Molecular Devices SpectraMax ABS).

Values were then normalized to SKU013.

RNA Extraction

Normalized cell suspensions (OD_{600} = 0.1) from SD replicate platings were diluted ~1:50

in 10 mL RPMI-Growth media, prepared from RPMI medium 1640 (Gibco) + MOPS (Fisher

Bioreagent) buffered to pH 7 and grown overnight at 35°C, 220 RPM to an OD_{600} = 2.0 $\pm$ 0.3.

Concentrated cultures were diluted in >20mL RPMI-Growth media to an OD_{600} = 0.2 then split

into two 10mL aliquots for direct comparison of planktonic growth and biofilm development.

Planktonic cells were grown over 4 hours in 50mL conical at 35°C, 220RPM and pelleted at 4000

RPM. For biofilm development cells were transferred to T75 flask pre-coated with 20% human

plasma in carbonate-bicarbonate buffer prepared as described above. Following 2 hr incubation

at 37°C, non-adhered cells were gently removed and fresh 10mL RPMI-Growth media was added

for an additional 4 hr incubation at 37°C. Biofilms were gently washed once with PBS and biofilms

were then removed by scraping in the presence of 10mL sterile PBS and pipette transfer to a

50mL conical tube for centrifuge pelleting. All pellets were flash frozen in liquid nitrogen and stored

at -80°C. RNA was isolated using the Invitrogen RiboPure™ Yeast kit (ThermoFisher Scientific™,

USA) per manufacturer's instructions.

RNA Sequencing

The Hartwell Center (SJCRH) prepared the RNA libraries using Illumina chemistries per

manufacturer's instructions and sequenced the RNA on an Illumina NextSeq (Illumina Inc., USA)

using the total stranded RNA-Seq application for paired end runs targeting 150bp reads.

Sequenced reads were analyzed with CLC Genomics Workbench version 24.0.1 (QIAGEN).

Imported reads were trimmed using default settings for failed reads and adaptor sequences and

then subsequently mapped to the *C. auris* B8441 reference genome (GenBank accession:

GCA_002759435.3) with paired reads counted as one and expression values set to RPKM.

Principal component analysis was utilized for the initial clustering assessment of biological

replicates and mismatch, insertion, and deletion costs were set to default parameters for RNA-

Seq Analysis 2.7. Differential expression was calculated using negative binomial GLM modeling

with whole transcriptome RNA-Seq defaults for normalization with Trimmed Mean of M values,

using Differential gene expression v1.4 (CLC genomics workbench, Qiagen). Comparisons were

made to the wildtype *SKU013* clinical isolate for each condition. Differential expression was

defined as fold changes ≥ 2 or ≤ -2 (FDR p-value ≤0.05)

Scanning Electron Microscopy

Sterile silicon chips were loaded into 24-well plate and coated overnight with 0.5mL 20%

human plasma in carbonate-bicarbonate buffer. Suspensions of SKU013, *Δrba1*::*RBA1*+_a , and

Δrba1_a were cultured in RPMI-Biofilm media as described above for biofilm production assays

with 0.5mL of cells loaded into the wells. Following 24 hour incubation, wells were gently washed

with 1mL PBS and biofilms were fixed with 0.1M sodium cacodylate buffer at pH 7.4 with 0.15%

alcian blue, 1% tannic acid, 2.5% glutaraldehyde, 2% paraformaldehyde for 2 hours and stored

at 4°C. Preparation, staining, and imaging the fixed biofilms was performed by the Electron

Microscopy Division of the Cell and Tissue Imaging Center (CTIC-EM) at SJCRH for SEM Fixed

biofilms were washed with 0.1M sodium cacodylate buffer + 0.15% alcian blue prior to osmium

tetroxide staining in the presence of 0.15% alcian blue. Imaging was performed on a Zeiss

GeminiSEM 460. Scale bars were embedded using ImageJ[49].

Polystyrene adhesion assay

Adhesion to flat-bottom 96-well non-tissue culture treated polystyrene plates (CS100-

701311, NEST SCIENTIFIC) was assessed in RPMI-Growth+ media, defined as RPMI medium

1640 + MOPS + 2% dextrose, buffered to pH 7. Adhesion to tissue culture treated plates

(TP92696, TPP Midwest Scientific), was assessed for RPMI-Growth+ media and RPMI-Biofilm

media in the presence and absence of human plasma, coated as described above. cells cultured

and plates were loaded as described for biofilm production assays. Following a 90-minute

incubation at 37°C, 250RPM, wells were gently washed once with PBS to remove non-adhered

cells. Adhered cells were stained with 100μL of a 10:1 solution of XTT sodium salt and phenazine

methosulfate (VWR International LLC). Absorbances at OD_{492} and OD_{630} were read after 60

minutes and values were normalized to SKU013.

Keratinocyte adhesion assay

Adhesion of *Candida auris* to human keratinocytes was determined as previously

described[30]. Human N/TERT keratinocytes[50] were grown in 96-well plates to complete confluence

in keratinocyte-SFM medium (Gibco, 10724-011) supplemented with 30 μg/ml bovine pituitary

extract, 0.2 ng/ml epidermal growth factor, and 0.31 mM $CaCl_2$. Overnight *C. auris* cultures were

diluted to an OD_{600} of 0.03, and 100 μl of this suspension was added to each well containing

keratinocytes. In parallel, an equal volume of culture was added to empty wells, centrifuged (210

g, 1 min), and imaged to determine input cell counts, while plates containing keratinocytes were

centrifuged under the same conditions and incubated at 37 °C with 5% CO_2 for 1 h. Non-adherent

yeast cells were then removed by washing three times with PBS. The remaining cells were fixed

in 4% formaldehyde and stained with FITC-conjugated anti-*Candida* antibody (Meridian

Bioscience, B65411F; 1:1000). Keratinocyte nuclei and cell bodies were labeled with Hoechst

(Thermo Fisher, 62249; 1:1000) and CellMask Deep Red (Thermo Fisher, H32721; 1:5000),

respectively.

Fluorescent images were acquired at 10X magnification using a BioTek Lionheart FX

microscope. The number of adherent *C. auris* cells was quantified using a customized CellProfiler

analysis pipeline[30, 51]. Numbers of input cells were determined from brightfield images pre-

processed in Fiji/ImageJ (v2.14.0)[49] using a custom macro for edge-based segmentation and

binary mask generation. The proportion of adherent to input cells was calculated to assess the

adhesion efficiency of each *C. auris* strain.

Catheter-Associated Urinary Tract Infection (CAUTI) Modeling

Female C57BL/6 mice (Jackson Laboratory) were subjected to transurethral implantation

of a silicone catheter followed by transurethral inoculation of the bladder lumen with ~5 × 10^5

colony forming units (CFU) in PBS, as previously described [36]. Mice were euthanized at 24 hours

post-infection by cervical dislocation after anesthesia inhalation and the catheter, bladder,

kidneys, spleen, and heart were aseptically harvested. Bladders were weighed and then

homogenized for CFUs. The other organs were homogenized, and catheters were cut into small

pieces before sonication for fungal CFU enumeration.

Statistics and Reproducibility

Statistical parameters and applied analysis methods including exact sample numbers (n)

and definitions for representative central value and spread (mean ± SEM, CFU quantifications;

representative modal value (n=3), minimum inhibitory concentrations) are annotated in

corresponding figure legends. Data from at least three experiments were pooled for each

adhesion and biofilm assay and twice for each *in vivo* mouse assay. One-way ANOVA tests and

two-tailed Mann-Whitney *U* tests were performed with GraphPad Prism 10 software (GraphPad

Software) for comparisons of described in-vitro assays and in vivo CAUTI experiments,

respectively.

For RNA sequencing, statistical analysis was performed using CLC Genomics Workbench

version 20.0 (QIAGEN) with paired reads counted as one and expression values set to RPKM.

Whole transcriptome differential gene expression analysis was performed with mismatch,

insertion, and deletion costs set to default parameters, and a Wald test was used to compare

SKU013 against Δ*rba1_a*. Differential expression determined by a fold change of ≥ 2 or ≤ -2 with

a False discovery rate (FDR) corrected p value ≤0.05.

ACKNOWLEDGMENTS

This work was supported in part by the St. Jude Children's Research Hospital Children's

Infection Defense Center and Center for Infectious Diseases Research awards granted to JMR,

by the National Institutes of Health (NIH)/ National Institute of Allergy and Infectious Diseases

(NIAID) grant U19-AI181767 awarded to TRP, and by the NIH/ NIAID F32 AI181164 awarded to

GZ. This work was supported by institutional funds from the University of Notre Dame (to FHST

and ALFM), and by grants from the Good Venture Foundation (Open Philanthropy) (to ALFM),

from the National Institutes of Health R01AI177875 and R21AI171742 (to FHST), and

R01DK128805 (to ALFM). This research included experiments conducted by the Hartwell Center

for Bioinformatics & Biotechnology, the St. Jude Center for Advanced Genome Engineering and

the Cell and Tissue Imaging Center - Electron Microscopy (CTIC-EM) which are supported in part

by ALSAC and the National Cancer Institute at the National Institutes of Health [P30 CA021765].

The funders had no role in study design, data collection and interpretation, or the decision to

submit the work for publication.

AUTHOR CONTIBUTIONS

Conceptualization, J.M.R., T.R.O., F.H.S.T., and A.L.F.M.; methodology, J.M.R., L.A.D., A.A.L.,

S.J.J., Q.Z., T.R.O., and A.L.F.M.; Investigation, L.A.D., A.A.L., S.J.J., Q.Z., Q.J.C., and J.M.R.;

writing—original draft, L.A.D. and J.M.R.; writing—review & editing, L.A.D., A.A.L., S.J.J., G.Z.,

M.A., Q.J.C., S.A., E.M., I.A., W.A., K.A., T.R.O., F.H.S.T., A.L.F.M., and J.M.R.; funding

acquisition J.M.R., T.R.O., F.H.S.T., A.L.F.M., and Q.Z.; supervision, T.R.O., F.H.S.T., A.L.F.M.,

and J.M.R..

FIGURES AND TABLES
673

674
675 **Figure 1. Clinical emergence and spread of *C. auris* isolates harboring mutations in *RBA1***
676 **during an outbreak of infections.** Phylogenetic analysis of 70 *C. auris* isolates from a
677 documented outbreak of infections in Kuwait. Branch length termination is indicated with black
678 boxes. Patient case numbers, culture sources, and *RBA1* mutations are indicated on the right.
679

680
Figure 2. *RBA1* loss-of-function mutations emerge during patient infection and are associated with increased biofilm production A) Timeline of patient cases 51 and 53 with *C. auris* isolation dates and antifungal treatment courses indicated, AMB: liposomal amphotericin B, CSF: caspofungin. Isolate numbers circled with yellow backgrounds indicating *RBA1* mutation. **B)** Visual difference in biofilm formation on sidewall of polystyrene culture tubes after overnight incubation in RPMI media. Arrow indicates visible biofilm ring. **C)** Biofilm production of *C. auris* isolates obtained from patient cases 51 and 53, formed over 48 hours on fibrinogen treated polystyrene in urine supplemented with 10% serum assessed by absorbance at 595 nm wavelength following crystal violet stain and normalized to the initial isolate SKU013. Isolates indicated with dark blue bars have a wildtype *RBA1* allele, while isolates with green bars have the *RBA1*E47* mutation. **D)** Biofilm production of *C. auris* clinical isolates harboring *RBA1*Q174*, *RBA1*Q225*, and *RBA1*C303S mutations and corresponding closest genetic relatives formed over 48 hours on fibrinogen treated polystyrene in urine media supplemented with 10% serum assessed by absorbance at 595 nm wavelength following crystal violet stain and normalized to SKU013. Isolates indicated with dark blue bars have a wildtype *RBA1* allele, while isolates with light blue bars have the *RBA1*Q174* mutation, isolates with red bars have the *RBA1*Q225* mutation, and isolates with orange bars have the *RBA1*C303S mutation.

Figure 3. Loss of *RBA1* enhances biofilm production in multiple conditions mimicking difference niche of infection. A) Biofilm production of *C. auris* isolates and *RBA1* strains formed over 48 hours on fibrinogen coated microtiter plates in human urine media supplemented with 10% human serum assessed by absorbance at 595 nm wavelength following crystal violet stain. Absorbance is normalized to SKU013 (*RBA1*^{WT}) with bars representing mean and standard error of the mean. Kruskal-Wallis statistical testing with Dunn's correction for multiple comparisons (GraphPad Prism) showed significant differences between *Δrba1_a, Δrba1_b*, and SKU018 (*RBA1*^{E47*}) when compared to SKU013 (*RBA1*^{WT}) for both adhesion and biofilm production (****P= <0.0001). **B)** Biofilm production of *C. auris* isolates and *RBA1* strains formed over 24 hours on human plasma coated microtiter plates in RPMI tissue culture media supplemented with 10% human serum assessed by absorbance at 595 nm wavelength following crystal violet stain. Absorbance is normalized to SKU013 (*RBA1*^{WT}) with bars representing mean and standard error of the mean. Kruskal-Wallis statistical testing with Dunn's correction for multiple comparisons (GraphPad Prism) showed significant differences between *Δrba1_a, Δrba1_b*, and SKU018 (*RBA1*^{E47*}) when compared to SKU013 (*RBA1*^{WT}) for both adhesion and biofilm production (****P= <0.0001).

718
719
Figure 4. Loss of *RBA1* promotes formation of robust biofilms with greater extracellular
matrix. A) Representative SEM images for SKU013 (*RBA1*^WT) biofilms formed over 24 hours in
RPMI media supplemented with 10% human serum following substrate treatment with human
plasma using the 'Silicone substrate–confocal microscopy" method. Black box in left panel
indicates area enlarged in right panel. **B)** Representative SEM images for *Δrba1* biofilms formed
over 24 hours in RPMI media supplemented with 10% human serum following substrate treatment
with human plasma using the 'Silicone substrate–confocal microscopy" method. Black box in left
panel indicates area enlarged in right panel.
728

Figure 5. Loss of *RBA1* increases expression of *C. auris* adhesins and virulence factors
under both planktonic and biofilm conditions. Volcano plot of differentially expressed genes
between clinical isolate SKU013 and *Δrba1* strain during exponential growth phase when cultured
under **A)** planktonic growth conditions (10mL RPMI media in 50mL conical tube incubated at 35°C
with 220RPM) and **B)** biofilm-forming conditions (10mL RPMI media in plasma-treated T75 flasks
for static incubation at 37°C). Genes involved in cell aggregation, biofilm formation, or drug
resistance are highlighted in blue, green, and purple, respectively. Dotted lines represent lower
limit equivalents for fold changes of ≥1.5 or ≤-1.5 on the x-axis, and a false discovery rate (FDR)
p-value ≤0.05 on the y-axis. **C)** Venn diagram comparing differentially expressed genes (≥1.5 or
≤-1.5 on the x-axis, and a false discovery rate p-value ≤0.05) exhibiting up-regulation or down-
regulation in the *Δrba1* strain under planktonic or biofilm growth conditions. **D)** Differential
expression of *C. auris* gene predicted to encode GPI-anchored proteins, cell surface adhesions,
proteins associated with fungal cell wall, aspartic-type endopeptidases, and transcriptional
regulators predicted to be orthologous to those known to to be involved in biofilm formation in
Candida albicans. Similar *C. albicans* genes are listed in gray and are shown in **bold** if predicted
to be orthologous. Genes which are differentially expressed (≥1.5 or ≤-1.5 on the x-axis, and a
false discovery rate p-value ≤0.05) are indicated with a black dot.

748
749
Figure 6. Loss of *RBA1* increases adhesion to both environmental surfaces and keratinocytes. Adhesion of *C. auris* isolates and *RBA1* strains to **A)** untreated polystyrene, **B)** tissue culture treated polystyrene, **C)** human plasma-coated tissue culture-treated polystyrene, and **D)** human plasma-coated tissue culture-treated polystyrene with 10% human serum supplement in RPMI base media after 90 minutes assessed by absorbance at 490 nm wavelength following XTT treatment. **E)** Representative immunofluorescence images showing adherence of SKU013 (*RBA1*^WT^), *Δrba1*, and *Δrba1::RBA1*^+^ cells to human N/TERT keratinocytes. Scale bar, 50µm. Magenta=keratinocyte; Blue=nuclei; Yellow=*C. auris*. **F)** Adherence of *C. auris* isolates and *RBA1* strains to human N/TERT keratinocytes as measured by the proportion of attached cells remaining after washing. Bars represent mean ± standard deviation for three biological replicates, statistical significance was determined using 1-way ANOVA followed by Dunnett's test

for multiple comparisons, *P ≤ 0.05. **G)** Representative immunofluorescence images showing
adherence of *C. auris* clinical isolates harboring *RBA1*^{Q174*}, *RBA1*^{Q225*}, and *RBA1*^{C303S} mutations
and corresponding closest genetic relatives to human N/TERT keratinocytes. Scale bar, 50µm.
Magenta=keratinocyte; Blue=nuclei; Yellow=*C. auris.* Representative immunofluorescence
images showing adherence of SKU013 (*RBA1*^{WT}), *Δrba1*, and *Δrba1::RBA1$^+$* cells to human
N/TERT keratinocytes. Scale bar, 50µm. Magenta=keratinocyte; Blue=nuclei; Yellow=*C. auris.* **H)**
Adherence of *C. auris* clinical isolates harboring *RBA1*^{Q174*}, *RBA1*^{Q225*}, and *RBA1*^{C303S} mutations
and corresponding closest genetic relatives to human N/TERT keratinocytes as measured by the
proportion of attached cells remaining after washing. Bars represent mean ± standard deviation
for three biological replicates, statistical significance was determined using 1-way ANOVA
followed by Dunnett's test for multiple comparisons, *P ≤ 0.05.
772

773
774

Figure 7. Loss of *RBA1* results in greater fungal burden in a murine model of catheter
associated urinary tract infection. Fungal burden of SKU013 and the derivative *Δrba1* strain
obtained from the catheter material or bladder, kidney, spleen, and heart tissues after infection in
a murine model of catheter associated urinary tract infection (CAUTI). Two-tailed Mann-Whitney
tests showed significant differences in fungal burden recovered from bladder tissue (*$P \leq 0.05$,
n=8 mice per group). CFU, colony forming units; LOD, lower limit of detection.
781

782
783
Figure 8. Identification of *RBA1* mutations emerging during numerous outbreaks of
***C. auris* infections worldwide. A)** Diagram depicting non-synonymous mutations in *RBA1*
identified by analyzing publicly available *C. auris* whole genome sequencing data. Green lines on
top of figure indicate early stop codons and frameshift mutations. Blue lines on bottom of figure
indicate missense mutations. N241H and *354Q were identified among all clinical isolates
classified as clade IV. DBD: DNA-binding domain. **B)** World map indicating locations of
documented outbreaks of *C. auris* infections where mutations in *RBA1* were identified from
publicly available data. Green fill indicates outbreaks involving isolates representing genetic Clade
I and blue fill indicates outbreaks involving isolates representing genetic clade IV. *RBA1* mutations
identified from publicly available sequencing of a global collection of *C. auris* isolates.

	Culture Source	SNPs vs SKU013	*RBA1*	Susceptibility		
				FLU	AmB	Echino
Patient 51						
SKU013	Endotracheal asp.	-	WT	**R**	S	S
SKU014	Endotracheal asp.	2	WT	**R**	S	S
SKU015	Endotracheal asp.	4	WT	**R**	S	S
SKU016	Endotracheal asp.	4	WT	**R**	S	S
SKU017	Endotracheal asp.	4	WT	**R**	S	S
SKU018	Urine	8	**E47***	**R**	S	**R**
SKU020	Blood	5	WT	**R**	S	S
Patient 53						
SKU022	Endotracheal asp.	6	**E47***	**R**	S	S
SKU023	Endotracheal asp.	6	**E47***	**R**	S	S
SKU026	Tracheal sec.	7	**E47***	**R**	S	S
SKU031	Urine	7	**E47***	**R**	**R**	S
SKU032	Urine	89	WT	**R**	S	S
SKU033	Urine	8	**E47***	**R**	**R**	S
SKU035	Urine	8	**E47***	**R**	S	**R**

796
797

Table 1. Description of isolates obtained from patient cases 51 and 53. Isolates name, culture source, number of SNPs in coding regions relative to the earliest isolate SKU013, *RBA1* genotype, and interpretation of susceptibility to fluconazole (FLU), amphotericin B (AmB), and echinocandins (Echino) based on current CDC tentative breakpoints. SNPs: single nucleotide polymorphisms.

804
805

REFERENCES

1. World Health Organization. *WHO fungal priority pathogens list to guide research, development and public health action*. (World Health Organization, Geneva, 2022).

2. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol Immunol* **53**, 41-44 (2009).

3. Lockhart SR*, et al.* Simultaneous Emergence of Multidrug-Resistant *Candida auris* on 3 Continents Confirmed by Whole-Genome Sequencing and Epidemiological Analyses. *Clin Infect Dis* **64**, 134-140 (2017).

4. European Centre for Disease Prevention and Control. *Survey on the epidemiological situation, laboratory capacity and preparedness for Candidozyma (Candida) auris, 2024.* (Publications Office of the European Union, 2025).

5. Sabino R, Veríssimo C, Pereira Á A, Antunes F. *Candida auris*, an Agent of Hospital-Associated Outbreaks: Which Challenging Issues Do We Need to Have in Mind? *Microorganisms* **8**, (2020).

6. Baker AD, Gold JAW, Forsberg K, Jones S, Lyman MM. Progression from *Candida auris* Colonization Screening to Clinical Case Status, United States, 2016-2023. *Emerg Infect Dis* **31**, 1613-1617 (2025).

7. Gülten E*, et al. Candida auris* vs. Non-*Candida auris* Candidemia in Critically Ill Patients: Clinical Outcomes, Risk Factors, and Mortality. *J Fungi (Basel)* **11**, (2025).

8. Arslan E*, et al.* Comparison of risk factors and mortality of candidemia by *Candida auris* versus *Candida* non-*auris* in intensive care units. *Infect Dis (Lond)*, 1-15 (2025).

9. Akinbobola AB, Kean R, Hanifi SMA, Quilliam RS. Environmental reservoirs of the drug-resistant pathogenic yeast *Candida auris*. *PLoS Pathog* **19**, e1011268 (2023).

10. Proctor DM*, et al.* Clonal *Candida auris* and ESKAPE pathogens on the skin of residents of nursing homes. *Nature* **639**, 1016-1023 (2025).

11. Pacilli M*, et al.* Regional Emergence of *Candida auris* in Chicago and Lessons Learned From Intensive Follow-up at 1 Ventilator-Capable Skilled Nursing Facility. *Clin Infect Dis* **71**, e718-e725 (2020).

847   12.   Chow NA*, et al.* Multiple introductions and subsequent transmission of multidrug-resistant
848   *Candida auris* in the USA: a molecular epidemiological survey. *The Lancet Infectious*
849   *Diseases* **18**, 1377-1384 (2018).

850
851   13.   Lyman M*, et al.* Worsening Spread of *Candida auris* in the United States, 2019 to 2021.
852   *Ann Intern Med* **176**, 489-495 (2023).

853
854   14.   Lyman M*, et al.* Notes from the Field: Transmission of Pan-Resistant and Echinocandin-
855   Resistant *Candida auris* in Health Care Facilities - Texas and the District of Columbia,
856   January-April 2021. *MMWR Morb Mortal Wkly Rep* **70**, 1022-1023 (2021).

857
858   15.   Rybak JM*, et al.* In vivo emergence of high-level resistance during treatment reveals the
859   first identified mechanism of amphotericin B resistance in *Candida auris*. *Clin Microbiol*
860   *Infect* **28**, 838-843 (2022).

861
862   16.   Codda G*, et al.* In vivo evolution to echinocandin resistance and increasing clonal
863   heterogeneity in *Candida auris* during a difficult-to-control hospital outbreak, Italy, 2019 to
864   2022. *Euro Surveill* **28**, (2023).

865
866   17.   Carolus H*, et al.* Genome-Wide Analysis of Experimentally Evolved *Candida auris* Reveals
867   Multiple Novel Mechanisms of Multidrug Resistance. *mBio* **12**, (2021).

868
869   18.   Rybak JM*, et al.* Mutations in *TAC1B*: a Novel Genetic Determinant of Clinical Fluconazole
870   Resistance in *Candida auris*. *mBio* **11**, (2020).

871
872   19.   La Bella AA, Santiago-Tirado FH, Flores-Mireles AL. *Candida auris* is emerging as a
873   prevalent urinary pathogen. *PLoS Pathog* **21**, e1013138 (2025).

874
875   20.   Chow NA*, et al.* Tracing the Evolutionary History and Global Expansion of *Candida auris*
876   Using Population Genomic Analyses. *mBio* **11**, (2020).

877
878   21.   Muñoz JF*, et al.* Clade-specific chromosomal rearrangements and loss of subtelomeric
879   adhesins in *Candida auris*. *Genetics* **218**, (2021).

880
881   22.   Barker KS*, et al.* Relative contributions of the *ERG11*[(VF125AL)] and *MRR1A*[(N647T)] mutations
882   to fluconazole resistance in Clade III *Candidozyma (Candida) auris* clinical isolates. *Clin*
883   *Microbiol Infect*, (2025).

884
23.   Rybak JM, Sharma C, Doorley LA, Barker KS, Palmer GE, Rogers PD. Delineation of the
Direct Contribution of *Candida auris ERG11* Mutations to Clinical Triazole Resistance.
Microbiol Spectr **9**, e0158521 (2021).

24. Mayr EM, Ramírez-Zavala B, Krüger I, Morschhäuser J. A Zinc Cluster Transcription
Factor Contributes to the Intrinsic Fluconazole Resistance of *Candida auris*. *mSphere* **5**,
(2020).

25. Massic L*, et al.* Acquired amphotericin B resistance attributed to a mutated *ERG3* in
Candidozyma auris. *Antimicrob Agents Chemother* **69**, e0060125 (2025).

26. Doorley LA, Meza-Perez V, Jones SJ, Rybak JM. A *Candidozyma (Candida) auris-*
Optimized Episomal Plasmid-Induced Cas9-Editing System Reveals the Direct Impact of
the S639F-Encoding *FKS1* Mutation. *J Infect Dis* **232**, e529-e536 (2025).

27. Spruijtenburg B*, et al.* Whole genome sequencing analysis demonstrates therapy-induced
echinocandin resistance in *Candida auris* isolates. *Mycoses* **66**, 1079-1086 (2023).

28. Asadzadeh M*, et al.* Molecular characterisation of *Candida auris* isolates from
immunocompromised patients in a tertiary-care hospital in Kuwait reveals a novel mutation
in *FKS1* conferring reduced susceptibility to echinocandins. *Mycoses* **65**, 331-343 (2022).

29. La Bella AA*, et al.* The catheterized bladder environment promotes Efg1- and Als1-
dependent *Candida albicans* infection. *Science Advances* **9**, eade7689 (2023).

30. Zhao G*, et al.* Adhesin Als4112 promotes *Candida auris* skin colonization through
interactions with keratinocytes and extracellular matrix proteins. *Nat Commun* **16**, 5673
(2025).

31. Santana DJ*, et al.* A *Candida auris*-specific adhesin, Scf1, governs surface association,
colonization, and virulence. *Science* **381**, 1461-1467 (2023).

32. de Assis LJ*, et al.* Nature of β-1,3-Glucan-Exposing Features on *Candida albicans* Cell
Wall and Their Modulation. *mBio* **13**, e0260522 (2022).

33. Hopke A, Nicke N, Hidu EE, Degani G, Popolo L, Wheeler RT. Neutrophil Attack Triggers
Extracellular Trap-Dependent *Candida* Cell Wall Remodeling and Altered Immune
Recognition. *PLoS Pathog* **12**, e1005644 (2016).

34. Yang M*, et al.* Control of β-glucan exposure by the endo-1,3-glucanase Eng1 in *Candida*
albicans modulates virulence. *PLoS Pathog* **18**, e1010192 (2022).

35. Louvet M*, et al.* Ume6-dependent pathways of morphogenesis and biofilm formation in
Candida auris. *Microbiol Spectr* **12**, e0153124 (2024).

36. Conover MS, Flores-Mireles AL, Hibbing ME, Dodson K, Hultgren SJ. Establishment and
Characterization of UTI and CAUTI in a Mouse Model. *J Vis Exp*, e52892 (2015).

932
933 37. Escandón P, *et al.* Notes from the Field: Surveillance for *Candida auris* - Colombia,
934 September 2016-May 2017. *MMWR Morb Mortal Wkly Rep* **67**, 459-460 (2018).

935
936 38. Barbian HJ, *et al. Candida auris* Outbreak and Epidemiologic Response in Burn Intensive
937 Care Unit, Illinois, USA, 2021-2023. *Emerg Infect Dis* **31**, 438-447 (2025).

938
939 39. Rhodes J, *et al.* Genomic epidemiology of the UK outbreak of the emerging human fungal
940 pathogen *Candida auris*. *Emerg Microbes Infect* **7**, 43 (2018).

941
942 40. Araúz AB, *et al.* Isolation of *Candida auris* from 9 patients in Central America: Importance
943 of accurate diagnosis and susceptibility testing. *Mycoses* **61**, 44-47 (2018).

944
945 41. Chowdhary A, *et al.* New clonal strain of Candida auris, Delhi, India. *Emerg Infect Dis* **19**,
946 1670-1673 (2013).

947
948 42. Gorzalski A, *et al.* The use of whole-genome sequencing and development of
949 bioinformatics to monitor overlapping outbreaks of Candida auris in southern Nevada.
950 *Front Public Health* **11**, 1198189 (2023).

951
952 43. Carolus H, *et al.* Acquired amphotericin B resistance leads to fitness trade-offs that can
953 be mitigated by compensatory evolution in *Candida auris. Nat Microbiol* **9**, 3304-3320
954 (2024).

955
956 44. Burrack LS, Todd RT, Soisangwan N, Wiederhold NP, Selmecki A. Genomic Diversity
957 across *Candida auris* Clinical Isolates Shapes Rapid Development of Antifungal
958 Resistance In Vitro and In Vivo. *mBio* **13**, e0084222 (2022).

959
960 45. Mishra A, Aggarwal A, Khan F. Medical Device-Associated Infections Caused by Biofilm-
961 Forming Microbial Pathogens and Controlling Strategies. *Antibiotics (Basel)* **13**, (2024).

962
963 46. Molina JJ, Flores-Mireles AL. CAUTIon - not all UTIs are the same. *Nat Rev Urol* **22**, 799-
964 814 (2025).

965
966 47. Ahmad S, Khan Z, Al-Sweih N, Alfouzan W, Joseph L. *Candida auris* in various hospitals
967 across Kuwait and their susceptibility and molecular basis of resistance to antifungal
968 drugs. *Mycoses* **63**, 104-112 (2020).

969
970 48. Gulati M, *et al.* In Vitro Culturing and Screening of *Candida albicans* Biofilms. *Curr Protoc*
971 *Microbiol* **50**, e60 (2018).

972
973 49. Schindelin J, *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
974 *Methods* **9**, 676-682 (2012).

976   50.   Dickson MA*, et al.* Human keratinocytes that express hTERT and also bypass a
977   p16(INK4a)-enforced mechanism that limits life span become immortal yet retain normal
978   growth and differentiation characteristics. *Mol Cell Biol* **20**, 1436-1447 (2000).

980   51.   Lamprecht MR, Sabatini DM, Carpenter AE. CellProfiler: free, versatile software for
981   automated biological image analysis. *Biotechniques* **42**, 71-75 (2007).

983