Genetic analysis of F1 cluster phages that infect *Mycobacterium smegmatis* identifies two distinct holin-like proteins that regulate the host lysis event

Richard S. Pollenz[1*], Kira Ruiz-Houston[2], Wynter Dean[1], Loc Nguyen[1]

[1]Department of Molecular Biosciences, University of South Florida, Tampa, Florida, United States of America

[2]Department of Microbiology, University of Wisconsin, Madison, Wisconsin, United States of America

[*]Corresponding Author:

E-mail: pollenz@usf.edu (RSP)

ABSTRACT

Phages Girr and NormanBulbieJr (NBJ) infect Gram-positive *Mycobacterium smegmatis* mc^2

155. Both phages contain conserved lysis cassettes that harbor two endolysin genes (*lysin A*

and *lysin B*) and two genes encoding transmembrane domain (TMD) holin-like proteins. The

first holin-like protein, termed LysF1a is 88 amino acids, has two TMDs and a predicted N-in-C-

in membrane topology. The second, termed LysF1b, has a single N-terminal TMD and a

predicted N-out-C-in topology making it distinct from the type III holins or spanins in size and

membrane topology. Deletion of *lysF1b* results in severe lysis defect phenotypes manifest by

reduced plaque size and changes to lysis timing in liquid culture. Deletion of both *lysF1a* and

lysF1b genes results in phages that show the same lysis phenotypes as the single *lysF1b*

deletion. Phages with only *lysF1b* are lysis competent and trigger lysis prematurely when

exposed to energy poisons while phages with *lysF1a* or *lysF1a/lysF1b* deletions do not trigger

prematurely. Deletion of genes upstream of the lysis cassette did not impact lysis phenotypes.

Lysis recovery mutants were isolated from phages lacking the *lysF1b* gene and these mutants

generated wild type plaque size but triggered lysis prematurely and showed ~65% reductions in

burst size. Genome sequencing identified different point mutations that mapped to TMD1 or the

C-terminal region of the *lysF1a* gene. Infection of an *M. smegmatis* strain that does not produce

lipomannan and lipoarabinomannan by either wild type phages or phages carrying the *lysF1b*

deletion showed modest plaque size increases but did not fully complement the lysis defect of

phages lacking the *lysF1b* gene. Collectively, the findings show that both LysF1a and LysF1b

proteins are required for efficient bacterial lysis by these F1 cluster phages. LysF1a does not

function as a pure antiholin but requires the expression of the LysF1b protein for efficient lysis

functioning.

INTRODUCTION

With an estimated 10^{31} different bacteriophage particles in the world, the pathway whereby dsDNA phages lyse their bacterial hosts after infection and replication is one of the most common cell fates on earth [1]. Extensive biochemical and genetic work with dsDNA phages that infect Gram-negative hosts has established a general three-step model for the triggering of the lysis pathway that involves three distinct sets of proteins. The first are endolysin/s that have defined catalytic activity to lyse the peptidoglycan (PG) layer [2]. The second are holin or antiholin proteins that have between 1-4 transmembrane domain helices (TMD). These proteins work together to, 1) control the timing of the lysis event following phage infection [3, 4], 2) assist with the destabilization of the membrane proton motive force (PMF) [3, 4] and 3) in the canonical holin models, generate large 1μm "pores" in the inner membrane (IM) to allow the exit of the endolysin enzymes to the periplasmic space where they can access and cleave the PG layer [5, 6]. Finally, there are secreted membrane proteins and lipoproteins termed spanins that are associated with the IM and outer membrane (OM). There proteins interact after the PG layer is cleaved to generates the lesion across the IM and OM that allows for the efficient release of the phage progeny [7-9].

dsDNA phages such as lambda, P2, and phage 21 all have genomes < 50,00bp and have a clustered lysis cassette that contain all the genes necessary to mediate the bacterial lysis event [10-12]. Phage lambda encodes the canonical type I holin protein that has 3TMDs and forms large oligomers that create 1μm holes in the IM to allow exit of the endolysin from the cytoplasm to the periplasmic space [12]. Phage 21 defines the type II pinholin lysis pathway and utilizes a 2TMD holin protein that forms ~1,000 heptamers in the IM. The pinholins create only nanometer size holes that are not big enough to allow exit of the endolysin, and they function to disrupt the PMF and trigger lysis. Thus, phages that encode pinholins also encode endolysins that carry secretion-arrest-release (SAR) signals and the SAR-endolysin is held in an inactive

state in the IM prior to triggering. The disruption of the PMF causes the release of the IM

tethered SAR-endolysin to the periplasmic space where it cleaves the PG [4, 13, 14]. In both

lambda and phage 21, a single open reading frame with two in-frame start codons (dual start

motif) encodes two holin proteins that only differ by the addition of 1-2 N-terminal amino acids

[10, 11, 14, 15]. The longer version of each holin functions as the inactive antiholin and in both

cases a percentage of the holin forms dimers with the antiholin that holds that holin in an

inactive state prior to triggering. Upon triggering, the antiholin assumes active holin function

greatly increasing the amount of active holin protein available for hole formation. In phages with

large genomes, such as P1 and T4, the holin, antiholin, endolysin, spanin and lysis inhibiter

(LIN) rI and rIII genes can be bioinformatically characterized throughout the genome and are not

in a defined cassette [10, 11, 16, 17]. The type III holin from phage T4 has a single N-terminal

TMD with an N-in-C-out topology with the bulk of the sequence oriented in the periplasmic

space [16, 17]. T4 phage encodes two different antiholins and LIN proteins that modulate the

holin lysis timing and participate in the lysis inhibition activity of phage T4 [16, 17]. Thus, a

common theme in the phage-mediated lysis pathway of lambda, P1, P2 and T4 is the utilization

of holins and antiholins to regulate the lysis event. Phage T7 on the other hand, has only 1

annotated holin and no identified antiholin and deletion of the holin is not lethal but results in

various lysis phenotypes suggesting that other unidentified proteins may also participate in the

lysis pathway [18]. Another variation of the lysis pathway has recently been shown for the

Enterococcal phage Mu [19]. Mu does not utilize canonical holins and antiholins in the bacterial

lysis pathway and instead uses a novel 1TMD protein termed a releasin protein to release the

SAR-endolysin from the IM so it can cleave the PG [19]. As with lambda and phage 21, Mu

contains canonical spanins to create the final lesion for progeny release once the PG has been

cleaved. It is speculated that Mu does not utilize the holin/antiholin regulated lysis mechanism

because it copies its genome via replicative transposition during the lytic cycle and has less

selective pressure due to transposition immunity [19]. This pathway adds yet another variation

in phage-mediated bacterial lysis and it is striking that all the *E. coli* phages that have been

rigorously studied utilize a similar set of tools to complete the lysis pathway through distinct

molecular mechanisms.

There are 1000s of annotated dsDNA phages that infect Gram-positive hosts, yet the

molecular mechanism that these pages use to lyse their hosts is not resolved to the level of the

E. coli phages. Although it is hypothesized that the lysis mechanism in these phages will utilize

a similar set of proteins as described for phages that infect Gram-negative bacteria, the bulk of

biochemical and genetic work has focused more on the enzymes that target the cell wall than on

the holins, possibly because these enzymes may be employed as antibiotic agents [20-22].

Gram-positive bacteria lack a formal OM and instead have a much more complex PG layer and

an extended structural cell wall components beyond the PG such as arabinogalactans and

mycolic acids [23-26]. Thus, addition of an endolysin to Gram-positive bacteria can result in cell

death. Bioinformatic, biochemical and genetic analysis has identified several types of phage-

encoded endolysins that target different regions of the cell wall. The first type of endolysin is

termed lysin A and targets distinct domains of the peptidoglycan [21, 27]. These enzymes are

highly modular, do not contain canonical SAR domains, and deletion studies show that the lysin

A is essential to phage propagation [28, 29]. It is unknown if the lysin A is released to the

peptidoglycan layer through holins since some studies show that exogenous expression of

selected lysin A enzymes can be cytotoxic to *M. smegmatis* [21, 30-33]. Many phages that infect

Gram-positive hosts also contain a second endolysin termed lysin B that targets the linkages in

the mycolic acid layer of the cell wall [28, 31, 34, 35]. It has been hypothesized that lysin B

provides a fitness advantage for optimal bacterial lysis and phage release since deletion of lysin

B results in viable phages with reduced plaque and burst size [28].

In phages that infect Gram-positive hosts the presence of genes that encode holin-like

proteins has primarily been characterized by bioinformatic analysis and many of the phages

contain more than one gene encoding a TMD protein that could serve as a holin or antiholin.

Currently there are reports on host lysis by phages that infect *Staphylococcus aureus* [36-38],

Bacillus sp. [39,40], *Lactococcal sp*. [41], *Oenococcus oeni* [42, 43], *Streptococcus sp* [44],

Gordonia rubripertincta [45-47] and *Mycobacterium sp*. [48-55]. The holin function for the

encoded TMD proteins in most of these reports is based primarily on the proximity of the TMD

gene/s near an endolysin. Where biochemical work is reported, the majority has been carried

out using exogenous expression of the TMD genes in *E. coli*. However, there are several

reports on phages that infect *Mycobacterium smegmatis* where gene deletion has been utilized

to evaluate lysis function of the predicted holin proteins.

The first study evaluated two genes encoding TMD proteins that are found directly

downstream of the *lysin A* and *lysin B* in the F1 cluster phage Ms6 [50-52]. Ms6 gene *26*

encodes an 88 amino acid protein with 2TMDs while gene *27* encodes a 124 amino acid protein

with 1TMD. Deletion of gene *26* resulted in reduced lysis timing, reduced plaque size but wild

type burst size, while deletion of gene *27* resulted in delayed lysis timing, increased plaque size

and increased burst size. However, there are limitations to these findings because holin function

was not verified using energy poisons and the mutant phages were not tested in liquid lysis

assays in the *M. smegmatis* host versus *E. coli*. In addition, the experiments to assess function

and interaction of the two proteins were carried out using recombinant *E. coli* expression

systems that are not physiologic. Thus, while it appears that both proteins are needed for the

lysis event, it is unclear how each is contributing to the lysis of the *M. smegmatis* host.

The second set of studies evaluated phage D29, a cluster A2 phage that also infects *M.*

smegmatis [53-56]. D29 has a lysis cassette with a *lysin A* (gene *10*), 2TMD holin (gene *11*) and

a *lysin B* (gene *12*). It is important to note that the 2TMD holin of D29 is not an ortholog to the

2TMD protein in Ms6 and differs significantly in amino acid sequence and size. The lysis

cassette that is also distinct from most other actinobacteriophages in that it has no other genes

encoding holin-like TMD proteins within or near the putative lysis cassette. Deletion of gene *11*

results in a viable phage with a different phenotype that the 2TMD deletion in Ms6 [53].

D29_Δ*11* phage shows a modest 30% reduced plaque size, slightly delayed lysis timing and

reduced burst size [53]. The lack of a more severe lysis phenotype is surprising given that gp11

is the only defined holin, especially since the D29 lysin A does not have a defined signal peptide

or SAR-like domain for export [32, 33]. However, it has been reported that the D29 lysin A may

be exported prior to lysis in a yet to be defined pathway that appears to be independent of the

holin [32, 33]. Alternatively, the modest lysis phenotype could also indicate that D29 has

additional holin-like genes that are not located within the defined "lysis cassette" region. Such a

scenario has recently been described in a bioinformatic study of the lysis cassettes of 77

phages that infect the Gram-positive *Gordonia rubripertincta* strain [46]. This study showed that

there was a high level of diversity of the lysis cassette gene organization and that putative holin-

like genes could be identified in genomic regions distal to the endolysin containing lysis

cassettes [46]. This study also showed that most of the phages evaluated had at least two holin-

like genes.

In summary, there is limited biochemical and genetic analysis of host lysis pathways by

phages that infect Gram-positive hosts, especially within the host bacteria. One of the most

intriguing findings from the annotation of phages that infect Gram-positive hosts is the presence

of multiple genes that encode predicted holin-like TMD proteins in the defined lysis cassettes.

Since antiholins have not been experimentally identified in phages that infect Gram-positive

hosts, one interpretation of these data are that holins and antiholins are expressed from distinct

genes as shown for *E. coli* phages P2 and T4 [16, 17, 57, 58]. It is also possible that some of

the TMD genes encode proteins with spanin-like functions that interact with the complex cell

wall of the Gram-positive bacteria. This hypothesis is supported by a report that identified a 177

amino acid protein termed LysZ, that is encoded by phages that infect *Corynebacterium*

glutamicum and may function to disrupt the integrity of the bacterial cell wall [59]. The current

study aims to evaluate how two predicted holin-like TMD proteins encoded by phages Girr and

NormanBulbieJr (NBJ) function in the lysis pathway of *Mycobacterium smegmatis* mc^2 155 by

utilizing recombineered phages and multiple lysis assays under physiological growth conditions.

MATERIALS AND METHODS

Bioinformatic analysis of transmembrane domains and AlphaFold3 analysis

The identification of transmembrane domains (TMD) and protein topologies was carried

out essentially as described by Pollenz et al., 2024 [46] utilizing Deep TMHMM (v1.0.24) [60,

TOPCONS (v2.0) [61], SignalP-5.0 and SignalP-6.0 [62], LipoP 1.0 [61] and HHpred

(databases: PDB_mmCIF70_30_Mar; Pfam-A_v37) [64]. All membrane proteins described in

this report had 100% consensus predictions of TMD domain location, number and protein

topology in all programs. Structural predictions of proteins and protein-protein interactions were

determined using AlphFold3 [65]. For single proteins, pTM scores >50% indicate high

confidence in the model predictions. For protein-protein interactions, ipTM scores <90% are

considered low confidence and should be used with caution [63]. All amino acid alignments

were performed using Clustal Omega [66].

Bacterial strains, growth and plaque assay

The list of bacterial strains, phage and plasmids utilized in this study is presented in

Table 1. *Mycobacterium smegmatis* mc^{2}155 [67] was used as wild type host for all phage

related experiments. *M. smegmatis* ΔmtpA and *M. smegmatis* ΔmtpA-comp are described in

[68]. *M. smegmatis* strains were propagated on Middlebrook 7H10 agar supplemented with 10%

Albumen Dextrose Complex (ADC; 5% albumen, 2% dextrose, 154mM NaCl) at 37^{o}C. To

generate saturated stock cultures, a single colony of *M. smegmatis* was propagated in 25ml of

Middlebrook 7H9 media supplemented with 10% ADC, 1mM CaCl$_2$, 0.5% glycerol and 0.05%

Table 1. Bacterial Strains, Phages and Plasmids

Bacterial Strains	Genotype/Features	Reference/Source
M. smegmatis	*Mycobacterium smegmatis* mc^2 155	67
M. smegmatis Δ*mptA*	*Mycobacterium smegmatis* mc^2 155	68

	lacking the mptA enzyme	
M. smegmatis ΔmptA-comp	*ΔmptA* strain carrying complementing mtpA expression plasmid	68
E. coli	*Escherichia coli* NEB5α F'I^Q	New England Biolabs

Bacteriophage	Genotype/Features	Source
Phage Girr_WT	Mycobacteriophage Girr	70. PhagesDB: https://phagesdb.org/phages/Girr
Phage Girr_*Δ26*	Girr_WT with deletion of gene *26*	This study
Phage Girr_*Δ29*-20	Girr_WT with deletion of gene *29* clone 20	This study
Phage Girr_*Δ29*-25	Girr_WT with deletion of gene *29* clone 25	This study
Phage Girr_*Δ27-30*	Girr_WT with deletion of genes 27-30	This study
Phage Girr_*ΔlysF1a*	Girr_WT with deletion of gene *34*	This study
Phage Girr_*ΔlysF1b*	Girr_WT with deletion of gene *35*	This study
Phage Girr_*ΔlysF1a/_ΔlysF1b*	Girr_WT with deletion of gene *34* and gene *35*	This study
Phage NBJ_WT	Mycobacteriophage NormanBulbieJr	71. PhagesDB: https://phagesdb.org/phages/NormanBulbieJr
Phage NBJ_*Δ29*	NBJ_WT with deletion of gene *29*	This study
Phage NBJ_*ΔlysF1a*	NBJ_WT with deletion of gene *33*	71
Phage NBJ_*ΔlysF1b*	NBJ_WT with deletion of gene *34*	This study
Phage NBJ_*ΔlysF1a/_ΔlysF1b*	NBJ_WT with deletion of gene 32 and gene *33*	This study
Phage D29_	Mycobacteriophage D29	56

Plasmids	Features	Source
pExTra-Girr_*29*	pExTra01::(Ptet-Girr_*34*-mcherry); Kan^R	70
pExTra-Girr_*34*	pExTra01::(Ptet-Girr_*34*-mcherry); Kan^R	70
pExTra-Girr_*35*	pExTra01::(Ptet-Girr_*35*-mcherry); Kan^R	70
pExTra-NBJ_*29*	pExTra01::(Ptet-NBJ_*29*-mcherry); Kan^R	71
pExTra-NBJ_*32*	pExTra01::(Ptet-NBJ_*32*-mcherry); Kan^R	71
pExTra-NBJ_*33*	pExTra01::(Ptet-NBJ_*33*-mcherry); Kan^R	71
pExTra-Waterfoul_*32*	pExTra01::(Ptet-Waterfoul_*32*-mcherry); Kan^R	69

pExTra-Waterfoul_32ΔN	pExTra01::(Ptet-Waterfoul_*32(truncation of N-terminal TMD)*-mcherry); Kan^R	This study
pExTra-Waterfoul_32ΔC-40	pExTra0 ::(Ptet-Waterfoul_*32(truncation of C-terminal 40aa)*-mcherry); Kan^R	This study
pJV53	Acetamide inducible plasmid with Ched9c gene *60-61*. Kan^R	73

Tween-80 with shaking (225rpm) at 37°C. For all liquid lysis and one-step experiments, 100ul of

saturated stock was added to 250ml baffled flasks containing 55ml of supplemented 7H9

without Tween. Cultures were shaken (225rpm) overnight at 37°C and utilized in experiments

the next day while in log phase growth (OD$_{600}$ <0.8). Titering experiments showed that an *M.*

smegmatis culture at an OD$_{600}$ of 1.0 was equal to ~6.5 x 10^7 CFU/ml. This value was utilized

when phages were infected at different MOIs. Plaque assays were completed by mixing the

appropriate amount of phage with 250ul of saturated *M. smegmatis* in Middlebrook 7H9 media

supplemented with 10% ADC, 1mM CaCl$_2$ and 0.5% glycerol. The mixture was vortexed and

incubated at 22°C for 10 minutes. Following the infection, the sample was mixed with 4-5ml of

top agar (Middlebrook 7H9, 1mM CaCl$_2$, 0.5% agar) and plated immediately onto 7H10 agar

plates. Plates were allowed to harden and placed in a 37°C bacterial incubator for 16-48 hrs.

Competent *M. smegmatis* cells for recombineering or for plasmid transfection and

complementation studies were prepared as described previously [29, 69-71]. *Escherichia coli*

NEB5α F′I^Q (New England Biolabs) was used for plasmid amplification and purification and

grown in LB broth or on LB agar supplemented with 50ug/ml kanamycin sulfate and.

Plasmid constructs, *M. smegmatis* transformation and complementation analysis

All PCR primers utilized in this study are shown in S1 Table 1. To generate the inducible

expression plasmids pExTra_Waterfoul_*32ΔN*, pExTra_Waterfoul_*32ΔC40*,

pExTra_Waterfoul_32 was PCR amplified with Q5 DNA polymerase (New England Biolabs)

using Waterfoul specific primers containing with pExTra homology regions (Table S1). Purified

PCR products (were ligated into linearized pExTra01 using isothermal assembly (NEB HiFi 2x

Master Mix) as described [67-69]. Recombinant plasmids were recovered by transformation of

E. coli NEB5α F′I^Q and colonies selected on LB agar supplemented with 50μg/μl kanamycin

sulfate. Plasmid DNA was purified using the GeneJet kit (Fisher) and all plasmids were

sequence-verified by Sanger sequencing using the pExTra Seq primer (Azenta). For generating

complementing strains, competent *M. smegmatis* cells were prepared as described [67-69] and

40μl mixed with 100ng plasmid DNA in a 1mm electroporation cuvette. The sample was

electroporated with 1800V and the time constant set to 5.0 msec. Following electroporation, the

sample was mixed with 950μl of supplemented 7H9 media without Tween and shaken (225rpm)

at 37^{o}C for 120 minutes. Following the recovery period, 10μl-100μl aliquots were spread on

7H10 agar plates supplemented with 10μg/μl kanamycin sulfate and incubated at 37^{o}C for 3-5

days. To generate cultures for plaque assay, single colonies of *M. smegmatis* carrying the

appropriate plasmid were harvested into 25ml of supplemented 7H9 media containing 10μg/μl

kanamycin sulfate. Cultures were shaken (225rpm) at 37^{o}C for 24-48 hours. Saturated cultures

at OD_{600} > 1.5 were centrifuged at 6200rpm for 10 minutes and the pellets resuspended in

supplemented 7H9 media (without Tween) to a final OD_{600} of 2.0. These cells were utilized for

phage infection and then mixed with 4.5ml top agar supplemented with 10ng/μl kanamycin

sulfate with or without 100ng/μl anhydrotetracycline (aTc). The mixture was added to 7H10 agar

containing 10μg/μl kanamycin sulfate supplemented with or without 100ng/μl aTc and incubated

at 37^{o}C for the times indicated in the results section.

Bacteriophage recombineering and genome sequencing

All gblocks and PCR primers utilized in this study are shown in Table S1. The deletion of

genes in phage Girr and NBJ were carried out using the Bacteriophage Recombineering of

Electroporated DNA (BRED) procedure as described [29, 72], with the following modifications.

Phage genomic DNA was purified using the Wizard DNA Clean up kit (Promega). dsDNA

recombineering substrates were synthesized as gblocks (Integrated DNA Technologies). To

generate µg quantities of dsDNA substrates, gblocks were solubilized in water to 10ng/µl and

PCR amplified with gblock specific primers using 2X Q5 master mix (New England Biolabs).

PCR products were purified using Zymoclean DNA recovery (Zymoresearch). To generate

recombinant phage, 200-400ng dsDNA substrate was mixed with 100ng phage genomic DNA

and incubated on ice for 10min. The DNA mixture was added to 100µl of recombineering *M.*

smegmatis in a 2mm electroporation cuvette and electroporated at 2500V with the time constant

set to 20.0 msec. Following electroporation, 900µl of supplemented 7H9 media was added to

the cells and they were shaken (225rpm) at 37^{o}C for 90 minutes. Following the recovery period,

the sample was mixed with 250µl of saturated *M. smegmatis* and plated with 5ml 7H9 top agar

onto 7H10 agar plates. Plates were incubated at 37^{o}C for 18-24 hours to recover primary

plaques. Primary plaques were picked into 120µl of phage buffer (10 mM Tris-HCl, pH 7.5; 10

mM MgSO$_4$, 68.5 mM NaCl; 1mM CaCl$_2$), vortexed, and incubated at 22^{o}C for 2-8 hrs. The

presence of mutant phage genomes was determined by Deletion Amplification Detection Assay

(DADA) PCR [29] using a 20µl reaction with: 1µl of plaque sample, 2µl of 0.2 µM DADA primer

mix (1:1 mix of For and Rev primers), 5µl glycerol (40%), 2µl water and 10µl OneTaq 2x Master

Mix (New England Biolabs). PCR conditions were: 94^{o}C for 1 minute followed by 35 cycles of

94^{o}C-1min/66^{o}C 1min and a final 66^{o}C step for 5 minutes. Positive primary plaque samples

were serially diluted and plated on wild type or complementing *M. smegmatis* as indicated.

Secondary plaques were picked as above and screened by standard PCR using gene specific

flanking primers. Pure populations of mutant phage were evaluated by full genome sequencing

as described [73, 74]. Briefly, Genomic DNA was used to create sequencing libraries with the

NEB XLEP-P1 kit. Sequencing was performed by the Pittsburgh Bacteriophage Institute and the

library run on an Illumina NextSeq 1000 instrument. Raw reads were trimmed with cutadapt 4.7

(using the option: –nextseq-trim 30) and filtered with skewer 0.2.2 (using the options: -q 20 -Q

30 -n -I 50) prior to assembly. The resulting sequences were assembled using Consed (v29.0)

with Unicycler (v5.0) and contigs checked for completeness, accuracy, and genome termini.

Final phage contigs were aligned to wild type phage to identify deletions and point mutations.

Plaque size analysis

Plaques were generated using the plaque assay procedure described above. For

experiments where plaque size would be evaluated between several different phages, identical

bacterial samples, incubation times, phage concentrations and top agar amounts were utilized.

Plaque size was evaluated using ImageJ software [75] and by manual measurement using

photos embedded in PowerPoint and custom rulers. In most experiments, results are presented

for an average of 50 plaques and the statistical significance of measurements determined by

unpaired t-test (https://www.graphpad.com/quickcalcs/ttest1/). Plaque volume was calculated by

the formula $\pi r^2 h$ where r is the radius of the plaque and h was set to 1mm for the top agar

thickness.

Liquid lysis assay and cell viability assay from liquid cultures

Liquid lysis assays were carried out essentially as described by Payne et al., 2009 [28]

with the following modifications. Briefly, overnight cultures of *M. smegmatis* in log phase growth

were diluted to an OD_{600} of 0.25-0.30 in supplemented 7H9 without Tween and 40ml of culture

was added to 250ml baffled flasks. High titer phage lysate was added to the cultures to create

an MOI of 10 and the mixture set at 37°C for 30 minutes without shaking. T_0 for all experiments

was the time of phage addition. After 30 minutes (T_{30}), the cultures were constantly shaken a

225rpm and aliquots of cells removed with a syringe and sterile needle from the shaking culture

without stopping the incubator. The OD_{600} of the sample was immediately recorded for each

aliquot using a GENESYS spectrophotometer (Fisher Scientific). In some experiments, KCN

(1M in water), or $CHCl_3$ (100%) was added directly to the shaking culture to produce a final

concentration of 10mM or 1%, respectively. All liquid lysis experiments were repeated at least

five separate times and representative experiments are presented.

To assess if there were viable cells surviving in the phage infected cultures, bacteria was

aliquoted from a single stock sample at an OD_{600} of 0.25-0.27 and infected at an MOI of 10 as

described above. An aliquot of culture was removed from uninfected and infected cultures at the

times specified in the experiment, evaluated at OD_{600} and serial diluted in 7H9 neat. 250µl

aliquots of the dilutions were spread on 7H10 agar plates and incubated at $37^{o}C$ until colonies

could be counted (4-5 days). To assure that surviving colonies all arose from infected cells and

were not the result of phage resistant *M. smegmatis* or contamination, lytic phage D29 [56] was

used as a positive control with an expected outcome of zero viable cells.

ATP assay

The presence of ATP in culture supernatants was determined utilizing the ENLIGHTEN

kit (Promega). Briefly, liquid lysis assays were carried out as described above. A 500ul aliquot of

culture was removed at indicated time points and immediately passed through a 0.2µ syringe

filter to remove bacteria. The sample was frozen at $-20^{o}C$ prior to analysis. The presence of ATP

in culture supernatants was determined by mixing 5µl with 50µl ENLIGHTEN reagent

(Promega), mixing for 5 seconds and then reading the integrated luminescence over 10

seconds using a GloMax luminometer (Promega). To assure that the readings were within the

linear range of the instrument an ATP standard curve was utilized.

One-step growth curves and burst size determination

Several experimental procedures have been utilized for one-step growth curves with

Gram-positive bacteria [33, 49, 51, 53]. In our hands, none of these procedures generated

consistent results or provided the ability to directly measure burst size. Thus, we developed a

one-step protocol that is illustrated in S2 Figure 1 and described below. To prepare the cells, the

OD_{600} of fresh overnight cultures of *M. smegmatis* in log growth were determined and 10ml

centrifuged at 3,000rpm for 10 minutes. The cells were resuspended in supplemented 7H9 to a

final concentration equal to an OD_{600} of 3.2/ml. 250µl of cells (~5 x 10^7 cfu) were added to

microfuge tubes and ~5 x 10^7 phage added to the cells to create an MOI of 1. The same amount

of phage was also added to 250µl of supplemented 7H9 media without cells to verify the titer

and determine the level of phage adsorption. The mixture was vortexed and set at 37^{o}C for 30

minutes without shaking. T_0 for all experiments was the time of phage addition to the cells. After

30 minutes, 750µl of supplemented 7H9 was added to all samples. Samples were vortexed and

centrifuged at 6,200rpm for 4 minutes to separate unabsorbed phage. The supernatant was

saved for analysis of titer and adsorption, and the cell pellet immediately resuspended in 1ml

supplemented 7H9, vortexed and diluted 1:10,000 into 70ml of supplemented 7H9 in baffled

250ml flasks. An aliquot was removed and 5µl, 12.5µl and 25µl used for plaque assay to

determine the number of infectious centers to determine the precise number of infected cells. A

second aliquot was removed and filtered through a 0.2µ syringe filter and 150µl utilized for

plaque assay to test for unabsorbed (free) phage at T_{45} minutes (the first one-step data point).

The flask was placed in a 37^{o}C incubator and constantly shaken at 225rpm for the duration of

the experiment. At specified time points, 500µl aliquots were removed from the shaking flasks

and each sample immediately passed through a 0.2µ syringe filter to remove cells and collect

free phage. The amount of phage in each sample was determined by performing plaque assays

using 150µl of serially diluted samples (usually 10^0 to 10^{-4} dilutions were sufficient). The phage

titer in the flask, expressed as PFU/ml was determined at each time point by counting the

plaques on each plate, dividing by the volume of phage used in the infection (150µl), multiplying

by the dilution factor and then multiplying by 10^3. The number of infectious centers in the

infected 70ml culture was determined by counting the plaques, dividing by the volume used for

the plaque assay and multiplying by 70,000. The original phage titer and the percentage of

adsorbed phage was determined by spot titers and plaque assay of the serially diluted

supernatants collected from the samples of phage with and without the added cells. Since the

number of cells and phage used in these experiments was held constant, this procedure

consistently produced ~3.5-5.5 x 10^7 infectious centers. This procedure also resulted in

essentially zero unabsorbed phage in the culture at the start of the one-step assay. Burst size at

all time points could be directly determined by dividing the total phage in the 70ml culture by the

number of infectious centers. All phages were evaluated through a minimum of three separate

assays. A representative data set showing the raw data for adsorption, infections centers and a

one-step assay is provided in S3 Figure 2 and S4 Figure 3.

RESULTS AND DISCUSSION

Topologies of Holin-like TMD Proteins Within the Lysis Cassettes of F1 Cluster Phages

Girr and NormanBulbieJr.

To investigate the contribution of predicted holin-like proteins in the lysis of *M.*

smegmatis, it was desirable to begin work in phages that had a defined lysis cassette and prior

wet-lab data. F1 cluster phages Girr and NormanBulbieJr (NBJ) were selected since both

phages have a define lysis cassette and a full genome cytotoxicity screen has been reported for

both phages [70, 71]. Girr and NBJ have an average nucleotide identity of 89.86% but only

share 69 of 102 genes. Both phages have a defined lysis cassette organization located

downstream of the *tape measure* and *minor tail* genes and encode a Lysin A enzyme predicted

to target the peptidoglycan layer and a Lysin B that targets the linkage of the arabinogalactins to

the mycolic acids (Figure 1A). The *lysin B* gene is directly followed by two genes encoding

proteins that have predicted TMDs (Girr *34* and *35* and NBJ *32* and *33*).

It was first important to bioinformatically evaluate the predicted topologies of these TMD

proteins and determine how they compared to previously characterized holins or other phage

proteins involved in bacterial lysis. Girr *34* and NBJ *32* encode small 80aa and 77aa proteins,

respectively, that have two predicted TMD regions and an N-in-C-in topology by all TMD

prediction programs (Figure 1B). The proteins are 97% identical and have cytoplasmic N and C-

terminal domains that each contain four charged amino acids. The presence of two TMDs and

the N-in-C-in topology is like the phage 21 S^{21} type II pinholin (Figure 1BC). However, the

presence of a pinholin is typically associated with a SAR-endolysin that is secreted to the

periplasmic space prior to lysis triggering [10, 11, 13]. Analysis of the Girr or NBJ Lysin A

proteins by the various TMD prediction programs and AlpahFold3 (S5 Figure 4), show no

evidence of SAR domains or signal sequences and this is consistent with previous analysis of

Lysin A proteins from phages that infect *M. smegmatis* [21]. Thus, it is critical to contrast several

structural differences of gp32 and gp34 to the canonical $S^{21}68$ pinholin and $S^{21}71$ antiholin.

First, the $S^{21}68$ holin protein has only 3 N-terminal amino acids prior to the TMD1 and

this feature is a critical to its function as a holin and allows the TMD1 to become cytoplasmic

(Figure 1BC) [11-15]. Second, the S^{21} gene has a dual start motif that produces both the $S^{21}68$

holin protein and the $S^{21}71$ antiholin after phage infection [11-15]. The difference in the two

proteins is the addition of the extra basic K residue to the N-terminal cytoplasmic domain of

$S^{21}71$ that limits the TMD1 from exiting the membrane to the same degree as the $S^{21}68$ holin

(Figure 1BC; [11-15]). Thus, the expression of $S^{21}71$ keeps a percentage of the $S^{21}68$ in an

inactive dimer until triggering at which time the $S^{21}71$ converts to an active holin [13-15, 76, 77].

Third, the $S^{21}68_{IRS}$ protein is an engineered version of $S^{21}68$ that adds the MRYIRS motif and 4

charged residues in the N-terminal cytoplasmic domain that creates a dominant negative

version of the antiholin that is poorly triggered to an active holin [76]. Neither Girr gene *34* nor

NBJ gene *32* have a dual start motif that generates multiple products of the same open reading

frame. Genes *34* and *32* encode a single protein product that has a predicted N-terminal

cytoplasmic domain of 13-16 amino acids of which four are charged and two others are polar.

Therefore, the N-terminal topology and the number of charged amino acids in Girr gp34 and

NBJ gp32 are more similar with the $S^{21}71$ antiholin and $S^{21}68_{IRS}$ and would predict that TMD1 of

these proteins remains embedded in the membrane. In addition, the TMDs of S^{21} both contain

GxxxG motifs that are predicted to contribute to homotypic and/or heterotypic helix interactions

of the monomers that facilitate the formation of the pinholes [76, 77]. While TMD1 and TMD2 of

Girr and NBJ have a high percentage of GA residues (Figure 1B), they lack defined GxxxG

domains found in S^{21}. Finally, although AlphaFold3 does not model membrane proteins as well

as soluble proteins, Girr gp34 was not modeled with high confidence into either a monomer or a

heptamer compared to $S^{21}68$ (S6 Figure 5). Taken together, this data suggests that Girr gp34 or

NBJ gp32 are unlikely to function as a type II holin. Based on the literature we will designate

these proteins as LysF1a due to a predicted function in lysis (lys) and being first identified in F1

cluster phages.

Since there is a second gene encoding a protein with a single TMD directly downstream

of the *lysF1a* gene, we hypothesized that this protein may also be involved in the lysis pathway.

Girr *35* and NBJ *33* both have -4bp ATGA overlaps to the upstream *lysF1a* gene and encode

98% identical 124aa proteins with a single N-terminal TMD and a predicted N-out-C-in topology

by all TMD prediction programs (Figure 2A). Both proteins have a 22 amino acid extracellular

domain followed by the single TMD. There are four basic residues following the TMD and

AlphaFold3 predicts an extended helical cytoplasmic C-terminal domain that contains ~50%

KRDE charged residues typical of holins [10, 11]. Figure 3B shows the AlphaFold3 models and

predicted topologies of representative proteins with a single TMD that are implicated in phage-

mediated bacterial lysis compared gp33 and gp35. Most pertinent is the type IIIa holin typified

by the T protein of phage T4 that forms large 1um holes like canonical type I holins [16, 17] and

the recently identified type IIIb holin identified in jumbophage PhiKZ [78]. Neither type III holin

has >20% amino acid identity to gp33 and gp35, and both are much larger with the T having the

opposite membrane topology with a globular extracellular domain critical to its holin function

(Figure 2B). Although the PhiKZ holin has an extended cytoplasmic domain, the protein has a

C-out-N-in topology and also has a globular cytoplasmic domain. Nether have the highly

charged cytoplasmic regions found in the proteins, It noteworthy that spanins also have a single

TMD and model with an extended helical prediction, but like the T holin, these proteins are also

oriented in a N-in-C-out topology that has been shown experimentally and is also predicted by

the TMD prediction programs (Figure 2B; [10, 11]). Interestingly, there is a recent report

regarding phages that infect the Gram-positive *Corynebacterium glutamicum* strain. The authors

have characterized a protein with 1 TMD that is termed LysZ and is proposed to interact with

lipid-anchored glycans on the extracellular side of the IM to promote efficient lysis [59].

However, like the type IIIa holin, LysZ is much larger than the Girr and NBJ proteins and it is

proposed to have a N-in-C-out topology that allows it to interact with the lipid-anchored glycans

[59]. Thus, Girr gp35 and NBJ gp33 are most similar in structure and membrane topology to the

recently identified phage Mu releasin protein (Figure 2B) [19]. Releasin however, is named due

to its function to release the Mu SAR-endolysin from the membrane in a holin independent

manner [19]. Mu also has <20% amino acid identity to gp33 and gp35 and does not possess the

highly charged cytoplasmic domain. Thus, of all the characterized proteins involved in lysis that

have 1 TMD, gp35 and gp33 appear to represent a novel class of proteins and will be termed

LysF1b since the gene is downstream of the *lysF1a*.

Deletion of Individual *lysF1a* and *lysF1b* Genes in F1 Phage are Viable and Results in

Reduced Plaque size

To evaluate the function of the *lysF1a* and *lysF1b* genes, the BRED recombineering

method was utilized to delete them in both Girr and NBJ as described in Methods. Several

primary plaques were positive for each deletion and pure populations of Girr_*ΔlysF1a*,

Girr_*ΔlysF1b*, NBJ_*ΔlysF1a*, and NBJ_*ΔlysF1b* were recovered from secondary plaques. Figure

3 shows the lysis cassette organizations of the Girr mutants and the PCR verification of the

gene deletions from clonal high titer lysates. The data for NBJ is shown in S7 Figure 6 and

deletion of gene *33* is also presented in Wise et al. [71]. The genome structure of all deletions

was confirmed by sequencing the immediate region of the deletion and by whole genome

sequencing to validate that there were no point mutations outside of the targeted gene deletion

region.

To assess the phenotypes of the mutant phage, equal concentrations of Girr_WT,

Girr_*ΔlysF1a* and Girr_*ΔlysF1b* were used to infect *M. smegmatis* and the total number of

recovered plaques and plaque sizes evaluated. All phages formed plaques with equivalent

efficiencies under the standard assay conditions described in Methods suggesting no defects in

adsorption (S8 Figure 7). However, both Girr_*ΔlysF1a* and Girr_*ΔlysF1b* presented with

significantly smaller plaques than Girr_WT (Figure 4) as did NBJ_*ΔlysF1a* and NBJr_*ΔlysF1a*

(S9 Figure 8). A summary of the plaque size data for both Girr and NBJ is presented in Table 2.

Deletion of the *lysF1a* gene from Girr or NBJ results in ~40% reduction in plaque diameter and

Table 2 Plaque sizes in WT Girr and *lysF1a* and *lysF1b* deletion mutants.

PHAGE	Plaque size (mm)	% of WT (diam.)	Plaque (vol, mm^3)	% of WT (plaque vol)
NBJ_WT	3.49 +/- 0.533	100	11.479	100
NBJ_*ΔlysF1b*	1.23 +/- 0.225*	35.24*	1.425*	12.42*
NBJ_*ΔlysF1a*	2.05 +/- 0.660*	58.73*	3.96*	34.50*
Girr_WT	3.88 +/- 0.731	100	14.188	100
Girr_*ΔlysF1b*	1.28 +/- 0.376*	32.98*	1.544*	10.88*
Girr_*ΔlysF1a*	2.31 +/- 0.719*	59.53*	5.029*	35.44*

451 *M. smegmatis* was infected with identical numbers of phage and plaque size determined after
452 36 hrs. of growth at 37°C. Plaque diameter was measured using ImageJ. Data is presented as
453 average +/- standard deviation for a minimum of 50 individual plaques. * = statistically significant
454 from Girr or NBJ WT phage *p<0.0001.*
455

a 65% reduction in plaque volume, while deletion of the *lysF1b* is more severe showing a ~65%

reduction in plaque diameter and near 90% reduction in plaque volume. To validate that these

plaque size reductions were directly related to the loss of *lysF1a* or *lysF1b* genes, *M. smegmatis*

containing complementing expression plasmids were infected with the mutant phages and

plaque size evaluated as described in Methods. Since both Girr LysF1b and NBJ LysF1b are

highly toxic when exogenously expressed in *M. smegmatis* [70, 71], complementation of the

lysF1b deletions first utilized Waterfoul gp32. Waterfoul is a K5 cluster phage with a canonical

lysis cassette and gene *32* encodes a 1TMD with the same general size and membrane

topology as Girr LysF1b (Figure 2D) and is not toxic when induced by aTc in *M. smegmatis* [67].

When M. *smegmatis* transfected with Waterfoul gp32 is infected with Girr_*ΔlysF1b* in the

presence of aTc, there is a significant increase in plaque size compared to untreated cells

(Figure 5A). Importantly, the increase in plaque size was not observed when the N-terminal

TMD domain or the highly charged C-terminal 40 amino acids of Waterfoul gp32 were truncated

(Figure 5A). Thus, the complementation by Waterfoul gp32 requires the full-length protein.

Recently it has been reported that the pTet promoter in pExTra is leaky and that it is possible to

complete complementation analysis in the absence of aTc [71]. Thus, *M. smegmatis* was

transfected with pExTra-Girr*35* and pExTra-NBJ*33* and the resulting cells infected with

Girr_Δ*lys1b* as described in Methods. Figure 5B shows that there is a significant increase in

plaque size when Girr_Δ*lys1b* is plated on the complementing strains expressing the LysF1b

proteins compared to cells with the base pExTra01 vector. These results support that the

reduced plaque size of Girr_Δ*lysF1b* is directly attributable to the loss of *lysF1b.* Similarly, the

wild type plaque size is restored when *M. smegmatis* harboring the pExTra-Girr_*34* plasmid is

infected with Girr_Δ*lysF1a* (Figure 6). Thus, the loss of either *lysF1* gene appears to create a

lysis defect phenotype that is like the phenotype when the *lysin B* is deleted in phage Giles and

the cell wall components remain a barrier to efficient phage release [28].

Deletion of Individual *lysF1* Genes Result in Distinct *M. smegmatis* Liquid Lysis

Phenotypes and ATP Release.

A main function of holins is the control of the lysis timing by the destruction of the proton

motive force (PMF) [10, 11]. To assess lysis timing of Girr, NBJ and the *lysF1* mutants, liquid

lysis assays were performed where the optical density (OD_{600}) of infected log growth *M.*

smegmatis cultures is measured over time. Importantly, in all studies, once phage had been

allowed to absorb for 30 minutes, cultures were continually shaken even while samples were

withdrawn so that lysis would not be prematurely triggered by changing the culture conditions

[11, 79]. The data show that cultures infected with Girr_WT grow steadily until ~110 minutes

post infection. Beginning at 110 minutes, there is a significant drop in the OD_{600} of the culture

(triggering event) until it reaches a baseline of OD_{600} 0.10 at about 200 minutes (Figure 7A). In

contrast, Girr_$\Delta lysF1a$ grows until ~130 minutes and then shows a reduction in OD_{600} until 300

minutes that does not reach the same OD_{600} baseline as Girr_WT even after 480 minutes of

incubation (Figure 7A). Strikingly, cultures infected with Girr_$\Delta lysF1b$ do not show a drop in

OD_{600} but appear to cease growing after ~200 minutes with only a slight decrease in OD_{600} over

the remaining growth period out to 480 minutes (Figure 7A). For NBJ_WT, OD_{600} begins to drop

after 130 minutes and like Girr_WT reaches a baseline of around OD_{600} 0.10 (Figure 7B).

NBJ_$\Delta lysF1a$ shows a delay in lysis timing like Girr_$\Delta lysF1a$, but the delay is more pronounced

as the OD_{600} does not begin to drop until ~180 minutes post infection (Figure 9B). Just like

Girr_$\Delta lysF1b$, NBJ_$\Delta lysF1b$ ceases to grow after ~200 minutes and does not show a significant

reduction in OD_{600} even after 480 minutes of growth (Figure 7B). Thus, in both phages, deletion

of *lysF1a* results in a triggering delay while deletion of *lysF1b* results in no defined triggering

event in this assay. It is interesting that Girr_WT and NBJ_WT show a ~20-minute difference in

triggering time and that a difference is also observed between Girr_$\Delta lysF1a$ and NBJ_$\Delta lysF1a.$

These results may be due to differences in how the lysis cassette genes are expressed since

Girr has an HNH endonuclease gene between the *lysin A* and *lysin B* (Figure 1A) and the

intergenic region between the *lysin B* gene and the *lysF1a* is 8bp in NBJ but 38bp in Girr (S10

Figure 9).

ATP release has been used as a measure of cell viability and membrane integrity

following phage infection [27, 53]. Figure 7C shows the ATP release from the same experiment

that is presented in Figure 7AB. As expected, ATP release for Girr_WT, NBJ_WT, Girr_$\Delta lysF1a$

and NBJ_$\Delta lysF1a$ generally mirrors the time at which lysis is triggered and the culture OD_{600}

begin to decline. ATP levels in the media also peak when the OD_{600} baseline is reached at about

300 minutes (Figure 7C). In contrast, there is delayed but gradual increase in the amount of

ATP released from the cultures infected with Girr_$\Delta lysF1b$ and NBJ_$\Delta lysF1b$ out to 300

minutes. Although the ATP level does not reach that of Girr_WT or NBJ_WT, the release of ATP

confirms that cell integrity has been comprised after infection with the *lysF1b* mutants. To

directly assess this contention, aliquots of the cultures were removed at 100- and 260-minutes

post infection, serial diluted and plated on 7H10 agar to assess colony growth as detailed in

Methods. Both Girr and NBJ are temperate phages, thus, it was expected that the infected

cultures would show a small percentage of surviving lysogens as reported in Wise et. al. 2025.

A representative data set with NBJ is shown in S11 Figure 10. Girr showed similar results. Cells

infected with NBJ_WT or NBJ_*ΔlysF1b* show >90% reduced colonies 100-minutes post

infection when compared to the uninfected culture that was essentially at the same OD_{600}. When

the cultures were reevaluated at 260-minutes post infection, the colony counts for both NBJ_WT

and NBJ_*ΔlysF1b* show a slight 1.4-fold rise due to lysogeny growth, but are still 90% reduced

in viable cells compared to the number at the start of the experiment. The lytic phage D29 kills

all cells in the culture, validating that the surviving cells are not the result of infection resistant

cells or contamination. Thus, this data clearly supports the hypothesis that the lack of a defined

OD_{600} decline in the *ΔlysF1b* infected cells is due to inefficient lysis and not due to having more

viable cells than those infected with Girr_WT.

Collectively, the important finding from this data set is that an F1 phage with only the

1TMD LysF1b protein can trigger the lysis event in the absence of the 2TMD lysF1a protein, but

that the lysF1a protein does not support significant bacterial lysis as measured by a drop in

OD_{600}. The lack of OD_{600} decline is not due to there being a high population of viable cells,

although there is a small percentage (<10% of the starting cell population) of surviving lysogens.

In fact, the liquid lysis results for Girr_*ΔlysF1b* and NBJ_*ΔlysF1b* is like the lethal *lysin A*

deletion in phage Giles where the cultures stopped growing after infection and the OD_{600} gently

rose and then plateaued but did not show a sharp triggering decline up through 360-minutes

post infection [28]. The similarity in these results suggest that the lysis defects in the *ΔlysF1b*

phage may be related to the inability of the endolysins to efficiently cleave the cell wall

components because they are either not efficiently released from the infected cell or are

exported but cannot be released from the membrane or activated in some manner.

Energy Poisons Cause Early Triggering of F 1Cluster Phages with Deletion of the *lysF1a*

Gene but not the *lysF1b* Gene in Liquid Lysis Assays

The results presented so far indicate that the expression of both TMD proteins are

required for phage viability, but phages that express only the LysF1b protein can support

bacterial lysis while phages that express only LysF1a are viable but severely lysis

compromised. The diagnostic assay for holin function is the ability to cause early triggering of an

infected culture by the addition of energy poisons (e.g., cyanide or the protonophore

dinitrophenol) due to the disruption of the proton motive force (PMF) [11, 79]. This type of assay

has been utilized in nearly all the pioneering work on holin function of phages that infect *E. coli.*

[10, 11, 19] and both Girr_WT and NBJ_WT can be triggered early if infected cells are treated

with CN (S12 Figure 11). To assess the effect of CN on the various mutants, *M. smegmatis* was

infected with Girr_WT or Girr_*ΔlysF1a* and the OD_{600} of the culture measured as described in

Methods. At 110 minutes after infection, KCN (final concentration of 10mM) was added directly

to one culture of Girr_*ΔlysF1a* and to a culture of uninfected cells. Figure 8A shows that

Girr_WT triggers at ~110 minutes while Girr_*ΔlysF1a* triggers at 130 minutes just as in Figure

9A. However, when KCN is added to a culture infected with Girr_*ΔlysF1a*, there is a drop in

OD_{600} beginning at ~120 minutes that clearly precedes the drop in the untreated Girr_*ΔlysF1a*

culture. KCN addition to uninfected cells results in immediate cessation of growth, but not a

significant drop in OD_{600}.To validate these results, studies were repeated with the NBJ mutants.

NBJ_WT triggers later than Girr_WT at ~130 minutes and declines to OD_{600} 0.10 by 300

minutes as expected (Figure 10B). NBJ_*ΔlysF1a* shows delayed triggering just as in Figure 8B

and addition of KCN to a culture infected with NBJ_*ΔlysF1a* at 120 minutes results in lysis

triggering by 130 minutes with a gradual reduction in OD_{600} that mirrors the NBJ_WT. The ability

of Girr_*ΔlysF1a* and NBJ_*ΔlysF1a* to trigger lysis, identifies the 1TMD LysF1b protein as a

putative holin and not a releasin, since energy poisons do not trigger lysis prematurely in phage

Mu [19].

570   The next set of studies focused on the phages with deletion of the 1TMD *lysF1b* genes

to determine if they could also be triggered by the addition of KCN. Figure 8C shows the results

for Girr_*ΔlysF1b*. As previously shown in Figure 8A, *M. smegmatis* infected with Girr_*ΔlysF1b*

does not exhibit a defined triggering event as the cells simply cease growing with a slight

reduction in OD_{600} out to 480 minutes. Addition of KCN to the Girr_*ΔlysF1b* culture at 180

minutes results in a modest 30% reduction of OD_{600} between 180-240 minutes compared to the

Girr_*ΔlysF1b* untreated culture. However, following this initial decline, the level of change in

OD_{600} generally mirrors that of the untreated Girr_*ΔlysF1b* culture and the KCN treated *M.*

smegmatis control. In contrast, addition of 1% $CHCl_3$ to the Girr_*ΔlysF1b* culture at 180 minutes,

results in an immediate precipitous drop in OD_{600} to the level of Girr_WT indicating that the

endolysin enzymes are present and active in the cells. The results for NBL_*ΔlysF1b* are shown

in Figure 8D and exhibit near identical results to Girr_*ΔlysF1b* in that addition of KCN cause a

modest drop during the first hour after addition, and then an OD_{600} decline that mirrors the

untreated NBJ_*ΔlysF1b* culture and the KCN treated *M smegmatis*. Addition of $CHCl_3$ results in

the same type of sharp decline in OD_{600} to the level of the NBJ_WT baseline. Collectively, these

results show that phages expressing only the 2TMD LysF1a protein are not efficiently triggered

following disruption of the PMF but that the cells have the requisite enzymes for lysis since lysis

occurs immediately after addition of 1% $CHCl_3$. The inability of the LysF1a proteins to be

triggered by KCN is like the $S^{21}71$ or $S^{21}68_{IRS}$ antiholins [13, 76, 77].

F1 Cluster Phage Deleted for *lysF1a* have Delayed Burst Timing While Phage Deleted for

***lysF1b* Deletion have Reduced Timing and Greatly Reduced Burst Size**

The previous results show that infection of *M. smegmatis* with either Girr_*ΔlysF1a* or

NBJ_*ΔlysF1a* results in a defined triggering (lysis) event that can be prematurely initiated by

addition of KCN. However, the results do not establish if phage have been efficiently released

from the lysed cells, the timing of the burst event and the burst size. Since it is not possible to

utilize lysogens to study phage release in *M. smegmatis*, one-step grow curves were utilized to

evaluate the various *lysF1a* and *lysF1b* mutant phages. The experimental set up for these

studies is fully described in the methods and representative raw data sets are also shown in S3

Figure 2 and S4 Figure 3. The one-step results from three different experiments utilizing Girr

phages are presented in Figure 9. Phage release from Girr_WT shows a lag time of ~180

minutes from the time of infection and then a gradual phage release for 90 minutes before

reaching a plateau at ~300 minutes (Figure 9A). The slow release of phage over a 90-minute

period is consistent with the liquid lysis assays that showed an ~60-minute timeline of OD_{600}

decline after triggering. This is an important observation because there should not be a sharp,

immediate lysis of all cells and accompanying phage release since the phage infections are not

synchronized but carried out over a 30-minute period and the efficient release of phage from

Gram-positive bacteria may also be hampered by the extensive cell wall. Girr_*ΔlysF1a* shows a

delayed lag time by ~20 minutes compared to Girr_WT and a more prolonged period of phage

release over 150 minutes before reaching a plateau. The overall burst size for Girr_WT over

three experiments was 277 +/- 81 PFU/cell. Girr_*ΔlysF1a* showed a lower burst of 255 +/- 75

PFU/cell but this was not statistically different from the for Girr_WT. In contrast, Girr_*ΔlysF1b*

shows a very small but defined burst at ~ 210 minutes that is best illustrated when the data is

plotted independent of the other phages (Figure 9B). The overall burst size for Girr_*ΔlysF1b* is

only ~29.2 +/- 36 PFU/cell suggesting very poor efficiency of phage release from the infected

cells. Overall, this data is consistent with the plaque size and liquid lysis results and supports

the hypothesis that *M. smegmatis* integrity is compromised when infected with Girr_*ΔlysF1b* but

the release of phage progeny is severely impacted. In addition, the near wild type burst size

observed in cells infected with Girr_*ΔlysF1a*, further support the holin function of the LysF1b

protein.

Deletion of both *lysF1a* and *lysF1b* is not Lethal to Phage Viability and Shows Similar

Phenotypes to the Δ*lysF1b* Mutants

Since deletion of either of the individual TMD genes resulted in viable phage, it was of

interest to test whether a double deletion of both TMD genes was viable. Both Girr and NBJ

were used for these studies and the NBJ data set is shown in Figure 10. BRED was utilized was

utilized to delete *lysF1b* from NBJ_*ΔlysF1a* DNA. Figure 11A/B shows the genomic context of

the double deletion and the PCR confirmation that both *lysF1a* and *lysF1b* is deleted. The

deletion was confirmed by Sanger sequencing of the deleted region and full genome

sequencing confirmed that there we no additional nucleotide changes to the genome. A viable

double deletion of *lysF1a* and *lysF1b* was also produced in phage Girr that showed identical

results to NBJ. Interestingly, while NBJ_*ΔlysF1a/ΔlysF1b* had dramatically reduced plaque size

compared to the NBJ_WT phage, the plaques were essentially the same size as the single

mutation NBJ_*ΔlysF1b* phage (Figure 10C). Liquid lysis data in Figure 10D show that NBJ_WT

triggers lysis as expected at ~130-140-minutes while NBJ_*ΔlysF1b* and the

NBJ_*ΔlysF1a/ΔlysF1b* phage causes reduced culture growth but no defined triggering.

Importantly, the addition of KCN was able to prematurely trigger the NBJ_WT but did not trigger

the NBJ_*ΔlysF1b* or trigger NBJ_*ΔlysF1a/ΔlysF1b.* The finding that the double deletion of

lysF1a and *lysF1b* is viable and has essentially the same phenotypes as the NBJ_*ΔlysF1b*,

support the hypothesis that the LysF1a protein does not contribute to lysis in the absence of

LysF1b. These findings also suggest that the F1 phage encoded lysins can be released from

the cell independent of LysF1a and LysF1b proteins as suggested for Mycobacteriophage D29

and others [21, 30-32],

Lysis Recovery Mutants Isolated from Girr_ΔlysF1b or NBJ_ΔlysF1b Show Wild Type

Plaque Size and have Point Mutations in the *lysF1a* Gene

Since both Girr_ΔlysF1b and NBJ_ΔlysF1b show significant plaque size defects it was

possible to screen for lysis recovery mutants (LRM) by assessing changes to the small plaque

size following multiple rounds of infection and replication. Figure 11A shows the presence of a

large plaque in a Girr_ΔlysF1b plaque assay after the second round of harvest and infection.

Table 3. Lysis Recovery Mutants (LRM) from ΔlysF1b Phages.

Phage	Type of mutation	Genetic location	Mutation Result
Girr_ΔlysF1b-LRM1	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_ΔlysF1b-LRM2	point mutation-nucleotide substitution	gene *35 (lysF1a)*	R19H
Girr_ΔlysF1b-LRM6	point mutation-nucleotide deletion	gene *35 (lysF1a)*	missense coding after A63 and premature stop at aa74: A63APRQPRNGFGD.STOP
Girr_ΔlysF1b-LRM7	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_ΔlysF1b-LRM8	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_ΔlysF1b-LRM11	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_ΔlysF1b-LRM12	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_ΔlysF1b-LRM14	point mutation-nucleotide substitution	gene *35 (lysF1a)*	G66D
Girr_ΔlysF1b-LRM15	point mutation-nucleotide insertion	gene *35 (lysF1a)*	stop after G66
Girr_ΔlysF1b-LRM16	point mutation-nucleotide substitution	gene *35 (lysF1a)*	A31T
NBJ_ΔlysF1b-LRM1	point mutation-nucleotide substitution	gene *33 (lysF1a)*	A68E
NBJ_ΔlysF1b-LRM2	point mutation-nucleotide substitution	gene *33 (lysF1a)*	A68E
NBJ_ΔlysF1b-LRM3	point mutation-nucleotide substitution	gene *33 (lysF1a)*	A68E

| NBJ_ΔlysF1b-LRM7 | point mutation-nucleotide substitution | gene *33* (*lysF1a*) | D9A |

When phages were recovered from the large plaque, serial diluted and replated, the wild type

plaque size was recovered across the full phage population (Figure 11B). This type of

experiment was repeated multiple times with Girr_ΔlysF1b and NBJ_ΔlysF1b and ten

independent Girr_ΔlysF1b-LRMs and four independent NBJ_ΔlysF1b-LRMs were recovered. All

Girr LRM mutants showed a large plaque phenotype compared to the parent Girr_ΔlysF1b (S13

Figure 12) and PCR analysis of the *lysF1a-lysF1b* genomic region confirmed that all the LRM

mutants contained the expected ΔlysF1b deletion (S13 Figure 12). To determine the location of

any nucleotide mutations within the genome of the LRM mutants, DNA was extracted and the

genomes were sequenced as detailed in Methods. In all cases, the Girr_ΔlysF1b-LRMs and

NBJ_ΔlysF1b-LRMs contained the expected *lysF1b* deletion but all phage also contained either

nucleotide insertions, deletions or substitutions within the *lysF1a* open reading frame that

altered the amino acid sequence of the LysF1a protein. A summary of all LRM mutants is

provided in Table 3 and a schematic showing the amino acid changes to the different

Girr_lysF1a-LRMs is presented in S14 Figure 13.

Six different mutations were identified in the ten Girr_ΔlysF1b-LRMs and two in the four

NBJ_ΔlysF1b-LRMs.The mutations mapped to three different domains of the LysF1a protein:

the N-terminal cytoplasmic domain, the first TMD or the beginning of the cytoplasmic C-terminal

domain. No mutants were recovered that impacted the second TMD or distal regions of the

cytoplasmic C-terminal domain. In the Girr LRM mutants, two different missense mutations were

found in TMD1 of LysF1a that result in R19H and A31T while one of the NBL LRM mutants

changed the charged cytoplasmic D9 to A. In addition, three G66D mutants were recovered in

the Girr LRMs, and three A68E mutants were recovered in the NBJ LRMs. All of these change a

small hydrophobic amino acid to a negatively charged amino acid just after the end of the TMD2

at the beginning of the cytoplasmic C-terminal domain. Finally, four mutants generate a

nonsense mutation that either truncates the Girr LysF1a protein after G66 or create a frame shift

after A63 that results in missense coding and a premature stop after amino acid 73. All LRM

mutations result in changes to the LyF1a protein that likely alter its conformation, thus, the

findings support that the LysF1a protein can in fact efficiently function in the lysis pathway and

that the protein does not exclusively function as an antiholin to the LysF1b as suggest earlier.

Importantly, plaques were also evaluated for LRM mutants after multiple rounds of plating of the

Δ*lysF1a/*Δ*lysF1b* double mutants, but even after six rounds of infection and plating, we failed to

identify any changes to plaque sizes, and this suggests that there are no other negative

regulators (putative antiholins) of the lysis pathway in these phages.

Lysis Recovery Mutants Exhibit Early Lysis and Reduced Burst Size

Since the LRMs generate a wild type plaque size (Figure 11A; S13 Figure 12), it was

pertinent to evaluate the lysis timing in liquid culture and determine lag time and burst size. *M.*

smegmatis was infected with Girr_WT, Girr_Δ*lysF1b* and three different Girr_Δ*lysF1b*-LRMs and

the OD_{600} of the culture measured as described in Methods. Figure 12A shows that Girr_WT

triggers lysis as expected at ~110-minutes while Girr_Δ*lysF1b* causes reduced culture growth

but no defined triggering as also shown in Figures 8 and 9. In stark contrast, Girr_Δ*lysF1b*-

LRM1*,* Girr_Δ*lysF1b*-LRM2 and Girr_Δ*lysF1b*-LRM16, all trigger lysis prematurely at ~90

minutes and ultimately reduce the OD_{600} to the same baseline as Girr_WT. When NBJ_WT,

NBJ_Δ*lysF1b*, NBJ_Δ*lysF1b*-LRM1 and NBJ_Δ*lysF1b*-LRM7 were tested in the liquid lysis

assay, the NBJ-LRMs also trigger lysis prematurely and reduce the OD_{600} to the same level as

NBJ_WT (Figure 11B). To validate the early triggering time and assess its impact on burst size,

one-step growth curves were completed as described in Methods. Figure 12C shows a

representative one-step experiment with Girr_WT and two LRM mutants. Phage release from

Girr_Δ*lysF1b*-LRM1 and Girr_Girr_Δ*lysF1b*-LRM2 is detected by 150 minutes and plateaus by

200-minutes, a time point at which Girr_WT has achieved <10% burst size. The two Girr LMRs

average burst size is also only 95.5 +/- 12.5 PFU/cell compared to 289.6 +/- 65 PFU/cell for the

Girr_WT. The reduced burst size is likely the result of lysis triggering well before all phage

progeny have been full assembled and is consistent with reports that have evaluated early

triggering in *E. coli* phages [11, 12, 76]. The current results also indicate that plaque size in our

experimental system is not a function of burst size since a >65% reduction in burst size is still

able to generate wild type-sized plaques (Figure 12A; S13 Figure 12). These findings add

further support to the hypothesis that the small plaque phenotypes of the various LysF1 mutants

is the result of inefficient cell wall lysis that hampers phage release.

Analysis of *lysF1b* Phage Plaque Sizes in *M. smegmatis* Lacking the Lipoarabinomannan

Layer of the Cell Wall

A recent study proposes a 1TMD protein termed LysZ may function as a spanin-like

protein to disrupt the integrity of the Gram-positive *Corynebacterium glutamicum* cell wall by

interacting with the lipoarabinomannan (LAM) layer [59]. In these studies, deletion of *lysZ* from

phages CL31 or Cog results in a small plaque phenotype that is rescued if phage were plated

on a *Corynebacterium glutamicum* strain that is defective in the synthesis the LAM. Since both

LysZ and LysF1b have a single TMD and deletion of the genes results in small plaques, it was

pertinent to determine whether an *M. smegmatis* strain lacking the mtpA enzyme required to

synthesize LAM (*M. smegmatis* ΔmtpA) [68], would complement the small plaque phenotype of

Girr_Δ*lysF1b*. Wild type *M. smegmatis*, *M. smegmatis* ΔmtpA, and *M. smegmatis* ΔmtpA-comp

(ΔmtpA carrying complementing mtpA expression plasmid), were grown to saturation and set to

OD_{600} of 2.0 prior to infection. Equal amounts of phage were used to infect and cells and

plaques evaluated after incubation for 48 hrs. at 37°C. Figure 13A shows that both wild type Girr

and Girr_Δ*lysF1b* exhibit a modest but significant increase in plaque size when infected into the

M. smegmatis ΔmtpA strain compared to the wild type *M. smegmatis* or the *M. smegmatis*

ΔmtpA-comp strain. Girr_WT plaques increased by 1.4-fold from 2.8mm to 3.9mm while

Girr_Δ*lysF1b* increased by 1.8-fold from 0.93mm to 1.7mm (Figure 13B). To account for the

possibility that the change is plaque size were caused by reduced cell growth and less *M.*

smegmatis ΔmtpA bacteria growing over the courses of the experiment, studies were repeated

using high density cultures at 2 x 10^8 CFU/infection. This level of bacteria will exacerbate small

plaque phenotypes since the increased cells enhance absorption and reduce phage diffusion

[28; 84-86]. In these studies, both Girr_WT and Girr_Δ*lysF1b* showed similar plaques size

increases as the low-density experiments verifying that *M. smegmatis* lacking a formal LAM

layer is generally less restrictive to lysis and phage release. Since both Girr_WT and

Girr_Δ*lysF1b* show increased plaque size and the plaques of Girr_Δ*lysF1b* do not reach

Girr_WT size, the results indicate that the lack of the LAM layer does not effectively complement

the deletion of *lyF1b*. This is consistent with the proposed holin function of the LysF1b and

supports the hypothesis that the LysF1b and LysZ proteins have distinct functions in the lysis

pathway in their respective bacterial hosts although holin function for LysZ has not been

formally evaluated.

Analysis of Additional Genes Proposed to be Involved in the Lysis Pathway of F1 Cluster

Phages

It is currently unknown how the various endolysins encoded by actinobacteriophages

gain access to the peptidoglycan layer following phage infection. Although exogenous

expression of the lysin A from some phages is cytotoxic to *M. smegmatis*, there is no evidence

of signal sequences or SAR domains in these proteins [21, 30--34]. However, *Mycobacterium*

sp. appear to have several different secretion systems that do not require a signal sequence

[80-83]. Additionally, it has been proposed that phage Ms6 encodes a small chaperone protein

(gp23) required for the export of Lysin A because the deletion of the gene results in a small

plaque phenotype [50, 52]. Since both phage Girr and NBJ encode a homolog to gp23 in the

same general genomic area upstream of the Lysin A (gene *29*; Figure 14A), it was of interest to

determine if deletion caused a lysis defect in Girr and NBJ. Figure 14D shows that Girr gp29 is

73 amino acids and shares 76.2% identity to Ms6 gp23 while NBJ gp29 shares 76 of 77 amino

acids with gp23. BRED was utilized to delete gene *29* from Girr and two independent Girr_Δ*29*

phage mutants were isolated (Figure 14B). Girr_Δ*29-20* and Girr__Δ*29-25* showed no lysis

defect when compared to Girr_WT in the liquid lysis assay (Figure 15C) and showed identical

plaque size to Girr_WT (Figure 14E). BRED deletion of NBJ *29* (NBJ_Δ*29*) also showed wild

type plaque size (Figure 14E) and no difference in liquid lysis when compared to NBJ_WT.

To explore the possible interaction of the Lysin A and gp29 further, the proteins were

modeled using AlphaFold3. The results presented in S15 Figure 14 to S17 Figure 16 show that

all models produced data with ipTM values <0.4 indicating a very low confidence in the

interactions of the soluble Lysin A and gp29 proteins. Thus, these results suggest that gp29

from phage Girr or NBJ is not involved in the lysis pathway. A plausible explanation for the lysis

defects observed in the Ms6 studies is that the method of deletion of gene *23* somehow

impacted the expression of Lysin A (gene *24*) to generate a lysis phenotype. Full genome

sequencing data of the Ms6 mutants was not presented or discussed [50, 52] and the

requirement of a phage encoded chaperone has not been identified in any other phage.

IMPLICATIONS AND FUTURE DIRECTIONS

The results presented in this report identify the novel LysF1a and LysF1b proteins as

holin-like components required for efficient lysis of *M. smegmatis* by F1 cluster phages. The

data support that LysF1a and LysF1b function together to time the lysis event and disrupt the

PMF, however, it appears that LysF1a requires the presence of the LysF1b protein to be active

as a putative holin. Holin function is based on several key findings. First, holin function for the

LysF1b protein is directly demonstrated by the ability of energy poisons to trigger premature

lysis in phages that do not express LysF1a (Figure 10). While it could be argued that the energy

poison might trigger lysis by releasing a lysin that was previously exported and tethered to the

outer membrane, this hypothesis is not supported by the fact that CN treatment does not trigger

lysis in the Δ*lysF1a*/Δ*lysF1b* double mutant. Second, holin function is supported for LysF1b

instead of releasin function because there is no defined SAR-endolysins in F1 cluster phages

and the energy poisons do not trigger early lysis by releasin in phage Mu [18]. Finally, holin

function is supported for LysF1a by the ability to recover LRM mutants in the Δ*lysF1b*

background of both Girr and NBJ that map to the LysF1a protein and create phage that trigger

lysis early but otherwise are fully lysis competent (Figure 12).

The 1TMD LysF1b proteins represent a new class of holins and appear to be found

predominantly in phages that infect Gram positive hosts. HHpred analysis shows that the

LysF1b proteins have a high probability hit to Pfam PF10874 (DUF2746) that contains 906

proteins. Within this Pfam are hits to phages that infect *Mycobacterium sp, Gordonia sp,* and

Streptomyces sp and hits to *Micromonospora sp, Rothia sp, Saccharothrix sp, Nocardia*

terpenica, Propionicmonas paludicola, Microbacterium talmoniae, Xylanimonas oleitrophica and

Actinacidiphilia glaucinigra where the later hits are likely to proteins that are encoded by

prophages within the host bacteria. To assess whether additional phages in the

Actinobacteriophage Database at PhagesDB.org express a LysF1b-like homolog [87], phages

infecting *M. smegmatis* were bioinformatically evaluated as described in Methods to identify the

lysis cassette and associated genes that encode proteins with 1TMD and 2TMD in the same

cassette. Of the ~2,600 phages in the database that infect *M. smegmatis*, ~50% (from 22 of the

32 clusters) encode a LysF1b-like protein with a predicted N-out-C-in membrane topology and

also have an associated gene encoding a protein with N-in-C-in 2TMD in a defined lysis

cassette (S18 Figure 17, S19 Figure 18). Of the remaining 50% of annotated *M. smegmatis*

phages, 30% encode a LysF1b-like homolog but do not have an associated gene that encodes

a 2TMD protein, but instead have a gene encoding a protein with 4TMD. Thus, ~80% of the

phages that infect *M. smegmatis* have a LysF1b homolog with the predicted N-out-C-in

topology. These lysF1a-like proteins are grouped to 20 different phams based on their amino

acid identity [88, 89]. Representative LysF1a-like proteins from each pham are presented in S20

Figure 19. The proteins differ in size from 92-150 amino acids, have several basic amino acids

after the TMD as highlighted in Figure 2 and have between 30%-50% DEKRH charged amino

acids in the C-terminal cytoplasmic domain (S20 Figure 19). Multiple sequence alignment

indicates minimal amino acid identity between the proteins (S21 Figure 20) and the alignment

also failed to identify any domains of high identity. Thus, while ~80% of the phages that infect

M. smegmatis in the PhagesDB.org database express a LysF1b-like protein, it is unclear

whether they all function like the LysF1b proteins from Girr and NBJ.

Since the F1 phages evaluated in this study can clearly perform lysis independent of the

LysF1a, and the double deletion of both *lysF1a* and *lysF1b* has the same phenotype as the

Δ*lysF1b*, what then is the function of the 2TMD LysF1a proteins in the lysis pathway? As

discussed throughout this report, a role of LysF1a as a type II pinholin is not supported by the

sequence data (Figure 1BC). In fact, the amino acid sequence data and lack of lysis function in

the absence of LysF1b, better supports LysF1a as an pure antiholin like P2 phage LysA [55,56].

However, deletion of the *lysF1a* gene results in a lysis triggering *delay* in both Girr_Δ*lysF1b* and

NBJ_ Δ*lysF1b* with the later delayed over 60 minutes compared to NBJ_WT (Figure 8). A lysis

delay is not consistent with the function of LysF1a as a pure antiholin to the LysF1b, since it

would be expected that deletion of the antiholin would result in *early* host lysis by Girr_Δ*lysF1a*

or Girr_Δ*lysF1a* since the LysF1b would not be inhibited from forming active multimers as

shown for lambda S_{105} and phage 21 $S^{21}68$ [10-12]. In this context, the key findings are 1) that

destabilization of the PMF alone is not sufficient to "activate" LysF1a into a lesion that supports

wild type lysis or phage release (Figure 8), and 2) that LRMs can be recovered in the Δ*lysF1b*

background that have point mutations LysF1a that result in a fully lysis competent LysF1a

protein that happens to trigger premature (i.e. unregulated) lysis (Figure 12). These findings

suggest that amino acid changes to the protein may induce conformational changes to the

LysF1a that allow it to fully support bacterial lysis and phage release in the absence of the

LysF1b. Thus, an attractive working model is that LysF1a is not an antiholin to LysF1b but

simply is not lysis active without LysF1b. Therefore, as expression of both proteins increases

following infection, they reach a concentration at which they can formally interact, and this

interaction facilitates conformational changes in the LysF1a that allow it to become active in

lysis. The two proteins then function together to facilitate the release of endolysins and/or phage

particles. Such a model would be like the holin/antiholin system in phage lambda and phage 21

where the phages express both active holins and inactive antiholins and triggering results in an

instantaneous conversion of all proteins to active holins [10-12].

Currently, it is unclear how the LysF1a and LysF1b proteins interact and whether both

proteins together actively participate in the creation a heteromeric lesion in the inner membrane.

Clearly, LysF1b alone can trigger lysis and initiate the process of phage release, but as stated

above, lysis timing and the efficiency of phage release are still not optimal. Phage SSP1

appears to have two distinct holins that support lysis when expressed exogenously, but it has

not been formally demonstrated that they both form a heteromeric lesion [40]. In addition, the

lysis pathway of phage P2 utilizes two distinct proteins, holin Y and antiholin LysA, but LysA

protein appears to be fully regulatory to holin Y and is not converted into an active holin like the

antiholins S_{107} or $S^{21}71$ [11, 57, 58]. Thus, the interaction of two distinct holin monomers that

have different membrane topologies will require direct study of LysF1a and LysF1b proteins

within membranes to resolve the structure. Such work could also directly address whether the

LysF1b also has interactions with cell wall components to facilitate phage release as suggested

for the LysZ proteins of *Corynebacterium* phages [59]. Whether the other Mycobacteriophages

that express different variants of the LysF1a and LysF1b proteins also utilize them in the same

way as Girr and NBJ will require additional deletion and complementation studies. The current

study therefore provides a framework to evaluate lysis under physiological conditions in Gram-

positive hosts. It is exciting to speculate that due to the presence of *lysF1*-like and *lysZ*-like

genes in so many phages, it may be that the phages that infect Gram-positive hosts use a

similar set of tools to complete the lysis pathway through distinct molecular mechanisms just as

observed for phages that infect *E. coli.*

LIMITATIONS

The focus of the current report was to genetically evaluate the function of LysF1a and

LysF1b proteins in the physiological context of phage infection in *M. smegmatis*. There are

several limitations to the current data set. First, Girr and NBJ are temperate phages and do not

kill the entire liquid cultures even at an MOI of 10:1. This is one reason that the culture OD_{600} in

our experiments does not drop below 0.10. Although we have shown that the residual surviving

bacteria is <10% of the starting culture and does not impact the ability to assess lysis timing, we

have a limited window of analysis of ~6-8 hrs. after infection before the lysogens increase the

OD_{600} and begin to interfere with the analysis. Thus, we don't have time points past 480 minutes

and it's possible that the various *lysF1b* mutants ultimately lyse the infected cells if allowed to

grow for 24 hrs. Second, due to the lysogens present in the experiments, the ability to evaluate

the infected bacteria via microscopy (especially electron microscopy) is challenging. Microscopy

data is therefore not presented since infected cells and lysogens can't be distinguished prior to

lysis and even after lysis the images have lysogens that interfere with the analysis. Third, the

LysF1 proteins are integral membrane proteins and AlphaFold3 modeling has reduced

confidence for membrane proteins and usually shows low confidence pTM scores of <0.5.

Additionally, using AlphaFold3 to assess protein-protein interactions for membrane proteins is

also poor with ipTM scores generally <0.5 (low confidence). Finally, the ability to exogenously

express LysF1a and LysF1b proteins from plasmids to assess possible interactions and

phenotypes in the *M. smegmatis* host is not possible due to the strong cytotoxicity of the LysF1b

[70, 71].

DATA AVAILABILITY STATEMENT

The authors affirm that all data necessary for confirming the conclusions of this article are represented fully within the article and its tables and figures. Full genome sequences of Girr and NBJ are available in GenBank through accession numbers MH669003 and MH399784, respectively. Extended data, including all original gel figures are posted to FileShare https://doi.org/10.6084/m9.figshare.31562248

ACKNOWLEDGMENTS

We are grateful to the members of the Science Education Alliance for their invaluable research support and for supplying phage and plasmid reagents, particularly Dainelle Heller, Viknesh Sivanathan, Graham Hatfull, Deborah Jacobs-Sera, Becky Garlena and Dan Russell. We also thank the Pittsburgh Bacteriophage Institute for genome sequencing of the phages described in this study. We want to thank A. McKitterick for helpful discussions regarding LysZ. We want to thank Y. Morita for generously sending the ΔmtpA *M. smegmatis* strains. We thank Welkin Pope for helpful discussions regarding infectious centers.

REFERENCS

1. Mushegian AR. Are There 10(31) Virus Particles on Earth, or More, or Fewer? J Bacteriol. 2020;202(9).

2. Oliveira H, Melo LD, Santos SB, Nobrega FL, Ferreira EC, Cerca N, et al. Molecular aspects and comparative genomics of bacteriophage endolysins. J Virol. 2013;87(8):4558-70.

3. Ryan GL, Rutenberg AD. Clocking out: modeling phage-induced lysis of Escherichia coli. J Bacteriol. 2007;189(13):4749-55.

4. Pang T, Fleming TC, Pogliano K, Young R. Visualization of pinholin lesions in vivo. Proc Natl Acad Sci U S A. 2013;110(22):E2054-63.

5. Savva CG, Dewey JS, Deaton J, White RL, Struck DK, Holzenburg A, et al. The holin of bacteriophage lambda forms rings with large diameter. Mol Microbiol. 2008;69(4):784-93.

6. Savva CG, Dewey JS, Moussa SH, To KH, Holzenburg A, Young R. Stable micron-scale
holes are a general feature of canonical holins. Mol Microbiol. 2014;91(1):57-65.
7. Summer EJ, Berry J, Tran TA, Niu L, Struck DK, Young R. Rz/Rz1 lysis gene equivalents in
phages of Gram-negative hosts. J Mol Biol. 2007;373(5):1098-112.
8. Rajaure M, Berry J, Kongari R, Cahill J, Young R. Membrane fusion during phage lysis. Proc
Natl Acad Sci U S A. 2015;112(17):5497-502.
9. Berry J, Rajaure M, Pang T, Young R. The spanin complex is essential for lambda lysis. J
Bacteriol. 2012;194(20):5667-74.
10. Young R. Phage lysis: three steps, three choices, one outcome. J Microbiol.
2014;52(3):243-58.
11. Cahill J, Young R. Phage Lysis: Multiple Genes for Multiple Barriers. Adv Virus Res.
2019;103:33-70.
12. Cahill J, Holt A, Theodore M, Moreland R, O'Leary C, Martin C, et al. Spatial and temporal
control of lysis by the lambda holin. mBio. 2024;15(2):e0129023.
13. Park T, Struck DK, Dankenbring CA, Young R. The pinholin of lambdoid phage 21: control
of lysis by membrane depolarization. J Bacteriol. 2007;189(24):9135-9.
14. Barenboim M, Chang CY, dib Hajj F, Young R. Characterization of the dual start motif of a
class II holin gene. Mol Microbiol. 1999;32(4):715-27.
15. Blasi U, Young R. Two beginnings for a single purpose: the dual-start holins in the
regulation of phage lysis. Mol Microbiol. 1996;21(4):675-82.
16. Ramanculov E, Young R. An ancient player unmasked: T4 rl encodes a t-specific antiholin.
Mol Microbiol. 2001;41(3):575-83.
17. Ramanculov E, Young R. Genetic analysis of the T4 holin: timing and topology. Gene.
2001;265(1-2):25-36.
18. Chamblee JS, Ramsey J, Chen Y, Maddox LT, Ross C, To KH, et al. Endolysin Regulation
in Phage Mu Lysis. mBio. 2022;13(3):e0081322.

19. Xu H, Bao X, Hong W, Wang A, Wang K, Dong H, Hou J, Govinden R, Deng B, Chenia HY.

Biological Characterization and Evolution of Bacteriophage T7-Δholin During the Serial

Passage Process. Font. Microbiology. 2021;12.

20. Raman SK, Siva Reddy DV, Jain V, Bajpai U, Misra A, Singh AK. Mycobacteriophages:

therapeutic approach for mycobacterial infections. Drug Discov Today.

2024;29(7):104049.

21. Payne KM, Hatfull GF. Mycobacteriophage endolysins: diverse and modular enzymes with

multiple catalytic activities. PLoS One. 2012;7(3):e34052.

22. Schmelcher M, Donovan DM, Loessner MJ. Bacteriophage endolysins as novel

antimicrobials. Future Microbiol. 2012;7(10):1147-71.

23. Arenskotter M, Broker D, Steinbach A. Biology of the metabolically diverse genus Gordonia.

Appl Environ Microbiol. 2004;70(6):3195-204.

24. Goodfellow M, Jones AL. Corynebacteriales ord. nov. Bergey's Manual of Systematics of

Archaea and Bacteria. 2015. p. 1-14.

25. Brennan PJ. Structure, function, and biogenesis of the cell wall of Mycobacterium

tuberculosis. Tuberculosis (Edinb). 2003;83(1-3):91-7.

26. Hart EM, Bernhardt TG. The mycomembrane. Curr Biol. 2025;35(3):R85-R6.

27. Catalao MJ, Pimentel M. Mycobacteriophage Lysis Enzymes: Targeting the Mycobacterial

Cell Envelope. Viruses. 2018;10(8).

28. Payne K, Sun Q, Sacchettini J, Hatfull GF. Mycobacteriophage Lysin B is a novel

mycolylarabinogalactan esterase. Mol Microbiol. 2009;73(3):367-81.

29. Marinelli LJ, Piuri M, Swigonova Z, Balachandran A, Oldfield LM, van Kessel JC, et al.

BRED: a simple and powerful tool for constructing mutant and recombinant

bacteriophage genomes. PLoS One. 2008;3(12):e3957.

30. Joshi H, Seniya SP, Suryanarayanan V, Patidar ND, Singh SK, Jain V. Dissecting the

structure-function relationship in lysozyme domain of mycobacteriophage D29-encoded

peptidoglycan hydrolase. FEBS Lett. 2017;591(20):3276-87.

31. Singh AK, Gangakhedkar R, Thakur HS, Raman SK, Patil SA, Jain V. Mycobacteriophage

D29 Lysin B exhibits promising anti-mycobacterial activity against drug-resistant

Mycobacterium tuberculosis. Microbiol Spectr. 2023;11(6):e0459722.

32. Pohane AA, Joshi H, Jain V. Molecular dissection of phage endolysin: an interdomain

interaction confers host specificity in Lysin A of Mycobacterium phage D29. J Biol Chem.

2014;289(17):12085-95.

33. Nair G, Jain V. An intramolecular cross-talk in D29 mycobacteriophage endolysin governs the

lytic cycle and phage-host population dynamics. Sci. Adv.2024;10(6):

34. Gil F, Catalao MJ, Moniz-Pereira J, Leandro P, McNeil M, Pimentel M. The lytic cassette of

mycobacteriophage Ms6 encodes an enzyme with lipolytic activity. Microbiology

(Reading). 2008;154(Pt 5):1364-71.

35. Gil F, Grzegorzewicz AE, Catalao MJ, Vital J, McNeil MR, Pimentel M. Mycobacteriophage

Ms6 LysB specifically targets the outer membrane of Mycobacterium smegmatis.

Microbiology (Reading). 2010;156(Pt 5):1497-504.

36. Loessner MJ, Gaeng S, Wendlinger G, Maier SK, Scherer S. The two-component lysis

system of Staphylococcus aureus bacteriophage Twort: a large TTG-start holin and an

associated amidase endolysin. FEMS Microbiol Lett. 1998;162(2):265-74.

36. Kwan T, Liu J, DuBow M, Gros P, Pelletier J. The complete genomes and proteomes of 27

Staphylococcus aureus bacteriophages. Proc Natl Acad Sci U S A. 2005;102(14):5174-

9.

38. Vybiral D, Takac M, Loessner M, Witte A, von Ahsen U, Blasi U. Complete nucleotide

sequence and molecular characterization of two lytic Staphylococcus aureus phages:

44AHJD and P68. FEMS Microbiol Lett. 2003;219(2):275-83.

39. Godinho LM, El Sadek Fadel M, Monniot C, Jakutyte L, Auzat I, Labarde A, et al. The

977   Revisited Genome of Bacillus subtilis Bacteriophage SPP1. Viruses. 2018;10(12).

978   40. Fernandes S, São-José C. Probing the function of the two holin-like proteins of

979   bacteriophage SPP1. Virology. 2017;500:184-189.

980   41. Escobedo S, Campelo AB, Wegmann U, Garcia P, Rodriguez A, Martinez B. Insight into the

981   Lytic Functions of the Lactococcal Prophage TP712. Viruses. 2019;11(10).

982   42. Sao-Jose C, Parreira R, Vieira G, Santos MA. The N-terminal region of the Oenococcus

983   oeni bacteriophage fOg44 lysin behaves as a bona fide signal peptide in Escherichia coli

984   and as a cis-inhibitory element, preventing lytic activity on oenococcal cells. J Bacteriol.

985   2000;182(20):5823-31.

986   43. Gindreau E, Lonvaud-Funel A. Molecular analysis of the region encoding the lytic system

987   from Oenococcus oeni temperate bacteriophage phi 10MC. FEMS Microbiol Lett.

988   1999;171(2):231-8.

989   44. Nelson D, Schuch R, Zhu S, Tscherne DM, Fischetti VA. Genomic sequence of C1, the first

990   streptococcal phage. J Bacteriol. 2003;185(11):3325-32.

991   45. Pope WH, Mavrich TN, Garlena RA, Guerrero-Bustamante CA, Jacobs-Sera D,

992   Montgomery MT, et al. Bacteriophages of Gordonia spp. Display a Spectrum of Diversity

993   and Genetic Relationships. mBio. 2017;8(4).

994   46. Pollenz RS, Bland J, Pope WH. Bioinformatic characterization of endolysins and holin-like

995   membrane proteins in the lysis cassette of phages that infect Gordonia rubripertincta.

996   PLoS One. 2022;17(11):e0276603.

997   47. McGarrah CEE, Algarin-Martinez ED, Cavasini MED, Correa V, Danielson DF, Dean WR, et

998   al. Isolation and Annotation of Azira, a CT Cluster Phage That Infects Gordonia

999   rubripertincta. Microbiol Resour Announc. 2023;12(7):e0034723.

48. Dedrick RM, Marinelli LJ, Newton GL, Pogliano K, Pogliano J, Hatfull GF. Functional

requirements for bacteriophage growth: gene essentiality and expression in

mycobacteriophage Giles. Mol Microbiol. 2013;88(3):577-89.

49. Sandoval AM, Abram AM, Alhabib ZM, Antonyan AS, Brikho SM, Buhay SI, et al. Complete

Genome Sequences of Cluster G Mycobacteriophage Darionha, Cluster A

Mycobacteriophage Salz, and Cluster J Mycobacteriophage ThreeRngTarjay. Microbiol

Resour Announc. 2020;9(20).

50. Catalao MJ, Gil F, Moniz-Pereira J, Pimentel M. Functional analysis of the holin-like proteins

of mycobacteriophage Ms6. J Bacteriol. 2011;193(11):2793-803.

51. Garcia M, Pimentel M, Moniz-Pereira J. Expression of Mycobacteriophage Ms6 lysis genes

is driven by two sigma(70)-like promoters and is dependent on a transcription

termination signal present in the leader RNA. J Bacteriol. 2002;184(11):3034-43.

52. Catalao MJ, Gil F, Moniz-Pereira J, Pimentel M. The mycobacteriophage Ms6 encodes a

chaperone-like protein involved in the endolysin delivery to the peptidoglycan. Mol

Microbiol. 2010;77(3):672-86.

53. Bavda VR, Jain V. Deciphering the Role of Holin in Mycobacteriophage D29 Physiology.

Front Microbiol. 2020;11:883.

54. Bavda VR, Yadav A, Jain V. Decoding the molecular properties of mycobacteriophage D29

Holin provides insights into Holin engineering. J Virol. 2021;95(10).

55. Kamilla S, Jain V. Mycobacteriophage D29 holin C-terminal region functionally assists in

holin aggregation and bacterial cell death. FEBS J. 2016;283(1):173-90.

56. Ford ME, Sarkis GJ, Belanger AE, Hendrix RW, Hatfull GF. Genome structure of

mycobacteriophage D29: implications for phage evolution. J Mol Biol. 1998;279(1):143-

64.

57. To KH, Dewey J, Weaver J, Park T, Young R. Functional analysis of a class I holin, P2 Y. J

Bacteriol. 2013;195(6):1346-55.

58. Christie GE, Calendar R. Bacteriophage P2. Bacteriophage. 2016;6(1):e1145782.

59. McKitterick AC, Lyerly EW, Bernhardt TG. Bacteriophages target membrane-anchored

glycopolymers to promote host cell lysis and progeny release. bioRxiv.

2025:2025.06.24.661397.

60. Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P, Nielsen H, et al.

DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural

networks. bioRxiv. 2022:2022.04.08.487609.

61. Tsirigos KD, Peters C, Shu N, Kall L, Elofsson A. The TOPCONS web server for consensus

prediction of membrane protein topology and signal peptides. Nucleic Acids Res.

2015;43(W1):W401-7.

62. Almagro Armenteros JJ, Tsirigos KD, Sonderby CK, Petersen TN, Winther O, Brunak S, et

al. SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat

Biotechnol. 2019;37(4):420-3.

63. Juncker AS, Willenbrock H, Von Heijne G, Brunak S, Nielsen H, Krogh A. Prediction of

lipoprotein signal peptides in Gram-negative bacteria. Protein Sci. 2003;12(8):1652-62.

64. Soding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology

detection and structure prediction. Nucleic Acids Res. 2005;33(Web Server

issue):W244-8.

65. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure

prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630(8016):493-

500.

66. Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez, R,

Butcher S. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024.

Nucleic acids research. 2024;52(W1):W521–W525.

67. Snapper SB, Melton RE, Mustafa S, Kieser T, Jacobs WR, Jr. Isolation and characterization

of efficient plasmid transformation mutants of Mycobacterium smegmatis. Mol Microbiol.

1990;4(11):1911-9.

68. Sparks IL, Kado T, Prithviraj M, Nijjer J, Yan J, Morita YS. Lipoarabinomannan mediates

localized cell wall integrity during division in mycobacteria. Nat Commun.

2024;15(1):2191.

69. Heller D, Amaya I, Mohamed A, Ali I, Mavrodi D, Deighan P, et al. Systematic

overexpression of genes encoded by mycobacteriophage Waterfoul reveals novel

inhibitors of mycobacterial growth. G3 (Bethesda). 2022;12(8).

70. Pollenz RS, Barnhill K, Biggs A, Bland J, Carter V, Chase M, et al. A genome-wide

cytotoxicity screen of cluster F1 mycobacteriophage Girr reveals novel inhibitors of

Mycobacterium smegmatis growth. G3 (Bethesda). 2024;14(5).

71. Wise B, Edwards K, Jirsa C, [student authors], Kanther M, Payne K, Pollenz RS, Heller DM.

Identification and characterization of host-modulating effectors encoded by the Cluster F1

mycobacteriophage NormanBulbieJr. bioRxiv 2025.

72. van Kessel JC, Hatfull GF. Mycobacterial recombineering. Methods Mol Biol. 2008;435:203-

15.

73. Gordon D, Abajian C, Green P. Consed: a graphical tool for sequence finishing. Genome

Res. 1998;8(3):195-202.

74. Russell DA. Sequencing, Assembling, and Finishing Complete Bacteriophage Genomes.

Methods Mol Biol. 2018;1681:109-25.

75. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.

Nat Methods. 2012;9(7):671-5.

76. Pang T, Park T, Young R. Mutational analysis of the S21 pinholin. Mol Microbiol.

2010;76(1):68-77.

77. Park T, Struck DK, Deaton JF, Young R. Topological dynamics of holins in programmed

bacterial lysis. Proc Natl Acad Sci U S A. 2006;103(52):19713-8.

78. Manohar P, Wan J, Ganser G, Sharp K, Young R. The Lysis cassette of jumbophage PhiKZ.

Sci Rep. 2026;16(1):5840.

79. Doermann AH. The intracellular growth of bacteriophages. I. Liberation of intracellular

bacteriophage T4 by premature lysis with another phage or with cyanide. J Gen Physiol.

1952;35(4):645-56.

80. Rigel NW, Gibbons HS, McCann JR, McDonough JA, Kurtz S, Braunstein M. The Accessory

SecA2 System of Mycobacteria Requires ATP Binding and the Canonical SecA1. J Biol

Chem. 2009;284(15):9927-9936.

81. Abdallah A, Gey van Pittius N, DiGiuseppe Champion P, Cox J, Luirink J, et. al. Type VII

secretion — mycobacteria show the way. Nat Rev Microbiol. 2007;5:883–891.

82. Bendtsen JD, Kiemer L, Fausbøll A, Brunak S. Non-classical protein secretion in bacteria.

BMC Microbiol. 2005;5:58.

83. Zhang Z, Dong L, Li X, Deng T, Wang Q. The PE/PPE family proteins of Mycobacterium

tuberculosis: evolution, function, and prospects for tuberculosis control. Front. Immunol.

2025;16:1606311.

84. Ameh EM, Tyrrel S, Harris JA, Pawlett M, Orlova EV, Ignatiou A, et al. Lysis Performance of

Bacteriophages with Different Plaque Sizes and Comparison of Lysis Kinetics After

Simultaneous and Sequential Phage Addition. Phage (New Rochelle). 2020;1(3):149-57.

85. Abedon ST, Yin J. Bacteriophage plaques: theory and analysis. Methods Mol Biol.

2009;501:161-74.

86. Dennehy JJ, Abedon ST, Turner PE. Host density impacts relative fitness of bacteriophage

Phi6 genotypes in structured habitats. Evolution. 2007;61(11):2516-27.

87. Russell DA, Hatfull GF. PhagesDB: the actinobacteriophage database. Bioinformatics.

2017;33(5):784-6.

88. Gauthier CH, Cresawn SG, Hatfull GF. 2022. PhaMMseqs: a new pipeline for constructing

phage gene phamilies using MMseqs2. G3 Genes Genomes Genetics.

2022;12(11):jkac233.

89. Gauthier CH, Hatfull GF. PhamClust: a phage genome clustering tool using proteomic

equivalence. mSystems. 2023;8(5):e00443-23.

1106

1107   **FIGURE LEGANDS**

1108

Figure 1. Lysis cassette organization of Girr and NormanBulbieJr (NBJ) and **amino acid**

sequence and topology of Girr and NBJ LysF1a proteins and comparison to phage 21 S^{21}

pinholin. **A.** Predicted protein coding genes are shown as colored boxes. The genomic location

is represented by the ruler, and the numbers represent the kilobase from nucleotide 1. Both

lysis cassette regions are located upstream of the *tape measure* and *minor tail* genes. Genes

above the genome are transcribed in the forward direct and genes below in the reverse

direction. Genes with the same color encode proteins that are grouped to the same pham. Pink

shading indicated nucleotide identity >90% in the shaded region. The *lysin A, lysin B,* 2TMD

lysF1a and 1TMD *lysF1b* genes are identified. NKF = genes with no know function; *HNH* = HNH

endonuclease. B. The amino acid sequences of Girr and NBJ LysF1a showing the location of

the predicted TMDs and charged amino acids. **C.** The amino acid sequence of phage 21 $S^{21}68$

holin, $S^{21}71$ antiholin and $S^{21}68_{IRS}$ showing the location of the predicted TMDs and charged

amino acids. **D.** Predicted membrane topology of Girr LysF1a, $S^{21}68_{IRS}$ and the proposed

topology of the $S^{21}68$ holin homodimer with cytoplasmic TMD1.

1123

Figure 2. Amino acid sequence and topology of lysis proteins with 1TMD. **A.** The amino

acid sequences of Girr LysF1b, NBJ LysF1b, Waterfoul gp32 and Mu releasin showing the

location of the predicted TMD and the charged amino acids. **B.** AlphaFold3 structural

predictions and membrane topologies of 1TMD proteins involved in lysis. pTM values: Mu, 0.51;

NBJ LysF1b, 0.40; Girr LysF1b, 0.41; Rz, 0.57; T, 0.74; phiKZ, 0.58. **C.** Predicted topologies of

Girr LysF1b and Waterfoul gp32.

1130

Figure 3. Deletion of Girr *lysF1a* and *lysF1b*. A. Genomic structure and PCR primer set

locations in Girr gene deletion mutants. Primer set 1 (P1) uses primers Girr-FL_*lysF1a* (F and

R) and generates a 755bp fragment in Girr_WT and 487bp with *lysF1a* deletion. Primer set 2

(P2) uses primers Girr-FL_*lysF1b* (F and R) and generates a 933bp fragment in Girr_WT and

594bp with *lysF1b* deletion. Primer set 3 (P3) uses the forward Girr-FL_*lysF1a* and reverse Girr-

FL_*lysF1b* and generates a 1189bp fragment in Girr_WT and either 921bp with *lysF1a* deletion

or 818b with *lysF1b* deletion. **B.** PCR verification of high titer lysates of Girr_WT, Girr_Δ*lysF1a*

and Girr_Δ*lysF1b* using the indicated primer sets. PCR band sizes are identified and show

correct deletions of the indicated genes.

1140

1141   **Figure 4 Plaque sizes in Girr_WT, Girr_Δ*lysF1a* and Girr_Δ*lysF1b*.** *M. smegmatis* was

1142   infected with identical numbers of the indicated phage as detailed in Methods and plaque size

1143   determined after 36 hrs. of growth at 37°C. Scale = 1cm. Plaque volume schematics are

1144   presented to illustrate the plaque size differences.

1145

Figure 5. Complementation of plaque phenotype in Girr_Δ*lysF1b*. A. Plaque assays were

completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1b* as described in Methods. Plaque

assays were completed with *M. smegmatis* complementing strains carrying the indicated

pExTra-Waterfoul expression plasmids using Girr_Δ*lysF1b* as described in Methods. Plaques

were evaluated following incubation at 37°C for 36 hrs. Plaques sizes were quantified using

ImageJ and the average diameter +/- standard deviation for > 50 plaques is shown. * =

statistically different from Girr_WT, p *<0.0001*. ** = statistically different from Girr_Δ*lysF1b*

infected on *M. smegmatis* with pExTra Waterfoul_*32* with no aTc, p *<0.0001*. Scale = 1cm. B.

Plaque assays were completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1b* as described

in Methods. Plaque assays were completed with *M. smegmatis* complementing strains carrying

the indicated pExTra-Waterfoul expression plasmids using Girr_Δ*lysF1b* as described in

Methods. Plaques were evaluated following incubation at 37°C for 48 hrs. Plaques sizes were

quantified using ImageJ and the average diameter +/- standard deviation for > 50 plaques is

shown * = statistically different from Girr_WT, *p <0.0001*. ** = statistically different from

Girr_Δ*lysF1b* infected on *M. smegmatis* with pExTra01, *p <0.0001*. Scale = 1cm.

1161

Figure 6. Complementation of plaque phenotype in Girr_Δ*lysF1a*. Plaque assays were

completed with *M. smegmatis* using Girr_WT or Girr_Δ*lysF1a* as described in Methods. Plaque

assays were completed with *M. smegmatis* complementing strains carrying pExTra-Girr_*34*

expression plasmid using Girr_Δ*lysF1a*as described in Methods. Plaques were evaluated

following incubation at 37°C for 48 hrs. Plaques sizes were quantified using ImageJ and the

average diameter +/- standard deviation for > 50 plaques is shown. * = statistically different from

Girr_WT, *p <0.0001*. ** = statistically different from Girr_Δ*lysF1a* infected on *M. smegmatis* with

pExTra-Girr_*34* with no aTc, *p <0.0001*. Scale = 1cm.

1170

Figure 7. Liquid lysis assay and ATP release from *M. smegmatis* after infection. At T_0 a log

growth culture of *M. smegmatis* at an OD_{600} of ~0.25 was infected with the indicated phage at

an MOI of 10:1 and incubated at 37°C for 30 minutes without shaking. Control cultures (no

phage) did not receive phage. At T_{30} minutes the culture was constantly shaken a 225rpm,

aliquots removed at the indicated times and bacterial growth evaluated at OD_{600}. Open white

arrow shows the 30-minute infection time. **A.** Representative results following infection with

Girr_WT, Girr_Δ*lysF1a* and Girr_Δ*lysF1b*. **B**. Representative results following infection with

NBJ_WT, NBJ_Δ*lysF1a* and NBJ_Δ*lysF1b.* **C.** Samples of culture were removed from the

cultures represented in A and B at the times indicated by the solid black triangle on the x axis.

The presence of ATP was evaluated as described in Methods.

1181

Figure 8. Energy poisons trigger early lysis by phage expressing LysF1b but not LysF1a.

At T_0 a log growth culture of *M. smegmatis* at an OD_{600} of ~0.25 was infected with the indicated

phage at an MOI of 10:1 and incubated at $37^{\circ}C$ for 30 minutes without shaking. Control cultures

(no phage) did not receive phage. At T_{30} minutes the culture was constantly shaken a 225rpm,

aliquots removed at the indicated times and bacterial growth evaluated at OD_{600}. **A**.

Representative results following infection with Girr_WT, and Girr_$\Delta lysF1a$. KCN (final

concentration of 10mM) was added to a culture infected with Girr_$\Delta lysF1a$ and an uninfected

culture at 110 minutes (open arrow). **B**. Representative results following infection with NBJ_WT,

and NBJ_$\Delta lysF1a$. KCN (final concentration of 10mM) was added to a culture infected with

NBJ_$\Delta lysF1a$ and uninfected culture at 120 minutes (open arrow). **C.** Representative results

following infection with Girr_WT, and Girr_$\Delta lysF1b$. KCN (final concentration of 10mM) or $CHCl_3$

(final concentration 1%) was added to cultures infected with Girr_$\Delta lysF1b$ and uninfected

cultures at 180 minutes (open arrow). **D**. Representative results following infection with

NBJ_WT, and NBJ_$\Delta lysF1b$. KCN (final concentration of 10mM) or $CHCl_3$ (final concentration

1%) was added to cultures infected with NBJ_$\Delta lysF1b$ and uninfected cultures at 180 minutes

(open arrow).

Figure 9. One-step growth curves of *M. smegmatis* infected with Girr_$\Delta lysF1a$ and

Girr_$\Delta lysF1b$. A. One-step growth curves using Girr_WT, Girr_$\Delta lysF1a$ and Girr_$\Delta lysF1b$ were

completed as described in Methods. Each time point represents the average and standard

deviation from 3 different experiments. **B**. The data for Girr_$\Delta lysF1b$ from panel A replotted with

a different Y axis than used in A. **C**. Burst size was determined as described in Methods. The

average bust size +/- standard deviation is from three different experiments and represents the

average of all points once the curve reached a plateau.

Figure 10. Deletion of both lysF1a and lysF1b in NBJ. A. Genomic structure and PCR primer

set locations in NBJ_WT and NBJ_ΔlysF1a/ΔlysF1b mutant. **B**. PCR verification of

lysF1a/lysF1b double deletion. Primer set 7 (P7) uses the forward NBJ-FL_*lysF1a* and reverse

NBJ-FL_*lysF1b* and generates a 1179bp fragment in NBJ_WT, a 985bp fragment in

NBJ_ΔlysF1a, 808bp fragment in NBJ_ΔlysF1b and a 580bp fragment in NBJ_ΔlysF1a/ΔlysF1b.

C. *M. smegmatis* was infected with identical numbers of the indicated phage as detailed in

Methods and plaque size determined after 36 hrs. of growth at 37^{o}C. Scale = 1cm. **D**. Liquid

lysis assay. At T$_0$ a log growth culture of *M. smegmatis* at an OD$_{600}$ of ~0.25 was infected with

the indicated phage at an MOI of 10:1 and incubated at 37^{o}C for 30 minutes without shaking

(indicated by white arrow). Control cultures (no phage) did not receive phage. At T$_{30}$ minutes the

culture was constantly shaken a 225rpm, aliquots removed at the indicated times and bacterial

growth evaluated at OD$_{600}$. A separate culture of non-infected cells and NBJ_WT was treated

with KCN (final concentration 10mm) at 100 minutes (black arrow). A separate culture of non-

infected cells, NBJ_ΔlysF1b and NBJ_ΔlysF1a/ΔlysF1b was treated with KCN (final

concentration 10mm) at 150 minutes (black arrow).

1222

Figure 11. Isolation of Girr_ΔlysF1b-LRM mutants. **A**. *M. smegmatis* was infected with

GirrΔlysF1b, plate lysates recovered from the plaque assay and *M. smegmatis* infected with the

recovered phage. Panel A shows a representative plaque assay following the second round of

infection. A large plaque is circled. **B**. Phage were isolated from the large plaque in A, serial

diluted and used for plaque assay. Plates were incubated at 37^{o}C for 36 hrs. **C**. Plaque assay

from Girr_WT phage. Plates were incubated at 37^{o}C for 36 hrs. Scale bar = 1cm in all images.

1229

Figure 12. Liquid lysis and one-step assay of Girr and NBJ LRM mutants. At T$_0$ a log

growth culture of *M. smegmatis* at an OD$_{600}$ of ~0.25 was infected with the indicated phage at

MOI 10:1 for 30 minutes at 37°C without shaking. Control cultures (no phage) did not receive

phage. At T_{30} minutes the culture was constantly shaken a 225rpm and aliquots removed over

time. *M. smegmatis* growth was evaluated at OD_{600}. **A**. Representative results following infection

with Girr_WT and indicated LRM mutants. **B**. Representative results following infection with

NBJ_WT and indicated LRM mutants **C**. One-step growth curves using Girr_WT and

Girr_Δ*lysF1b*-LRM1 and Girr_Δ*lysF1b*-LRM2 were completed as described in Methods. Each

time point represents the average and standard deviation from 3 different experiments.

1239

Figure 13. Analysis of *lysF1b* phage plaque sizes in *M. smegmatis* lacking the

lipoarabinomannan layer. A. Representative plaque assays using Wild type *M. smegmatis, M.*

smegmatis ΔmtpA-comp or *M. smegmatis* ΔmtpA cultures infected with the same amount of

Girr_WT or Girr_Δ*lysF1b* and incubated for 48 hrs. at 37°C. Ruler = 1cm. **B**. Plaque sizes were

determined as described in Methods. The average plaque diameter +/- standard deviation for 25

plaques is shown.

1246

Figure 14. Analysis of NKF gene *29* in Girr and NBJ. A. Gene organization of Girr, NBJ and

Ms6 lysis cassette region. Predicted protein coding genes are shown as colored boxes. The

genomic location is represented by the ruler, and the numbers represent the kilobase from

nucleotide 1. Genes above the genome are transcribed in the forward direct and genes below in

the reverse direction. Genes with the same color encode proteins that are grouped to the same

pham. Pink shading indicated nucleotide identity >90% in the shaded region. *NKF* = Girr_*29*,

NBJ_*29* and Ms6_*23*. **B**. PCR analysis of Girr_Δ*29*-20 and Girr_Δ*29*-25. **C**. Liquid lysis assay of

the indicated phages was carried out as described in Methods. **D**. Amino acid alignment of Girr

gp29, NBJ gp29 and Ms6 gp23 by Clustal Omega. **E**. Plaque assay of Girr_WT, Girr_Δ*29-20*,

Girr Δ*29-25*, NBJ_WT, and NBJ_Δ*29.* Ruler = 1cm. Plaque sizes were determined as described

in Methods. The average plaque diameter +/- standard deviation for 25 plaques is shown.

1258

1259

SUPPLEMENTAL INFORMATION

S1 Table 1. DNA gBLOCKS and Primers.

S2 Figure 1. One-step growth curves schematic protocol.

S3 Figure 2. One-step example data set: titers, adsorption and infectious centers.

S4 Figure 3. One-step example data set: Girr_WT.

S5 Figure 4. AlphaFold3 predictions of Girr and NBJ Lysin A.

S6 Figure 5. AlphaFold3 predictions of Girr LysF1a and phage 21 $S^{21}68$.

S7 Figure 6. Deletion of NBJ *lysF1a* and *lysF1b.*

S8 Figure 7. Infection and adsorption of Girr_WT and TMD mutants.

S9 Figure 8. Plaque sizes in NBJ_WT and *lysF1a* and *lysF1b* deletion mutants.

S10 Figure 9. Genomic context of lysis cassette genes in Girr and NBJ.

S11 Figure 10. Energy poisons and $CHCl_3$ treatment trigger early lysis by Girr_WT and

NBJ_WT.

S12 Figure 11. Cell viability of *M. smegmatis* infected with NBJ phages.

S13 Figure 12. PCR analysis of Girr LRM variants and plaques.

S14 Figure 13. Schematic of Girr_lysF1a-LRM amino acid sequences and membrane topology.

S15 Figure 14. AlphaFold3 analysis of Ms6 chaperone and Lysin A interactions.

S16 Figure 15. AlphaFold3 analysis of Girr gp29 and lysin A interactions.

S17 Figure 16. AlphaFold3 analysis of NBJ gp29 and Lysin A interactions.

S18 Figure 17. Part I: Identification of genes that encode *lysF1a* and *LysF1b* homologs in

phages that infect *M. smegmatis.*

S19 Figure 18. Part II: Identification of genes that encode *lysF1a* and *LysF1b* homologs in

phages that infect *M. smegmatis.*

S20 Figure 19. Identification of LysF1b homologs in phages that infect *M. smegmatis.*

S21 Figure 20. . Percent identity of LysF1b homologs

A.

MIWESVREAVDAAYQPDDGIDL**IGLLIIGLPSTIAAIGTGIVGV**LTV**R**GQ**RK**G**R**ERARQIDAKTDEIHEQTVNTHDTNMRDDLDEIRDLVRDGFKQIQRDIGGLREELRTERLERIEGDKRRD Girr gp35 LysF1b

MIWESVREAVDAAYQPDDGIJDL**IGLLIIGLPSTIAAIGTGIVGV**LTV**R**GQ**RK**G**R**ERARRIDAKTYEIHEQTVNTHDTNMRDDLDEIRDLVRDGFKQIQRDIGGLREELRTERLERIEGDKRRD NBJ gp33 LysF1b

MTGSAVSALPSD**WYGLAGLAVVNAAGLLALFM****K**AR**G****HK**QVANS**H**STNLRDDIDGLRDHIVELGRDIGGMRTDVGQLRGEVRDDRTHTHHRFRRIEDKLDELAGRGARESG Waterfoul gp32

M**DLISVLALWPYLLPVVAGGAVWA**M**RR**SFASTER**VE**RLENRMTEMETRYASIPGTEDVHEMRLQIAELSGDIRVLSQRVQSFSHQLELLLENAVN**R**SNS Mu releasin

NH₂ —————————[**TMD1**]—[**basic**]—[**KRHDE residues**]— COOH

Green = TMD domain	Basic region after TMD	+ Charged	- Charged

B.

Mu releasin

Girr gp35 LysF1b

NBJ gp33 LysF1b

T4
T
Type IIIa holin

phiKZ
Type IIIb holin

Lambda
Rz
spanin

extracellular/periplasmic

cytoplasmic

30/80 (31%) charged

24/75 (32%) charged

40/79 (50%) charged

C.

Girr gp35 LysF1b (124aa)

Waterfoul gp32 (111aa)

(1)M
extracellular
(22)
TMD
cytoplasmic
(45)
COOH
(124)
40/79 (50%) charged

M(1)
(12)
TMD
(36)
COOH
(111)
36/75 (48%) charged

Plaque volume comparisons

10mm

Girr_WT

Girr_ΔlysF1a

Girr_ΔlysF1b

Girr_WT

Girr_ΔlysF1a

Girr_ΔlysF1b

Girr_WT
M. Smegmatis
4.56 +/- 0.65 mm

Girr_*ΔlysF1a*
M. Smegmatis
pExTra-Girr*34* + *aTc*
**4.15 +/- 0.55 mm

Girr_*ΔlysF1a*
M. Smegmatis
*2.79 +/- 0.55mm

Girr_*ΔlysF1a*
M. Smegmatis
pExTra-Girr*34 no aTc*
2.83 +/- 0.353 mm

A.

Legend:
- ● Girr_WT
- ▲ Girr_ΔlysF1b
- □ Girr_ΔlysF1a

Y-axis: T_N/T_{45} (x 10^4)
X-axis: Time After Start of Infection (minutes)

B.

Legend:
- ▲ Girr_ΔlysF1b

Y-axis: T_N/T_{45} (x 10^4)
X-axis: Time After Start of Infection (minutes)

C.

phage	Burst Size PFU/cell
Girr_WT	277 +/- 81
Girr_ΔlysF1a	255 +/- 75
Girr_ΔlysF1b	*29.2 +/- 36

*statistically significant from WT phage *p<0.0001*.

A.

NBJ_WT

NBJ_ΔlysF1a/ΔlysF1b

Primer set 7 (P7): NBJ-FL_lysF1a F and Girr-FL_lysF1b R

B.

Primer set 7

C.

NBJ_WT
3.10 +/- 0.41nm

NBJ_ΔlysF1b
*0.98 +/- 0.19nm

NBJ_ΔlysF1a/ΔlysF1b
*0.93 +/- 0.18nm

D.

○ No phage
◐ No phage + CN
□ NBJ_WT
● NBJ_WT + CN
△ NBJ_ΔlysF1b
▲ NBJ_ΔlysF1b + CN
◇ NBJ_ΔlysF1a/ ΔlysF1b
◆ NBJ_ΔlysF1a/ ΔlysF1b + CN

A.

Girr_ΔlysF1b

B.

Girr_ΔlysF1b-LRM
3.01 +/- 0.26mm

C.

Girr_WT
2.94 +/- 0.25mm

A.

M. Smegmatis Girr_WT	ΔmtpA-Comp Girr_WT	ΔmtpA Girr_WT

M. Smegmatis Girr_Δ*lysF1b*	ΔmtpA-Comp Girr_Δ*lysF1b*	ΔmtpA Girr_Δ*lysF1b*

B.

*   * *p <0.0001* to WT
*   ** *p <0.0001* to *M smeg* WT
*   *** *p <0.0001* to *M smeg* Δ*lysF1b*

A.

Girr

NKF NKF NKF *NKF* NKF

27 30

26 28 29 31 *lysin A* 32 *HNH* 33 *lysin B* 34 *lysF1a* 35 *lysF1b* 36

25 26 27 28 29 30

Ms6

NKF NKF NKF *NKF* NKF

19 21 24 *lysin A* 26 *lysF1a* 28

20 22 23 25 *lysin B* 27 *lysF1b*

20 21 22 23 24 25

NBJ

NKF NKF NKF *NKF* NKF

24 27 30 *lysin A* 32 *lysF1a* 34

26 28 29 31 *lysin B* 33 *lysF1b*

25 26 27 28 29

NKF

B.

M P4 P3 P4 P3 P3 P4

1000bp
500bp
100bp

Girr_WT Girr_WT Girr_Δ29-20 Girr_Δ29-20 Girr_Δ29-25 Girr_Δ29-25

▶◀ Primer set 3 (P3): Girr-FL_lysF1a F and Girr-FL_lysF1b R

▶◀ Primer set 4 (P4): Girr-FL_29 F & R

C.

○ No phage
● Girr_WT
▲ Girr_Δ29-20
△ Girr_Δ29-25

OD600 vs Time After Start of Infection (minutes)

D.

Girr_29 MDRLGIILLKLLGPLADRIADRIADRITENLPDLSDLDDQIVAKLPDLTNLPEQVINIIGRLPRFPFLLGGKP

Ms6_23   MDRLGIVLLKLFGPLADRIADRIADRIAENLPDLSNLDDQIVAKLPDLTNLPAQVMDIIDSALRSIPVLGGILGSKR

NBJ_29   MDRLGIILLKLFGPLADRIADRIADRITENLPDLSNLDDQIVAKLPDLTNLPAQVMDIIDGALRSIPVLGGILGSKR

E.

Girr_WT
3.11 +/- 0.12 nm

Girr_Δ29-20
2.98 +/- 0.16 nm

Girr_Δ29-25
3.02 +/- 0.14 nm

NBJ_WT
2.25 +/- 0.25 nm

NBJ_Δ29
2.15 +/- 0.09 nm